Control of a type III-Dv CRISPR-Cas system by the transcription factor RpaB

and interaction of its transcribed leader with the DEAD-box RNA helicase CrhR

Raphael Bilger[1], Angela Migur[1†], Alexander Wulf[2,3], Claudia Steglich[1], Henning

Urlaub[2,3] and Wolfgang R. Hess[1*]

[1]University of Freiburg, Faculty of Biology, Genetics and Experimental Bioinformatics,

Schänzlestr. 1, D-79104 Freiburg, Germany

[2]Bioanalytics Research Group, Department of Clinical Chemistry, University Medical

Centre, D-37075, Göttingen, Germany;

[3]Bioanalytical Mass Spectrometry Group, Max Planck Institute for Biophysical

Chemistry, D-37077 Göttingen, Germany;

[†]Present address: RNA Synthetic Biology Group, Helmholtz Institute for RNA-based

Infection Research (HIRI), Würzburg, Germany

*Corresponding author: Wolfgang R. Hess; Phone: 0049-761-2032796; Fax: 0049-

761-2032745; Email: wolfgang.hess@biologie.uni-freiburg.de

Keywords: CRISPR-Cas systems, CrhR DEAD-box RNA helicase, CRISPR leader,

cyanobacteria, regulation of CRISPR-Cas gene expression; RNA structure,

transcription factor RpaB

Abbreviations: Cas proteins, CRISPR-associated proteins; CRISPR, Clustered

Regularly Interspaced Short Palindromic Repeats; crRNAs, CRISPR RNAs; CrhR,

cyanobacterial RNA helicase Redox; HLR1 motif, high light regulatory 1 motif; HNS,

Histone-like Nucleoid Structuring Protein; RpaB, regulator of phycobilisome
association B; TSS, transcriptional start site

ABSTRACT

CRISPR-Cas systems in bacteria and archaea provide powerful defense against
phages and other foreign genetic elements. The principles of CRISPR-Cas activity are
well understood, but less is known about how their expression is regulated. The
cyanobacterium *Synechocystis* sp. PCC 6803 encodes three different CRISPR-Cas
systems. The expression of one of them, a type III-Dv system, responds to changes
in environmental conditions, such as nitrogen starvation or varying light intensities.
Here, we found that the promoter of the respective six-gene *cas* operon is controlled
by the light- and redox-responsive transcription factor RpaB. RpaB binds to an HLR1
motif located 53 to 70 nt upstream of the transcription start site, resulting in
transcriptional activation at low light intensities. However, the strong promoter driving
transcription of the cognate repeat-spacer array is not controlled by RpaB. Instead, we
found that the 125 nt transcribed leader is bound by the redox-sensitive RNA helicase
CrhR. Cross-linking coupled to mass spectrometry analysis revealed six residues
involved in the CrhR-RNA interaction. Of these, L103, F104, H225, and C371 were
predicted to be on the surface of a dimeric CrhR model, while C184 was not and P443
in the very C-terminal region could not be assigned to a structural element. These
results show that the expression of the CRISPR-Cas system is linked to the redox
status of the photosynthetic cyanobacterial cell at two different levels. While RpaB
affects transcription, CrhR interacts with the transcribed leader post-transcription.
These results highlight the complex interplay between a CRISPR-Cas system and its
host cell.

Introduction

CRISPR–*cas* systems encode RNA-based adaptive and inheritable immune systems in many archaea and bacteria[1,2], which are highly diverse and were classified into two classes, six types and 33 subtypes[3]; however, new subtypes are still being discovered. Type III CRISPR-Cas systems, characterized by the presence of the signature gene *cas10*, exist in 34% and 25% of archaeal and bacterial genomes that encode CRISPR-Cas loci, respectively[4]. Type III systems are further classified into the five subtypes A to E[3,5,6]. Although detailed insights have been obtained regarding the molecular mechanisms and peculiarities of the different types of CRISPR-Cas systems, knowledge about their regulation of expression has remained fragmentary.

In the subtype I-E system of *E. coli*, regulation by transcription factors has been demonstrated. The DNA-binding protein HNS (Histone-like Nucleoid Structuring Protein) acts as a repressor by inhibiting the expression of crRNA and *cas* genes[7]. As an antagonist of HNS, LeuO activates the expression of *cas* genes, thereby enhancing the resistance against invading DNA[8]. Finally, a signaling cascade was identified that involves the BaeSR two-component regulatory system, which senses envelope stress (e.g., phage attack) by the membrane-localized kinase BaeS. Once activated, BaeS phosphorylates the cytoplasmic transcription factor BaeR[9], which, among others, activates the expression of *cas* genes in *E. coli*[10].

In the thermophilic archaeon *Sulfolobus islandicus*, expression of the type I-A CRISPR locus is regulated by Csa3a and Csa3b, two transcriptional regulators containing CARF and HTH domains. While Csa3a activates the expression of adaptation genes and the CRISPR array[11], interference genes are repressed by Csa3b. Repression is achieved in the absence of viral infection by co-binding of the cascade complex[12].

Further regulatory mechanisms have been described for CRISPR/Cas systems in

Pectobacterium atrosepticum[13] and *Pseudomonas aeruginosa*[14,15].

Cyanobacteria are the only prokaryotes whose physiology is based on oxygenic

photosynthesis, making them immensely important primary producers. Field studies

have shown that both cyanobacterial cell counts and the number of co-infecting

bacteriophages (cyanophages) can be very high, with up to 50% of all cyanobacteria

estimated to be infected at any one time[16] and likely affect cyanobacterial

biogeography and biogeochemistry at the scale of oceanic subregions[17]. Accordingly,

active defense mechanisms can be expected in cyanobacteria.

The unicellular cyanobacterium *Synechocystis* sp. PCC 6803 (from here:

Synechocystis 6803) is a model for the CRISPR biology of cyanobacteria. It possesses

three separate and complete CRISPR-Cas subtype I-D (CRISPR1), III-Dv (CRISPR2),

and III-Bv (CRISPR3) systems that are highly expressed and active in interference

assays[3,18–24]. From these, the CRISPR2 system is of particular interest as it has

recently been suggested to function as a protein-assisted ribozyme[24].

Each of the three CRISPR-*cas* loci in *Synechocystis* 6803 is associated with one gene

that has been suggested as a regulator (genes *sll7009*, *sll7062* and *sll7078*[18]). Indeed,

deletion of *sll7009*, encoding a putative WYL domain protein, led to increased

accumulation of crRNAs in the CRISPR1 system, but did not change the crRNA levels

of the other two systems[25]. This result was consistent with the observation that CARF

and WYL domain regulatory proteins are widely distributed ligand-binding specific

regulators of CRISPR-*cas* systems[26]. Sll7062 differs from the other two possible

regulators by the presence of an N-terminal CARF7 family domain fused to a RelE

RNase domain, a setup characteristic of Csm6 proteins. Csm6 proteins are not

transcription factors but CRISPR-associated RNases that are activated by cyclic

oligoadenylate (cOA)-mediated signaling[27]. Accordingly, Sll7062 was renamed

SyCsm6 when its activity was tested upon production as a recombinant protein,

together with the CARF-HEPN domain protein SyCsx1 (Slr7061)[28]. Therefore, the

CRISPR2 system actually lacks an obvious candidate for a regulatory gene in its

vicinity.

However, when we characterized the regulon controlled by the transcription factor

RpaB we noticed the possible involvement of a host genome-encoded factor in

CRISPR2 regulation[29]. RpaB ("regulator of phycobilisome association B", Slr0947) is

an OmpR-type transcription factor that was predicted to control more than 150

promoters via binding to the HLR1 ("high light regulatory 1") motif, a pair of imperfect

8-nt long direct repeats (G/T)TTACA(T/A) (T/A) separated by two random nucleotides.

RpaB mediates the activation of transcription when the HLR1 motif is located in a

window 45 to 66 nt upstream of the transcription start site (TSS), whereas all other

locations mediate repression[29]. The results showed that RpaB is a transcription factor

of central importance for the light- and redox-dependent remodeling of the

photosynthetic apparatus and many associated pathways. Surprisingly, there was also

a predicted binding site in the promoter that drives the transcription of the *cas* gene

operon of CRISPR2, the III-D system in *Synechocystis* 6803, but this was not

investigated further.

Another protein with a central role in light- and redox-dependent responses in

Synechocystis 6803 is the cyanobacterial RNA helicase Redox (CrhR)[30]. CrhR

(Slr0083) is the single DEAD-box RNA helicase in *Synechocystis* 6803 that is capable

of altering RNA secondary structures by catalyzing double-stranded RNA unwinding

as well as annealing[31]. The molecular effects of *crhR* deletion or inactivation have

been studied at the transcriptome[32,33] and proteome levels[34], and several attempts

have been made to identify the RNA targets of CrhR directly[35].

Here, we applied a further approach to pulldown RNA interacting with CrhR expressed

as a recombinant protein, and found that the transcribed leader of the type III-Dv

CRISPR-Cas system was co-purified. Therefore, we investigated the possible

regulatory impact of the host genome-encoded transcription factor RpaB on the

expression of the CRISPR2 system, and described and validated the interaction of

CrhR with the transcribed leader of the repeat-spacer array of the same system.

Results

Expression of the type III-Dv CRISPR2 system in* Synechocystis *6803 is affected

by environmental conditions

In our previous analysis of the distribution of putative HLR1 binding sites for the

transcription factor RpaB in *Synechocystis* 6803, one site was predicted in the

CRISPR2 *cas* gene promoter, however, this had not been studied further[29]. This

promoter drives the transcription of six genes, *sll7067* to *sll7062*, into a single

transcriptional unit (TU)[36]. Therefore, these six genes constitute an operon. They

encode a Cas10, a Cas7-Cas5-Cas11 fusion, a Cas7-2x, a Csx19, a Cas7 with an

insertion, and the SyCsm6 protein (**Figure 1A**). Technically, two TUs, TU7058 and

TU7063, were defined for the CRISPR2 *cas10* promoter because it contains two TSSs

(at positions 62704 and 62807 on the reverse strand)[36]. Our previous genome-wide

mapping of TSSs using differential RNA-Seq indicated the regulated expression of this

operon. High numbers of reads were found for TU7058 under most tested growth

conditions, but relatively lower numbers were recorded after transferring the cultures

to high light (470 µmol photons m^{-2} s^{-1} for 30 min), and no reads at all if cultures were

incubated in the dark for 12 h[36].

The respective repeat spacer array is transcribed on the forward strand, starting from

a single TSS approximately 6 kb away from the *cas* gene operon (**Figure 1A**). To

explore the possible differential accumulation of leader and CRISPR RNAs (crRNAs),

total RNA samples obtained from cultures grown under the same ten conditions as

those previously used for differential RNA-Seq were analyzed by Northern

hybridization. We used two probes, either complementary to the CRISPR leader RNA

and the first two spacers and repeats or to spacers 1 to 4. The first probe detected a

major signal of approximately 150 nt (**Figure 1B**, left panel), which matches the length

of the leader (125 nt;[18]) plus the length to the cleavage site within the first repeat (27

nt;[19]) and two weaker signals matching the lengths of a repeat-spacer unit of ~72 nt

and the finally processed spacer 1 of 44 nt. The second probe detected the same ~150

nt precursor transcript due to the overlap in the repeat, but revealed the strongest

signals for repeat-spacer units 2 and 3, which are somewhat longer (~75 to 77 nt) then

other units. Their accumulation was highly condition-dependent. The strongest signals

were obtained with the samples from cultures exposed to cold stress, stationary

phase, N, and C starvation, whereas the signals were weaker in samples from cultures

exposed to heat shock or high light and were not detected in samples from cultures

incubated in the dark for 12 h (**Figure 1B**, right panel). These results matched the

observations made for differential transcript accumulation of the CRISPR2 *cas10*

operon based on differential RNA-Seq.

Transcriptional regulation of the CRISPR2 cas 10 *promoter*

The observed differential accumulation of *cas* gene operon-mRNAs and crRNAs may

be due to differential transcription, post-transcriptional regulation, or both. Prediction

of a putative HLR1 motif in the CRISPR2 *cas10* promoter indicated possible

transcriptional regulation. This HLR1 motif is located -70 nt to -53 nt from the TSS of

TU7063 and -172 to -155 nt from the TSS of TU7058. To examine it's possible

relevance, we cloned the CRISPR2 *cas* gene 5´UTR and the promoter region (+122

to -203 with regard to the TSS of TU7058), including the HLR1 motif (native promoter,

P_{nat}) upstream of the *luxAB* reporter gene in the vector pILA[37]. As a control, we

mutated the HLR1 motif by substituting four nucleotides with guanosines (mutated

promoter, P_{mut}). Initial P_{nat} activity was measured under low-light conditions. Promoter

activity was measured again after a 4-hour incubation under high light, where we

observed a decrease in the measured activity to the level of the no-promoter control

(P_{less}). After transfer back to low light, P_{nat} activity increased significantly over time,

reaching an approximately ten-fold luminescence level after 120 min (**Figure 2A**). In

contrast, the P_{mut} promoter carrying the mutated HLR1 motif exhibited a basal level of

bioluminescence, similar to the control strain carrying promoterless (P_{less}) *luxAB*

genes, even after transfer back to low light. This indicates the importance of mutated

nucleotides in the recognition and binding of RpaB to the promoter. When we exposed

the cells after the initial 4 h continuously to high light (**Figure 2B**) or added the electron

transfer inhibitor DCMU (**Figure 2C**), bioluminescence remained at a basal level with

P_{nat}, P_{mut}, and P_{less} for the duration of the experiment. This effect might be specifically

related to RpaB and a change in the redox status as the P_{syr9} promoter used for control

showed modest activity under high light and no negative influence of added DCMU.

These results suggested that the CRISPR2 *cas10* promoter is regulated by a redox-

dependent mechanism involving the HLR1 motif. Furthermore, these results are

consistent with the prediction that RpaB positively regulates this promoter because it

is known to dissociate from its binding motif HLR1 under high light[38].

RpaB binds to native CRISPR2* cas10 *promoter but not to mutated HLR1 site

We then validated the prediction that RpaB regulates the transcription of the CRISPR2

effector complex. Therefore, RpaB from *Synechocystis* 6803, fused to a C-terminal

6×histidine tag, was expressed in *E. coli* DE3 and purified using nickel

chromatography (**Figure S1**). For the electrophoretic mobility shift assay (EMSA),

increasing amounts of purified RpaB were incubated with 0.5 pmol of Cy3 labelled

DNA probes carrying either the wild type or the mutated HLR1 motif (P_{nat} and P_{mut}). As

a positive control, we used the *psbA2* promoter P_{psbA2}, which was previously

characterized and shown to contain a functional HLR1 motif[39].

For the P_{nat} and P_{psbA2} fragments, a band shift was observed with 50 pmol recombinant

6×His-RpaB. For the P_{mut} fragment with four substituted bases within the HLR1 motif,

the highest amount of added 6×His-RpaB (250 pmol) was not sufficient to induce a

band shift (**Figure 2D**).

Taken together, these results strongly suggest that the redox-dependant transcription

factor RpaB positively regulates the transcription of the CRISPR2 effector complex

under low light by binding to the HLR1 site. This suggests that the expression of the

CRISPR2 *cas10* complex is activated under low-light conditions by RpaB and

deactivated under high light when RpaB binding is lost. We wondered what this would

mean to the accumulation of crRNAs. Moreover, upon acclimation to high light, RpaB

regains DNA-binding activity[29]. Therefore, we performed another experiment in which

we extended the time at high light to 6 h, transferred cells to nitrogen starvation conditions, added the electron transport inhibitors DCMU or DBMIB to the cultures, and analysed the accumulation of CRISPR2 leader and crRNAs. Northern hybridization against spacers 1 - 4 showed several bands ranging from approximately 250 nt (pre-crRNA) to 72 nt (**Figure 3A**), which corresponds to a crRNA precursor[18]. At approximately 150 nt, we observed a double band corresponding to the partially processed pre-crRNAs as in **Figure 1B**. Because the spacers differ in sequence and length, the intermediate cleavage products had slightly different sizes. Contrary to the results in **Figure 1B**, we observed no decrease but a slight increase in the accumulation of pre-crRNA after six hours of exposure to high light, which indicated that the cells had acclimated to the new environmental conditions (**Figure 3A**). In nitrogen-depleted medium, we observed a decrease in the pre-crRNA accumulation after six hours, consistent with **Figure 1B** and previous transcriptome analysis results[36]. To test the impact of changes in redox conditions on pre-crRNA accumulation, we added the photosynthesis inhibitor DCMU (3-(3,4-dichlorophenyl)-1,1-dimethylurea) or the cytochrome b_6f complex inhibitor DBMIB (2,5-dibromo-3methyl-6-isopropyl benzoquinone) to our cultures. Here, we observed a weaker accumulation of the lower double bands at 150 nt. Furthermore, in the presence of DBMIB, mature crRNAs (< 80 nt) vanished almost completely, which is consistent with the overall decrease in spacer transcript accumulation in the presence of the inhibitors. To test if spacer and leader accumulation differed, we hybridized the same membrane against the leader transcript (**Figure 3B**). The signal ran at approximately 150 nt, matching the previously estimated length of 125 nt for the transcribed leader plus the length of the first repeat up to the first Cas6 cleavage site of 29 nt[18]. The accumulation of the leader was similar to that of spacer repeats under high light and nitrogen

depletion conditions (**Figure 3A**). The addition of DCMU greatly reduced the accumulation of the leader compared to the standard (low light) conditions, and the addition of DBMIB resulted in the loss of the leader transcript signal. The observed effects of DCMU and DBMIB could be explained by a general inhibitory effect on RNA synthesis. To test this possibility, we hybridized a probe for *atpT* mRNA, which was previously found to be strongly induced by the addition of DCMU or DBMIB [40,41]. Both inhibitors upregulated the accumulation of *atpT* mRNA, demonstrating that transcription was not inhibited globally (**Figure 3C**).

These results suggest that the stability of the CRISPR2 leader transcript is linked to the redox status of the plastoquinone pool.

The CRISPR2 array promoter is highly active

We observed a decreased accumulation of the spacer-repeat and leader transcripts when the cells were exposed to high light of 470 μmol photons m^{-2} s^{-1} for 30 min (**Figure 1B**). To determine whether the promoter itself might be regulated, similar to the promoter of the *cas* gene operon, we cloned the region -1 nt to -100 nt from its TSS, fused it to a synthetic ribosome binding site into the pILA vector, and integrated the construct in *Synechocystis 6803* as for the CRISPR2 *cas10* promoter constructs. **Figure 3D** shows the *luxAB* reporter assay with the CRISPR2 array promoter. The cultures were exposed to high light for 30 min before being returned to low light to avoid acclimation to the increased light intensity. The measured bioluminescence was extremely high, reaching up to 18,000 units, the highest measured activity in a comparison of five different promoters (**Figure S2**).

Moreover, we did not observe a decrease in the bioluminescence signal after exposing the cultures to high light for 30 min, but rather a further increase in bioluminescence.

These results suggested that the CRISPR2 array promoter is not influenced by

environmental light conditions and that the observed regulation of leader and crRNA

transcripts was caused by another mechanism.

In vitro pull-downs reveal the CRISPR2 leader RNA is interacting with CrhR

Because we found no evidence for RpaB control of the crRNA promoter, we

considered preliminary results that indicated the involvement of the DEAD-box RNA

helicase CrhR as another possible factor. CrhR mediates light- and redox-dependent

responses in *Synechocystis* 6803[30]. We used CrhR produced as a recombinant

protein in *E. coli*. Two *E. coli* strains expressing recombinant His-tagged native CrhR

or CrhR$_{K57A}$ with enhanced RNA binding due to the K57A substitution within the ATP-

binding motif were constructed[35]. The *crhR*$_{WT}$ and *crhR*$_{K57A}$ reading frames were

cloned into the pQE-70 vector upstream of a segment encoding a 6×His-tag and

transformed into *E. coli* M15. Expression of His-tagged proteins was induced using 1

mM IPTG. The ~55 kDa proteins corresponding to His-tagged CrhR and CrhR$_{K57A}$

were detected three hours after induction with IPTG, purified via HiTrap Talon crude

column (Cytiva) chromatography, and eluted with a step gradient of imidazole

concentrations (**Figure S3**).

The recombinant proteins were incubated with *Synechocystis* 6803 total RNA,

subjected to co-immunoprecipitation (co-IP) and three cDNA libraries were prepared

from the bound RNA, representing RNA interacting with recombinant CrhR,

recombinant CrhR$_{K57A}$, and total RNA for background control. The experiment was

performed in biological duplicates. The total number of reads obtained from the single-

end Illumina sequencing are listed in **Table S1**. The reads were trimmed, the adapter

contaminants were filtered out with cutadapt, and then mapped to the *Synechocystis*

6803 chromosome and plasmids using bowtie2[42]. Using the PEAKachu peak caller[43]

39 peaks were identified in the CrhR library (**Figure 4, Table 1**), and 41 peaks were

called with the RNA obtained from CrhR$_{K57A}$ (**Figure 4, Table 2**) which met a log$_2$FC

≥1 and adjusted p-value ≥0.05. The peaks mapped to positions on the chromosome

and the plasmids pSYSA, pSYSM and pSYSX. Of these, 24 peaks obtained from the

pull-downs with CrhR and 25 peaks from pull-downs with CrhR$_{K57A}$ were shared

between the two libraries, among them the CRISPR2 leader RNA (**Figure 4**). Both

RNA helicase CrhR and the CrhR$_{K57A}$ mutant showed strong interactions with their

own mRNA, consistent with previous results on its autoregulatory features[44]. In

addition to the leader, several crRNAs of the CRISPR2 array were also enriched in

CrhR co-IP (**Table 1**).

The most highly enriched transcripts for CrhR$_{K57A}$ were asRNAs to the genes *sll0169*,

slr2000 and *slr1494*, encoding the DUF4101 and DnaJ-domain-containing protein

Sll0169, the S-layer homology domain-containing protein Slr2000 and an ABC

transporter subunit (**Table 2**).

Because the CRISPR2 leader RNA was enriched in co-IPs with both proteins, EMSA

experiments were performed to confirm the interaction between CRISPR2 leader and

CrhR. For this, the CRISPR2 leader RNA was synthesized by T7 RNA polymerase *in*

vitro and used as an RNA substrate. For transcript synthesis, a DNA fragment with

coordinates 68373 - 68498 on pSYSA was amplified using the primers

EMSA_CRISPR2LeadeR-T7_Fw (carrying a T7 promoter sequence followed by two

Gs) and EMSA_CRISPR2LeadeR-T7_Rv. The resulting 128 nt transcript was Cy3

labelled. Binding of 2 pmol of Cy3-labeled transcripts to various amounts of purified

recombinant His-tagged CrhR or CrhR$_{K57A}$ ranging from 1 to 50 pmol was performed

in the presence of poly dI-dC in high molar excess to the transcripts as a competitor

to confirm the specificity of the RNA-protein interaction. A gel shift of the CRISPR2

leader was observed upon the addition of only 1 pmol of CrhR (**Figure 5**). We

concluded that the CRISPR2 leader transcript was strongly bound by both CrhR and

CrhR$_{K57A}$.

Effect of Δ*crhR *mutation and redox stress conditions on CRISPR2 leader and

crRNA accumulation

We next studied the effect of environmental stress conditions on *Synechocystis* 6803

wild type and the Δ*crhR* mutant. Cells were cultivated under standard growth

conditions (low light and 30°C) and exposed to either 20°C or high light, followed by

recovery under low light. Total RNA was extracted and hybridized with probes against

CRISPR2 leader or spacers 1-4. When testing the wild type and the mutant under

standard and cold conditions (**Figure 6A**), we observed a lower CRISPR2 leader

accumulation in Δ*crhR* than in the wild type. We analyzed the signal intensities

normalized to 5S control RNA and observed that in the wild type strain, the leader

transcript decreased by approximately 40% at 20°C (**Figure 6B**). In Δ*crhR*, we

observed similar amounts of CRISPR2 leader transcripts under both conditions, but

generally lower amounts than in the wild type strain. Contrary to the wild type, the

CRISPR2 leader transcript did not seem to be affected by the change in temperature

in Δ*crhR*. These results indicated that CrhR, on the one hand, had a basal stabilizing

effect on the CRISPR2 leader transcript accumulation, but that it had a destabilizing

effect during temperature downshifts.

When testing the influence of high light on CRISPR2 leader and repeat spacer array

transcript accumulation, we observed a rapid decrease in the accumulation of both

transcripts after exposure to high light for 5 min (**Figure 6C and D**). The same

observations were made after 30 min under high-light conditions. For recovery, the

cultures were again exposed to low light, and cultivation was continued for 2 h. After

the recovery phase, the number of leader and repeat-spacer transcripts was similar to

that before high light exposure. These results suggest a rapid degradation of the

leader and spacer transcripts by an unknown mechanism and show a fast adaptation

to a change in the redox status in the cell.

Determination of CrhR amino acid residues interacting with the CRISPR2 leader

To confirm the interaction unambiguously and to identify the amino acid residues of

CrhR interacting with the CRISPR2 leader, CrhR was cross-linked to CRISPR2 leader

RNA *in vitro*. In total, we obtained 12 cross-linked peptide fragments for two replicates

using UV and CrhR$_{K57A}$, one for CrhR as well as 3 peptides for CrhR$_{K57A}$ using the

chemical cross-linker 1,2,3,4-diepoxybutane (**Figure 7A)**. The amino acid residues

cross-linked to the RNA were determined as described previously[45] and are shown in

Figure 7B. None of the cross-linked amino acid residues were located in the known

conserved motifs of DEAD-box RNA helicases (**Figure S4**). We used Alphafold 2[46,47]

to predict the three-dimensional structure of CrhR. Consistent with recent reports that

CrhR exists in solution predominantly as a homodimer[48], AlphaFold modeled it as a

dimer and predicted alpha helices and beta folds in the most conserved part of the

protein. No structure was predicted for the C-terminal section of the protein, consistent

with a lack of sequence conservation (**Figure 7C**). The dimeric structure is consistent

with the homodimeric structure of *Geobacillus stearothermophilus* RNA helicase

CshA, the closest homologue of CrhR (43.57% sequence identity) for which the

structure has been resolved[49]. By analyzing the model, we found that the amino acid

residues L103, F104, H225, and C371 were located on the surface of CrhR, whereas

the amino acid residue C184 was not. We could not draw a conclusion about the

possible location of the amino acid residue P443 because the modelling failed for the

65 C-terminal residues.

Discussion

The main role of CRISPR/Cas systems is the defense against encountered phages.

Therefore, their constitutive expression might be naively expected. However, there is

mounting evidence that some CRISPR/Cas systems are regulated at the

transcriptional level, such as in *E. coli*[7−10], or *Sulfolobus islandicus*[11,12]. Regulation in

response to environmental factors, such as temperature or high cell density, when the

risk of phage infection is high, was found for *Pseudomonas aeruginosa*, in which

LasI/R and RhlI/R, two autoinducer pairs from the quorum sensing pathway, promote

the expression of the type I-F CRISPR/Cas system[14,15]. Furthermore, resource

availability can also strongly influence *cas* gene expression. The cAMP receptor

protein (CRP) binds to DNA in the presence of its co-factor cAMP, the level of which

depends on the availability of glucose in the environment. In the phytopathogen

Pectobacterium atrosepticum, CRP increases the expression of type I-F *cas* genes

when glucose is scarce[13], whereas *cas* transcription is negatively regulated by the

cAMP-CRP complex in the type I-E system of *E. coli* when glucose is available[50].

Here, we show that RpaB, a DNA-binding response regulator, controls the

transcription of the subtype III-Dv *cas* operon in the cyanobacterium *Synechocystis*

6803. RpaB is a redox-responsive transcription factor that is highly conserved in

cyanobacteria, and is a key regulator of light acclimation[51]. RpaB controls a large panel

of genes relevant for photosynthesis, photoprotection, membrane transport, and

more[29]. Analysis of the distribution of the HLR1 binding motif of RpaB in *Synechocystis*

6803 showed that RpaB functions as an activator under low-light conditions when the

HLR1 motif is located at positions -66 to -45 to the TSS and as a repressor if located

elsewhere in the promoter[29]. The finding that the abundance of crRNAs for the III-Dv

system in *Synechocystis* 6803 varies greatly between different environmental

conditions can therefore partially be explained by the control of the *cas* gene promoter

through RpaB binding to HLR1 at an activating position. The availability of Cas

proteins certainly can be limiting for the formation of Cas complexes and the protection

of the crRNAs bound to them. However, we were puzzled that the repeat-spacer array

promoter, albeit very strong, not only lacked an HLR1 motif but also showed in reporter

gene assays even a slightly higher activity at high light, contrary to the *cas* gene

promoter.

This led us consider the interaction between CrhR and the 125 nt transcribed leader[18].

The CRISPR leader is usually understood as a longer region, containing the

promoter[2,52], regulatory sequence elements important for adaptation[53–55] and TSS of

the repeat-spacer array. CRISPR leaders have mostly been studied for their roles in

spacer acquisition in the genome. However, they may also play an important role in

the post-transcriptional regulation of pre-crRNA and affect crRNA maturation and

interference. The sRNA-dependent post-transcriptional regulation of a CRISPR array

was identified in *P. aeruginosa*, where binding of the sRNA PhrS to the leader of a

type I-F system repressed the Rho-dependent termination of CRISPR array

transcription[56].

We show that this transcribed leader also exists as a distinct sRNA in the cell and that

the accumulation of CRISPR2 leader and crRNAs is strongly affected by the cellular

redox status. We found that this leader RNA was highly enriched in *in vitro* co-IPs with

recombinant CrhR and CrhR$_{K57A}$. We confirmed the leader-CrhR (and -CrhR$_{K57A}$)

interaction by EMSA and identified the interacting amino acid residues by protein-RNA

cross-linking coupled to mass spectrometry analysis[45,57]. The cross-linked residues,

L103/F104, H225, C371, C184 and P443, were not previously described to be involved

in the interactions between DEAD-box RNA helicases and their substrates[58].

However, these residues are in line with calculations of UV cross-linking efficiencies

for different amino acids that, among others, included phenylalanine (F), histidine (H)

and proline (P) found here[59]. Moreover, the systematic analysis of interactions

between mutagenized RNA- and protein variants suggested that π-stacking

interactions between aromatic amino acids (such as Y, F or H) and guanosine or

uridine residues are important for cross-linking, also for flanking amino acids[60],

whereas cysteine is prone to cross-linking due to its high reactivity[57]. Four of the here

identified 6 amino acids match these criteria and L103 is flanked by aromatic amino

acids on both sides (**Figure S4**). Moreover, with the exception of C184 they were all

predicted to be on the surface of a dimeric CrhR model (**Figure 7C**). Thus, both

aspects are consistent with the possible involvement of these residues in RNA

recognition and binding and indicate potential for further analyses in the future.

RNA helicases are enzymes that can modify RNA structures. Therefore, they are

associated with all aspects of RNA metabolism, such as regulation of gene expression,

RNA maturation and decay, transcription or the packaging of RNA into

ribonucleoprotein particles[61,62], processes that are also relevant for the formation of

CRISPR-Cas complexes. The expression of *crhR* is regulated by the redox status of

the electron transport chain[30] and becomes strongly enhanced in response to

downshifts in temperature[63]. CrhR plays a role in the modulation of multiple metabolic

pathways during cold acclimation[34] and is indispensable for energy redistribution and

the regulation of photosystem stoichiometry at low temperatures[64]. Consistent with

these physiological functions, CrhR is localized to the thylakoid membrane, but also

co-sedimented with degradosome and polysome complexes[65]. Our data showed

decreased leader and crRNA accumulation upon shifts to high light or low nitrogen,

which was most pronounced upon addition of the inhibitor DBMIB, suggesting that

these constitute redox stress effects. A redox component in the expression of

CRISPR-Cas systems has not been previously shown. However, such regulation is of

great importance for cyanobacteria, which are the only prokaryotes that perform

oxygenic photosynthesis. In fact, it was found that phage adsorption to the

cyanobacterial host, replication, modulation of host cell metabolism, and survival in

the environment following lysis all exhibited light-dependent components[66].

Indeed, transcriptional control of *cas* gene transcription through RpaB and the

recruitment of the DEAD-box RNA helicase CrhR by the transcribed leader is

consistent with this mode of regulation (**Figure 8**). The involvement of CrhR adds to

recent reports on a connection between components of the degradosome and the type

III CRISPR-Cas machinery[21,67]. The parallel control of *cas* gene transcription by the

transcription factor RpaB and the effect of CrhR on CRISPR leader and crRNA

accumulation highlight the intriguing complexity of CRISPR-Cas regulation in the

context of the host cell.

Materials and Methods

Strains and growth conditions

Cultures of wild type and different mutant strains of *Synechocystis* 6803 were grown

at 30°C in liquid BG11 medium[68] supplemented with 20 mM TES (N-[Tris-

(hydroxymethyl)-methyl]-2-aminoethane sulphonic acid) under continuous illumination

with white light of 50 µmol photons m^{-2} s^{-1} without shaking unless otherwise mentioned.

The flasks were aerated with ambient air through a glass tube and a sterile filter for

constant and fast growth. To induce gene expression from Cu^{2+}-responsive P*petE*, 2

µM $CuSO_4$ was added to exponentially growing cells in BG11 medium without Cu^{2+}.

Mutant strains of *Synechocystis* 6803 were grown in the presence of appropriate

antibiotics at the following concentrations: spectinomycin (Sp) 20 µg/mL and

kanamycin (Km) 50 µg/mL for the ΔcrhR/FLAG, Δ*crhR*/FLAG-CrhR, and Δ*crhR*/FLAG-

CrhR$_{K57A}$ strains. For the cold and high light stress experiments, *Synechocystis* 6803

strains were cultivated at 30°C under continuous white light (30-50µmol m^{-2} s^{-1}) and

shaking until reaching OD_{750nm} = 0.600. For cold stress, the cultures were then split

into two groups: one group was cultivated at 30°C, and the second group was placed

in a water bath and kept at 20°C with ice for 2 h. Under high light conditions, if not

indicated otherwise, the cells were exposed to 300 µmol m^{-2} s^{-1} for 5 or 30 min. For

recovery, the cells were placed back to low light conditions for 2 h.

E. coli strains were grown in liquid LB medium (10 g l^{-1} bacto-tryptone, 5 g l^{-1} bacto

yeast extract, 10 g l^{-1} NaCl), with continuous agitation or on agar-solidified (1.5% [w/v]

Bacto agar) LB, supplemented with appropriate antibiotics, at 37 °C.

RNA isolation

Synechocystis 6803 cells were collected by vacuum filtration on hydrophilic

polyethersulfone filters (Pall Supor®-800, 0.8 µm), transferred to a tube containing 1

mL of PGTX buffer[69], snap-frozen in liquid nitrogen and stored at -80 °C until further

use. RNA was extracted as described[35], and the RNA concentration was determined

using a NanoDrop ND-1000 spectrophotometer (Peqlab).

Recombinant protein expression and purification

E. coli M15 was transformed with the vectors pQE70:crhR-6×His and pQE70:crhR$_{K57A}$-6×His for overexpression of recombinant His-tagged proteins CrhR and CrhR$_{K57A}$. Overnight cultures were diluted 1:100 in fresh LB medium supplemented with ampicillin and kanamycin, and grown to an OD$_{600}$ of 0.7. Protein expression was induced by adding isopropyl-β-D-thiogalactopyranoside (IPTG; 1 mM final concentration). Three hours after IPTG induction, the cells were harvested by centrifugation at 6,000 × g for 10 min at room temperature. Cell pellets were resuspended in lysis buffer (50 mM NaH$_2$PO$_4$ (pH 8), 1 M NaCl, 10% glycerol, 15 mM imidazole, cOmplete™ Protease Inhibitor Cocktail (Roche)) and lysed using the One Shot constant cell disruption system (Constant Systems Limited, United Kingdom) at 2.4 kbar. Cell debris was pelleted by centrifugation at 13,000 × g for 30 min at 4°C and the lysate was filtered through 0.45 µm Supor-450 filters (Pall). Recombinant proteins were immobilized on a HiTrap Talon crude 1 mL column (GE Healthcare), equilibrated with buffer A (50 mM NaH$_2$PO$_4$ (pH 8), 500 mM NaCl), and eluted with elution buffer B (50 mM NaH$_2$PO$_4$ (pH 8), 500 mM imidazole, 500 mM NaCl).

Expression of recombinant 6×His-tagged RpaB in Rosetta (DE3)+pLysS was performed as described by Kadowaki *et al.*, 2016[38], using phosphate buffered TB medium instead of 2 × YT medium. Purification of RpaB followed the same protocol[38] using Precellys 24 (Bertin technologies) to disrupt the cells and 0.5 mL bedded Ni-NTA Agarose beads (Qiagen GmbH) to bind 6×His-tagged proteins. Protein concentration was calculated using Bradford assay. The protein samples were mixed with the Coomassie Plus (Bradford) Assay Reagent (Thermo Fisher Scientific) in a 96 well plate. The absorption at 595 nm was measured using a Victor[3] 1420 multilabel

518 plate reader (Perkin Elmer). Protein concentration was calculated based on a bovine

519 serum albumin calibration curve.

In vitro His-tag affinity purification and RNA pulldown

Recombinant proteins were isolated from *E. coli* M15 strains via precipitation on

Dynabeads™ magnetic beads (125 µl), which bind histidine-tagged proteins. To prove

the coupling of His-tagged proteins to the beads, an aliquot (5%) of a sample

containing beads was washed after protein pulldown from *E. coli* M15 and used for

SDS-PAGE analysis. The beads coupled with the His-tagged protein were further

incubated in 25 mM Tris-HCl RNA elution buffer containing 2M NaCl in order to

eliminate contaminating RNA molecules from *E. coli*, washed in 1× TBS, and

incubated in the cell lysate of *Synechocystis* 6803 wild type for 20 min. RNA from wild-

type *Synechocystis* co-precipitated together with the recombinant His-tagged proteins

was eluted in the same RNA elution buffer and further utilized for the generation of

libraries for Illumina sequencing.

CrhR-CRISPR2 leader RNA cross-linking and enrichment of cross-linked peptide-RNA heteroconjugates

We used 10 min of UV irradiation at 254 nm to covalently cross-link approximately 1

nmol of the complex formed between CRISPR2 leader RNA and CrhR protein in a

volume of 100 µL in buffer containing 50 mM NaH_2PO_4, 300 mM NaCl, and 250 mM

imidazole (pH 8.0) as described[21]. Subsequently, cross-linked peptide-RNA

heteroconjugates were enriched according to our previously established workflow[45,57].

We ethanol-precipitated the samples and resuspended the pellet in buffer containing

4 M urea and 50 mM Tris-HCl (pH 7.9). The urea concentration was then decreased

to 1 M by adding 5 vol of 50 mM Tris-HCl (pH 7.9). The RNA was hydrolyzed by adding

1 μg RNase A and T1 (Ambion, Applied Biosystems) at 52°C for 2 h, followed by

digestion with benzonase at 37°C for 1 h and trypsin (Promega) digestion overnight at

the same temperature. To remove the non-cross-linked RNA fragments and to desalt

the sample, it was passed through a C18 column (Dr. Maisch GmbH), followed by

enrichment of the cross-linked peptides over an in TiO_2 column (GL Sciences)

according to existing protocols[45], but using 10 μm TiO_2 beads as described[21]. Samples

were then dried, resuspended in 5% v/v acetonitrile and 1% v/v formic acid, and

subjected to liquid chromatography and mass spectrometry analysis.

Analysis by mass spectrometry

A nano-liquid chromatography system (Dionex, Ultimate 3000, Thermo Fisher

Scientific) coupled with a Q-Exactive HF instrument (Thermo Fisher Scientific)[45] was

used for liquid chromatography and mass spectrometry analysis. Online ESI-MS was

performed in data-dependent mode using a TOP20 HCD method. All precursor and

fragment ions were scanned in the Orbitrap, and the resulting spectra were measured

with high accuracy (< 5 ppm) at both the MS and MS/MS levels. A dedicated database

search tool was used for data analysis[57].

Promoter activity assay

The promoter region and 5´UTR of the CRISPR2 *cas* gene operon was PCR-amplified

with primer pairs prom_*cas10*_luxAB_fw and prom_*cas10*_luxAB_rev to amplify the

wild-type promoter and primer pairs prom_*cas10*_mut_luxAB_fw and

prom_*cas10*_luxAB_rev to substitute the ACAA motif in the conserved HLR1 site with

a GGGG motif. For the CRISPR2 array promoter, we cloned the 100 base pair region

upstream of the transcription start site (68274-68373) with primers

Prom_CRISPR2_fw and Prom_CRISPR2_RBS_rev to fuse the promoter with an

artificial ribosome binding site[70]. The pILA backbone, containing a promoterless *luxAB*

gene[37], was amplified in three parts with the primer pairs pILA_1_fw (or

pILA_1_RBS_fw for CRISPR2 array promoter) /pILA_1_rev, pILA_2_fw/ pILA_2_rev,

and pILA_3_fw/ pILA_3_rev. Primers were designed with overlaps to adjacent

fragments. PCR fragments were assembled using AQUA cloning[71] and transformed

into *E. coli* DH5alpha. The constructs were named pILA-$P_{CRISPR2\_cas10nat}$ and pILA-

$P_{CRISPR2\_cas10mut}$, respectively. Sequenced constructs were transformed into

Synechocystis 6803 carrying the *luxCDE* operon, encoding the enzymes for the

synthesis of decanal[72]. Segregation of the constructs was achieved by transferring

single clones to new BG11-0.75% Kobe Agar plates containing increasing

concentrations of kanamycin (10-50 µg/mL). Full segregation was verified by PCR

using primers pIGA-fw and pIGA_rev, and sequencing. The clones with segregated

pILA construct were grown in BG11 supplemented with 50 µg/µL kanamycin, 10 µg/µL

chloramphenicol and 10 mM glucose under continuous light (30-50 µmol photons m^{-2}

s^{-1}) and shaking until reaching mid-log phase ($OD_{750 nm}$ = 0.7-0.8). Cultures were

diluted to $OD_{750 nm}$ = 0.4 prior to exposure to high light conditions (300 µmol photons

$m^{-2} s^{-1}$) for four hours. Afterwards, the cells were placed back in low light (40-50 µmol

$m^{-2} s^{-1}$). As shown in **Figure 2C**, DCMU was added to the cells during the high-light

phase (50 µM final concentration). As shown in **Figure 3B**, the cells were kept under

high light after the initial 4 h of exposure and were not switched back to low light again.

Bioluminescence was measured *in vivo* by using a VICTOR[3] multiplate reader

(PerkinElmer) at total light counts per second. Cell suspensions (100 µL) were

measured in a white 96-well plate (CulturePlate^TM-96, PerkinElmer). Bioluminescence

was measured before and after exposure to high light and every 30-60 min during

recovery under low light. Next, we exposed the cells to both high and low light. In

preliminary tests, we noticed that cellular production of decanal was not sufficient to

monitor bioluminescence *in vivo*. Therefore, we added 2 µL of decanal prior to the

measurements. A strain carrying the promotorless *luxAB* gene served as a negative

control. A strain carrying the P_{Syr9}::*luxAB* construct was used as a control strain.

Technical triplicates were measured. Statistical relevance was calculated using a 2

tailed t-test in Excel (Microsoft).

Electrophoretic mobility shift assay (EMSA)

For binding of RpaB with wild-type and mutated HLR1 motifs, regions of interest were

PCR-amplified using primers EMSA_P*cas10*_HLR1_fw and

EMSA_P*cas10*_HLR1_rev, labeled with Cyanine 3 at its 5´end, from pILA-

$P_{CRISPR2\_cas10}$ constructs. The HLR1 motif from the *psbA2* promoter was used as a

positive control and amplified using the primer pair EMSA_P*psbA2*_fw and

EMSA_P*psbA2*_rev. Different amounts of eluted 6×His-tagged RpaB (0-250 pmol)

were mixed with 0.5 pmol of Cy3 labelled DNA target in binding buffer (20 mM HEPES-

NaOH pH 7.6; 40 mM KCl; 0.05 mg/mL BSA; 5% glycerol; 0.1 mM MnCl2; 1 mM DTT;

0.05 µg/µL poly (dIdC)). The reaction mixture was incubated for 30 min at room

temperature in the dark to protect from light. Electrophoresis was performed in a 3%

agarose-0.5 × TBE gel. The gel was run for 60 min in the dark at 80 V and 4°C. The

signals were visualized with a Laser Scanner Typhoon FLA 9500 (GE Healthcare)

using a green-light laser and Cy3 filter.

Binding of CrhR or CrhR$_{K57A}$ to 0.2 pmol or 2 pmol of Cy3 labelled RNA was performed

in buffer containing 20 mM HEPES-KOH (pH 8.3), 3 mM MgCl2, 1 mM DTT, 500 µg/mL

BSA. As a substrate competitor, 1 µg of LightShift poly (dIdC) (Thermo Fisher

Scientific) was added. The reactions were incubated at room temperature for 15 min

prior to loading on 2% agarose-TAE gels.

Library preparation for RNA-seq

Total RNA was Turbo DNase (Thermo Fisher Scientific) treated, purified, and size-

separated using the RNA Clean & Concentrator-5 Kit (Zymo Research) and treated

with 5'-polyphosphatase (Epicentre) as described[35]. RNA was phosphorylated at the

5' end by T4 polynucleotide kinase (NEB) and ligated to the 5' adapter (**Table S2**). A

ThermoScript Reverse Transcriptase Kit (Invitrogen) was used for cDNA synthesis

and amplified with Phusion High-Fidelity DNA polymerase (Thermo Fisher Scientific)

using PCR primers 1 and 2 (**Table S2**). PCR conditions were 98°C for 30 s, followed

by 18 cycles of denaturation at 98°C for 10 s, primer annealing at 60°C for 30 s, and

extension for 15 s at 72°C, and a final extension step at 72°C for 2 min. The ExoSAP-

IT PCR Product Cleanup Reagent (Thermo Fisher Scientific) was used for primer

removal, and the samples were further purified with the NucleoSpin® Gel and PCR

Clean-up Kit and eluted with 20 µL of NE buffer. A 10 µL aliquot of each prepared DNA

library was sequenced on an Illumina sequencer.

RNA-seq data analysis

RNA-seq data analysis was performed using tools installed in usegalaxy.eu. The

paired-end or single-end reads were trimmed, adapters and reads shorter than 14 nt

were filtered out by cutadapt 1.16[73]. Mapping was performed on the chromosome and

plasmids of *Synechocystis* 6803 by bowtie2 2.3.4.3 with the parameters for paired-

end reads: -I 0 -X 500 --fr --no-mixed --no-discordant --very-sensitive[42]. Unmapped

reads were filtered. Peak calling of the mapped reads was performed using PEAKachu

0.1.0.2, with the parameters: --pairwise_replicates --norm_method deseq --

mad_multiplier 2.0 –fc_cutoff 1 --padj_threshold 0.05.

Data availability

The RNA-seq data have been deposited in the SRA database

https://www.ncbi.nlm.nih.gov/sra/ and are openly available under the accession

numbers SRX6451369 to SRX6451374. All mass spectrometry proteomics datasets

analyzed during this study are available at the Proteomics Identifications Database

(PRIDE, at https://www.ebi.ac.uk/pride/) under the project accession number

PXD047440.

Funding

This work was supported by the German Research Foundation priority program

SPP2141 "Much more than Defence: The Multiple Functions and Facets of CRISPR-

Cas" (grants HE 2544/14-2 to WRH and UR225/7-2 to HU).

Acknowledgments

We thank Richard Reinhardt and his team, Cologne, Germany, for sequence analyses.

The support by Sergey Moshkovskiy and Olexandr Dybkov in interpreting mass

spectrometry data is greatly appreciated. We thank Monika Raabe, Göttingen,

Ingeborg Scholz and Viktoria Reimann, Freiburg, Germany, for the expert technical

assistance, and George Owttrim, Edmonton, Canada, for discussions about CrhR.

Conflict of interest

The authors declare the absence of conflicts of interest.

AUTHOR CONTRIBUTIONS

W.R.H. designed the work. Protein-RNA cross-linking experiments and identification

of cross-linked peptide-RNA bonds were performed by A.W. and H.U., the analyses

of RpaB effects, CRISPR2 leader and repeat-spacer accumulation were performed by

R.B. All other CrhR-related experiments were carried out by A.M., the setup of cDNA

libraries and the analysis of pull-down results were performed by A.M. and C.S. A.M.,

R.B. and W.R.H. wrote the paper with contributions from all authors.

Tables

Table 1. RNA enriched from *Synechocystis* 6803 by *in vitro* pulldown using

recombinant CrhR as bait. The experiment was performed in biological duplicates.

Acronyms: asRNA, antisense RNA; Chr, chromosome.

Peak start	Peak end	S	log$_2$FC	p$_{adj}$	Annotation	Description
Plasmid pSYSA						
69586	69663	+	4.57	1.25E-07	CRISPR2 array	S15R16
68651	68827	+	2.66	9.90E-08	CRISPR2 array	R3S3R4S4R5
68370	68531	+	2.42	3.47E-09	CRISPR2 leader	Transcribed leader of subtype III-Dv CRISPR Cas system and R1
69040	69125	+	1.89	1.56E-04	CRISPR2 array	R7S7R8
68828	68996	+	1.44	3.08E-03	CRISPR2 array	S5R6S6R7
69207	69355	+	1.33	2.71E-02	CRISPR2 array	S10R11S11R12
Plasmid pSYSM						
28098	28318	+	2.38	1.42E-03	sll5026-as	asRNA to DNA phosphorothioation-associated putative methyltransferase
Plasmid pSYSX						
82429	82556	+	2.12	6.29E-05	ssr6089-gene	Small 72 aa hypothetical protein
27704	27866	+	1.36	1.58E-02	ssr6030-gene	ORF and 3' UTR of small 72 aa hypothetical protein
Chr						
2888522	2888828	+	5.99	9.24E-54	slr0083-gene	DEAD-box RNA helicase CrhR
2887947	2888283	+	5.90	2.20E-60	slr0083-gene	DEAD-box RNA helicase CrhR
2888829	2889112	+	5.22	3.52E-45	slr0083-gene	DEAD-box RNA helicase CrhR
1446281	1446359	-	5.11	1.26E-10	slr2000-as	asRNA to the gene of hypothetical protein
2887606	2887893	+	4.89	2.76E-42	slr0083-gene	DEAD-box RNA helicase CrhR
2347288	2347369	+	3.06	5.96E-17	ncr1160-ncRNA	Ncr1160 (SyR11)
1242318	1242389	-	3.02	4.28E-07	sll1757-5'UTR	asRNA to gene of hypothetical protein
467144	467292	+	2.75	8.07E-15	smr0005-5'UTR, gene	5'UTR and ORF of photosystem I subunit XII
2082308	2082609	+	2.34	6.55E-07	slr0442-gene	Hypothetical protein
1167431	1167719	+	2.09	2.10E-04	smr0009-gene	Photosystem II PsbN protein
3250506	3250676	+	2.07	2.25E-06	ncr1600-ncRNA	5'UTR of *rnj* encoding RNase J
991513	991704	-	2.06	3.55E-05	sll1463-gene	ATP-dependent zinc metalloprotease FtsH4
342867	343104	+	1.83	3.15E-03	ncr0150-ncRNA; slr0974-gene	Nr0150; ORF of initiation factor IF-3 (*infC*)
1654914	1655020	+	1.78	9.43E-04	ssr2194-gene	Hypothetical protein

1997946	1998013	-	1.73	2.26E-02	slr0935-as-asRNA	asRNA to gene encoding hypothetical protein
2839588	2839742	+	1.70	2.20E-02	ssr1407-gene	Gene encoding hypothetical protein
1141835	1141991	-	1.62	1.69E-03	ssl1633-5'UTR, gene	5'UTR and ORF of high light-inducible polypeptide HliC
1558635	1558806	-	1.60	1.54E-02	ssl3769-gene; ncl0710-ncRNA	Hypothetical protein; Ncl0710
2290635	2290796	+	1.56	6.75E-03	ssr0536-5'UTR, gene	5'UTR and ORF of hypothetical protein
3516172	3516339	-	1.56	4.78E-03	sll0430-gene	HtpG, heat shock protein 90, molecular chaperone
3204827	3204897	-	1.53	6.96E-03	sll0505-gene	Hypothetical protein
822850	822979	+	1.42	8.23E-03	slr1890-gene	ORF of bacterioferritin
2082626	2082803	+	1.39	1.59E-02	slr0442-gene	Hypothetical protein
631992	632169	-	1.33	9.24E-04	ssl1911-5'UTR, gene	5'UTR and ORF of glutamine synthetase inactivating factor IF7
1905897	1906083	-	1.31	3.45E-02	sll1434-5'UTR, gene	5'UTR and ORF of hypothetical protein
2028142	2028326	+	1.29	4.39E-02	sll0905-as-asRNA	asRNA to gene *maf*
1052427	1052589	+	1.24	3.21E-03	ssr2062-5'UTR, gene	5'UTR and ORF of hypothetical protein
2270222	2270394	-	1.14	2.26E-02	ncl1130-ncRNA	Ncl1130
117577	117740	-	1.12	2.65E-02	sll0704-5'UTR, gene	5'UTR and ORF of cysteine desulfurase

687 **Table 2.** RNA enriched from *Synechocystis* 6803 in an *in vitro* pulldown assay using

688 recombinant His-CrhR$_{K57A}$ as bait. The experiment was performed in biological

689 duplicates.

Replicon	Peak start	Peak end	Strand	log2 FC	padj	Annotation	Description
Plasmid pSYSA							
pSYSA	68370	68525	+	1.66	5.47E-09	CRISPR2 leader	Transcribed leader of the subtype III-Dv CRISPR Cas system
pSYSA	100746	101032	+	1.07	0.045406	slr7104-gene	5'UTR and ORF of transposase
Plasmid pSYSM							
pSYSM	28097	28309	+	2.03	0.005485	sll5026-as	asRNA to DNA phosphorothioation-associated putative methyltransferase
pSYSM	81234	81442	+	1.33	0.006107	slr5082-gene	ORF of Rpn family recombination-promoting nuclease/putative transposase
pSYSM	108252	108558	+	1.21	0.032833	slr5118-gene	ORF of Rpn family recombination-promoting nuclease/putative transposase
pSYSM	112774	113051	+	1.16	0.033782	slr5124-gene	ORF of Rpn family recombination-promoting nuclease/putative transposase
Plasmid pSYSX							
pSYSX	82424	82546	+	1.47	0.000739	ssr6089	Small 72 aa hypothetical protein
Chr							
NC_000911	2888558	2888865	+	8.76	0	slr0083-gene	DEAD-box RNA helicase CrhR
NC_000911	2888152	2888426	+	8.22	0	slr0083-gene	DEAD-box RNA helicase CrhR
NC_000911	1796717	1796808	+	8.19	3.42E-06	sll1879-as	asRNA to the gene of two-component response regulator ycf55
NC_000911	2887878	2888146	+	7.96	0	slr0083-gene	DEAD-box RNA helicase CrhR
NC_000911	2888867	2889139	+	7.87	0	slr0083-gene, 3'UTR	ORF and 3'UTR of CrhR
NC_000911	2887594	2887822	+	6.78	0	slr0083-gene	DEAD-box RNA helicase CrhR
NC_000911	2315608	2315718	+	4.63	5.47E-09	sll0169-as	asRNA to the gene of cell division protein Ftn2 homolog
NC_000911	1446283	1446373	-	4.56	1.49E-12	slr2000-as	asRNA to gene of hypothetical protein
NC_000911	45835	46078	-	3.98	5.47E-09	slr1494-as	asRNA to the gene of putative multidrug resistance family ABC transporter
NC_000911	1242313	1242398	-	2.58	2.87E-08	sll1757-5'UTR	ORF of hypothetical protein
NC_000911	342865	343096	+	2.02	2.34E-05	ncr0150-ncRNA; slr0974-gene	Ncr0150; ORF of initiation factor IF-3 (InfC)
NC_000911	2732766	2732918	-	1.74	0.009472	sll0184-5'UTR, gene	5'UTR and ORF of group 2 RNA polymerase sigma factor SigC
NC_000911	1167395	1167724	+	1.71	0.000146	smr0009-gene	ORF of photosystem II PsbN protein
NC_000911	467114	467324	+	1.62	6.96E-22	smr0005-5'UTR, gene	5'UTR and ORF of photosystem I subunit XII
NC_000911	2082349	2082581	+	1.62	0.000109	slr0442-gene	ORF of hypothetical protein

NC_000911	1807144	1807311	-	1.56	0.010578	sll1873-5'UTR, gene	5'UTR and ORF of hypothetical protein
NC_000911	117580	117742	-	1.42	1.54E-08	sll0704-5'UTR, gene	5'UTR and ORF of cysteine desulfurase
NC_000911	2347290	2347369	+	1.39	2.70E-06	ncr1160-ncRNA	Ncr1160
NC_000911	822850	823003	+	1.37	0.000606	slr1890-gene	ORF of bacterioferritin
NC_000911	2028141	2028320	+	1.36	0.005349	sll0905-as	asRNA to the gene maf
NC_000911	336220	336557	-	1.32	0.002837	sll0933-gene; ssl1784-gene	ORF of hypothetical protein; ORF of 30S ribosomal protein S15
NC_000911	3204824	3204940	-	1.16	0.004903	sll0505-gene	ORF of hypothetical protein
NC_000911	1919637	1919895	-	1.13	0.004242	sll1096-5'UTR, gene	5'UTR and ORF of 30S ribosomal protein S12
NC_000911	3250459	3250679	+	1.11	0.001265	ncr1600-ncRNA	Ncr1600 (5'UTR of rnj)
NC_000911	2290587	2290787	+	1.09	0.041919	ssr0536-5'UTR, gene	5' UTR and ORF of hypothetical protein (84 aa)
NC_000911	1654915	1655063	+	1.06	0.039957	ssr2194-gene	Small 65 aa hypothetical protein
NC_000911	1052426	1052590	+	1.05	5.10E-08	ssr2062-5'UTR, gene	5'UTR and ORF of hypothetical protein
NC_000911	1211732	1211980	-	0.99	0.009472	sll1774-gene	ORF of 260 aa hypothetical protein
NC_000911	2224954	2225263	-	0.91	0.031283	sll1863-gene	ORF of 107 aa hypothetical protein
NC_000911	69119	69384	+	0.90	0.036436	slr1119-gene	ORF of 233 aa hypothetical protein
NC_000911	2644699	2644967	+	0.89	0.000729	slr0628-5'UTR, gene	5'UTR and ORF of *rpsN*, 30S ribosomal protein S14
NC_000911	632000	632168	-	0.83	9.03E-05	ssl1911-5'UTR, gene	5'UTR and ORF of glutamine synthetase inactivating factor IF7
NC_000911	862001	862259	-	0.81	0.011675	ssl2233-5'UTR, gene	5'UTR and ORF of *rpsT*, 30S ribosomal protein S20
NC_000911	458684	458969	-	0.57	0.014432	sll1515-gene	Orf of 149 aa hypothetical protein

693   **Figures**

694

695   **Figure 1. Organization of the subtype III-Dv (CRISPR2) locus in *Synechocystis***

696   **6803 and the influence of environmental conditions**. **A.** The subtype III-Dv

697   CRISPR-Cas system is located on the pSYSA plasmid. Several *cas* genes are located

698   upstream of the CRISPR array, which consists of a 125 nt long leader and 56 spacers

699   of 34-46 nt in length, interspaced by 37 nt long repeats (grey squares). Arrows in

700   yellow illustrate *cas* genes encoding proteins for the adaptation module, in blue *cas6*

701   genes, and in purple genes encoding the effector complex. Accessory genes are

702   indicated by green arrows and genes encoding hypothetical proteins are in light grey.

The transcriptional start sites (TSS) of the CRISPR array and the effector complex
operon are marked by bent red arrows and the localization of antisense RNA probes
used for Northern hybridization is indicated by straight narrow arrows in black. **B.**
Influence of different environmental conditions on the accumulation of leader and
crRNAs including spacer 1 and 2 (left panel) and spacers 1 to 4 (right panel). For
Northern hybridization, ^{32}P-labeled transcript probes, as indicated in panel (A), were
used after separation of 12 µg RNA each isolated from cultures grown under 10
different conditions on a 10% urea-polyacrylamide gel. Exp. (exponential phase), stat.
(stationary phase), 42°C (heat stress), 18°C (cold stress), dark (darkness), HL, high
light (470 µmol photons $m^{-2} s^{-1}$ for 30 min), -CO_2 (limitation in inorganic carbon supply),
-N (nitrogen limitation), -P (phosphorus limitation) and -Fe (iron limitation). The
membrane and 5S rRNA hybridization to control equal loading are from Figure 2A in
publication[74].
716

Figure 2. RpaB control of the CRISPR2 *cas10* gene promoter. A. Activity of the

CRISPR2 *cas10* promoter under continuous low light after exposure to high light. **B.**

Activity of the CRISPR2 *cas10* promoter after shift to high light. **C.** Activity of the

CRISPR2 *cas10* promoter under continuous low light in the presence of DCMU after

exposure to high light. *Synechocystis* strains were transformed with pILA constructs

with a promoterless *luxAB* (P_{less}), the sRNA Syr9 promoter (P_{syr9}), the wild type (P_{wt})

and modified *cas10* promoter mutated in its HLR1 site (P$_{mut}$). The bioluminescence of

100 µL culture aliquots was measured using a Victor^3 multiplate reader. Data are

presented as the mean ± SD from three independent experiments. **D.** Electrophoretic

mobility shift assays (EMSA) to test the binding of purified His-RpaB (**Figure S1**) to

CRISPR2 *cas10* promoter fragments containing the native (P$_{nat}$) or mutated HLR1

sequence (P$_{mut}$). The HLR1-containing *psbA2* promoter fragment (P$_{psbA2}$) was used as

positive control. Substituted bases in the P$_{mut}$ fragment are highlighted in red. 0.5 pmol

Cy3-labeled DNA fragments of 80 nt length were incubated for 15 min in the dark with

His-RpaB at the indicated concentrations. Samples were separated on 0.5 x TBE and

3% agarose gels. The arrows represent imperfect repeats at the HLR1 site. HL: high

light; LL: low light.

Figure 3. Effect of addition of electron transport inhibitors on CRISPR leader accumulation. *Synechocystis* 6803 wild type was cultivated under low light (LL), high light (HL), nitrogen-depleted BG11 medium (-N), and in the presence of the electron transport inhibitors DCMU or DBMIB (50 µM each). Cells were harvested after 6 h of incubation. Total RNA was extracted, and 10 µg per lane was loaded onto an 8 M urea-10% PAA gel. Single-stranded RNA probes were hybridized against spacers 1-4 of CRISPR2 (A), CRISPR2 leader (B), and the *atpT* mRNA (C). The relative amounts of CRISPR2 leader transcripts were normalized to 5S rRNA intensity and quantified. **D.** Activity of the CRISPR2 array promoter under continuous low-light conditions or after exposure to high light.

Figure 4 RNA target enrichment in the *in vitro* RNA pulldown from *Synechocystis* 6803. A. Recombinant His-tagged CrhR was used. **B.** CrhR$_{K57A}$ RNA helicase mutant was used. The peaks obtained with PEAKachu are shown in the MA plot. Based on two biological replicates, 39 CrhR and 41 CrhR$_{K57A}$ peaks were identified with a statistically significant enrichment (padj < 0.05, log$_2$FC > 0. Common peaks for CrhR and CrhR$_{K57A}$ are shown in yellow, and the unique peaks are shown in green.

Figure 5. Interaction of CrhR wild type and mutant K57A with the CRISPR2 leader. A. Enrichment of CRISPR2 leader in the *in vitro* RNA pulldown from *Synechocystis* 6803 using recombinant His-tagged CrhR and CrhR$_{K57A}$. Colored plots show the read coverage of the enriched RNAs. The experiment was performed in two biological replicates. CRISPR2 repeats and identified peaks are represented by magenta and blue boxes, respectively. **B.** EMSA showing binding of CrhR (upper panel) and CrhR$_{K57A}$ (lower panel) to the CRISPR2 leader RNA. EMSA was performed with 2 pmol (81 ng) Cy3 labelled CRISPR2 leader RNA and various amounts of purified His-tagged CrhR or CrhR$_{K57A}$, in the presence of 1 μg of competitor poly dI-dC. Representative results from two independent experiments are shown.

Figure 6. CRISPR2 leader and repeat-spacer accumulation in
***Synechocystis* 6803 wild type and Δ*crhR* under cold and light stress conditions.**
A. Impact of *crhR* deletion on CRISPR2 leader transcript accumulation. Wild type and

Δ*crhR* were cultivated at 30°C and shifted to 20°C for 2 h. Total RNA was extracted

and 10 µg were loaded per lane on an 8 M urea-10% PAA gel. The strains were tested

for accumulation of the CRISPR2 leader transcript by hybridization using a transcript

oligonucleotide. Hybridization of 5S rRNA is shown as the control of equal loading. A

representative of biological duplicates is shown. **B.** Relative amounts of CRISPR2

leader transcripts were normalized to 5S rRNA intensity and quantified.

Ten µg of *Synechocystis* wild type and Δ*crhR*, cultivated under low light (LL) or high

light (HL), with or without recovery (R), were separated on a 10% denaturing

polyacrylamide gel. ^{32}P-labeled transcript probe specific to **C**. spacer1 – spacer4 or **D**.

CRISPR2 leader were hybridized. Hybridization of 5S rRNA is shown for control of

equal loading.

A

sample	XL_position	XL_method	RNA_adduct
mt_DEB_Rep1	QQIEVCTIPNR	DEB	U+C4H6O2
mt_DEB_Rep2	QQIEVCTIPNR	DEB	G+C4H6O2
mt_DEB_Rep2	QQIEVCTIPNR	DEB	U+C4H6O2
mt_UV_Rep1	QTACFSATM(Oxidation)PR	UV	U
mt_UV_Rep1	IEQQLYHVPR	UV	CU
mt_UV_Rep1	IEQQLYHVPR	UV	U
mt_UV_Rep1	QQIEVCTIPNR	UV	U
mt_UV_Rep1	SDWEVPEVDFNKPVLR	UV	U
mt_UV_Rep1	LFILNVYGGQSIER	UV	U
mt_UV_Rep2	IEQQLYHVPR	UV	U
mt_UV_Rep2	QQIEVCTIPNR	UV	U
mt_UV_Rep2	IEQQLYHVPR	UV	UU
mt_UV_Rep2	SDWEVPEVDFNKPVLR	UV	U
mt_UV_Rep2	LFILNVYGGQSIER	UV	U
mt_UV_Rep2	LFILNVYGGQSIER	UV	UU
wt_UV_Rep2	QQIEVCTIPNR	UV	U-H2O1-H1O3P1

B

```
MTNTLTSTFADLGLSEKRCQLLADIGFEAPTQIQTE
AIPLLLSGRDMLAQSQTGTGKTAAFALPLMDRIDPE
GDLQALILTPTRELAQQVAEAMKDFSHERRLFILNV
YGGQSIERQIRSLERGVQIVVGTPGRVIDLIDRKKL
KLETIQWVVLDEADEMLSMGFIDDVKTILRKTPPTR
QTACFSATMPREIKELVNQFLNDPALVTVKQTQSTP
TRIEQQLYHVPRGWSKAKALQPILEMEDPESAIIFV
RTKQTAADLTSRLQEAGHSVDEYHGNLSQSQRERLV
HRFRDGKIKLVVATDIAARGLDVNNLSHVVNFDLPD
NAETYIHRIGRTGRAGKTGKAIALVEPIDRRLLRSI
ENRLKQQIEVCTIPNRSQVEAKRIEKLQEQLKEALT
GERMASFLPLVRELSDEYDAQAIAAAALQMIYDQSC
PHWMKSDWEVPEVDFNKPVLRRGRNAGGGQNKSGGG
YQGKPGKPRRSSGGRRPAYSDRQQ
```

Figure 7. Cross-linking of CrhR to CRISPR2 leader RNA. A. Overview on the peptide fragments and RNA adducts detected in the two replicate samples with CrhR_K57A (mt) or CrhR (wt). **B.** Sequence of CrhR. The amino acid residues of CrhR_K57A cross-linked by UV treatment at 254 nm to the CRISPR2 leader are highlighted in red. The respective detected peptide fragments are underlined and in boldface letters. The QQIEVcTIPNR peptide was also detected for CrhR and in

addition by chemical cross-linking using 1,2,3,4-diepoxybutane[75] instead of UV

treatment. **C.** Structure of a CrhR homodimer (amino acids 9 - 427), predicted by

Alphafold2[46,47]. CrhR amino acid residues cross-linked to the CRISPR2 leader

transcript and identified by LC-MS are highlighted in red. The cross-linked CrhR

residues in the context of conserved sequence segments and previously identified

functionally relevant domains is given in **Figure S4**.

Figure 8. Multi-level redox control of CRISPR2 expression. The transcription factor RpaB binds to its HLR1 motif (red rectangle) under low light, initiating the expression of the *cas* gene operon (purple arrow), encoding the effector complex of the subtype III-Dv system in *Synechocystis* 6803. Under high light conditions, the change in the redox status of the photosystems, located in the thylakoid membrane (green arcs), leads RpaB to dissociate from its HLR1 motif, resulting in repression of the transcription of the *cas* gene operon (purple arrows). At the post-transcriptional level, high light conditions lead to a rapid decrease in the CRISPR2 leader and repeat-spacer transcript. The DEAD-box RNA helicase CrhR recognizes the leader transcript. The attachment of helicase to the CRISPR2 leader transcript is temperature dependent. At low temperature (20°C), CrhR binds to the leader transcript, whereas at higher temperature (30°C), it inhibits this interaction. HL: high light; LL: low light; TM: Thylakoid membrane.

References

1. Makarova, K.S., Grishin, N.V., Shabalina, S.A., Wolf, Y.I., and Koonin, E.V. (2006). A putative RNA-interference-based immune system in prokaryotes: computational analysis of the predicted enzymatic machinery, functional analogies with eukaryotic RNAi, and hypothetical mechanisms of action. Biol. Direct *1*, 7.

2. Brouns, S.J.J., Jore, M.M., Lundgren, M., Westra, E.R., Slijkhuis, R.J.H., Snijders, A.P.L., Dickman, M.J., Makarova, K.S., Koonin, E.V., and Oost, J. van der (2008). Small CRISPR RNAs guide antiviral defense in prokaryotes. Science *321*, 960–964.

3. Makarova, K.S., Wolf, Y.I., Iranzo, J., Shmakov, S.A., Alkhnbashi, O.S., Brouns, S.J.J., Charpentier, E., Cheng, D., Haft, D.H., Horvath, P., et al. (2020). Evolutionary classification of CRISPR-Cas systems: a burst of class 2 and derived variants. Nat. Rev. Microbiol. *18*, 67–83. 10.1038/s41579-019-0299-x.

4. Makarova, K.S., Wolf, Y.I., Alkhnbashi, O.S., Costa, F., Shah, S.A., Saunders, S.J., Barrangou, R., Brouns, S.J.J., Charpentier, E., Haft, D.H., et al. (2015). An updated evolutionary classification of CRISPR-Cas systems. Nat. Rev. Microbiol. *13*, 722–736.

5. Özcan, A., Krajeski, R., Ioannidi, E., Lee, B., Gardner, A., Makarova, K.S., Koonin, E.V., Abudayyeh, O.O., and Gootenberg, J.S. (2021). Programmable RNA targeting with the single-protein CRISPR effector Cas7-11. Nature *597*, 720–725. 10.1038/s41586-021-03886-5.

6. van Beljouw, S.P.B., Haagsma, A.C., Rodríguez-Molina, A., van den Berg, D.F., Vink, J.N.A., and Brouns, S.J.J. (2021). The gRAMP CRISPR-Cas effector is an RNA endonuclease complexed with a caspase-like peptidase. Science *373*, 1349–1353. 10.1126/science.abk2718.

7. Pul, Ü., Wurm, R., Arslan, Z., Geissen, R., Hofmann, N., and Wagner, R. (2010). Identification and characterization of *E. coli* CRISPR-cas promoters and their silencing by H-NS. Mol. Microbiol. *75*, 1495–1512.

8. Westra, E.R., Pul, Ü., Heidrich, N., Jore, M.M., Lundgren, M., Stratmann, T., Wurm, R., Raine, A., Mescher, M., and Van Heereveld, L. (2010). H-NS-mediated repression of CRISPR-based immunity in *Escherichia coli* K12 can be relieved by the transcription activator LeuO. Mol. Microbiol. *77*, 1380–1393.

9. MacRitchie, D.M., Buelow, D.R., Price, N.L., and Raivio, T.L. (2008). Two-component signaling and gram negative envelope stress response systems. Adv. Exp. Med. Biol. *631*, 80–110. 10.1007/978-0-387-78885-2_6.

10. Perez-Rodriguez, R., Haitjema, C., Huang, Q., Nam, K.H., Bernardis, S., Ke, A., and DeLisa, M.P. (2011). Envelope stress is a trigger of CRISPR RNA-mediated DNA silencing in *Escherichia coli*. Mol. Microbiol. *79*, 584–599. 10.1111/j.1365-2958.2010.07482.x.

11. Liu, T., Liu, Z., Ye, Q., Pan, S., Wang, X., Li, Y., Peng, W., Liang, Y., She, Q., and Peng, N. (2017). Coupling transcriptional activation of CRISPR–Cas system and DNA repair genes by Csa3a in *Sulfolobus islandicus*. Nucleic Acids Res. *45*, 8978–8992. 10.1093/nar/gkx612.

12. He, F., Vestergaard, G., Peng, W., She, Q., and Peng, X. (2017). CRISPR-Cas type I-A Cascade complex couples viral infection surveillance to host

861 transcriptional regulation in the dependence of Csa3b. Nucleic Acids Res. *45*,
862 1902–1913. 10.1093/nar/gkw1265.

13. Patterson, A.G., Chang, J.T., Taylor, C., and Fineran, P.C. (2015). Regulation of
the Type I-F CRISPR-Cas system by CRP-cAMP and GalM controls spacer
acquisition and interference. Nucleic Acids Res. *43*, 6038–6048.
10.1093/nar/gkv517.

14. Høyland-Kroghsbo, N.M., Paczkowski, J., Mukherjee, S., Broniewski, J., Westra,
E., Bondy-Denomy, J., and Bassler, B.L. (2017). Quorum sensing controls the
Pseudomonas aeruginosa CRISPR-Cas adaptive immune system. PNAS *114*,
131–135.

15. Høyland-Kroghsbo, N.M., Muñoz, K.A., and Bassler, B.L. (2018). Temperature, by
controlling growth rate, regulates CRISPR-Cas activity in *Pseudomonas
aeruginosa*. mBio *9*, e02184-18. 10.1128/mBio.02184-18.

16. Murray, A.G., and Eldridge, P.M. (1994). Marine viral ecology: incorporation of
bacteriophage into the microbial planktonic food web paradigm. J. Plankton Res.
16, 627–641. 10.1093/plankt/16.6.627.

17. Carlson, M.C.G., Ribalet, F., Maidanik, I., Durham, B.P., Hulata, Y., Ferrón, S.,
Weissenbach, J., Shamir, N., Goldin, S., Baran, N., et al. (2022). Viruses affect
picocyanobacterial abundance and biogeography in the North Pacific Ocean. Nat.
Microbiol *7*, 570–580. 10.1038/s41564-022-01088-x.

18. Scholz, I., Lange, S.J., Hein, S., Hess, W.R., and Backofen, R. (2013). CRISPR-
Cas systems in the cyanobacterium *Synechocystis* sp. PCC6803 exhibit distinct
processing pathways involving at least two Cas6 and a Cmr2 protein. PLoS ONE
8, e56470.

19. Reimann, V., Alkhnbashi, O.S., Saunders, S.J., Scholz, I., Hein, S., Backofen, R.,
and Hess, W.R. (2017). Structural constraints and enzymatic promiscuity in the
Cas6-dependent generation of crRNAs. Nucleic Acids Res. *45*, 915–925.

20. Kieper, S.N., Almendros, C., Behler, J., McKenzie, R.E., Nóbrega, F.L., Haagsma,
A.C., Vink, J.N.A., Hess, W.R., and Brouns, S.J.J. (2018). Cas4 facilitates PAM-
compatible spacer selection during CRISPR adaptation. Cell Rep. *22*, 3377–338.

21. Behler, J., Sharma, K., Reimann, V., Wilde, A., Urlaub, H., and Hess, W.R. (2018).
The host-encoded RNase E endonuclease as the crRNA maturation enzyme in a
CRISPR–Cas subtype III-Bv system. Nature Microbiol. *3*, 367–377.
10.1038/s41564-017-0103-5.

22. Scholz, I., Lott, S.C., Behler, J., Gärtner, K., Hagemann, M., and Hess, W.R.
(2019). Divergent methylation of CRISPR repeats and cas genes in a subtype I-D
CRISPR-Cas-system. BMC Microbiol. *19*, 147.1-147.11.

23. McBride, T.M., Schwartz, E.A., Kumar, A., Taylor, D.W., Fineran, P.C., and
Fagerlund, R.D. (2020). Diverse CRISPR-Cas complexes require independent
translation of small and large subunits from a single gene. Mol. Cell *80*, 971–979.

24. Schwartz, E., Bravo, J., Ahsan, M., Macias, L., McCafferty, C., Dangerfield, T.,
Walker, J., Brodbelt, J., Palermo, G., Fineran, P., et al. (2023). Type III CRISPR-
Cas effectors act as protein-assisted ribozymes during RNA cleavage. Res Sq,
rs.3.rs-2837968. 10.21203/rs.3.rs-2837968/v1.

25. Hein, S., Scholz, I., Voß, B., and Hess, W.R. (2013). Adaptation and modification
of three CRISPR loci in two closely related cyanobacteria. RNA Biol. *10*, 852–864.

26. Makarova, K.S., Anantharaman, V., Grishin, N.V., Koonin, E.V., and Aravind, L.
(2014). CARF and WYL domains: ligand-binding regulators of prokaryotic defense
systems. Front. Genet. *5*, 102.

27. Garcia-Doval, C., Schwede, F., Berk, C., Rostøl, J.T., Niewoehner, O., Tejero, O.,
Hall, J., Marraffini, L.A., and Jinek, M. (2020). Activation and self-inactivation
mechanisms of the cyclic oligoadenylate-dependent CRISPR ribonuclease Csm6.
Nat. Commun. *11*, 1596. 10.1038/s41467-020-15334-5.

28. Ding, J., Schuergers, N., Baehre, H., and Wilde, A. (2022). Enzymatic properties
of CARF-domain proteins in *Synechocystis* sp. PCC 6803. Front. Microbiol *.13*,
1046388. 10.3389/fmicb.2022.1046388.

29. Riediger, M., Kadowaki, T., Nagayama, R., Georg, J., Hihara, Y., and Hess, W.R.
(2019). Biocomputational analyses and experimental validation identify the regulon
controlled by the redox-responsive transcription factor RpaB. iScience *15*, 316–
331. 10.1016/j.isci.2019.04.033.

30. Kujat, S.L., and Owttrim, G.W. (2000). Redox-regulated RNA helicase expression.
Plant Physiol. *124*, 703–714.

31. Chamot, D., Colvin, K.R., Kujat-Choy, S.L., and Owttrim, G.W. (2005). RNA
structural rearrangement via unwinding and annealing by the cyanobacterial RNA
helicase, CrhR. J. Biol. Chem. *280*, 2036–2044.

32. Prakash, J.S.S., Krishna, P.S., Sirisha, K., Kanesaki, Y., Suzuki, I., Shivaji, S., and
Murata, N. (2010). An RNA helicase, CrhR, regulates the low-temperature-
inducible expression of heat-shock genes *groES, groEL1* and *groEL2* in
Synechocystis sp. PCC 6803. Microbiology (Reading, Engl.) *156*, 442–451.

33. Georg, J., Rosana, A.R.R., Chamot, D., Migur, A., Hess, W.R., and Owttrim, G.W.
(2019). Inactivation of the RNA helicase CrhR impacts a specific subset of the
transcriptome in the cyanobacterium *Synechocystis* sp. PCC 6803. RNA Biol. *16*,
1205–1214.

34. Rowland, J.G., Simon, W.J., Prakash, J.S.S., and Slabas, A.R. (2011). Proteomics
reveals a role for the RNA helicase *crhR* in the modulation of multiple metabolic
pathways during cold acclimation of *Synechocystis* sp. PCC6803. J. Proteome
Res. *10*, 3674–3689.

35. Migur, A., Heyl, F., Fuss, J., Srikumar, A., Huettel, B., Steglich, C., Prakash, J.S.S.,
Reinhardt, R., Backofen, R., Owttrim, G.W., et al. (2021). The temperature-
regulated DEAD-box RNA helicase CrhR interactome: Autoregulation and
photosynthesis-related transcripts. J. Exp. Bot. *72*, 7564–7579.
10.1093/jxb/erab416.

36. Kopf, M., Klähn, S., Scholz, I., Matthiessen, J.K.F., Hess, W.R., and Voß, B.
(2014). Comparative analysis of the primary transcriptome of *Synechocystis* sp.
PCC 6803. DNA Res. *21*, 527–539.

37. Kunert, A., Hagemann, M., and Erdmann, N. (2000). Construction of promoter
probe vectors for *Synechocystis* sp. PCC 6803 using the light-emitting reporter
systems Gfp and LuxAB. J. Microbiol. Methods *41*, 185–194.

38. Kadowaki, T., Nagayama, R., Georg, J., Nishiyama, Y., Wilde, A., Hess, W.R., and
Hihara, Y. (2016). A feed-forward loop consisting of the response regulator RpaB
and the small RNA PsrR1 controls light acclimation of photosystem I gene
expression in the cyanobacterium *Synechocystis* sp. PCC 6803. Plant Cell Physiol.
57, 813–823.

39. Eriksson, J., Salih, G.F., Ghebramedhin, H., and Jansson, C. (2000). Deletion
mutagenesis of the 5′ *psbA2* region in *Synechocystis* 6803: Identification of a
putative cis element involved in photoregulation. Mol. Cell Biol. Res. Comm. *3*,
292–298. 10.1006/mcbr.2000.0227.

40. Song, K., Baumgartner, D., Hagemann, M., Muro-Pastor, A.M., Maaß, S., Becher,
D., and Hess, W.R. (2022). AtpΘ is an inhibitor of F0F1 ATP synthase to arrest
ATP hydrolysis during low-energy conditions in cyanobacteria. Curr. Biol. *32*, 136-
148.e5. 10.1016/j.cub.2021.10.051.

41. Song, K., Hagemann, M., Georg, J., Maaß, S., Becher, D., and Hess, W.R. (2022).
Expression of the cyanobacterial FoF1 ATP synthase regulator AtpΘ depends on
small DNA-binding proteins and differential mRNA stability. Microbiol Spectr. *10*,
e0256221. 10.1128/spectrum.02562-21.

42. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie
2. Nat Methods *9*, 357–359. 10.1038/nmeth.1923.

43. Bischler, T. (2022). PEAKachu.

44. Rosana, A.R.R., Whitford, D.S., Migur, A., Steglich, C., Kujat-Choy, S.L., Hess,
W.R., and Owttrim, G.W. (2020). RNA helicase-regulated processing of the
Synechocystis rimO-crhR operon results in differential cistron expression and
accumulation of two sRNAs. J. Biol. Chem. *295*, 6372–6386.
10.1074/jbc.RA120.013148.

45. Sharma, K., Hrle, A., Kramer, K., Sachsenberg, T., Staals, R.H.J., Randau, L.,
Marchfelder, A., van der Oost, J., Kohlbacher, O., Conti, E., et al. (2015). Analysis
of protein-RNA interactions in CRISPR proteins and effector complexes by UV-
induced cross-linking and mass spectrometry. Methods *89*, 138–148.

46. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly
accurate protein structure prediction with AlphaFold. Nature *596*, 583–589.
10.1038/s41586-021-03819-2.

47. Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger,
M. (2022). ColabFold: making protein folding accessible to all. Nat. Methods *19*,
679–682. 10.1038/s41592-022-01488-1.

48. Whitman, B.T., Murray, C.R.A., Whitford, D.S., Paul, S.S., Fahlman, R.P., Glover,
M.J.N., and Owttrim, G.W. (2022). Degron-mediated proteolysis of CrhR-like
DEAD-box RNA helicases in cyanobacteria. J. Biol. Chem. *298*, 101925.
10.1016/j.jbc.2022.101925.

49. Huen, J., Lin, C.-L., Golzarroshan, B., Yi, W.-L., Yang, W.-Z., and Yuan, H.S.
(2017). Structural Insights into a unique dimeric DEAD-box helicase CshA that
promotes RNA decay. Structure *25*, 469–481. 10.1016/j.str.2017.01.012.

50. Yang, C.-D., Chen, Y.-H., Huang, H.-Y., Huang, H.-D., and Tseng, C.-P. (2014).
CRP represses the CRISPR/Cas system in *Escherichia coli*: evidence that
endogenous CRISPR spacers impede phage P1 replication. Mol. Microbiol. *92*,
1072–1091. 10.1111/mmi.12614.

51. Wilde, A., and Hihara, Y. (2016). Transcriptional and posttranscriptional regulation
of cyanobacterial photosynthesis. Biochim. Biophys.Acta *1857*, 296–308.

52. Lillestøl, R.K., Shah, S.A., Brügger, K., Redder, P., Phan, H., Christiansen, J., and
Garrett, R.A. (2009). CRISPR families of the crenarchaeal genus *Sulfolobus*:
bidirectional transcription and dynamic properties. Mol. Microbiol. *72*, 259–272.

53. Erdmann, S., and Garrett, R.A. (2012). Selective and hyperactive uptake of foreign
DNA by adaptive immune systems of an archaeon via two distinct mechanisms.
Mol. Microbiol. *85*, 1044–1056.

54. Díez-Villaseñor, C., Guzmán, N.M., Almendros, C., García-Martínez, J., and
Mojica, F.J.M. (2013). CRISPR-spacer integration reporter plasmids reveal distinct
genuine acquisition specificities among CRISPR-Cas I-E variants of *Escherichia*
coli. RNA Biol. *10*, 792–802.

55. Yosef, I., Shitrit, D., Goren, M.G., Burstein, D., Pupko, T., and Qimron, U. (2013).
DNA motifs determining the efficiency of adaptation into the *Escherichia coli*
CRISPR array. PNAS *110*, 14396–14401.

56. Lin, P., Pu, Q., Wu, Q., Zhou, C., Wang, B., Schettler, J., Wang, Z., Qin, S., Gao,
P., Li, R., et al. (2019). High-throughput screen reveals sRNAs regulating crRNA
biogenesis by targeting CRISPR leader to repress Rho termination. Nat. Commun.
10, 3728. 10.1038/s41467-019-11695-8.

57. Kramer, K., Sachsenberg, T., Beckmann, B.M., Qamar, S., Boon, K.-L., Hentze,
M.W., Kohlbacher, O., and Urlaub, H. (2014). Photo-cross-linking and high-
resolution mass spectrometry for assignment of RNA-binding sites in RNA-binding
proteins. Nat. Methods *11*, 1064–1070. 10.1038/nmeth.3092.

58. Linder, P., and Jankowsky, E. (2011). From unwinding to clamping - the DEAD box
RNA helicase family. Nat. Rev. Mol. Cell Biol. *12*, 505–516.

59. Shchepachev, V., Bresson, S., Spanos, C., Petfalski, E., Fischer, L., Rappsilber,
J., and Tollervey, D. (2019). Defining the RNA interactome by total RNA-
associated protein purification. Mol. Syst. Biol. *15*, e8689

60. Knörlein, A., Sarnowski, C.P., de Vries, T., Stoltz, M., Götze, M., Aebersold, R.,
Allain, F.H.-T., Leitner, A., and Hall, J. (2022). Nucleotide-amino acid π-stacking
interactions initiate photo cross-linking in RNA-protein complexes. Nat. Commun.
13, 2719. 10.1038/s41467-022-30284-w.

61. Bourgeois, C.F., Mortreux, F., and Auboeuf, D. (2016). The multiple functions of
RNA helicases as drivers and regulators of gene expression. Nat. Rev. Mol. Cell
Biol. *17*, 426–438.

62. Khemici, V., and Linder, P. (2018). RNA helicases in RNA decay. Biochem. Soc.
Trans. *46*, 163–172.

63. Rosana, A.R.R., Chamot, D., and Owttrim, G.W. (2012). Autoregulation of RNA
helicase expression in response to temperature stress in *Synechocystis* sp. PCC
6803. PLoS ONE *7*, e48683.

64. Sireesha, K., Radharani, B., Krishna, P.S., Sreedhar, N., Subramanyam, R.,
Mohanty, P., and Prakash, J.S.S. (2012). RNA helicase, CrhR is indispensable for
the energy redistribution and the regulation of photosystem stoichiometry at low
temperature in *Synechocystis* sp. PCC6803. Biochim. Biophys. Acta *1817*, 1525–
1536.

65. Rosana, A.R.R., Whitford, D.S., Fahlman, R.P., and Owttrim, G.W. (2016).
Cyanobacterial RNA helicase CrhR localizes to the thylakoid membrane region
and cosediments with degradosome and polysome complexes in *Synechocystis*
sp. strain PCC 6803. J. Bacteriol. *198*, 2089–2099. 10.1128/JB.00267-16.

66. Clokie, M.R.J., and Mann, N.H. (2006). Marine cyanophages and light. Environ.
Microbiol. *8*, 2074–2082. 10.1111/j.1462-2920.2006.01171.x.

67. Chou-Zheng, L., and Hatoum-Aslan, A. (2019). A type III-A CRISPR-Cas system
employs degradosome nucleases to ensure robust immunity. Elife *8*, e45393.

68. Stanier, R.Y., Deruelles, J., Rippka, R., Herdman, M., and Waterbury, J.B. (1979).
Generic assignments, strain histories and properties of pure cultures of
cyanobacteria. Microbiology *111*, 1–61. 10.1099/00221287-111-1-1.

69. Pinto, F., Thapper, A., Sontheim, W., and Lindblad, P. (2009). Analysis of current
and alternative phenol based RNA extraction methodologies for cyanobacteria.
BMC Mol. Biol. *10*, 79. 10.1186/1471-2199-10-79.

70. Kelly, C.L., Taylor, G.M., Hitchcock, A., Torres-Méndez, A., and Heap, J.T. (2018).
A rhamnose-inducible system for precise and temporal control of gene expression
in cyanobacteria. ACS Synth. Biol. *7*, 1056–1066. 10.1021/acssynbio.7b00435.

71. Beyer, H.M., Gonschorek, P., Samodelov, S.L., Meier, M., Weber, W., and
Zurbriggen, M.D. (2015). AQUA cloning: a versatile and simple enzyme-free
cloning approach. PLOS ONE *10*, e0137652.

72. Klähn, S., Baumgartner, D., Pfreundt, U., Voigt, K., Schön, V., Steglich, C., and
Hess, W.R. (2014). Alkane biosynthesis genes in cyanobacteria and their
transcriptional organization. Front. Bioeng. Biotechnol. *2*, 24.
10.3389/fbioe.2014.00024.

73. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnet j. *17*, 10. 10.14806/ej.17.1.200.

74. Zhan, J., Steglich, C., Scholz, I., Hess, W.R., and Kirilovsky, D. (2021). Inverse
regulation of light harvesting and photoprotection is mediated by a 3'-end-derived
sRNA in cyanobacteria. Plant Cell *33*, 358–380. 10.1093/plcell/koaa030.

75. Bäumert, H.G., Sköld, S.-E., and Kurland, C.G. (1978). RNA-protein
neighbourhoods of the ribosome obtained by crosslinking. Eur. J. Biochem. *89*,
353–359. 10.1111/j.1432-1033.1978.tb12536.x.

1073