Development of Pial Collaterals by Extension of Pre-existing Artery Tips

Suraj Kumar, Niloufer Shanavas, Swarnadip Ghosh, Vinayak Sivaramakrishnan,

Manish Dwari, Soumyashree Das#

National Centre for Biological Sciences, Tata Institute of Fundamental Research,

Bengaluru, KA 560065, India

Corresponding author (soumyashree@ncbs.res.in)

Key Words

Pial Collateral Artery, Artery Endothelial Cell, Vascular Endothelial Growth Factor

Receptor-2, Intra-vital Imaging, Genetic Lineage Tracing

SUMMARY

Pial collaterals provide protection from ischemic damage and improve prognosis of

stroke patients. The origin or precise sequence of events underlying pial collateral

development is unclear and has prevented clinicians from adapting new

vascularization and regeneration therapies. We use genetic lineage tracing and intra-

vital imaging of mouse brains at cellular resolution to show that during

embryogenesis, pial collateral arteries develop from extension and anastomoses of

pre-existing artery tips, in a VegfR2 dependent manner. Our data demonstrate that

an arterial receptor, Cxcr4, earlier shown to drive artery cell migration and coronary

collateral development, is dispensable for formation and maintenance of pial

collateral arteries. Our study reveals that collateral arteries of the brain are built by a

unique mechanism, distinct from that of the heart.

INTRODUCTION

Pial (leptomeningeal) collateral arteries are an extensive network of inter-artery anastomoses found in the pial layer of the brain. In humans, the extent of pial collateral circulation is a predictor of stroke outcome.[1,2] Stroke occurs when blood supply is obstructed to a part of the brain and is one of the leading causes of morbidity worldwide (World Health Organization).[3] Pial collaterals serve as an alternate route of blood flow and alleviate the symptoms of ischemic stroke by bypassing vascular occlusions. Thus, understanding mechanisms which drive the origin and remodelling of pial collateral arteries is of immense clinical relevance.

Results from preclinical studies indicate wide-spread variation in the pial collateral artery network among individuals and organs.[4] These variations are a result of differential expression and function of proteins associated with Vascular endothelial growth factor (Vegf) signalling axis.[5–7] Different mouse strains demonstrate differential Vegfa expression level which corelate with their pial collateral density.[8] Similarly, global overexpression of Vegf ligand increases number of native pial collaterals and improve collateral perfusion post-injury.[5,6] The receptor mediating the function of Vegf ligand during the formation of pial collaterals is Vegf receptor-2 (VegfR2).[6] A global knockdown of Vegf or VegfR2 during embryogenesis and in a time-frame when pial collaterals are expected to form, results in reduction in their number and diameter.[6] Thus, Vegf is a key driver of pial collateral formation and a critical factor which regulates variation in extent of pial vascular network in mice.

The vital role of Vegf in murine pial collateral development was discovered almost a decade ago.[5,6] However, it is still unclear how the various molecular players of Vegf pathway orchestrate the formation or remodelling of pial collateral network. In

mice, it is hypothesized that an *arteriole-specific sprouting angiogenesis like* process

leads to the formation of the foremost pial collaterals.[6] However, molecular evidence

supporting the identity of these sprouting cells (if arteries or capillaries) is missing.

Additionally, the cellular origin or fate of these sprouts is unknown. Finally, the

possibility of cooperation from other vascular processes, such as arterialization,

arteriogenesis or artery-reassembly, in building pial collaterals, needs to be

thoroughly investigated.

In this study, we use genetic lineage tracing in conjunction with live imaging of

pial vasculature of transgenic mouse embryos and adults to understand mechanisms

by which Vegf regulates pial collateral development and maintenance. Our data

indicate that the cellular source of pial collateral arteries is primarily pre-existing

artery endothelial cells (ECs) which build the first network of pial collaterals at

embryonic day (e)15.5 via VegfR2-dependent artery tip extension. While arterial

VegfR2 was essential for formation of pial collaterals *de novo*, it was dispensable for

its maintenance through adulthood. Arterial Cxcr4 (C-X-C chemokine receptor type-

4), which is critical for coronary artery formation in mice[9] was dispensable during pial

collateral formation and its maintenance. Together, our data indicate that artery

endothelial cells build collateral arteries by utilizing distinct vascular mechanisms in

heart and brain.

RESULTS

Pial collateral arteries form during embryogenesis

We developed a whole brain imaging technique to reliably visualize all pial

arteries (Figure **S1A**, **B**). Due to lack of unique markers, accurately identifying pial

collateral arteries is exclusively dependent on their location. Pial collateral arteries

are artery segments which interconnect the ends of cerebral arteries and can be

seen in the watershed region (Figure **S1B**). To acquire this spatial information, it was

imperative to image the entire pial artery network. Our technique allowed us to

precisely identify pial collateral arteries. We specifically focused on collateral arteries

which connected and received blood flow from Anterior Cerebral Artery (ACA) and

Middle Cerebral Artery (MCA), as the coverage of this arterial network is most

extensive.

We first investigated the developmental stage when pial collaterals first

appeared. Mouse brains were harvested at different developmental time points and

whole brain immunostaining was performed for Connexin-40 (Cx40) protein, a known

marker for mature artery endothelial cell. Analyses of pial arterial network in both

hemispheres revealed the appearance of MCA at embryonic day e12.5 followed by

ACA at e13.5 (Figure **S1C** and **S2A**, **B**). At e14.5, both cerebral arteries were

present with no arterial connections between them (figure **S2C**). Subsequently, the

first arterial anastomoses appeared at e15.5 (Figure **S2D**) which then remodelled to

expand in number through birth/postnatal day (P)0 (Figure **S1C** and **S2D-F**). This

method confirmed that both MCA and ACA are present at e14.5 (Figure **S2G**) and

the foremost pial collaterals appear at e15.5 (Figure **S2H**).

To confirm the above results, we next used a knock-in mouse line ($Cx40^{eGFP/+}$)

where eGFP expression was driven by $Cx40$ promoter activity. Brains from different

developmental stages confirmed our findings from Cx40 immunostaining. MCA

appeared at e12.5, ACA appeared at e14.5 and collateral arteries between MCA and

ACA appear at e15.5 (Figure **S3A-F** and **S1C**). eGFP$^+$ cerebral arteries (MCA and

ACA) were formed by e14.5 (Figure **S3G**) and the first eGFP$^+$ collateral artery

between MCA and ACA appeared at e15.5 (Figure **S3H**). Thus, using two

independent methods— Cx40 immunostaining and detecting its reporter activity, we demonstrate that mouse pial collateral arteries form during development, and appear precisely at e15.5 (Figure **S1C**).

Intravital imaging reveals arterial origin of pial collaterals by artery tip extension

Despite several hypotheses, there is lack of compelling data in support of vascular mechanisms which drive pial collateral development. Pial collaterals appear at late embryonic stages. At this developmental stage, rapid vascular changes make it difficult to distinguish collateral-specific events from overall vascular growth, thus making it challenging to draw information from fixed brain samples. Additionally, late embryonic stages cannot be cultured *ex vivo* due to extremely demanding metabolic and physiological needs. Thus, we custom-developed an imaging paradigm to perform intra-vital imaging of pial layer at late embryonic stages (e16-e17) (Figure **1A**), a developmental stage where the brain is actively building pial collaterals. Briefly, embryos of the desired developmental time points were dissected out (with their intact yolk sac) from the anesthetized pregnant dams while still connected to the mother via their umbilical cord. Embryos were maintained at 37˚C and immobilized at appropriate orientation to aid imaging. The embryos were subjected to stereoscopic or confocal imaging for several hours. This imaging set up allowed visualization of the entire hemisphere— MCA, ACA, and connected collateral arteries (Figure **1B**).

Pial collaterals appear only after the primary branches of MCA and ACA have developed (Figure **S1C**). We hypothesized that the branches of these cerebral arteries contribute towards formation of pial collaterals. To test this hypothesis,

Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5 mice were used to label artery

ECs with TdTomato and all endothelial cells with eGFP. To activate TdTomato,

Tamoxifen was administered at e13.5, when both MCA and ACA are present and

would be labelled in 2 days, [9,10] by e15.5. Imaging of endothelial cells derived from

arteries was performed at e16-e17. This experimental strategy allowed visualization

of cellular dynamics specific to artery cells derived from pre-existing arteries (Figure

1C) and excluded the possibility of capturing other vascular processes.

We performed intra-vital time-lapse imaging of pial arteries and investigated if

pial collaterals originate from pre-existing artery endothelial cells. We focused on the

watershed region: an initial artery-free region which spans the distal ends of both

arterial trees (MCA and ACA) and is later occupied by pial collaterals (Figure **S1C**

and **1D-F**). Stereoscopic images of e16.5 brains revealed that branches from MCA

and ACA extend (arrowheads, Figure **1E**) and anastomose to build collaterals in the

pial watershed (brackets, Figure **1E**). Additionally, TdTomato expression in artery

tips (orange arrow, Figure **1E**) and collaterals (green arrow, Figure **1E**) increased

over time. These changes were executed within a short span of 80 minutes, and

could not have been captured in fixed brain samples.

We next investigated artery cell dynamics associated with the growth fronts of

the two cerebral arteries (Figure **1F**, **G**). At e16.5, the distal branches of the

developing MCA extended and anastomosed to build an artery network *de novo*

(arrowheads, Figure **1G**). This is the first evidence demonstrating that pre-existing

cerebral artery tips build pial collaterals. We then performed intra-vital time-lapse

confocal imaging of brains at e16, a stage prior to when we observed collateral

artery networks being built *de novo* (Figure **1E**, **G**). Confocal imaging at e16 provided

the necessary cellular resolution to quantitatively assess artery-tip extensions from

MCA and ACA branches (Figure **1H-J**). As expected, we clearly observed artery tip

extension of MCA branch (Figure **1H**, **I**) and the ACA branch (Figure **1H**, **J**) within an

hour. Additionally, we noted several artery-specific movements within the pial layer.

This included the appearance of single artery endothelial cells (arrows, Figure **S4A**,

B), movement of these single artery cells towards an artery (Figure **S4C**, **D**), and

emergence of arterial sprouts (arrowheads, Figure **S4A**, **B**). Most interestingly, all

artery cell movements were observed along microvascular tracks which appeared

structurally more organized/tubular than the surrounding capillaries (arrowheads,

Figure **1K** and Figure **S4A**, **E**). These "microvascular tracks" were distinct from the

surrounding capillary plexus in that they formed direct connections between two

arteries but were not entirely populated by TdTomato[+] artery derived endothelial

cells (Figure **1K** and Figure **S4A**, **E**). Together, for the first time, we provide direct

evidence for the vascular origin of pial collaterals— pre-existing MCA/ACA artery tips

grow on distinct microvascular paths and connect with each other to build pial

collaterals *de novo* (Figure **1L**).

Artery ECs are the major cellular contributors of pial collaterals during

embryogenesis

We next assessed the vascular-lineage composition of pial collaterals at a

later stage when the collateral artery network has fully formed. This would allow us

to: 1) determine if the artery cell contribution to nascent pial collaterals observed at

e16.5 (Figure **1E**) persisted in later stages and 2) check contribution from

microvascular lineage which is evident in newly formed collaterals in heart[9,11] and

skeletal muscle.[12] Genetic lineage tracing with *Cx40CreER; Rosa26[TdTomato]* and

ApjCreER; Rosa26[TdTomato] mice were performed where TdTomato expression could

be conditionally induced, with Tamoxifen, specifically in artery ECs or capillaries

respectively. Tamoxifen was administered to timed pregnant dams at e14.5 when

both MCA and ACA arteries are present (Figure **2A**, **S2G**, **S3G**). Embryos were

analysed at e18.5 or P0 when the pial collateral artery network has completely

formed (Figure **2A**, **S2H**, **S3H**). Whole embryonic brains were immunostained for

Cx40 and segments of collateral arteries between MCA and ACA were analysed for

lineage contribution (Figure **2B-J**).

We assessed the extent of lineage contribution from artery or capillary ECs

towards pial collaterals. As expected, all pre-existing artery ECs were labelled and

traced in *Cx40CreER; Rosa26^{TdTomato}* embryos (Figure **2C**, **D**, **G**, **I**). Whole brain

imaging of the embryo (Figure **2C**) allowed identification of pial collateral arteries

present between MCA and ACA in the watershed (Figure **2D**). Individual collateral

segments as identified by Cx40 immunostaining were analysed for presence of

TdTomato$^+$ artery ECs (Figure **2D**, **G**, **I**). This analysis revealed that 77% of the

Cx40$^+$ collateral area to be TdTomato$^+$ (Figure **2B**). In contrast to this data, a similar

analysis in an independent experiment with *ApjCreER; Rosa26^{TdTomato}* mice (Figure

2E, **F**, **H**, **J**) showed only 31.6% of the collateral area to be TdTomato$^+$, i.e. traced

from capillaries (Figure **2B**). Thus, artery ECs are the major contributors of pial

collateral arteries during development.

Arterial VegfR2 mediates artery tip extension and is essential for pial collateral artery development

We analysed similarities, if any, in the molecular programs used by artery

endothelial cells during collateral development in the heart and brain. Post-injury,

neonatal coronary artery ECs undergo extensive migration and proliferation to build

new coronary collateral arteries in the watershed. The processes of artery cell

migration and proliferation are heavily dependent on Cxcr4[9] and VegfR2[13]

respectively. To assess the functions of Cxcr4 and VegfR2 in pial collateral

development, we deleted the respective genes at e14.5 and assessed the extent of

pial collateral network at e18.5/P0 by whole brain imaging (Figure **3A**). Surprisingly,

deletion of arterial *Cxcr4* did not affect the pial collateral network at e18.5 (Figure **3B**,

control=8, artery specific knockout for *Cxcr4*=13) or in adulthood (Figure **S5**, N=5 per

genotype). This data suggests that a Cxcr4-independent vascular mechanism

facilitates pial collateral development and its maintenance.

We next assessed if pial collateral artery development is dependent on

arterial VegfR2. Whole brain imaging of control and *VegfR2* artery specific knockout

embryos showed differences in collateral connectivity in the watershed, and required

systematic quantification (Figure **3C**-**F**). The extensive connectivity of pial arteries

across the hemisphere (including watershed) provide several possible collateral

paths for the blood to flow. These artery-artery connections are innumerable. We

developed a method that allowed us to compare the pial artery coverage between

the control and *VegfR2* artery EC specific knockout brains quantitatively (Figure **3C**).

Intensity line-profiles were generated across the width of the hemisphere (Figure **3C**

i, **ii**). The identified peaks per line-profile was plotted across the width of the

hemisphere (Figure **3C iii**). The data was fit to a gaussian function and the fitted

curves were used as representative of the arterial coverage per hemisphere in each

mouse (Figure **3C iv**). The average of the fitted curves represented one genotype

(grey curve, Figure **3C iv**). Quantification of pial artery coverage showed significantly

different distributions between control and *VegfR2* knockout brains (KS test, D

value=0.569, p value=2.2066×10^{-141}) (Figure **3E**). Peak arterial coverage was

observed at the centre of the hemisphere and watershed was defined as a region ±

500µm from the centre (Figure **3E**). The loss of arterial coverage in the watershed

was evident in the knockout (Figure **3E**) and was quantified as area under the curve

(Figure **3E**, **F**). This area under the curve represented the extent of collateral artery

network in the watershed and was significantly lower in brains without arterial

VegfR2 (Figure **3F**, unpaired t-test, p value= 0.0001). Thus, depletion of *VegfR2* from

arteries reduced the extent of pial collaterals formed during embryogenesis.

We next investigated the possible mechanisms by which arterial VegfR2

facilitated pial collateral formation. *VegfR2* was deleted from artery ECs at e14.5,

and analyses of pial collateral network was performed at e16.5 (Figure **3G**), a

timepoint when collaterals are beginning to form via artery tip extensions (Figure **1D-**

G). Quantification showed fewer TdTomato$^+$ collateral arteries (i.e. TdTomato

contiguous or Td^{con}) without arterial *VegfR2* (Figure **3H**). This indicated that initiation

of collateral development at an earlier timepoint, e16.5, also depends on arterial

VegfR2. This VegfR2-dependent collateral phenotype (Figure **3H**) was independent

of artery cell proliferation which was minimal at this stage, for both control and

VegfR2-deleted brains (data not shown). Close analyses of collaterals in control

brains showed that TdTomato$^+$ artery cells occupied a major length of microvascular

tracks connecting branches of MCA and ACA (Figure **3I**). This observation was in

contrast to *VegfR2*-depleted brains where TdTomato$^+$ artery cells sparsely populated

the microvascular tracks (Figure **3J**). Thus, arterial VegfR2 regulates pial collateral

formation by promoting artery growth along microvascular tracks.

We investigated if endothelial cells of capillary lineage populate pial collaterals

via Dach1; a transcriptional factor that drives arterialization of capillaries into

coronary arteries during cardiac development in mice.[14] As observed in developing

mouse hearts,[14,15] expression of Dach1 was higher in capillaries as compared to

arteries (Figure **S6A**). *Dach1* was deleted from capillaries at e14.5 and the extent of

pial collateral network was assessed at e18.5 (Figure **S6B**). Interestingly, no

significant difference was observed in the number of Cx40$^+$ collaterals in the *Dach1*

capillary-EC specific knockout brains (Figure **S6C**). Thus, in contrast to heart, the

formation of pial collaterals during embryogenesis is independent of capillary Dach1

activity.

Pial collaterals do not remodel in adulthood

We next investigated the cellular dynamics associated with fully formed and

mature pial collateral arteries between MCA and ACA in adult mice. For this, we

utilized an extensive longitudinal in vivo imaging paradigm that allowed us to get

single cell resolution of vascular structures. 3-5 months old *Cx40CreER;*

Rosa26^{TdTomato}; *Tie2::eGFP::Claudin5* mice underwent craniotomy and an optical

window of 4mm was implanted on one of the hemispheres between the cranial

landmarks—bregma and lambda (Figure **4A**, **B**). In these transgenic mice, pre-

existing pial arteries were labelled with TdTomato upon Tamoxifen administration

(Figure **4C**), and constitutively expressing GFP marked all endothelial cells. 5

consecutive doses of Tamoxifen resulted in 95.4% and 86.8% recombination in

proximal and distal arteries respectively, but a lower 65.3% recombination in the pial

collateral regions (Figure **4D**). A lower recombination efficiency in the pial collateral

arteries resulted in a patchy reporter pattern comprising of a mixture of TdTomato$^+$

and TdTomato$^-$ cells (Figure **4D**). Like in developmental stages (Figure **2B**), artery

ECs populate adult pial collaterals to a greater extent as compared to other vascular

lineages. This ultimately allowed us to most accurately track the position of individual

cells within a given collateral artery.

An optical window allows longitudinal imaging. Imaging the collateral zone

(watershed between MCA and ACA) for a month (Figure **4E**), resulted in high quality

images with single cell resolution (Figure **4E-H**). Morphology of the pial vessels

alongside reporter activity in endothelial cells were instrumental in identifying pial

arteries, veins and capillaries (Figure **4E**, **I**). High resolution images of artery

endothelial cells (both TdTomato$^+$ GFP$^+$ and TdTomato$^-$ GFP$^+$) within a collateral

artery segment across many collaterals and optical windows were analysed (Figure

4J, **K**). Data showed that artery endothelial cells do not remodel or reposition

themselves within a given collateral artery segment (Figure **4J-M**). Cellular matrices

representative of the relative position of collateral artery cells suggested that under

physiological conditions, every artery cell within a mature/ adult pial collateral

maintains its contact with the same neighbouring cells (Figure **4L**, **M**). Thus,

surprisingly, the cellular organization within pial collaterals and the overall collateral

structure was extremely stable. For the first time our data, with cellular resolution,

demonstrates the static nature of artery endothelial cells during homeostasis.

Arterial VegfR2 is dispensable for pial collateral maintenance

We next asked if arterial VegfR2 is necessary for the maintenance of adult

pial collaterals using adult mice deficient in arterial *VegfR2*. Using the same in vivo

imaging paradigm mentioned above, pial collateral segments in artery endothelial

cell specific *VegfR2* knockout mice were imaged at cellular resolution and followed

for a month with imaging time-points separated by 2 weeks (Figure **5A**). Longitudinal

imaging through an optical window (Figure **5B**) yielded a map of vascular network

which could be followed for months (Figure **5C**, **D**). Of note, our qualitative

observation suggests that *VegfR2* depleted pial collaterals had smaller diameters.

High magnification images of watershed between MCA and ACA (Figure **5E-G**)

allowed us to analyse individual *VegfR2^{-/-}* artery endothelial cells (TdTomato$^+$)

residing on adult collateral arteries (Figure **5H-J**). Long term imaging of the same

collateral regions across many collaterals and mice showed no significant changes in

cellular positions relative to their neighbouring cells (Figure **5K**-**M** and Figure **S7**).

Thus, arterial VegfR2 is dispensable for maintenance of mature pial collateral

arteries. Together, in this study, we highlight the essential role of arterial VegfR2 in

promoting artery-tip extension and pial collateral development during embryogenesis

(Figure **5N**); a function dispensable in adulthood.

DISCUSSION

In the present study, we investigate artery cell dynamics during collateral artery

formation and maintenance in mouse brain. Firstly, using an artery-specific

molecular marker, we dissect out where and when pial collaterals appear during

embryogenesis. Secondly, we develop a live imaging method in mouse embryos to

provide conclusive evidence that pre-existing arteries grow on pre-determined

microvascular tracks which then connect to build pial collaterals *de novo*. Thirdly, by

combining mouse genetics with whole brain confocal imaging at cellular resolution,

we show that an angiogenic vascular receptor VegfR2 facilitates artery-tip extension

on microvascular tracks and aids origin of pial collaterals. Finally, we show that

Cxcr4, a receptor essential for Artery Reassembly and formation of coronary

collateral arteries *de novo*, is dispensable for development of pial collaterals. Thus,

two fundamentally distinct mechanisms contribute to the origin of coronary and pial

collaterals.

Organ-specific mechanisms build collaterals

For the first time, we provide direct proof that a vascular mechanism driven

specifically by artery endothelial cells leads to the origin of pial collaterals. Like in the

heart, this artery specific process is regulated by arterial VegfR2, but is distinct from

Artery Reassembly where pre-existing artery cells dissociate from resident

coronaries, migrate, de-differentiate/proliferate and coalesce to build coronary

collaterals.[9,13] Instead, developing pial arteries extend along microvascular tracks

and connect with other artery tips to build new collateral artery paths. Furthermore, in

heart, coronary collateral formation is regulated by arterial Cxcr4 which drives artery

cell migration during Artery Reassembly. In contrast, arterial Cxcr4 had no impact on

pial collateral formation or its maintenance. Unlike heart, pial artery endothelial cells

contribute minimally via proliferation. Evidence for negligible artery cell proliferation

prompts us to hypothesize that artery endothelial cells at the distal ends experience

a forward push from the proximal parts of the artery. This forward pressure could be

generated due to incorporation of new cells or expansion of existing cells at the

proximal artery segments.

The need for functional collaterals is most during an ischemic condition such

as an artery occlusion, and can significantly impact tissue function. The differences

in factors which regulate origin of coronary and pial collaterals, are potentially driven

by the metabolic needs of these tissues. While cardiomyocytes can survive hypoxia

for few hours,[16] neurons are more susceptible to low levels of oxygen.[17] Higher

hypoxia sensitivity of brain tissue may lead to formation of pial collaterals during

embryogenesis as a protective measure, while coronary collaterals form only upon

induction of an arterial occlusion. Furthermore, collateral arteries in skeletal

muscle,[12,18] intestine,[19] and hind limb[20] can also undergo arteriogenesis.[21] Identifying

collateral-building vascular mechanisms across organs could reveal clinically

relevant information and open avenues for therapeutics.

Arteries and capillaries collaborate to build collaterals

We demonstrate the formation of collateral arteries, in real time, from pre-

existing arterial lineage as observed[9,10,13,22] or proposed[6] earlier. While this

manuscript was in progress, Perovic et al reported that two separate vascular

lineages populate pial collaterals.[10] We also observed that well-formed pial

collaterals are majorly populated by artery cells, and minimally by capillaries. Intra-

vital imaging of embryonic brains reveal that pre-existing artery cells extend along

defined microvascular paths to connect and build pial collaterals. These artery

specific events are executed within a short span, are difficult to capture using static

images and can be observed only by live imaging.

We cannot exclude the contribution of other artery-driven mechanisms. We

observed single artery cells migrating towards pre-existing arteries; coalescence of

these single cells with arteries could lead to arterial growth. Alternatively, the

migration and incorporation of single artery cells at the proximal ends of the artery

could indirectly contribute towards distal elongation of arteries. The dynamics of

single artery cell movements could also establish the branching pattern of the pial

artery network; the precise mechanisms for which are unknown.

We show that pre-existing artery tip extensions occur along a distinct

supporting structure. In the heart, single artery endothelial cells migrate by hitch-

hiking on to the capillary networks during the formation of new collaterals. On the

contrary, our study in the embryonic brains reveal the presence of a unique (Tie2[+])

vascular segment on which the arteries extend. These microvascular tracks are

distinct from the surrounding capillary mesh in that they anastomose ACA and MCA

arterial branches but do not show any artery specification. We speculate that the

microvascular tracks form during the specific time window (e16-e17) which coincides

with the period when majority of collaterals appear. Thus, these tracks may play a

crucial role in pial collateral formation. We hypothesize that the cellular origin of

these microvascular tracks is of capillary in nature. Elucidating the underlying

mechanism governing their formation and their molecular identity is an exciting

avenue for future investigation.

Our data about pre-defined microvascular tracks guiding the growth of pial

arteries suggest that the microvasculature could shape the artery network of the

brain. However, the microvascular contribution towards pial collaterals via

arterialization is less likely, as deletion of *Dach1*, a regulator of arterialization,[14,23]

had minimal effect on pial collateral network during embryogenesis. We propose that

as artery tips extend and get connected in the watershed, microvascular cells are

"trapped" within pial collaterals. This could generate a mixed lineage distribution[10]

within pial collaterals—both in embryos and adults. Together, multiple mechanisms

could be actively building pial vasculature during embryogenesis. That being said,

possible contribution from endothelial progenitor cells or peri-vascular cell types

(pericytes, smooth muscle cells) towards pial collateral formation cannot be entirely

excluded.

Distinct mechanism for collateral maintenance

Given that collateral arteries have a unique hemodynamic niche,[24] it would be

interesting to investigate which molecular programs maintain or remodel pial

collaterals in an adult brain. During embryogenesis, pial collaterals are built *de novo*,

in an artery-free zone and remodel as and when blood flows through them. In

contrast, mature collaterals remodel in response to differential blood flow patterns

and oxygen demands of their surrounding tissue. Postnatal remodelling of pial

collaterals is dependent on recruitment of smooth muscle cells and exhibits tortuous

morphology.[24] These fundamental differences could explain the requirement for

distinct molecular programs for building new collaterals and maintaining old ones.

This is evident in our study; VegfR2 controlled pial artery network during

embryogenesis but was dispensable in adulthood. That being said, arterial VegfR2

may regulate other aspects of vascular changes those require activation of

angiogenic pathways[25] such as remodelling induced by hypoxia, ischemia or aging.

We did not notice any remodelling events within adult pial collaterals, in presence or

absence of arterial Cxcr4. Together, adult pial collaterals are maintained via

mechanisms independent of two key angiogenic receptors in arteries—VegfR2 and

Cxcr4. It will be interesting to test if adult collateral maintenance is independent of

signals originating from an arterial niche. It is possible that surrounding

microvasculature plays a more prominent role in maintenance of pial arteries and

collaterals in adults. Investigating the molecular identity of the microvasculature

surrounding pial collaterals could reveal new vascular mechanisms key to collateral

maintenance.

MATERIALS AND METHODS

Mouse lines

Mouse lines	Source/ References
Rosa26^TdTomato	The Jackson Laboratory, stock number 007909
Cx40CreER	Miquerol et al. 2015[26]

ApjCreER	Chen et al. 2014[15]
VegfR2 flox	The Jackson Laboratory, stock number 018977
Cx40[eGFP/+]	Miquerol et al. 2004[27]
Tie2::eGFP::Claudin5	Knowland et al. 2014[28], Lutz et al. 2017[29]
Cxcr4 flox	The Jackson Laboratory, stock number 008767
Dach1 flox	Chang et al. 2017[14]

All experiments were performed on mice with mixed strain background. All mice

were housed and bred at the National Centre for Biological Sciences animal facility

in accordance with the Institutional Animal Ethics Committee (IAEC). All procedures

were performed with approval from IAEC. Data shown in this study is compiled from

age-matched males and females from several litters.

Genotyping primer information

Primer (F, forward; R, Reverse)	Sequence (5' → 3')
Cre F	GCATTACCGGTCGATGCAACGAGTGATGAG
Cre R	GAGTGAACGAACCTGGAAATCAGTGCG
TdTomato mutant F	CTGTTCCTGTACGGCATGG
TdTomato mutant R	GGCATTAAAGCAGCGTATCC
TdTomato wild-type F	AAGGGAGCTGCAGTGGAGTA
TdTomato wild-type R	CCGAAAATCTGTGGGAAGTC
VegfR2 flox F	CCACAGAACAACTCAGGGCTA
VegfR2 flox R	GGGAGCAAAGTCTCTGGAAA

GFP F	GCACGACTTCTTCAAGTCCGCCATGCC
GFP R	GCGGATCTTGAAGTTCACCTTGATGCC
Cxcr4 flox F	CCACCCAGGACAGTGTGACTCTAA
Cxcr4 flox R	GATGGGATTTCTGTATGAGGATTAGC
Dach1 flox F	CTCCTGAAGATGAGGAGCTCACCCC
Dach1 flox R	AACAATTCTTGTCCTTCACGTGCCC

Visualization of pial arteries in fixed samples

Harvesting whole brains/heads from mouse embryos

Depending on the developmental stages and size of the embryos, pial collateral

arteries were visualized either in whole brains (e18.5 or P0) or using whole heads

(e12.5- e16.5).

e12.5-e15.5: To perform the whole head explant at e12.5-e16.5 stages, the pregnant

females were euthanized through carbon dioxide overdose and a Y-shape incision

was made on the abdomen. Intact embryonic sacs were dissected out. Small

punctures were made on the embryo sac with the help of microforceps. The whole

embryo sac was fixed in 4% Paraformaldehyde (PFA) for 2 hours with gentle shaking

at 4°C. Following fixation, the embryo sac was washed at 4°C with gentle shaking.

This washing step was performed with 1X PBS and repeated 4 times every 15-20

mins. The embryos were then extracted from the embryo sacs and the heads were

removed with the help of microforceps.

e16.5: The embryonic explant at e16.5 is performed similarly. However, for

Tie2::eGFP::Claudin5 genotype, the loose skin on top of the brain was removed to

aid immunostaining. Whole heads were imaged within 4 hrs, prior to PFA fixation to

avoid loss of GFP fluorescence signal.

e18.5: To perform the whole brain explant at e18.5 stage, the timed pregnant

females were euthanized as above. The skull of each embryo was gently removed

using the microforceps. The embryonic brains were scooped out with the help of a

microspatula and fixed in 4% PFA at 4°C for 2.5 hours. Agitation was avoided at this

stage to preserve tissue integrity of embryonic brains. The fixed embryonic brains

were then washed with 1X PBS three times at 15 minutes intervals with gentle

shaking at 4°C.

P0: Neonatal explants at P0 were performed by euthanizing the pups through carbon

dioxide overdose followed by decapitation. The skull was removed using scissors

and microforceps. The brains were scooped out using the microspatula. The brains

were then fixed with 4% PFA at 4°C for 2 hrs followed by three washes with 1X PBS

at an interval of 15 mins.

Whole brain/head immunostaining of mouse embryos

e12.5-e15.5: The embryonic heads (e12.5 - e15.5) were incubated in 0.1% PBST

(0.1% TritonX-100 in 1X PBS) for 2.5 hours to permeabilize the tissue, followed by

incubation with primary antibodies overnight at 4°C with gentle shaking. Heads were

then washed with 1X PBS, changing the buffer every 1-2 hours for 1 day. This step

was followed by incubation with secondary antibodies made in 0.1% PBST. Samples

were then washed every 1-2 hours for 1 day with 1X PBS. Following immunostaining

the brains are immersed in an anti-fade mounting media (from Jackson

ImmunoResearch Laboratories Inc.) and imaged immediately.

The embryonic and neonatal brains were incubated with primary antibody (made in

1X PBS) overnight, at 4°C with gentle shaking. Samples were then washed with 1X

PBS, changing the buffer every 1-2 hours for 1 day and incubated with Alexa Fluor

conjugated secondary antibodies (made in 1X PBS). Finally, samples were washed

every 1-2 hours for 1 day with 1X PBS and stored in mounting media or imaged

immediately.

Antibodies used are: Cx40 (Catalog number: Cx40-A, Alpha Diagnostics, 1:500

dilution), Dach1 (Catalog number: 10194-1-AP, Proteintech, 1:500 dilution), VegfR2

(Catalog number: AF644, R&D Systems, 1:250). All secondary antibodies were

procured from Invitrogen and used at a dilution of 1:500.

Imaging whole brains/heads of mouse embryos

The mice heads and brains from different developmental time-points were imaged

using Olympus FV-3000 upright or inverted confocal microscope by placing the

sample in a 2-well chamber Catalog number: 154461; Nunc™ Lab-Tek™ II Chamber

Slide™ System; Thermofisher Scientific. Whole brains and heads were scanned

using 4X and 10X air objectives, and multiple stacks were captured. The captured

image stacks were processed in ImageJ and stitched together using Adobe

Photoshop software.

The embryonic heads at e16.5 was imaged using Olympus FV-3000 5L Upright

microscope. The pre-fixed embryonic heads were attached to a 35mm plastic petri

dish using agarose gel prior to imaging. The whole head imaging was performed

using 4X air objective and watershed images were captured with 20X water

dipping/immersion objective and 1.5X optical zoom with brains immersed in 1X PBS.

Visualization of pial arteries in live mice

In vivo imaging of embryonic pial arteries

Immobilization of live embryo: *Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5*

pregnant dams with embryos at developmental stages between embryonic day

(e)16-e17 were anesthetized with a subcutaneous injection of Ketamine-Xylazine

(Ketamine (80 mg/kg); Xylazine (5 mg/kg)) cocktail. The dams were then placed in a

supine position on a regulated heating pad (TC1000, CWE) that was positioned on

top of a 3D-printed baseplate (as shown in Figure **1A**). A small incision was placed

along the midline of the abdomen. With a gentle retraction of the abdominal skin and

wall, a single uterine horn within the abdominal cavity was isolated. The uterus was

surgically opened and a single embryo along with the yolk sac was isolated and

placed into a 35mm petri dish. The embryo still connected to the dam was

immobilized by submerging the bottom part with 1% agarose, as shown in Figure **1A**.

The immobilized embryo is then submerged in dissection media (DMEM/F12 + 10%

FBS, Catalog number: 21041025, Thermo Fisher Scientific), and the media is

maintained at 37°C using a custom-made Arduino controlled media heating and

circulating device.

<u>Live embryonic imaging:</u> The setup along with anesthetized dam was placed under

Macro Zoom microscope MVX10-Olympus for large scale imaging and under

FV3000 upright confocal Microscope-Olympus for high resolution imaging using 20X

water dipping objective. In Macro Zoom Microscopy, the entire observable depth was

scanned whilst obtaining videos of 1 minute spanning a duration >2 hours in a time-

lapse manner. In confocal microscopy, the entire observable depth was scanned

with a step size of 1 micron. A total of 4 embryos were imaged under the Macro

zoom microscope and 5 embryos were imaged using confocal microscope. Out of

these embryos, the artery cell dynamics was captured in 2 brains from each

category.

<u>Image processing:</u> Image stacks obtained from Macro zoom microscopy was

registered using ImageJ plugin (*StackReg*) and the registered stack was de-

convolved using ImageJ plugin (*DeconvolutionLab2*) and finally maximum intensity

projection was obtained as shown in Figure **1D**. The image stacks obtained from the

confocal microscopy were processed similarly except the deconvolution step.

The MCA and ACA growth fronts were automatically traced and the length of the

traced path was measured using Simple Neurite Tracer (SNT). The distance of

single cells from an artery was measured manually using ImageJ.

In vivo imaging adult pial arteries

<u>Craniotomy:</u> *Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5 and*

VegfR2^{flox/flox};Cx40CreER; Rosa26^{TdTomato};Tie2::eGFP::Claudin5 adult mice (3-5

months) underwent craniotomy to obtain optical access to the pial layer of the brain.

Mice were anesthetized under 2% Isoflurane (Abbott, North Chicago, IL, USA) and

were placed on a stereotaxic mount (Item number 51547; Stoelting, Wood Dale, IL,

USA) with the skull held tightly using ear bars. The mice were kept anesthetized with

2% Isoflurane using a nose-cone. 5mg/kg Dexamethasone in 1X PBS was

administered subcutaneously and the mice kept warm using a heating pad (TC1000,

CWE). The eyes were covered with an eye ointment and the hair above the skull

removed using a hair trimmer. The surgical area was thoroughly wiped clean with

70% ethanol. Sterile surgical tools were used for the surgery. A small incision was

made using a scalpel blade and the exposed skin was removed. The connective

tissue covering the skull in the exposed area was scraped off using scalpel blade

and Q-tips. Using a high-speed electrical drill (burr of diameter ~0.25mm), a window

of ~4mm diameter was carefully etched on the skull. The skull within the window was

eventually detached from the rest of the skull after repeated drilling and was carefully

lifted using a pair of fine forceps (Catalog number: 11255-20, Dumont #55) without

damaging the dura. During the process of drilling, the skull was constantly irrigated

with cortex buffer (Concentration; in mM: NaCl–125, KCl–5, glucose–10, HEPES–10,

$CaCl_2$–2 and $MgCl_2$–2. pH set to 7.35).[30] The exposed dura was aspirated with

cortex buffer until there is no bleeding and the window looked clear. An optical plug

made up of two 4 mm coverslip and one 5 mm coverslip (CS-5R & CS-4R; Warner

Instruments, Hamden, CT, USA) glued together (with NOA 68, Norland Optical

Adhesive) was placed on the exposed dura. The optical plug was gently pressed

using a toothpick such that the 5mm cover slip was touching the skull bone and then

cyano-acrylic glue was applied across the perimeter of the 5mm coverslip attaching it

to the skull. At this stage we had an optical window of 4mm. An aluminium head bar

with circular hole of 8mm was placed onto the skull aligned to the previously made

optical window. Next, dental cement (Catalog number 56877, 3M Relyx U200) was

applied over the exposed skull using a sterile toothpick while being careful to not get

it on the optical window. Once the dental cement was completely dry, the animal was

placed on a warm pad in a clean recovery cage. Animals undergoing craniotomy

were maintained on meloxicam (2 mg/kg) and enrofloxacin (5 mg/Kg body weight)

for 3-4 days until full recovery. Mice were further allowed to recover for at least 2

weeks post-surgery before the start of any experiment. Mice with an optical window

can be imaged longitudinally for 2-3 months until the optical window starts to lose its

clarity.

Live Adult Imaging: To perform in vivo imaging, mice were anesthetized through

intraperitoneal injection of Ketamine (80mg/Kg body weight) – Xylazine (5mg/Kg

body weight) cocktail. The eyes were covered with an eye ointment and the mice

were placed onto a custom-made baseplate and immobilized using the implanted

metal head plate to this baseplate. A heating pad (37°C) was placed under the

mouse to maintain the body temperature. Subsequently, the baseplate was placed

on a motorized XY-table of an upright confocal microscope (FV3000 upright confocal

Microscope-Olympus). The entire optical window was imaged using 10X air objective

(0.4 NA) and selected pial regions with collateral arteries were imaged with 20X

water dipping objective (0.5 NA). A total of N=3 adult mice per group (control and

VegfR2 artery endothelial specific knockout) were imaged longitudinally for a period

of 1 month.

<u>Image processing</u>: The 10X image stacks obtained from the confocal microscopy of

the cranial window were registered using ImageJ plugin (*StackReg*) and maximum

intensity projection was obtained. The region of interest from the 20X image stacks

were cropped out slice by slice and stitched in Adobe Photoshop to avoid the

overlaying microvascular network in order to focus on the pial collaterals.

Quantifications

Lineage contribution towards pial collaterals during embryogenesis

Cx40 immunostained pial collateral segments from either *Cx40CreER;*

Rosa26^{Tdtomato} or *ApjCreER; Rosa26^{TdTomato}* brains were imaged at 10X with 2.12

optic zoom. Regions of interest were manually drawn to mark the collateral artery

segments. Lineage contribution was determined by quantifying TdTomato$^+$ coverage

on the collateral segment. This was performed by thresholding the images with Otsu

algorithm in ImageJ. The area fraction of the TdTomato$^+$ area per collateral segment

was calculated using the Measure tool in ImageJ. Contribution of the artery and

capillary derived ECs in collaterals was quantified by calculating the percentage of

collateral area covered by TdTomato$^+$ signal. 45 collateral segments (of an

approximate length of 600-800μm) from 7 *Cx40CreER*-lineage traced brains and 57

collateral segments from 8 *ApjCreER*-lineage traced brains were analysed.

Number of collateral arteries per hemisphere

In *Cx40CreER* TdTomato labelled brains, the collateral number was quantified based

on the number of contiguous TdTomato$^+$ vessels (Td-contiguous) connecting MCA

and ACA. Data shown in Figure **3H** was summarized from 2 control and 3 artery-

specific *VegfR2* knockout. Data shown in Figure **S2H** was summarized from e13.5

(N=5), e14.5 (N=5), e15.5 (N=5), e18.5 (N=5) and P0 (N=5) embryos and neonates.

Data shown in Figure **S3H** was summarized from e13.5 (N=3), e14.5 (N=3), e15.5

(N=2), e18.5 (N=2) and P0 (N=5) embryos and neonates. Data shown in Figure **S6C**

is quantified by counting number of Cx40 immunostained collateral arteries between

MCA and ACA, and is summarized from 4 control and 7 *Dach1* capillary specific

knockouts. An unpaired t-test was done to test significance statistically.

Efficiency of TdTomato recombination

Segments of collateral arteries, proximal arteries (PA) and distal arteries (DA) from

the image stacks taken with 20X objective were obtained from live adult imaging of

Cx40CreER; *Rosa26^{TdTomato}*; *Tie2::eGFP::Claudin5* mice. TdTomato$^+$ and TdTomato$^-$

cells were counted in each artery segment. Each segment was of an approximate

length of 600-800μm.The recombination efficiency was measured as ratio of

TdTomato$^+$ cells and total number of cells in the segment. A total of 10 collateral

segments, 10 PA segments and 10 DA segments from 4 mice were analysed to

obtain the recombination efficiency of TdTomato. One-way ANOVA was done to test

significance statistically (mean comparison Tukey test, PA vs. collateral, $p<0.0001$;

DA vs. collateral, $p<0.0001$; PA vs. DA, $p=0.08$).

Pial artery coverage in whole brains and in the collateral zone

The image stacks were obtained using confocal imaging of whole embryonic brains

with 10X objective, stitched using ImageJ plugin (*Stitching*) and max intensity

projection (MIP) was obtained. The MIPs were used for further analysis wherein one

complete hemisphere was cropped out and aligned vertically to the major axis of its

own elliptical boundary. Using a custom ImageJ macro, a line of the length of the

major axis was scanned across the width of the image and intensity profile for every

pixel along the horizontal axis was generated. Total number of local maxima (peaks)

for each intensity profile was obtained using the built-in ImageJ macro function

Array.findMaxima (array, tolerance). The tolerance used was 2 times the average

background intensity. The number of peaks thus obtained was plotted across the

width and the plot was fitted to a Gaussian function using OriginPro 2023b. The fitted

curve was used as a representative of the arterial coverage per hemisphere. The fit

curves obtained from various mice were aligned based on the Gaussian peaks and

averaged to get the representative curves for various experimental groups. We

considered a region of 1mm (-0.5mm to +0.5 mm) across the Gaussian peak as the

collateral zone (watershed). The area under the curve within this region was used as

a measure for the extent of the collateral network. 9 hemispheres from 9 embryos

(e18.5) were analysed per group (control and artery-specific knockout for *VegfR2*).

To test the statistical significance of the differences observed in pial artery coverage

(Figure **3E**), a Kolmogorov-Smirnov (KS) test was done. D value= 0.569, p value=

2.2066 X 10^{-141}. To test the statistical significance of the area under the curve (pial

collateral artery network shown in Figure **3F**), an unpaired t-test was done

(p=0.0001).

AUTHORS CONTRIBUTION

SD conceived the idea and designed the experiments. SK performed intravital

imaging of embryos and adults. NS, SG, VS, MD performed other embryonic

experiments. SD, SK, NS, SG wrote the manuscript. All authors contributed to

editing the manuscript.

ACKNOWLEDGEMENTS

We thank the following facilities for their technical support: Animal Care and

Resource Center at NCBS, Mouse Genome Engineering Facility at NCBS, Central

Imaging and Flow Cytometry Facility at NCBS. We sincerely thank Dr. Dritan Agalliu

(Columbia University) and Dr. Kristy Red-Horse (Stanford University) for generously

sharing *Tie2::eGFP::Claudin5* and *Dach1$^{fl/fl}$* mice respectively. We thank our funding

sources. This work is supported by NCBS/TIFR core funding and DBT/Wellcome

Trust India Alliance Intermediate Fellowship to SD (IA/I/20/2/505205). SK is

supported by DBT/Wellcome Trust India Alliance Intermediate Fellowship to SD. NS

and VS were supported by DBT INSPIRE Scholarship for Higher Education (SHE).

DECLARATION OF INTERESTS

None

REFERENCES

1. Lima, F.O., Furie, K.L., Silva, G.S., Lev, M.H., Camargo, E.C.S., Singhal, A.B., Harris, G.J., Halpern, E.F., Koroshetz, W.J., Smith, W.S., et al. (2010). The pattern of leptomeningeal collaterals on CT angiography is a strong predictor of long-term functional outcome in stroke patients with large vessel intracranial occlusion. Stroke *41*, 2316–2322. 10.1161/STROKEAHA.110.592303.

2. Maas, M.B., Lev, M.H., Ay, H., Singhal, A.B., Greer, D.M., Smith, W.S., Harris, G.J., Halpern, E., Kemmling, A., Koroshetz, W.J., et al. (2009). Collateral vessels on CT angiography predict outcome in acute ischemic stroke. Stroke *40*, 3001–3005. 10.1161/STROKEAHA.109.552513.

675   3.   World Health Organisation. The top 10 causes of death

676   https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death.

677   4.   Zhang, H., Prabhakar, P., Sealock, R., and Faber, J.E. (2010). Wide genetic

678   variation in the native pial collateral circulation is a major determinant of

679   variation in severity of stroke. J. Cereb. Blood Flow Metab. *30*, 923–934.

680   10.1038/jcbfm.2010.10.

681   5.   Clayton, J.A., Chalothorn, D., and Faber, J.E. (2008). Vascular endothelial

682   growth factor-a specifies formation of native collaterals and regulates collateral

683   growth in ischemia. Circ. Res. *103*, 1027–1036.

684   10.1161/CIRCRESAHA.108.181115.

685   6.   Lucitti, J.L., MacKey, J.K., Morrison, J.C., Haigh, J.J., Adams, R.H., and Faber,

686   J.E. (2012). Formation of the collateral circulation is regulated by vascular

687   endothelial growth factor-a and a disintegrin and metalloprotease family

688   members 10 and 17. Circ. Res. *111*, 1539–1550.

689   10.1161/CIRCRESAHA.112.279109.

690   7.   Lucitti, J.L., Sealock, R., Buckley, B.K., Zhang, H., Xiao, L., Dudley, A.C., and

691   Faber, J.E. (2016). Variants of Rab GTPase-Effector Binding Protein-2 Cause

692   Variation in the Collateral Circulation and Severity of Stroke. Stroke *47*, 3022–

693   3031. 10.1161/STROKEAHA.116.014160.

694   8.   Chalothorn, D., Clayton, J.A., Zhang, H., Pomp, D., and Faber, J.E. (2007).

695   Collateral density, remodeling, and VEGF-A expression differ widely between

696   mouse strains. Physiol. Genomics. *30*, 179–191.

697   10.1152/PHYSIOLGENOMICS.00047.2007.

9.   Das, S., Goldstone, A.B., Wang, H., Farry, J., D'Amato, G., Paulsen, M.J.,

Eskandari, A., Hironaka, C.E., Phansalkar, R., Sharma, B., et al. (2019). A

Unique Collateral Artery Development Program Promotes Neonatal Heart

Regeneration. Cell *176*, 1128-1142.e18. 10.1016/J.CELL.2018.12.023.

10. Perovic, T., Hollfinger, I., Mayer, S., Lips, J., Dopatka, M., Harms, C., and

Gerhardt, H. (2023). Pial collaterals develop through mosaic colonization of

capillaries by arterial and microvascular endothelial cells. bioRxiv,

2023.10.09.561542. 10.1101/2023.10.09.561542.

11. Jamaiyar, A., Juguilon, C., Wan, W., Richardson, D., Chinchilla, S., Gadd, J.,

Enrick, M., Wang, T., McCabe, C., Wang, Y., et al. (2022). The essential role

for endothelial cell sprouting in coronary collateral growth. J. Mol. Cell. Cardiol.

165, 158–171. 10.1016/j.yjmcc.2022.01.005.

12. Gabhann, F. Mac, and Peirce, S.M. (2010). Collateral Capillary Arterialization

following Arteriolar Ligation in Murine Skeletal Muscle. Microcirculation *17*,

333–347. 10.1111/J.1549-8719.2010.00034.X.

13. Arolkar, G., Kumar, S.K., Wang, H., Gonzalez, K.M., Kumar, S., Bishnoi, B.,

Rios Coronado, P.E., Woo, Y.J., Red-Horse, K., and Das, S. (2023).

Dedifferentiation and Proliferation of Artery Endothelial Cells Drive Coronary

Collateral Development in Mice. Arterioscler. Thromb. Vasc. Biol. *43*, 1455–

1477. 10.1161/ATVBAHA.123.319319.

14. Chang, A.H., Raftrey, B.C., D'Amato, G., Surya, V.N., Poduri, A., Chen, H.I.,

Goldstone, A.B., Woo, J., Fuller, G.G., Dunn, A.R., et al. (2017). DACH1

stimulates shear stress-guided endothelial cell migration and coronary artery

721  growth through the CXCL12-CXCR4 signaling axis. Genes. Dev. *31*, 1308–

722  1324. 10.1101/GAD.301549.117/-/DC1.

723 15. Chen, H.I., Sharma, B., Akerberg, B.N., Numi, H.J., Kivelä, R., Saharinen, P.,

724  Aghajanian, H., McKay, A.S., Bogard, P.E., Chang, A.H., et al. (2014). The

725  sinus venosus contributes to coronary vasculature through VEGFC-stimulated

726  angiogenesis. Development *141*, 4500–4512. 10.1242/dev.113639.

727 16. Scott Budinger, G.R., Duranteau, J., Chandel, N.S., and Schumacker, P.T.

728  (1998). Hibernation during Hypoxia in Cardiomyocytes ROLE OF

729  MITOCHONDRIA AS THE O_2 SENSOR. J. Biol. Chem. *273*, 3320–3326.

730  10.1074/jbc.273.6.3320.

731 17. Rink, C., and Khanna, S. (2011). Significance of Brain Tissue Oxygenation and

732  the Arachidonic Acid Cascade in Stroke. https://home.liebertpub.com/ars *14*,

733  1889–1903. 10.1089/ARS.2010.3474.

734 18. Mannion, J.D., Buckman, P.D., Magno, M.G., and Dimeo, F. (1992). Collateral

735  blood flow from skeletal muscle to normal myocardium. J. Surg. Res. *53*, 578–

736  587. 10.1016/0022-4804(92)90258-2.

737 19. Crissinger, K.D., and Neil Granger, D. (1988). Characterization of Intestinal

738  Collateral Blood Flow in the Developing Piglet. Pediatr. Res. *24*, 473–476.

739  10.1203/00006450-198810000-00011.

740 20. Kochi, T., Imai, Y., Takeda, A., Watanabe, Y., Mori, S., Tachi, M., and

741  Kodama, T. (2013). Characterization of the Arterial Anatomy of the Murine

742  Hindlimb: Functional Role in the Design and Understanding of Ischemia

743  Models. PLoS One *8*, e84047. 10.1371/JOURNAL.PONE.0084047.

21. Gao, Y., Aravind, S., Patel, N.S., Fuglestad, M.A., Ungar, J.S., Mietus, C.J., Li,

S., Casale, G.P., Pipinos, I.I., and Carlson, M.A. (2020). Collateral

Development and Arteriogenesis in Hindlimbs of Swine After Ligation of

Arterial Inflow. J. Surg. Res. *249*, 168–179. 10.1016/J.JSS.2019.12.005.

22. He, L., Liu, Q., Hu, T., Huang, X., Zhang, H., Tian, X., Yan, Y., Wang, L.,

Huang, Y., Miquerol, L., et al. (2016). Genetic lineage tracing discloses

arteriogenesis as the main mechanism for collateral growth in the mouse

heart. Cardiovasc. Res. *109*, 419–430. 10.1093/CVR/CVW005.

23. Raftrey, B., Williams, I.M., Rios Coronado, P.E., Fan, X., Chang, A.H., Zhao,

M., Roth, R., Trimm, E., Racelis, R., D'amato, G., et al. (2021). Dach1 Extends

Artery Networks and Protects Against Cardiac Injury. Circ. Res. *129*, 702–716.

10.1161/CIRCRESAHA.120.318271.

24. Zhang, H., Chalothorn, D., and Faber, J.E. (2019). Collateral Vessels Have

Unique Endothelial and Smooth Muscle Cell Phenotypes. Int. J. Mol. Sci. *20*,

3608. 10.3390/IJMS20153608.

25. Stanca Melincovici, C., Boşca, A.B., Şuşman, S., Mărginean, M., Mihu, C.,

Istrate, M., Moldovan, I.-M., Roman, A.L., and Mihu, C.M. (2018). Vascular

endothelial growth factor (VEGF)-key factor in normal and pathological

angiogenesis. Rom. J. Morphol. Embryol. *59*, 455-467.

26. Miquerol, L., Thireau, J., Bideaux, P., Sturny, R., Richard, S., and Kelly, R.G.

(2015). Endothelial plasticity drives arterial remodeling within The endocardium

after myocardial infarction. Circ. Res. *116*, 1765–1771.

10.1161/CIRCRESAHA.116.306476.

27.   Miquerol, L., Meysen, S., Mangoni, M., Bois, P., van Rijen, H.V.M., Abran, P.,

Jongsma, H., Nargeot, J., and Gros, D. (2004). Architectural and functional

asymmetry of the His–Purkinje system of the murine heart. Cardiovasc. Res.

63, 77–86. 10.1016/j.cardiores.2004.03.007.

28.   Knowland, D., Arac, A., Sekiguchi, K.J., Hsu, M., Lutz, S.E., Perrino, J.,

Steinberg, G.K., Barres, B.A., Nimmerjahn, A., and Agalliu, D. (2014).

Stepwise Recruitment of Transcellular and Paracellular Pathways Underlies

Blood-Brain Barrier Breakdown in Stroke. Neuron *82*, 603–617.

10.1016/j.neuron.2014.03.003.

29.   Lutz, S.E., Smith, J.R., Kim, H., Bates, J.M., Gandhi, S.P., and Agalliu, D.

(2017). Caveolin1 Is Required for Th1 Cell Infiltration, but Not Tight Junction

Remodeling, at the Blood-Brain Barrier in Autoimmune Neuroinflammation.

CellReports *21*, 2104–2117. 10.1016/j.celrep.2017.10.094.

30.   Holtmaat, A.J.G.D., Trachtenberg, J.T., Wilbrecht, L., Shepherd, G.M., Zhang,

X., Knott, G.W., and Svoboda, K. (2005). Transient and Persistent Dendritic

Spines in the Neocortex In Vivo. Neuron *45*, 279–291.

10.1016/j.neuron.2005.01.003.

Main figures and figure legends

Figure 1: Live imaging of embryonic pial vasculature

(**A**) Intra-vital imaging set up to visualize the developing pial vasculature in mouse

embryos. **i** heating pad to maintain body temperature of anaesthetized pregnant dam

at 37°C. **ii** custom-built media heating device which also circulates dissection media

at 37°C **iii** shows macro-zoom or confocal microscopy performed on live,

immobilized embryos.

(**B**) Representative schematic of an embryonic brain. Grey and Red show cerebral

arteries and pial collaterals respectively.

(**C**) Experimental design to genetically label and trace the dynamics of pre-existing

artery endothelial cells in a developing pial layer.

(**D-G**) Images obtained through macro-zoom fluorescent microscopy of e16.5

embryos. In black are cells derived from pre-existing artery endothelial cells. Images

were taken by focusing in and out to capture the entire curvature of the embryonic

brain. **D** shows an entire left hemisphere. **E** represents watershed region spanned by

MCA and ACA artery tips and imaged at time=0 minutes, 20 minutes and 80 minutes

respectively. Arrows and arrowheads show growing ends of pre-existing artery tips

from MCA and ACA. Brackets mark regions where collaterals appear. **F** shows inset

in **D**. **E** and **G** show insets in **F**. **G** represents the MCA growth front imaged at time=0

minutes, 20 minutes and 80 minutes respectively. * shows a superficial yolk sac

vessel which photobleaches with time. Arrowheads mark extension and

anastomoses of MCA artery tips.

(**H-J**) Images obtained through confocal microscopy from e16 embryos. **H** shows pial

watershed and the cerebral arteries spanning the watershed. **I, J** show MCA and

ACA growth fronts respectively imaged at time=0 minutes and 60 minutes. In black

are cells derived from pre-existing artery endothelial cells. Images shown in **I** are

maximum intensity projections from 50 optical slices and capture the entire width of

the artery. Images shown in **J** are maximum intensity projections from 40 optical

slices and capture the entire width of the artery * represents an internal landmark

from where the length measurements were done.

(**K**) Confocal images of e17 pial layer showing presence of TdTomato[+] pre-existing

artery endothelial cells (arrowheads) on eGFP[+] microvascular tracks. Arrowheads

point to TdTomato[+] artery cells (red) on microvascular tracks (green).

(**L**) A model proposing extension of artery tips along microvascular tracks which

leads to pial collateral formation during embryogenesis. Red arrows point to the

direction of artery tip elongation. MCA, Middle cerebral artery; ACA, Anterior cerebral

artery; e, embryonic day; Tam, Tamoxifen; EC, endothelial cell; YSV, yolk sac

vessel; brch., branch; coll, collateral; MV, microvascular. Scale bars: **D**, **F** 1mm; **E**, **G**

200 µm; **H**, 500 µm; **I**, **J**, **K** 100 µm.

Figure 2: Cellular origin of pial collaterals

(**A**) Experimental design to assess lineage contribution from artery ECs and capillary

ECs towards pial collaterals at e18.5/ P0.

(**B**) Graph shows the average percent area of collateral segments covered with

TdTomato⁺ cells.

(**C**) Representative image of whole right hemisphere from *Cx40CreER*;

Rosa26^{TdTomato} e18.5 embryos. Black shows TdTomato⁺ arteries.

(**D**) Boxed region of interest with a collateral segment from the watershed area

shown in **C**. The pial collateral is lineage traced for artery ECs (red) and

immunostained with Cx40 (green).

(**E**) Representative image of whole right hemisphere from *ApjCreER*; *Rosa26^{TdTomato}*

e18.5 embryos. Black shows Cx40 immunostaining.

(**F**) Boxed region of interest with a collateral segment from the watershed area

shown in **E**. The pial collateral is lineage traced for capillary ECs (magenta) and

immunostained with Cx40 (cyan).

(**G**, **I**) Collateral segment from **D** is (**G**) immunostained for Cx40 and (**I**) lineage

traced with TdTomato.

(**H**, **J**) Collateral segment from **F** is (**H**) immunostained for Cx40 and (**J**) lineage

traced with TdTomato. Arrowheads point to TdTomato⁺ cells from Apj⁺ lineage on the

collateral segment. e, embryonic day, P, postnatal day; MCA, Middle cerebral artery;

ACA, Anterior cerebral artery; brch., branch; EC, endothelial cell. Scale bars: **C** and

E, 500µm; **D** and **F**, 100µm

A

Cxcr4^{L/L}; Cx40CreER; Rosa26^{TdTomato}
and
VegfR2^{L/L}; Cx40CreER; Rosa26^{TdTomato}

B Arterial depletion of Cxcr4
does not change pial collateral artery network

Cx40CreER;Rosa26^{TdTomato}   *Cxcr4^{L/L};Cx40CreER;Rosa26^{TdTomato}*

C
Quantification of pial artery coverage
in watershed

D Arterial VegfR2 regulates formation of
pial collateral artery network

Cx40CreER;Rosa26^{TdTomato}   *VegfR2^{L/L};Cx40CreER;Rosa26^{TdTomato}*

E Reduced artery coverage
in absence of arterial VegfR2

F Reduced collateral artery network
in absence of arterial VegfR2

Cx40CreER;Rosa26^{TdTomato}
VegfR2^{L/L};Cx40CreER;Rosa26^{TdTomato}

VegfR2-null artery tips fail to extend and anastomose

G *Cx40CreER; Rosa26^{TdTomato}*
or
Cx40CreER; Rosa26^{TdTomato}
Tie2::eGFP::Claudin5

H Reduced number of
Td^{con} collaterals
without arterial VegfR2

I *Cx40CreER; Rosa26^{TdTomato} Tie2::eGFP::Claudin5*
TdTomato (pre-existing artery ECs)   eGFP (all ECs)

Figure 3: VegfR2-dependent formation of pial collateral arteries

(**A**) Experimental design to lineage label and trace pre-existing artery endothelial cells in pial layer, with or without arterial *VegfR2* and *Cxcr4* at e18.5/ P0.

(**B**) Representative confocal images of whole brains with and without arterial *Cxcr4*. Black shows pre-existing artery endothelial cells. Control= 8, artery specific knockout for *Cxcr4*= 13.

(**C**) Summary of method to quantify pial artery coverage across the entire brain including watershed.

(**D**) Confocal images of whole brains with and without arterial *VegfR2*. Black shows pre-existing artery endothelial cells.

(**E**) Quantification of overall pial artery coverage with and without arterial *VegfR2* quantified using method shown in **C**. KS test performed on the distributions generated D value= 0.569, p value= 2.2066×10^{-141}.

(**F**) Quantification of pial collateral artery network with and without arterial *VegfR2* quantified using method shown in **C**. Unpaired t-test, p=0.0001.

(**G**) Experimental design to assess the effect of *VegfR2* deletion from arteries at e16.5. (**H**) Quantification of pial collateral arteries contiguously populated with TdTomato[+] artery cells, with and without arterial *VegfR2*. (**I**, **J**) Confocal images of pial watershed region showing TdTomato-contiguous pial collateral arteries (Td[con]) in red and microvasculature in green. Arrowheads show part of microvascular track not occupied by TdTomato[+] artery cell, arrows point to artery tips on microvascular tracks. MCA, Middle cerebral artery; ACA, Anterior cerebral artery; e, embryonic day; P, postnatal day; Tam, Tamoxifen; EC, endothelial cell; Td[con], TdTomato-contiguous;

883   AU, Arbitrary Unit; aEC, artery endothelial cell; KO, knockout. Scale bars: **B**, **D**,

884   500µm; **I**, **J**, 100µm.

Intra-vital imaging of pial collaterals in adult mice

A Dorsal view of cranial window

B Side view of cranial window

C *Cx40CreER; Rosa26^TdTomato*
Tie2::eGFP::Claudin5

D TdTomato Recombination efficiency (%)

Cx40CreER; Rosa26^TdTomato *Tie2::eGFP::Claudin5*
TdTomato (pre-existing artery ECs) eGFP (all ECs)

E watershed

F t=0 days

G t=14 days

H t=28 days

I Pial arteries — Pial veins
PA, proximal artery   coll, collateral
DA, distal artery

J t=0 days | t=14 days | t=28 days

K t=0 days | t=14 days | t=28 days

L t=0 days | t=14 days | t=28 days

M t=0 days | t=14 days | t=28 days

Figure 4: Longitudinal live-imaging of adult pial collaterals

(**A**, **left**) Dorsal view of the cranial window created with the attached headbar on adult mouse head. (**A**, **right**) Brightfield image of the 4mm cranial window obtained with a stereoscope shows the pial vasculature. Dotted rectangle encompasses MCA, ACA and the watershed between them and is imaged at high resolution as shown in **E**.

(**B**) Schematic of the side view of the cranial window showing the 3-tier optical plug inserted into the skull giving access to pial layer.

(**C**) Experimental design for longitudinal imaging of pial vasculature in adult mice.

(**D**) Quantification of TdTomato reporter recombination in pial collateral arteries and different segments of cerebral arteries. One-way ANOVA was done to test significance statistically (mean comparison Tukey test, PA vs. collateral, $p<0.0001$; DA vs. collateral, $p<0.0001$; PA vs. DA, $p=0.08$).

(**E-K**) Images of pial watershed with pre-existing artery endothelial cells labelled in TdTomato (red) and all endothelial cells labelled in eGFP (green). (**E**) Confocal image from optical window created for longitudinal imaging of *Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5* mouse as shown in Figure **4A**. The image shows both pial and dural vessels. (**F-K**) Confocal images of pial collateral arteries from cranial window shown in **E**, across 3 timepoints, each 2 weeks apart. (**I**) Pial layer segmented out from **E**. red and blue show pial arteries and veins respectively. **J**, **K** are the insets shown in **F-H**.

(**L**, **M**) Cellular distribution of TdTomato+ artery cells and eGFP+ endothelial cells in collateral artery segments shown in **J** and **K** respectively. * represents a reference

913   cell. MCA, middle cerebral artery; ACA, anterior cerebral artery; d, days; Tam,

914   Tamoxifen; PA, proximal artery; DA, distal artery; coll, collateral; brch., branch; EC,

915   endothelial cell. Scale bars: **E**, 200μm; **J**, **K** 25μm

A *VegfR2^{L/L};Cx40CreER; Rosa26^{TdTomato}*
Tie2::eGFP::Claudin5

Craniotomy

14d | dose 1 dose 2 dose 3 dose 4 dose 5 | 7d 14d 14d

5d

label pre-existing arteries + *VegfR2* deletion

Tam

t=0d t=14d t=28d
intra-vital imaging

B 4mm diameter cranial window

MCA brch. — ACA brch.

C

C watershed
TdTomato (pre-existing artery ECs)
eGFP (all ECs)

ACA brch.

MCA brch.

E-G

collateral

D Pial arteries Pial veins

collateral

MCA brch.

ACA brch.

collateral

MCA-ACA collateral artery

E t=0 days   **F** t=14 days   **G** t=28 days

H   I   J

H t=0 days   **I** t=14 days   **J** t=28 days

K   **L**   **M**

N VegfR2 regulates pial collateral development by artery tip extension

development (embryogenesis)
VegfR2-driven artery-tip extension

maintenance (adulthood)
VegfR2/Cxcr4 independent process

pial collateral arteries

flow from MCA   flow from ACA

artery lineage   non-artery lineage

Figure 5: Analyses of role of arterial VegfR2 in pial collateral maintenance

(A) Experimental design to visualize artery cell dynamics in pial layer of adult mice without arterial *VegfR2*.

(**B**) Brightfield image of cranial window showing pial vasculature. The dotted rectangle encompasses branches of MCA, ACA and the watershed between them.

(**C**) Confocal image of cranial window created in mice without arterial *VegfR2* as shown in **B**. Red shows all artery cells, green shows all endothelial cells.

(**D**) Pial layer segmented out from **C** shows pial arteries and veins in red and blue respectively.

(**E**-**G**) Confocal images of collateral artery from three timepoints, each 2 weeks apart as shown in the boxed region in **C**.

(**H**-**J**) Confocal images of collateral artery segments shown in **E**-**G**.

(**K**-**M**) Cellular distribution of TdTomato⁺ artery cells and eGFP⁺ endothelial cells in collateral artery segments shown in **H**-**J**. * represents a reference cell.

(**N**) Schematic summarizing the role of arterial VegfR2 in development and maintenance of pial collateral arteries. While arterial VegfR2 facilitates tip extension of pre-existing arteries during development, it is dispensable for pial collateral maintenance in adulthood. Pial collaterals, both in developing and mature brains, are majorly populated by endothelial cells of artery-lineage. MCA, middle cerebral artery; ACA, anterior cerebral artery; d, days; Tam, Tamoxifen; EC, endothelial cell; brch., branch. Scale bars: **C**, 150 µm; **E**-**G**, 50µm; **H**-**J**, 20µm.

Supplementary figures and figure legends

Development of cerebral artery trees and pial collaterals in embryonic mouse brains

Supplemental Figure 1: Visualization of pial arteries and collateral arteries

Schematic summarizes key steps of immunostaining methods used to image pial artery network in (**A**) mouse embryonic heads and (**B**) mouse embryonic/neonatal brains with confocal microscope. (**C**) Schematic shows findings from **A** and **B** in one hemisphere. Grey represents cerebral arteries, Red represents collateral artery connections between cerebral arteries. Brackets show watershed area (WSA), MCA appears at e12.5, ACA appears at e13.5, and the foremost pial collateral arteries appear at e15.5. These pial collaterals then increase in number till e18.5/P0. e, embryonic; P, postnatal day; MCA, middle cerebral artery; ACA, anterior cerebral artery.

Cx40 immunostaining

Supplemental Figure 2: Detection of pial collaterals with whole brain immunostaining

(**A-F**) Representative images of right hemispheres from developing mouse embryos immunostained for Connexin-40 (Cx40). Cx40 marks all artery endothelial cells. MCA appears at e12.5, ACA appears at e13.5 and collaterals between MCA and ACA appear at e15.5. Arrowheads point to collateral arteries between MCA and ACA. (**G**) Quantification of Cx40 immunostained brains from different developmental stages reveals that both MCA and ACA have formed by e14.5. (**H**) Quantification of Cx40$^+$ collateral arteries per hemisphere shows that the first pial collaterals appear at e15.5. e, embryonic; P, postnatal day; MCA, middle cerebral artery; ACA, anterior cerebral artery; brch., branch. Scale bars: 500μm

left hemisphere

e12.5
(MCA appears)

e13.5

right hemisphere

C e14.5
(ACA appears)

D e15.5
(collateral arteries
appear)

E e18.5

F P0

G MCA and ACA artery trees form by e14.5

H Pial collateral arteries appear at e15.5

Supplemental Figure 3: Detection of pial collaterals with *Cx40* reporter activity

(**A-F**) Representative images of brains from developing *Cx40eGFP/+* mouse embryos. Connexin-40 (Cx40) marks all artery endothelial cells. MCA appears at e12.5, ACA appears at e14.5 and collaterals between MCA and ACA appear at e15.5. Arrowheads point to collateral arteries between MCA and ACA. (**G**) Quantification of *Cx40eGFP/+* brains from different developmental stages reveals that both MCA and ACA have formed by e14.5. (**H**) Quantification of eGFP$^+$ collateral arteries per hemisphere shows that the first pial collaterals appear at e15.5. e, embryonic; P, postnatal day; MCA, middle cerebral artery; ACA, anterior cerebral artery; brch., branch. Scale bars: 500µm

Supplemental Figure 4: Visualization of pial artery cell dynamics on

microvascular tracks (**A**) Confocal image showing watershed region between MCA

and ACA of a live e16 embryo. (**B**, **C**) Insets from **A** showing growth of ACA in 60

mins. In **B**, arrowhead point to an artery tip emerging within 60mins. Arrow points to

single artery cells. In **C**, arrow shows direction of migration of single artery cell.

Dotted line is the length of micro-vessel between the single cell and the pial artery. *

shows a reference point from which measurements were done. (**D**) Confocal image

of another e16 embryonic collateral zone showing single artery endothelial cell

migration towards the pial collateral within 110mins. Dotted line is the length of

micro-vessel between the single cell and the pial collateral. Arrow shows direction of

single artery cell migration. (**E**) Inset from **A** showing presence of TdTomato$^+$ artery

endothelial cells (yellow arrowheads) on eGFP$^+$ microvascular tracks (magenta

arrows). MCA, Middle Cerebral Artery; ACA, Anterior Cerebral Artery; MV,

Microvascular; Td, TdTomato; EC, Endothelial cell. Scale bars: **A**, 500µm; **B-E**,

100µm.

1004

A   *Cxcr4^{L/L};Cx40CreER; Rosa26^{TdTomato}*
Tie2::eGFP::Claudin5

artery-EC specific deletion of *Cxcr4*
does not impact adult pial collateral artery network

B   *Cx40CreER;Rosa26^{TdTomato}*

C   *Cxcr4^{L/L};Cx40CreER;Rosa26^{TdTomato}*

1005

1006

1007

1008

1009

1010

1011

1012

1013

Supplemental Figure 5: Pial artery coverage does not change without arterial

Cxcr4 (**A**) Experimental set up to assess the effect of arterial *Cxcr4* deletion on pial

artery coverage in adults. (**B**, **C**) Representative confocal images of whole adult

brains (from N=5 for each genotype), genetically lineage traced for pre-existing

artery endothelial cells, (**B**) with or (**C**) without arterial *Cxcr4*. ACA, anterior cerebral

artery; MCA, middle cerebral artery. Scale bars: 1mm

A

Dach1 is highly expressed in pial capillary ECs at P0

No effect on collateral development with deletion of *Dach1* from capillary ECs

Supplemental Figure 6: Dach1-independent pial collateral development (**A**)

Dach1 expression (green) in capillaries and arteries in P0 brains. Dotted line marks

the TdTomato⁺ (red) arterial boundary. VegfR2 marks all vasculature. DAPI marks

nuclei. (**B**) Experimental design showing Tamoxifen injection at e14.5, and analyses

of collateral numbers in e18.5/P0 brains. (**C**) Quantification of Cx40⁺ pial collaterals

between MCA and ACA in control (*Dach1ᶠˡ/ᶠˡ*) and *ApjCreER*; *Dach1ᶠˡ/ᶠˡ* (Dach1 cap-

EC KO) brains. e, embryonic; cap-EC, capillary endothelial cell; KO, knockout. Scale

bar: 20µm.

1048

Intra-vital imaging of VegfR2 deficient pial collaterals in adult mice

VegfR2$^{L/L}$ Cx40CreER; Rosa26^{TdTomato}
Tie2::eGFP::Claudin5
TdTomato (pre-existing artery ECs) eGFP (all ECs)

1049

1050

1051

1052

1053

1054

Supplemental Figure 7: VegfR2-independent maintenance of pial collaterals

Confocal images of the entire cranial window from 3 different mice shows that the overall pial artery network remains unchanged without arterial *VegfR2*. MCA, Middle Cerebral Arteries, ACA, Anterior Cerebral Arteries, Coll, Pial collateral arteries between MCA and ACA. Scale bars: 500µm.