Development of Pial Collaterals by Extension of Pre-existing Artery Tips

Suraj Kumar, Niloufer Shanavas, Swarnadip Ghosh, Vinayak Sivaramakrishnan,

Manish Dwari, Soumyashree Das[#]

National Centre for Biological Sciences, Tata Institute of Fundamental Research,

Bengaluru, KA 560065, India

[#] Corresponding author (soumyashree@ncbs.res.in)

Key Words

Pial Collateral Artery, Artery Endothelial Cell, Vascular Endothelial Growth Factor

Receptor-2, Intra-vital Imaging, Genetic Lineage Tracing

SUMMARY

Pial collaterals provide protection from ischemic damage and improve prognosis of

stroke patients. The origin or precise sequence of events underlying pial collateral

development is unclear and has prevented clinicians from adapting new

vascularization and regeneration therapies. We use genetic lineage tracing and intra-

vital imaging of mouse brains at cellular resolution to show that during

embryogenesis, pial collateral arteries develop from extension and anastomoses of

pre-existing artery tips, in a VegfR2 dependent manner. This process of artery tip

extension occurs on pre-determined microvascular tracks. Our data demonstrate that

an arterial receptor, Cxcr4, earlier shown to drive artery cell migration and coronary

collateral development, is dispensable for formation and maintenance of pial

collateral arteries. Our study reveals that collateral arteries of the brain are built by a

unique mechanism, distinct from that of the heart.

INTRODUCTION

Pial (leptomeningeal) collateral arteries are an extensive network of inter-artery anastomoses found in the pial layer of the brain. In humans, the extent of pial collateral circulation is a predictor of stroke outcome.[1,2] Stroke occurs when blood supply is obstructed to a part of the brain and is one of the leading causes of morbidity worldwide (World Health Organization).[3] Pial collaterals serve as an alternate route of blood flow and alleviate the symptoms of ischemic stroke by bypassing vascular occlusions. Thus, understanding mechanisms which drive the origin and remodelling of pial collateral arteries is of immense clinical relevance.

Results from preclinical studies indicate wide-spread variation in the pial collateral artery network among individuals and organs.[4] These variations are a result of differential expression and function of proteins associated with Vascular endothelial growth factor (Vegf) signalling axis.[5–7] Different mouse strains demonstrate differential Vegfa expression level which corelate with their pial collateral density.[8] Similarly, global overexpression of Vegf ligand increases number of native pial collaterals and improve collateral perfusion post-injury.[5,6] The receptor mediating the function of Vegf ligand during the formation of pial collaterals is Vegf receptor-2 (VegfR2).[6] A global knockdown of Vegf or VegfR2 during embryogenesis and in a time-frame when pial collaterals are expected to form, results in reduction in their number and diameter.[6] Thus, Vegf is a key driver of pial collateral formation and a critical factor which regulates variation in extent of pial vascular network in mice.

The vital role of Vegf in murine pial collateral development was discovered almost a decade ago.[5,6] However, it is still unclear how the various molecular players of Vegf pathway orchestrate the formation or remodelling of pial collateral network. In

mice, it is hypothesized that an *arteriole-specific sprouting angiogenesis like* process

leads to the formation of the foremost pial collaterals.[6] However, molecular evidence

supporting the identity of these sprouting cells (if arteries or capillaries) is missing.

Additionally, the cellular origin or fate of these sprouts is unknown. Finally, the

possibility of cooperation from other vascular processes, such as arterialization,

arteriogenesis or artery-reassembly, in building pial collaterals, needs to be

thoroughly investigated.

In this study, we use genetic lineage tracing in conjunction with live imaging of

pial vasculature of transgenic mouse embryos and adults to understand mechanisms

by which Vegf regulates pial collateral development and maintenance. Our data

indicate that the cellular source of pial collateral arteries is primarily pre-existing

artery endothelial cells (ECs) which build the first network of pial collaterals at

embryonic day (e)15.5 via VegfR2-dependent artery tip extension. The artery tips

extend on pre-determined microvascular tracks which then connect and build pial

collaterals. While arterial VegfR2 was essential for formation of pial collaterals *de*

novo, it was dispensable for its remodelling or maintenance through adulthood.

Arterial Cxcr4 (C-X-C chemokine receptor type-4), which is critical for coronary artery

formation in mice[9] was dispensable during pial collateral formation and its

maintenance. Together, our data indicate that artery endothelial cells build collateral

arteries by utilizing distinct vascular mechanisms in heart and brain.

RESULTS

Pial collateral arteries form during embryogenesis

We developed a whole brain imaging technique to reliably visualize all pial

arteries (Figures **S1A** and **S1B**). Due to lack of unique markers, accurately

identifying pial collateral arteries is exclusively dependent on their location. Pial

collateral arteries are artery segments which interconnect the ends of cerebral

arteries and can be seen in the watershed region (Figure **S1B**). To acquire this

spatial information, it was imperative to image the entire pial artery network. Our

technique allowed us to precisely identify pial collateral arteries. We specifically

focused on collateral arteries which connected and received blood flow from Anterior

Cerebral Artery (ACA) and Middle Cerebral Artery (MCA), as the coverage of this

arterial network is most extensive.

We first investigated the developmental stage when pial collaterals first

appeared. Mouse brains were harvested at different developmental time points and

whole brain immunostaining was performed for Connexin-40 (Cx40) protein, a known

marker for mature artery endothelial cell. Analyses of pial arterial network in both

hemispheres revealed the appearance of MCA at embryonic day e12.5 followed by

ACA at e13.5 (Figures **S1C**, **S2A** and **S2B**). At e14.5, both cerebral arteries were

present with no arterial connections between them (Figure **S2C**). Subsequently, the

first arterial anastomoses appeared at e15.5 (Figure **S2D**) which then remodelled to

expand in number through birth/postnatal day (P)0 (Figures **S1C** and **S2D**-**S2F**). This

method confirmed that both MCA and ACA are present at e14.5 (Figure **S2G**) and

the foremost pial collaterals appear at e15.5 (Figure **S2H**).

To confirm the above results, we next used a knock-in mouse line (*Cx40eGFP/+*)

where eGFP expression was driven by *Cx40* promoter activity. Brains from different

developmental stages confirmed our findings from Cx40 immunostaining. MCA

appeared at e12.5, ACA appeared at e14.5 and collateral arteries between MCA and

ACA appear at e15.5 (Figures **S3A**-**S3F** and **S1C**). eGFP+ cerebral arteries (MCA

and ACA) were formed by e14.5 (Figure **S3G**) and the first eGFP+ collateral artery

between MCA and ACA appeared at e15.5 (Figure **S3H**). Thus, using two

independent methods— Cx40 immunostaining and detecting its reporter activity, we

demonstrate that mouse pial collateral arteries form during development, and appear

precisely at e15.5 (Figure **S1C**).

Intra-vital imaging reveals arterial origin of pial collaterals by artery tip
extension

Despite several hypotheses, there is lack of compelling data in support of

vascular mechanisms which drive pial collateral development. Pial collaterals appear

at late embryonic stages. At this developmental stage, rapid vascular changes make

it difficult to distinguish collateral-specific events from overall vascular growth, thus

making it challenging to draw information from fixed brain samples. Additionally, late

embryonic stages cannot be cultured *ex vivo* due to extremely demanding metabolic

and physiological needs. Thus, we custom-developed an imaging paradigm to

perform intra-vital imaging of pial layer at late embryonic stages (e16-e17) (Figure

1A), a developmental stage where the brain is actively building pial collaterals.

Briefly, embryos of the desired developmental time points were dissected out (with

their intact yolk sac) from the anesthetized pregnant dams while still connected to

the mother via their umbilical cord. Embryos were maintained at 37˚C and

immobilized at appropriate orientation to aid imaging. The embryos were subjected

to stereoscopic or confocal imaging for several hours. This imaging set up allowed

visualization of the entire hemisphere— MCA, ACA, and connected collateral arteries

(Figure **1B**).

Pial collaterals appear only after the primary branches of MCA and ACA have

developed (Figure **S1C**). We hypothesized that the branches of these cerebral

arteries contribute towards formation of pial collaterals. To test this hypothesis,

Cx40CreER; Rosa26[TdTomato]; Tie2::eGFP::Claudin5 mice were used to label artery

ECs with TdTomato and all endothelial cells with eGFP. To activate TdTomato,

Tamoxifen was administered at e13.5, when both MCA and ACA are present and

would be labelled in 2 days, [9,10] by e15.5. Imaging of endothelial cells derived from

arteries was performed at e16-e17. This experimental strategy allowed visualization

of cellular dynamics specific to artery cells derived from pre-existing arteries (Figure

1C) and excluded the possibility of capturing other vascular processes.

We performed intra-vital time-lapse imaging of pial arteries and investigated if

pial collaterals originate from pre-existing artery endothelial cells. We focused on the

watershed region: an initial artery-free region which spans the distal ends of both

arterial trees (MCA and ACA) and is later occupied by pial collaterals (Figures **S1C**

and **1D-1F**). Stereoscopic images of e16.5 brains revealed that branches from MCA

and ACA extend (arrowheads, Figure **1E**) and anastomose to build collaterals in the

pial watershed (brackets, Figure **1E**). Additionally, TdTomato expression in artery

tips (orange arrow, Figure **1E**) and collaterals (green arrow, Figure **1E**) increased

over time. These changes were executed within a short span of 80 minutes, and

could not have been captured in fixed brain samples.

We next investigated artery cell dynamics associated with the growth fronts of

the two cerebral arteries (Figures **1F** and **1G**). At e16.5, the distal branches of the

developing MCA extended and anastomosed to build an artery network *de novo*

(arrowheads, Figure **1G**). This is the first evidence demonstrating that pre-existing

cerebral artery tips build pial collaterals. We then performed intra-vital time-lapse

confocal imaging of brains at e16, a stage prior to when we observed collateral

artery networks being built *de novo* (Figures **1E** and **1G**). Confocal imaging at e16

provided the necessary cellular resolution to quantitatively assess artery-tip

extensions from MCA and ACA branches (Figures **1H**-**1J**). As expected, we clearly

observed artery tip extension of MCA branch (Figures **1H** and **1I**) and the ACA

branch (Figure **1H**, **J**) within an hour. Additionally, we noted several artery-specific

movements within the pial layer. This included the appearance of single artery

endothelial cells (arrows, Figures **1K** and **1L**), movement of these single artery cells

towards an artery (Figures **1K**, **1M** and **S4**), and emergence of arterial sprouts

(arrowheads, Figures **1K** and **1L**). Together, for the first time, we provide direct

evidence for the vascular origin of pial collaterals— pre-existing MCA/ACA artery tips

extend and connect with each other to build pial collaterals *de novo*.

Artery tip extension occurs on pre-determined microvascular tracks

We next investigated if other endothelial cell sub-types participate in, and/or

facilitate, pial artery tip extensions during collateral development. Live imaging of pial

watershed showed that all artery cell movements, almost always, occurred along

microvascular (MV) tracks which appeared structurally more organized/tubular than

the surrounding capillaries (arrowheads, Figures **2A** and **2B**). These "microvascular

tracks" were distinct from the surrounding capillary plexus in that they formed direct

connections between two arteries, but were not necessarily entirely populated by

TdTomato$^+$ artery derived endothelial cells (Figure **2B**). We hypothesized that

development of MV tracks precedes artery tip extensions and tested two potential

mechanisms by which MV tracks could assist artery tip extensions during pial

collateral development (Figure **2C**). Firstly, complete formation of MV tracks between

two arteries at e16 could be a pre-requisite for initiation of artery tip extension, as

often observed at e16. (Figures **2Ci-iv**). Alternatively, MV tracks could appear first,

but continuously grow/remodel to allow artery tip extension (Figures **2Ci**, **v**, **vi**) and

subsequent pial collateral formation (Figure **2Civ**). If the former was true, the

number/length of MV tracks in a given watershed would remain unchanged between

e16 and e16.5, however the TdTomato coverage of these MV tracks would be higher

at e16.5 (Figures **2Cii-iii**). If the latter mechanism was true, at e16.5, the length of

MV tracks would be higher, and the TdTomato coverage of these MV tracks would

increase or remain unchanged (Figures **2Cv**, **vi**). To test these two potential

mechanisms, *Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5* mouse line was

used, where MV tracks can be clearly visualized with eGFP reporter (Figure **2B**).

Confocal imaging of e16 (Figure **2D**) and e16.5 (Figure **2I**) whole brains, at single

cell resolution allowed examination of watershed regions at both developmental

time-points (Figures **2E-2H** and **2K-2N**). Next, robust quantification of MV tracks for

each watershed region was performed (Figures **2H**, **2N** and **2J**). Amongst eGFP⁺

vessels, an MV track has a distinct morphology from the surrounding capillary plexus

and can be identified as eGFP⁺ tubular structures connecting the distal artery

branches (Figures **2F** and **2L**). We identified a eGFP⁺ vessel as an MV track if 1) it is

a continuous tubular structure and 2) is connected to a TdTomato⁺ artery at least at

one end. These MV tracks were traced on multiple z-slices, using Simple Neurite

Tracer (Figures **2H** and **2N**), and next, the extent of TdTomato coverage was

calculated for total length of MV tracks per watershed (Figures **2H**, **2N** and **2O**).

Quantifications show that while the total length of MV tracks was increased by 1.5

fold in e16.5 (Figures **2H**, **2N** and **2J**), the TdTomato coverage of these MV tracks

did not change between e16 and e16.5 brains (Figures **2H**, **2N** and **2O**). These

results support the second mechanism (Figures **2C-i→v→vi→iv**); thus, suggesting

that pial collateral development is accomplished by simultaneous growth of MV

tracks and artery-tips on them.

Artery ECs are the major cellular contributors of pial collaterals during

embryogenesis

We next assessed the vascular-lineage composition of pial collaterals at a

later stage when the collateral artery network has fully formed. This would allow us

to: 1) determine if the artery cell contribution to nascent pial collaterals observed at

e16.5 (Figure **1E**) persisted in later stages and 2) check contribution from

microvascular lineage which is evident in newly formed collaterals in heart[9,11] and

skeletal muscle.[12] Genetic lineage tracing with *Cx40CreER; Rosa26^{TdTomato}* and

ApjCreER; Rosa26^{TdTomato} mice were performed where TdTomato expression could

be conditionally induced, with Tamoxifen, specifically in artery ECs or capillaries

respectively. Tamoxifen was administered to timed pregnant dams at e14.5 when

both MCA and ACA arteries are present (Figures **3A**, **S2G** and **S3G**). Embryos were

analysed at e18.5 or P0 when the pial collateral artery network has completely

formed (Figures **3A**, **S2H** and **S3H**). Whole embryonic brains were immunostained

for Cx40 and segments of collateral arteries between MCA and ACA were analysed

for lineage contribution (Figures **3B**-**3J**).

We assessed the extent of lineage contribution from artery or capillary ECs

towards pial collaterals. As expected, all pre-existing artery ECs were labelled and

traced in *Cx40CreER; Rosa26^{TdTomato}* embryos (Figures **3C**, **3D**, **3G** and **3I**). Whole

brain imaging of the embryo (Figure **3C**) allowed identification of pial collateral

arteries present between MCA and ACA in the watershed (Figure **3D**). Individual

collateral segments as identified by Cx40 immunostaining were analysed for

presence of TdTomato+ artery ECs (Figures **3D**, **3G** and **3I**). This analysis revealed

that 77% of the Cx40+ collateral area to be TdTomato+ (Figure **3B**). In contrast to this

222 data, a similar analysis in an independent experiment with *ApjCreER; Rosa26^TdTomato*

223 mice (Figures **3E**, **3F**, **3H** and **3J**) showed only 31.6% of the collateral area to be

224 TdTomato⁺, i.e. traced from capillaries (Figure **3B**). Thus, artery ECs are the major

225 contributors of pial collateral arteries during development.

227 Arterial VegfR2 mediates artery tip extension and is essential for pial collateral

228 artery development

We analysed similarities, if any, in the molecular programs used by artery

endothelial cells during collateral development in the heart and brain. Post-injury,

neonatal coronary artery ECs undergo extensive migration and proliferation to build

new coronary collateral arteries in the watershed. The processes of artery cell

migration and proliferation are heavily dependent on Cxcr4[9] and VegfR2[13]

respectively. To assess the functions of Cxcr4 and VegfR2 in pial collateral

development, we deleted the respective genes at e14.5 and assessed the extent of

pial collateral network at e18.5/P0 by whole brain imaging (Figure **4A**). Surprisingly,

deletion of arterial *Cxcr4* did not affect the pial collateral network at e18.5 (Figure **4B**,

control=8, artery specific knockout for *Cxcr4*=13) or in adulthood (Figure **S5**, N=5 per

genotype). This data suggests that a Cxcr4-independent vascular mechanism

facilitates pial collateral development and its maintenance.

We next assessed if pial collateral artery development is dependent on

arterial VegfR2. Whole brain imaging of control and *VegfR2* artery specific knockout

embryos showed differences in collateral connectivity in the watershed, and required

systematic quantification (Figures **4C-4F**). The extensive connectivity of pial arteries

across the hemisphere (including watershed) provide several possible collateral

paths for the blood to flow. These artery-artery connections are innumerable. We

developed a method that allowed us to compare the pial artery coverage between

the control and *VegfR2* artery EC specific knockout brains quantitatively (Figure **4C**).

Intensity line-profiles were generated across the width of the hemisphere (Figures

4C i, **ii**). The identified peaks per line-profile was plotted across the width of the

hemisphere (Figure **4C iii**). The data was fit to a Gaussian function and the fitted

curves were used as representative of the arterial coverage per hemisphere in each

mouse (Figures **4C iv**). The average of the fitted curves represented one genotype

(grey curve, Figure **4C iv**). Quantification of pial artery coverage showed significantly

different distributions between control and *VegfR2* knockout brains (KS test, D

value=0.569, p value=2.2066 X 10^{-141}) (Figure **4E**). Peak arterial coverage was

observed at the centre of the hemisphere and watershed was defined as a region ±

500μm from the centre (Figure **4E**). The loss of arterial coverage in the watershed

was evident in the knockout (Figure **4E**) and was quantified as area under the curve

(Figures **4E** and **4F**). This area under the curve represented the extent of collateral

artery network in the watershed and was significantly lower in brains without arterial

VegfR2 (Figure **4F**, unpaired t-test, p value= 0.0001). Thus, depletion of *VegfR2* from

arteries reduced the extent of pial collaterals formed during embryogenesis.

We next investigated the possible mechanisms by which arterial VegfR2

facilitated pial collateral formation. *VegfR2* was deleted from artery ECs at e14.5,

and analyses of pial collateral network was performed at e16.5 (Figure **4G**), a

timepoint when collaterals are beginning to form via artery tip extensions (Figures

1D-1G)**.** Quantification showed fewer TdTomato$^+$ collateral arteries (i.e. TdTomato

contiguous or Td^{con}) without arterial *VegfR2* (Figure **4H**). This indicated that initiation

of collateral development at an earlier timepoint, e16.5, also depends on arterial

VegfR2. This VegfR2-dependent collateral phenotype (Figure **4H**) was independent

of artery cell proliferation which was minimal at this stage, for both control and

VegfR2-deleted brains (data not shown). Close analyses of collaterals in control

brains showed that TdTomato$^+$ artery cells occupied a major length of microvascular

tracks connecting branches of MCA and ACA (Figure **4I**). This observation was in

contrast to *VegfR2*-depleted brains where TdTomato$^+$ artery cells sparsely populated

the microvascular tracks (Figure **4J**). Thus, arterial VegfR2 regulates pial collateral

formation by promoting artery growth along microvascular tracks.

We investigated if endothelial cells of capillary lineage populate pial collaterals

via Dach1; a transcriptional factor that drives arterialization of capillaries into

coronary arteries during cardiac development in mice.[14] As observed in developing

mouse hearts,[14,15] expression of Dach1 was higher in capillaries as compared to

arteries (Figure **S6A**). *Dach1* was deleted from capillaries at e14.5 and the extent of

pial collateral network was assessed at e18.5 (Figure **S6B**). Interestingly, no

significant difference was observed in the number of Cx40$^+$ collaterals in the *Dach1*

capillary-EC specific knockout brains (Figure **S6C**). Thus, in contrast to heart, the

formation of pial collaterals during embryogenesis is independent of capillary Dach1

activity.

Pial collaterals do not remodel in adulthood

We asked the question if arterial VegfR2 could modulate neonatal remodelling

events. To address this, arterial VegfR2 was deleted in arteries at P0, and whole

brain imaging was performed at P2 and P7 (Figure **S7A**). The arterial deletion of

VegfR2 was robust at P7 (Figures **S7B** and **S7C**). Analyses of whole P2 (Figures

S7D-**S7G**) and P7 (Figures **S7H**-**S7K**) hemispheres revealed no significant

differences in arterial coverage with depletion of arterial VegfR2 (Figures **S7G, K**).

Thus, arterial VegfR2 is dispensable for neonatal pial artery and collateral artery

remodelling.

We next investigated the cellular dynamics associated with fully formed and

mature pial collateral arteries between MCA and ACA in adult mice. For this, we

utilized an extensive longitudinal in vivo imaging paradigm that allowed us to get

single cell resolution of vascular structures. 3-5 months old *Cx40CreER;*

*Rosa26^TdTomato^*; *Tie2::eGFP::Claudin5* mice underwent craniotomy and an optical

window of 4mm was implanted on one of the hemispheres between the cranial

landmarks—bregma and lambda (Figures **5A** and **5B**). In these transgenic mice, pre-

existing pial arteries were labelled with TdTomato upon Tamoxifen administration

(Figure **5C**), and constitutively expressing GFP marked all endothelial cells. 5

consecutive doses of Tamoxifen resulted in 95.4% and 86.8% recombination in

proximal and distal arteries respectively, but a lower 65.3% recombination in the pial

collateral regions (Figure **5D**). A lower recombination efficiency in the pial collateral

arteries resulted in a patchy reporter pattern comprising of a mixture of TdTomato$^+$

and TdTomato$^-$ cells (Figure **5D**). Like in developmental stages (Figure **3B**), artery

ECs populate adult pial collaterals to a greater extent as compared to other vascular

lineages. This ultimately allowed us to most accurately track the position of individual

cells within a given collateral artery.

An optical window allows longitudinal imaging. Imaging the collateral zone

(watershed between MCA and ACA) for a month (Figure **5E**), resulted in high quality

images with single cell resolution (Figures **5E-5H**). Morphology of the pial vessels

alongside reporter activity in endothelial cells were instrumental in identifying pial

arteries, veins and capillaries (Figures **5E** and **5I**). High resolution images of artery

endothelial cells (both TdTomato$^+$ GFP$^+$ and TdTomato$^-$ GFP$^+$) within a collateral

artery segment across many collaterals and optical windows were analysed (Figures

5J and **5K**). Data showed that artery endothelial cells do not remodel or reposition

themselves within a given collateral artery segment (Figures **5J**-**5M**). Cellular

matrices representative of the relative position of collateral artery cells suggested

that under physiological conditions, every artery cell within a mature/ adult pial

collateral maintains its contact with the same neighbouring cells (Figures **5L** and

5M). Thus, surprisingly, the cellular organization within pial collaterals and the overall

collateral structure was extremely stable. For the first time our data, with cellular

resolution, demonstrates the static nature of artery endothelial cells during

homeostasis.

Arterial VegfR2 is dispensable for pial collateral maintenance

We next asked if arterial VegfR2 is necessary for the maintenance of adult

pial collaterals using adult mice deficient in arterial *VegfR2*. Using the same in vivo

imaging paradigm mentioned above, pial collateral segments in artery endothelial

cell specific *VegfR2* knockout mice were imaged at cellular resolution and followed

for a month with imaging time-points separated by 2 weeks (Figure **6A**). Longitudinal

imaging through an optical window (Figure **6B**) yielded a map of vascular network

which could be followed for months (Figures **6C** and **6D**). Of note, our qualitative

observation suggests that *VegfR2* depleted pial collaterals had smaller diameters.

High magnification images of watershed between MCA and ACA (Figures **6E**-**6G**)

allowed us to analyse individual *VegfR2*[-/-] artery endothelial cells (TdTomato[+])

residing on adult collateral arteries (Figures **6H**-**6J**). Long term imaging of the same

collateral regions across many collaterals and mice showed no significant changes in

cellular positions relative to their neighbouring cells (Figures **6K**-**6M** and **S8**). Thus,

arterial VegfR2 is dispensable for maintenance of mature pial collateral arteries.

Together, in this study, we highlight the essential role of arterial VegfR2 in promoting

artery-tip extension and pial collateral development during embryogenesis (Figure

6N); a function dispensable in adulthood.

DISCUSSION

In the present study, we investigate artery cell dynamics during collateral artery

formation and maintenance in mouse brain. Firstly, using an artery-specific

molecular marker, we dissect out where and when pial collaterals appear during

embryogenesis. Secondly, we develop a live imaging method in mouse embryos to

provide conclusive evidence that pre-existing arteries grow on pre-determined

microvascular tracks which then connect to build pial collaterals *de novo.* Thirdly, by

combining mouse genetics with whole brain confocal imaging at cellular resolution,

we show that an angiogenic vascular receptor VegfR2 facilitates artery-tip extension

on microvascular tracks and aids origin of pial collaterals. Finally, we show that

Cxcr4, a receptor essential for Artery Reassembly and formation of coronary

collateral arteries *de novo*, is dispensable for development of pial collaterals. Thus,

two fundamentally distinct mechanisms contribute to the origin of coronary and pial

collaterals.

Organ-specific mechanisms build collaterals

For the first time, we provide direct proof that a vascular mechanism driven

specifically by artery endothelial cells leads to the origin of pial collaterals. Like in the

heart, this artery specific process is regulated by arterial VegfR2, but is distinct from

Artery Reassembly where pre-existing artery cells dissociate from resident

coronaries, migrate, de-differentiate/proliferate and coalesce to build coronary

collaterals.[9,13] Instead, developing pial arteries extend along microvascular tracks

and connect with other artery tips to build new collateral artery paths. Furthermore, in

heart, coronary collateral formation is regulated by arterial Cxcr4 which drives artery

cell migration during Artery Reassembly. In contrast, arterial Cxcr4 had no impact on

pial collateral formation or its maintenance. Unlike heart, pial artery endothelial cells

contribute minimally via proliferation. Evidence for negligible artery cell proliferation

prompts us to hypothesize that artery endothelial cells at the distal ends experience

a forward push from the proximal parts of the artery. This forward pressure could be

generated due to incorporation of new cells or expansion of existing cells at the

proximal artery segments.

The need for functional collaterals is most during an ischemic condition such

as an artery occlusion, and can significantly impact tissue function. The differences

in factors which regulate origin of coronary and pial collaterals, are potentially driven

by the metabolic needs of these tissues. While cardiomyocytes can survive hypoxia

for few hours,[16] neurons are more susceptible to low levels of oxygen.[17] Higher

hypoxia sensitivity of brain tissue may lead to formation of pial collaterals during

embryogenesis as a protective measure, while coronary collaterals form only upon

induction of an arterial occlusion. Furthermore, collateral arteries in skeletal

muscle,[12,18] intestine,[19] and hind limb[20] can also undergo arteriogenesis.[21] Identifying

collateral-building vascular mechanisms across organs could reveal clinically

relevant information and open avenues for therapeutics.

Arteries and capillaries collaborate to build collaterals

We demonstrate the formation of collateral arteries, in real time, from pre-

existing arterial lineage as observed[9,10,13,22] or proposed[6] earlier. While this

manuscript was in progress, Perovic et al reported that two separate vascular

lineages populate pial collaterals.[10] We also observed that well-formed pial

collaterals are majorly populated by artery cells, and minimally by capillaries. Intra-

vital imaging of embryonic brains reveal that pre-existing artery cells extend along

defined microvascular paths to connect and build pial collaterals. These artery

specific events are executed within a short span, are difficult to capture using static

images and can be observed only by live imaging.

We cannot exclude the contribution of other artery-driven mechanisms. We

observed single artery cells migrating towards pre-existing arteries; coalescence of

these single cells with arteries could lead to arterial growth. Alternatively, the

migration and incorporation of single artery cells at the proximal ends of the artery

could indirectly contribute towards distal elongation of arteries. The dynamics of

single artery cell movements could also establish the branching pattern of the pial

artery network; the precise mechanisms for which are unknown.

We show that pre-existing artery tip extensions occur along a distinct

supporting structure. In the heart, single artery endothelial cells migrate by hitch-

hiking on to the capillary networks during the formation of new collaterals. On the

contrary, our study in the embryonic brains reveal the presence of a unique (Tie2+)

vascular segment on which the arteries extend. These microvascular tracks are

distinct from the surrounding capillary mesh in that they anastomose ACA and MCA

arterial branches but do not show any artery specification. We speculate that the

microvascular tracks form during the specific time window (e16-e17) which coincides

with the period when majority of collaterals appear. Thus, these tracks may play a

crucial role in pial collateral formation. We hypothesize that the cellular origin of

these microvascular tracks is of capillary in nature. Elucidating the underlying

mechanism governing their formation and their molecular identity is an exciting

avenue for future investigation.

Our data about pre-defined microvascular tracks guiding the growth of pial

arteries suggest that the microvasculature could shape the artery network of the

brain. However, the microvascular contribution towards pial collaterals via

arterialization is less likely, as deletion of *Dach1*, a regulator of arterialization,[14,23]

had minimal effect on pial collateral network during embryogenesis. We propose that

as artery tips extend and get connected in the watershed, microvascular cells are

"trapped" within pial collaterals. This could generate a mixed lineage distribution[10]

within pial collaterals—both in embryos and adults. Together, multiple mechanisms

could be actively building pial vasculature during embryogenesis. That being said,

possible contribution from endothelial progenitor cells or peri-vascular cell types

(pericytes, smooth muscle cells) towards pial collateral formation cannot be entirely

excluded.

Distinct mechanism for collateral maintenance

Given that collateral arteries have a unique hemodynamic niche,[24] it would be

interesting to investigate which molecular programs maintain or remodel pial

collaterals in an adult brain. During embryogenesis, pial collaterals are built *de novo*,

in an artery-free zone and remodel as and when blood flows through them. In

contrast, mature collaterals remodel in response to differential blood flow patterns

and oxygen demands of their surrounding tissue. Postnatal remodelling of pial

collaterals is dependent on recruitment of smooth muscle cells and exhibits tortuous

morphology.[24] These fundamental differences could explain the requirement for

distinct molecular programs for building new collaterals and maintaining old ones.

This is evident in our study; VegfR2 controlled pial artery network during

embryogenesis but was dispensable in adulthood. That being said, arterial VegfR2

may regulate other aspects of vascular changes those require activation of

angiogenic pathways[25] such as remodelling induced by hypoxia, ischemia or aging.

We did not notice any remodelling events within adult pial collaterals, in presence or

absence of arterial Cxcr4. Together, adult pial collaterals are maintained via

mechanisms independent of two key angiogenic receptors in arteries—VegfR2 and

Cxcr4. It will be interesting to test if adult collateral maintenance is independent of

signals originating from an arterial niche. It is possible that surrounding

microvasculature plays a more prominent role in maintenance of pial arteries and

collaterals in adults. Investigating the molecular identity of the microvasculature

surrounding pial collaterals could reveal new vascular mechanisms key to collateral

maintenance.

MATERIALS AND METHODS

Mouse lines

Mouse lines	Source/ References
Rosa26^{TdTomato}	The Jackson Laboratory, stock number 007909
Cx40CreER	Miquerol et al. 2015[26]
ApjCreER	Chen et al. 2014[15]
VegfR2 flox	The Jackson Laboratory, stock number 018977
Cx40$^{eGFP/+}$	Miquerol et al. 2004[27]
Tie2::eGFP::Claudin5	Knowland et al. 2014[28], Lutz et al. 2017[29]
Cxcr4 flox	The Jackson Laboratory, stock number 008767
Dach1 flox	Chang et al. 2017[14]

All experiments were performed on mice with mixed strain background. All mice

were housed and bred at the National Centre for Biological Sciences animal facility

in accordance with the Institutional Animal Ethics Committee (IAEC). All procedures

were performed with approval from IAEC. Data shown in this study is compiled from

age-matched males and females from several litters.

Genotyping primer information

Primer (F, forward; R, Reverse)	Sequence (5' → 3')
Cre F	GCATTACCGGTCGATGCAACGAGTGATGAG
Cre R	GAGTGAACGAACCTGGAAATCAGTGCG
TdTomato mutant F	CTGTTCCTGTACGGCATGG
TdTomato mutant R	GGCATTAAAGCAGCGTATCC
TdTomato wild-type F	AAGGGAGCTGCAGTGGAGTA
TdTomato wild-type R	CCGAAAATCTGTGGGAAGTC
VegfR2 flox F	CCACAGAACAACTCAGGGCTA
VegfR2 flox R	GGGAGCAAAGTCTCTGGAAA
GFP F	GCACGACTTCTTCAAGTCCGCCATGCC
GFP R	GCGGATCTTGAAGTTCACCTTGATGCC
Cxcr4 flox F	CCACCCAGGACAGTGTGACTCTAA
Cxcr4 flox R	GATGGGATTTCTGTATGAGGATTAGC
Dach1 flox F	CTCCTGAAGATGAGGAGCTCACCCC
Dach1 flox R	AACAATTCTTGTCCTTCACGTGCCC

Visualization of pial arteries in fixed samples

Harvesting whole brains/heads from mouse embryos

Depending on the developmental stages and size of the embryos, pial collateral arteries were visualized either in whole brains (e18.5 or P0) or using whole heads (e12.5- e16.5).

e12.5-e15.5: To perform the whole head explant at e12.5-e15.5 stages, the pregnant females were euthanized through carbon dioxide overdose and a Y-shape incision was made on the abdomen. Intact embryonic sacs were dissected out. Small punctures were made on the embryo sac with the help of microforceps. The whole embryo sac was fixed in 4% Paraformaldehyde (PFA) for 2 hours with gentle shaking at 4°C. Following fixation, the embryo sac was washed at 4°C with gentle shaking. This washing step was performed with 1X PBS and repeated 4 times every 15-20 mins. The embryos were then extracted from the embryo sacs and the heads were removed with the help of microforceps.

e16-e16.5: The embryonic explants at e16/e16.5 are performed similarly. However, for *Tie2::eGFP::Claudin5* genotype, the loose skin on top of the brain was removed to aid immunostaining. Whole heads were imaged within 4 hrs, prior to PFA fixation to avoid loss of GFP fluorescence signal.

e18.5: To perform the whole brain explant at e18.5 stage, the timed pregnant females were euthanized as above. The skull of each embryo was gently removed using the microforceps. The embryonic brains were scooped out with the help of a microspatula and fixed in 4% PFA at 4°C for 2.5 hours. Agitation was avoided at this

stage to preserve tissue integrity of embryonic brains. The fixed embryonic brains

were then washed with 1X PBS three times at 15 minutes intervals with gentle

shaking at 4°C.

P0: Neonatal explants at P0 were performed by euthanizing the pups through carbon

dioxide overdose followed by decapitation. The skull was removed using scissors

and microforceps. The brains were scooped out using the microspatula. The brains

were then fixed with 4% PFA at 4°C for 2 hrs followed by three washes with 1X PBS

at an interval of 15 mins.

Whole brain/head immunostaining of mouse embryos

e12.5-e15.5: The embryonic heads (e12.5 - e15.5) were incubated in 0.1% PBST

(0.1% TritonX-100 in 1X PBS) for 2.5 hours to permeabilize the tissue, followed by

incubation with primary antibodies overnight at 4°C with gentle shaking. Heads were

then washed with 1X PBS, changing the buffer every 1-2 hours for 1 day. This step

was followed by incubation with secondary antibodies made in 0.1% PBST. Samples

were then washed every 1-2 hours for 1 day with 1X PBS. Following immunostaining

the brains are immersed in an anti-fade mounting media (from Jackson

ImmunoResearch Laboratories Inc.) and imaged immediately.

The embryonic and neonatal brains were incubated with primary antibody (made in

1X PBS) overnight, at 4°C with gentle shaking. Samples were then washed with 1X

PBS, changing the buffer every 1-2 hours for 1 day and incubated with Alexa Fluor

conjugated secondary antibodies (made in 1X PBS). Finally, samples were washed

every 1-2 hours for 1 day with 1X PBS and stored in mounting media or imaged

immediately.

Antibodies used are: Cx40 (Catalog number: Cx40-A, Alpha Diagnostics, 1:500

dilution), Dach1 (Catalog number: 10194-1-AP, Proteintech, 1:500 dilution), VegfR2

(Catalog number: AF644, R&D Systems, 1:250). All secondary antibodies were

procured from Invitrogen and used at a dilution of 1:500.

Imaging whole brains/heads of mouse embryos

The mice heads and brains from different developmental time-points were imaged

using Olympus FV-3000 upright or inverted confocal microscope by placing the

sample in a 2-well chamber (Catalog number: 154461; Nunc™ Lab-Tek™ II

Chamber Slide™ System; Thermofisher Scientific). Whole brains and heads were

scanned using 4X and 10X air objectives, and multiple stacks were captured. The

captured image stacks were processed in ImageJ and stitched together using Adobe

Photoshop software.

The embryonic heads at e16/e16.5 were imaged using Olympus FV-3000 5L Upright

microscope. The pre-fixed embryonic heads were attached to a 35mm plastic petri

dish using agarose gel prior to imaging. The whole head imaging was performed

using 4X air objective and watershed images were captured with 10X air objective or

20X water dipping/immersion objective and 1.5X optical zoom with brains immersed

in 1X PBS.

Visualization of pial arteries in live mice

In vivo imaging of embryonic pial arteries

Immobilization of live embryo: *Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5*
pregnant dams with embryos at developmental stages between embryonic day
(e)16-e17 were anesthetized with a subcutaneous injection of Ketamine-Xylazine
(Ketamine (80 mg/kg); Xylazine (5 mg/kg)) cocktail. The dams were then placed in a
supine position on a regulated heating pad (TC1000, CWE) that was positioned on
top of a 3D-printed baseplate (as shown in Figure **1A**). A small incision was placed
along the midline of the abdomen. With a gentle retraction of the abdominal skin and
wall, a single uterine horn within the abdominal cavity was isolated. The uterus was
surgically opened and a single embryo along with the yolk sac was isolated and
placed into a 35mm petri dish. The embryo still connected to the dam was
immobilized by submerging the bottom part with 1% agarose, as shown in Figure **1A**.
The immobilized embryo is then submerged in dissection media (DMEM/F12 + 10%
FBS, Catalog number: 21041025, Thermo Fisher Scientific), and the media is
maintained at 37°C using a custom-made Arduino controlled media heating and
circulating device.

Live embryonic imaging: The setup along with anesthetized dam was placed under
Macro Zoom microscope MVX10-Olympus for large scale imaging and under
FV3000 upright confocal Microscope-Olympus for high resolution imaging using 20X
water dipping objective. In Macro Zoom Microscopy, the entire observable depth was
scanned whilst obtaining videos of 1 minute spanning a duration >2 hours in a time-
lapse manner. In confocal microscopy, the entire observable depth was scanned
with a step size of 1 micron. A total of 4 embryos were imaged under the Macro

zoom microscope and 5 embryos were imaged using confocal microscope. Out of

these embryos, the artery cell dynamics was captured in 2 brains from each

category.

Image processing: Image stacks obtained from Macro zoom microscopy was

registered using ImageJ plugin (*StackReg*) and the registered stack was de-

convolved using ImageJ plugin (*DeconvolutionLab2*) and finally maximum intensity

projection was obtained as shown in Figure **1D**. The image stacks obtained from the

confocal microscopy were processed similarly except the deconvolution step.

The MCA and ACA growth fronts were automatically traced and the length of the

traced path was measured using Simple Neurite Tracer (SNT v4.1.13). The distance

of single cells from an artery was measured manually using ImageJ.

In vivo imaging adult pial arteries

Craniotomy: *Cx40CreER; Rosa26^{TdTomato}*; *Tie2::eGFP::Claudin5 and*

VegfR2$^{flox/flox}$;Cx40CreER; Rosa26^{TdTomato};Tie2::eGFP::Claudin5 adult mice (3-5

months) underwent craniotomy to obtain optical access to the pial layer of the brain.

Mice were anesthetized under 2% Isoflurane (Abbott, North Chicago, IL, USA) and

were placed on a stereotaxic mount (Item number 51547; Stoelting, Wood Dale, IL,

USA) with the skull held tightly using ear bars. The mice were kept anesthetized with

2% Isoflurane using a nose-cone. 5mg/kg Dexamethasone in 1X PBS was

administered subcutaneously and the mice kept warm using a heating pad (TC1000,

CWE). The eyes were covered with an eye ointment and the hair above the skull

removed using a hair trimmer. The surgical area was thoroughly wiped clean with

70% ethanol. Sterile surgical tools were used for the surgery. A small incision was

made using a scalpel blade and the exposed skin was removed. The connective

tissue covering the skull in the exposed area was scraped off using scalpel blade

and Q-tips. Using a high-speed electrical drill (burr of diameter ~0.25mm), a window

of ~4mm diameter was carefully etched on the skull. The skull within the window was

eventually detached from the rest of the skull after repeated drilling and was carefully

lifted using a pair of fine forceps (Catalog number: 11255-20, Dumont #55) without

damaging the dura. During the process of drilling, the skull was constantly irrigated

with cortex buffer (Concentration; in mM: NaCl–125, KCl–5, glucose–10, HEPES–10,

$CaCl_2$–2 and $MgCl_2$–2. pH set to 7.35).[30] The exposed dura was aspirated with

cortex buffer until there is no bleeding and the window looked clear. An optical plug

made up of two 4 mm coverslip and one 5 mm coverslip (CS-5R & CS-4R; Warner

Instruments, Hamden, CT, USA) glued together (with NOA 68, Norland Optical

Adhesive) was placed on the exposed dura. The optical plug was gently pressed

using a toothpick such that the 5mm cover slip was touching the skull bone and then

cyano-acrylic glue was applied across the perimeter of the 5mm coverslip attaching it

to the skull. At this stage we had an optical window of 4mm. An aluminium head bar

with circular hole of 8mm was placed onto the skull aligned to the previously made

optical window. Next, dental cement (Catalog number 56877, 3M Relyx U200) was

applied over the exposed skull using a sterile toothpick while being careful to not get

it on the optical window. Once the dental cement was completely dry, the animal was

placed on a warm pad in a clean recovery cage. Animals undergoing craniotomy

were maintained on meloxicam (2 mg/kg) and enrofloxacin (5 mg/Kg body weight)

for 3-4 days until full recovery. Mice were further allowed to recover for at least 2

weeks post-surgery before the start of any experiment. Mice with an optical window

can be imaged longitudinally for 2-3 months until the optical window starts to lose its

clarity.

<u>Live Adult Imaging</u>: To perform in vivo imaging, mice were anesthetized through

intraperitoneal injection of Ketamine (80mg/Kg body weight) – Xylazine (5mg/Kg

body weight) cocktail. The eyes were covered with an eye ointment and the mice

were placed onto a custom-made baseplate and immobilized using the implanted

metal head plate to this baseplate. A heating pad (37°C) was placed under the

mouse to maintain the body temperature. Subsequently, the baseplate was placed

on a motorized XY-table of an upright confocal microscope (FV3000 upright confocal

Microscope-Olympus). The entire optical window was imaged using 10X air objective

(0.4 NA) and selected pial regions with collateral arteries were imaged with 20X

water dipping objective (0.5 NA). A total of N=3 adult mice per group (control and

VegfR2 artery endothelial specific knockout) were imaged longitudinally for a period

of 1 month.

<u>Image processing</u>: The 10X image stacks obtained from the confocal microscopy of

the cranial window were registered using ImageJ plugin (*StackReg*) and maximum

intensity projection was obtained. The region of interest from the 20X image stacks

were cropped out slice by slice and stitched in Adobe Photoshop to avoid the

overlaying microvascular network in order to focus on the pial collaterals.

Quantifications

Lineage contribution towards pial collaterals during embryogenesis

Cx40 immunostained pial collateral segments from either *Cx40CreER;*

Rosa26^Tdtomato or *ApjCreER; Rosa26^TdTomato* brains were imaged at 10X with 2.12

optic zoom. Regions of interest were manually drawn to mark the collateral artery

segments. Lineage contribution was determined by quantifying TdTomato$^+$ coverage

on the collateral segment. This was performed by thresholding the images with Otsu

algorithm in ImageJ. The area fraction of the TdTomato$^+$ area per collateral segment

was calculated using the Measure tool in ImageJ. Contribution of the artery and

capillary derived ECs in collaterals was quantified by calculating the percentage of

collateral area covered by TdTomato$^+$ signal. 45 collateral segments (of an

approximate length of 600-800μm) from 7 *Cx40CreER*-lineage traced brains and 57

collateral segments from 8 *ApjCreER*-lineage traced brains were analysed.

Number of collateral arteries per hemisphere

In *Cx40CreER*+ (TdTomato) labelled brains, the collateral number was quantified

based on the number of contiguous TdTomato$^+$ vessels (Td-contiguous) connecting

MCA and ACA. Data shown in Figure **4H** was summarized from 2 control and 3

artery-specific *VegfR2* knockout. Data shown in Figure **S2H** was summarized from

e13.5 (N=5), e14.5 (N=5), e15.5 (N=5), e18.5 (N=5) and P0 (N=5) embryos and

neonates. Data shown in Figure **S3H** was summarized from e13.5 (N=3), e14.5

(N=3), e15.5 (N=2), e18.5 (N=2) and P0 (N=5) embryos and neonates. Data shown

in Figure **S6C** is quantified by counting number of Cx40 immunostained collateral

arteries between MCA and ACA, and is summarized from 4 control and 7 *Dach1*

capillary specific knockouts. An unpaired t-test was done to test significance

statistically.

Efficiency of TdTomato recombination

Segments of collateral arteries, proximal arteries (PA) and distal arteries (DA) from

the image stacks taken with 20X objective were obtained from live adult imaging of

Cx40CreER;*Rosa26*^{TdTomato};*Tie2::eGFP::Claudin5* mice. TdTomato$^+$ and TdTomato$^-$

cells were counted in each artery segment. Each segment was of an approximate

length of 600-800µm.The recombination efficiency was measured as ratio of

TdTomato$^+$ cells and total number of cells in the segment. A total of 10 collateral

segments, 10 PA segments and 10 DA segments from 4 mice were analysed to

obtain the recombination efficiency of TdTomato. One-way ANOVA was done to test

significance statistically (mean comparison Tukey test, PA vs. collateral, $p<0.0001$;

DA vs. collateral, $p<0.0001$; PA vs. DA, $p=0.08$).

Pial artery coverage in whole brains and in the collateral zone

The image stacks were obtained using confocal imaging of whole embryonic brains

with 10X objective, stitched using ImageJ plugin (*Stitching*) and max intensity

projection (MIP) was obtained. The MIPs were used for further analysis wherein one

complete hemisphere was cropped out and aligned vertically to the major axis of its

own elliptical boundary. Using a custom ImageJ macro, a line of the length of the

major axis was scanned across the width of the image and intensity profile for every

pixel along the horizontal axis was generated. Total number of local maxima (peaks)

for each intensity profile was obtained using the built-in ImageJ macro function

Array.findMaxima (array, tolerance). The tolerance used was 2 times the average

background intensity. The number of peaks thus obtained was plotted across the

width and the plot was fitted to a Gaussian function using OriginPro 2023b. The fitted

curve was used as a representative of the arterial coverage per hemisphere. The fit

curves obtained from various mice were aligned based on the Gaussian peaks and

averaged to get the representative curves for various experimental groups. We

considered a region of 1mm (-0.5mm to +0.5 mm) across the Gaussian peak as the

collateral zone (watershed). The area under the curve within this region was used as

a measure for the extent of the collateral network. 9 hemispheres from 9 embryos

(e18.5) were analysed per group (control and artery-specific knockout for *VegfR2*).

To test the statistical significance of the differences observed in pial artery coverage

(Figure **4E**), a Kolmogorov-Smirnov (KS) test was done. D value= 0.569, p value=

2.2066 X 10^{-141}. To test the statistical significance of the area under the curve (pial

collateral artery network shown in Figure **4F**), an unpaired t-test was done

(p=0.0001).

Similar analyses were performed in Figure **S7** with N= 3 per group. Unpaired t-test

was performed.

Microvascular tracks and their coverage with TdTomato⁺ arteries per

watershed

To analyse the Microvascular (MV) tracks, the watershed regions of the e16 (9

watershed from n=9 embryos) and e16.5 (7 watershed from n=5 embryos) brains

were imaged using 10X air objective at 1.5X optical zoom. The region of interest

(ROI) of comparable width (600-800 microns) within the watershed was taken for

further analyses. The eGFP⁺ MV tracks were traced using Simple Neurite Tracer

(SNT v4.1.13).The total MV length and area of the ROI were measured using SNT

and ImageJ respectively, to calculate the total MV length per unit area of the

watershed. Further, to measure TdTomato coverage of the MV tracks, the

TdTomato⁻ vessel segments on the MV track were traced and their lengths measured

using SNT to quantify the fraction of TdTomato coverage. Both, the MV tracks and

TdTomato coverage were quantified by going through multiple z-slices. An unpaired

t-test was done to test significance statistically (p=0.4085 for %TdTomato coverage

in the MV tracks and p=0.009 for normalized total MV length per watershed).

AUTHORS CONTRIBUTION

715 SD conceived the idea and designed the experiments. SK performed intravital

716 imaging of embryos and adults. NS, SG, VS, MD, SK performed embryonic

717 experiments. SD, SK, NS, SG wrote the manuscript. All authors contributed to

718 editing the manuscript.

ACKNOWLEDGEMENTS

720 We thank the following facilities for their technical support: Animal Care and

721 Resource Center at NCBS, Mouse Genome Engineering Facility at NCBS, Central

722 Imaging and Flow Cytometry Facility at NCBS. We sincerely thank Dr. Dritan Agalliu

723 (Columbia University) and Dr. Kristy Red-Horse (Stanford University) for generously

724 sharing *Tie2::eGFP::Claudin5* and *Dach1$^{fl/fl}$* mice respectively.

FUNDING

726 This work is supported by NCBS/TIFR core funding and DBT/Wellcome Trust India

727 Alliance Intermediate Fellowship to SD (IA/I/20/2/505205). Suraj K is supported by

728 DBT/Wellcome Trust India Alliance Intermediate Fellowship to SD. NS and VS were

729 supported by DST INSPIRE Scholarship for Higher Education (SHE).

DECLARATION OF INTERESTS

731 None

REFERENCES

1. Lima, F.O., Furie, K.L., Silva, G.S., Lev, M.H., Camargo, E.C.S., Singhal, A.B., Harris, G.J., Halpern, E.F., Koroshetz, W.J., Smith, W.S., et al. (2010). The pattern of leptomeningeal collaterals on CT angiography is a strong predictor of long-term functional outcome in stroke patients with large vessel intracranial occlusion. Stroke *41*, 2316–2322. 10.1161/STROKEAHA.110.592303.

2. Maas, M.B., Lev, M.H., Ay, H., Singhal, A.B., Greer, D.M., Smith, W.S., Harris, G.J., Halpern, E., Kemmling, A., Koroshetz, W.J., et al. (2009). Collateral vessels on CT angiography predict outcome in acute ischemic stroke. Stroke *40*, 3001–3005. 10.1161/STROKEAHA.109.552513.

3. World Health Organisation. The top 10 causes of death https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death.

4. Zhang, H., Prabhakar, P., Sealock, R., and Faber, J.E. (2010). Wide genetic variation in the native pial collateral circulation is a major determinant of variation in severity of stroke. J. Cereb. Blood Flow Metab. *30*, 923–934. 10.1038/jcbfm.2010.10.

5. Clayton, J.A., Chalothorn, D., and Faber, J.E. (2008). Vascular endothelial growth factor-a specifies formation of native collaterals and regulates collateral growth in ischemia. Circ. Res. *103*, 1027–1036. 10.1161/CIRCRESAHA.108.181115.

6. Lucitti, J.L., MacKey, J.K., Morrison, J.C., Haigh, J.J., Adams, R.H., and Faber, J.E. (2012). Formation of the collateral circulation is regulated by vascular endothelial growth factor-a and a disintegrin and metalloprotease family

members 10 and 17. Circ. Res. *111*, 1539–1550.

10.1161/CIRCRESAHA.112.279109.

7.   Lucitti, J.L., Sealock, R., Buckley, B.K., Zhang, H., Xiao, L., Dudley, A.C., and

Faber, J.E. (2016). Variants of Rab GTPase-Effector Binding Protein-2 Cause

Variation in the Collateral Circulation and Severity of Stroke. Stroke *47*, 3022–

3031. 10.1161/STROKEAHA.116.014160.

8.   Chalothorn, D., Clayton, J.A., Zhang, H., Pomp, D., and Faber, J.E. (2007).

Collateral density, remodeling, and VEGF-A expression differ widely between

mouse strains. Physiol. Genomics. *30*, 179–191.

10.1152/PHYSIOLGENOMICS.00047.2007.

9.   Das, S., Goldstone, A.B., Wang, H., Farry, J., D'Amato, G., Paulsen, M.J.,

Eskandari, A., Hironaka, C.E., Phansalkar, R., Sharma, B., et al. (2019). A

Unique Collateral Artery Development Program Promotes Neonatal Heart

Regeneration. Cell *176*, 1128-1142.e18. 10.1016/J.CELL.2018.12.023.

10.   Perovic, T., Hollfinger, I., Mayer, S., Lips, J., Dopatka, M., Harms, C., and

Gerhardt, H. (2024). Pial collaterals develop through mosaic colonization of

capillaries by arterial and microvascular endothelial cells. Elife 13.

10.7554/ELIFE.93746.1

11.   Jamaiyar, A., Juguilon, C., Wan, W., Richardson, D., Chinchilla, S., Gadd, J.,

Enrick, M., Wang, T., McCabe, C., Wang, Y., et al. (2022). The essential role

for endothelial cell sprouting in coronary collateral growth. J. Mol. Cell. Cardiol.

165, 158–171. 10.1016/j.yjmcc.2022.01.005.

12.   Gabhann, F. Mac, and Peirce, S.M. (2010). Collateral Capillary Arterialization

following Arteriolar Ligation in Murine Skeletal Muscle. Microcirculation *17*,

333–347. 10.1111/J.1549-8719.2010.00034.X.

13.   Arolkar, G., Kumar, S.K., Wang, H., Gonzalez, K.M., Kumar, S., Bishnoi, B.,

Rios Coronado, P.E., Woo, Y.J., Red-Horse, K., and Das, S. (2023).

Dedifferentiation and Proliferation of Artery Endothelial Cells Drive Coronary

Collateral Development in Mice. Arterioscler. Thromb. Vasc. Biol. *43*, 1455–

1477. 10.1161/ATVBAHA.123.319319.

14.   Chang, A.H., Raftrey, B.C., D'Amato, G., Surya, V.N., Poduri, A., Chen, H.I.,

Goldstone, A.B., Woo, J., Fuller, G.G., Dunn, A.R., et al. (2017). DACH1

stimulates shear stress-guided endothelial cell migration and coronary artery

growth through the CXCL12-CXCR4 signaling axis. Genes. Dev. *31*, 1308–

1324. 10.1101/GAD.301549.117/-/DC1.

15.   Chen, H.I., Sharma, B., Akerberg, B.N., Numi, H.J., Kivelä, R., Saharinen, P.,

Aghajanian, H., McKay, A.S., Bogard, P.E., Chang, A.H., et al. (2014). The

sinus venosus contributes to coronary vasculature through VEGFC-stimulated

angiogenesis. Development *141*, 4500–4512. 10.1242/dev.113639.

16.   Scott Budinger, G.R., Duranteau, J., Chandel, N.S., and Schumacker, P.T.

(1998). Hibernation during Hypoxia in Cardiomyocytes ROLE OF

MITOCHONDRIA AS THE O_2 SENSOR. J. Biol. Chem. *273*, 3320–3326.

10.1074/jbc.273.6.3320.

17.   Rink, C., and Khanna, S. (2011). Significance of Brain Tissue Oxygenation and

the Arachidonic Acid Cascade in Stroke. https://home.liebertpub.com/ars *14*,

1889–1903. 10.1089/ARS.2010.3474.

18.   Mannion, J.D., Buckman, P.D., Magno, M.G., and Dimeo, F. (1992). Collateral

blood flow from skeletal muscle to normal myocardium. J. Surg. Res. *53*, 578–

587. 10.1016/0022-4804(92)90258-2.

19.   Crissinger, K.D., and Neil Granger, D. (1988). Characterization of Intestinal

Collateral Blood Flow in the Developing Piglet. Pediatr. Res. *24*, 473–476.

10.1203/00006450-198810000-00011.

20.   Kochi, T., Imai, Y., Takeda, A., Watanabe, Y., Mori, S., Tachi, M., and

Kodama, T. (2013). Characterization of the Arterial Anatomy of the Murine

Hindlimb: Functional Role in the Design and Understanding of Ischemia

Models. PLoS One *8*, e84047. 10.1371/JOURNAL.PONE.0084047.

21.   Gao, Y., Aravind, S., Patel, N.S., Fuglestad, M.A., Ungar, J.S., Mietus, C.J., Li,

S., Casale, G.P., Pipinos, I.I., and Carlson, M.A. (2020). Collateral

Development and Arteriogenesis in Hindlimbs of Swine After Ligation of

Arterial Inflow. J. Surg. Res. *249*, 168–179. 10.1016/J.JSS.2019.12.005.

22.   He, L., Liu, Q., Hu, T., Huang, X., Zhang, H., Tian, X., Yan, Y., Wang, L.,

Huang, Y., Miquerol, L., et al. (2016). Genetic lineage tracing discloses

arteriogenesis as the main mechanism for collateral growth in the mouse

heart. Cardiovasc. Res. *109*, 419–430. 10.1093/CVR/CVW005.

23.   Raftrey, B., Williams, I.M., Rios Coronado, P.E., Fan, X., Chang, A.H., Zhao,

M., Roth, R., Trimm, E., Racelis, R., D'amato, G., et al. (2021). Dach1 Extends

Artery Networks and Protects Against Cardiac Injury. Circ. Res. *129*, 702–716.

10.1161/CIRCRESAHA.120.318271.

24.   Zhang, H., Chalothorn, D., and Faber, J.E. (2019). Collateral Vessels Have

Unique Endothelial and Smooth Muscle Cell Phenotypes. Int. J. Mol. Sci. *20*,

3608. 10.3390/IJMS20153608.

25.   Stanca Melincovici, C., Boşca, A.B., Şuşman, S., Mărginean, M., Mihu, C.,

Istrate, M., Moldovan, I.-M., Roman, A.L., and Mihu, C.M. (2018). Vascular

endothelial growth factor (VEGF)-key factor in normal and pathological

angiogenesis. Rom. J. Morphol. Embryol. *59*, 455-467.

26.   Miquerol, L., Thireau, J., Bideaux, P., Sturny, R., Richard, S., and Kelly, R.G.

(2015). Endothelial plasticity drives arterial remodeling within The endocardium

after myocardial infarction. Circ. Res. *116*, 1765–1771.

10.1161/CIRCRESAHA.116.306476.

27.   Miquerol, L., Meysen, S., Mangoni, M., Bois, P., van Rijen, H.V.M., Abran, P.,

Jongsma, H., Nargeot, J., and Gros, D. (2004). Architectural and functional

asymmetry of the His–Purkinje system of the murine heart. Cardiovasc. Res.

63, 77–86. 10.1016/j.cardiores.2004.03.007.

28.   Knowland, D., Arac, A., Sekiguchi, K.J., Hsu, M., Lutz, S.E., Perrino, J.,

Steinberg, G.K., Barres, B.A., Nimmerjahn, A., and Agalliu, D. (2014).

Stepwise Recruitment of Transcellular and Paracellular Pathways Underlies

Blood-Brain Barrier Breakdown in Stroke. Neuron *82*, 603–617.

10.1016/j.neuron.2014.03.003.

29.   Lutz, S.E., Smith, J.R., Kim, H., Bates, J.M., Gandhi, S.P., and Agalliu, D.

(2017). Caveolin1 Is Required for Th1 Cell Infiltration, but Not Tight Junction

Remodeling, at the Blood-Brain Barrier in Autoimmune Neuroinflammation.

CellReports *21*, 2104–2117. 10.1016/j.celrep.2017.10.094.

30. Holtmaat, A.J.G.D., Trachtenberg, J.T., Wilbrecht, L., Shepherd, G.M., Zhang,

X., Knott, G.W., and Svoboda, K. (2005). Transient and Persistent Dendritic

Spines in the Neocortex In Vivo. Neuron *45*, 279–291.

10.1016/j.neuron.2005.01.003.

Main figures and figure legends

Figure 1: Live imaging of embryonic pial vasculature

(**A**) Intra-vital imaging set up to visualize the developing pial vasculature in mouse embryos. **i** heating pad to maintain body temperature of anaesthetized pregnant dam at 37°C. **ii** custom-built media heating device which also circulates dissection media at 37°C **iii** shows macro-zoom or confocal microscopy performed on live, immobilized embryos.

(**B**) Representative schematic of an embryonic brain. Grey and Red show cerebral arteries and pial collaterals respectively.

(**C**) Experimental design to genetically label and trace the dynamics of pre-existing artery endothelial cells in a developing pial layer.

(**D-G**) Images obtained through macro-zoom fluorescent microscopy of e16.5 embryos. In black are cells derived from pre-existing artery endothelial cells. Images were taken by focusing in and out to capture the entire curvature of the embryonic brain. **D** shows an entire left hemisphere. **E** represents watershed region spanned by MCA and ACA artery tips and imaged at time=0 minutes, 20 minutes and 80 minutes respectively. Arrows and arrowheads show growing ends of pre-existing artery tips from MCA and ACA. Brackets mark regions where collaterals appear. **F** shows inset in **D**. **E** and **G** show insets in **F**. **G** represents the MCA growth front imaged at time=0 minutes, 20 minutes and 80 minutes respectively. * shows a superficial yolk sac vessel which photobleaches with time. Arrowheads mark extension and anastomoses of MCA artery tips.

(**H-J**) Images obtained through confocal microscopy from e16 embryos. **H** shows pial watershed and the cerebral arteries spanning the watershed. **I**, **J** show MCA and ACA growth fronts respectively imaged at time=0 minutes and 60 minutes. In black

are cells derived from pre-existing artery endothelial cells. Images shown in **I** are

maximum intensity projections from 50 optical slices and capture the entire width of

the artery. Images shown in **J** are maximum intensity projections from 40 optical

slices and capture the entire width of the artery. * represents an internal landmark

from where the length measurements were done.

(**K-M**) Images obtained through confocal microscopy from e16 embryos. (**L, M**)

Insets from **K** showing artery cell growth of developing ACA in 60 mins. (**L**)

Arrowhead points to an artery tip emerging within 60mins. Arrow points to a region

where single artery cells appear. (**M)** Arrow shows direction of migration of single

artery cell. Dotted line is the length of micro-vessel between the single cell and the

pial artery. MCA, Middle cerebral artery; ACA, Anterior cerebral artery; e, embryonic

day; Tam, Tamoxifen; EC, endothelial cell; YSV, yolk sac vessel; brch., branch; coll,

collateral; MV, microvascular. Scale bars: **D**, **F**, 1mm; **E**, **G**, 200 µm; **H**, **K**, 500 µm; **I**,

J, **L**, **M**, 100 µm.

A
e16

MCA — watershed — B — ACA

B
TdTomato (artery cell lineage)
eGFP (all ECs)

eGFP
TdTomato

Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5

to MCA — MV track
Td+ EC
MV track
MV track
Td+ EC — to ACA

eGFP
MV track
Td+ EC
MV track

C
Potential mechanisms leading to pial collateral fromation

capillaries — microvascular tracks — pre-existing artery tips

Watershed e14.5
i

MV tracks form between two artery tips
inception of MV tracks

ii — e16 — v

artery tip extension
artery tip extension — MV track remodelling

iii — e16.5 — vi

collateral
iv
e18.5/ P0

Growth of MV tracks and arteries occur concurrently

Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5

Tam at e14.5

D
e16
TdTomato (artery cell lineage) eGFP (all ECs)
E-H

I
e16.5
K-N

E
to ACA ← watershed → to MCA
TdTomato (artery cell lineage) eGFP (all ECs)

F
all ECs

G
artery cells

H
skeleton of MV tracks
with or without TdTomato+ cells

J
Total length of MV track per watershed
p=0.009
**
Normalized MV track length per watershed area (AU)
2.0
1.5
1.0
0.5
e16   e16.5

O
Td coverage of MV tracks per watershed
ns
% coverage of Td+ cells
e16   e16.5

K
to ACA ← watershed → to MCA
TdTomato (artery cell lineage) eGFP (all ECs)

L
all ECs

M
artery cells

N
skeleton of MV tracks
with or without TdTomato+ cells

Figure 2: Analyses of development of MV (microvascular) tracks and artery

tips during pial collateral development

(**A**) Pial watershed and the cerebral arteries spanning the watershed in e16 mouse

brain.

(**B**) Inset from **A** showing TdTomato⁺ artery endothelial cells (yellow arrowheads) on

eGFP⁺ microvascular tracks (magenta arrows).

(**C**) Two potential mechanisms for pial collateral formation. Hypothesis 1: (i) An

artery-free e14.5 watershed region (ii) is first occupied by microvascular tracks

(green) connecting two arteries (red) by e16. (iii) Artery tips then start extending on

pre-determined MV tracks at e16.5. Arrows indicate direction of artery tip migration.

(iv) This process leads to pial collateral formation. Hypothesis 2: Alternatively, (v)

while MV tracks start appearing at e16, (vi) they actively remodel and grow in length

to allow artery tip extensions in a directional manner. This leads to formation of pial

collaterals.

(**D-N**) Concurrent MV track growth and artery tip extension. (**D and I**) Whole brain

images of e16 and e16.5 embryos, respectively, from *Cx40CreER; Rosa26^TdTomato*;

Tie2:eGFP:Claudin5 mice, where TdTomato (red) marks pre-existing artery

endothelial cells and eGFP (green) marks all endothelial cells. **E-H** is inset shown in

D. (**E**) Higher magnification confocal image of watershed region (715 μm width)

between ACA and MCA. All endothelial cells are in green and artery endothelial cells

in red. (**F**) eGFP⁺ vessels in the selected watershed from e16 watershed shown in **E**.

(**G**) Pre-existing artery cells in the watershed from e16 watershed shown in **E**. (**H**)

Skeleton of all the eGFP⁺ microvascular tracks traced in the e16 watershed (shown

in **E**) through multiple z-slices, using Simple Neurite Tracer. Traced MV tracks

occupied by TdTomato⁺ artery endothelial cells are shown in red and MV tracks not

occupied with TdTomato is shown in cyan. (**J**) Quantification of total length of

microvascular tracks per watershed area (p=0.009; unpaired t test) and (**O**)

Quantification of percentage TdTomato coverage of microvascular tracks in

watershed regions from e16 (9 watersheds from n=9 embryos) and e16.5 (7

watersheds from n=5 embryos) brains (p=0.4085; unpaired t test). **K-N** is inset

shown in **I**. (**K**) Higher magnification confocal image of watershed region (715 μm

width) between ACA and MCA. All endothelial cells are in green and artery

endothelial cells in red. (**L**) eGFP$^+$ vessels in the selected watershed from e16

watershed shown in **K**. (**M**) Pre-existing artery cells in the watershed from e16

watershed shown in **K**. (**N**) Skeleton of all the eGFP$^+$ microvascular tracks traced in

the e16 watershed (shown in **K**) through multiple z-slices, using Simple Neurite

Tracer. Traced MV tracks occupied by TdTomato$^+$ artery endothelial cells are shown

in red and MV tracks not occupied with TdTomato is shown in cyan. Tam,

Tamoxifen; MCA, Middle Cerebral Artery; ACA, Anterior Cerebral Artery; MV,

Microvascular; Td, TdTomato; e, embryonic day; EC, Endothelial cell; AU, Arbitrary

units. Scale bar: A, 500μm; B, 100μm; D and I, 500μm; E and K, 100μm.

Figure 3: Cellular origin of pial collaterals

(**A**) Experimental design to assess lineage contribution from artery ECs and capillary ECs towards pial collaterals at e18.5/ P0.

(**B**) Graph shows the average percent area of collateral segments covered with TdTomato$^+$ cells.

(**C**) Representative image of whole right hemisphere from *Cx40CreER*; *Rosa26^{TdTomato}* e18.5 embryos. Black shows TdTomato$^+$ arteries.

(**D**) Boxed region of interest with a collateral segment from the watershed area shown in **C**. The pial collateral is lineage traced for artery ECs (red) and immunostained with Cx40 (green).

(**E**) Representative image of whole right hemisphere from *ApjCreER*; *Rosa26^{TdTomato}* e18.5 embryos. Black shows Cx40 immunostaining.

(**F**) Boxed region of interest with a collateral segment from the watershed area shown in **E**. The pial collateral is lineage traced for capillary ECs (magenta) and immunostained with Cx40 (cyan).

(**G**, **I**) Collateral segment from **D** is (**G**) immunostained for Cx40 and (**I**) lineage traced with TdTomato.

(**H**, **J**) Collateral segment from **F** is (**H**) immunostained for Cx40 and (**J**) lineage traced with TdTomato. Arrowheads point to TdTomato$^+$ cells from Apj$^+$ lineage on the collateral segment. e, embryonic day, P, postnatal day; MCA, Middle cerebral artery; ACA, Anterior cerebral artery; brch., branch; EC, endothelial cell; Tam, Tamoxifen.

Scale bars: **C** and **E**, 500µm; **D** and **F**, 100µm

A

Cxcr4^{L/L}; Cx40CreER; Rosa26^{TdTomato}
and
VegfR2^{L/L}; Cx40CreER; Rosa26^{TdTomato}

B Arterial depletion of Cxcr4
does not change pial collateral artery network

Cx40CreER;Rosa26^{TdTomato}   *Cxcr4^{L/L};Cx40CreER;Rosa26^{TdTomato}*

C

Quantification of pial artery coverage
in watershed

D Arterial VegfR2 regulates formation of
pial collateral artery network

Cx40CreER;Rosa26^{TdTomato}   *VegfR2^{L/L};Cx40CreER;Rosa26^{TdTomato}*

E Reduced artery coverage
in absence of arterial VegfR2

F Reduced collateral artery network
in absence of arterial VegfR2

Cx40CreER;Rosa26^{TdTomato}
VegfR2^{L/L};Cx40CreER;Rosa26^{TdTomato}

VegfR2-null artery tips fail to extend and anastomose

G

Cx40CreER; Rosa26^{TdTomato}
or
Cx40CreER; Rosa26^{TdTomato}
Tie2::eGFP::Claudin5

with or without arterial *VegfR2*

H Reduced number of
Td^{con} collaterals
without arterial VegfR2

I *Cx40CreER; Rosa26^{TdTomato} Tie2::eGFP::Claudin5*
TdTomato (pre-existing artery ECs) eGFP (all ECs)

J

Figure 4: VegfR2-dependent formation of pial collateral arteries

(**A**) Experimental design to lineage label and trace pre-existing artery endothelial cells in pial layer, with or without arterial *VegfR2* and *Cxcr4* at e18.5/ P0.

(**B**) Representative confocal images of whole brains with and without arterial *Cxcr4*. Black shows pre-existing artery endothelial cells. Control= 8, artery specific knockout for *Cxcr4*= 13.

(**C**) Summary of method to quantify pial artery coverage across the entire brain including watershed.

(**D**) Confocal images of whole brains with and without arterial *VegfR2*. Black shows pre-existing artery endothelial cells.

(**E**) Quantification of overall pial artery coverage with and without arterial *VegfR2* quantified using method shown in **C**. KS test performed on the distributions generated D value= 0.569, p value= 2.2066×10^{-141}.

(**F**) Quantification of pial collateral artery network with and without arterial *VegfR2* quantified using method shown in **C**. Unpaired t-test, p=0.0001.

(**G**) Experimental design to assess the effect of *VegfR2* deletion from arteries at e16.5. (**H**) Quantification of pial collateral arteries contiguously populated with TdTomato$^+$ artery cells, with and without arterial *VegfR2*. (**I**, **J**) Confocal images of pial watershed region showing TdTomato-contiguous pial collateral arteries (Td^{con}) in red and microvasculature in green. Arrowheads show part of microvascular track not occupied by TdTomato$^+$ artery cell, arrows point to artery tips on microvascular tracks. MCA, Middle cerebral artery; ACA, Anterior cerebral artery; e, embryonic day; P, postnatal day; Tam, Tamoxifen; EC, endothelial cell; Td^{con}, TdTomato-contiguous;

994   AU, Arbitrary Unit; aEC, artery endothelial cell; KO, knockout. Scale bars: **B**, **D**,

995   500µm; **I**, **J**, 100µm.

Intra-vital imaging of pial collaterals in adult mice

 Figure 5: Longitudinal live-imaging of adult pial collaterals

(**A**, **left**) Dorsal view of the cranial window created with the attached headbar on

adult mouse head. (**A**, **right**) Brightfield image of the 4mm cranial window obtained

with a stereoscope shows the pial vasculature. Dotted rectangle encompasses MCA,

ACA and the watershed between them and is imaged at high resolution as shown in

E.

(**B**) Schematic of the side view of the cranial window showing the 3-tier optical plug

inserted into the skull giving access to pial layer.

(**C**) Experimental design for longitudinal imaging of pial vasculature in adult mice.

(**D**) Quantification of TdTomato reporter recombination in pial collateral arteries and

different segments of cerebral arteries. One-way ANOVA was done to test

significance statistically (mean comparison Tukey test, PA vs. collateral, $p<0.0001$;

DA vs. collateral, $p<0.0001$; PA vs. DA, $p=0.08$).

(**E-K**) Images of pial watershed with pre-existing artery endothelial cells labelled in

TdTomato (red) and all endothelial cells labelled in eGFP (green). (**E**) Confocal

image from optical window created for longitudinal imaging of *Cx40CreER;*

Rosa26^{TdTomato}; Tie2::eGFP::Claudin5 mouse as shown in Figure **4A**. The image

shows both pial and dural vessels. (**F-K**) Confocal images of pial collateral arteries

from cranial window shown in **E**, across 3 timepoints, each 2 weeks apart. (**I**) Pial

layer segmented out from **E**. red and blue show pial arteries and veins respectively.

J, **K** are the insets shown in **F-H**.

(**L**, **M**) Cellular distribution of TdTomato⁺ artery cells and eGFP⁺ endothelial cells in

collateral artery segments shown in **J** and **K** respectively. * represents a reference

cell. MCA, middle cerebral artery; ACA, anterior cerebral artery; d, days; Tam,

Tamoxifen; PA, proximal artery; DA, distal artery; coll, collateral; brch., branch; EC,

endothelial cell. Scale bars: **E**, 200µm; **J**, **K** 25µm

1027

1028

1029

1030

Figure 6: Analyses of role of arterial VegfR2 in pial collateral maintenance

(A) Experimental design to visualize artery cell dynamics in pial layer of adult mice without arterial *VegfR2*.

(**B**) Brightfield image of cranial window showing pial vasculature. The dotted rectangle encompasses branches of MCA, ACA and the watershed between them.

(**C**) Confocal image of cranial window created in mice without arterial *VegfR2* as shown in **B**. Red shows all artery cells, green shows all endothelial cells.

(**D**) Pial layer segmented out from **C** shows pial arteries and veins in red and blue respectively.

(**E-G**) Confocal images of collateral artery from three timepoints, each 2 weeks apart as shown in the boxed region in **C**.

(**H-J**) Confocal images of collateral artery segments shown in **E-G**.

(**K-M**) Cellular distribution of TdTomato$^+$ artery cells and eGFP$^+$ endothelial cells in collateral artery segments shown in **H-J**. * represents a reference cell.

(**N**) Schematic summarizing the role of arterial VegfR2 in development and maintenance of pial collateral arteries. While arterial VegfR2 facilitates tip extension of pre-existing arteries during development, it is dispensable for pial collateral maintenance in adulthood. Pial collaterals, both in developing and mature brains, are majorly populated by endothelial cells of artery-lineage. MCA, middle cerebral artery; ACA, anterior cerebral artery; d, days; Tam, Tamoxifen; EC, endothelial cell; brch., branch. Scale bars: **C**, 150 μm; **E-G**, 50μm; **H-J**, 20μm.

Supplementary figures and figure legends

Development of cerebral artery trees and pial collaterals in embryonic mouse brains

1054

1055

1056

1057

1058

1059

Supplemental Figure 1: Visualization of pial arteries and collateral arteries

Schematic summarizes key steps of immunostaining methods used to image pial artery network in (**A**) mouse embryonic heads and (**B**) mouse embryonic/neonatal brains with confocal microscope. (**C**) Schematic shows findings from **A** and **B** in one hemisphere. Grey represents cerebral arteries, Red represents collateral artery connections between cerebral arteries. Brackets show watershed area (WSA), MCA appears at e12.5, ACA appears at e13.5, and the foremost pial collateral arteries appear at e15.5. These pial collaterals then increase in number till e18.5/P0. e, embryonic; P, postnatal day; MCA, middle cerebral artery; ACA, anterior cerebral artery.

Supplemental Figure 2: Detection of pial collaterals with whole brain immunostaining

(**A-F**) Representative images of right hemispheres from developing mouse embryos immunostained for Connexin-40 (Cx40). Cx40 marks all artery endothelial cells. MCA appears at e12.5, ACA appears at e13.5 and collaterals between MCA and ACA appear at e15.5. Arrowheads point to collateral arteries between MCA and ACA. (**G**) Quantification of Cx40 immunostained brains from different developmental stages reveals that both MCA and ACA have formed by e14.5. (**H**) Quantification of Cx40$^+$ collateral arteries per hemisphere shows that the first pial collaterals appear at e15.5. e, embryonic; P, postnatal day; MCA, middle cerebral artery; ACA, anterior cerebral artery; brch., branch. Scale bars: 500μm

left hemisphere

right hemisphere

Supplemental Figure 3: Detection of pial collaterals with *Cx40* reporter activity

(**A-F**) Representative images of brains from developing *Cx40eGFP/+* mouse embryos.

Connexin-40 (Cx40) marks all artery endothelial cells. MCA appears at e12.5, ACA

appears at e14.5 and collaterals between MCA and ACA appear at e15.5.

Arrowheads point to collateral arteries between MCA and ACA. (**G**) Quantification of

Cx40eGFP/+ brains from different developmental stages reveals that both MCA and

ACA have formed by e14.5. (**H**) Quantification of eGFP+ collateral arteries per

hemisphere shows that the first pial collaterals appear at e15.5. e, embryonic; P,

postnatal day; MCA, middle cerebral artery; ACA, anterior cerebral artery; brch.,

branch. Scale bars: 500µm

embryo1

1105

Supplemental Figure 4: Visualization of pial artery cell dynamics on

microvascular tracks Confocal image of e16 embryonic collateral zone showing

single artery endothelial cell migration towards the pial collateral within 110mins.

Dotted line is the length of micro-vessel between the single cell and the pial

collateral. Arrow shows direction of single artery cell migration. MCA, Middle

Cerebral Artery; ACA, Anterior Cerebral Artery. Tamoxifen was injected at e13.5.

Scale bars: **A**, 100μm.

1113

A

Cxcr4^{L/L};Cx40CreER; Rosa26^{TdTomato}
Tie2::eGFP::Claudin5

artery-EC specific deletion of *Cxcr4*
does not impact adult pial collateral artery network

B *Cx40CreER;Rosa26^{TdTomato}*

C *Cxcr4^{L/L};Cx40CreER;Rosa26^{TdTomato}*

1114

1115

1116

1117

1118

1119

1120

1121

1122

 Supplemental Figure 5: Pial artery coverage does not change without arterial

 Cxcr4 (**A**) Experimental set up to assess the effect of arterial *Cxcr4* deletion on pial

 artery coverage in adults. (**B**, **C**) Representative confocal images of whole adult

 brains (from N=5 for each genotype), genetically lineage traced for pre-existing

 artery endothelial cells, (**B**) with or (**C**) without arterial *Cxcr4*. ACA, anterior cerebral

 artery; MCA, middle cerebral artery; Tam, Tamoxifen. Scale bars: 1mm

1129

1130

1131

1132

1133

1134

1135

1136

1137

A

Dach1 is highly expressed in pial capillary ECs at P0

No effect on collateral development with deletion of *Dach1* from capillary ECs

B

ApjCreER; Dach1fl/fl

C

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

Supplemental Figure 6: Dach1-independent pial collateral development (**A**)

Dach1 expression (green) in capillaries and arteries in P0 brains. Dotted line marks

the TdTomato⁺ (red) arterial boundary. VegfR2 marks all vasculature. DAPI marks

nuclei. (**B**) Experimental design showing Tamoxifen injection at e14.5, and analyses

of collateral numbers in e18.5/P0 brains. (**C**) Quantification of Cx40⁺ pial collaterals

between MCA and ACA in control (*Dach1^fl/fl*) and *ApjCreER*; *Dach1^fl/fl* (Dach1 cap-

EC KO) brains. e, embryonic; cap-EC, capillary endothelial cell; KO, knockout; Tam,

Tamoxifen. Scale bar: 20µm.

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

Depletion of arterial VegfR2 does not affect neonatal pial collateral artery network

A *Cx40CreER; Rosa26^{TdTomato}* or *VegfR2^{L/L};Cx40CreER; Rosa26^{TdTomato}*

B P7, *Cx40CreER; Rosa26^{TdTomato}*

C P7, *VegfR2^{L/L};Cx40CreER; Rosa26^{TdTomato}*

P2 hemispheres

D *Cx40CreER; Rosa26^{TdTomato}*

E *VegfR2^{L/L};Cx40CreER; Rosa26^{TdTomato}*

F Quantification of arterial coverage

G Quantification of AUC (arterial coverage in watershed)

Cx40CreER;Rosa26^{TdTomato}
VegfR2^{L/L};Cx40CreER;Rosa26^{TdTomato}

P7 hemispheres

H *Cx40CreER; Rosa26^{TdTomato}*

I *VegfR2^{L/L};Cx40CreER; Rosa26^{TdTomato}*

J Quantification of arterial coverage

K Quantification of AUC (arterial coverage in watershed)

Cx40CreER;Rosa26^{TdTomato}
VegfR2^{L/L};Cx40CreER;Rosa26^{TdTomato}

1170

1171

1172

1173

1174

1175

Supplemental Figure 7: VegfR2-independent remodelling of pial collaterals

(**A**) Experimental design. (**B**) Confocal imaging of whole brains immunostained for

VegfR2 shows presence of the protein in pial arteries. Boxed region in left image is

magnified as the right image. Arrows point to VegfR2 protein expression in MCA

branches. (**C**) Confocal imaging of whole brains immunostained for VegfR2 shows

absence of the protein in pial arteries (enclosed in dotted lines). Boxed region in left

image is magnified as the right image. (**D**, **E**) Confocal whole P2 brain images

showing TdTomato$^+$ arteries (**D**) with or (**E**) without VegfR2. (**F**, **G**) shows

quantification of (**F**) pial artery coverage across the entire P2 hemisphere or (**G**)

collateral artery coverage in the watershed, in brains with or without arterial VegfR2.

Unpiared t-test was performed with p value= 0.5. (**H**, **I**) Confocal whole P7 brain

images showing TdTomato$^+$ arteries (**H**) with or (**I**) without VegfR2. (**J, K**) shows

quantification of (**J**) pial artery coverage across the entire P7 hemisphere or (**K**)

collateral artery coverage in the watershed, in brains with or without arterial VegfR2.

Unpiared t-test was performed with p value= 0.63. Scale bars: 500 µm

1191

1192

Intra-vital imaging of VegfR2 deficient pial collaterals in adult mice

VegfR2^{L/L} Cx40CreER; Rosa26^{TdTomato}
Tie2::eGFP::Claudin5
TdTomato (pre-existing artery ECs) eGFP (all ECs)

1193

1194

1195

1196

1197

1198

Supplemental Figure 8: VegfR2-independent maintenance of pial collaterals

Confocal images of the entire cranial window from 3 different mice shows that the overall pial artery network remains unchanged without arterial *VegfR2*. MCA, Middle Cerebral Arteries, ACA, Anterior Cerebral Arteries, Coll, Pial collateral arteries between MCA and ACA. Scale bars: 500µm.