Development of Pial Collaterals by Extension of Pre-existing Artery Tips

Suraj Kumar, Niloufer Shanavas, Swarnadip Ghosh, Vinayak Sivaramakrishnan,

Manish Dwari, Soumyashree Das[#]

National Centre for Biological Sciences, Tata Institute of Fundamental Research,

Bengaluru, KA 560065, India

[#] Corresponding author (soumyashree@ncbs.res.in)

Key Words

Pial Collateral Artery, Artery Endothelial Cell, Vascular Endothelial Growth Factor

Receptor-2, Intra-vital Imaging, Genetic Lineage Tracing

SUMMARY

Pial collaterals provide protection from ischemic damage and improve prognosis of

stroke patients. The origin or precise sequence of events underlying pial collateral

development is unclear and has prevented clinicians from adapting new

vascularization and regeneration therapies. We use genetic lineage tracing and intra-

vital imaging of mouse brains at cellular resolution to show that during

embryogenesis, pial collateral arteries develop from extension and anastomoses of

pre-existing artery tips in a VegfR2 dependent manner. Our data demonstrate that

an arterial receptor, Cxcr4, earlier shown to drive artery cell migration and coronary

collateral development, is dispensable for formation and maintenance of pial

collateral arteries. Our study reveals that collateral arteries of the brain are built by a

unique mechanism, distinct from that of the heart.

INTRODUCTION

Pial (leptomeningeal) collateral arteries are an extensive network of inter-artery anastomoses found in the pial layer of the brain. In humans, the extent of pial collateral circulation is a predictor for stroke outcome.[1,2] Stroke occurs when blood supply is obstructed to a part of the brain and is one of the leading causes of morbidity worldwide (World Health Organization).[3] Pial collaterals serve as an alternate route of blood flow and alleviate the symptoms of ischemic stroke by bypassing vascular occlusions. Thus, understanding mechanisms which drive the origin and remodelling of pial collateral arteries is of immense clinical relevance.

Results from preclinical studies indicate wide-spread variation in the pial collateral artery network among individuals and organs.[4] These variations are a result of differential expression and function of proteins associated with Vascular endothelial growth factor (Vegf) signalling axis.[5–7] Different mouse strains demonstrate differential Vegfa expression level which corelate with their pial collateral density.[8] Similarly, global overexpression of Vegf ligand increases number of native pial collaterals and improve collateral perfusion post-injury.[5,6] The receptor mediating function of Vegf ligand during the formation of pial collaterals is Vegf receptor-2 (VegfR2).[6] A global knockdown of Vegf or VegfR2 during embryogenesis and in a time-frame when pial collaterals are expected to form, results in reduction in their number and diameter.[6] Thus, Vegf is a key driver of pial collateral formation and a critical factor which regulates variation in extent of pial vascular network in mice.

The vital role of Vegf in murine pial collateral development was discovered almost a decade ago.[5,6] However, it is still unclear how the various molecular players of Vegf pathway orchestrate the formation or remodelling of pial collateral network. In

mice, it is hypothesized that an *arteriole-specific sprouting angiogenesis like* process

leads to the formation of the foremost pial collaterals.[6] However, molecular evidence

supporting the identity of these sprouting cells (if arteries or capillaries) is missing.

Additionally, the cellular origin or fate of these sprouts is unknown. Finally, the

possibility of cooperation from other vascular processes, such as arterialization,

arteriogenesis or artery-reassembly, in building pial collaterals, needs to be

thoroughly investigated.

In this study, we use genetic lineage tracing in conjunction with live imaging of

pial vasculature of transgenic mouse embryos and adults to understand mechanisms

by which Vegf regulates pial collateral development and maintenance. Our data

indicate that the cellular source of pial collateral arteries is primarily pre-existing

artery endothelial cells (ECs) which build the first network of pial collaterals at

embryonic day (e) 15.5 via VegfR2-dependent artery tip extension. While arterial

VegfR2 was essential for formation of pial collaterals *de novo*, it was dispensable for

its maintenance through adulthood. Arterial Cxcr4 (C-X-C chemokine receptor type-

4), which is critical for coronary artery formation in mice[9] was dispensable during pial

collateral formation and its maintenance. Together, our data indicate that artery

endothelial cells build collateral arteries by adapting distinct vascular mechanisms in

heart and brain.

RESULTS

Pial collateral arteries form during embryogenesis

We developed a whole brain imaging technique to reliably visualize all pial

arteries (Figure **S1A**, **B**). Due to lack of unique markers, accurately identifying pial

collateral arteries is exclusively dependent on their location. Pial collateral arteries

are artery segments which interconnect the ends of cerebral arteries and can be

seen in the watershed region (Figure **S1B**). To acquire this spatial information, it was

imperative to image the entire pial network. Our technique allowed us to precisely

identify pial collateral arteries. We specifically focused on collateral arteries which

connected and received blood flow from Anterior Cerebral Artery (ACA) and Middle

Cerebral Artery (MCA) as the coverage of this arterial network is most extensive.

We first investigated the developmental stage when pial collaterals first

appeared. Mouse brains were harvested at different developmental time points and

whole brain immunostaining was performed for Connexin-40 (Cx40) protein, a known

marker for mature artery endothelial cell. Analyses of pial arterial network in both

hemispheres revealed the appearance of MCA at embryonic day e12.5 followed by

ACA at e13.5 (Figure **S1C** and **S2**). At e14.5, both cerebral arteries were present

with no arterial connections between them. Subsequently, the first arterial

anastomoses appeared at e15.5 which then remodelled to expand in number

through birth/postnatal day (P)0 (Figure **S1C** and **S2A-F**). This method confirmed

that both MCA and ACA are present at e14.5 (Figure **S2G**) and the foremost pial

collaterals appear at e15.5 (Figure **S2H**).

To confirm the above results, we next used a knock-in mouse line (*Cx40*$^{eGFP/+}$)

where eGFP expression was driven by *Cx40* promoter activity. Brains from different

developmental stages confirmed our findings from Cx40 immunostaining. MCA

appeared at e12.5, ACA appeared at e14.5 and collateral arteries between MCA and

ACA appear at e15.5 (Figure **S3A-F** and **S1C**). eGFP$^+$ cerebral arteries (MCA and

ACA) were formed by e14.5 (Figure **S3G**) and the first eGFP$^+$ collateral artery

between MCA and ACA appeared at e15.5 (Figure **S3H**). Thus, using two

independent methods— Cx40 immunostaining and detecting its reporter activity, we

demonstrate that mouse pial collateral arteries form during development, and appear

precisely at e15.5 (Figure **S1C**).

Intravital imaging reveals arterial origin of pial collaterals by artery tip

extension along microvascular tracks

Despite several hypotheses, there is lack of compelling data in support of

vascular mechanisms which drive pial collateral development. Pial collaterals appear

at late embryonic stages. At this developmental stage rapid vascular changes make

it difficult to distinguish collateral-specific events from overall vascular growth, thus

making it challenging to draw information from fixed brain samples. Additionally, late

embryonic stages cannot be cultured *ex vivo* due to extremely demanding metabolic

and physiological needs. Thus, we custom-developed an imaging paradigm to

perform intra-vital imaging of pial layer at late embryonic stages (e16-e17) (Figure

1A), a developmental stage where the brain is actively building pial collaterals.

Briefly, embryos of the desired developmental time points were dissected out (with

their intact yolk sac) from the anesthetized pregnant mice while still connected to the

mother via their umbilical cord. Embryos were maintained at 37˚C and immobilized at

appropriate orientation to aid imaging. The embryos were subjected to stereoscopic

or confocal imaging for several hours. This imaging set up allowed visualization of

the entire hemisphere— MCA, ACA, and connected collateral arteries (Figure **1B**).

Pial collaterals appear only after the primary branches of MCA and ACA have

developed (Figure **S1C**). We hypothesized that the branches of these cerebral

arteries contribute towards formation of pial collaterals. To test this hypothesis,

Cx40CreER; Rosa26^{TdTomato}; Tie2:: eGFP::Claudin5 mice were used to label artery

ECs with TdTomato and all endothelial cells with eGFP. To activate TdTomato,

Tamoxifen was administered at e13.5, when both MCA and ACA are present and

would be labelled in 2 days, [9,10] by e15.5. Imaging of endothelial cells derived from

arteries was performed at e16-e17. This experimental strategy allowed visualization

of cellular dynamics specific to artery cells derived from pre-existing arteries (Figure

1C) and excluded the possibility of capturing other vascular processes.

We performed intra-vital time-lapse imaging of pial arteries and investigated if

pial collaterals originate from pre-existing artery endothelial cells. We focused on the

watershed region: an initial artery-free region which spans the distal ends of both

arterial trees (MCA and ACA) and is later occupied by pial collaterals (Figure **S1C**

and **1D-F**). Stereoscopic images of e16.5 brains revealed that branches from MCA

and ACA extend (arrowheads, Figure **1E**) and anastomose to build collaterals in the

pial watershed (brackets, Figure **1E**). Additionally, TdTomato expression in artery

tips (orange arrow, Figure **1E**) and collaterals (green arrow, Figure **1E**) increased

over time. These changes were executed within a short span of 80 minutes, and

could not have been captured in fixed brain samples.

We next investigated artery cell dynamics associated with the growth fronts of

the two cerebral arteries (Figure **1F**, **G**). At e16.5, the distal branches of the

developing MCA extended and anastomosed to build an artery network *de novo*

(arrowheads, Figure **1G**). This is the first evidence demonstrating that pre-existing

cerebral artery tips build pial collaterals. We then performed intra-vital time-lapse

confocal imaging of brains at e16, a stage prior to when we observed collateral

artery networks being built de novo (Figure **1E**, **G**). Confocal imaging at e16 provided

the necessary cellular resolution to quantitatively assess artery-tip extensions from

MCA and ACA branches (Figure **1H-J**). As expected, we clearly observed artery tip

extension of MCA branch (Figure **1H**, **I**) within an hour. Additionally, we noted

several artery-specific movements within the pial layer. This included the

appearance of single artery endothelial cells (arrows, Figure **S4A**, **B**), movement of

these single artery cells towards an artery (Figure **1J**, Figure **S4C**), and emergence

of arterial sprouts (arrowheads, Figure **S4A**, **B**). Most interestingly, all artery cell

movements were observed along microvascular tracks which appeared structurally

more organized/tubular than the surrounding capillaries (arrowheads, Figure **1K** and

Figure **S4A**, **D**). These "microvascular tracks" were distinct from the surrounding

capillary plexus in that they formed direct connections between two arteries but were

not entirely populated by TdTomato⁺ artery derived endothelial cells (Figure **1K** and

Figure **S4A**, **D**). Together, for the first time, we provide direct evidence for the

vascular origin of pial collaterals— pre-existing MCA/ACA artery tips grow on distinct

microvascular paths and connect with each other to build pial collaterals *de novo*

(Figure **1L**).

Artery ECs are the major cellular contributors of pial collaterals during

embryogenesis

We next assessed the vascular-lineage composition of pial collaterals at a

later stage when the collateral artery network has fully formed. This would allow us

to: 1) determine if the artery cell contribution to nascent pial collaterals observed at

e16.5 (Figure **1E**) persisted in later stages and 2) check contribution from

microvascular lineage which is evident in newly formed collaterals in heart[9,11] and

skeletal muscle.[12] Genetic lineage tracing with *Cx40CreER Rosa26^TdTomato* and

ApjCreER Rosa26^TdTomato mice were performed where TdTomato expression could

be conditionally induced, with Tamoxifen, specifically in artery ECs or capillaries

respectively. Tamoxifen was administered to timed pregnant mice at e14.5 when

both MCA and ACA arteries are present (Figure **2A**, **S2G**, **S3G**). Embryos were

analysed at e18.5 or P0 when the pial collateral artery network has completely

formed (Figure **2A**, **S2H**, **S3H**). Whole embryonic brains were immunostained for

Cx40 and segments of collateral arteries between MCA and ACA were analysed for

lineage contribution (Figure **2B-J**).

We assessed the extent of lineage contribution from artery or capillary ECs

towards pial collaterals. As expected, all pre-existing artery ECs were labelled and

traced in *Cx40CreER; Rosa26^{TdTomato}* embryos (Figure **2C**, **D**, **G**, **I**). Whole brain

imaging of the embryo (Figure **2C**) allowed identification of pial collateral arteries

present between MCA and ACA in the watershed (Figure **2D**). Individual collateral

segments as identified by Cx40 immunostaining were analysed for presence of

TdTomato$^+$ artery ECs (Figure **2D**, **G**, **I**). This analysis revealed that 77% of the

Cx40$^+$ collateral area to be TdTomato$^+$ (Figure **2B**). In contrast to this data, a similar

analysis in an independent experiment with *ApjCreER; Rosa26^{TdTomato}* mice (Figure

2E, **F**, **H**, **J**) showed only 31.6% of the collateral area to be TdTomato$^+$, i.e. traced

from capillaries (Figure **2B**). Thus, artery ECs are the major contributors of pial

collateral arteries during development.

Arterial VegfR2 mediates artery tip extension and is essential for pial collateral artery development

We analysed similarities, if any, in the molecular programs used by artery

endothelial cells during collateral development in the heart and brain. Post-injury,

neonatal coronary artery ECs undergo extensive migration and proliferation to build

new coronary collateral arteries in the watershed. The processes of artery cell

migration and proliferation are heavily dependent on Cxcr4[9] and VegfR2[13]

respectively. To assess the functions of Cxcr4 and VegfR2 in pial collateral

development, we deleted the respective genes at e14.5 and assessed the extent of

pial collateral network at e18.5/P0 by whole brain imaging (Figure **3A**). Surprisingly,

deletion of arterial *Cxcr4* did not affect the pial collateral network at e18.5 (Figure **3B**,

control=8, artery specific knockout for *Cxcr4*=13) or in adulthood (Figure **S5**, N=5 per

genotype). This data suggests that a Cxcr4-independent vascular mechanism

facilitates pial collateral development and its maintenance.

We next assessed if pial collateral artery development is dependent on

arterial VegfR2. Whole brain imaging of control and *VegfR2* artery specific knockout

embryos showed differences in collateral connectivity in the watershed, and required

systematic quantification (Figure **3C**-**F**). The extensive connectivity of pial arteries

across the hemisphere (including watershed) provide several possible collateral

paths for the blood to flow. These artery-artery connections are innumerable. We

developed a method that allowed us to compare the pial artery coverage between

the control and *VegfR2* artery EC specific knockout brains quantitatively (Figure **3C**).

Intensity line-profiles were generated across the width of the hemisphere (Figure **3C**

i, **ii**). The identified peaks per line-profile was plotted across the width of the

hemisphere (Figure **3C iii**). The data was fit to a gaussian function and the fitted

curves were used as representative of the arterial coverage per hemisphere in each

mouse (Figure **3C iv**). The average of the fitted curve represented one mouse brain

(grey curve, Figure **3C iv**). This curve represented the artery coverage per

hemisphere. Quantification of pial artery coverage showed significantly different

distributions between control and *VegfR2* knockout brains (KS test, D value=0.569, p

value=2.2066×10^{-141}) (Figure **3E**). Peak arterial coverage was observed at the

centre of the hemisphere and watershed was defined as a region ± 500µm from the

centre (Figure **3E**). The loss of arterial coverage in the watershed was evident in the

knockout (Figure **3E**) and was quantified as area under the curve (Figure **3E**, **F**).

This area under the curve represented the extent of collateral artery network in the

watershed and was significantly lower in brains without arterial *VegfR2* (Figure **3F**,

unpaired t-test, p value= 0.0001). Thus, *VegfR2*-depleted arteries were unable to

synthesize pial collaterals during embryogenesis.

We next investigated the possible mechanisms by which arterial VegfR2

facilitated pial collateral formation. *VegfR2* was deleted from artery ECs at e14.5,

and analyses of pial collateral network was performed at e16.5 (Figure **3G**), a

timepoint when collaterals are beginning to form via artery tip extensions (Figure **1D-**

G). Quantification showed fewer TdTomato[+] collateral arteries (i.e. TdTomato

continuous or Td[con]) without arterial *VegfR2* (Figure **3H**), indicating that initiation of

collateral development at an earlier timepoint, e16.5, also depends on arterial

VegfR2. This VegfR2-dependent collateral phenotype (Figure **3H**) was independent

of artery cell proliferation which was minimal at this stage, for both control and

VegfR2-deleted brains (data not shown). Close analyses of collaterals in control

brains showed that TdTomato[+] artery cells occupied a major length of microvascular

tracks connecting branches of MCA and ACA (Figure **3I**). This observation was in

contrast to *VegfR2*-depleted brains where TdTomato[+] artery cells sparsely populated

the microvascular tracks (Figure **3J**). Thus, arterial VegfR2 regulates pial collateral

formation by promoting artery growth along microvascular tracks.

We investigated if endothelial cells of capillary lineage populate pial collaterals

via Dach1; a transcriptional factor that drives arterialization of capillaries into

coronary arteries during cardiac development in mice.[14] As observed in developing

mouse hearts,[14,15] expression of Dach1 was higher in capillaries as compared to

arteries (Figure **S6A**). *Dach1* was deleted from capillaries at e14.5 and the extent of

pial collateral network was assessed at e18.5 (Figure **S6B**). Interestingly, no

significant difference was observed in the number of Cx40$^+$ collaterals in the *Dach1*

capillary-EC specific knockout brains (Figure **S6C**). This suggests that the vascular

mechanism which leads capillaries to populate pial collaterals is independent of

Dach1 activity.

Pial collaterals do not remodel in adulthood

We next investigated the cellular dynamics associated with fully formed and

mature pial collateral arteries between MCA and ACA in adult mice. For this, we

utilized an extensive longitudinal in vivo imaging paradigm that allowed us to get

single cell resolution of vascular structures. 3-5 months old *Cx40CreER;*

Rosa26^{TdTomato};Tie2::eGFP::Claudin5 mice underwent craniotomy and an optical

window of 4mm was implanted on one of the hemispheres between the cranial

landmarks—bregma and lambda (Figure **4A**, **B**). In these transgenic mice, pre-

existing pial arteries were labelled with TdTomato upon Tamoxifen administration

(Figure **4C**), and constitutively expressing GFP marked all endothelial cells. 5

consecutive doses of Tamoxifen resulted in 95.4% and 86.8% recombination in

proximal and distal arteries respectively, but a lower 65.3% recombination in the pial

collateral regions (Figure **4D**). A lower recombination efficiency in the pial collateral

arteries resulted in a patchy reporter pattern comprising of a mixture of TdTomato$^+$

and TdTomato$^-$ cells (Figure **4D**). Like in developmental stages (Figure **2B**), artery

ECs populate adult pial collaterals to a greater extent as compared to other vascular

lineages. This ultimately allowed us to most accurately track the position of individual

cells within a given collateral artery.

An optical window allows longitudinal imaging. Imaging the collateral zone

(watershed between MCA and ACA) for a month (Figure **4E**), resulted in high quality

images with single cell resolution (Figure **4E-H**). Morphology of the pial vessels

alongside reporter activity in endothelial cells were instrumental in identifying pial

arteries, veins and capillaries (Figure **4E**, **I**). High resolution images of artery

endothelial cells (both TdTomato$^+$ GFP$^+$ and TdTomato$^-$ GFP$^+$) within a collateral

artery segment across many collaterals and optical windows were analysed (Figure

4J, **K**). Data showed that artery endothelial cells do not remodel or reposition

themselves within a given collateral artery segment (Figure **4J-M**). Cellular matrices

representative of the relative position of collateral artery cells suggested that under

physiological conditions, every artery cell within a mature/ adult pial collateral

maintains its contact with the same neighbouring cells (Figure **4L**, **M**). Thus, for the

first time our data, with cellular resolution, demonstrates the static nature of artery

endothelial cells during homeostasis.

Arterial VegfR2 is dispensable for pial collateral maintenance

We next asked if arterial VegfR2 is necessary for the maintenance of adult

pial collaterals using adult mice deficient in arterial *VegfR2*. Using the same in vivo

imaging paradigm mentioned above, pial collateral segments in artery endothelial

cell specific *VegfR2* knockout mice were imaged at cellular resolution and followed

for a month with imaging time-points separated by 2 weeks (Figure **5A**). Longitudinal

imaging through an optical window (Figure **5B**) yielded a map of vascular network

which could be followed for months (Figure **5C**, **D**). Of note, our qualitative

observation suggests that *VegfR2* depleted pial collaterals had smaller diameters.

High magnification images of watershed between MCA and ACA (Figure **5E-G**)

allowed us to analyse individual *VegfR2$^{-/-}$* artery endothelial cells (TdTomato$^+$)

residing on adult collateral arteries (Figure **5H-J**). Long term imaging of the same

collateral regions across many collaterals and mice showed no significant changes in

cellular positions relative to the neighbouring cells (Figure **5K-M** and Figure **S7**).

Thus, arterial VegfR2 is dispensable for maintenance of mature pial collateral

arteries. Together, in this study, we highlight the essential role of arterial VegfR2 in

promoting artery-tip extension and pial collateral development during embryogenesis

(Figure **5N**); a function dispensable in adulthood.

DISCUSSION

In the present study, we investigate artery cell dynamics during collateral artery

formation and maintenance in mouse brain. Firstly, using an artery-specific

molecular marker, we dissect out where and when pial collaterals appear during

embryogenesis. Secondly, we develop a live imaging method of mouse embryos to

provide conclusive evidence that pre-existing arteries grow and connect to build pial

collaterals *de novo*. Thirdly, by combining mouse genetics with whole brain confocal

imaging at cellular resolution, we show an angiogenic vascular receptor VegfR2,

facilitates artery-tip extension on microvessels and aids origin of pial collaterals.

Finally, we show that Cxcr4, a receptor essential for Artery Reassembly and

formation of coronary collateral arteries *de novo*, was dispensable for development

of pial collaterals. Thus, two fundamentally distinct mechanisms contribute to the

origin of coronary and pial collaterals.

Organ-specific mechanisms build collaterals

For the first time, we provide direct proof that a vascular mechanism driven

specifically by artery endothelial cells leads to the origin of pial collaterals. Like in the

heart, this artery specific process is regulated by arterial VegfR2, but is distinct from

Artery Reassembly where pre-existing artery cells dissociate from resident

coronaries, migrate, de-differentiate/proliferate and coalesce to build coronary

collaterals.[9,13] Unlike heart, pial artery endothelial cells provide minimal contribution

via proliferation. Instead, developing pial arteries extend along microvascular tracks

and connect with other artery tips to build new collateral artery paths. These

endothelial cells which migrate or elongate along the microvasculature could arise

from proximal parts of the cerebral arteries. Additionally, arterial Cxcr4 had no impact

on pial collateral formation or its maintenance. This is in contrast to coronary

collateral formation where arterial Cxcr4 drives artery cell migration during Artery

Reassembly. The need for functional collaterals is most during an artery occlusion,

and can significantly impact tissue function. The similarities and differences between

the origin of coronary and pial collaterals could be driven by the metabolic needs of

the tissue. While cardiomyocytes can survive hypoxia for few hours, neurons are

more susceptible to low levels of oxygen.[16,17] The differences in sensitivity to hypoxia

could be the reason for formation of pial collaterals during embryogenesis while

coronary collaterals form only upon arterial occlusion. Furthermore, collateral arteries

found in skeletal muscle,[11,18] intestine,[19] and hind limb[20] can undergo

arteriogenesis.[21] Identifying collateral-building vascular mechanisms across organs

could reveal clinically relevant information and open avenues for therapeutics.

Old arteries build new arteries

We demonstrate the formation of collateral arteries from pre-existing arterial

lineage as observed[9,10,13,22] or proposed[6] earlier. While this manuscript was in

progress, Perovic et al reported that two separate vascular lineages populate pial

collaterals.[10] We also observed that well-formed pial collaterals are majorly

populated by artery cells, and minimally by capillaries. Intra-vital imaging of

embryonic brains reveal that pre-existing artery cells extend along defined

microvascular paths to connect and build pial collaterals. These artery specific

events are executed within hours, are difficult to capture using static images and can

be observed only by live imaging. We cannot exclude the contribution of other artery-

driven mechanisms, such as migration of single artery cells from proximal segments

of cerebral arteries and their incorporation into pre-existing artery tips leading to

further artery elongation. The capillary contribution towards pial collaterals via

arterialization is less likely as deletion of *Dach1*, a regulator of arterialization,[14,23] had

minimal effect on pial collateral network during embryogenesis. We propose that as

artery tips extend and get connected in the watershed, microvascular cells are

"trapped" within pial collaterals. This could generate a mixed lineage distribution[10]

within pial collaterals—both in embryos and adults.

We also observed artery cells migrating towards pre-existing arteries;

coalescence of these single cells with arteries could lead to arterial growth as

observed during Artery Reassembly in heart. Thus, multiple mechanisms could be

actively building pial vasculature during embryogenesis. That being said, possible

contribution from endothelial progenitor cells or peri-vascular cell types (pericytes,

smooth muscle cells) towards pial collateral formation cannot be entirely excluded.

Given that collateral arteries have a unique hemodynamic niche,[24] it would be

interesting to investigate which molecular programs maintain or remodel pial

collaterals in an adult brain.

Distinct mechanism for collateral maintenance

During embryogenesis, pial collaterals are built *de novo*, in an artery-free

zone and remodel as and when blood flows through them. In contrast mature

collaterals remodel in response to differential blood flow pattern and oxygen

demands of surrounding tissue. Postnatal remodelling of pial collaterals is dependent

on recruitment of smooth muscle cells and exhibits tortuous morphology.[24] These

fundamental differences could explain the requirement for distinct molecular

programs for building new collaterals and maintaining old ones. This is evident in our

study; VegfR2 controlled pial artery network during embryogenesis but was

dispensable in adulthood. That being said, arterial VegfR2 may regulate other

aspects of vascular changes those require activation of angiogenic pathways[25] such

as remodelling induced by hypoxia, ischemia or aging. We did not notice any

remodelling events within adult pial collaterals, in presence or absence of arterial

Cxcr4. Together, adult pial collaterals are maintained via mechanisms independent

of two key angiogenic receptors in arteries—VegfR2 and Cxcr4. It will be interesting

to test if adult collateral maintenance is independent of signals originating from

arterial niche. It is possible that surrounding microvasculature plays a greater role in

maintenance of pial arteries and collaterals in adults. Investigating the molecular

identity of the microvasculature surrounding pial collaterals could reveal new

vascular mechanisms key to collateral maintenance.

MATERIALS AND METHODS

Mouse lines

Mouse lines	Source/ References
Rosa26[TdTomato]	The Jackson Laboratory, stock number 007909
Cx40CreER	Miquerol et al. 2015[26]

ApjCreER	Chen et al. 2014[15]
VegfR2 flox	The Jackson Laboratory, stock number 018977
Cx40$^{eGFP/+}$	Miquerol et al. 2004[27]
Tie2::eGFP::Claudin5	Knowland et al. 2014[28], Lutz et al. 2017[29]
Cxcr4 flox	The Jackson Laboratory, stock number 008767
Dach1 flox	Chang et al. 2017[14]

All experiments were performed on mice with mixed strain background. All mice were

housed and bred at the National Centre for Biological Sciences animal facility (in

accordance with the Institutional Animal Ethics Committee). Tamoxifen regimes for

individual experiments are explained in the respective experimental sections (see

below). Data shown in this study is compiled from age-matched males and females

from several litters.

Genotyping primer information

Primer (F, forward; R, Reverse)	Sequence (5' → 3')
Cre F	GCATTACCGGTCGATGCAACGAGTGATGAG
Cre R	GAGTGAACGAACCTGGAAATCAGTGCG
Cre Internal positive control F	CTAGGCCACAGAATTGAAAGATCT
Cre Internal positive control R	GTAGGTGGAAATTCTATCATCC
TdTomato mutant F	CTGTTCCTGTACGGCATGG
TdTomato mutant R	GGCATTAAAGCAGCGTATCC
TdTomato wild-type F	AAGGGAGCTGCAGTGGAGTA
TdTomato wild-type R	CCGAAAATCTGTGGGAAGTC

VegfR2 flox F	CCACAGAACAACTCAGGGCTA
VegfR2 flox R	GGGAGCAAAGTCTCTGGAAA
GFP F	GCACGACTTCTTCAAGTCCGCCATGCC
GFP R	GCGGATCTTGAAGTTCACCTTGATGCC
Cxcr4 flox F	CCACCCAGGACAGTGTGACTCTAA
Cxcr4 flox R	GATGGGATTTCTGTATGAGGATTAGC
Dach1 flox F	CTCCTGAAGATGAGGAGCTCACCCC
Dach1 flox R	AACAATTCTTGTCCTTCACGTGCCC

Visualization of pial arteries in fixed samples

Harvesting whole brains/heads from mouse embryos

Depending on the developmental stages and size of the embryos, pial collateral

arteries were visualized either in whole brains (e18.5 or P0) or using whole heads

(e12.5- e16.5).

e12.5-e16.5: To perform the whole head explant at e12.5-e16.5 stages, the pregnant

females were euthanized as mentioned above. Small punctures were made on the

embryo sac with the help of microforceps. The whole embryo sac was fixed in 4%

Paraformaldehyde (PFA) for 2 hours with gentle shaking at 4°C. Following fixation,

the embryo sac was washed at 4°C with gentle shaking. This washing step was

performed with 1X PBS ad repeated 4 times every 15-20 mins interval. The embryos

were then extracted from the embryo sacs and the heads were removed with the

help of microforceps. Finally, specifically for e16.5, the loose skin on top of the brain

was removed to aid immunostaining.

e18.5: To perform the whole brain explant at e18.5 stage, the timed pregnant

females were euthanized through carbon dioxide overdose and a Y-shape incision

was made on the abdomen. First the embryo sac was dissected out followed by

careful removal of embryos from their sacs. The skull of each embryo was gently

removed using the microforceps. The embryonic brains were scooped out with the

help of a microspatula and fixed in 4% PFA at 4°C for 2.5 hours. Agitation was

avoided at this stage to preserve tissue integrity of embryonic brains. The fixed

embryonic brains are then washed with 1X PBS three times at 15 minutes intervals

with gentle shaking at 4°C.

P0: Neonatal explants at P0 were performed by euthanizing the pups through carbon

dioxide overdose followed by decapitation. The skull was removed using scissors

and microforceps. The brains were scooped out using the microspatula. The brains

were then fixed using 4%PFA at 4°C for 2 hrs followed by three washes with 1X PBS

at an interval of 15 mins.

Whole brain/head immunostaining of mouse embryos

e12.5-e15.5: The embryonic heads (e12.5 - e15.5) were incubated in 0.1% PBST

(0.1% TritonX-100 in 1X PBS) for 2.5 hours to permeabilize the tissue, followed by

incubation with primary antibodies overnight at 4°C with gentle shaking. Heads were

then washed with 1X PBS, changing the buffer every 1-2 hours for 1 day. This step

was followed by incubation with secondary antibodies made in 0.1% PBST. Samples

were then washed every 1-2 hours for 1 day with 1X PBS. Following immunostaining

the brains are immersed in an anti-fade mounting media (from Jackson

ImmunoResearch Laboratories Inc.) and imaged immediately.

<u>e16.5:</u> The embryonic explant at e16.5 is performed similarly. However, for

Tie2::eGFP::Claudin5 genotype, whole heads were imaged within 4 hrs, prior to PFA

fixation to avoid loss of GFP fluorescence signal.

The embryonic and neonatal brains were incubated with primary antibody (made in

1X PBS) overnight, at 4°C with gentle shaking. Samples were then washed with 1X

PBS, changing the buffer every 1-2 hours for 1 day and incubated with Alexa Fluor

conjugated secondary antibodies (made in 1X PBS). Finally, samples were washed

every 1-2 hours for 1 day with 1X PBS and stored in mounting media or imaged

immediately.

Antibodies used are: Cx40 (Catalog number: Cx40-A, Alpha Diagnostics, 1:500

dilution), Dach1 (Catalog number: 10194-1-AP, Proteintech, 1:500 dilution), VegfR2

(Catalog number: AF644, R&D Systems, 1:250). All secondary antibodies were

procured from Invitrogen and used at a dilution of 1:500.

Imaging whole brain/heads of mouse embryos

The mice heads and brains were imaged using Olympus FV-3000 inverted confocal

microscope by placing the sample in a 2-well chamber Catalog number: 154461;

Nunc™ Lab-Tek™ II Chamber Slide™ System; Thermofisher Scientific. Whole

brains and heads were scanned using 4X and 10X air objectives, and digital images

of multiple stacks for each scanned area were captured. The captured images were

processed in ImageJ and stitched together using Adobe Photoshop software.

The embryonic heads at e16.5 was imaged using Olympus FV-3000 5L Upright

microscope. The pre-fixed embryonic heads were attached to a 35mm plastic petri

dish using agarose gel prior to imaging. The whole head imaging was performed

using 4X air objective and watershed images were captured with 20X water

dipping/immersion objective and 1.5X optical zoom with brains immersed in 1X PBS.

<u>Visualization of pial arteries in live mice</u>

In vivo imaging of embryonic pial arteries

<u>Immobilization of live embryo:</u> *Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5*

Pregnant dams with embryos at developmental stages between embryonic day

(e)15.5-e16.5 were anesthetized with a subcutaneous injection of Ketamine-Xylazine

(Ketamine (0.05 mg/kg); Xylazine (0.5 mg/kg)) cocktail. The dams were then placed

in a supine position on a regulated heating pad (TC1000, CWE) that was positioned

on top of a 3D-printed baseplate (as shown in Figure 1B). A small incision was

placed along the midline of the abdomen. With a gentle retraction of the abdominal

skin and wall, a single uterine horn within the abdominal cavity was isolated. The

uterus was surgically opened and a single embryo along with the yolk sac was

isolated and placed into a 35mm petri dish. The embryo still connected to the dam

was immobilized by submerging the bottom part with 1% agarose, as shown in figure

1B. The immobilized embryo is then submerged in dissection media (DMEM/F12 +

10% FBS, Catalog number: 21041025, Thermo Fisher Scientific), and the media is

maintained at 37°C using a custom-made Arduino controlled media heating and

circulating device.

<u>Live embryonic imaging:</u> The setup along with anesthetized dam was placed under

Macro Zoom microscope MVX10-Olympus for large scale imaging and under

FV3000 upright confocal Microscope-Olympus for high resolution imaging using 20X

water dipping objective. In Macro Zoom Microscopy, the entire observable depth was

scanned whilst obtaining videos of 1 minute spanning a duration of 1-2 hours in a

time-lapse manner. In confocal microscopy, the entire observable depth was

scanned with a step size of 1 micron. A total of 4 embryos were imaged under the

Macro zoom microscope and 5 embryos were imaged using confocal microscope.

Out of these embryos, the artery cell dynamics was captured in 2 brains from each

category.

<u>Image processing:</u> Image stacks obtained from Macro zoom microscopy was

registered using ImageJ plugin (*StackReg*) and the registered stack was de-

convolved using ImageJ plugin (*DeconvolutionLab2*) and finally maximum intensity

projection was obtained as shown in figure 1C. The image stacks obtained from the

confocal microscopy were processed similarly except the deconvolution step.

In vivo imaging adult pial arteries

<u>Craniotomy:</u> *Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5 and*

VegfR2^{flox};Cx40CreER; Rosa26^{TdTomato};Tie2::eGFP::Claudin5 adult mice (>2

months) underwent craniotomy to obtain optical access to the pial layer of the brain.

Mice were anesthetized under 2% Isoflurane (Abbott, North Chicago, IL, USA) and

were placed on a stereotaxic mount (Item number 51547; Stoelting, Wood Dale, IL,

USA) with the skull held tightly using ear bars. The mice were kept anesthetized with

2% Isoflurane using a nose-cone. 5mg/kg Dexamethasone in 1X PBS was

administered subcutaneously and the mice kept warm using a heating pad (TC1000, CWE). The eyes of the animals were covered with an eye ointment and the hair above the skull removed using a hair trimmer. The surgical area was thoroughly wiped clean with 70% ethanol. Sterile surgical tools are used for the surgery. A small incision was made using a scalpel blade and the exposed skin was removed. The connective tissue covering the skull in the exposed area was scraped off using scalpel blade and Q-tips. Using a high-speed electrical drill (burr of diameter ~0.25mm), a window of ~4mm diameter was carefully etched on the skull. The skull within the window was eventually detached from the rest of the skull after repeated drilling and carefully lifted using a pair of fine forceps (Catalog number: 11255-20, Dumont #55) without damaging the dura. During the process of drilling the skull was constantly irrigated with cortex buffer (Concentration; in mM: NaCl–125, KCl–5, glucose–10, HEPES–10, $CaCl_2$–2 and $MgCl_2$–2. pH set to 7.35).[30] The exposed dura was aspirated with cortex buffer until there is no bleeding and the window looked clear. An optical plug made up of two 4 mm coverslip and one 5 mm coverslip (CS-5R & CS-4R; Warner Instruments, Hamden, CT, USA) glued together (with NOA 68, Norland Optical Adhesive) was placed on the exposed dura. The optical plug was gently pressed using a toothpick such that the 5mm cover slip was touching the skull bone and then cyano-acrylic glue was applied across the perimeter of the 5mm coverslip attaching it to the skull. At this stage we had an optical window of 4mm. An aluminium head bar with circular hole of 8mm was placed onto the skull aligned to the previously made optical window. Next, dental cement (Catalog number 56877, 3M Relyx U200) was applied over the exposed skull using a sterile toothpick while being careful to not get it on the optical window. Once the dental cement was completely dry, the animal was placed on a warm pad in a clean recovery cage.

Animals undergoing craniotomy were maintained on meloxicam (2 mg/kg) and

enrofloxacin (5 mg/Kg body weight) for 3-4 days until full recovery. Mice were further

allowed to recover for at least 2 weeks post-surgery before the start of any

experiment. Mice with an optical window can be imaged longitudinally for 2-3 months

until the optical window starts to lose its clarity.

Live Adult Imaging: To perform in vivo imaging, mice were anesthetized through

intraperitoneal injection of Ketamine (80mg/Kg body weight) – Xylazine (5mg/Kg

body weight) cocktail. The eyes were covered with an eye ointment and the mice

were placed onto a custom-made baseplate and immobilized using the implanted

metal head plate to this baseplate. A heating pad (37°C) was placed under the

mouse to maintain the body temperature. Subsequently, the baseplate was placed

on a motorized XY-table of an upright confocal microscope (FV3000 upright confocal

Microscope-Olympus). The entire optical window was imaged using 10X air objective

(0.4 NA) and selected pial regions with collateral arteries were imaged with 20X

water immersion objective (0.5 NA). A total of N=3 adult mice per group (control and

VegfR2 artery endothelial specific knockout) were imaged longitudinally for a period

of a month.

Image processing: The 10X image stacks obtained from the confocal microscopy of

the cranial window were registered using ImageJ plugin (*StackReg*) and maximum

intensity projection was obtained. The region of interest from the 20X image stacks

were cropped out slice by slice and stitched in Adobe Photoshop to avoid the

overlaying microvascular network in order to focus on the pial collaterals.

Quantifications

Lineage contribution towards pial collaterals during embryogenesis

Cx40 immunostained pial collateral segments from either *Cx40CreER; Rosa26*^{*Tdtomato*} or *ApjCreER; Rosa26*^{*TdTomato*} brains were imaged at 10X with 2.12 optic zoom. Regions of interest were manually drawn to mark the collateral artery segments. Lineage contribution was determined by quantifying TdTomato$^+$ coverage on the collateral segment. This was performed by thresholding the images with Otsu-Red tool in ImageJ. The area fraction of the TdTomato$^+$ area per collateral segment was calculated using the Measure tool in ImageJ. Contribution of the artery and capillary derived ECs in collaterals was quantified by calculating the percentage of collateral area covered by TdTomato$^+$ signal. 45 collateral segments (of an approximate length or 600-800μm) from 7 *Cx40CreER*-lineage traced brains and 57 collateral segments from 8 *ApjCreER*-lineage traced brains were analysed.

Number of collateral arteries per hemisphere

In *Cx40CreER* TdTomato labelled brains, the collateral number was quantified based on the number of continuous TdTomato$^+$ vessels (Td-continuous) connecting MCA and ACA. Data shown in Figure **3H** was summarized from 2 control and 3 artery-specific VegfR2 knockout. Data shown in Figure **S2H** was summarized from e13.5 (N=3), e14.5 (N=3), e15.5 (N=2), e18.5 (N=2) and P0 (N=5) embryos and neonates. Data shown in Figure **S6C** is quantified by counting number of Cx40 immunostained collateral arteries between MCA and ACA, and is summarized from 4 control and 7 *Dach1* capillary specific knockouts. An unpaired t-test was done to test significance statistically.

Efficiency of Cre recombination

Segments of collateral arteries, proximal arteries (PA) and distal arteries (DA) from

the image stacks taken with 20X objective were obtained from live adult imaging of

Cx40CreER;*Rosa26^{TdTomato}*;*Tie2::eGFP::Claudin5* mice. Tdtomato$^+$ and Tdtomato$^-$,

cells were counted in each artery segment. Each segment was of an approximate

length of 600-800µm.The recombination efficiency was measured as ratio of

TdTomato$^+$ cells and total number of cells in the segment. A total of 10 collateral

segments, 10 PA segments and 10 DA segments from 4 mice were analysed to

obtain the recombination efficiency of Cre. One-way ANOVA was done to test

significance statistically (Tukey test, p=0.0001).

Pial artery coverage in whole brains and in the collateral zone

The image stacks were obtained using confocal imaging of whole embryonic brains

with 10X objective, stitched using ImageJ plugin (stitching) and max intensity

projection (MIP) was obtained. The MIPs were used for further analysis wherein one

complete hemisphere was cropped out and aligned vertically to the major axis of its

own elliptical boundary. Using a custom ImageJ macro, a line of the length of the

major axis was scanned across the width of the image and intensity profile for every

pixel along the horizontal axis was generated. Total number of local maxima (peaks)

for each intensity profile was obtained using the built-in ImageJ macro function

Array.findMaxima (array, tolerance). The tolerance used was 2 times the average

background intensity. The number of peaks thus obtained was plotted across the

width and the plot was fitted to a Gaussian function using Origin(Pro) 2023b. The

fitted curve was used as a representative of the arterial coverage per hemisphere.

The fit curves obtained from various mice were aligned based on the Gaussian

peaks and averaged to get the representative curves for various experimental

groups. We considered a region of 1mm (-0.5mm to +0.5 mm) across the Gaussian peak as the collateral zone (watershed). The area under the curve within this region was used as a measure for the extent of the collateral network. 9 hemispheres from 9 embryos (e18.5) were analysed per group (control and artery-specific knockout for VegfR2). To test the statistical significance of the differences observed in pial artery coverage (Figure **3E**), a Kolmogorov-Smirnov (KS) test was done. D value= 0.569, p value= 2.2066 X 10^{-141}. To test the statistical significance of the area under the curve (pial collateral artery network shown in Figure **3F**), an unpaired t-test was done.

AUTHORS CONTRIBUTION

SD conceived the idea and designed the experiments. SK performed intravital imaging of embryos and adults. NS, SG, VS, MD performed other embryonic experiments. SD, SK, NS, SG wrote the manuscript. All authors contributed to editing the manuscript.

ACKNOWLEDGEMENTS

We thank the following facilities for their technical support: Animal Care and Resource Center at NCBS, Mouse Genome Engineering Facility at NCBS, Central Imaging and Flow Cytometry Facility at NCBS. We sincerely thank Dr. Dritan Agalliu (Columbia University) and Dr. Kristy Red-Horse (Stanford University) for generously sharing *Tie2::eGFP::Claudin5* and *Dach1$^{fl/fl}$* mice respectively.

FUNDING

This work is supported by NCBS/TIFR core funding and DBT/Wellcome Trust India Alliance Intermediate Fellowship to SD (IA/I/20/2/505205). Suraj K is supported by

DBT/Wellcome Trust India Alliance Intermediate Fellowship to SD. NS and VS were

supported by DBT INSPIRE Scholarship for Higher Education (SHE).

DECLARATION OF INTERESTS

None

REFERENCES

1.	Lima, F.O., Furie, K.L., Silva, G.S., Lev, M.H., Camargo, E.C.S., Singhal, A.B.,

Harris, G.J., Halpern, E.F., Koroshetz, W.J., Smith, W.S., et al. (2010). The

pattern of leptomeningeal collaterals on CT angiography is a strong predictor

of long-term functional outcome in stroke patients with large vessel intracranial

occlusion. Stroke *41*, 2316–2322. 10.1161/STROKEAHA.110.592303.

2.	Maas, M.B., Lev, M.H., Ay, H., Singhal, A.B., Greer, D.M., Smith, W.S., Harris,

G.J., Halpern, E., Kemmling, A., Koroshetz, W.J., et al. (2009). Collateral

vessels on CT angiography predict outcome in acute ischemic stroke. Stroke

40, 3001–3005. 10.1161/STROKEAHA.109.552513.

3.	The top 10 causes of death. https://www.who.int/news-room/fact-

sheets/detail/the-top-10-causes-of-death.

4.	Zhang, H., Prabhakar, P., Sealock, R., and Faber, J.E. (2010). Wide genetic

variation in the native pial collateral circulation is a major determinant of

variation in severity of stroke. Journal of Cerebral Blood Flow & Metabolism

30, 923–934. 10.1038/jcbfm.2010.10.

5.	Clayton, J.A., Chalothorn, D., and Faber, J.E. (2008). Vascular endothelial

growth factor-a specifies formation of native collaterals and regulates collateral

growth in ischemia. Circ Res *103*, 1027–1036.

10.1161/CIRCRESAHA.108.181115.

6.   Lucitti, J.L., MacKey, J.K., Morrison, J.C., Haigh, J.J., Adams, R.H., and Faber,

J.E. (2012). Formation of the collateral circulation is regulated by vascular

endothelial growth factor-a and a disintegrin and metalloprotease family

members 10 and 17. Circ Res *111*, 1539–1550.

10.1161/CIRCRESAHA.112.279109.

7.   Lucitti, J.L., Sealock, R., Buckley, B.K., Zhang, H., Xiao, L., Dudley, A.C., and

Faber, J.E. (2016). Variants of Rab GTPase-Effector Binding Protein-2 Cause

Variation in the Collateral Circulation and Severity of Stroke. Stroke *47*, 3022–

3031. 10.1161/STROKEAHA.116.014160.

8.   Chalothorn, D., Clayton, J.A., Zhang, H., Pomp, D., and Faber, J.E. (2007).

Collateral density, remodeling, and VEGF-A expression differ widely between

mouse strains. Physiol Genomics *30*, 179–191.

10.1152/physiolgenomics.00047.2007.

9.   Das, S., Goldstone, A.B., Wang, H., Farry, J., D'Amato, G., Paulsen, M.J.,

Eskandari, A., Hironaka, C E., Phansalkar, R., Sharma, B., et al. (2019). A

Unique Collateral Artery Development Program Promotes Neonatal Heart

Regeneration. Cell *176*, 1128-1142.e18. 10.1016/J.CELL.2018.12.023.

10.   Perovic, T., Hollfinger, I., Mayer, S., Lips, J., Dopatka, M., Harms, C., and

Gerhardt, H. (2023). Pial collaterals develop through mosaic colonization of

capillaries by arterial and microvascular endothelial cells. bioRxiv,

2023.10.09.561542. 10.1101/2023.10.09.561542.

11.   Gabhann, F. Mac, and Peirce, S.M. (2010). Collateral Capillary Arterialization

following Arteriolar Ligation in Murine Skeletal Muscle. Microcirculation *17*,

333–347. 10.1111/J.1549-8719.2010.00034.X.

12.   Jamaiyar, A., Juguilon, C., Wan, W., Richardson, D., Chinchilla, S., Gadd, J.,

Enrick, M., Wang, T., McCabe, C., Wang, Y., et al. (2022). The essential role

for endothelial cell sprouting in coronary collateral growth. J Mol Cell Cardiol

165, 158–171. 10.1016/j.yjmcc.2022.01.005.

13.   Arolkar, G., Kumar, S.K., Wang, H., Gonzalez, K.M., Kumar, S., Bishnoi, B.,

Rios Coronado, P.E., Woo, Y.J., Red-Horse, K., and Das, S. (2023).

Dedifferentiation and Proliferation of Artery Endothelial Cells Drive Coronary

Collateral Development in Mice. Arterioscler Thromb Vasc Biol *43*, 1455–1477.

10.1161/ATVBAHA.123.319319.

14.   Chang, A.H., Raftrey, B.C., D'Amato, G., Surya, V.N., Poduri, A., Chen, H.I.,

Goldstone, A.B., Woo, J., Fuller, G.G., Dunn, A.R., et al. (2017). DACH1

stimulates shear stress-guided endothelial cell migration and coronary artery

growth through the CXCL12-CXCR4 signaling axis. Genes Dev *31*, 1308–

1324. 10.1101/GAD.301549.117/-/DC1.

15.   Chen, H.I., Sharma, B., Akerberg, B.N., Numi, H.J., Kivelä, R., Saharinen, P.,

Aghajanian, H., McKay, A.S., Bogard, P.E., Chang, A.H., et al. (2014). The

sinus venosus contributes to coronary vasculature through VEGFC-stimulated

angiogenesis. Development *141*, 4500–4512. 10.1242/dev.113639.

16.   Scott Budinger, G.R., Duranteau, J., Chandel, N.S., and Schumacker, P.T.

(1998). Hibernation during Hypoxia in Cardiomyocytes ROLE OF

697  MITOCHONDRIA AS THE O 2 SENSOR*. Journal of Biological Chemistry

698  *273*, 3320–3326. 10.1074/jbc.273.6.3320.

699 17. Rink, C., and Khanna, S. (2011). Significance of Brain Tissue Oxygenation and

700  the Arachidonic Acid Cascade in Stroke. https://home.liebertpub.com/ars *14*,

701  1889–1903. 10.1089/ARS.2010.3474.

702 18. Mannion, J.D., Buckman, P.D., Magno, M.G., and Dimeo, F. (1992). Collateral

703  blood flow from skeletal muscle to normal myocardium. Journal of Surgical

704  Research *53*, 578–587. 10.1016/0022-4804(92)90258-2.

705 19. Crissinger, K.D., and Neil Granger, D. (1988). Characterization of Intestinal

706  Collateral Blood Flow in the Developing Piglet. Pediatr Res *24*, 473–476.

707  10.1203/00006450-198810000-00011.

708 20. Kochi, T., Imai, Y., Takeda, A., Watanabe, Y., Mori, S., Tachi, M., and

709  Kodama, T. (2013). Characterization of the Arterial Anatomy of the Murine

710  Hindlimb: Functional Role in the Design and Understanding of Ischemia

711  Models. PLoS One *8*, e84047. 10.1371/JOURNAL.PONE.0084047.

712 21. Gao, Y., Aravind, S., Patel, N.S., Fuglestad, M.A., Ungar, J.S., Mietus, C.J., Li,

713  S., Casale, G.P., Pipinos, I.I., and Carlson, M.A. (2020). Collateral

714  Development and Arteriogenesis in Hindlimbs of Swine After Ligation of

715  Arterial Inflow. Journal of Surgical Research *249*, 168–179.

716  10.1016/J.JSS.2019.12.005.

717 22. He, L., Liu, Q., Hu, T., Huang, X., Zhang, H., Tian, X., Yan, Y., Wang, L.,

718  Huang, Y., Miquerol, L., et al. (2016). Genetic lineage tracing discloses

arteriogenesis as the main mechanism for collateral growth in the mouse

heart. Cardiovasc Res *109*, 419–430. 10.1093/CVR/CVW005.

23. Raftrey, B., Williams, I.M., Rios Coronado, P.E., Fan, X., Chang, A.H., Zhao,

M., Roth, R., Trimm, E., Racelis, R., D'amato, G., et al. (2021). Dach1 Extends

Artery Networks and Protects Against Cardiac Injury. Circ Res *129*, 702–716.

10.1161/CIRCRESAHA.120.318271.

24. Zhang, H., Chalothorn, D., and Faber, J.E. (2019). Collateral Vessels Have

Unique Endothelial and Smooth Muscle Cell Phenotypes. International Journal

of Molecular Sciences 2019, Vol. 20, Page 3608 *20*, 3608.

10.3390/IJMS20153608.

25. Stanca Melincovici, C., Boşca, A.B., Şuşman, S., Mărginean, M., Mihu, C.,

Istrate, M., Moldovan, I.-M., Roman, A.L., and Mihu, C.M. (2018). Vascular

endothelial growth factor (VEGF)-key factor in normal and pathological

angiogenesis. Rom J Morphol Embryol *59*.

26. Miquerol, L., Thireau, J., Bideaux, P., Sturny, R., Richard, S., and Kelly, R.G.

(2015). Endothelial plasticity drives arterial remodeling within The endocardium

after myocardial infarction. Circ Res *116*, 1765–1771.

10.1161/CIRCRESAHA.116.306476.

27. Miquerol, L., Meysen, S., Mangoni, M., Bois, P., van Rijen, H.V.M., Abran, P.,

Jongsma, H., Nargeot, J., and Gros, D. (2004). Architectural and functional

asymmetry of the His–Purkinje system of the murine heart. Cardiovasc Res

63, 77–86. 10.1016/j.cardiores.2004.03.007.

28.   Knowland, D., Arac, A., Sekiguchi, K.J., Hsu, M., Lutz, S.E., Perrino, J.,

742   Steinberg, G.K., Barres, B.A., Nimmerjahn, A., and Agalliu, D. (2014).

743   Stepwise Recruitment of Transcellular and Paracellular Pathways Underlies

744   Blood-Brain Barrier Breakdown in Stroke. Neuron *82*, 603–617.

745   10.1016/j.neuron.2014.03.003.

746   29.   Lutz, S.E., Smith, J.R., Kim, H., Bates, J.M., Gandhi, S.P., and Agalliu, D.

747   (2017). Caveolin1 Is Required for Th1 Cell Infiltration, but Not Tight Junction

748   Remodeling, at the Blood-Brain Barrier in Autoimmune Neuroinflammation.

749   CellReports *21*, 2104–2117. 10.1016/j.celrep.2017.10.094.

750   30.   Holtmaat, A.J.G.D., Trachtenberg, J.T., Wilbrecht, L., Shepherd, G.M., Zhang,

751   X., Knott, G.W., and Svoboda, K. (2005). Transient and Persistent Dendritic

752   Spines in the Neocortex In Vivo. Neuron *45*, 279–291.

753   10.1016/j.neuron.2005.01.003.

Main figures and figure legends

Intra-vital time-lapse imaging setup for visualization of embryonic pial arteries

Genetic lineage tracing
pial artery endothelial cells

Pial collateral arteries develop via extension of pre-existing artery tips
TdTomato (cells derived from pre-exsiting arteries)

Figure 1: Live imaging developing pial vasculature

(**A**) Intra-vital imaging set up to visualize the developing pial vasculature in mouse

embryos. **i** heating pad to maintain body temperature of anaesthetized pregnant

mouse at 37°C. **ii** custom built media heating device which also circulates dissection

media at 37°C **iii** shows macro-zoom or confocal microscopy performed on live,

immobilized embryos.

(**B**) Representative schematic of an embryonic brain. Grey and Red show cerebral

arteries and pial collaterals respectively.

(**C**) Experimental design to genetically label and trace the dynamics of pre-existing

artery endothelial cells in a developing pial layer.

(**D-G**) Images obtained through macro-zoom fluorescent microscopy of e16.5

embryos. In black are cells derived from pre-existing artery endothelial cells. Images

were taken by focusing in and out to capture the entire curvature of the embryonic

brain. **D** shows an entire left hemisphere. **E** represents watershed region spanned

with MCA and ACA artery tips and imaged at time=0 minutes, 20 minutes and 80

minutes respectively. Arrows and arrowheads show growing ends of pre-existing

artery tips from MCA and ACA. Brackets mark regions where collaterals appear. **F**

shows inset in **D**. **E** and **G** show insets in **F**. **G** represents the MCA growth front

imaged at time=0 minutes, 20 minutes and 80 minutes respectively. * shows a

superficial yolk sac vessel which photobleaches with time.

(**H-J**) Images obtained through confocal microscopy from e16 embryos. **H** shows pial

watershed and the cerebral arteries spanning the watershed. **I**, **J** show MCA and

ACA growth fronts respectively imaged at time=0 minutes and 60 minutes. In black

are cells derived from pre-existing artery endothelial cells. Images are maximum

intensity projections from 50 optical slices and capture the entire width of the artery.

Dotted line shows the length of micro-vessel connecting the single artery cell to the

base of the emergent sprout. * represents an internal landmark from where the

length measurements were done.

(**K**) Confocal images of e17 pial layer showing presence of TdTomato[+] pre-existing

artery endothelial cells (arrowheads) on GFP[+] microvascular tracks. Arrowheads

point to TdTomato[+] artery cells (red) on microvascular tracks (green).

(**L**) A model proposing extension of artery tips along microvascular tracks which

leads to pial collateral formation during embryogenesis. Grey arrows point to the

direction of artery tip elongation. MCA, Middle cerebral artery; ACA, Anterior cerebral

artery; e, embryonic day; Tam, Tamoxifen; EC, endothelial cell; YSV, yolk sac

vessel; brch., branch; coll, collateral; MV, microvascular. Scale bars: **D**, **F** 1mm; **E**, **G**

200 µm; **H**, 500 µm; **I**, **J**, **K** 100 µm.

Figure 2: Cellular origin of pial collaterals

(**A**) Experimental design to assess lineage contribution from artery ECs and capillary ECs towards pial collaterals at e18.5/ P0.

(**B**) Graph shows the average percent area of collateral segments covered with TdTomato$^+$ cells.

(**C**) Representative image of whole right hemisphere from *Cx40CreER*; *Rosa26^{TdTomato}* e18.5 embryos. Black shows TdTomato$^+$ arteries.

(**D**) Boxed region of interest with a collateral segment from the watershed area shown in **C**. The pial collateral is lineage traced for artery ECs (red) and immunostained with Cx40 (green).

(**E**) Representative image of whole right hemisphere from *ApjCreER*; *Rosa26^{TdTomato}* e18.5 embryos. Black shows Cx40 immunostaining.

(**F**) Boxed region of interest with a collateral segment from the watershed area shown in **E**. The pial collateral is lineage traced for capillary ECs (red) and immunostained with Cx40 (green).

(**G**, **I**) Collateral segment from **D** is (**G**) immunostained for Cx40 and (**I**) lineage traced with TdTomato.

(**H**, **J**) Collateral segment from **F** is (**H**) immunostained for Cx40 and (**J**) lineage traced with TdTomato. Arrowheads point to TdTomato$^+$ cells from Apj$^+$ lineage on the collateral segment. e, embryonic day, P, postnatal day; MCA, Middle cerebral artery; ACA, Anterior cerebral artery; brch., branch; EC, endothelial cell. Scale bars: **C** and **E**, 500µm; **E** and **F**, 100µm

Figure 3: VegfR2-dependent formation of pial collateral arteries

(**A**) Experimental design to lineage label and trace pre-existing artery endothelial cells in pial layer, with or without arterial *VegfR2* and *Cxcr4* at e18.5/ P0.

(**B**) Representative confocal images of whole brains with and without arterial *Cxcr4*. Black shows pre-existing artery endothelial cells. Control= 8, artery specific knockout for *Cxcr4*= 13.

(**C**) Summary of method for quantifying complex pial artery coverage across the entire brain and in the watershed.

(**D**) Confocal images of whole brains with and without arterial *VegfR2*. Black shows pre-existing artery endothelial cells.

(**E**) Quantification of overall pial artery coverage without arterial *VegfR2* quantified using method shown in **C**. KS test performed on the distribution generated D value= 0.569, p value= 2.2066×10^{-141}.

(**F**) Quantification of pial collateral artery network without arterial *VegfR2* quantified using method shown in **C**. Unpaired t-test, p=0.0001.

(**G-J**) Analysis of e16.5 brains. (**G**) Experimental design. (**H**) Quantification of pial collateral arteries continuously populated with TdTomato⁺ artery cells. (**I**, **J**) Confocal images of pial watershed region showing TdTomato-continuous pial collateral arteries (Td^{con}) in red and microvasculature in green. Arrowheads show part of microvascular track not occupied by TdTomato⁺ artery cell, arrows point to artery tips on microvascular tracks. MCA, Middle cerebral artery; ACA, Anterior cerebral artery; e, embryonic day; P, postnatal day; Tam, Tamoxifen; EC, endothelial cell; Td^{con},

848   TdTomato-continuous; AU, Arbitrary Unit. Scale bars: **B**, 500µm; **D**, 500µm; **I, J**,

849   100µm.

Intra-vital imaging of pial collaterals in adult mice

A Dorsal view of cranial window

B Side view of cranial window

C Cx40CreER; Rosa26^TdTomato
Tie2::eGFP::Claudin5

D TdTomato Recombination efficiency (%)

Cx40CreER; Rosa26^TdTomato Tie2::eGFP::Claudin5
TdTomato (pre-existing artery ECs) eGFP (all ECs)

E watershed

F t=0 days **G** t=14 days **H** t=28 days

I
Pial arteries
Pial veins
PA, proximal artery coll, collateral
DA, distal artery

J t=0 days t=14 days t=28 days

K t=0 days t=14 days t=28 days

L t=0 days t=14 days t=28 days

M t=0 days t=14 days t=28 days

Figure 4: Longitudinal live imaging of adult pial collaterals

(**A**, **B**) Dorsal and side view of the cranial window created on adult mouse head.

(**C**) Experimental design for longitudinal imaging of pial vasculature in adult mice.

(**D**) Quantification of TdTomato reporter recombination in pial collateral arteries and different segments of cerebral arteries. One-way ANOVA, Tukey test, p=0.0001

(**E-K**) Images of pial watershed with pre-existing artery endothelial cells labelled in TdTomato (red) and all endothelial cells labelled in eGFP (green).

(**E**) Confocal images from optical window created for longitudinal imaging of *Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5* mice. The image shows both pial and dural vessels.

(**F-K**) Confocal images of pial collateral arteries from cranial window shown in E, across 3 timepoints, each 2 weeks apart. (**I**) Pial layer segmented out from **E**. red and blue show pial arteries and veins respectively. **J**, **K** are the insets shown in **F-H**.

(**L**, **M**) Cellular distribution of TdTomato+ artery cells and eGFP+ endothelial cells in collateral artery segments shown in **J** and **K** respectively. * represents a reference cell. MCA, middle cerebral artery; ACA, anterior cerebral artery; d, days; Tam, Tamoxifen; PA, proximal artery; DA, distal artery; coll, collateral; brch., branch; EC, endothelial cell. Scale bars: **E**, 1mm; **J**, **K** 25μm

A *VegfR2^{L/L};Cx40CreER; Rosa26^{TdTomato}*
Tie2::eGFP::Claudin5

B 4mm diameter cranial window

C watershed
TdTomato (pre-existing artery ECs)
eGFP (all ECs)

D Pial arteries   Pial veins

MCA-ACA collateral artery

E t=0 days   **F** t=14 days   **G** t=28 days

H t=0 days   **I** t=14 days   **J** t=28 days

K   **L**   **M**

N VegfR2 regulates pial collateral development by artery tip extension

development (embryogenesis)
VegfR2-driven artery-tip extension

maintenance (adulthood)
VegfR2/Cxcr4 independent process

pial collateral arteries

flow from MCA   flow from ACA

artery lineage   non-artery lineage

Figure 5: Analyses of role of arterial VegfR2 in pial collateral maintenance

(**A**) Experimental design to visualize artery cell dynamics in pial layer of adult mice without arterial *VegfR2*.

(**B**) White-field image of cranial window.

(**C**) Confocal image of cranial window created in mice without arterial *VegfR2*. Red shows all artery cells, green shows all endothelial cells.

(**D**) Pial layer segmented out from C shows pial arteries and veins in red and blue.

(**E-G**) Confocal images of collateral artery from three timepoints, each 2 weeks apart.

(**H-J**) Confocal images of collateral artery segments shown in E-G.

(**K-M**) Cellular distribution of TdTomato+ artery cells and eGFP$^+$ endothelial cells in collateral artery segments shown in **H-J**. * represents a reference cell.

(**N**) Schematic summarizing the role of arterial VegfR2 in development and maintenance of pial collateral arteries. While arterial VegfR2 facilitates tip extension of pre-existing arteries during development, it is dispensable for pial collateral maintenance in adulthood. Pial collaterals, both in developing and mature brains, are majorly populated by endothelial cells of artery-lineage. MCA, middle cerebral artery; ACA, anterior cerebral artery; d, days; Tam, Tamoxifen; EC, endothelial cell; brch., branch. Scale bars: **C**, 1mm; **E-G**, 50µm; **H-J**, 20µm.

Supplementary figures and figure legends

Development of cerebral artery trees and pial collaterals in embryonic mouse brains

Supplemental Figure 1: Visualization of pial arteries and collateral arteries

Schematic summarizes key steps of immunostaining methods used to image pial artery network in (**A**) mouse embryonic heads and (**B**) mouse embryonic/neonatal brains with confocal microscope. (**C**) Schematic shows findings from **A** and **B** in one hemisphere. Grey represents cerebral arteries, Red represents collateral artery connections between cerebral arteries. Brackets show watershed area (WSA), MCA appears at e12.5, ACA appears at e13.5, and the foremost pial collateral arteries appear at e15.5. These pial collaterals then increase in number till e18.5/P0. e, embryonic; P, postnatal day; MCA, middle cerebral artery; ACA, anterior cerebral artery.

Supplemental Figure 2: Detection of pial collaterals with whole brain immunostaining

(**A-F**) Representative images of right hemispheres from developing mouse embryos immunostained for Cx40. Cx40 marks all arteries. MCA appears at e12.5, ACA appears at e13.5 and collaterals between MCA and ACA appear at e15.5. Arrowheads point to collateral arteries between MCA and ACA. (**G**) Quantification of Cx40 immunostained brains from different developmental stages reveals that both MCA and ACA have formed by e14.5. (**H**) Quantification of Cx40$^+$ collateral arteries per hemisphere shows that the first pial collaterals appear at e15.5. e, embryonic; P, postnatal day; MCA, middle cerebral artery; ACA, anterior cerebral artery; brch., branch. Scale bars: 500µm

left hemisphere

e12.5
(MCA appears)

e13.5

right hemisphere

e14.5
(ACA appears)

e15.5
(collateral arteries
appear)

e18.5

P0

G MCA and ACA artery trees form by e14.5

H Pial collateral arteries appear at e15.5

Supplemental Figure 3: Detection of pial collaterals with *Cx40* reporter activity

(**A-F**) Representative images of brains from developing *Cx40$^{eGFP/+}$* mouse embryos. Cx40 marks all arteries. MCA appears at e12.5, ACA appears at e14.5 and collaterals between MCA and ACA appear at e15.5. Arrowheads point to collateral arteries between MCA and ACA. (**G**) Quantification of *Cx40$^{eGFP/+}$* brains from different developmental stages reveals that both MCA and ACA have formed by e14.5. (**H**) Quantification of eGFP$^+$ collateral arteries per hemisphere shows that the first pial collaterals appear at e15.5. e, embryonic; P, postnatal day; MCA, middle cerebral artery; ACA, anterior cerebral artery; brch., branch. Scale bars: 500µm

Supplemental Figure 4: Visualization of pial artery cell dynamics on

microvascular tracks (**A**) Confocal image showing watershed region between MCA

and ACA of a live e16 embryo. (**B**) Inset from **A** showing growth of ACA in 60 mins.

Arrowhead point to an artery tip emerging at 60mins from imaging. Arrow points to

single artery cells. (**C**) Confocal image of e16 collateral zone showing single artery

endothelial cell migration towards the pial collateral in 110mins. Dotted line is the

length of micro-vessel between the single cell and the pial collateral. Magenta arrow

shows direction of single artery cell migration. (**D**) Inset from **A** showing presence of

TdTomato$^+$ artery endothelial cells (yellow arrowheads) on eGFP$^+$ microvascular

tracks (magenta arrows). MCA, Middle Cerebral Artery; ACA, Anterior Cerebral

Artery; MV, Microvascular; Td, TdTomato; EC, Endothelial cell. Scale bars: **A**,

500µm; **B-D**, 100µm.

A

Cxcr4^{L/L} .Cx40CreER; Rosa26^{TdTomato}
Tie2::eGFP::Claudin5

artery EC specific deletion of *Cxcr4*
does not impact adult pial collateral artery network

B *Cx40CreER;Rosa26^{TdTomato}*   **C** *Cxcr4^{L/L};Cx40CreER;Rosa26^{TdTomato}*

966   **Supplemental Figure 5: Pial artery coverage without arterial *Cxcr4* (A)**

967   Experimental set up to assess the effect of arterial *Cxcr4* deletion on pial artery

968   coverage in adults. (**B**, **C**) Confocal images of whole adult brains genetically lineage

969   traced for pre-existing artery endothelial cells, (**B**) with or (**C**) without arterial *Cxcr4*.

970   Scale bars: 1mm

A

Dach1 is highly expressed in pial capillary ECs at P0

No effect on collateral development with deletion of *Dach1* from capillary ECs

Supplemental Figure 6: Dach1-independent pial collateral development (**A**)

Dach1 expression in capillaries and arteries in P0 brains. (**B**) Experimental design

showing Tamoxifen injection at e14.5, and analyses of collateral numbers in e18.5

brains. Quantification of (**C**) Cx40$^+$ pial collaterals between MCA and ACA in control

(*Dach1$^{fl/fl}$*) and *ApjCreER*; *Dach1$^{fl/fl}$* (Dach1 cap-EC KO) brains. e, embryonic; SMA,

α-smooth muscle actin, cap-EC, capillary endothelial cell; KO, knockout. Scale bar:

20μm.

Intra-vital imaging of VefR2 deficient pial collaterals in adult mice

VegfR2^{L/L} Cx40CreER; Rosa26^{TdTomato}
Tie2::eGFP::Claudin5
TdTomato (pre-existing artery ECs) eGFP (all ECs)

1000

1001

1002

1003

1004

Supplemental Figure 7: VegfR2-independent maintenance of pial collaterals

Confocal images of the entire cranial window from 3 different mice shows that the overall pial artery network remains unchanged without arterial *VegfR2*. MCA, Middle Cerebral Arteries, ACA, Anterior Cerebral Arteries, Coll, Pial collateral arteries between MCA and ACA. Scale bars: 500µm.