Development of Pial Collaterals by Extension of Pre-existing Artery Tips

Suraj Kumar, Swarnadip Ghosh, Niloufer Shanavas, Vinayak Sivaramakrishnan,

Manish Dwari, Soumyashree Das[1][*]

National Centre for Biological Sciences, Tata Institute of Fundamental Research,

Bengaluru, KA 560065, India

[1] Lead contact

[*] Correspondence: soumyashree@ncbs.res.in

Key Words

Pial Collateral Artery, Artery Endothelial Cell, Vascular Endothelial Growth Factor

Receptor-2, Intra-vital Imaging, Genetic Lineage Tracing

SUMMARY

Pial collaterals provide protection from ischemic damage and improve prognosis of stroke patients. The origin or precise sequence of events underlying pial collateral development is unclear and has prevented clinicians from adapting new vascularization and regeneration therapies. We use genetic lineage tracing and intra-vital imaging of mouse brains at cellular resolution to show that during embryogenesis, pial collateral arteries develop from extension and anastomoses of pre-existing artery tips, in a VegfR2 dependent manner. This process of artery tip extension occurs on pre-determined microvascular tracks. Our data demonstrate that an arterial receptor, Cxcr4, earlier shown to drive artery cell migration and coronary collateral development, is dispensable for formation and maintenance of pial collateral arteries. Our study shows that collateral arteries of the brain are built by a unique mechanism, distinct from that of the heart.

INTRODUCTION

Pial (leptomeningeal) collateral arteries are an extensive network of inter-artery anastomoses found in the pial layer of the brain. In humans, the extent of pial collateral circulation is a predictor of stroke outcome.[1–3] Stroke occurs when blood supply is obstructed to a part of the brain and is one of the leading causes of morbidity worldwide (World Health Organization).[4] Pial collaterals serve as an alternate route of blood flow and alleviate the symptoms of ischemic stroke by bypassing vascular occlusions. Thus, understanding mechanisms which drive the origin and remodelling of pial collateral arteries is of immense clinical relevance.

Results from preclinical studies indicate wide-spread variation in the pial collateral artery network among individuals and organs.[5] These variations are a result of differential expression and function of proteins associated with Vascular endothelial growth factor (Vegf) signalling axis.[6–8] Different mouse strains demonstrate differential Vegfa expression level which corelate with their pial collateral density.[9] Similarly, global overexpression of Vegf ligand increases number of native pial collaterals and improve collateral perfusion post-injury.[6,7] The receptor mediating the function of Vegf ligand during the formation of pial collaterals is Vegf receptor-2 (VegfR2).[7] A global knockdown of Vegf or VegfR2 during embryogenesis and in a time-frame when pial collaterals are expected to form, results in reduction in their number and diameter.[7] Thus, Vegf is a key driver of pial collateral formation and a critical factor which regulates variation in extent of pial vascular network in mice.

The vital role of Vegf in murine pial collateral development was discovered almost a decade ago.[6,7] However, it is still unclear how the various molecular players of Vegf pathway orchestrate the formation or remodelling of pial collateral network. In

mice, it is hypothesized that an *arteriole-specific sprouting angiogenesis like* process

leads to the formation of the foremost pial collaterals.[7] However, molecular evidence

supporting the identity of these sprouting cells (if arteries or capillaries) is missing.

Additionally, the cellular origin or fate of these sprouts is unknown. Finally, the

possibility of cooperation from other vascular processes, such as arterialization,

arteriogenesis or artery-reassembly, in building pial collaterals, needs to be

thoroughly investigated.

In this study, we use genetic lineage tracing in conjunction with live imaging of

pial vasculature of transgenic mouse embryos and adults to understand mechanisms

by which Vegf regulates pial collateral development and maintenance. Our data

indicate that the cellular source of pial collateral arteries is primarily pre-existing

artery endothelial cells (ECs) which build the first network of pial collaterals at

embryonic day (e)15.5 via VegfR2-dependent artery tip extension. The artery tips

extend on pre-determined microvascular tracks which then connect and build pial

collaterals. While arterial VegfR2 was essential for formation of pial collaterals *de*

novo, it was dispensable for its pruning or maintenance through adulthood. Arterial

Cxcr4 (C-X-C chemokine receptor type-4), which is critical for coronary artery

formation in mice[10] was dispensable during pial collateral formation and its

maintenance. Together, our data indicate that artery endothelial cells build collateral

arteries by utilizing distinct vascular mechanisms in heart and brain.

RESULTS

Pial collateral arteries form during embryogenesis

We developed a whole brain imaging technique to reliably visualize all pial

arteries (Figures **S1A** and **S1B**). Due to lack of unique markers, accurately

identifying pial collateral arteries is exclusively dependent on their location. Pial

collateral arteries are artery segments which interconnect the ends of cerebral

arteries and can be seen in the watershed region (Figure **S1B**). To acquire this

spatial information, it was imperative to image the entire pial artery network. Our

technique allowed us to precisely identify pial collateral arteries. We specifically

focused on collateral arteries which connected and received blood flow from Anterior

Cerebral Artery (ACA) and Middle Cerebral Artery (MCA), as the coverage of this

arterial network is most extensive.

We first investigated the developmental stage when pial collaterals first

appeared. Mouse brains were harvested at different developmental time points and

whole brain immunostaining was performed for Connexin-40 (Cx40) protein, a known

marker for mature artery endothelial cell. Analyses of pial arterial network in both

hemispheres revealed the appearance of MCA at embryonic day (e)12.5 followed by

ACA at e13.5 (Figures **S1C**, **S2A** and **S2B**). At e14.5, both cerebral arteries were

present with no arterial connections between them (Figure **S2C**). Subsequently, the

first arterial anastomoses appeared at e15.5 (Figure **S2D**) which then remodelled to

expand in number through birth/postnatal day (P)0 (Figures **S1C** and **S2D**-**S2F**). This

method confirmed that both MCA and ACA are present at e14.5 (Figure **S2G**) and

the foremost pial collaterals appear at e15.5 (Figure **S2H**).

To confirm the above results, we next used a knock-in mouse line (*Cx40^{eGFP/+}*)

where eGFP expression was driven by *Cx40* promoter activity. Brains from different

developmental stages confirmed our findings from Cx40 immunostaining. MCA

appeared at e12.5, ACA appeared at e14.5 and collateral arteries between MCA and

ACA appear at e15.5 (Figures **S3A**-**S3F** and **S1C**). eGFP⁺ cerebral arteries (MCA

and ACA) were formed by e14.5 (Figure **S3G**) and the first eGFP⁺ collateral artery

between MCA and ACA appeared at e15.5 (Figure **S3H**). Thus, using two

independent methods— Cx40 immunostaining and detecting its reporter activity, we

demonstrate that mouse pial collateral arteries form during development, and appear

precisely at e15.5 (Figure **S1C**).

Intra-vital imaging reveals arterial origin of pial collaterals by artery tip

extension

Despite several hypotheses, there is lack of compelling data in support of

vascular mechanisms which drive pial collateral development. Pial collaterals appear

at late embryonic stages. At this developmental stage, rapid vascular changes make

it difficult to distinguish collateral-specific events from overall vascular growth, thus

making it challenging to draw information from fixed brain samples. Additionally, late

embryonic stages cannot be cultured *ex vivo* due to extremely demanding metabolic

and physiological needs. Thus, we custom-developed an imaging paradigm to

perform intra-vital imaging of pial layer at late embryonic stages (e16-e17) (Figure

1A), a developmental stage where the brain is actively building pial collaterals.

Briefly, embryos of the desired developmental time points were dissected out (with

their intact yolk sac) from the anesthetized pregnant dams while still connected to

the mother via their umbilical cord. Embryos were maintained at 37˚C and

immobilized at appropriate orientation to aid imaging. The embryos were subjected

to stereoscopic or confocal imaging for several hours. This imaging set up allowed

visualization of the entire hemisphere— MCA, ACA, and connected collateral arteries

(Figure **1B**).

Pial collaterals appear only after the primary branches of MCA and ACA have

developed (Figure **S1C**). We hypothesized that the branches of these cerebral

arteries contribute towards formation of pial collaterals. To test this hypothesis,

Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5 mice were used to label artery

ECs with TdTomato and all endothelial cells with eGFP. To activate TdTomato,

Tamoxifen was administered at e13.5, when both MCA and ACA are present and

would be labelled in 2 days, [10,11] by e15.5. Imaging of endothelial cells derived from

arteries was performed at e16-e17. This experimental strategy allowed visualization

of cellular dynamics specific to artery cells derived from pre-existing arteries (Figure

1C) and excluded the possibility of capturing other vascular processes.

We performed intra-vital time-lapse imaging of pial arteries and investigated if

pial collaterals originate from pre-existing artery endothelial cells. We focused on the

watershed region: an initial artery-free region which spans the distal ends of both

arterial trees (MCA and ACA) and is later occupied by pial collaterals (Figures **S1C**

and **1D-1F**). Stereoscopic images of e16.5 brains revealed that branches from MCA

and ACA extend (arrowheads, Figure **1E**) and anastomose to build collaterals in the

pial watershed (brackets, Figure **1E**). Additionally, TdTomato expression in artery

tips (orange arrow, Figure **1E**) and collaterals (green arrow, Figure **1E**) increased

over time. These changes were executed within a short span of 80 minutes, and

could not have been captured in fixed brain samples.

We next investigated artery cell dynamics associated with the growth fronts of

the two cerebral arteries (Figures **1F** and **1G**). At e16.5, the distal branches of the

developing MCA extended and anastomosed to build an artery network *de novo*

(arrowheads, Figure **1G**). This is the first evidence demonstrating that pre-existing

cerebral artery tips build pial collaterals. We then performed intra-vital time-lapse

confocal imaging of brains at e16, a stage prior to when we observed collateral

artery networks being built *de novo* (Figures **1E** and **1G**). Confocal imaging at e16

provided the necessary cellular resolution to quantitatively assess artery-tip

extensions from MCA and ACA branches (Figures **1H**-**1J**). As expected, we clearly

observed artery tip extension of MCA branch (Figures **1H** and **1I**) and the ACA

branch (Figures **1H** and **1J**) within an hour. Additionally, we noted several artery-

specific movements within the pial layer. This included the appearance of single

recombined artery endothelial cells (arrows, Figures **1K** and **1L**), movement of these

single recombined artery cells towards an artery (Figures **1K**, **1M** and **S4A**), and

emergence of arterial sprouts (arrowheads, Figures **1K**, **1L and S4B**-**S4D**).

Together, we provide direct evidence for the vascular origin of pial collaterals— pre-

existing MCA/ACA artery tips extend and connect with each other to build pial

collaterals *de novo*.

Artery tip extension occurs on pre-determined microvascular tracks

We next investigated if other endothelial cell sub-types participate in, and/or

facilitate, pial artery tip extensions during collateral development. To investigate the

same, *Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5* mouse line was used.

Tie2::eGFP::Claudin5 is a transgenic mouse strain which expresses a fusion protein

of eGFP with Claudin5 (a tight junction protein present in all ECs in the central

nervous system). This eGFP-Claudin5 protein expression is under the control of *Tie2*

promoter/enhancer, making it visible in all endothelial cells.[12,13] Thus, in *Cx40CreER*;

Rosa26^{TdTomato}; Tie2::eGFP::Claudin5 mice, TdTomato marked artery ECs while

eGFP marked all ECs. Live imaging of pial watershed showed that all artery cell

movements, almost always, occurred along microvascular (MV) tracks which

appeared structurally more organized/tubular than the surrounding capillaries

(arrowheads, Figures **2A**, **2B** and **S5**). These MV tracks were distinct from the

surrounding capillary plexus in that they formed direct connections between two

arteries, but were not necessarily entirely populated by TdTomato$^+$ artery derived

endothelial cells (Figure **2B**). Additionally, our data show that these MV tracks

originated from Apj$^+$ capillary/venous endothelial cells (Figures **S6A-S6H**), expressed

microvascular marker Endomucin (Figures **S6I-S6L**) and lacked arterial marker Cx40

(Figures **S6M-S6O**). Together, our data provide support for the non-arterial nature of

the MV tracks (Figure **S6P**), and hence, the intermixing of TdTomato$^+$ and

TdTomato$^-$ cells, are not due to uneven Tamoxifen recombination of *Cx40CreER*. We

hypothesized that development of MV tracks precedes artery tip extensions and

tested two potential mechanisms by which MV tracks could assist artery tip

extensions during pial collateral development (Figure **2C**). Firstly, complete formation

of MV tracks between two arteries at e16 could be a pre-requisite for initiation of

artery tip extension, as often observed at e16. (Figures **2Ci-iv**). Alternatively, MV

tracks could appear first, but continuously grow/remodel to allow artery tip extension

(Figures **2Ci**, **v**, **vi**) and subsequent pial collateral formation (Figure **2Civ**). If the

former was true, the number/length of MV tracks in a given watershed would remain

unchanged between e16 and e16.5, however the TdTomato coverage of these MV

tracks would be higher at e16.5 (Figures **2Cii-iii**). If the latter mechanism was true, at

e16.5, the length of MV tracks would be higher, and the TdTomato coverage of these

MV tracks would increase or remain unchanged (Figures **2Cv**, **vi**). To test these two

potential mechanisms, *Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5* mouse

line was used, where MV tracks can be clearly visualized with eGFP reporter

(Figures **2B** and **S5**). Confocal imaging of e16 (Figure **2D**) and e16.5 (Figure **2I**)

whole brains, at single cell resolution allowed examination of watershed regions at

both developmental time-points (Figures **2E-2H** and **2K-2N** respectively). Next,

robust quantification of MV tracks for each watershed region was performed (Figures

2H, **2N** and **2J**). Amongst eGFP⁺ vessels, an MV track has a distinct morphology

from the surrounding capillary plexus and can be identified as eGFP⁺ tubular

structures connecting the distal artery branches (Figures **2F** and **2L**). We identified

an eGFP⁺ vessel as a MV track if 1) it is a continuous tubular structure and 2) is

connected to a TdTomato⁺ artery at least at one end. These MV tracks were traced

on multiple z-slices, using Simple Neurite Tracer (Figures **2H** and **2N**), and next, the

extent of TdTomato coverage was calculated for total length of MV tracks per

watershed (Figures **2H**, **2N** and **2O**). Quantifications show that while the total length

of MV tracks was increased by 1.5 fold in e16.5 (Figures **2H**, **2N** and **2J**), the

TdTomato coverage of these MV tracks did not change between e16 and e16.5

brains (Figures **2H**, **2N** and **2O**). These results support the second mechanism

(Figures **2C-i→v→vi→iv**); thus, suggesting that pial collateral development is

accomplished by simultaneous growth of MV tracks and artery-tips on them.

Artery ECs are the major cellular contributors of pial collaterals during

embryogenesis

We next assessed the vascular-lineage composition of pial collaterals at a

later stage when the collateral artery network has fully formed. This would allow us

to: 1) determine if the artery cell contribution to nascent pial collaterals observed at

e16.5 (Figure **1E**) persisted in later stages and 2) check contribution from

microvascular lineage which is evident in newly formed collaterals in heart[10,14] and

skeletal muscle.[15] Genetic lineage tracing with *Cx40CreER; Rosa26^TdTomato* and

ApjCreER; Rosa26^TdTomato mice were performed independently, where TdTomato

expression could be conditionally induced, with Tamoxifen, specifically in artery ECs

or capillaries respectively. Tamoxifen was administered to timed pregnant dams at e14.5 when both MCA and ACA arteries are present (Figures **3A**, **S2G** and **S3G**). Embryos were analysed at e18.5 or P0 when the pial collateral artery network has completely formed (Figures **3A**, **S2H** and **S3H**). Whole embryonic brains were immunostained for Cx40 and segments of collateral arteries between MCA and ACA were analysed for lineage contribution (Figures **3B**-**3J**).

We assessed the extent of lineage contribution from artery or capillary ECs towards pial collaterals in two separate experiments. As expected, all pre-existing artery ECs were labelled and traced in *Cx40CreER; Rosa26^{TdTomato}* embryos (Figures **3C**, **3D**, **3G** and **3I**). Whole brain imaging of the embryo (Figure **3C**) allowed identification of pial collateral arteries present between MCA and ACA in the watershed (Figure **3D**). Individual collateral segments as identified by Cx40 immunostaining were analysed for presence of TdTomato$^+$ artery ECs (Figures **3D**, **3G** and **3I**). This analysis revealed that 77% of the Cx40$^+$ collateral area to be TdTomato$^+$ (Figure **3B**). In contrast to this data, a similar analysis in an independent experiment with *ApjCreER; Rosa26^{TdTomato}* mice (Figures **3E**, **3F**, **3H** and **3J**) showed only 31.6% of the collateral area to be TdTomato$^+$, i.e. traced from capillaries (Figure **3B**). Thus, artery ECs are the major contributors of pial collateral arteries during development.

Arterial VegfR2 mediates artery tip extension and is essential for pial collateral artery development

We analysed similarities, if any, in the molecular programs used by artery endothelial cells during collateral development in the heart and brain. Post-injury, neonatal coronary artery ECs undergo extensive migration and proliferation to build new coronary collateral arteries in the watershed. The processes of artery cell

migration and proliferation are heavily dependent on Cxcr4[10] and VegfR2[16]

respectively. To assess the functions of Cxcr4 and VegfR2 in pial collateral

development, we deleted the respective genes at e14.5 and assessed the extent of

pial collateral network at e18.5/P0 by whole brain imaging (Figure **4A**). Surprisingly,

deletion of arterial *Cxcr4* did not affect the pial collateral network at e18.5 (Figures

4B, S7A and **S7B**) or in adulthood (Figures **S7C**-**S7E**). This data suggests that a

Cxcr4-independent vascular mechanism facilitates pial collateral development and

its maintenance.

We next assessed if pial collateral artery development is dependent on

arterial VegfR2. Whole brain imaging of control and *VegfR2* artery specific knockout

embryos showed differences in collateral connectivity in the watershed, and required

systematic quantification (Figures **4C**-**4F**). The extensive connectivity of pial arteries

across the hemisphere (including watershed) provides several possible collateral

paths for the blood to flow. These artery-artery connections are innumerable. We

developed a method that allowed us to compare the pial artery coverage between

the control and *VegfR2* artery EC specific knockout brains quantitatively (Figure **4C**).

Intensity line-profiles were generated across the width of the hemisphere (Figures

4C i, **ii**). The identified peaks per line-profile was plotted across the width of the

hemisphere (Figure **4C iii**). The data was fit to a Gaussian function and the fitted

curves were used as representative of the arterial coverage per hemisphere in each

mouse (Figures **4C iv**). The average of the fitted curves represented one genotype

(grey curve, Figure **4C iv**). Quantification of pial artery coverage showed significantly

different distributions between control and *VegfR2* knockout brains (KS test, D

value=0.569, p value=2.2066×10^{-141}) (Figure **4E**). Peak arterial coverage was

observed at the centre of the hemisphere and watershed was defined as a region ±

500μm from the centre (Figure **4E**). The loss of arterial coverage in the watershed

was evident in the knockout (Figure **4E**) and was quantified as area under the curve

(Figures **4E** and **4F**). This area under the curve represented the extent of collateral

artery network in the watershed and was significantly lower in brains without arterial

VegfR2 (Figure **4F**). Thus, depletion of *VegfR2* from arteries reduced the extent of

pial collaterals formed during embryogenesis.

We next investigated the possible mechanisms by which arterial VegfR2

facilitated pial collateral formation. *VegfR2* was deleted from artery ECs at e14.5,

and analyses of pial collateral network was performed at e16.5 (Figure **4G**), a

timepoint when collaterals are beginning to form via artery tip extensions (Figures

1D-1G). A thorough analyses of the pial watershed at e16.5 was performed to

assess the extent of MV tracks and TdTomato (artery-lineage) coverage (Figures

S8A and **S8B**). Close analyses of collaterals in control brains showed that

TdTomato[+] artery cells occupied a major length of microvascular tracks connecting

branches of MCA and ACA (Figure **4H**). In contrast to the control brains, VegfR2-

depleted brains showed reduced TdTomato[+] artery cell coverage of the

microvascular tracks (Figures **4I, 4J,** and **S8B**). Of note, without arterial VegfR2, the

length of MV tracks also reduced (Figure **S8C**), without affecting the density of

capillaries (Figures **S8D**, **S8E**). Quantification of collateral arteries showed fewer

TdTomato contiguous or Td[con] collateral arteries, without arterial VegfR2 (Figure **4K**).

This indicated that initiation of collateral development at an earlier timepoint, e16.5,

also depends on arterial VegfR2. This VegfR2-dependent collateral phenotype

(Figures **4J** and **4K**) was independent of artery cell proliferation which was minimal

at this stage, for both control and VegfR2-deleted brains (data not shown).

Interestingly, this was in contrast to the cell cycle state of ischemic coronaries, which

is 10-fold higher than pial arteries.[10,16] Together, arterial VegfR2 regulates pial

collateral formation by promoting artery extensions along microvascular tracks.

We next investigated if the presence of microvascular cells on pial collaterals

(Figure **3B**) is a result of arterialization, i.e., differentiation of capillary cells into artery

endothelial cells. To address this question, we assessed the potential role of Dach1,

a transcriptional factor, which drives arterialization of cardiac capillary cells into

coronary artery endothelial cells, during development and post-injury.[17] As observed

in developing mouse hearts,[18,19] expression of Dach1 was higher in pial capillaries

as compared to pial arteries (Figure **S9A**). Next, *Dach1* was deleted from capillaries

using *ApjCreER* at e14.5, when both cerebral arteries are present but a pial

collateral network is absent. Whole brains were then analysed for pial collateral

network at e18.5 (Figure **S9B**). We observed a robust deletion of *Dach1* from

capillary ECs (Figures **S9C**-**S9D**). However, no significant difference was observed

in the number of Cx40$^+$ collaterals in the *Dach1* capillary-EC specific knockout brains

(Figure **S9E**). Thus, in contrast to coronary artery development,[19] the formation of

pial collaterals during embryogenesis is independent of capillary Dach1 activity.

Pial collaterals do not remodel in adulthood

We asked the question if arterial VegfR2 could modulate neonatal pruning

events. To address this, *VegfR2* was deleted in arteries at P0, and whole brain

imaging was performed at P2, P7 and P14 (Figure **S10A**). The arterial deletion of

VegfR2 was robust at P7 (Figures **S10B**-**S10C**). Analyses of whole P2 (Figures

S10D-**S10G**), P7 (Figures **S10H**-**S10K**) and P14 (Figures **S10L**-**S10O**) hemispheres

revealed no significant differences in arterial coverage with depletion of arterial

VegfR2 (Figures **S10G, S10K** and **S10O**). Thus, arterial VegfR2 is dispensable for

neonatal pial artery and collateral artery pruning.

We next investigated the cellular dynamics associated with fully formed and

mature pial collateral arteries between MCA and ACA in adult mice. For this, we

utilized an extensive longitudinal in vivo imaging paradigm that allowed us to get

single cell resolution of vascular structures. 3-5 months old *Cx40CreER;*

Rosa26^{TdTomato}; Tie2::eGFP::Claudin5 mice underwent craniotomy and an optical

window of 4mm was implanted on one of the hemispheres between the cranial

landmarks—bregma and lambda (Figures **5A** and **5B**). In these transgenic mice, pre-

existing pial arteries were labelled with TdTomato upon Tamoxifen administration

(Figure **5C**), and constitutively expressing eGFP marked all endothelial cells. 5

consecutive doses of Tamoxifen resulted in 95.4% and 86.8% recombination in

proximal and distal arteries respectively, but a lower 65.3% recombination in the pial

collateral regions (Figures **5D** and **S11**). A lower recombination efficiency in the pial

collateral arteries resulted in a patchy reporter pattern comprising of a mixture of

TdTomato$^+$ and TdTomato$^-$ cells (Figure **5D**). Like in developmental stages (Figure

3B), artery ECs populate adult pial collaterals to a greater extent as compared to

other vascular lineages. This ultimately allowed us to most accurately track the

position of individual cells within a given collateral artery.

An optical window allows longitudinal imaging. Imaging the collateral zone

(watershed between MCA and ACA) for a month (Figure **5E**), resulted in high quality

images with single cell resolution (Figures **5E-5H**). Morphology of the pial vessels

alongside reporter activity in endothelial cells were instrumental in identifying pial

arteries, veins and capillaries (Figures **5E** and **5I**). High resolution images of artery

endothelial cells (both TdTomato$^+$ eGFP$^+$ and TdTomato$^-$ eGFP$^+$) within a collateral

artery segment across many collaterals and optical windows were analysed (Figures

5J and **5K**). Data showed that artery endothelial cells do not remodel or reposition

themselves within a given collateral artery segment (Figures **5J**-**5M**). Cellular

matrices representative of the relative position of collateral artery cells suggested

that under physiological conditions, every artery cell within a mature/ adult pial

collateral maintains its contact with the same neighbouring cells (Figures **5L** and

5M). Thus, surprisingly, the cellular organization within pial collaterals and the overall

collateral structure was extremely stable. Together, our data with cellular resolution,

demonstrate the static nature of artery endothelial cells during homeostasis.

To investigate if cells of pial collaterals reposition in response to hypoxic

conditions, we next subjected the mice to 7% atmospheric oxygen. We utilized a

similar imaging paradigm as shown in Figure **5**. Longitudinal imaging of cranial

windows allowed visualization of pial vasculature at cellular resolution (Figures

S12A-**S12B**). *Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5* mice were used to

label/trace pre-existing arteries, followed by exposure to low (7%) atmospheric

oxygen (Figure **S12C**). High resolution long-term imaging of every cranial window

revealed that diameter of collateral arteries between MCA and ACA increased with

low tissue oxygen (Figures **S12D**-**S12F**). This increase in diameter was not due to an

increase in artery cell number (Figure **S12G**), but because of an increase in size of

individual artery cell (Figures **S12H**-**S12J**). Together, artery endothelial cells within

pial collateral arteries do not reorganize, even under hypoxic conditions.

Arterial VegfR2 is dispensable for pial collateral maintenance

We next asked if arterial VegfR2 is necessary for the maintenance of adult

pial collaterals using adult mice deficient in arterial *VegfR2*. Using the same in vivo

imaging paradigm mentioned above, pial collateral segments in artery endothelial

cell specific *VegfR2* knockout mice were imaged at cellular resolution and followed

for a month with imaging time-points separated by 2 weeks (Figure **6A**). Longitudinal

imaging through an optical window (Figure **6B**) yielded a map of vascular network

which could be followed for months (Figures **6C** and **6D**). Of note, our qualitative

observation suggests that *VegfR2* depleted pial collaterals had smaller diameters.

High magnification images of watershed between MCA and ACA (Figures **6E**-**6G**)

allowed us to analyse individual $VegfR2^{-/-}$ artery endothelial cells (TdTomato$^+$)

residing on adult collateral arteries (Figures **6H**-**6J**). Similar to wild-type (Figure **S11**),

VegfR2 deleted proximal arteries showed very high TdTomato recombination (Figure

S13A). Long term imaging of the same collateral regions across many collaterals

and mice showed no significant changes in cellular positions relative to their

neighbouring cells (Figures **6K**-**6M** and **S13B**). Thus, arterial VegfR2 is dispensable

for maintenance of mature pial collateral arteries. Together, in this study, we

highlight the essential role of arterial VegfR2 in promoting artery tip extension and

pial collateral development during embryogenesis (Figure **6N**); a function

dispensable in adulthood.

DISCUSSION

In the present study, we investigate artery cell dynamics during collateral artery

formation and maintenance in mouse brain. Firstly, using an artery-specific

molecular marker, we dissect out where and when pial collaterals appear during

embryogenesis. Secondly, we develop a live imaging method in mouse embryos to

provide conclusive evidence that pre-existing arteries grow on pre-determined

microvascular tracks which then connect to build pial collaterals *de novo.* Thirdly, by

combining mouse genetics with whole brain confocal imaging at cellular resolution,

we show that an angiogenic vascular receptor, VegfR2, facilitates artery-tip

extension on microvascular tracks and aids origin of pial collaterals. Finally, we show

that Cxcr4, a receptor essential for artery reassembly and formation of coronary

collateral arteries *de novo*, is dispensable for development of pial collaterals. Thus,

two fundamentally distinct mechanisms contribute to the origin of coronary and pial

collaterals.

Organ-specific mechanisms build collaterals

In this study, we provide direct support that a vascular mechanism driven

specifically by artery endothelial cells leads to the origin of pial collaterals. Like in the

heart, this artery specific process is regulated by arterial VegfR2, but is distinct from

artery reassembly, where pre-existing artery cells dissociate from resident

coronaries, migrate, de-differentiate/proliferate and coalesce to build coronary

collaterals.[10,16] Instead, developing pial arteries extend along microvascular tracks

and connect with other artery tips to build new collateral artery paths. Furthermore, in

heart, coronary collateral formation is regulated by arterial Cxcr4 which drives artery

cell migration during artery reassembly. In contrast, arterial Cxcr4 had no impact on

pial collateral formation or its maintenance. Unlike heart, pial artery endothelial cells

contribute minimally via proliferation. Evidence for negligible artery cell proliferation

within the pre-existing pial arteries prompts us to hypothesize that formation of pial

collaterals during embryogenesis is minimally dependent on proliferation of pre-

existing artery ECs. That being said, the underlying mechanism leading to extension

of distal artery tips remain unclear. It is possible that distal artery endothelial cells

experience a forward push from the proximal segments of the artery. This forward

pressure could be generated due to incorporation of new cells. Accommodating

these additional cells within an artery would require further cellular organization

throughout the length of the artery, and could lead to distal extension of the artery

tips. Finally, the impact of blood flow on cellular organization within pre-existing

arteries and/or its extension, remains an open question for future investigation.

The need for functional collaterals is most during an ischemic condition such

as an artery occlusion, and can significantly impact tissue function. The differences

in factors which regulate origin of coronary and pial collaterals, are potentially driven

by the metabolic needs of these tissues. While cardiomyocytes can survive hypoxia

for few hours,[20] neurons are more susceptible to low levels of oxygen.[21] Higher

hypoxia sensitivity of brain tissue may lead to formation of pial collaterals during

embryogenesis as a protective measure, while coronary collaterals form only upon

induction of an arterial occlusion. Furthermore, collateral arteries in skeletal

muscle,[15,22] intestine,[23] and hind limb[24] can also undergo arteriogenesis.[25] Identifying

collateral-building vascular mechanisms across organs could reveal clinically

relevant information and open avenues for therapeutics.

Arteries and capillaries collaborate to build collaterals

We demonstrate the formation of collateral arteries, in real time, from pre-

existing arterial lineage as observed[10,11,16,26] or proposed[7] earlier. While this

manuscript was in progress, Perovic et al reported that two separate vascular

lineages populate pial collaterals.[11] We also observed that well-formed pial

collaterals are majorly populated by artery cells, and minimally by capillaries. Intra-

vital imaging of embryonic brains reveal that pre-existing artery cells extend along

defined microvascular paths to connect and build pial collaterals. These artery

specific events are executed within a short span, are difficult to capture using static

images and can be observed only by live imaging.

We cannot exclude the contribution of other artery-driven mechanisms. We

observed single recombined artery cells migrating towards pre-existing arteries;

coalescence of these cells with arteries could lead to arterial growth. Alternatively,

the migration and incorporation of single artery cells at the proximal ends of the

artery could indirectly contribute towards distal elongation of arteries. The dynamics

of single artery cell movements could also establish the branching pattern of the pial

artery network; the precise mechanisms for which are unknown.

We show that pre-existing artery tip extensions occur along a distinct

supporting structure. In the heart, single artery endothelial cells migrate by hitch-

hiking on to the capillary networks during the formation of new collaterals. On the

contrary, our study in the embryonic brains reveals the presence of a unique (Tie2$^+$)

vascular segment on which the arteries extend. These microvascular tracks are

distinct from the surrounding capillary mesh in that they anastomose ACA and MCA

arterial branches but do not show any artery specification. We speculate that the

microvascular tracks form during the specific time window (e16-e17) which coincides

with the period when majority of collaterals appear. Thus, these tracks may play a

crucial role in pial collateral formation. Based on our Apj lineage data, we

hypothesize that the cellular origin of these microvascular tracks is of capillary in

nature. However, we cannot completely exclude the possibility that some tracks

(guiding vessels for artery tip extensions) could also originate from mixed lineages.

Thus, though capillaries do not massively contribute to the ECs of the pial collateral,

they could create/remodel into MV tracks, to indirectly facilitate development of pial

collaterals. Elucidating the underlying mechanism governing MV track formation is

an exciting avenue for future investigation. Additionally, determining their precise

molecular identity will allow creating genetic tools to assess if MV tracks are

indispensable for development of pial collaterals.

Our data about pre-defined microvascular tracks guiding the growth of pial

arteries suggest that the microvasculature could shape the artery network of the

brain. However, the microvascular contribution towards pial collaterals via

arterialization is less likely, as deletion of *Dach1*, a regulator of arterialization in

heart,[17,19] had minimal effect on pial collateral network during embryogenesis. This

data contrasts the results from heart studies where deletion of *Dach1* from all

endothelial cells using the *Tie2Cre* driver, had a massive impact on coronary artery

development during mouse embryogenesis.[19] Thus, we propose that as artery tips

extend and get connected in the watershed, microvascular cells are "trapped" within

pial collaterals. This could generate a mixed lineage distribution[11] within pial

collaterals—both in embryos and adults. Together, multiple mechanisms could be

actively building pial vasculature during embryogenesis. That being said, possible

contribution from endothelial progenitor cells or peri-vascular cell types (pericytes,

smooth muscle cells) towards pial collateral formation cannot be entirely excluded.

Distinct mechanism for collateral maintenance

We observed Cx40$^-$ ECs co-existed with Cx40$^+$ artery cells within a single

adult collateral segment. The source of Cx40$^-$ cells within a mature pial collateral is

unclear. Our data explicitly show that unlike pial capillaries, artery ECs (Cx40$^+$ or

Cx40$^-$) within adult pial collaterals preserve their cellular location and appear to be

static structures, at least during homeostasis. Thus, it is less likely that (Cx40$^-$)

microvascular cells such as capillary ECs would get incorporated into the collaterals.

We speculate that the (TdTomato⁻) Cx40⁻ cells are artery ECs of a different

molecular origin and/or function. Determining the source or need for such molecular

heterogeneity could be key to identifying special arterial roles in context of injury

such as hypoxia or cerebral artery occlusions.

Pial collateral arteries undergo outward remodelling when subjected to

hypoxia or ischemia.[27] Studies show that artery endothelial cells of pial collaterals

proliferate to support this outward remodelling.[11,27] In contrast, our longitudinal

imaging data provide evidence that the number (or position) of artery cells within a

pial collateral segment does not change in response to hypoxia. It is possible that a

longer exposure to hypoxia or if allowed to recover from these hypoxic conditions,

the pial collateral artery cells would undergo proliferation leading to an increase in

collateral diameter. Thus, cellular response to ischemia or hypoxia could be a multi-

layered phenomenon, involving more than one mechanism.

Given that collateral arteries have a unique hemodynamic niche,[28] it would be

interesting to investigate which molecular programs maintain or remodel pial

collaterals in an adult brain. During embryogenesis, pial collaterals are built *de novo*,

in an artery-free zone and remodel as and when blood flows through them. In

contrast, mature collaterals remodel in response to differential blood flow patterns

and oxygen demands of their surrounding tissue. Postnatal remodelling of pial

collaterals is dependent on recruitment of smooth muscle cells and exhibits tortuous

morphology.[28] These fundamental differences could explain the requirement for

distinct molecular programs for building new collaterals and maintaining old ones.

This is evident in our study; VegfR2 controlled pial artery network during

embryogenesis but was dispensable in adulthood. That being said, arterial VegfR2

may regulate other aspects of vascular changes those require activation of

angiogenic pathways,[29] such as remodelling induced by hypoxia, ischemia or aging.

We did not notice any remodelling events within adult pial collaterals, in presence or

absence of arterial Cxcr4. Together, adult pial collaterals are maintained via

mechanisms independent of two key angiogenic receptors in arteries—VegfR2 and

Cxcr4. It will be interesting to test if adult collateral maintenance is independent of

signals originating from an arterial niche. It is possible that surrounding

microvasculature plays a more prominent role in maintenance of pial arteries and

collaterals in adults. Investigating the molecular identity of the microvasculature

surrounding pial collaterals could reveal new vascular mechanisms key to collateral

maintenance.

LIMITATIONS OF THE STUDY

The experimental designs of this study do not allow us to exclude the contribution of

other cellular mechanisms associated with pial collateral formation during

embryogenesis. Additionally, we are unable to comment on the molecular nature of

artery cell heterogeneity observed in adult pial collaterals and its role (if any) on

recovery from ischemia.

MATERIALS AND METHOD

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-Cx40	Alpha Diagnostics Int.	Catalog# Cx40-A; RRID:AB_1616128

Anti-Dach1	Proteintech	Catalog# 10194-1-AP; RRID:AB_2069179
Anti-VegfR2	R&D Systems	Catalog# AF644; RRID:AB_355500
Anti-Endomucin	Thermo Fisher Scientific	Catalog# 14-5851-82; RRID:AB_891527
Anti-Actin, alpha-Smooth Muscle Monoclonal Antibody	Sigma-Aldrich	Catalog# C6198; RRID:AB_476856
Alexa fluor-conjugated antibodies (488, 555, 633)	Thermo Fisher Scientific	N/A
Chemicals, peptides, and recombinant proteins		
Tamoxifen	Sigma-Aldrich	Catalog# T5648-1G
Corn oil	Sigma-Aldrich	Catalog# T5648-5G
PFA	Thermo Fisher Scientific	Catalog# A11313.0E
Fetal bovine serum	HiMedia	Catalog# RM10432-500ML
DMEM/F12 media	Thermo Fisher Scientific	Catalog# 21041025
TritonX-100	Sigma-Aldrich	Catalog# X100-100ML
DAPI	Sigma-Aldrich	Catalog# D9542-5MG
Experimental models: Organisms/strains		
Rosa26^{TdTomato}	The Jackson Laboratory	Stock# 007909
Cx40CreER	Miquerol et al.[30]	N/A
ApjCreER	Chen et al.[18]	N/A
VegfR2 flox	The Jackson Laboratory	Stock# 018977
Cx40^{eGFP/+}	Miquerol et al.[31]	N/A

Tie2::eGFP::Claudin5	Knowland et al.[12]; Lutz et al.[13]	N/A
Cxcr4 flox	The Jackson Laboratory	Stock# 008767
Dach1 flox	Chang et al.[19]	N/A
Oligonucleotides		
Cre Forward primer (5' → 3') GCATTACCGGTCGATGCAACGAGTGATGAG	Arolkar and Kumar et al.[16]	N/A
Cre Reverse primer (5' → 3') GAGTGAACGAACCTGGAAATCAGTGCG	Arolkar and Kumar et al.[16]	N/A
TdTomato mutant Forward primer (5' → 3') CTGTTCCTGTACGGCATGG	The Jackson Laboratory	N/A
TdTomato mutant Reverse primer (5' → 3') GGCATTAAAGCAGCGTATCC	The Jackson Laboratory	N/A
TdTomato wild-type Forward primer (5' → 3') AAGGGAGCTGCAGTGGAGTA	The Jackson Laboratory	N/A
TdTomato wild-type Reverse primer (5' → 3') CCGAAAATCTGTGGGAAGTC	The Jackson Laboratory	N/A
VegfR2 flox Forward primer (5' → 3') CCACAGAACAACTCAGGGCTA	The Jackson Laboratory	N/A
VegfR2 flox Reverse primer (5' → 3') GGGAGCAAAGTCTCTGGAAA	The Jackson Laboratory	N/A
GFP Forward primer (5' → 3') GCACGACTTCTTCAAGTCCGCCATGCC	Arolkar and Kumar et al.[16]	N/A
GFP Reverse primer (5' → 3') GCGGATCTTGAAGTTCACCTTGATGCC	Arolkar and Kumar et al.[16]	N/A
Cxcr4 flox Forward primer (5' → 3') CCACCCAGGACAGTGTGACTCTAA	The Jackson Laboratory	N/A
Cxcr4 flox Reverse primer (5' → 3') GATGGGATTTCTGTATGAGGATTAGC	The Jackson Laboratory	N/A
Dach1 flox Forward primer (5' → 3') CTCCTGAAGATGAGGAGCTCACCCC	Chang et al.[19]	N/A
Dach1 flox Reverse primer (5' → 3') AACAATTCTTGTCCTTCACGTGCCC	Chang et al.[19]	N/A
Software and algorithms		
ImageJ	NIH (https://imagej.nih.gov/ij/)	RRID:SCR_003070

GraphPad Prism 9	GraphPad Software (http://www.graphpad.com/ scientific-software/prism/)	RRID:SCR_002798
Adobe Photoshop	Adobe (https://www.adobe.com/products/ photoshop.html)	RRID:SCR_014199
Adobe Illustrator	Adobe (https://www.adobe.com/products/ photoshop.html)	RRID:SCR_010279
OriginPro 2024b	OriginLab (https://www.originlab.com/index.aspx ?go=PRODUCTS/ Origin)	RRID:SCR_014212
Other		
Anti-fade mounting media	Jackson ImmunoResearch Laboratories Inc	N/A
Dental cement	3M Relyx U200	Catalog# 56877
Norland Optical Adhesive 68	Norland Products	Catalog# NOA 81,
2-well chamber slide	Nunc™ Lab-Tek™ (Thermo Fisher Scientific)	Catalog# 154461
5 mm coverslip	Warner Instruments	Catalog# CS-5R, CS-4R

536 **RESOURCE AVAILABILITY**

537 **Lead contact**

538 Further information and requests for resources and reagents should be directed to

539 and will be fulfilled by the lead contact, Soumyashree Das

540 (soumyashree@ncbs.res.in).

Materials availability

This study did not generate new unique reagents.

Data and code availability

- All data reported in the paper will be made available by the lead contact upon reasonable request.
- This paper does not report original code.
- Any additional information required to reanalyse the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Mice

All experiments were performed on mice with mixed strain background. All mice were housed and bred at the National Centre for Biological Sciences animal facility in accordance with the Institutional Animal Ethics Committee (IAEC). All procedures were performed with approval from IAEC. Data shown in this study is compiled from age-matched males and females from several litters.

METHOD DETAILS

Visualization of pial arteries in fixed samples

1. **Harvesting whole brains/heads from mouse embryos**

Depending on the developmental stages and size of the embryos, pial collateral arteries were visualized either in whole brains (e18.5 or P0) or using whole heads (e12.5- e16.5).

e12.5-e15.5: To perform the whole head explant at e12.5-e15.5 stages, the pregnant

females were euthanized through carbon dioxide overdose and a Y-shape incision

was made on the abdomen. Intact embryonic sacs were dissected out. Small

punctures were made on the embryo sac with the help of microforceps. The whole

embryo sac was fixed in 4% Paraformaldehyde (PFA) for 2 hours with gentle shaking

at 4°C. Following fixation, the embryo sac was washed at 4°C with gentle shaking.

This washing step was performed with 1X PBS and repeated 4 times every 15-20

mins. The embryos were then extracted from the embryo sacs and the heads were

removed with the help of microforceps.

e16-e16.5: The embryonic explants at e16/e16.5 are performed similarly. However,

for *Tie2::eGFP::Claudin5* genotype, the loose skin on top of the brain was removed

to aid immunostaining. Whole heads were imaged within 4 hrs, prior to PFA fixation

to avoid loss of GFP fluorescence signal.

e18.5: To perform the whole brain explant at e18.5 stage, the timed pregnant

females were euthanized as above. The skull of each embryo was gently removed

using the microforceps. The embryonic brains were scooped out with the help of a

microspatula and fixed in 4% PFA at 4°C for 2.5 hours. Agitation was avoided at this

stage to preserve tissue integrity of embryonic brains. The fixed embryonic brains

were then washed with 1X PBS three times at 15 minutes intervals with gentle

shaking at 4°C.

P0: Neonatal explants at P0 were performed by euthanizing the pups through carbon

dioxide overdose followed by decapitation. The skull was removed using scissors

and microforceps. The brains were scooped out using the microspatula. The brains

were then fixed with 4% PFA at 4°C for 2 hrs followed by three washes with 1X PBS

at an interval of 15 mins.

2. Whole brain/head immunostaining of mouse embryos

e12.5-e15.5: The embryonic heads (e12.5 - e15.5) were incubated in 0.1% PBST

(0.1% TritonX-100 in 1X PBS) for 2.5 hours to permeabilize the tissue, followed by

incubation with primary antibodies overnight at 4°C with gentle shaking. Heads were

then washed with 1X PBS, changing the buffer every 1-2 hours for 1 day. This step

was followed by incubation with secondary antibodies made in 0.1% PBST. Samples

were then washed every 1-2 hours for 1 day with 1X PBS. Following immunostaining

the brains are immersed in an anti-fade mounting media (from Jackson

ImmunoResearch Laboratories Inc.) and imaged immediately.

The embryonic and neonatal brains were incubated with primary antibody (made in

1X PBS) overnight, at 4°C with gentle shaking. Samples were then washed with 1X

PBS, changing the buffer every 1-2 hours for 1 day and incubated with Alexa Fluor

conjugated secondary antibodies (made in 1X PBS). Finally, samples were washed

every 1-2 hours for 1 day with 1X PBS and stored in mounting media or imaged

immediately.

Antibodies used are: Cx40 (Catalog number: Cx40-A, Alpha Diagnostics, 1:500

dilution), Dach1 (Catalog number: 10194-1-AP, Proteintech, 1:500 dilution), VegfR2

(Catalog number: AF644, R&D Systems, 1:250 dilution), Endomucin (Catalog

number: 14-5851-82; Thermo Fisher Scientific, 1:500 dilution) and alpha-smooth

muscle actin (Catalog number: C6198, Sigma-Aldrich, 1:500 dilution). All secondary

antibodies were procured from Invitrogen and used at a dilution of 1:500.

3. Imaging whole brains/heads of mouse embryos

The mice heads and brains from different developmental time-points were imaged using Olympus FV-3000 upright or inverted confocal microscope by placing the sample in a 2-well chamber (Catalog number: 154461; Nunc™ Lab-Tek™ II Chamber Slide™ System; Thermo Fisher Scientific). Whole brains and heads were scanned using 4X and 10X air objectives, and multiple stacks were captured. The captured image stacks were processed in ImageJ and stitched together using Adobe Photoshop software.

The embryonic heads at e16/e16.5 were imaged using Olympus FV-3000 5L Upright microscope. The pre-fixed embryonic heads were attached to a 35mm plastic petri dish using agarose gel prior to imaging. The whole head imaging was performed using 4X air objective and watershed images were captured with 10X air objective or 20X water dipping/immersion objective and 1.5X optical zoom with brains immersed in 1X PBS.

Visualization of pial arteries in live mice

1. In vivo imaging of embryonic pial arteries

1.1 Immobilization of live embryo

Cx40CreER; Rosa26^{TdTomato}; *Tie2::eGFP::Claudin5* pregnant dams with embryos at developmental stages between embryonic day (e)16-e17 were anesthetized with a subcutaneous injection of Ketamine-Xylazine (Ketamine (80 mg/kg); Xylazine (5 mg/kg)) cocktail. The dams were then placed in a supine position on a regulated heating pad (TC1000, CWE) that was positioned on top of a 3D-printed baseplate (as shown in Figure **1A**). A small incision was placed along the midline of the

abdomen. With a gentle retraction of the abdominal skin and wall, a single uterine

horn within the abdominal cavity was isolated. The uterus was surgically opened and

a single embryo along with the yolk sac was isolated and placed into a 35mm petri

dish. The embryo still connected to the dam was immobilized by submerging the

bottom part with 1% agarose, as shown in Figure **1A**. The immobilized embryo is

then submerged in dissection media (DMEM/F12, catalog number: 21041025,

Thermo Fisher Scientific + 10% FBS, catalog number: RM10432-500ML, HiMedia),

and the media is maintained at 37°C using a custom-made Arduino controlled media

heating and circulating device.

1.2 Live embryonic imaging

The setup along with anesthetized dam was placed under Macro Zoom microscope

MVX10-Olympus for large scale imaging and under FV3000 upright confocal

Microscope-Olympus for high resolution imaging using 20X water dipping objective.

In Macro Zoom Microscopy, the entire observable depth was scanned whilst

obtaining videos of 1 minute spanning a duration >2 hours in a time-lapse manner. In

confocal microscopy, the entire observable depth was scanned with a step size of 1

micron. A total of 4 embryos were imaged under the Macro zoom microscope and 5

embryos were imaged using confocal microscope. Out of these embryos, the artery

cell dynamics was captured in 2 brains from each category.

1.3 Image processing

Image stacks obtained from Macro zoom microscopy was registered using ImageJ

plugin (*StackReg*) and the registered stack was de-convolved using ImageJ plugin

(*DeconvolutionLab2*) and finally maximum intensity projection was obtained as

shown in Figure **1D**. The image stacks obtained from the confocal microscopy were

processed similarly except the deconvolution step.

The MCA and ACA growth fronts were automatically traced and the length of the

traced path was measured using Simple Neurite Tracer (SNT v4.1.13). The distance

of single cells from an artery was measured manually using ImageJ.

2. In vivo imaging adult pial arteries

2.1 Craniotomy

Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5 and VegfR2^{L/L};Cx40CreER;

Rosa26^{TdTomato}; Tie2::eGFP::Claudin5 adult mice (3-5 months) underwent

craniotomy to obtain optical access to the pial layer of the brain. Mice were

anesthetized under 2% Isoflurane (Abbott, North Chicago, IL, USA) and were placed

on a stereotaxic mount (Item number 51547; Stoelting, Wood Dale, IL, USA) with the

skull held tightly using ear bars. The mice were kept anesthetized with 2% Isoflurane

using a nose-cone. 5mg/kg Dexamethasone in 1X PBS was administered

subcutaneously and the mice kept warm using a heating pad (TC1000, CWE). The

eyes were covered with an eye ointment and the hair above the skull removed using

a hair trimmer. The surgical area was thoroughly wiped clean with 70% ethanol.

Sterile surgical tools were used for the surgery. A small incision was made using a

scalpel blade and the exposed skin was removed. The connective tissue covering

the skull in the exposed area was scraped off using scalpel blade and Q-tips. Using

a high-speed electrical drill (burr of diameter ~0.25mm), a window of ~4mm diameter

was carefully etched on the skull. The skull within the window was eventually

detached from the rest of the skull after repeated drilling and was carefully lifted

using a pair of fine forceps (Catalog number: 11255-20, Dumont #55) without

damaging the dura. During the process of drilling, the skull was constantly irrigated

with cortex buffer (Concentration; in mM: NaCl–125, KCl–5, glucose–10, HEPES–10,

$CaCl_2$–2 and $MgCl_2$–2. pH set to 7.35).[32] The exposed dura was aspirated with

cortex buffer until there is no bleeding and the window looked clear. An optical plug

made up of two 4 mm coverslip and one 5 mm coverslip (CS-5R & CS-4R; Warner

Instruments, Hamden, CT, USA) glued together (with NOA 81, Norland Optical

Adhesive) was placed on the exposed dura. The optical plug was gently pressed

using a toothpick such that the 5mm cover slip was touching the skull bone and then

cyano-acrylic glue was applied across the perimeter of the 5mm coverslip attaching it

to the skull. At this stage we had an optical window of 4mm. An aluminium head bar

with circular hole of 8mm was placed onto the skull aligned to the previously made

optical window. Next, dental cement (Catalog number: 56877, 3M Relyx U200) was

applied over the exposed skull using a sterile toothpick while being careful to not get

it on the optical window. Once the dental cement was completely dry, the animal was

placed on a warm pad in a clean recovery cage. Animals undergoing craniotomy

were maintained on meloxicam (2 mg/kg) and enrofloxacin (5 mg/Kg body weight)

for 3-4 days until full recovery. Mice were further allowed to recover for at least 2

weeks post-surgery before the start of any experiment. Mice with an optical window

can be imaged longitudinally for 2-3 months until the optical window starts to lose its

clarity.

2.2 Live Adult Imaging

To perform in vivo imaging, mice were anesthetized through intraperitoneal injection

of Ketamine (80mg/Kg body weight) – Xylazine (5mg/Kg body weight) cocktail. The

eyes were covered with an eye ointment and the mice were placed onto a custom-

made baseplate and immobilized using the implanted metal head plate to this

baseplate. A heating pad (37°C) was placed under the mouse to maintain the body

temperature. Subsequently, the baseplate was placed on a motorized XY-table of an

upright confocal microscope (FV3000 upright confocal Microscope-Olympus). The

entire optical window was imaged using 10X air objective (0.4 NA) and selected pial

regions with collateral arteries were imaged with 20X water dipping objective (0.5

NA). A total of N=3 adult mice per group (control and *VegfR2* artery endothelial

specific knockout) were imaged longitudinally for a period of 1 month.

2.3 Hypoxia regime

The *Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5* mice (N=3) were exposed to

hypoxia by gradually lowering the fraction of inspired oxygen (FiO$_2$) by 1% per day

over a period of 2 weeks to acclimatize them, and then maintained at 7% FiO$_2$ for an

additional 2 weeks (Figure **S12C**). The hypoxia chamber (Biospherix Ltd.) was

continuously supplied with nitrogen gas as needed to maintain the desired oxygen

levels, and CO$_2$ levels were kept below 600 ppm throughout the experiment by

changing the number of open holes (0.7 cm diameter) provided on each of the walls.

Oxygen and CO$_2$ levels were constantly monitored using inbuilt sensors (Biospherix;

ProOx Model P360 and ProCO2 Model P120ppm).

2.4 Image processing

721 The 10X image stacks obtained from the confocal microscopy of the cranial window

722 were registered using ImageJ plugin (*StackReg*) and maximum intensity projection

723 was obtained. The region of interest from the 20X image stacks were cropped out

724 slice by slice and stitched in Adobe Photoshop to avoid the overlaying microvascular

725 network in order to focus on the pial collaterals.

QUANTIFICATION AND STATISTICAL ANALYSIS

Number of collateral arteries per hemisphere

In *Cx40CreER*[+] (TdTomato) labelled brains, the collateral number was quantified based on the number of contiguous TdTomato[+] vessels (Td-contiguous) connecting MCA and ACA. Data shown in Figure **4K** was summarized from 2 control and 3 artery-specific *VegfR2* knockout. Data shown in Figure **S2H** was summarized from e13.5 (N=7), e14.5 (N=9), e15.5 (N=6), e18.5 (N=6) and P0 (N=5) embryos and neonates. Data shown in Figure **S3H** was summarized from e13.5 (N=3), e14.5 (N=3), e15.5 (N=2), e18.5 (N=2) and P0 (N=5) embryos and neonates. Data shown in Figure **S9E** is quantified by counting number of Cx40 immunostained collateral arteries between MCA and ACA, and is summarized from N=4 control and N=7 capillary specific *Dach1* knockouts. An unpaired *t* test was done to test significance statistically. Data were plotted using GraphPad Prism and presented as mean ± standard deviation. The p values in statistical analyses have been described as follow: ns (not significant), $p > 0.05$; $^*p < 0.05$; $^{**}p < 0.01$; $^{***}p < 0.001$; and $^{****}p < 0.0001$.

Microvascular tracks and their coverage with TdTomato[+] arteries per watershed

To analyse the Microvascular (MV) tracks, the watershed regions of the e16 (9 watershed from N=9 embryos) and e16.5 (7 watershed from N=5 embryos) brains were imaged using 10X air objective at 1.5X optical zoom. The region of interest (ROI) of comparable width (600-800 microns) within the watershed was taken for further analyses. The eGFP[+] MV tracks were traced using Simple Neurite Tracer (SNT v4.1.13). The total MV length and area of the ROI were measured using SNT

and ImageJ respectively, to calculate the total MV length per unit area of the

watershed. Further, to measure TdTomato coverage of the MV tracks, the

TdTomato⁻ vessel segments on the MV track were traced and their lengths measured

using SNT to quantify the fraction of TdTomato coverage. Both, the MV tracks and

TdTomato coverage were quantified by going through multiple z-slices and data

were presented as mean ± standard deviation. Two-tailed unpaired *t* test was done

to test statistical significance using GraphPad Prism software (p=0.4085 for

%TdTomato coverage in the MV tracks in Figure 2O and p=0.009 for normalized

total MV length per watershed in Figure 2J). The p values in statistical analyses have

been described as follow: ns (not significant), p > 0.05; *p < 0.05; **p < 0.01; ***p <

0.001; and ****p < 0.0001.

Lineage contribution towards pial collaterals during embryogenesis

Cx40 immunostained pial collateral segments from either *Cx40CreER;*

Rosa26^{TdTomato} or *ApjCreER; Rosa26^{TdTomato}* brains were imaged at 10X with 2.12

optic zoom. Regions of interest were manually drawn to mark the collateral artery

segments. Lineage contribution was determined by quantifying TdTomato⁺ coverage

on the collateral segment (Figure **3**). This was performed by thresholding the images

with Otsu algorithm in ImageJ. The area fraction of the TdTomato⁺ area per collateral

segment was calculated using the Measure tool in ImageJ. Contribution of the artery

and capillary derived ECs in collaterals was quantified by calculating the percentage

of collateral area covered by TdTomato⁺ signal. 45 collateral segments (of an

approximate length of 600-800μm) from 7 *Cx40CreER*-lineage traced brains

(biological replicate) and 57 collateral segments from 8 *ApjCreER*-lineage traced

brains were analysed. The data were presented as mean ± standard deviation using

GraphPad Prism.

Pial artery coverage in whole brains and in the collateral zone

The image stacks were obtained using confocal imaging of whole embryonic brains

with 10X objective, stitched using ImageJ plugin (*Stitching*) and max intensity

projection (MIP) was obtained. The MIPs were used for further analysis wherein one

complete hemisphere was cropped out and aligned vertically to the major axis of its

own elliptical boundary. Using a custom ImageJ macro, a line of the length of the

major axis was scanned across the width of the image and intensity profile for every

pixel along the horizontal axis was generated. Total number of local maxima (peaks)

for each intensity profile was obtained using the built-in ImageJ macro function

Array.findMaxima (array, tolerance). The tolerance used was 2 times the average

background intensity. The number of peaks thus obtained was plotted across the

width and the plot was fitted to a Gaussian function using OriginPro 2024b software.

The fitted curve was used as a representative of the arterial coverage per

hemisphere. The fit curves obtained from various mice were aligned based on the

Gaussian peaks and averaged to get the representative curves for various

experimental groups. We considered a region of 1mm (-0.5mm to +0.5 mm) across

the Gaussian peak as the collateral zone (watershed). The area under the curve

within this region was used as a measure for the extent of the collateral network. 9

hemispheres from 9 embryos (e18.5) were analysed per group (control and artery-

specific knockout for *VegfR2*). To test the statistical significance of the differences

observed in pial artery coverage (Figure **4E**), a Kolmogorov-Smirnov (KS) test was

done. D value= 0.569, p= 2.2066 X 10^{-141}. To test the statistical significance of the

area under the curve (pial collateral artery network shown in Figure **4F**), two-tailed

unpaired *t* test was done (p=0.0001) using GraphPad Prism. Similar analyses were

performed in Figures **S7** and **S10**. Two-tailed unpaired *t* test was performed using

the same. In all cases, data were presented as mean ± standard deviation and each

dot in Figures **4F, S10G, S10K** and **S10O** represents a brain (biological replicate).

The p values in statistical analyses have been described as follow: ns (not

significant), $p > 0.05$; $^{*}p < 0.05$; $^{**}p < 0.01$; $^{***}p < 0.001$; and $^{****}p < 0.0001$.

Efficiency of TdTomato recombination

Segments of collateral arteries, proximal arteries (PA) and distal arteries (DA) from

the image stacks taken with 20X objective were obtained from live adult imaging of

Cx40CreER; *Rosa26^{TdTomato}*; *Tie2::eGFP::Claudin5* mice. TdTomato^{+} and TdTomato^{-}

cells were counted in each artery segment (Figure **5D**) using ImageJ. Each segment

was of an approximate length of 600-800µm.The recombination efficiency was

measured as ratio of TdTomato^{+} cells and total number of cells in the segment. A

total of 10 collateral segments, 10 PA segments and 10 DA segments from 4 mice

were analysed to obtain the recombination efficiency of TdTomato. One-way ANOVA

was done to test significance statistically (Tukey's mean comparison test, PA vs.

collateral, $p<0.0001$; DA vs. collateral, $p<0.0001$; PA vs. DA, $p=0.08$). GraphPad

Prism was used to visualize the data as mean ± standard deviation. The p values in

statistical analyses have been described as follow: ns (not significant), $p > 0.05$; $^{*}p <$

0.05; $^{**}p < 0.01$; $^{***}p < 0.001$; and $^{****}p < 0.0001$.

Impact of arterial VegfR2 deletion on collateral formation

To assess the effect of arterial VegfR2 deletion on pial collateral development,

VegfR2$^{L/L}$; Cx40CreER; *Rosa26^{TdTomato}*; *Tie2::eGFP::Claudin5* mouse strain was

used. The watershed regions of e16.5 brains from control (7 watersheds from N=5

embryos) and knockout (5 watersheds from N=5 embryos) were imaged using 10X

air objective at 1.5X optical zoom. Microvascular tracks and their coverage with

TdTomato^{+} arterial cells were quantified using ImageJ (Figures **4K** and **S8A-S8C**)

following above-mentioned method. Capillary density was measured by thresholding

the images and obtaining the area fraction using ImageJ (Figures **S8D-S8E**). Each

dot in Figure S8E represents brain (biological replicate). For all cases, data were

plotted as mean ± standard deviation and two-tailed unpaired t test was done to test

the statistical significance using GraphPad Prism software (p=0.0004 for normalized

total MV length per watershed, p=0.0004 for %TdTomato coverage, and p=0.73 for

capillary density). The p values in statistical analyses have been described as follow:

ns (not significant), $p > 0.05$; $^{*}p < 0.05$; $^{**}p < 0.01$; $^{***}p < 0.001$; and $^{****}p < 0.0001$.

Quantification for deletion of *Dach1*

Dach1 deletion from the capillaries was validated in brains at embryonic day 18.5 by

immunostaining with antibodies for Dach1, VegfR2, and SMA along with nuclear

marker DAPI (Figure **S9C**). Dach1 expression was quantified manually using ImageJ

by calculating corrected total cell fluorescence (CTCF) from individual DAPI^{+}

capillary nucleus. 5 regions of interest (ROI) from each of the control (N=2) and 4

ROIs from each knockout brains (N=3 biological replicates) were used for analyses.

The data were plotted as mean ± standard deviation and two-tailed unpaired *t* test

was done to test the statistical significance using GraphPad Prism (p<0.0001). The p

values in statistical analyses have been described as follow: ns (not significant), p >

0.05; *p < 0.05; **p < 0.01; ***p < 0.001; and ****p < 0.0001.

AUTHORS CONTRIBUTION

SD conceived the idea and designed the experiments. SK performed intravital imaging of embryos and adults. SK, SG, NS, VS, MD performed embryonic experiments. SD, SK, SG, NS wrote the manuscript. All authors contributed to editing the manuscript.

ACKNOWLEDGEMENTS

We thank the following facilities for their technical support: Animal Care and Resource Center at NCBS, Mouse Genome Engineering Facility at NCBS, Central Imaging and Flow Cytometry Facility at NCBS. We sincerely thank Dr. Dritan Agalliu (Columbia University) and Dr. Kristy Red-Horse (Stanford University) for generously sharing *Tie2::eGFP::Claudin5* and *Dach1$^{fl/fl}$* mice respectively.

FUNDING

This work is supported by the Department of Atomic Energy, Government of India, Project Identification No. RTI 4006, and DBT/Wellcome Trust India Alliance Intermediate Fellowship to SD (IA/I/20/2/505205). SK is supported by DBT/Wellcome Trust India Alliance Intermediate Fellowship to SD. NS and VS were supported by DST INSPIRE Scholarship for Higher Education (SHE).

DECLARATION OF INTERESTS

None

REFERENCES

1. Lima, F.O., Furie, K.L., Silva, G.S., Lev, M.H., Camargo, E.C.S., Singhal, A.B., Harris, G.J., Halpern, E.F., Koroshetz, W.J., Smith, W.S., et al.

(2010). The pattern of leptomeningeal collaterals on CT angiography is a

strong predictor of long-term functional outcome in stroke patients with

large vessel intracranial occlusion. Stroke *41*, 2316–2322.

https://doi.org/10.1161/STROKEAHA.110.592303.

2.   Maas, M.B., Lev, M.H., Ay, H., Singhal, A.B., Greer, D.M., Smith, W.S.,

Harris, G.J., Halpern, E., Kemmling, A., Koroshetz, W.J., et al. (2009).

Collateral vessels on CT angiography predict outcome in acute ischemic

stroke. Stroke *40*, 3001–3005.

https://doi.org/10.1161/STROKEAHA.109.552513.

3.   Binder, N.F., El Amki, M., Glück, C., Middleham, W., Reuss, A.M.,

Bertolo, A., Thurner, P., Deffieux, T., Lambride, C., Epp, R., et al. (2024).

Leptomeningeal collaterals regulate reperfusion in ischemic stroke and

rescue the brain from futile recanalization. Neuron *112*, 1456-1472.e6.

https://doi.org/https://doi.org/10.1016/j.neuron.2024.01.031.

4.   The top 10 causes of death https://www.who.int/news-room/fact-

sheets/detail/the-top-10-causes-of-death.

5.   Zhang, H., Prabhakar, P., Sealock, R., and Faber, J.E. (2010). Wide

genetic variation in the native pial collateral circulation is a major

determinant of variation in severity of stroke. Journal of Cerebral Blood

Flow & Metabolism *30*, 923–934. https://doi.org/10.1038/jcbfm.2010.10.

6.   Clayton, J.A., Chalothorn, D., and Faber, J.E. (2008). Vascular

endothelial growth factor-a specifies formation of native collaterals and

regulates collateral growth in ischemia. Circ Res *103*, 1027–1036.

https://doi.org/10.1161/CIRCRESAHA.108.181115.

7.   Lucitti, J.L., MacKey, J.K., Morrison, J.C., Haigh, J.J., Adams, R.H., and
Faber, J.E. (2012). Formation of the collateral circulation is regulated by
vascular endothelial growth factor-a and a disintegrin and
metalloprotease family members 10 and 17. Circ Res *111*, 1539–1550.
https://doi.org/10.1161/CIRCRESAHA.112.279109.

8.   Lucitti, J.L., Sealock, R., Buckley, B.K., Zhang, H., Xiao, L., Dudley,
A.C., and Faber, J.E. (2016). Variants of Rab GTPase-Effector Binding
Protein-2 Cause Variation in the Collateral Circulation and Severity of
Stroke. Stroke *47*, 3022–3031.
https://doi.org/10.1161/STROKEAHA.116.014160.

9.   Chalothorn, D., Clayton, J.A., Zhang, H., Pomp, D., and Faber, J.E.
(2007). Collateral density, remodeling, and VEGF-A expression differ
widely between mouse strains. Physiol Genomics *30*, 179–191.
https://doi.org/10.1152/PHYSIOLGENOMICS.00047.2007.

10.   Das, S., Goldstone, A.B., Wang, H., Farry, J., D'Amato, G., Paulsen,
M.J., Eskandari, A., Hironaka, C.E., Phansalkar, R., Sharma, B., et al.
(2019). A Unique Collateral Artery Development Program Promotes
Neonatal Heart Regeneration. Cell *176*, 1128-1142.e18.
https://doi.org/10.1016/J.CELL.2018.12.023.

11.   Perovic, T., Hollfinger, I., Mayer, S., Lips, J., Dopatka, M., Harms, C.,
and Gerhardt, H. (2024). Pial collaterals develop through mosaic
colonization of capillaries by arterial and microvascular endothelial cells.
Elife *13*, RP93746. https://doi.org/https://doi.org/10.7554/eLife.93746.1.

12. Knowland, D., Arac, A., Sekiguchi, K.J., Hsu, M., Lutz, S.E., Perrino, J.,

Steinberg, G.K., Barres, B.A., Nimmerjahn, A., and Agalliu, D. (2014).

Stepwise Recruitment of Transcellular and Paracellular Pathways

Underlies Blood-Brain Barrier Breakdown in Stroke. Neuron *82*, 603–

617. https://doi.org/https://doi.org/10.1016/j.neuron.2014.03.003.

13. Lutz, S.E., Smith, J.R., Kim, H., Bates, J.M., Gandhi, S.P., and Agalliu,

D. (2017). Caveolin1 Is Required for Th1 Cell Infiltration, but Not Tight

Junction Remodeling, at the Blood-Brain Barrier in Autoimmune

Neuroinflammation. CellReports *21*, 2104–2117.

https://doi.org/10.1016/j.celrep.2017.10.094.

14. Jamaiyar, A., Juguilon, C., Wan, W., Richardson, D., Chinchilla, S.,

Gadd, J., Enrick, M., Wang, T., McCabe, C., Wang, Y., et al. (2022). The

essential role for endothelial cell sprouting in coronary collateral growth.

J Mol Cell Cardiol *165*, 158–171.

https://doi.org/https://doi.org/10.1016/j.yjmcc.2022.01.005.

15. Gabhann, F. Mac, and Peirce, S.M. (2010). Collateral Capillary

Arterialization following Arteriolar Ligation in Murine Skeletal Muscle.

Microcirculation *17*, 333–347. https://doi.org/10.1111/J.1549-

8719.2010.00034.X.

16. Arolkar, G., Kumar, S.K., Wang, H., Gonzalez, K.M., Kumar, S., Bishnoi,

B., Rios Coronado, P.E., Woo, Y.J., Red-Horse, K., and Das, S. (2023).

Dedifferentiation and Proliferation of Artery Endothelial Cells Drive

Coronary Collateral Development in Mice. Arterioscler Thromb Vasc Biol

43, 1455–1477. https://doi.org/10.1161/ATVBAHA.123.319319.

17. Raftrey, B., Williams, I.M., Rios Coronado, P.E., Fan, X., Chang, A.H.,
Zhao, M., Roth, R., Trimm, E., Racelis, R., D'amato, G., et al. (2021).
Dach1 Extends Artery Networks and Protects Against Cardiac Injury.
Circ Res *129*, 702–716.
https://doi.org/10.1161/CIRCRESAHA.120.318271.

18. Chen, H.I., Sharma, B., Akerberg, B.N., Numi, H.J., Kivelä, R.,
Saharinen, P., Aghajanian, H., McKay, A.S., Bogard, P.E., Chang, A.H.,
et al. (2014). The sinus venosus contributes to coronary vasculature
through VEGFC-stimulated angiogenesis. Development *141*, 4500–
4512. https://doi.org/10.1242/dev.113639.

19. Chang, A.H., Raftrey, B.C., D'Amato, G., Surya, V.N., Poduri, A., Chen,
H.I., Goldstone, A.B., Woo, J., Fuller, G.G., Dunn, A.R., et al. (2017).
DACH1 stimulates shear stress-guided endothelial cell migration and
coronary artery growth through the CXCL12-CXCR4 signaling axis.
Genes Dev *31*, 1308–1324. https://doi.org/10.1101/GAD.301549.117/-
/DC1.

20. Scott Budinger, G.R., Duranteau, J., Chandel, N.S., and Schumacker,
P.T. (1998). Hibernation during Hypoxia in Cardiomyocytes ROLE OF
MITOCHONDRIA AS THE O 2 SENSOR*. Journal of Biological
Chemistry *273*, 3320–3326. https://doi.org/10.1074/jbc.273.6.3320.

21. Rink, C., and Khanna, S. (2011). Significance of Brain Tissue
Oxygenation and the Arachidonic Acid Cascade in Stroke.
https://home.liebertpub.com/ars *14*, 1889–1903.
https://doi.org/10.1089/ARS.2010.3474.

22.   Mannion, J.D., Buckman, P.D., Magno, M.G., and Dimeo, F. (1992).

963   Collateral blood flow from skeletal muscle to normal myocardium.

964   Journal of Surgical Research *53*, 578–587.

965   https://doi.org/https://doi.org/10.1016/0022-4804(92)90258-2.

966   23.   Crissinger, K.D., and Neil Granger, D. (1988). Characterization of

967   Intestinal Collateral Blood Flow in the Developing Piglet. Pediatr Res *24*,

968   473–476. https://doi.org/10.1203/00006450-198810000-00011.

969   24.   Kochi, T., Imai, Y., Takeda, A., Watanabe, Y., Mori, S., Tachi, M., and

970   Kodama, T. (2013). Characterization of the Arterial Anatomy of the

971   Murine Hindlimb: Functional Role in the Design and Understanding of

972   Ischemia Models. PLoS One *8*, e84047.

973   https://doi.org/10.1371/JOURNAL.PONE.0084047.

974   25.   Gao, Y., Aravind, S., Patel, N.S., Fuglestad, M.A., Ungar, J.S., Mietus,

975   C.J., Li, S., Casale, G.P., Pipinos, I.I., and Carlson, M.A. (2020).

976   Collateral Development and Arteriogenesis in Hindlimbs of Swine After

977   Ligation of Arterial Inflow. Journal of Surgical Research *249*, 168–179.

978   https://doi.org/10.1016/J.JSS.2019.12.005.

979   26.   He, L., Liu, Q., Hu, T., Huang, X., Zhang, H., Tian, X., Yan, Y., Wang, L.,

980   Huang, Y., Miquerol, L., et al. (2016). Genetic lineage tracing discloses

981   arteriogenesis as the main mechanism for collateral growth in the mouse

982   heart. Cardiovasc Res *109*, 419–430.

983   https://doi.org/10.1093/CVR/CVW005.

984   27. Zhang, H., Rzechorzek, W., Aghajanian, A., and Faber, J.E. Hypoxia

985   induces de novo formation of cerebral collaterals and lessens the

986   severity of ischemic stroke. https://doi.org/10.1177/0271678X20924107.

987   28. Zhang, H., Chalothorn, D., and Faber, J.E. (2019). Collateral Vessels

988   Have Unique Endothelial and Smooth Muscle Cell Phenotypes.

989   International Journal of Molecular Sciences 2019, Vol. 20, Page 3608

990   *20*, 3608. https://doi.org/10.3390/IJMS20153608.

991   29. Stanca Melincovici, C., Boşca, A.B., Şuşman, S., Mărginean, M., Mihu,

992   C., Istrate, M., Moldovan, I.-M, Roman, A.L., and Mihu, C.M. (2018).

993   Vascular endothelial growth factor (VEGF)-key factor in normal and

994   pathological angiogenesis. Rom J Morphol Embryol *59*.

995   30. Miquerol, L., Thireau, J., Bideaux, P., Sturny, R., Richard, S., and Kelly,

996   R.G. (2015). Endothelial plasticity drives arterial remodeling within The

997   endocardium after myocardial infarction. Circ Res *116*, 1765–1771.

998   https://doi.org/10.1161/CIRCRESAHA.116.306476.

31. Miquerol, L., Meysen, S., Mangoni, M., Bois, P., van Rijen, H.V.M.,

Abran, P., Jongsma, H., Nargeot, J., and Gros, D. (2004). Architectural

and functional asymmetry of the His–Purkinje system of the murine

heart. Cardiovasc Res *63*, 77–86.

https://doi.org/10.1016/j.cardiores.2004.03.007.

32. Holtmaat, A.J.G.D., Trachtenberg, J.T., Wilbrecht, L., Shepherd, G.M.,

Zhang, X., Knott, G.W., and Svoboda, K. (2005). Transient and

Persistent Dendritic Spines in the Neocortex In Vivo. Neuron *45*, 279–

291. https://doi.org/10.1016/j.neuron.2005.01.003.

Main figures legends

 Figure 1: Live imaging of embryonic pial vasculature

 (**A**) Intra-vital imaging set up to visualize the developing pial vasculature in mouse

 embryos. **i** heating pad to maintain body temperature of anaesthetized pregnant dam

 at 37°C. **ii** custom-built media heating device which also circulates dissection media

 at 37°C **iii** shows macro-zoom or confocal microscopy performed on live,

 immobilized embryos.

 (**B**) Representative schematic of an embryonic brain. Grey and Red show cerebral

 arteries and pial collaterals respectively.

 (**C**) Experimental design to genetically label and trace the dynamics of pre-existing

 artery endothelial cells in a developing pial layer.

 (**D-G**) Images obtained through macro-zoom fluorescent microscopy of e16.5

 embryos. In black are cells derived from pre-existing artery endothelial cells. Images

 were taken by focusing in and out to capture the entire curvature of the embryonic

 brain. **D** shows an entire left hemisphere. **E** represents watershed region spanned by

 MCA and ACA artery tips and imaged at time=0 minutes, 20 minutes and 80 minutes

 respectively. Arrows and arrowheads show growing ends of pre-existing artery tips

 from MCA and ACA. Brackets mark regions where collaterals appear. **F** shows inset

 in **D**. **E** and **G** show insets in **F**. **G** represents the MCA growth front imaged at time=0

 minutes, 20 minutes and 80 minutes respectively. * shows a superficial yolk sac

 vessel which photobleaches with time. Arrowheads mark extension and

 anastomoses of MCA artery tips.

 (**H-J**) Images obtained through confocal microscopy from e16 embryos. **H** shows pial

 watershed and the cerebral arteries spanning the watershed. **I**, **J** show MCA and

 ACA growth fronts respectively imaged at time=0 minutes and 60 minutes. In black

are cells derived from pre-existing artery endothelial cells. Images shown in **I** are

maximum intensity projections from 50 optical slices and capture the entire width of

the artery. Images shown in **J** are maximum intensity projections from 40 optical

slices and capture the entire width of the artery. * represents an internal landmark

from where the length measurements were done.

(**K-M**) Images obtained through confocal microscopy from e16 embryos. (**L**, **M**)

Insets from **K** showing artery cell growth of developing ACA in 60 mins. (**L**)

Arrowhead points to an artery tip emerging within 60mins. Arrow points to a region

where single recombined artery cells appear. (**M)** Arrow shows direction of migration

of single recombined artery cell. Dotted line is the length of micro-vessel between the

single recombined cell and the pial artery. MCA, Middle cerebral artery; ACA,

Anterior cerebral artery; e, embryonic day; Tam, Tamoxifen; EC, endothelial cell;

YSV, yolk sac vessel; brch., branch; coll, collateral; MV, microvascular. Scale bars:

D, **F**, 1mm; **E**, **G**, 200 µm; **H**, **K**, 500 µm; **I**, **J**, **L**, **M**, 100 µm.

A e16

MCA — watershed
B
ACA

B TdTomato (artery cell lineage)
eGFP (all ECs)

Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5

eGFP
TdTomato

to MCA
MV track
Td+ EC
MV track
MV track
Td+ EC
to ACA

eGFP
MV track
Td+ EC
MV track

C Potential mechanisms leading to pial collateral fromation

Watershed e14.5
i

MV tracks form between two arteries
ii   e16   v   inception of MV tracks

artery tip extension
iii   e16.5   vi   artery tip extension / MV track remodelling

collateral
iv   e18.5/ P0

capillaries   microvascular tracks   pre-existing artery tips

Growth of MV tracks and arteries occur concurrently

Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5

D e16
TdTomato (artery cell lineage) eGFP (all ECs)
Tam at e14.5

I e16.5
K-N

to ACA → ← to MCA
watershed

E TdTomato (artery cell lineage) eGFP (all ECs)

F all ECs

G artery cells

H skeleton of MV tracks with or without TdTomato+ cells

J Total length of MV track per watershed
p=0.009 **
Normalized MV track length per watershed area (AU)
2.0
1.5
1.0
0.5
e16   e16.5

O Td coverage of MV tracks per watershed
ns
% coverage of Td+ cells
e16   e16.5

to ACA → ← to MCA
watershed

K TdTomato (artery cell lineage) eGFP (all ECs)

L all ECs

M artery cells

N skeleton of MV tracks with or without TdTomato+ cells

1050

1051

Figure 2: Analyses of development of MV (microvascular) tracks and artery tips during pial collateral development

(**A**) In vivo imaging of live mouse embryo at e16 shows the cerebral arteries spanning the pial watershed in mouse brain.

(**B**) Inset from **A** showing TdTomato$^+$ artery endothelial cells (yellow arrowheads) on eGFP$^+$ microvascular tracks (arrows). Yellow arrows point to microvascular tracks partially or completely occupied by TdTomato$^+$ artery ECs. Magenta arrows point to microvascular tracks at the growing front of the TdTomato$^+$ labelled arteries.

(**C**) Two potential mechanisms for pial collateral formation. Hypothesis 1: (i) An artery-free e14.5 watershed region (ii) is first occupied by microvascular tracks (magenta) connecting two arteries (cyan) by e16. (iii) Artery tips then start extending on pre-determined MV tracks at e16.5. Arrows indicate direction of artery tip migration. (iv) This process leads to pial collateral formation. Hypothesis 2: Alternatively, (v) while MV tracks start appearing at e16, (vi) they actively remodel and grow in length to allow artery tip extensions in a directional manner. This leads to formation of pial collaterals. Cyan represents pial artery, magenta represents MV track.

(**D-N**) Concurrent MV track growth and artery tip extension. (**D and I**) Whole brain images of e16 and e16.5 embryos respectively, from *Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5* mice, where TdTomato (cyan) marks pre-existing artery endothelial cells and eGFP (magenta) marks all endothelial cells. **E-H** is inset shown in **D**. (**E**) Higher magnification confocal image of watershed region (715 µm width) between ACA and MCA. All endothelial cells are in magenta and artery endothelial cells in cyan. (**F**) eGFP$^+$ vessels in the selected watershed from e16 watershed shown in **E**. (**G**) Pre-existing artery cells in the watershed from e16 watershed shown

in **E**. (**H**) Skeleton of all the eGFP$^+$ microvascular tracks traced in the e16 watershed (shown in **E**) through multiple z-slices, using Simple Neurite Tracer. Traced MV tracks occupied by TdTomato$^+$ artery endothelial cells are shown in cyan and MV tracks not occupied with TdTomato is shown in yellow. (**J**) Quantification of total length of microvascular tracks per watershed area (Two-tailed unpaired t test, p=0.009, **p < 0.01) and (**O**) Quantification of percentage TdTomato coverage of microvascular tracks in watershed regions from e16 (9 watersheds from N=9 embryos) and e16.5 (7 watersheds from N=5 embryos) brains. Data shown as mean ± standard deviation. (Two-tailed unpaired t test, p=0.4085, ^{ns} non-significant, p>0.05). **K-N** is inset shown in **I**. (**K**) Higher magnification confocal image of watershed region (715 µm width) between ACA and MCA. All endothelial cells are in magenta and artery endothelial cells in cyan. (**L**) eGFP$^+$ vessels in the selected watershed from e16 watershed shown in **K**. (**M**) Pre-existing artery cells in the watershed from e16 watershed shown in **K**. (**N**) Skeleton of all the eGFP$^+$ microvascular tracks traced in the e16 watershed (shown in **K**) through multiple z-slices, using Simple Neurite Tracer. Traced MV tracks occupied by TdTomato$^+$ artery endothelial cells are shown in cyan and MV tracks not occupied with TdTomato is shown in yellow. Tam, Tamoxifen; MCA, Middle Cerebral Artery; ACA, Anterior Cerebral Artery; MV, Microvascular; Td, TdTomato; e, embryonic day; EC, Endothelial cell; AU, Arbitrary units. Scale bar: A, 500µm; B, 100µm; D and I, 500µm; E and K, 100µm.

A

Cx40CreER; Rosa26^{TdTomato}
or
ApjCreER; Rosa26^{TdTomato}

MCA and ACA present
pial collaterals appear
trace artery/cap-EC lineages into collaterals

e14.5   e15.5   e16.5   e18.5   P0

Tam

label pre-existing arteries

whole brain confocal imaging & lineage analyses

B Pial collaterals predominantly originate from pre-existing artery ECs

Cx40CreER; Rosa26^{TdTomato}   *ApjCreER; Rosa26^{TdTomato}*

Average area (%) per vessel segment

77%   *Cx40*-lineage

31.6%   *Apj*-lineage

Cx40CreER; Rosa26^{TdTomato}
C TdTomato   **D** watershed — Cx40 Cx40-lineage

ACA brchs.   MCA

ApjCreER; Rosa26^{TdTomato}
E Cx40   **F** watershed — Cx40 Apj-lineage

ACA brchs.   MCA

G pial collateral artery segment   Cx40

H pial collateral artery segment   Cx40

I Cx40-lineage

J Apj-lineage

1098

1099

Figure 3: Cellular origin of pial collaterals

(**A**) Experimental design to assess lineage contribution from artery ECs and capillary ECs towards pial collaterals at e18.5/ P0.

(**B**) Graph shows the average percent area of collateral segments covered with TdTomato⁺ cells in two independent genetic lineage tracing experiments using (left) *Cx40CreER* (N=7 biological replicate) and (right) *ApjCreER* (N=8 biological replicate) mouse lines respectively. The whiskers in the boxplot extends from minimum to maximum value.

(**C**) Representative image of whole right hemisphere from *Cx40CreER*; *Rosa26^{TdTomato}* e18.5 embryos. Black shows TdTomato⁺ arteries.

(**D**) Boxed region of interest with a collateral segment from the watershed area shown in **C**. The pial collateral is lineage traced for artery ECs (cyan) and immunostained with Cx40 (magenta).

(**E**) Representative image of whole right hemisphere from *ApjCreER*; *Rosa26^{TdTomato}* e18.5 embryos. Black shows Cx40 immunostaining.

(**F**) Boxed region of interest with a collateral segment from the watershed area shown in **E**. The pial collateral is lineage traced for capillary ECs (cyan) and immunostained with Cx40 (magenta).

(**G**, **I**) Collateral segment from **D** is (**G**) immunostained for Cx40 and (**I**) lineage traced with TdTomato.

(**H**, **J**) Collateral segment from **F** is (**H**) immunostained for Cx40 and (**J**) lineage traced with TdTomato. Arrowheads point to TdTomato⁺ cells from Apj⁺ lineage on the collateral segment. e, embryonic day, P, postnatal day; MCA, Middle cerebral artery; ACA, Anterior cerebral artery; brch., branch; EC, endothelial cell; Tam, Tamoxifen.

Scale bars: **C** and **E**, 500µm; **D** and **F**, 100µm

Figure 4: VegfR2-dependent formation of pial collateral arteries

(**A**) Experimental design to lineage label and trace pre-existing artery endothelial cells in pial layer, with or without arterial *VegfR2* and *Cxcr4* at e18.5/ P0.

(**B**) Representative confocal images of whole brains with and without arterial *Cxcr4*. Black shows pre-existing artery endothelial cells. Control, N= 8 and artery specific knockout for *Cxcr4*, N= 13.

(**C**) Summary of method to quantify pial artery coverage across the entire brain including watershed.

(**D**) Confocal images of whole brains with and without arterial *VegfR2*. Black shows pre-existing artery endothelial cells.

(**E**) Quantification of overall pial artery coverage with and without arterial *VegfR2* quantified using method shown in **C**. KS test performed on the distributions generated D value= 0.569, p value= 2.2066 X 10^{-141}.

(**F**) Quantification of pial collateral artery network with and without arterial *VegfR2* quantified using method shown in **C**. N=9 biological replicates for each group. Two-tailed unpaired *t* test, p=0.0001.

(**G**) Experimental design to assess the effect of *VegfR2* deletion from arteries at e16.5.

(**H**, **I**) Confocal images of pial watershed region showing TdTomato-contiguous pial collateral arteries (Td^{con}) in magenta and microvasculature in cyan. Arrowheads show part of microvascular track not occupied by TdTomato$^+$ artery cell, arrows point to artery tips on microvascular tracks.

(**J**) Quantification of TdTomato$^+$ artery cell coverage of pre-existing MV tracks with

(N=7 biological replicates) or without (N=5 biological replicates) arterial *VegfR2*.

Two-tailed unpaired *t* test, p=0.0004.

(**K**) Quantification of pial collateral arteries contiguously populated with TdTomato$^+$

artery cells, with (N=2) and without (N=3 individual brains) arterial *VegfR2*. Two-

tailed unpaired *t* test, p=0.0124. Each dot in F, J, and K represents a biological

replicate. Data shown as mean ± standard deviation. *p < 0.05 and ***p < 0.001.

MCA, Middle cerebral artery; ACA, Anterior cerebral artery; e, embryonic day; P,

postnatal day; Tam, Tamoxifen; EC, endothelial cell; Td^{con}, TdTomato-contiguous;

AU, Arbitrary Unit; aEC, artery endothelial cell; KO, knockout. Scale bars: **B**, **D**,

500µm; **H**, **I**, 100µm.

1160

Intra-vital imaging of pial collaterals in adult mice

1161

Figure 5: Longitudinal live-imaging of adult pial collaterals

(**A**, **left**) Dorsal view of the cranial window created with the attached headbar on

adult mouse head. (**A**, **right**) Brightfield image of the 4mm cranial window obtained

with a stereoscope shows the pial vasculature. Dotted rectangle encompasses MCA,

ACA and the watershed between them and is imaged at high resolution as shown in

E.

(**B**) Schematic of the side view of the cranial window showing the 3-tier optical plug

inserted into the skull giving access to pial layer.

(**C**) Experimental design for longitudinal imaging of pial vasculature in adult mice.

(**D**) Quantification of TdTomato reporter recombination in pial collateral arteries and

different segments of cerebral arteries. Each dot represents individual vessel. One-

way ANOVA was done to test significance statistically (Tukey's mean comparison

test, PA vs. collateral, $p<0.0001$; DA vs. collateral, $p<0.0001$; PA vs. DA, $p=0.08$).

$^{****}p < 0.0001$.

(**E-K**) Images of pial watershed with pre-existing artery endothelial cells labelled in

TdTomato (magenta) and all endothelial cells labelled in eGFP (cyan). (**E**) Confocal

image from optical window created for longitudinal imaging of *Cx40CreER;*

Rosa26^{TdTomato}; Tie2::eGFP::Claudin5 mouse as shown in Figure **4A**. The image

shows both pial and dural vessels. (**F-K**) Confocal images of pial collateral arteries

from cranial window shown in **E**, across 3 timepoints, each 2 weeks apart. (**I**) Pial

layer segmented out from **E**. red and blue show pial arteries and veins respectively.

J, **K** are the insets shown in **F-H**.

(**L**, **M**) Cellular distribution of TdTomato+ artery cells and eGFP+ endothelial cells in

collateral artery segments shown in **J** and **K** respectively. * represents a reference

cell. MCA, middle cerebral artery; ACA, anterior cerebral artery; d, days; Tam,

Tamoxifen; PA, proximal artery; DA, distal artery; coll, collateral; brch., branch; EC,

endothelial cell. Scale bars: **E**, 200µm; **J**, **K** 25µm

A *VegfR2^{L/L};Cx40CreER; Rosa26^{TdTomato}*
Tie2::eGFP::Claudin5

Craniotomy

5d

dose 1 dose 2 dose 3 dose 4 dose 5

label
pre-existing
arteries
+
VegfR2
deletion

14d   7d   14d   14d

Tam

t=0d   t=14d   t=28d
intra-vital imaging

B 4mm diameter cranial window

MCA brch.   ACA brch.

C

C watershed
TdTomato (pre-existing artery ECs)
eGFP (all ECs)

MCA brch.   E-G   ACA brch.

collateral

D Pial arteries   Pial veins

collateral

MCA brch.   ACA brch.

collateral

MCA-ACA collateral artery

E t=0 days   **F** t=14 days   **G** t=28 days

H   I   J

H t=0 days   **I** t=14 days   **J** t=28 days

K   **L**   **M**

N VegfR2 regulates pial collateral development by artery tip extension

development
(embryogenesis)
VegfR2-driven
artery-tip extension

maintenance
(adulthood)
VegfR2/Cxcr4
independent process

pial collateral arteries

flow from MCA   flow from ACA

artery lineage   non-artery lineage

1189

1190

1191

Figure 6: Analyses of role of arterial VegfR2 in pial collateral maintenance

(A) Experimental design to visualize artery cell dynamics in pial layer of adult mice without arterial *VegfR2*.

(**B**) Brightfield image of cranial window showing pial vasculature. The dotted rectangle encompasses branches of MCA, ACA and the watershed between them.

(**C**) Confocal image of cranial window created in mice without arterial *VegfR2* as shown in **B**. Magenta shows all artery cells, cyan shows all endothelial cells.

(**D**) Pial layer segmented out from **C** shows pial arteries and veins in red and blue respectively.

(**E-G**) Confocal images of collateral artery from three timepoints, each 2 weeks apart as shown in the boxed region in **C**.

(**H-J**) Confocal images of collateral artery segments shown in **E-G**.

(**K-M**) Cellular distribution of TdTomato$^+$ artery cells and eGFP$^+$ endothelial cells in collateral artery segments shown in **H-J**. * represents a reference cell.

(**N**) Schematic summarizing the role of arterial VegfR2 in development and maintenance of pial collateral arteries. While arterial VegfR2 facilitates tip extension of pre-existing arteries during development, it is dispensable for pial collateral maintenance in adulthood. Pial collaterals, both in developing and mature brains, are majorly populated by endothelial cells of artery-lineage. MCA, middle cerebral artery; ACA, anterior cerebral artery; d, days; Tam, Tamoxifen; EC, endothelial cell; brch., branch. Scale bars: **C**, 150 µm; **E-G**, 50µm; **H-J**, 20µm.

Supplementary figures and figure legends

Supplemental Figure 1: Visualization of pial arteries and collateral arteries,

related to Figure 1

Schematic summarizes key steps of immunostaining methods used to image pial

artery network in (**A**) mouse embryonic heads and (**B**) mouse embryonic/neonatal

brains with confocal microscope. (**C**) Schematic shows findings from **A** and **B** in one

hemisphere. Grey represents cerebral arteries, Red represents collateral artery

connections between cerebral arteries. Brackets show watershed area (WSA), MCA

appears at e12.5, ACA appears at e13.5, and the foremost pial collateral arteries

appear at e15.5. These pial collaterals then increase in number till e18.5/P0. e,

embryonic day; P, postnatal day; MCA, middle cerebral artery; ACA, anterior

cerebral artery.

1229

1232   Supplemental Figure 2: Detection of pial collaterals with whole brain

1233   immunostaining, related to Figure 1

(**A-F**) Representative images of right hemispheres from developing mouse embryos

immunostained for Connexin-40 (Cx40). Cx40 marks all artery endothelial cells.

MCA appears at e12.5, ACA appears at e13.5 and collaterals between MCA and

ACA appear at e15.5. Arrowheads point to collateral arteries between MCA and

ACA. (**G**) Quantification of Cx40 immunostained brains from different developmental

stages reveals that both MCA and ACA have formed by e14.5. (**H**) Quantification of

Cx40$^+$ collateral arteries per hemisphere shows that the first pial collaterals appear at

e15.5. N=7 (e13.5), 9 (e14.5), 6 (e15.5), 6 (e18.5) and 5 (P0 brain). Each dot

represents an individual brain. e, embryonic day; P, postnatal day; MCA, middle

cerebral artery; ACA, anterior cerebral artery; brch., branch. Scale bars: 500μm

Cx40 eGFP/+

left hemisphere

e12.5 (MCA appears)

A — MCA

B e13.5 — MCA

right hemisphere

C e14.5 (ACA appears) — MCA, ACA

D e15.5 (collateral arteries appear) — MCA, ACA

E e18.5 — ACA, MCA brch

F P0 — ACA, MCA brch

G MCA and ACA artery trees form by e14.5

H Pial collateral arteries appear at e15.5

1244

1245

Supplemental Figure 3: Detection of pial collaterals with *Cx40* reporter activity,

related to Figure 1

(**A-F**) Representative images of brains from developing *Cx40eGFP/+* mouse embryos.

Connexin-40 (Cx40) marks all artery endothelial cells. MCA appears at e12.5, ACA

appears at e14.5 and collaterals between MCA and ACA appear at e15.5.

Arrowheads point to collateral arteries between MCA and ACA. (**G**) Quantification of

Cx40eGFP/+ brains from different developmental stages reveals that both MCA and

ACA have formed by e14.5. (**H**) Quantification of eGFP+ collateral arteries per

hemisphere shows that the first pial collaterals appear at e15.5. N=3 (e13.5), 3

(e14.5), 2 (e15.5), 2 (e18.5) and 5 (P0 brain). Each dot represents an individual

brain. e, embryonic day; P, postnatal day; MCA, middle cerebral artery; ACA,

anterior cerebral artery; brch., branch. Scale bars: 500μm

1258

1259

Supplemental Figure 4: Visualization of pial artery cell dynamics on microvascular tracks, related to Figure 1

(**A**) Confocal image of e16 embryonic collateral zone showing single recombined artery endothelial cell migration towards the pial collateral within 110mins. Dotted line is the length of micro-vessel between the single recombined cell and the pial collateral. Arrow shows direction of single recombined artery cell migration. (B) Stereo-image of a whole e16 brain showing both hemispheres and the pial watersheds. Insets C and D show artery growth fronts of MCA and ACA respectively. (C) Time-lapse stereo images show growth of MCA branch within 150 minutes. (D) Time-lapse stereo images show appearance of an ACA branch within 90 minutes. MCA, middle cerebral artery; ACA, anterior cerebral artery. Tamoxifen was injected at e13.5. Scale bars: **A**, **B** 100µm; **C**, **D**, 50µm

Supplemental Figure 5: Analyses of MV tracks in pial watershed, related to Figure 2

(**A**, **B**) Confocal images of pial watershed area from *Cx40CreER*; *Rosa26^TdTomato*; *Tie2::eGFP::Claudin5* brains, show TdTomato+ arteries in cyan and eGFP+ microvasculature in magenta. Microvascular (MV) tracks are shown using arrowheads. Orange arrowheads point to MV tracks not covered by artery cells. Green arrowheads point to MV tracks partially or completely covered by artery cells. Tamoxifen was administered at e13.5, and brains were analysed at e16. Scale bars: 100μm

ApjCreER; Rosa26^{TdTomato} Tam at e13.5, Analyses at e16
Apj$^+$ capillary/venous origin of MV tracks

A Pial watershed **B** Cx40 Apj-lineage **C** Cx40 **D** Apj-lineage

E Cx40 **F** Cx40 Apj-lineage **G** Cx40 **H** Apj-lineage

Cx40CreER; Rosa26^{TdTomato}; Tie2::eGFP::Claudin5
Tam at e13.5, Analyses at e16

I Pial watershed
MV tracks express
microvascular endomucin
TdTomato eGFP endomucin

J TdTomato **K** eGFP **L** endomucin

M Pial watershed
MV tracks lack
arterial marker, Cx40
Cx40 eGFP

N eGFP **O** Cx40

P MV tracks have
capillary/venous origin,
retain microvascular properties
and lack arterial features

Cx40$^-$ Endo$^+$ Apj-lineage
MV track capillary arteries

1284

1285

1286

Supplemental Figure 6: Molecular characterization of MV track during pial

collateral formation, related to Figure 2

(**A-H**) Confocal images of pial watershed from *ApjCreER; Rosa26^{TdTomato}* embryos.

MV tracks (arrowheads) lack expression of artery endothelial cell marker Cx40 and

show expression of TdTomato, an indicator for *Apj*-lineage. (**I-L**) Confocal images

show expression of Endomucin protein, a microvascular marker, on TdTomato⁻

eGFP⁺ MV tracks. Arrowheads point to MV tracks. (**M-O**) Confocal images of pial

watershed from *Cx40CreER*; *Rosa26^{TdTomato}*; *Tie2::eGFP::Claudin5* brains show lack

of expression of Cx40 protein, an artery EC marker, on TdTomato⁻ eGFP⁺ MV tracks.

Arrowheads point to MV tracks. (**P**) A cartoon of pial watershed with

Endomucin⁺Cx40⁻ MV tracks. Cyan represents arteries, magenta represents MV

tracks. For all experiments, Tamoxifen was administered at e13.5, and brains were

analysed at e16. Scale bars: 100µm

artery-EC specific deletion of *Cxcr4*
does not impact formation or maintenance of pial collateral artery network

Supplemental Figure 7: Pial artery coverage does not reduce without arterial

***Cxcr4*, related to Figure 4**

(**A**) Quantification of overall pial artery coverage with and without arterial *Cxcr4*. KS

test performed on the distributions generated D value= 0.1944, p value= 0.0001. (**B**)

Quantification of pial collateral artery network with and without arterial *Cxcr4* (N=3 for

control and N=5 for knockout). Each dot represents a biological replicate. Data

shown as mean ± standard deviation. Two-tailed unpaired *t* test, p=0.606. (**C**)

Experimental set up to assess the effect of arterial *Cxcr4* deletion on pial artery

1311   coverage in adults. (**D**, **E**) Representative confocal images of whole adult brains

1312   (from N=5 for each genotype), genetically lineage traced for pre-existing artery

1313   endothelial cells, (**D**) with or (**E**) without arterial *Cxcr4*. ACA, anterior cerebral artery;

1314   MCA, middle cerebral artery; Tam, Tamoxifen. Scale bars: 1mm

1315

Cx40CreER; Rosa26^{TdTomato};Tie2::eGFP::Claudin5 TdTomato (artery cell lineage) eGFP (all ECs)

A

to ACA ← watershed → to MCA

all ECs · artery cells · skeleton of MV tracks with or without TdTomato+ cells

VegfR2^{+/+}

B

to ACA ← watershed → to MCA

all ECs · artery cells · skeleton of MV tracks with or without TdTomato+ cells

VegfR2^{L/L}

C Total length of MV track per watershed

Normalized MV track length per watershed area (AU)

***

VegfR2^{+/+} *VegfR2^{L/L}*

Cx40CreER; Rosa26^{TdTomato} Tie2::eGFP::Claudin5

D No changes in pial capillary density without arterial *VegfR2*

Cx40CreER; Rosa26^{TdTomato} Tie2::eGFP::Claudin5

VegfR2^{L/L} Cx40CreER; Rosa26^{TdTomato} Tie2::eGFP::Claudin5

E Area coverage by eGFP+ capillary ECs (%)

p=0.73

VegfR2^{+/+} *VegfR2^{L/L}*

Cx40CreER; Rosa26^{TdTomato} Tie2::eGFP::Claudin5

1316

1317

Supplemental Figure 8: Analyses of MV tracks in pial watershed, without

arterial VegfR2, related to Figure 4

(**A**, **B**) Higher magnification confocal images of e16.5 watershed region between

ACA and MCA (**A**) with or (**B**) without arterial *VegfR2*. All endothelial cells are in

magenta and artery endothelial cells in cyan. Tamoxifen was injected at e14.5.

Skeleton of all the eGFP⁺ microvascular tracks traced in the e16.5 watershed

through multiple z-slices, using Simple Neurite Tracer. Traced MV tracks occupied

by TdTomato⁺ artery endothelial cells are shown in cyan and MV tracks not occupied

with TdTomato is shown in yellow. (**C**) Quantification of total length of microvascular

tracks per watershed area at e16.5 with or without arterial *VegfR2* (control: 7

watersheds from N=5 embryos, artery endothelial cell specific knockout for *VegfR2*:

5 watersheds from N=5 embryos). Two-tailed unpaired *t* test, p=0.0004. Each dot

represents a selected watershed region from embryonic brain. Data shown as mean

± standard deviation. (**D**) Confocal images of eGFP⁺ capillary ECs (magenta) in

e16.5 pial watershed with or without arterial *VegfR2*. (**E**) Quantification of capillary

EC density with (N=3 brains) or without (N=5 brains) arterial *VegfR2*. Two-tailed

unpaired *t* test, p=0.73. Each dot represents a biological replicate. Data shown as

mean ± standard deviation. ***p < 0.001. Tam, Tamoxifen; MCA, middle cerebral

artery; ACA, anterior cerebral artery; MV, Microvascular; e, embryonic day; EC,

endothelial cell; AU, Arbitrary units. Scale bars: **A**, **B**, 100 µm; **D**, 50µm.

A
Dach1 is highly expressed in pial capillary ECs at P0

TdTomato VegfR2 Dach1

capillaries

artery

TdTomato

VegfR2

Dach1

DAPI

capillaries

No effect on collateral development with deletion of *Dach1* from capillary ECs

B
ApjCreER; Dach1$^{fl/fl}$

MCA and ACA appear

pial collaterals appear

e14.5   e15.5   e16.5   e18.5/P0

Tam

deletion of Dach1 from Apj$^+$ capillaries

immunostaining, whole brain confocal imaging, analyses of knockout efficiency & collateral numbers

C
Dach1 SMA VegfR2   Dach1   SMA   VegfR2   DAPI

Control

Dach1 cap-EC KO

D
Normalized Dach1 expression (AU)

p<0.0001

****

Control   Dach1 cap-EC KO

E
No. of Cx40+ collaterals per hemisphere

p=0.349

control   Dach1 cap-EC KO

1338

1339

Supplemental Figure 9: Dach1-independent pial collateral development, related to Figure 4

(**A**) Dach1 expression (green) in capillaries and arteries in P0 brains. Dotted line marks the TdTomato⁺ (cyan) arterial boundary. VegfR2 marks all vasculature. DAPI marks nuclei. (**B**) Experimental design showing Tamoxifen injection at e14.5, and analyses of collateral numbers in e18.5/P0 brains. (**C**) Confocal images of pial vasculature show absence of Dach1 from capillary endothelial cells. (**D**) Quantification of *Dach1* deletion from pial capillary endothelial cells. N=2 for control and N=3 for knockout. Two-tailed unpaired *t* test, p < 0.0001. (**E**) Quantification of Cx40⁺ pial collaterals between MCA and ACA in control (*Dach1^{fl/fl}*, N=4) and *ApjCreER*; *Dach1^{fl/fl}* (Dach1 cap-EC KO, N=7) brains. Two-tailed Welch's *t* test, p=0.349. Each dot in **D** and **E** represents a biological replicate. Data shown as mean ± standard deviation. ****p < 0.0001. e, embryonic day; cap-EC, capillary endothelial cell; KO, knockout; Tam, Tamoxifen. Scale bar: A, 20μm; C, 25μm

1354

1355

Supplemental Figure 10: VegfR2-independent pruning of neonatal pial collaterals, related to Figure 5

(**A**) Experimental design. (**B**) Confocal imaging of whole brains immunostained for VegfR2 shows presence of the protein in pial arteries. Boxed region in left image is magnified as the right image. Arrows point to VegfR2 protein expression in MCA branches. (**C**) Confocal imaging of whole brains immunostained for VegfR2 shows absence of the protein in pial arteries (enclosed in dotted lines). Boxed region in left image is magnified as the right image. (**D**, **E**) Confocal whole P2 brain images showing TdTomato$^+$ arteries (**D**) with or (**E**) without *VegfR2*. (**F**, **G**) shows quantification of (**F**) pial artery coverage across the entire P2 hemisphere or (**G**) collateral artery coverage in the watershed, in brains with or without arterial VegfR2. Two-tailed unpaired *t* test was performed with p= 0.5. (**H**, **I**) Confocal whole P7 brain images showing TdTomato$^+$ arteries (**H**) with or (**I**) without *VegfR2*. (**J, K**) shows quantification of (**J**) pial artery coverage across the entire P7 hemisphere or (**K**) collateral artery coverage in the watershed, in brains with or without arterial *VegfR2*. Two-tailed unpaired *t* test was performed with p= 0.63. (**L**, **M**) Confocal whole P14 brain images showing TdTomato$^+$ arteries (**L**) with or (**M**) without *VegfR2*. (**N**, **O**) shows quantification of (**N**) pial artery coverage across the entire P14 hemisphere or (**O**) collateral artery coverage in the watershed, in brains with or without arterial *VegfR2*. For all the plots, each dot (N=3 for each group) represents individual mouse brain. Two-tailed unpaired *t* test was performed with p= 0.29. P, postnatal day; KO, knockout; Tam, Tamoxifen. Scale bars: 500μm

1378

1379

Supplementary Figure 11: *In vivo* **imaging of adult proximal arteries, related to**

Figure 5

(**A-D**) Images of pial watershed with pre-existing artery endothelial cells labelled in

TdTomato (cyan) and all endothelial cells labelled in eGFP (magenta). (**A**) Confocal

image from optical window created for longitudinal imaging of *Cx40CreER;*

Rosa26^{TdTomato}; Tie2::eGFP::Claudin5 mouse as shown in Figure **5A**. The image

shows both pial and dural vessels. (**B**) Pial layer segmented out from **A**. red and blue

show pial arteries and veins respectively. (**C**, **D**) Confocal images of pial proximal

arteries from boxed regions shown in **A**. **C** and **D** are proximal arteries connected to

MCA and ACA respectively. Arrow shows a TdTomato negative cell. MCA, middle

cerebral artery; ACA, anterior cerebral artery; PA, proximal artery; DA, distal artery;

col, collateral; brch., branch. Scale bar: A, 200µm; C, D, 50µm

1392

Supplemental Figure 12: Hypoxia induces increase in size of pial collateral

cells, related to Figure 5

(**A, B**) Cranial windows showing pial and dural vessels from mice (**A**) under

normoxia and (**B**) hypoxia. (**C**) Experimental design using *Cx40CreER;*

Rosa26^TdTomato; Tie2::eGFP::Claudin5. (**D**, **E**) Insets shown in **A** and **B** respectively.

Pial collateral artery segments from (**D**) normoxic and (**E**) hypoxic conditions. (**F**)

Quantification of diameter of pial collateral arteries in before and during hypoxia.

Two-tailed unpaired *t* test, p=0.029. (**G**) Quantification of total number of cells within

a pial collateral artery before and during hypoxia. Two-tailed unpaired *t* test, p=0.99.

(**J**) Quantification of average cell size before and during hypoxia. Two-tailed

unpaired *t* test, p=0.046. (**H**, **I**) Cells marked in **D**, and **E** respectively. A

representative cell from normoxic and hypoxic pial collateral segment. In **F**, **G**, and **J**,

each dot represents a biological replicate (N=3). Data shown as mean ± standard

deviation. *p < 0.05. MCA, middle cerebral artery; ACA, anterior cerebral artery; col,

collateral artery; Hx, Hypoxia; d, day; O_2, oxygen. Scale bars: **A**, **B**, 200μm; **D**, **E**,

50μm.

Intra-vital imaging of VegfR2 deficient pial collaterals in adult mice

VegfR2^{L/L} Cx40CreER; Rosa26^{TdTomato}
Tie2::eGFP::Claudin5
TdTomato (pre-existing artery ECs) eGFP (all ECs)

1409

1410

1411

Supplemental Figure 13: VegfR2-independent maintenance of pial collaterals, related to Figure 6

(**A**) Confocal image of a proximal artery from VegfR2 artery specific knockout adult mouse shown in Figure **6**. Arrows point to TdTomato⁻ cell. (**B**) Confocal images of the entire cranial window from 3 different mice show that the overall pial artery network remains unchanged without arterial *VegfR2*. MCA, middle cerebral arteries, ACA, anterior cerebral arteries, Coll, collateral arteries between MCA and ACA. Scale bars: **A**, 100µm; **B**, 500µm.