The response to ketogenic diet in colorectal cancer is mediated by lasting functional alterations of the

gut microbiome

Mina Tsenkova[1], Madita Brauer[1,2], Marat Kasakin[3], Susheel Bhanu Busi[3,4], Maryse Schmoetten[1], Anthoula

Gaigneaux[5], Vitaly Pozdeev[1], Marianne Meyers[1], Eric Koncina[1], Diran Herebian[6], Fabien Rodriguez[1],

Martine Schmitz[1], Laura de Nies[3], Ertan Mayatepek[6], Serge Haan[1], Carine de Beaufort[7], Jérôme Graas[8],

Thorsten Cramer[9], Carole Linster[3], Paul Wilmes[3,5], Elisabeth Letellier[1]

[1] Molecular Disease Mechanisms Group, Department of Life Sciences and Medicine, Faculty of Science,

Technology and Medicine, University of Luxembourg, Esch-sur-Alzette, Luxembourg

[2] Institute for Advanced Studies, University of Luxembourg, Esch-sur-Alzette, Luxembourg

[3] Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Esch-sur-Alzette, Luxembourg

[4] UK Centre for Ecology and Hydrology, Wallingford, United Kingdom

[5] Department of Life Sciences and Medicine, Faculty of Science, Technology and Medicine, University of

Luxembourg, Esch-sur-Alzette, Luxembourg

[6] Department of General Pediatrics, Neonatology and Pediatric Cardiology, Medical Faculty and University

Hospital Düsseldorf, Heinrich Heine University, Düsseldorf, Germany

[7] Pediatric Clinic, Centre Hospitalier de Luxembourg, Luxembourg, Luxembourg

[8] Centre d'Investigation et d'Epidémiologie Clinique, Luxembourg Institute of Health, Hesperange,

Luxembourg

[9] Molecular Tumor Biology, Department of General, Visceral and Transplantation Surgery, RWTH

University Hospital Aachen, Germany

Abstract

Colorectal cancer (CRC) patients have been shown to possess an altered gut microbiome. Diet is

a known microbiome modulator. An attenuating effect of the ketogenic diet (KD) on CRC cell growth has

been previously observed, however the role of the gut microbiome in driving this effect remains unknown.

Here, we describe a reduced colonic tumor burden upon KD consumption in a CRC mouse model with a

humanized microbiome. Importantly, we demonstrate a causal relationship through microbiome

transplantation into germ-free mice, whereby alterations in gut microbial function were maintained in the

absence of continued selective pressure from the KD. Stearic acid is identified as a putative microbiome-

derived anti-cancer metabolite. Taken together, the beneficial effects of the KD are mediated by the gut

microbiome and may be dependent on a pre-existing "non-responder" or "responder" gut microbiome

profile. These results have important implications for future clinical trials evaluating the effects of the

ketogenic diet on human CRC.

Introduction

Worldwide, colorectal cancer (CRC) is the third most prevalent cancer, with over 1.9 million new cases recorded in 2020[1]. CRC incidence and mortality rates have been associated with various lifestyle-related risk factors, in particular dietary habits. Recent years have seen a rise in CRC incidence in persons aged 50 years and under, further underlining the importance of studying these factors[2]. The ketogenic diet (KD) has been shown to possess anti-cancer properties in the context of CRC, and many studies have focused on the importance of the restriction of energy to slow the growth of rapidly proliferating cells and anti-cancer signaling through ketone bodies[3]. One recently published study investigated the effects of the KD on CRC development and described a new mechanism through which KD exerts its anti-cancer effects, namely through the ketone body β-hydroxybutyrate acting on the surface receptor Hcar2, inducing the transcriptional regulator *Hopx*, thus altering gene expression and inhibiting cell proliferation[4]. These studies often focus on the effects of KD on the cancer cells themselves, and rarely on other cells of the tumor microenvironment (TME), such as immune cells[5,6] and cancer-associated fibroblasts (CAFs)[7], or on external factors, such as the gut microbiome. Specifically in CRC, cancer progression is accompanied by a state of dysbiosis of the gut microbiome[8]. The study of the role of the different bacteria in CRC is of particular interest, considering the affected tissue type is in contact with the gut microbiome. Until recently, few studies had investigated the effects of the KD on the gut microbiome in health and disease[9], but this has now changed, with a number of new studies published over the last three years[6,8]. These studies remain largely descriptive regarding dietary-induced gut microbiome changes and the role of the gut microbiome in maintaining the reported anti-cancer effects of the KD in the context of CRC has not yet been investigated.

Results

KD leads to reduced colonic tumor burden and inflammatory response in CRC

To study the relevance of the gut microbiome in CRC development in a model relevant to human

disease, we first established a representative healthy human microbiome by mixing fecal samples from

five healthy donors (5HD; Extended Data Fig. 1a for patient metadata). The achieved 5HD mix

recapitulated the compositions of the individual donor samples, as evidenced by family, genus and species

level analysis of whole genome shotgun sequencing (WGS, Extended Data Fig. 1b-d, respectively), while

fecal metabolite levels varied across samples (Extended Data Fig. 1e). We performed fecal microbial

transplantation (FMT) of the 5HD mix into germ-free (GF) and antibiotic-treated specific-pathogen-free

(SPF) mice in an AOM/DSS CRC model (Fig. 1a and 1b, details can be found in the methods section)

followed by a switch from standard to ketogenic diet after the last cycle of DSS. No significant difference

in body weight between the standard diet (SD) and ketogenic diet (KD) groups was observed throughout

the experiments, although dips in mouse body weight were observed following the antibiotic treatment

and following each cycle of DSS (Extended Data Fig. 2a and 2b). There was no difference in colon length

between groups (Extended Data Fig. 2c, GF). KD-fed mice exhibited a lower colonic tumor burden (Fig. 1c

(GF) and Fig. 1d (SPF)) and a smaller tumor size (Fig. 1e, 1f and 1g (SPF)) than SD-fed mice.

Characterization of full-length lamina propria (LP), mesenteric lymph node (MLN) and spleen

immune cell profiles (Extended Data Fig. 2d and 2e) showed a reduction in almost all LP CD4$^+$ T cell

populations in KD-fed mice. Only LP natural killer T cells and type 3 innate lymphoid cells (GF LP) were

increased – both have been associated to gut homeostasis and health[10]. Finally, the expression of both

subunits of calprotectin (a biomarker of inflammation used in inflammatory bowel disease[11]) was reduced

in the spleens of GF KD-fed mice (Extended Data Fig. 2f and 2g). Nevertheless, it remains unclear whether

the reduction in immune cell populations reflects reduced inflammation induced by the KD or is rather a

direct consequence of the reduced tumor development.

To further assess the importance of KD in modulating inflammation in CRC, we employed a low-

grade inflammatory model, in which mice received four doses of AOM and were randomized into one of

three subgroups - GF, 5HD mix FMT, single CRC patient FMT, each with a KD and SD branch (Extended

Data Fig. 3a, 3b (5HD) and 3c (CRC)). A single CRC donor was used for colonization, rather than a mix of

multiple samples, due to the observed dissimilarities of microbial profiles between patients in our in-

house CRC stool collection. Despite the induction of ketosis in KD-fed mice (Extended Data Fig. 3d), we

observed no significant differences in colonic tumor burden between groups (Extended Data Fig. 3e), nor

in immune cell profiles (Extended Data Fig. 3f), suggesting not only that the KD specifically exerted its anti-

cancer effect in an inflammatory setting, independent of ketosis, likely by reducing inflammation, but also

that the fecal microbiome from this CRC patient did not exhibit a higher potential for CRC induction in this

setting, in contrast to observations made by Wong et al.[12]. Of note, long-term KD feeding induced fatty

liver and fibrosis in all sub-groups (Extended Data Fig. 3g), and as previously reported for long-term KD

consumption[13]. Altogether, these results suggest that the KD exerts an anti-tumorigenic effect in CRC,

potentially by modulating the immune system.

KD induces a distinct host metabolic profile

Ordination analyses based on the underlying β-diversity among host plasma samples collected

from animals undergoing ketogenic dietary intervention in the AOM/DSS inflammatory model of CRC,

revealed clustering based on dietary condition (Extended Data Fig. 4a (SPF) an Extended Data Fig. 4b (GF)).

β-HB plasma levels were elevated in KD-fed mice (Extended Data Fig. 4c (SPF) and 4d (GF)), confirming

ketosis. KD-fed mice presented altered amino acid, fatty acid, sugar, carbohydrate, vitamin, and bile acid

metabolism (Supp. Table 1 and Extended Data Fig. 4e (SPF), Supp. table 2 and Supp Fig. 4f (GF).

Furthermore, mannose was increased in KD-fed mice (Supp. Table 1 (SPF) and 2(GF)). Interestingly,

mannose has been reported to possess antitumor properties[14], impairing tumor cell growth *in vitro* and

in vivo[15], and targeting tumor-associated macrophages[16]. While treatment with β-hydroxybutyrate

showed no effect, treatment with the ketone bodies acetone and sodium acetoacetate tended to reduce

the differentiation of naive T cells into Th17 cells (Extended Data Fig. 4g and 4h), which play an important

role in maintaining gut homeostasis[17] and are affected by KD in a dose-dependent manner[5]. Treatment

with a commercially available lipid mixture mimicking KD (containing arachidonic, linoleic, linolenic,

myristic, oleic, palmitic and stearic acid) led to a dose-dependent reduction in Th17 differentiation

(Extended Data Fig. 4i), as did treatment with some individual fatty acids, such as palmitic and oleic acid

(Extended Data Fig. 4j), suggesting that Th17 cells are susceptible to fatty acid modulation, but that

different fatty acids may exhibit different effects. Collectively, these data indicate that KD induces

systemic functional metabolic changes in the host, which are not limited to the induction of ketosis.

The anti-cancer effects of the KD are mediated by the gut microbiome

The gut microbiome may affect CRC development either by directly interacting with host cells

(attachment, invasion and translocation), or indirectly, through bacterial metabolism (genotoxins, SCFAs

and others)[18]. We first excluded bacterial translocation as a primary factor in the KD-mediated phenotype,

as mucus layer thickness (Extended Data Fig. 5a and 5b (GF)) and LPS levels in plasma (Extended Data Fig.

5c (GF) and 5d (SPF)) remained unaltered upon KD consumption, in line with Ang et al.[5]. To test the causal

role of the gut microbiome in our experimental setup, we performed a similar experiment in humanized

gut microbiome GF mice, however we housed the mice in open-top cages in a gnotobiotic flexible-film

isolator to allow for microbial exchange between groups (Extended Data Fig. 5e, 5f and 5g). Despite

confirmed ketosis, no difference in colonic tumor burden (Extended Data Fig. 5h), nor in colonic tumor

size (Extended Data Fig. 5i and 5j) was observed between groups. Furthermore, fecal microbial

compositions were similar in both groups (Extended Data Fig. 5k), with only a few taxa differentially

abundant between groups (Extended Data Fig. 5l), suggesting that microbial exchange between mice on

KD and SD was sufficient to negate the beneficial effects of the KD. Most importantly, we next performed

cecal microbial transplantation (CMT) of samples collected from the SPF diet mice (Fig. 1d) into a next

generation of SD-fed GF mice undergoing the same cancer model (Fig. 2a, Extended Data Fig. 5m).

Strikingly, KD-CMT mice, exhibited a lower colonic tumor burden than SD-CMT mice (Fig. 2b). Average

colonic tumor size per mouse did not differ between groups (Fig. 2c), however, KD-CMT mice exhibited a

smaller proportion of large tumors (Fig. 2d and e). Altogether, our results demonstrate the causal role of

the gut microbiome in mediating the previously observed KD-driven anticancer effect.

KD induces a distinct fecal microbial profile

Ordination analyses based on the underlying β-diversity among fecal samples collected from KD-

and SD-fed AOM/DSS mice revealed clustering of 5HD mix used for FMT with the individual healthy donor

stool samples and with the murine stool samples collected three days after FMT (TG, Extended Data Fig.

6a (SPF) and Extended Data Fig. 6d (GF)), confirming the validity of the 5HD mix used for transplant and

its engraftment. Furthermore, we observed increasing differences between KD and SD samples over time

(Extended Data Fig. 6b (SPF) and Extended Data Fig. 6e (GF)). Early time points cluster together (T0,

collected at diet change), while later time points diverge (T1-T4). Differential analysis revealed

Akkermansia spp., bacteria which can suppress colorectal tumorigenesis through modulation of the

immune system[19], *Intestimonas* spp*., Lachnoclostridium* spp*., Bilophila* spp. (Extended Data Fig. 6c (SPF)),

Phocea spp., *Faecalitalea* spp. and others (Extended Data Fig. 6f (GF), Supp. Table 3 (SPF) and 4 (GF)) to

be more abundant in KD, while *Bifidobacterium* spp*., Turicibacter* spp. (SPF)*, Barnesiella* spp*.,*

Eisenbergiella spp*., Ruminococcus 1* and *2* spp. (GF) and others were depleted. Furthermore, the

differential abundance of various members of *Barnesiellaceae* spp*., Erysipelotrichaceae* spp*.,*

Lachnospiraceae spp. and *Ruminococcaceae* spp. suggested that dietary composition affects bacterial

members of the same families in different manners.

Investigation of the engrafted gut microbial communities at experimental endpoint revealed

distinct KD- and SD-associated microbiomes (Fig. 3a and 3b, Extended Data Fig. 6g, 6h (GF), 6i and 6j), as

well as KD-CMT- and SD-CMT-associated microbiomes (Fig. 3a and 3b, Extended Data Fig. 6i and 6j).

Although a selective pressure of the KD was observed, gut microbial composition differed between cecal

donor and recipient mice, likely due to the absence of continued selective pressure from the KD, or loss

of more fragile and hard-to-engraft bacteria (Fig. 3a). Metagenome-assembled genomes (MAGs),

representing abundant members of the microbial community, allowed us to achieve species-level

resolution. Noteworthy, some MAGs annotated as CRC-associated bacteria, such as *Rikenellaceae* spp.

(light blue) and *Muribaculaceae* spp. (dark blue, Fig. 3c, Extended Data Fig. 6k (GF)), were detected in SD

groups but not the KD groups. Interestingly, various *Alistipes* spp. were differentially abundant in all

experiments. Two MAGs (*Alistipes* and *COE* species) were only identified in KD-CMT, while 12 MAGs were

only identified in SD-CMT, among which *CAG-873*, *CAG-48* and *Parabacteroides distasonis* (Fig. 3c), also

identified in the donor SPF dietary experiment.

Distinct fecal microbial functional profiles are maintained upon FMT

Furthermore, as microbiome-mediated effects are based on the functional and metabolic

capacities of different microbes, we also investigated KD-induced functional changes. β-diversity analysis

based on KEGG and Pfam ordinations of SPF fecal samples revealed a clear separation between groups

(Extended Data Fig. 7a (KEGG) and 7b (Pfam)), suggesting selective pressure of the diet on the intestinal

microbiota. This could also be observed in the CMT experiment, where KD-CMT clustered separately from

SD-CMT, despite being maintained on SD (Extended Data Fig. 7c (KEGG) and 7d (Pfam)). Both SPF KD-fed

and KD-CMT mice exhibited a higher number of less abundant genes/Pfams than more abundant

genes/Pfams, suggesting a KD-induced functional depletion (Extended Data Fig. 7e (SPF, KEGG), 7f (CMT,

KEGG), 7g (CMT, KEGG) and 7h (CMT, Pfam)). Accordingly, gene set enrichment (Fig. 3d and Extended Data

Fig. 7i) and overrepresentation analysis (Fig. 3e) of identified KEGG orthologs revealed differences in

carbohydrate metabolism between the treatment groups in the SPF diet and CMT experiments, such as

reduced starch and sucrose metabolism under ketogenic conditions. In line with this, genes assigned to

Pfams associated with carbohydrate metabolism accounted for a large proportion of over- and under-

represented genes under ketogenic conditions (Extended Data Fig. 7j (SPF) and Extended Data Fig. 7k

(CMT)). Furthermore, we found differential abundance of several CAZymes between dietary conditions,

which are known to play a crucial role in the building and breaking down of complex carbohydrates for

use by the organism[20], with an overall increase in glycosyltransferases (GT; Extended Data Fig. 7l). Filtering

the SPF diet and CMT datasets for KEGGs, Pfams and CAZymes, which were simultaneously enriched or

depleted, yielded a list of 15 genes with varying functions, of which five were more abundant and ten

were less abundant in ketogenic conditions (Extended Data Fig. 7m, Supplementary Table 5). Altogether,

this data underlines the selective pressure of the KD on the gut microbiome, and its capacity to shape

long-term overall gut microbial function.

The microbial metabolite stearate is elevated in ketogenic conditions and exhibits anticancer

effects

In order to investigate the link between functional alterations of the gut microbiome and the

observed cancer phenotype, we investigated fecal metabolic profiles. Analysis of fecal metabolites

revealed distinct clustering of samples by condition in both the SPF dietary and CMT experiments (Fig. 4a

and 4b, respectively). Mice in the ketogenic conditions presented altered amino acid, fatty acid,

carbohydrate and SCFA profiles (Supp. Tables 6 and 7), albeit to a lesser extent in the CMT experiment. A

single detected long-chain fatty acid, namely octadecanoic acid (also known as stearic acid), was elevated

in KD (Fig. 4c) and KD-CMT (Fig. 4e) mice. Strikingly, stearic acid levels correlated with colonic tumor

burden in the SPF dietary experiment (Fig. 4e), with a similar trend observed in the CMT experiment,

suggesting a functional relevance of long-chain fatty acid metabolism in CRC. In addition, associations

between the abundances of all identified KEGG orthologs and tumor counts in the SPF dietary experiment

identified a correlation between colonic tumor burden and some lipid-metabolism-associated KEGGs,

such as FabH, 3-oxoacyl-[acyl-carrier-protein] synthase III (K00648), FabB, 3-oxoacyl-[acyl-carrier-protein]

synthase I (K00647), FabZ, 3-hydroxyacyl-[acyl-carrier-protein] dehydratase (K02372), and FabF, 3-

oxoacyl-[acyl-carrier-protein] synthase II (K09458) (Fig. 4f and Extended Data Fig. 8a). Indeed, core fatty

acid biosynthesis KOs (Extended Data Fig. 8b) were less abundant in the ketogenic diet group of the SPF

diet experiment, except for the different enoyl-carrier ACPs and the thioesterases, of which one or more

can be found in most bacterial genomes (Extended Data Fig. 8c). Reflecting their essential function, the

core enzymes of the fatty acid synthesis pathway correlating with tumor counts were found in both

Bacteroidetes and Firmicutes genomes. In KD, contigs carrying K00647, K00648, K09458 and K02372 were

assigned to *Akkermansia* spp., *Flavonifractor* spp. and uncharacterized Firmicutes genera, whereas many

SD contigs were assigned to Bacteroidetes genera (*Bacteroides* spp*., Parabacteroides* spp*., Muribaculum*

spp. and uncharacterized Bacteroidetes; Extended Data Fig. 8d).

Given that different bacterial taxa are known to produce and consume different fatty acids, which

are used for membrane biogenesis and as signaling molecules, we further investigated which bacterial

taxa may be responsible for the release of fatty acids and which bacteria may be able to uptake free fatty

acids. Interestingly, detection of long-chain fatty acids in different bacterial cultures (Extended Data Fig.

8e; re-analysis of the publicly available dataset from the Sonnenburg lab[21]), revealed that – in line with

the shifts in the microbial communities observed under KD (Fig. 4h and Extended Data Fig. 9f) - the

Ruminococcaceae and *Lachnospiraceae* families, both belonging to the class Clostridia, were consistent

producers of stearic acid, while *Tanerellaceae* and *Bacteroidaceae,* both belonging to the class

Bacteroidia, were consistent consumers (Fig. 4g). The *Clostridiaceae* family was composed of both

producers and consumers of stearic acid (Fig. 4g). Importantly, acyl-CoA thioesterase I TesA (K10804), a

key thioesterase responsible for the release of fatty acids from their acyl carrier, alongside the fatty acid

kinase FakA (K07030) used by Gram-positive bacteria to activate exogenous fatty acids, were inversely

correlated with colonic tumor burden in the SPF dietary experiment (Fig. 4i). A larger proportion of K10804

was assigned to Clostridia in the ketogenic condition, and Bacteroidia in the standard condition in both

experiments (Fig. 4i), indicating that the lower levels of luminal stearate detected in standard conditions

may be due to the increased abundance of Bacteroidia, which, although they harbor the gene for fatty

acid release, are known stearate consumers[21]. K07030 was largely attributed to Clostridia and other

Firmicutes (Fig. 4i), suggesting that these bacteria can uptake exogenous fatty acids (for example from the

ketogenic diet) and might further elongate them into other long-chain fatty acids for downstream use or

release. The observed contrast between the abundance of fatty acid release KOs and actual

production/consumption of stearic acid by different bacteria as detected in the Sonnenburg dataset

suggests that different bacteria contribute differently to the overall fatty acid pool. Therefore, despite no

difference in the overall abundance of K10804 and K07030 between dietary conditions (Extended Data

Fig. 8c), when assigned to stearate-producing bacterial families (as detected in the Sonnenburg dataset),

these KOs may result in increased levels of stearate, as in the ketogenic conditions, while when assigned

to stearate-consuming families, they may result in a decrease in stearate levels. To assess the putative

anti-cancer properties of the long-chain fatty acid of interest identified in the dietary and CMT

experiments, we treated various CRC cell lines with increasing concentrations of stearic acid. We observed

a dose-dependent reduction in cell growth (Fig. 4j, 4k, 4l, Extended Data Fig. 8g, 8h, 8i), highlighting stearic

acid as an anti-cancer metabolite.

Altogether, these results suggest that KD consumption can lead to lasting functional effects on

the gut microbiome, affecting the bacterial secretome, and persisting even when the selective pressure

of the KD itself is no longer present.

Discussion

In this study we identified the gut microbiome as a driving factor in the anti-cancer effect of KD in

CRC. We observed that KD alters gut microbial function, namely bacterial carbohydrate metabolism and

the fatty acid pool. Upon KD consumption and following microbial transplantation, we observed an

increase in fecal octadecanoic acid, a long-chain fatty-acid, which exhibited anti-cancer properties *in vitro*.

The anticancer properties of the ketone body β-HB have been reported in several instances[3,4],

however, we observed the absence of a KD-induced anticancer effect in the absence of robust gut

microbial changes, despite confirmed induction of ketosis in these mice. Thus, ketosis alone seems to be

insufficient to ensure CRC suppression, similarly to other studies showing no KD benefits in the context of

cancer, despite the induction of ketosis[3,22].

Our results instead suggest that the anti-cancer effect of KD is based on gut microbial function,

rather than composition. The importance of microbial function rather than microbial composition of the

intestinal flora can be observed in other contexts as well. In melanoma, three different studies

demonstrated that the microbiome was responsible for ensuring the response to checkpoint inhibitor

therapy[23–25]. Interestingly however, these studies highlighted distinct bacterial species underlying this

response, suggesting that microbiome-encoded functions rather than individual bacterial species are

important in driving therapy response.

In regard to our own study, the depletion of dietary carbohydrates is known to provide a growth

advantage for certain bacteria, such as *Alistipes* spp., *Bilophila* spp. and mucin-degraders like *Akkermanisa*

spp., whereas *Bifidobacterium* spp. and some *Bacteroides* spp. are typically depleted[5,26], most likely due

to their preference for dietary carbohydrates over host glycans[27–29]. The shift from taxa dependent on

dietary carbohydrates taxa to taxa able to use alternative energy sources is reflected in the CAZyme profile

shift in KD and KD-CMT mice. For instance, the glycosylhydrolase GH101, enriched in all three of our

metagenomic datasets, is mucin-specific, and is required to cleave core-1 *O*-linked glycans[30]. In contrast,

the glycosylhydrolases GH50, present in the genomes of many *Bacteroides* spp. and *Proteobacteria* spp.,

and GH128, amongst others present in the genomes of the strongly reduced *CAG-485* and *CAG-873*

members, are specific for dietary carbohydrates[31] and accordingly found in reduced abundance following

KD intervention. These specific effects clearly demonstrate how diet systematically shapes the taxonomic

composition of the microbiome, driven and accompanied by a functional adaptation of the microbiome

to the new environment.

Although still an ongoing debate, the link between mucin-degrading bacteria and gut barrier

function is well-described. While overgrowth of mucin-degrading bacteria are known to damage the

epithelium[32], microbial mucin metabolism is considered beneficial as it is known to change the

glycosylation profile of the intestine[33,34], and increase mucin turnover[35]. Indeed, no mucus thinning could

be observed in KD-fed mice.

Analysis of the fecal metabolomes in the dietary intervention and CMT experiments identified

differences in the fatty acid pool. Amongst others, microbiome-derived fatty acids have an

immunomodulatory potential[36,37]. Interestingly, both *A. muciniphila* and *R. gnavus* produced higher levels

of palmitic and stearic acid when grown in the presence of mucins, linking bacterial fatty acid metabolism

to mucin physiology. In addition to their essential role in membrane biogenesis, fatty acids produced by

bacteria have signaling functions and play an essential role in host-microbe interactions[36]. The fatty acid

profile is highly species-, or even strain-, and substrate-dependent and various microbial-derived fatty

acids have different effects linking bacterial fatty acid metabolism to host health and disease[36].

Stearic acid, a long-chain fatty-acid, can originate from the diet, the host[38] and the gut

microbiome[39]. Stearic acid is normally desaturated by the host enzyme SDC1[40], is downregulated upon

reduced glucose intake and has been linked to inflammation[41–43]. Recently, Kindt et al. showed that

microbiome-derived acetate activated SDC1 and promoted fatty acid desaturation[44]. Furthermore, certain

members of the gut microbiome can directly affect stearic acid levels by consuming it as an exogenous

fatty acid for membrane lipid synthesis[21], producing it *de novo*[45], transforming it from oleic acid, or as a

product of mono- and polyunsaturated acid degradation[46,47]. In our experiments, as fecal stearic acid

levels were increased not only upon KD consumption, but also in KD-CMT, it is likely that the shift in the

microbial community is contributing to the changes in the fatty acid pool. Importantly, potential stearate-

producing members were enriched by the KD, whereas consumers were depleted (Fig. 4g and 4h).

Furthermore, several studies investigating the properties of stearic acid have demonstrated its

anti-cancer effects, both *in vitro* and *in vivo*. Stearic acid inhibited colony formation in different human

cancer cell lines[48], reduced tumor burden in rat model of mammary carcinoma[49], possibly by inducing

apoptosis via signaling through protein kinase C[50]. Finally, a recent study was able to not only validate the

suppression of colorectal adenocarcinoma by stearic acid, but also its synergy with 5-FU treatment[51]. This

effect might not be limited to stearic acid and its implications for human health and CRC in particular need

further investigation.

Based on our observations in the context of this study, the gut microbiome plays a central role in

ensuring the anti-cancer effects of the KD in the context of a murine model of inflammation-driven CRC.

Nevertheless, within this experiment, we can observe a more pronounced anti-cancer phenotype for

some mice than for others. This might indicate that the initial gut microbiome profile may determine

whether an individual is a "responder" or a "non-responder" to the KD in the context of CRC. The concept

of "responders" and non-responders" has been explored, albeit to a relatively limited extent, in epilepsy

patients who undergo KD therapy for the reduction of seizures[52]. Furthermore, the response to some

cancer treatments, such as immune checkpoint inhibition depend on gut microbial composition[53], and

ketogenic dietary regimens may be able to enhance the effects of these therapies[8]. Future studies

identifying biomarkers to predict response to dietary intervention based on the resident gut microbiome

and studies aiming to transform a "non-responder" microbiome into a "responder" microbiome would be

a significant step forward towards personalized patient care.

Figure 1. Ketogenic diet leads to lower colonic tumor incidence and smaller tumor size in an inflammatory model of CRC. | a. Schematic representation of the dietary AOM/DSS experimental setup in the GF (top) and SPF (bottom) facilities (created with BioRender.com). **b.** Compositions of the KD and SD, with % of calories from the indicated source. **c.** Colonic tumor burden in GF KD- and SD-fed mice. **d-f.** Colonic tumor burden (**d**), normalized average colonic tumor surface area per mouse (**e**) and tumor surface area distribution (small= first tertile, medium=second tertile, big=third tertile; Chi-squared test χ2 (2, N = 87) = 3.4, p = 0.184; **f**) in SPF KD- and SD-fed mice. Data in (**c**) shows n=6 mice per condition from one experiment. Data shows n=8 and 9 mice in the KD-fed condition and the SD-fed conditions respectively, pooled from two independent experiments in the SPF facility. Two-tailed Mann-Whitney test, **p<0.01, ***p<0.001. **g.** Representative images of the distal portion of the colon from one KD- and one SD-fed mouse in the SPF facility. Scale bar = 1mm.

Figure 2. The tumor-inhibitory effect of ketogenic diet is mediated by the gut microbiome. | a.
Schematic representation of the cecal microbial transfer experimental setup in the GF facility (created
with BioRender.com). **b-e.** Colonic tumor burden (**b**), average colonic tumor surface area per mouse (in
pixels) (**c**), and tumor surface area distribution (small= first tertile, medium=second tertile, big=third
tertile; Chi-squared test χ2 (2, N = 384) = 10.3, p = 0.00586; **d**), in KC and SC recipient mice. n=16 KC
recipient mice and n=13 SC recipient mice. Cecal matter from n=8 different donor mice per condition was
used for CMT. Data shown as mean±SD in (**b**) and (**c**). Two-tailed Mann Whitney U test, **p<0.01,
***p<0.001 in (**b**) and (**c**). **e.** Representative images of the distal portion of the colon from one KC and SC
recipient mouse. Scale bar = 1mm.

Figure 3. Ketogenic diet alters gut microbiome composition and functions. | a-c. β-diversity as assessed

by NMDS (95% confidence interval) of Euclidean distance (**a**), relative abundance of bacterial families (**b**),

and highly characteristic bacterial MAGs (detected in at least six mice in one condition but not more than

three mice in the other (diet experiment); detected in at least four mice in one condition but not more

than three mice in the other (CMT experiment), **c**) detected in KD, SD, KC and SC murine fecal samples at

endpoint, as analyzed by WGS. In (**c**), *Rikenellaceae* are highlighted in light blue and *Muribaculaceae* in

dark blue. **d-e.** Overrepresentation (KEGG, input threshold log2FC > 2, **d**) and gene set enrichment analysis

(GSEA, KEGG) of differentially abundant genes (threshold p.adj. < 0.05, **e**) between KD, SD, KC and SC fecal

samples at endpoint. In (**d**), data point size corresponds to gene ratio. Data shows results from n=8 SPF

KD-fed mice, n=9 SPF SD-fed mice, pooled from two independent experiments, and n=8 KC and n=8 SC

mice from one experiment.

Figure 4. The microbial metabolite stearate is elevated in ketogenic conditions and exhibits anticancer

effects. | a-b. PCA (95% confidence interval) of identified metabolites from fecal samples from SPF KD-

and SD-fed mice (**a**) and from KC and SC recipient mice (**b**), as analyzed by united GC-MS and LC-MS. **c-d.**

Scaled octadecanoic acid (stearic acid) levels detected in fecal samples from SPF KD- and SD-fed mice(**d**)

and from KC and SC recipient mice (**e**), as detected by LC-MS. **e.** Correlation of colonic tumor burden with

fecal octadecanoic acid levels in SPF KD- and SD-fed mice (red) and in KC and SC recipient mice (blue).

Two-sided Pearson correlation testing was used. **f.** Correlation analysis (Pearson) between KO abundance

and colonic tumor burden in KD- and SD-fed mice. Vertical blue dotted lines represent thresholds for

correlation coefficients (r) above and below 0.65 and -0.65, respectively. The horizontal gray dotted line

represents the threshold for statistical significance -log10(0.05). KOs involved in fatty acid metabolism

appear labelled on the volcano plot and are represented in different colors (pink=synthesis,

orange=activation and blue=β-oxidation). Of those, significantly correlated KOs are labelled in light blue

font. **g.** Levels of stearic acid measured in bacterial cultures (in relation to sterile growth medium, shown

as $\log_2$) of the Sonnenburg dataset. Members of the class Clostridia are listed in red, while members of

the class Bacteroidia are listed in blue. Total read counts detected in murine fecal samples from KD-, SD-

fed, KC and SC recipient mice at endpoint of members listed in bolded text are shown in (**h**). **h.** Relative

abundance (total read counts) of stearate-producing and -consuming families of interest detected in

murine fecal samples from KD-, SD-fed, KC and SC recipient mice at endpoint selected from (**g**). **i.**

Correlation (Pearson) between colonic tumor burden and the selected KOs of interest K07030 and K10804

(**top**); Relative taxonomic assignment of contigs encoding for the selected KOs of interest K07030 and

K10804 at family level (**bottom**) detected in murine fecal samples from KD-, SD-fed, KC and SC recipient

mice at endpoint. Members of the class Bacteroidia are shaded in blue, while members of the class

Clostridia are shaded in red. **j-l.** Scaled HCT 116 (**j**), HT29 (**k**) and Caco2 (**l**) confluence at 72 hours after

treatment with stearic acid or corresponding controls (100µM, 150 µM, 200 µM, 250 µM, 300 µM). Data

shows n=3 (for HCT116 and HT29) and n=2 (for Caco2) independent experiments (indicated by different

datapoint shapes), each with eight technical replicates per condition. Two-sided Mann-Whitney test in (**c**)

and (**d**); Wilcoxon signed-rank test with Bonferroni correction in (**f**); ns p > 0.05, * p ≤ 0.05, ** p ≤ 0.01,

*** p ≤ 0.001, **** p ≤ 0.0001. Two-way ANOVA with multiple comparisons in (**j**), (**k**) and (**l**); **p<0.01,

***p<0.001. Data shown in (**a**), (**c**), (**e, red**), (**f**), (**h, red and black**) and (**i**) are from n=9 SPF SD-fed and n=8

SPF KD-fed mice, pooled from two independent experiments. Data shown in (**b**), (**d**), (**e, blue**) and (**h, gray**

and orange) are from n=8 mice per condition.

Figure 5. Functional ketogenic-diet-induced changes in the gut microbiome are responsible for its anti-

cancer properties.

Ketogenic diet consumption leads to lasting functional changes in the gut microbiome, such as alterations

in bacterial energy metabolism and in the long-chain fatty acid pool. The altered gut microbiome is

sufficient to reduce CRC development.

Extended data figure 1. | a. Metadata of fecal sample donors. M=male, F=female, N/A=not applicable,

UNKN=unknown. **b-d.** Relative abundance of the top 20 most abundant bacterial families (**b**), genera (**c**)

and species (**d**) in human stool samples (CLR units). **e.** Relative metabolite levels detected in human fecal

samples, analyzed by untargeted LC-MS.

Extended data figure 2. | a-b. Mouse body weight (grams) over time (weeks) in the SPF (**a**) and GF (**b**)

dietary experiments. Weight was not recorded between week five and ten as no scale was available in the

facility. **c.** Colon length (cm) in the GF dietary experiment at experimental endpoint. **d-e.** Heatmap of KD-

induced changes in the immune cell compartment in relation to SD (log2foldchange) in the SPF (**h**) and GF

(**i**) diet experiments. Stars overlaid on the heatmaps indicate *p<0.05, least squared means analysis. **f-g.**

Normalized S100A8 (**f**) and S100A9 (**g**) expression in the spleens of mice from the GF diet experiment, as

determined by qPCR. Data is shown as mean±SD in (**a**), (**b**), (**c**), (**f**), and (**g**). Mann-Whitney U in (**c**), (**f**), and

(**g**), * p<0.05, ** p<0.01, *** p<0.001. Data in (**a**) shows n=8 and 9 mice in the KD-fed condition and the

SD-fed condition respectively, pooled from two independent experiments in the SPF facility. Data in (**b**),

(**c**), (**d**), and (**e**) shows n=4 mice in the SD-fed condition and n=6 mice in the KD-fed condition from one

experiment in the GF facility. Data in (**f**) and (**g**) shows n=4 mice per condition from one experiment in the

GF facility.

Extended data figure 3. | a. Schematic representation of the dietary AOM experimental setup in the GF

facility (created with BioRender.com). **b-c.** Mouse body weight (grams) over time (days) in mice

administered 5HD (**b**) and CRC (**c**). **d.** Ketosis levels in plasma (mmol/L), measured using a handheld

ketometer. **e.** Colonic tumor burden at endpoint. **f.** Heatmap of immune cell population phenotyping of

intra-epithelial lymphocytes (IEL), lamina propria (LP), mesenteric lymph nodes (MLN) and spleens of KD-

fed mice in relation to SD-fed mice in GF mice (GF), in mice gavaged with a mixture of stool samples from

five healthy donors (5HD), and in mice gavaged with stool from a single CRC patient (CRC). Least-squared

means analysis was performed on each combination of tissue and cell immune cell type using a linear

model, however no statistically significant differences were found. **g.** One representative image of a

section of liver stained with H&E per condition. Scale bar=500μm. Data in (**b**), (**c**), (**d**) and (**e**) shown as

mean±SD. Mann-Whitney U test in (**d**) and (**e**), not significant (ns), * p<0.05, ** p<0.01, *** p<0.001. Data

in (**b**), (**c**), (**d**), (**e**) and (**f**) shows n=7 and 6 mice in the GF SD and KD conditions respectively, n=6 and 4

mice in the 5HD SD and KD conditions respectively, and n=4 and 5 mice in the CRC SD and KD conditions

respectively.

Extended data figure 4. | a-b. PCA (95% confidence interval) of plasma metabolites from mice in the SPF

(**a**) and GF (**b**) diet experiments, as analyzed by united GC-MS and LC-MS analyses (**a**) and GC-MS analysis

alone (**b**). **c-d.** Relative intensity of β-HB in plasma from mice in the SPF (**c**) and GF (**d**) diet experiments,

as analyzed by GC-MS. **e-f.** Log$_2$foldchange of identified plasma bile acids in KD-fed mice in relation to SD-

fed mice in the SPF (**e**) and GF (**f**) facilities, as analyzed UHPLC-MS. **g.** Schematic representation of the

experimental setup of a Th17 cell differentiation assay (created with BioRender.com). **h-j.** Th17

differentiation upon treatment with β-HB (10mM, 50mM, 100mM), acetone (1mM, 10mM, 100mM) or

sodium acetoacetate (10mM, 100mM, 250mM) in (**h**), a lipid mixture (1%, 2%, 5%) in (**i**), BSA-complexed

palmitic and oleic acid (10μm) in (**j**). Data in (**c**), (**d**), (**h**), (**i**) and (**j**) is shown as mean±SD. Two-tailed Mann-

Whitney U test in (**c**) and (**d**); Two-tailed unpaired T test with FDR correction in (**e**) and (**f**); Paired ordinary

one-way ANOVA in (**h**), (**i**) and (**j**); ns=not significant, * p<0.05, ** p<0.01, *** p<0.001. Paired ordinary

one-way ANOVA comparing each set of ketone body dilutions to the control yielded no statistical

significance in (**h**). Data in (**a**), (**c**) and (**e**) shows n=9 mice in the SD-fed condition and n=8 mice in the KD-

fed condition. Data in (**b**), (**d**) and (**f**) shows n=4 mice in the SD-fed condition and n=5 mice in the KD-fed

condition. n=6 mice in (**c**) and (**d**); n=5 mice in (**e**). Abbreviations: not detected (n.d.), cholic acid (CA),

hyocholic acid (HCA), chenodeoxycholic acid (CDCA), deoxycholic acid (DCA), 7-oxo-deoxycholic acid (7-

oxo-DCA), murideoxycholic acid (MDCA), lithocholic acid (LCA), ursodeoxycholic acid (UDCA),

hyodeoxycholic acid (HDCA), muricholic acids (α-MCA, β-MCA and ω-MCA), taurocholic acid (TCA),

taurohyocholic acid (THCA), taurochenodeoxycholic acid (TCDCA), taurodeoxycholic acid (TDCA),

taurolithocholic acid (TLCA), tauroursodeoxycholic acid (TUDCA), taurohyodeoxycholic acid (THDCA),

tauro-α-muricholic acid (T-α-MCA), tauro-β-muricholic acid (T-β-MCA) and tauro-ω-muricholic acid (T-ω-

MCA).

Extended data figure 5. | a-b. Colonic inner mucus layer thickness (in pixels, **a**) and representative images of Alcian Blue staining of colonic cryosections (**b**) in GF KD-fed and SD-fed mice. **c-d.** LPS (endotoxin units (EU) per milliliter) in plasma from GF (**c**) and SPF (**d**) KD-fed mice and SD-fed mice measured by ELISA. Panels **a-d** relate to experiments shown in Fig. 1. **e.** Schematic representation of the experimental setup in the GF facility, in which mice were housed in open-top cages in a gnotobiotic flexible-film isolator (created with Biorender.com). **f.** Ketosis levels in mouse plasma (mmol/L), measured using a handheld ketometer. **g.** Mouse body weight (grams) over time (weeks). **h-j.** Colonic tumor burden (**h**), average tumor surface area per mouse (pixels, **i**) and representative images of colonic tumors, scale bar=1mm (**j**) from mice at experimental endpoint. **k.** β-diversity as assessed by PCoA of Bray-Curtis distance in mouse fecal samples collected at two time points. **l.** Differentially abundant bacterial families (coefficient, CLR units) in mouse fecal samples, estimated using a generalized linear model on CLR data (p.adj<0.05 (Benjamini-Hochberg correction), max. abundance>0.05), with the standard diet condition as the reference. **m.** Mouse body weight (grams) over time (days) in the CMT experiment. Data is shown as mean±SD in (**a**), (**c**), (**d**), (**f**), (**g**), (**h**), (**i**) and (**m**). Two-tailed Mann-Whitney U test in (**a**), (**c**), (**d**), (**f**), (**h**) and (**i**); * p<0.05, ** p<0.01, *** p<0.001. Data in (**a**), (**c**) and (**d**) shows n=4 mice per condition. Data in (**f**), (**g**), (**h**), (**i**), (**k**) and (**l**) shows n=9 mice in the SD-fed condition and n=8 mice in the KD-fed condition, pooled from two independent experiments. Data in (**m**) shows 16 mice per condition.

Extended data figure 6. | a-f. β-diversity as assessed by PCoA (95% confidence interval) of Bray-Curtis

distance in five individual healthy donor stool samples, in a mix of the five healthy donor stool samples

(5HD mix) and in murine fecal samples three days after gavage (TG, **a** and **d**); β-diversity as assessed by

PCoA (95% confidence interval) of Bray-Curtis distance in murine fecal samples collected over time after

diet change (**b** and **e**); Abundance of selected genera which are differentially abundant at at least one

time point (using a generalized linear model on CLR, **c** and **f**) in fecal samples collected over time after diet

change from SD- and KD-fed mice in the SPF (**a**, **b** and **c**) and GF (**d**, **e**, and **f**) dietary experiments (T0 at

diet change, T1-T4 weeks after diet change), as analyzed by 16s rRNA sequencing. **g.** PCA of WGS

sequencing of fecal samples collected at endpoint from GF KD and SD-fed mice. **h.** Relative abundance of

bacterial families (**h**) detected in fecal samples from GF KD- and SD-fed mice at endpoint, as analyzed by

WGS. **i-j.** Relative abundance of bacterial genera (**i**) and orders (**h**) detected in fecal samples collected at

endpoint from GF and SPF KD- and SD-fed mice, as well as KC and SC recipient mice, as analyzed by WGS.

k. Highly characteristic bacterial MAGs (detected in at least three mice in one condition but not more than

two mice in the other), detected in GF KD- and SD-fed murine fecal samples at endpoint, as analyzed by

WGS. Data shows samples collected from n=9 SPF SD-fed mice and n=8 SPF KD-fed mice, pooled from two

independent experiments in (**a**), (**b**), (**c**), (**i**) and (**j**); n= 8 KC and n=8 SC recipient mice from one experiment

in (**i**) and (**j**); n=3 GF SD and n=6 GF KD mice from one experiment in (**d**), (**e**), (**f**), (**g**), (**i**), (**j**) and (**k**).

Extended data figure 7. | a-d. PCA (95% confidence interval) of the predicted functional features of the

gut microbiome (KEGG, **a** and **c**) and (Pfam, **b** and **d**) of SPF KD- and SD-fed mice (**a** and **b**) and KC and SC

recipient mice (**c** and **d**) at endpoint. **e-h.** Differences in gene expression profiles mapped to the KEGG (**e**

and **f**) and Pfam (**g** and **h**) databases detected in fecal samples from KD- and SD-fed mice (**e** and **g**), and KC

and SC recipient mice (**f** and **h**), at endpoint. Blue data points in (**g**) and (**h**) represent carbohydrate-

metabolism-related Pfams, solid data points represent Pfams above both defined thresholds, while empty

datapoints represent Pfams below these thresholds. Pfams named in (**h**) overlap with those named in (**g**).

i. Overrepresentation analysis (KEGG, threshold log2FC > 2, p.adj. < 0.05) of genes detected in fecal

samples collected at endpoint from SPF KD- and SD-fed mice, and KC and SC recipient mice. Gene sets

related to energy metabolism are shown as circular data points, while all other gene sets are shown as

triangular data points. Data point size corresponds to gene ratio. **j-k.** Differences in gene expression

profiles mapped to the DBcan database detected in fecal samples from KD- and SD-fed mice (**j**), and KC

and SC recipient mice (**k**) at endpoint. **l.** Abundance of main CAZyme families (summed transcripts per

million (TPM)) in fecal samples from KD- and SD-fed mice, and KC and SC recipient mice, at endpoint.

Boxplots show medians with 1^{st} and 3^{rd} quantiles. The whiskers from the hinges to the smallest/largest

values represent 1.5*inter-quartile range (IQR). Wilcoxon signed-rank test with Holm correction in (**f**), ns

$p > 0.05$, * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$. Abbreviations: auxiliary activities (AA),

carbohydrate binding molecule (CBM), glycoside hydrolase (GH), glycosyl transferase (GT), polysaccharide

lyase (PL) and S-layer homology (SLH) families. **m.** Abundance (log2foldchange) of genes simultaneously

enriched or depleted in the KEGG, Pfam and CAZyme databases in fecal samples from SPF KD- and SD-fed

mice, and KC and SC recipient mice, at endpoint. In (**e**), (**f**), (**g**), (**h**), (**j**) and (**k**), the horizontal lines

correspond to a three-fold increase (right) or decrease (left) in KD in relation to SD, in KC in relation SC.

The vertical lines correspond to p<0.05, the defined cutoff for statistical significance (DESeq2). Red data

points represent statistically significantly (p<0.05) differentially expressed genes with three-fold up- or

down-regulation. Data shows n=8 SPF KD-fed mice and n=9 mice SPF SD-fed mice, pooled from two

independent experiments, and n=8 KC and n=8 SC recipient mice from one experiment.

Extended data figure 8. | a. Correlation (Pearson) between colonic tumor burden of KD and SD-fed mice and the selected KOs of interest K00647, K00648, K02372 and K09458. **b.** Schematic representation of type II bacterial fatty acid synthesis created with BioRender, adapted from Yao et Rock (2017)[39]. Core enzymes are shown with their corresponding KOs. The abundance of bolded KOs correlates with colonic tumor burden in KD- and SD-fed mice. Abbreviations: acetyl-CoA carboxylase enzyme complex (AccABCD), β-ketoacyl-ACP synthase 1 (FabB), malonyl-CoA:ACP transacylase (FabD), β-ketoacyl-ACP synthase II (FabF), β-ketoacyl-ACP reductase (FabG), β-ketoacyl-ACP synthase III (FabH), enoyl-ACP reductase enzyme isoforms (FabI, FabK, FabL and FabV), β-hydroxyacyl-ACP dehydratase (FabZ), long-chain-fatty-acid-CoA ligase (FadD), long-chain fatty acid transport protein (FadL), fatty acid kinase (FakA, FakB), acyl-ACP thioesterase (FatA), iron transport protein (FatB), thioesterase 1/protease 1/lysophospholipase L1 (TesA), phosphate acyltransferase (PlsX), CoenzymeA (CoA), acyl carrier protein (ACP). **c-d.** Abundance of selected KOs of interest (summed transcripts per million (TPM), **c**) and relative taxonomic assignment of contigs encoding for the selected KOs of interest at genus level (**d**) in fecal samples from KD- and SD-fed mice, and KC and SC recipient mice, at endpoint. In (**c**) and (**d**), the abundance of KOs highlighted in red correlates with colonic tumor burden in KD- and SD-fed mice. **e.** Levels of oleic (left) and palmitic (right) acid measured in bacterial cultures (in relation to sterile growth medium) of the Sonnenburg dataset. **f.** Relative abundance (total read counts) of selected families of interest detected in murine fecal samples from KD-, SD-fed, KC and SC recipient mice at endpoint**.** Members of the class Clostridia are listed in red, while members of the class Bacteroidia are listed in blue. **g-i.** HCT 116 (**g**), HT29 (**h**) and Caco2 (**i**) proliferation over time after treatment with stearic acid or control (100μM, 150 μM, 200 μM, 250 μM, 300 μM). Data shows mean of n=3 independent experiments (**g** and **h**), or n=2 independent experiments (**i**), each with eight technical replicates per condition. In (**c**) and (**f**), boxplots show medians with 1st and 3rd quantiles. The whiskers from the hinges to the smallest/largest values represent 1.5*inter-quartile range (IQR). Wilcoxon signed-rank test with Bonferroni correction in (**f**), ns p > 0.05, * p ≤ 0.05, ** p ≤

0.01, *** p ≤ 0.001, **** p ≤ 0.0001. Data shown in (**a**), (**c, red and black**), (**e, red**), (**d, first and third**

columns), (**f, red and black**) and (**i**) are from n=9 SPF SD-fed and n=8 SPF KD-fed mice, pooled from two

independent experiments. Data shown in (**c, gray and orange**), (**d, second and fourth columns**), and (**f,**

gray and orange) are from n=8 mice per condition.

Extended data figure 9. | a. Representative gating strategy of dendritic cells. **b.** Representative gating

strategy of macrophages. **c.** Representative gating strategy of regulatory T cells. **d.** Representative gating

strategy of CD8[+] T cells. **e.** Representative gating strategy of T helper 1 (subpanel 1) and T helper 2 cells

(subpanel 2).

Extended data figure 10. | a. Representative gating strategy of innate lymphoid cells (ILC) - ILC1 (subpanel

1), ILC2 (subpanel 2), ILC3 (subpanel 3). Transcription factors and cytokines are stained in two separate

batches. **b.** Representative gating strategy of natural killer (subpanel 1) and natural killer T (subpanel 2)

cells. **c.** Representative gating strategy of T helper 17 cells – transcription-factor-based phenotyping

(subpanel 1), cytokine-based phenotyping (subpanel 2) and the combined strategy (subpanel 3).

Acknowledgments

We acknowledge the contributions of the following people in the context of this project and thank

them for their work: Dominik Ternes, Rubens Begaj, Martin Nurmik and Cedric Gilson (MDM group,

DLSM) for assistance during organ processing at experimental endpoint; Ysaline Bossicard (MDM group,

DLSM) for tissue cryosectioning; Stephanie Willieme, Eric Grant and Mahesh Desai (LIH) for processing of

murine fecal samples; Xiangyi Dong, Floriane Gavotto and Christian Jäger (LCSB Metabolomics Platform)

for metabolite profile acquisition and raw data analysis; Aurélien Ginolhac (DLSM) for contributions to

data pre-processing; Pau Perez Escriva (DLSM) for bacterial culturing and critical reading of the

manuscript; Giulia Cesi (DLSM) for her contributions to obtaining funding; Djalil Coowar, Jennifer Behm,

Julie Virolle, Cloe Ordrenneau and all other members of the Animal Facility Team (LCSB); Sara Gomez and

Eliane Klein (DLSM) for maintenance of organoid cultures and organoid experimentation; Rashi Halder for

fecal sample 16s and WGS sequencing. The sequencing experiments presented in this paper were carried

out at the LCSB sequencing platform (RRID: SCR_021931) at the University of Luxembourg. We also thank

the patients who contributed their samples to the national colorectal cancer cohort from which we

obtained the samples used in this study, as well as Christelle Bahlawane (CIEC), Eve Herkenne (CIEC),

Laetitia Garcia (CIEC), Ingrid Vreja (CIEC), and nurses from CIEC for contributing to this cohort. We thank

Dr. Barbara Kofler for her feedback and discussions during the course of the project. The experiments

presented in this paper were carried out using the high-performance computing facilities at the University

of Luxembourg (https://hpc.uni.lu)[54]. BioRender (BioRender.com) was used for graphical illustration.

Funding

This work was supported by the Luxembourg National Research Fund (FNR) (grant nos.

CORE/C16/BM/11282028 (E.L.), PoC/18/12554295 (E.L.), PRIDE17/11823097 (M.T.), PRIDE/11823097

(L.d.N.), and CORE/15/BM/10404093 (P.W.)), by the Luxembourg National Research Fund and the

Fondation Cancer Luxembourg (grant no. CORE/C20/BM/14591557 (E.L.)) as well as by the Fondation du

Pélican de Mie and Pierre Hippert-Faber under the aegis of the Fondation de Luxembourg ('Pelican Grant';

M.T. and M.M.), a FNRS-Télévie grant to M.M., no. 7.4565.21-40007364), an Internal Research Project at

the University of Luxembourg (MiDiCa—integrated analysis of the effects of microbiome-diet interactions

on human colorectal adenocarcinoma enterocytes; E.L., P.W. and S.H.), the Fondation Cancer and the

Fondation Kriibskrank Kanner Luxembourg (V.I.P), the Action LIONS Vaincre le Cancer Luxembourg and a

European Research Council grant under the European Union's Horizon 2020 research and innovation

programme (grant agreement no. 863664 to P.W.). This project was also supported by the Doctoral School

in Science and Engineering (M.T., M.K., M.M. and L.d.N.) and the Department of Life Sciences and

Medicine at the University of Luxembourg. The funders had no role in study design, data collection and

analysis, decision to publish or preparation of the manuscript.

612   **References**

613   1.   Colorectal cancer statistics. *WCRF International* https://www.wcrf.org/cancer-trends/colorectal-

614   cancer-statistics/.

615   2.   Akimoto, N. *et al.* Rising incidence of early-onset colorectal cancer - a call to action. *Nat. Rev. Clin.*

616   *Oncol.* **18**, 230–243 (2021).

617   3.   Weber, D. D. *et al.* Ketogenic diet in the treatment of cancer – Where do we stand? *Mol. Metab.* **33**,

618   102–121 (2019).

619   4.   Dmitrieva-Posocco, O. *et al.* β-Hydroxybutyrate suppresses colorectal cancer. *Nature* **605**, 160–165

620   (2022).

621   5.   Ang, Q. Y. *et al.* Ketogenic Diets Alter the Gut Microbiome Resulting in Decreased Intestinal Th17

622   Cells. *Cell* **181**, 1263-1275.e16 (2020).

623   6.   Kong, C. *et al.* Ketogenic diet alleviates colitis by reduction of colonic group 3 innate lymphoid cells

624   through altering gut microbiome. *Signal Transduct. Target. Ther.* **6**, 154 (2021).

625   7.   Wei, R. *et al.* Ketogenesis Attenuates KLF5-Dependent Production of CXCL12 to Overcome the

626   Immunosuppressive Tumor Microenvironment in Colorectal Cancer. *Cancer Res.* **82**, 1575–1588

627   (2022).

628   8.   Ferrere, G. *et al.* Ketogenic diet and ketone bodies enhance the anticancer effects of PD-1 blockade.

629   *JCI Insight* **6**, 145207 (2021).

630   9.   Paoli, A. *et al.* Ketogenic Diet and Microbiota: Friends or Enemies? *Genes* **10**, 534 (2019).

631   10.   Cairo, C. & Webb, T. J. Effective Barriers: The Role of NKT Cells and Innate Lymphoid Cells in the Gut.

632   *J. Immunol. Baltim. Md 1950* **208**, 235–246 (2022).

633   11.   Jukic, A., Bakiri, L., Wagner, E. F., Tilg, H. & Adolph, T. E. Calprotectin: from biomarker to biological

634   function. *Gut* **70**, 1978–1988 (2021).

12. Wong, S. H. *et al.* Gavage of Fecal Samples From Patients With Colorectal Cancer Promotes

Intestinal Carcinogenesis in Germ-Free and Conventional Mice. *Gastroenterology* **153**, 1621-1633.e6

(2017).

13. Mu, C. *et al.* Probiotics counteract hepatic steatosis caused by ketogenic diet and upregulate AMPK

signaling in a model of infantile epilepsy. *EBioMedicine* **76**, 103838 (2022).

14. Nan, F. *et al.* Mannose: A Sweet Option in the Treatment of Cancer and Inflammation. *Front.*

Pharmacol. **13**, (2022).

15. Gonzalez, P. S. *et al.* Mannose impairs tumour growth and enhances chemotherapy. *Nature* **563**,

719–723 (2018).

16. Liu, Q., Li, X., Zhang, H. & Li, H. Mannose Attenuates Colitis-Associated Colorectal Tumorigenesis by

Targeting Tumor-Associated Macrophages. *J. Cancer Prev.* **27**, 31–41 (2022).

17. De Simone, V., Pallone, F., Monteleone, G. & Stolfi, C. Role of TH17 cytokines in the control of

colorectal cancer. *Oncoimmunology* **2**, e26617 (2013).

18. Ternes, D. *et al.* Microbiome in Colorectal Cancer: How to Get from Meta-omics to Mechanism?

Trends Microbiol. **28**, 401–423 (2020).

19. Fan, L. *et al.* A. Muciniphila Suppresses Colorectal Tumorigenesis by Inducing TLR2/NLRP3-Mediated

M1-Like TAMs. *Cancer Immunol. Res.* **9**, 1111–1124 (2021).

20. Wardman, J. F., Bains, R. K., Rahfeld, P. & Withers, S. G. Carbohydrate-active enzymes (CAZymes) in

the gut microbiome. *Nat. Rev. Microbiol.* **20**, 542–556 (2022).

21. Han, S. *et al.* A metabolomics pipeline for mechanistic interrogation of the gut microbiome. *Nature*

595, 415–420 (2021).

22. Mao, T. *et al.* Elevated serum β-hydroxybutyrate, a circulating ketone metabolite, accelerates

colorectal cancer proliferation and metastasis via ACAT1. *Oncogene* **42**, 1889–1899 (2023).

658 23. Gopalakrishnan, V. *et al.* Gut microbiome modulates response to anti-PD-1 immunotherapy in

659   melanoma patients. *Science* **359**, 97–103 (2018).

660 24. Matson, V. *et al.* The commensal microbiome is associated with anti-PD-1 efficacy in metastatic

661   melanoma patients. *Science* **359**, 104–108 (2018).

662 25. Routy, B. *et al.* Gut microbiome influences efficacy of PD-1-based immunotherapy against epithelial

663   tumors. *Science* **359**, 91–97 (2018).

664 26. Olson, C. A. *et al.* The Gut Microbiota Mediates the Anti-Seizure Effects of the Ketogenic Diet. *Cell*

665   **173**, 1728-1741.e13 (2018).

666 27. cis-Encoded Small RNAs, a Conserved Mechanism for Repression of Polysaccharide Utilization in

667   Bacteroides | Journal of Bacteriology. https://journals.asm.org/doi/10.1128/jb.00381-16.

668 28. Pudlo, N. A. *et al.* Symbiotic Human Gut Bacteria with Variable Metabolic Priorities for Host Mucosal

669   Glycans. *mBio* **6**, 10.1128/mbio.01282-15 (2015).

670 29. Milani, C. *et al.* Bifidobacteria exhibit social behavior through carbohydrate resource sharing in the

671   gut. *Sci. Rep.* **5**, 15782 (2015).

672 30. Luis, A. S. & Hansson, G. C. Intestinal mucus and their glycans: A habitat for thriving microbiota. *Cell*

673   *Host Microbe* **31**, 1087–1100 (2023).

674 31. Drula, E. *et al.* The carbohydrate-active enzyme database: functions and literature. *Nucleic Acids*

675   *Res.* **50**, D571–D577 (2022).

676 32. Desai, M. S. *et al.* A dietary fiber-deprived gut microbiota degrades the colonic mucus barrier and

677   enhances pathogen susceptibility. *Cell* **167**, 1339-1353.e21 (2016).

678 33. van der Lugt, B. *et al.* Akkermansia muciniphila ameliorates the age-related decline in colonic mucus

679   thickness and attenuates immune activation in accelerated aging Ercc1−/Δ7 mice. *Immun. Ageing*

680   **16**, 6 (2019).

34. Gamage, H. K. A. H. *et al.* Changes in dietary fiber intake in mice reveal associations between colonic

mucin O-glycosylation and specific gut bacteria. *Gut Microbes* **12**, 1802209 (2020).

35. Johansson, M. E. V. *et al.* Normalization of Host Intestinal Mucus Layers Requires Long-Term

Microbial Colonization. *Cell Host Microbe* **18**, 582–592 (2015).

36. Brown, E. M., Clardy, J. & Xavier, R. J. Gut microbiome lipid metabolism and its impact on host

physiology. *Cell Host Microbe* **31**, 173–186 (2023).

37. Yamada, S. *et al.* Gut microbiota-mediated generation of saturated fatty acids elicits inflammation in

the liver in murine high-fat diet-induced steatohepatitis. *BMC Gastroenterol.* **17**, 136 (2017).

38. Guillou, H., Zadravec, D., Martin, P. G. P. & Jacobsson, A. The key roles of elongases and desaturases

in mammalian fatty acid metabolism: Insights from transgenic mice. *Prog. Lipid Res.* **49**, 186–199

(2010).

39. Yao, J. & Rock, C. O. Exogenous fatty acid metabolism in bacteria. *Biochimie* **141**, 30–39 (2017).

40. Flowers, M. T. & Ntambi, J. M. Role of stearoyl-coenzyme A desaturase in regulating lipid

metabolism. *Curr. Opin. Lipidol.* **19**, 248–256 (2008).

41. Lien, E. C. *et al.* Low glycaemic diets alter lipid metabolism to influence tumour growth. *Nature* **599**,

302–307 (2021).

42. Zhou, X. *et al.* Stearoyl-CoA Desaturase-Mediated Monounsaturated Fatty Acid Availability Supports

Humoral Immunity. *Cell Rep.* **34**, (2021).

43. Grajchen, E. *et al.* Fatty acid desaturation by stearoyl-CoA desaturase-1 controls regulatory T cell

differentiation and autoimmunity. *Cell. Mol. Immunol.* **20**, 666–679 (2023).

44. Kindt, A. *et al.* The gut microbiota promotes hepatic fatty acid desaturation and elongation in mice.

Nat. Commun. **9**, 3760 (2018).

45. Kim, K. S. *et al.* Genome-wide multi-omics analysis reveals the nutrient-dependent metabolic

features of mucin-degrading gut bacteria. *Gut Microbes* **15**, 2221811 (2023).

46. McKain, N., Shingfield, K. J. & Wallace, R. J. Metabolism of conjugated linoleic acids and 18 : 1 fatty

acids by ruminal bacteria: products and mechanisms. *Microbiology* **156**, 579–588 (2010).

47. Microbial Production of Conjugated Linoleic Acid and Conjugated Linolenic Acid Relies on a

Multienzymatic System | Microbiology and Molecular Biology Reviews.

https://journals.asm.org/doi/10.1128/mmbr.00019-18.

48. Fermor, B. F. *et al.* Fatty acid composition of normal and malignant cells and cytotoxicity of stearic,

oleic and sterculic acids in vitro. *Eur. J. Cancer Oxf. Engl. 1990* **28A**, 1143–1147 (1992).

49. Habib, N. A. *et al.* Stearic acid and carcinogenesis. *Br. J. Cancer* **56**, 455–458 (1987).

50. Evans, L. M., Cowey, S. L., Siegal, G. P. & Hardy, R. W. Stearate preferentially induces apoptosis in

human breast cancer cells. *Nutr. Cancer* **61**, 746–753 (2009).

51. Mitchel, J. *et al.* Computational Identification of Stearic Acid as a Potential PDK1 Inhibitor and In

Vitro Validation of Stearic Acid as Colon Cancer Therapeutic in Combination with 5-Fluorouracil.

Cancer Inform. **20**, 11769351211065979 (2021).

52. Zhang, Y. *et al.* Altered gut microbiome composition in children with refractory epilepsy after

ketogenic diet. *Epilepsy Res.* **145**, 163–168 (2018).

53. Naqash, A. R. *et al.* The role of gut microbiome in modulating response to immune checkpoint

inhibitor therapy in cancer. *Ann. Transl. Med.* **9**, 1034 (2021).

54. Varrette, S., Bouvry, P., Cartiaux, H. & Georgatos, F. Management of an academic HPC cluster: The

UL experience. in *2014 International Conference on High Performance Computing & Simulation*

(HPCS) 959–967 (IEEE, 2014). doi:10.1109/HPCSim.2014.6903792.

725