Adaptive Regularized Tri-Factor Non-Negative Matrix Factorization for Cell Type Deconvolution

Tianyi Liu[1]   Chuwen Liu[1]   Quefeng Li[1,*] Xiaojing Zheng[1,2,*]   Fei Zou[1,3,*]

[1]Department of Biostatistics, The University of North Carolina at Chapel Hill

[2]Department of Pediatrics, The University of North Carolina at Chapel Hill

[3]Department of Genetics, The University of North Carolina at Chapel Hill

Abstract

Motivation: Accurate deconvolution of cell types from bulk gene expression is crucial for understanding cellular compositions and uncovering cell-type specific differential expression and physiological states of diseased tissues. Existing deconvolution methods have limitations, such as requiring complete cellular gene expression signatures or neglecting partial biological information. Moreover, these methods often overlook varying cell-type mRNA amounts, leading to biased proportion estimates. Additionally, they do not effectively utilize valuable reference information from external studies, such as means and ranges of population cell-type proportions.

Results: To address these challenges, we introduce an Adaptive Regularized Tri-factor non-negative matrix factorization approach for deconvolution (ARTdeConv). We rigorously establish the numerical convergence of our algorithm. Through benchmark simulations, we demonstrate the superior performance of ARTdeConv compared to state-of-the-art reference-free methods. In a real-world application, our method accurately estimates cell proportions, as evidenced by the nearly perfect Pearson's correlation between ARTdeConv estimates and flow cytometry measurements in a dataset from a trivalent influenza vaccine study. Moreover, our analysis of ARTdeConv estimates in COVID-19 patients reveals patterns consistent with important immunological phenomena observed in other studies.

Availability and implementation: The proposed method, ARTdeConv, is implemented as an R package and can be accessed on GitHub for researchers and practitioners at https://github.com/gr8lawrence/ARTDeConv.

Keywords: Cell-type deconvolution, Convergence analysis, Multiplicative update algorithm, Non-negative matrix factorization, RNA sequencing, Single cell data

---

*Co-corresponding authors

1. Introduction

Heterogeneity in cell-type proportions exists across biological samples. Neglecting this heterogeneity in bulk gene expression can introduce bias into subsequent analyses such as differential expression analyses. Conversely, acknowledging and accounting for this heterogeneity has shown clear advantages, yielding more accurate survival time predictions and tumor type classifications (Elloumi et al., 2011; Avila Cobos et al., 2020).

While laboratory techniques like flow cytometry or immunohistochemistry are available for physically sorting cells into cell types and quantifying their abundances, they are often limited by the availability of cell samples, the particularity of antibodies to separate cells, as well as labor and time investments (Newman et al., 2015; Spitzer and Nolan, 2016). Cell type deconvolution (referred to as "deconvolution" in this article), a computational process aimed at digitally separating heterogeneous mixture signals into their constituent components, has been critical in expediting the estimation of cell-type proportions from bulk gene expression data, such as RNA sequencing (RNA-seq). Recent years have seen the emergence of several deconvolution methods extensive applications in the field of computational biology. These methods can generally be categorized into two groups: reference-based and reference-free, depending on whether they require individual cell-type gene expression signatures, referred to as a signature matrix, as prior knowledge (Li and Wu, 2019; Avila Cobos et al., 2020; Jin and Liu, 2021).

Gene signatures can be derived from either single-cell RNA-seq (scRNA-seq) data or sorted bulk RNA profiles of individual cell types and used as reference (Chen et al., 2018). Although gene signatures have been successfully established for some cell types, acquiring those for other cell types might be labor-intensive or even unfeasible (Morris and Kopetz, 2016; Wigerblad et al., 2022). Neglecting to incorporate gene signatures for a prevalent cell type can induce biases in the proportional estimates of other cell types within reference-based deconvolution (Avila Cobos et al., 2020). On the other hand, reference-free techniques enable the unsupervised estimation of cell-type proportions, but at the cost of disregarding information embedded in known gene signatures (Zaitsev et al., 2019; Chen et al., 2020). A deconvolution method that can utilize partial reference gene signatures presents a plausible compromise.

Moreover, external information on the population cell-type proportions may also help with estimating cell-type proportions. For example, in blood samples, the proportion means (or medians) and ranges of major cell types can are available from complete blood count (CBC) tests. However, deconvolution techniques that effectively incorporate such information are currently lacking in the field.

During the pre-processing of data, cell-type mRNA amounts, denoting the quantities of RNA molecules contained within an individual cell, are often lost after library normalization. Zaitsev et al. (2019) found that failures to account for mRNA amounts produced biased cell-type proportion estimates. However, many deconvolution methods assume a generative model where the loss of this information could not be accounted for (Avila Cobos et al., 2020). Therefore, incorporating these quantities into the generative model can be pivotal to deconvolution accuracy.

In this article, we propose an *A*daptive *R*egularized *T*ri-factor non-negative matrix factorization method for *deConv*olution (ARTdeConv). It addresses the three outstanding issues discussed above: utilizing partial reference gene signatures, incorporating external information on cell-type proportions, and accounting for cell-type mRNA amounts. Our contribution includes deriving and implementing a multiplicative update (MU) algorithm

Figure 1: A schema of the ARTdeConv workflow.

for solving ARTdeConv, proving the algorithm's convergence, and demonstrating its merits by simulations and real data analysis. A schematic representation of ARTdeConv's workflow is shown in Figure 1. An R package implementing ARTdeConv is also available on GitHub.

2. Method

2.1. Notation

Let $\mathbb{R}_+$ and $\mathbb{R}_{++}$ denote the set of non-negative and positive real numbers respectively. $m, n, K \in \mathbb{R}_{++}$ are used to denote the number of genes, samples, and cell types with $K \leq \min(m, n)$ in deconvolution. Let $K_0 < K$ be the number of cell types for which we have reference gene expression available. Let $\boldsymbol{Y} \in \mathbb{R}_+^{m \times n}$ be the matrix of bulk gene expression and $\boldsymbol{\Theta} \in \mathbb{R}_+^{m \times K}$ be the full gene signature matrix. Let $\boldsymbol{s} \in \mathbb{R}_{++}^K$ be a vector of relative cell-type mRNA amounts, and $\boldsymbol{P} \in \mathbb{R}_+^{K \times n}$ the proportion matrix. Let $\boldsymbol{\epsilon} \in \mathbb{R}^{m \times n}$ be the random error matrix. Unless otherwise specified, we denote $\boldsymbol{\theta}_k$ as the k-th column of $\boldsymbol{\Theta}$. We use $\boldsymbol{A}^\top$ to denote the transpose of $\boldsymbol{A}$. For any $m \times n$ matrix $\boldsymbol{A}$, its Frobenius norm is denoted as $\|\boldsymbol{A}\|_F = \sqrt{\operatorname{tr}(\boldsymbol{A}^\top \boldsymbol{A})} = \sqrt{\sum_{i=1}^m \sum_{j=1}^n A_{ij}^2}$. For two matrices $\boldsymbol{A}, \boldsymbol{B}$ of the same dimensions, $\boldsymbol{A} \odot \boldsymbol{B}$ denotes the element-wise product of $\boldsymbol{A}$ and $\boldsymbol{B}$ and $\frac{\boldsymbol{A}}{\boldsymbol{B}}$ denotes their element-wise quotient.

2.2. Model

We first propose the following tri-factor generative model

$$\boldsymbol{Y} = \boldsymbol{\Theta}\text{diag}(\boldsymbol{s})\boldsymbol{P} + \boldsymbol{\epsilon}, \tag{1}$$

which extends the canonical model $\boldsymbol{Y} = \boldsymbol{\Theta}\boldsymbol{P} + \boldsymbol{\epsilon}$ (Avila Cobos et al., 2020). The difference between them is $\text{diag}(\boldsymbol{s})$, a diagonal matrix of cell-type mRNA amounts. In practice, normalization, for example by library sizes, is frequently employed to alleviate between-sample technical artifacts in sequencing data (Hafemeister and Satija, 2019; Lytal et al., 2020). During this process, information regarding the quantities of RNA molecules packed within different kinds of cells sometimes gets lost. Failures to recover these quantities lead to documented biases in estimating cell-type proportions (Zaitsev et al., 2019). Should this occurs, $\text{diag}(\boldsymbol{s})$ can account for the RNA molecule quantities. Otherwise, $\text{diag}(\boldsymbol{s})$ would be close to the identity matrix $\boldsymbol{I}_K$. We notice that (1) is similar to the model of MuSiC, a fully reference-based method by Wang et al. (2019), where each sample is accorded an additional scaling factor c_i to account for between-sample measurement variation in bulk gene expression.

For performing deconvolution based on (1), ARTdeConv assumes prior knowledge of a partial signature matrix $\boldsymbol{\Theta}_{K_0} \in \mathbb{R}_+^{m \times K_0}$ as well as $\boldsymbol{Y}$. Then, an objective function is defined:

$$f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P}) = \frac{1}{2mn}\|\boldsymbol{Y} - \boldsymbol{\Theta}\text{diag}(\boldsymbol{s})\boldsymbol{P}\|_F^2 + \frac{\alpha_1}{2}R_1 + \frac{\alpha_2}{2}R_2 + \frac{\beta}{2}R_3, \tag{2}$$

where R_1, R_2, and R_3 are regularizers to be explained later and α_1, α_2, and β are their tuning parameters. We discuss the selection of optimal tuning parameters in Section 2.4. $\boldsymbol{\Theta}$, $\boldsymbol{s}$, and $\boldsymbol{P}$ are then estimated via

$$(\hat{\boldsymbol{\Theta}}, \hat{\boldsymbol{s}}, \hat{\boldsymbol{P}}) = \underset{\boldsymbol{\Theta}\in\mathbb{R}_+^{m\times K}, \ \boldsymbol{s}\in\mathbb{R}_{++}^K, \ \boldsymbol{P}\in\mathbb{R}_+^{K\times n}}{\text{argmin}} f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P}). \tag{3}$$

The main objective here is to obtain an estimate of the proportion matrix $\hat{\boldsymbol{P}}$. To follow a common practice of deconvolution and simplify the algorithm (Mohammadi et al., 2017), ARTdeConv does not directly constrain each column of $\boldsymbol{P}$ to sum to one during the estimation process. Instead, it obtains an unconstrained estimate $\hat{\boldsymbol{P}}$ and then re-normalizes its columns to have the unit sum.

Let $\boldsymbol{\Theta}_0 = \begin{bmatrix} \boldsymbol{\Theta}_{K_0} & \boldsymbol{0} & \cdots & \boldsymbol{0} \end{bmatrix} \in \mathbb{R}_+^{m\times K}$ and $\boldsymbol{\Delta}$ be an $m \times K$ matrix such that $\Delta_{jk} = I(k \leq K_0)$ for $1 \leq j \leq m$. The squared Frobenius distance between the estimated $\boldsymbol{\Theta}$ and $\boldsymbol{\Theta}_0$ is penalized through $R_1 = \|\boldsymbol{\Delta} \odot (\boldsymbol{\Theta} - \boldsymbol{\Theta}_0)\|_F^2$.

Though we present a special case where $\boldsymbol{\Theta}_{K_0}$ occupies the first K_0 columns of $\boldsymbol{\Theta}$, by re-defining $\boldsymbol{\Delta}$ correspondingly, $\boldsymbol{\Theta}_{K_0}$ can occupy any K_0 columns, covering all structures of prior knowledge on the signature matrix.

On the other hand, R_1 is not a strictly convex function of $\boldsymbol{\Theta}$, making the guarantee of the numerical convergence of ARTdeConv's algorithm difficult (more details on this later). To atone for this, we make $R_2 = \|(\boldsymbol{J}_{m\times K} - \boldsymbol{\Delta}) \odot \boldsymbol{\Theta}\|_F^2$, where $\boldsymbol{J}_{m\times K}$ is an $m \times K$ matrix of 1s. It can be shown that any positive linear combination of R_1 and R_2 is strictly convex with respect to $\boldsymbol{\Theta}$.

Information on the cell-type proportions such as means (or medians) and ranges in a population frequently exists in external data and is accessible online, for example through complete blood counts of leukocytes (Takami et al., 2021). Let m_k denote the reference mean (or median) of cell type k's population proportion and let r_k denote its range.

To incorporate it into our deconvolution method, we propose adding another penalty $R_3 = \sum_{k=1}^{K}(1/r_k)\|\boldsymbol{p}_k^\top - m_k\mathbf{1}_n\|_2^2$, where $\boldsymbol{p}_k^\top$ is the k-th row of $\boldsymbol{P}$. In R_3, m_k acts as a pivot, and the deviation from which is penalized for the estimated $\boldsymbol{p}_k^\top$. Meanwhile, $1/r_k$ acts as a weight so the proportion estimates of cell types with wider ranges are less penalized for departing from their respective m_k. Letting $\boldsymbol{M}$ be a $K \times n$ matrix such that $\boldsymbol{M} = \begin{bmatrix} m_1\mathbf{1}_n & m_2\mathbf{1}_n & \cdots & m_K\mathbf{1}_n \end{bmatrix}^\top$ and $\boldsymbol{\rho} = \mathrm{diag}(r_1, r_2, \cdots, r_K)$, we can write R_3 in a matrix form $R_3 = \|\boldsymbol{\rho}^{-1/2}(\boldsymbol{P} - \boldsymbol{M})\|_F^2$.

2.3. Multiplicative Update Algorithm

A multiplicative update (MU) algorithm is proposed to solve (3). The MU algorithm was originally designed to solve the canonical bi-factor non-negative matrix factorization (NMF) problem in Lee and Seung (2000) and was extensively studied by Lin (2007). Technically, it is similar to the majorization-minimization algorithm of Lange (2016). It can be readily extended to solving a multi-factor regularized NMF problem like the one for ARTdeConv.

Let t be a non-negative integer denoting the current number of iteration. The multiplicative update steps are derived by finding the gradients of a set of auxiliary functions for each row of $\boldsymbol{\Theta}^t$, each column of $\boldsymbol{P}^t$, and $\boldsymbol{s}^t$. These update steps are:

$$\boldsymbol{\Theta}^{t+1} = \boldsymbol{\Theta}^t \odot \frac{\boldsymbol{Y}(\boldsymbol{P}^t)^\top \mathrm{diag}(\boldsymbol{s}^t) + mn\alpha_1\boldsymbol{\Delta} \odot \boldsymbol{\Theta}_0}{\boldsymbol{\Theta}^t\mathrm{diag}(\boldsymbol{s}^t)\boldsymbol{P}^t(\boldsymbol{P}^t)^\top\mathrm{diag}(\boldsymbol{s}^t) + mn\left\{\alpha_1\boldsymbol{\Delta} \odot \boldsymbol{\Theta}^t + \alpha_2(\boldsymbol{J}_{m\times K} - \boldsymbol{\Delta}) \odot \boldsymbol{\Theta}^t\right\}};$$
$$(4)$$

$$\boldsymbol{P}^{t+1} = \boldsymbol{P}^t \odot \frac{\mathrm{diag}(\boldsymbol{s}^t)(\boldsymbol{\Theta}^{t+1})^\top\boldsymbol{Y} + mn\beta\boldsymbol{\rho}^{-1}\boldsymbol{M}}{\mathrm{diag}(\boldsymbol{s}^t)(\boldsymbol{\Theta}^{t+1})^\top\boldsymbol{\Theta}^{t+1}\mathrm{diag}(\boldsymbol{s}^t)\boldsymbol{P}^t + mn\beta\boldsymbol{\rho}^{-1}\boldsymbol{P}^t};$$
$$(5)$$

$$\boldsymbol{s}^{t+1} = \boldsymbol{s}^t \odot \frac{\boldsymbol{u}^t}{\boldsymbol{Z}^t\boldsymbol{s}^t},$$
$$(6)$$

where $u_k^t = \mathrm{tr}\left\{\boldsymbol{Y}^\top\boldsymbol{\theta}_k^{t+1}(\boldsymbol{p}^{t+1})_k^\top\right\}$ and $Z_{kl}^t = \left\{(\boldsymbol{\theta}^{t+1})_k^\top\boldsymbol{\theta}_l^{t+1}\right\}\left\{(\boldsymbol{p}^{t+1})_k^\top\boldsymbol{p}_l^{t+1}\right\}$ for $k, l = 1, 2, \cdots, K$.

Mathematical details for finding the auxiliary functions and the update steps can be found in Supplementary Material Section B. The pseudo-code of the MU algorithm is shown in Algorithm 1.

Algorithm 1 The MU Algorithm For Solving ARTdeConv

    Initialize $\boldsymbol{\Theta}^0 \in \mathbb{R}_+^{m\times K}$ such that $R_1 = 0$, $\boldsymbol{P}^0 \in \mathbb{R}_+^{K\times n}$, $\boldsymbol{s}^0 \in \mathbb{R}_{++}^K$. Fix α_1, α_2, and $\beta > 0$.

  Repeat
    Update $\boldsymbol{\Theta}^t$ to $\boldsymbol{\Theta}^{t+1}$ via (4);
    Update $\boldsymbol{P}^t$ to $\boldsymbol{P}^{t+1}$ via (5);
    Update $\boldsymbol{s}^t$ to $\boldsymbol{s}^{t+1}$ via (6);
  Until
$$\frac{|f(\boldsymbol{\Theta}^{t+1}, \boldsymbol{s}^{t+1}, \boldsymbol{P}^{t+1}) - f(\boldsymbol{\Theta}^t, \boldsymbol{s}^t, \boldsymbol{P}^t)|}{f(\boldsymbol{\Theta}^t, \boldsymbol{s}^t, \boldsymbol{P}^t)} < \delta,$$

where δ is a small tolerance parameter and f defined as in (2). Define the final outputs as $(\hat{\boldsymbol{\Theta}}, \hat{\boldsymbol{s}}, \hat{\boldsymbol{P}})$. Re-normalize $\hat{\boldsymbol{P}}$ by dividing each column of by its sum.

The MU algorithm has two advantages (under the assumptions discussed in Supplementary Material Section A). First, $\boldsymbol{\Theta}^t$, $\boldsymbol{P}^t$, and $\boldsymbol{s}^t$ remain non-negative throughout all iterations if their initial values are non-negative. It is recommended that $\boldsymbol{\Theta}^0$ is set to satisfy $R_1 = 0$, so any zero in the known partial signature matrix will remain zero. Second, the MU algorithm can be shown to achieve numerical convergence. If we enforce $\mathrm{diag}(\boldsymbol{s}^t) = \boldsymbol{I}_K$ and set $\alpha_1 = \alpha_2 = \beta = 0$, our MU algorithm is reduced to that of Lee and Seung (2000) for the canonical NMF. Due to the non-linearity of the objective function (2), it is not guaranteed that a global minimizer can be produced. Following the precedence of other NMF-based deconvolution software, we recommend running the MU algorithm with multiple restarts and choosing the one with the smallest Frobenius norm of the residuals on the estimated bulk expression (Gaujoux and Seoighe, 2010, 2013). We set 10 restarts as the default in our implementation.

We now describe the default initialization provided in the implementation of ARTdeConv. First, the first K_0 columns of $\boldsymbol{\Theta}^0$ were used as $\boldsymbol{\Theta}^*_{K_0}$ to satisfy $R_1 = 0$. Each initial value in the rest of the columns θ_{jl}^0 is generated independently by $\theta_{jl}^0 \sim |N(\bar{y}_j, \hat{\sigma}_j^2)|$, where $\bar{y}_j$ and $\hat{\sigma}_j^2$ are the mean and variance of the bulk expression values for gene j across all n samples. Then, for each cell type k of sample i, p_{ki}^0 is generated from a $N(m_k, 0.1)$ distribution. After this, negative entries in $\boldsymbol{P}^0$ are corrected to 0.01. Lastly, $\boldsymbol{s}^0$ is initialized by solving

$$\boldsymbol{s}^0 = \operatorname*{argmin}_{\boldsymbol{s} \in \mathbb{R}^K} \frac{1}{2} \|\boldsymbol{Y}^* - \boldsymbol{\Theta}^0 \mathrm{diag}(\boldsymbol{s}) \boldsymbol{P}^0\|_F^2.$$

Negative values in $\boldsymbol{s}^0$ will be replaced by the mean of positive values.

2.4. Tuning Parameter Selection

A grid search using a B-fold cross-validation (CV) is designed for selecting tuning parameters. Since R_2 is designed to make the objective function in (2) strictly convex, α_2 is advised to be fixed at a minuscule value. In the implementation of ARTdeConv, $\alpha_2 = 10^{-12}$ is the default value.

To begin, $\boldsymbol{Y}$ is randomly divided into B different folds, each containing $1/B$ columns of $\boldsymbol{Y}$. Two grids $\mathfrak{A}_1, \mathfrak{B} \subset \mathbb{R}_{++}$ containing respective candidates values for α_1 and β are declared. Next, for one $\alpha_1 \in \mathfrak{A}_1$, one $\beta \in \mathfrak{B}$ and a fold $b \in \{1, 2, \ldots, B\}$, the columns in the fold are held out as the test set $\boldsymbol{Y}_{\text{test}}^{(b)}$. The rest of the columns in $\boldsymbol{Y}$ are used as the training set $\boldsymbol{Y}_{\text{train}}^{(b)}$, upon which an ARTdeConv solution $(\tilde{\boldsymbol{\Theta}}^{(b)}, \tilde{\boldsymbol{s}}^{(b)}, \tilde{\boldsymbol{P}}^{(b)})$ is obtained.

Then, since it is assumed that n samples should share the same $\boldsymbol{\Theta}$ and $\boldsymbol{s}$ in (1), the estimated proportions $\hat{\boldsymbol{P}}^{(b)}$ on the test set $\boldsymbol{Y}_{\text{test}}^{(b)}$ is obtained by

$$\hat{\boldsymbol{P}}^{(b)} = \operatorname*{argmin}_{\boldsymbol{P} \in \mathbb{R}_+^{K \times n/B}} \|\boldsymbol{Y}_{\text{test}}^{(b)} - \tilde{\boldsymbol{\Theta}}^{(b)} \mathrm{diag}(\tilde{\boldsymbol{s}})^{(b)} \boldsymbol{P}\|_F^2.$$

Computationally, this is accomplished by solving a non-negative least square problem using the R package `nnls`. The CV error given α_1 and β is

$$\mathrm{Err}(\alpha_1, \beta) = \frac{1}{B} \sum_{b=1}^{B} \|\boldsymbol{Y}_{\text{test}}^{(b)} - \tilde{\boldsymbol{\Theta}}^{(b)} \mathrm{diag}(\tilde{\boldsymbol{s}})^{(b)} \hat{\boldsymbol{P}}^{(b)}\|_F^2.$$

Finally, we select the best tuning parameters α_1^* and β^* that minimize $\mathrm{Err}(\alpha_1, \beta)$.

2.5. Convergence Analysis

In this section, we prove the numerical convergence of Algorithm 1. The main result depends on two reasonable assumptions in Supplementary Material Section A. Consequently, the following theorem holds for Algorithm 1 for solving the problem in (3).

Theorem 1 (Convergence of Algorithm 1 to a Stationary Point). *Under the technical assumptions listed in Supplementary Material Section A, the sequence $\{(\boldsymbol{\Theta}^t, \boldsymbol{s}^t, \boldsymbol{P}^t)\}_{t=1}^{\infty}$ in Algorithm 1 converges to a stationary point of $f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P})$.*

The proof of Theorem 1 relies on showing that it is a special case of the block successive upper-bound minimization (BSUM) algorithm given in Razaviyayn et al. (2013). Thus, the convergence is guaranteed by related BSUM theories. Doing so involved proving that the sub-level set $\mathcal{X}^0 = \{(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P}) : f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P}) \leq f(\boldsymbol{\Theta}^0, \boldsymbol{s}^0, \boldsymbol{P}^0)\}$ is compact and the objective function in (3) is coercive, as well as demonstrating the majorization-minimization (MM) properties and strong convexity of the auxiliary functions used to derive the update steps in Algorithm 1. Details on the definitions of coercive functions and MM properties, as well as the proof of Theorem 1 are relegated to Supplementary Material Section C.

3. Results

3.1. Performance in Deconvolving Pseudo-Bulk Samples

To assess the deconvolution performance of ARTdeConv in comparison to alternative methods, we conducted a benchmarking simulation study by evaluating it against three distinct reference-free methods: NMF (Gaujoux and Seoighe, 2010), debCAM (Chen et al., 2020), and LINSEED (Zaitsev et al., 2019).

We focused on reference-free methods because reference-based methods are inapplicable under the assumption of partial knowledge on the signature matrix. In addition, we also included results of randomly guessing the proportions as a bottom line for deconvolution performance: each proportion was independently guessed from a $U(0, 1)$ distribution and the resulting proportion matrix was then re-normalized to make each column have the unit sum.

We generated a set of pseudo-bulk expression matrices using methodologies similar to those outlined in other previous studies (Avila Cobos et al., 2020) by the following formula

$$\boldsymbol{Y}^* = \boldsymbol{\Theta}^* \mathrm{diag}(\boldsymbol{s}^*)\boldsymbol{P}^* + \boldsymbol{\epsilon}. \tag{7}$$

In (7), $K = 5$ hypothetical cell types, CT1 - CT5, were constructed with known partial signatures on $K_0 = 4$ of them (CT1 - CT4).

In total, $n = 100$ samples were simulated on $\boldsymbol{P}^*$. To investigate the effects of the true relative abundance of CT5 in bulk tissue samples on deconvolution results, three classes of $\boldsymbol{P}^*$ were generated, representing when CT5 was rare, uniform, and extra compared to other cell types in the tissue. Reference medians $(m_1, \ldots, m_5)$ and ranges $(r_1, \ldots, r_5)$ were obtained directly from each row of $\boldsymbol{P}^*$.

The expression values in the true signature matrix $\boldsymbol{\Theta}^*$ were simulated row-by-row (i.e., gene-by-gene), controlled by a parameter γ, which dictated how cell-type-specific each gene was on average. Each simulated gene was more likely to be cell-type-specific when $\gamma = 1$ compared to when $\gamma = 0$. The expression of $M = 2000$ genes was first created

in a matrix $\boldsymbol{\Theta}^*_{\text{full}}$. Then, from these genes, a subset of $m = 500$ highly cell-type-specific genes called marker genes were selected for ARTdeConv and NMF. The method for the selection is described in details in Supplementary Material Section D.2. The expression of those selected genes was stored in $\boldsymbol{\Theta}^*$. The matrix $\boldsymbol{\Theta}^*_{K_0}$ consisting of the first four columns of $\boldsymbol{\Theta}^*$ was used as the partial signature matrix for ARTdeConv. On the other hand, debCAM and LINSEED searched for marker genes differently from $\boldsymbol{\Theta}^*_{\text{full}}$ by looking for simplicial vertices of the vector space spanned by normalized bulk gene expression (Zaitsev et al., 2019; Chen et al., 2020).

The matrix of errors $\boldsymbol{\epsilon}$ was first generated for all of the $M = 2000$ simulated genes. Beginning with $(\boldsymbol{\Theta}^*_{\text{full}}, \boldsymbol{s}^*, \boldsymbol{P}^*)$, the error-free full bulk expression matrix was calculated as $\boldsymbol{\Theta}^*_{\text{full}}\text{diag}(\boldsymbol{s}^*)\boldsymbol{P}^*$. The relative strength of the errors (noises) to the mean expression of genes (signals) was controlled by another parameter σ. Two levels of noises with $\sigma = 0.1$ or 10 (low vs. high) were introduced to evaluate the robustness of the methods to added noises. The final bulk expression matrix of all simulated genes was calculated using $\boldsymbol{Y}^*_{\text{full}} = \boldsymbol{\Theta}^*_{\text{full}}\text{diag}(\boldsymbol{s}^*)\boldsymbol{P}^* + \boldsymbol{\epsilon}$. From this, a sub-matrix $\boldsymbol{Y}^*$ corresponding to the genes in $\boldsymbol{\Theta}^*$ was selected. A detailed description of the probability distributions and their parameters in the generation of $\boldsymbol{\Theta}^*$, $\boldsymbol{s}^*$, $\boldsymbol{P}^*$, and $\boldsymbol{\epsilon}$ can be found in Supplementary Material Section D.1.

Having generated all necessary parts, 100 simulations were conducted for each combination of γ, σ, and CT-5 abundance class, where the deconvolution performance of benchmarked methods was reported. The tuning grid and algorithm parameters for ARTdeConv are described in Supplementary Material Section D.3. Due to a lack of identifiability, estimated proportions from NMF, debCAM, and LINSEED needed to be manually matched to different cell types, whose details were relegated to Supplementary Material Section D.4.

Performance metrics for evaluating deconvolution results included the following: (a) $\frac{1}{n}cor(\boldsymbol{p}^*_i, \hat{\boldsymbol{p}}_i)$, (b) $\frac{1}{5n}\sum_{i=1}^{n}\sum_{k=1}^{5}|p^*_{ki} - \hat{p}_{ki}|$, (c) $cor(\boldsymbol{p}^{*T}_5, \hat{\boldsymbol{p}}^{\top}_5)$, (d) $\frac{1}{n}\sum_{i=1}^{n}|p^*_{5i} - \hat{p}_{5i}|$, where cor denotes the Pearson's correlation between two vectors. (a) and (b) delineated the overall deconvolution accuracy, while (c) and (d) described the deconvolution accuracy for CT5, the missing cell type.

Overall, ARTdeConv had better performance than other benchmarked reference-free methods in recovering the correct proportions regardless of the level of cell-type specificity of signature genes, the level of additive noises, and the relative abundance of the missing cell type to other cell types (Figures 2-3). As expected, debCAM performed better when the signature genes were highly cell-type specific ($\gamma = 1$) while NMF did so when the true cell-type proportions were uniform. LINSEED, known to be insensitive to cell-type mRNA amounts, consistently had the worst performance of all benchmarked methods except in a few cases and were at times worse than random guesses. ARTdeConv also produced better estimation of CT5's proportions on average compared to the other methods, even when it was relatively rare in the bulk samples compared to other cell types (Figures 4-5).

We further remark that, in the simulations described above, $\boldsymbol{\Theta}^*$ included knowledge of marker genes for CT5. In practice, obtaining the marker genes of cell types without reference expression is, while not as straightforward, achievable through methods such as in Wang et al. (2016) and Qiu et al. (2021). Furthermore, deconvolution is completely feasible even without well defined marker genes for those cell types. See for instance the analyses in Sections 3.2 and 3.3.

Figure 2: Mean sample-wise Pearson correlation between true proportions and estimated cell-type proportions through five benchmarked methods.

Figure 3: Mean absolute deviation (MAD) between true proportions and estimated cell-type proportions through five benchmarked methods.

Figure 4: Pearson correlation between true proportions and estimated proportions of CT5 through five benchmarked methods.

Figure 5: Mean absolute deviation (MAD) between true proportions and estimated proportions of CT5 through five benchmarked methods.

3.2. ARTdeConv Accurately Estimates Cell Type Proportions of PBMC from Bulk Gene Expression in a Human Influenza Vaccine Study

To exemplify the application of ARTdeConv to real-world data, we utilized a dataset sourced from Hoek et al. (2015). This study examined alterations in gene expression within peripheral blood mononuclear cells (PBMCs) and sorted cell lines by magnetic-sorting (MACS). Blood samples were obtained from healthy volunteer subjects who were administered a trivalent inactivated influenza vaccine (TIV). Both bulk PBMCs and sorted PBMC cell lines from two enrolled subjects ("HD30" and "HD31") vaccinated with a single dose of 2011–2012 seasonal TIV were collected at four different time points: before vaccination (Day 0), and at Day 1, 3, and 7 post-vaccination. These samples were obtained and processed in fresh conditions for RNA-seq analysis.

Deconvolution was performed on eight bulk PBMC samples whose gene expression were measured in transcript per million (TPM). Gene expression in TPM from the sorted PBMC cell lines of the two subjects on Day 0 were used to construct the partial signature matrix. The study characterized four distinct PBMC cell types: T cells, B cells, natural killer (NK) cells, and monocytes. Additionally, there were other unspecified cell types collectively termed as "others". Detailed data processing steps could be referred to at Supplementary Material Section E.1.

The means and ranges of the cell-types in question were obtained through the reference values in Kleiveland (2015). For each cell type, the proportion means were calculated by taking the average of the upper and lower bounds of the reference values, while the ranges were calculated by taking the difference between the two bounds. For the "others" cell type, the mean was calculated by subtracting the means of all other cell types from one, and the range the difference between the upper and lower possible proportions of the other cell types. The exact values of the means and ranges are reported in Supplementary Material Table 1. The vector of ranges was further normalized such that it has unit L_2 norm before deconvolution.

The PBMC percentages on Day 0 were measured by flow cytometry and reported by Racle et al. (2017), which were used as the ground truths for evaluating the performance of ARTdeConv. Indeed, ARTdeConv achieved a notable degree of precision in estimating cell-type proportions on Day 0 as compared to the flow cytometry measurements (Figure 6). Namely, the estimated proportions for HD30 displayed a Pearson correlation of 0.992 with and a mean absolute deviation (MAD) of 0.027 from the ground truths. Similarly, a Pearson correlation of 0.977 was observed for HD31, along with a MAD of 0.049.

We also estimated proportions of PBMC cell types across all time points utilizing the same partial signature matrix (Figure 7). On HD30, a decline of T cell abundances before Day 1 and an increase between Days 1 and 3 were observed, followed by a slight decline until Day 7. An increase of monocyte abundances before and a decrease after Day 1 were also seen. Both trends are consistent with the profile of a virus shedder of the H1N1 virus in Rahil et al. (2020) with a small time shift. The time shift could be attributed to differences in viral strains and strengths between Rahil et al. (2020) and Hoek et al. (2015), as the former is a study on real H1N1 patients. The lack of notable changes in the cell-type proportions of HD31 resembles more to the profile of a non-virus shedder. More information on the viral progression of these two subjects is needed to confirm these observations.

We wish to point out that normalizing gene expression by TPM is known for losing

Figure 6: Estimated PBMC cell-type proportions from deconvolution using ARTdeConv versus flow cytometry measurements for the two subjects on Day 0.

Figure 7: Estimated proportions of PBMC cell types for the two subjects across all time points from deconvolution using ARTdeConv.

the information on cell-type mRNA amounts, which was corrected by ARTdeConv via diag($\hat{s}$). When re-running ARTdeConv on the same samples with diag(s) forced to be the identity matrix, we observed less accurate estimates of proportions (Supplementary Material Figure 1), corroborating the need for the correction of mRNA amounts in this scenario. Additional details regarding the tuning grid and algorithm parameter setup of this analysis can also be found at Supplememtary Material Section E.2.

3.3. ARTdeConv Reveals Changes in Key PBMC Cell Type Proportion in COVID-19 Patients

We performed another extensive deconvolution analysis on PBMC bulk samples gathered from 17 healthy controls and 14 patients with COVID-19 diagnosis recruited for a systemic immunity assessment against COVID-19 infections in humans by Arunachalam et al. (2020). The study also classified the COVID-19 severity of infected patients. Patients in the study were designated with three levels of severity: moderate, severe, and Intensity Care Unit-hospitalized (ICU). Details regarding the patient recruitment and the classification of COVID-19 severity are in Supplementary Material Section F.1.

Bulk RNA-seq data downloaded using NCBI GEO accession number GSE152418 were utilized to perform deconvolution for estimating the proportions of the four major PBMC cell types: T cell, B cell, natural killer (NK) cell, and monocyte. The authors also included single cell RNA-seq data on separate independent blood samples from five healthy controls and seven COVID-19 patients. Of these, all healthy and six COVID-19 infected subjects had both bulk and single cell expression recorded. The single cell data were downloaded using GEO accession GSE155673 and were used to construct the partial signature matrix. Complete descriptions of the bulk and single cell data and their processing, as well as the creation of gene signature matrices, can be found in Supplementary Material Section F.1.

Deconvolution was performed separately on samples from healthy controls and COVID-19 patients using bulk and signature matrices that matched the disease status. Since we were unable to obtain the mean and range proportions of the cell types involved separately for healthy and diseased populations, we opted to use the same set of numbers in Supplementary Table 1 for running ARTdeConv. Detailed descriptions of the tuning grid and other ARTdeConv parameters can be found in Supplementary Material Section F.2.

The complete results of deconvolved cell type proportions from ARTdeConv are shown in Figure 8 (boxplots) as well as Supplementary Figures 2 and 3 (bar charts). We observed that in COVID-19 samples, the proportions of T cells were notably lower than in the healthy control samples (Figure 8). Among the diseased samples, T cell depletion were found in several severe and ICU samples (Figure 9). This could be explained by T lymphopenia, commonly observed on blood samples of COVID-19 patients with severe symptoms but less frequently on those from patients with mild to moderate symptoms, as a result of the immunological responses of T cells to COVID-19 (Chen and John Wherry, 2020). It was also observed that Severe or ICU samples exhibited higher monocyte abundances in PBMC (Figure 9). Similar trends have also been observed by Zhang et al. (2022) in blood samples of patients with severe COVID-19.

Arunachalam et al. (2020) used the abundances of cells from the single-cell samples as a surrogate measurement for true abundances (except for dendritic cells, which were manually enriched in the single-cell samples). The relative deconvolved abundances of T cells, B cells, NK cells, and monocytes were then compared against the relative abundances of single cells. ARTdeConv demonstrated satisfactory accuracy in deconvolving cell-type

Figure 8: Boxplots for estimated PBMC cell-type proportions for healthy control and COVID-19 infected samples using ARTdeConv.

Figure 9: T cell proportions of the COVID-19 infected were estimated lower for severe/ICU patients while monocyte proportions were estimated higher.

Figure 10: Scatterplots of ARTdeConv estimated vs single cell proportions of PBMCs on subjects with both bulk and single cell samples.

abundances (Figure 10), achieving a Pearson's correlation of 0.860 among healthy control samples (MAD = 0.098) and 0.717 among COVID-19 samples (MAD = 0.103).

While we observed a slightly lower accuracy among COVID-19 patients compared to healthy controls, the COVID-19 patients spanned several disease severity classes and various duration of infection. These factors might increase the variation of gene expression, both on the cell type and the bulk levels, making deconvolution more challenging. The analysis also identified one outlier in healthy controls, S066 (Supplementary Figure 3). Given the typical high abundances of T cells in human PBMC samples, it was surprising that ARTdeConv did not detect any T cell in this sample, while attributed monocytes as the most abundant cell type. After further explorations in the expression of *CD3* and *CD14* genes, which are experimentally confirmed marker genes of T cells and classical monocytes respectively (Kleiveland, 2015), we discovered exceptionally low bulk *CD3* expression (Supplemetary Figure 4) and high *CD14* expression (Supplemetary Figure 5) in the sample from S066, which were consistent with this result.

4.   Discussion

In this paper, we introduce ARTdeConv, an innovative deconvolution approach. An important feature of ARTdeConv is its adoption of a tri-factor model, which integrates the cell-type mRNA amounts during the deconvolution process. ARTdeConv offers enhanced flexibility compared to reference-based methods, as it accommodates cell types whose reference gene expression are not known, while presents advantages over reference-free methods by incorporating the partial signature matrix. Moreover, the method makes effective use of reference information on proportion means and ranges derived from external studies.

Additionally, we derive the MU algorithm for ARTdeConv and present a theorem that establishes the convergence of this algorithm to stationary points. This proof is derived by casting ARTdeConv's algorithm as a special case of the BSUM algorithm introduced by Razaviyayn et al. (2013).

In the simulation studies, ARTdeConv showcased superior deconvolution accuracy over reference-free methods in benchmarking both overall and on the cell type with missing reference expression. The analysis of real data further underscores the reliability of ARTdeConv, especially when cell-type mRNA amount information is obscured.

An argument can be made that obtaining precise reference information on proportions demands additional efforts in practice. However, the advantages of integrating this information are substantial. Without it, the task of associating estimated proportions with their respective cell types can prove challenging. In fact, as demonstrated in Section 3.1, we had to perform manual matching of results for all reference-free methods, which becomes unfeasible if multiple cell types lack reference expression.

While ARTdeConv provides a guarantee of convergence, it does not assure that the estimated proportions are globally optimal. This particular challenge is not unique to ARTdeConv and is encountered in other reference-free methods such as NMF. To counter this, we recommend employing marker genes for the deconvolution process and considering multiple restarts of ARTdeConv. Other methods like debCAM mitigates this issue by assuming that cell-type-specific genes can be found for each involved cell type and solving the deconvolution using a different mathematical framework. However, their performance can diminish under scenarios that deviate from their ideal assumption, which is a common occurrence. Even extensive efforts in optimizing signature matrices struggle to identify purely cell-type-specific marker genes for all cell types (Newman et al., 2015).

When we were performing the real data analyses, the datasets from Hoek et al. (2015) and Arunachalam et al. (2020) offer an added advantage, as both the expression of bulk samples and the partial signature matrix are collected from independent samples in the same study and approximately concurrently. This minimizes the influence of most technical artifacts. In situations where bulk and cell-type reference expression are derived from different studies, additional caution against those technical artifacts should be exercised during data pre-processing prior to deconvolution.

Software Availability

ARTdeConv is available as an R package for download from the GitHub repository: https://github.com/gr8lawrence/ARTDeConv.

Acknowledgements

The authors gratefully acknowledge the support of National Institute of Allergy and Infectious Diseases at the National Institutes of Health through the grant U19AI144181 and that of National Institute on Aging through the grant R01AG073259.

References

S. Anders and W. Huber. Differential expression analysis for sequence count data. *Nature Precedings*, pages 1–1, 2010.

P. S. Arunachalam, F. Wimmers, C. K. P. Mok, R. A. P. M. Perera, M. Scott, T. Hagan, N. Sigal, Y. Feng, L. Bristow, O. Tak-Yin Tsang, D. Wagh, J. Coller, K. L. Pellegrini, D. Kazmin, G. Alaaeddine, W. S. Leung, J. M. C. Chan, T. S. H. Chik, C. Y. C. Choi, C. Huerta, M. Paine McCullough, H. Lv, E. Anderson, S. Edupuganti, A. A. Upadhyay, S. E. Bosinger, H. T. Maecker, P. Khatri, N. Rouphael, M. Peiris, and B. Pulendran. Systems biological assessment of immunity to mild versus severe COVID-19 infection in humans. *Science*, 369(6508):1210–1220, Sept. 2020.

F. Avila Cobos, J. Alquicira-Hernandez, J. E. Powell, P. Mestdagh, and K. De Preter. Benchmarking of cell type deconvolution pipelines for transcriptomics data. *Nature communications*, 11(1):1–14, 2020.

B. Chen, M. S. Khodadoust, C. L. Liu, A. M. Newman, and A. A. Alizadeh. Profiling tumor infiltrating immune cells with cibersort. *Cancer Systems Biology: Methods and Protocols*, pages 243–259, 2018.

L. Chen, C.-T. Wu, N. Wang, D. M. Herrington, R. Clarke, and Y. Wang. debcam: a bioconductor r package for fully unsupervised deconvolution of complex tissues. *Bioinformatics*, 36(12):3927–3929, 2020.

Z. Chen and E. John Wherry. T cell responses in patients with COVID-19. *Nat. Rev. Immunol.*, 20(9):529–536, Sept. 2020.

F. Elloumi, Z. Hu, Y. Li, J. S. Parker, M. L. Gulley, K. D. Amos, and M. A. Troester. Systematic bias in genomic classification due to contaminating non-neoplastic tissue in breast tumor samples. *BMC medical genomics*, 4(1):1–12, 2011.

R. Gaujoux and C. Seoighe. A flexible r package for nonnegative matrix factorization. *BMC bioinformatics*, 11(1):1–9, 2010.

R. Gaujoux and C. Seoighe. Cellmix: a comprehensive toolbox for gene expression deconvolution. *Bioinformatics*, 29(17):2211–2212, 2013.

C. Hafemeister and R. Satija. Normalization and variance stabilization of single-cell rna-seq data using regularized negative binomial regression. *Genome biology*, 20(1): 1–15, 2019.

K. L. Hoek, P. Samir, L. M. Howard, X. Niu, N. Prasad, A. Galassie, Q. Liu, T. M. Allos, K. A. Floyd, Y. Guo, et al. A cell-based systems biology assessment of human blood to monitor immune responses after influenza vaccination. *PloS one*, 10(2):e0118528, 2015.

H. Jin and Z. Liu. A benchmark for rna-seq deconvolution analysis under dynamic testing environments. *Genome biology*, 22:1–23, 2021.

J. Kim, Y. He, and H. Park. Algorithms for nonnegative matrix and tensor factorizations: A unified view based on block coordinate descent framework. *Journal of Global Optimization*, 58(2):285–319, 2014.

C. R. Kleiveland. *Peripheral Blood Mononuclear Cells*. Springer, 2015.

K. Lange. *MM optimization algorithms*. SIAM, 2016.

D. Lee and H. S. Seung. Algorithms for non-negative matrix factorization. *Advances in neural information processing systems*, 13, 2000.

Z. Li and H. Wu. Toast: improving reference-free cell composition estimation by cross-cell type differential analysis. *Genome biology*, 20(1):1–17, 2019.

C.-J. Lin. On the convergence of multiplicative update algorithms for nonnegative matrix factorization. *IEEE Transactions on Neural Networks*, 18(6):1589–1596, 2007.

N. Lytal, D. Ran, and L. An. Normalization methods on single-cell rna-seq data: an empirical survey. *Frontiers in genetics*, 11:41, 2020.

S. Mohammadi, N. Zuckerman, A. Goldsmith, and A. Grama. A critical survey of deconvolution methods for separating cell types in complex tissues. *Proc. IEEE*, 105(2): 340–366, Feb. 2017.

J. S. Morris and S. Kopetz. Tumor microenvironment in gene signatures: critical biology or confounding noise? *Clinical Cancer Research*, 22(16):3989–3991, 2016.

A. M. Newman, C. L. Liu, M. R. Green, A. J. Gentles, W. Feng, Y. Xu, C. D. Hoang, M. Diehn, and A. A. Alizadeh. Robust enumeration of cell subsets from tissue expression profiles. *Nature methods*, 12(5):453–457, 2015.

Y. Qiu, J. Wang, J. Lei, and K. Roeder. Identification of cell-type-specific marker genes from co-expression patterns in tissue samples. *Bioinformatics*, 37(19):3228–3234, 2021.

J. Racle, K. de Jonge, P. Baumgaertner, D. E. Speiser, and D. Gfeller. Simultaneous enumeration of cancer and immune cell types from bulk tumor gene expression data. *elife*, 6:e26476, 2017.

Z. Rahil, R. Leylek, C. M. Schürch, H. Chen, Z. Bjornson-Hooper, S. R. Christensen, P. F. Gherardini, S. S. Bhate, M. H. Spitzer, G. K. Fragiadakis, et al. Landscape of coordinated immune responses to h1n1 challenge in humans. *The Journal of clinical investigation*, 130(11):5800–5816, 2020.

M. Razaviyayn, M. Hong, and Z.-Q. Luo. A unified convergence analysis of block successive minimization methods for nonsmooth optimization. *SIAM Journal on Optimization*, 23 (2):1126–1153, 2013.

M. H. Spitzer and G. P. Nolan. Mass cytometry: single cells, many features. *Cell*, 165(4): 780–791, 2016.

A. Takami, S. Watanabe, Y. Yamamoto, H. Miyachi, Y. Bamba, M. Ohata, S. Mishima, H. Kubota, A. Nishiura, T. Inaba, M. Enomoto, T. Mitsuhashi, K. Nakanishi, R. Miura, E. Nonaka, K. Shimbo, Y. Yatomi, K. Tohyama, and Japanese Society for Laboratory Hematology subcommittee on Standardization of Blood Cell Morphology (JSLH-SBCM). Reference intervals of white blood cell parameters for healthy adults in japan. *Int. J. Lab. Hematol.*, 43(5):948–958, Oct. 2021.

N. Wang, E. P. Hoffman, L. Chen, L. Chen, Z. Zhang, C. Liu, G. Yu, D. M. Herrington, R. Clarke, and Y. Wang. Mathematical modelling of transcriptional heterogeneity identifies novel markers and subpopulations in complex tissues. *Scientific reports*, 6(1): 18909, 2016.

X. Wang, J. Park, K. Susztak, N. R. Zhang, and M. Li. Bulk tissue cell type deconvolution with multi-subject single-cell expression reference. *Nature communications*, 10(1):1–9, 2019.

G. Wigerblad, Q. Cao, S. Brooks, F. Naz, M. Gadkari, K. Jiang, S. Gupta, L. O'Neil, S. Dell'Orso, M. J. Kaplan, et al. Single-cell analysis reveals the range of transcriptional states of circulating human neutrophils. *bioRxiv*, 2022.

K. Zaitsev, M. Bambouskova, A. Swain, and M. N. Artyomov. Complete deconvolution of cellular mixtures based on linearity of transcriptional signatures. *Nature communications*, 10(1):1–16, 2019.

Y. Zhang, S. Wang, H. Xia, J. Guo, K. He, C. Huang, R. Luo, Y. Chen, K. Xu, H. Gao, J. Sheng, and L. Li. Identification of monocytes associated with severe COVID-19 in the PBMCs of severely infected patients through Single-Cell transcriptome sequencing. *Engineering*, 17:161–169, Oct. 2022.

Supplementary Material

A. Technical Assumptions

Before presenting the main results, we state the technical assumptions.

Assumption 1 (Initial conditions). *(i)* $\forall k$, $\exists j_k \in \{1, \ldots, m\}$ *such that* $\theta_{j_k k}^0 > 0$; *(ii)* $\forall k$, $\exists i_k \in \{1, \ldots, n\}$ *such that* $p_{k i_k}^0 > 0$. *(iii)* $\forall i' \in \{i_1, i_2, \ldots, i_K\}$, $\forall j$, $Y_{ji'} > 0$; $\forall j' \in \{j_1, j_2, \ldots, j_K\}$, $\forall i$, $Y_{j'i} > 0$.

Assumption 2 (Boundedness). *(i) For any* $1 \le j \le m$ *and* $1 \le i \le n$, $Y_{ji} \le M_Y < \infty$. *(ii) For any* $1 \le j \le m$ *and* $1 \le k \le K_0$, $\Theta_{0_{jk}} \le M_\Theta < \infty$.

These assumptions are reasonable in practice. For every cell type with established reference expression, it is safe to assume that there exist some genes that display non-zero cell-type-specific expression, thereby fulfilling Assumption 1(*i*). For cell types lacking reference expression, an initial value in $\mathbf{\Theta}^0$ can be straightforwardly set to a positive value, thereby also satisfying the same assumption. Assumption 1(*ii*) is automatically satisfied due to how $\boldsymbol{P}^0$ is initialized in Algorithm 1. Assumption 1(*iii*) plays a pivotal role in ensuring that positive initialized values remain positive throughout the iterative process (see Supplementary Material Section B for details). This assumption is easy to verify and holds true unless the bulk expression matrix is excessively sparse, which is uncommon in practice. Assumption 2 is a mild assumption on upper bounds of gene expression.

B. The MU Steps

B.1. A Preliminary Lemma

Similar to Lee and Seung (2000), $f(\mathbf{\Theta}, \boldsymbol{s}, \boldsymbol{P})$ is minimized using block-wise auxiliary functions that are quadratic over-estimators, which then leads to the MU steps in (4), (5), and (6). The claims that these updates will keep the positive initialized values positive will be investigated at the end of this section. An auxiliary function is defined next.

Definition 1 (Auxiliary function). *Given a vector space* V *and* $\boldsymbol{v}, \boldsymbol{v}' \in V$, *a function* $g(\boldsymbol{v}|\boldsymbol{v}') : V \times V \mapsto \mathbb{R}$ *is an auxiliary function for the function* $f(\boldsymbol{v}) : V \mapsto \mathbb{R}$ *at* $\boldsymbol{v}'$ *if the following conditions are satisfied:*

$$g(\boldsymbol{v}|\boldsymbol{v}') \ge f(\boldsymbol{v}'), \forall \boldsymbol{v} \in V, \ and \ g(\boldsymbol{v}'|\boldsymbol{v}') = f(\boldsymbol{v}').$$

As we shall see, the vector blocks in the auxiliary functions would be each row of $\mathbf{\Theta}$, each column of $\boldsymbol{P}$, and $\boldsymbol{s}$. To find the specific auxiliary functions for $f(\mathbf{\Theta}, \boldsymbol{s}, \boldsymbol{P})$, a slightly modified form of a technical lemma given by Lin (2007) is needed and stated next. Its proof is also given in the same paper and thus omitted.

Lemma 1. *Given a positive semi-definite matrix* $\boldsymbol{Q} \in \mathbb{R}_+^{b \times b}$ *and* $\boldsymbol{w} \in \mathbb{R}_+^b$, *let* $I \subseteq \{1, \ldots, b\}$ *be a set of indices such that* $w_k > 0$ *for any* $k \in I$, $\boldsymbol{Q}_{II} \in \mathbb{R}_{++}^{|I| \times |I|}$, *and* $\boldsymbol{w}_I \in \mathbb{R}_{++}^{|I|}$ *be the sub-matrix of* $\boldsymbol{Q}$ *and the sub-vector of* $\boldsymbol{w}$ *with indices in* I, *where* $|I|$ *denotes the cardinality*

of I. Then,

$$diag\left\{\frac{(\boldsymbol{Q}_{II}\boldsymbol{w}_I)}{\boldsymbol{w}_I}\right\} - \boldsymbol{Q}_{II} = \begin{bmatrix} \frac{(\boldsymbol{Q}_{II}\boldsymbol{w}_I)_1}{w_{I1}} & & \\ & \ddots & \\ & & \frac{(\boldsymbol{Q}_{II}\boldsymbol{w}_I)_{|I|}}{w_{I|I|}} \end{bmatrix} - \boldsymbol{Q}_{II}$$

is always positive semi-definite.

With Lemma 1, the auxiliary functions and multiplicative update (MU) steps are ready to be derived.

B.2. Derivations of The MU Steps

In addition to those in Section 2.1, we introduce two additional notation: for two square matrices of the same dimension, we write $\boldsymbol{A} \succ \boldsymbol{B}$ and $\boldsymbol{A} \succeq \boldsymbol{B}$ if $\boldsymbol{A} - \boldsymbol{B}$ is positive-definite and positive-semidefinite, respectively.

The auxiliary function and update step for each row $\boldsymbol{\theta}_j^\top$ is derived and then combined for $j = 1, \ldots, m$ into the form of (4). For simplicity in notations, the index j is omitted in subscripts. The iteration number t is also omitted for $\boldsymbol{s}$ and $\boldsymbol{P}$ since both are treated as fixed. That is, we denote $\boldsymbol{y}_j^\top$ as $\boldsymbol{y}^\top$, $\boldsymbol{\theta}_j^\top$ as $\boldsymbol{\theta}^\top$, and $\boldsymbol{\delta}_j^\top$ as $\boldsymbol{\delta}$. We also define $\boldsymbol{\delta}_j^c = \boldsymbol{J} - \boldsymbol{\delta}_j$ for all j. Thus, for the j-th row of $\boldsymbol{\Theta}$, we can define the sub-problem of $f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P})$ as

$$\underset{\boldsymbol{\theta} \in \mathbb{R}_+^K}{\operatorname{argmin}} \frac{1}{2mn} \|\boldsymbol{y}^\top - \boldsymbol{\theta}^\top \operatorname{diag}(\boldsymbol{s})\boldsymbol{P}\|_2^2 + \frac{\alpha_1}{2}\|\boldsymbol{\delta}^\top \odot (\boldsymbol{\theta}^\top - \boldsymbol{\theta}_0^\top)\|_2^2 + \frac{\alpha_2}{2}\|\boldsymbol{\delta}^{c\top} \odot \boldsymbol{\theta}^\top\|_2^2 + C_1(\boldsymbol{s}, \boldsymbol{P}) \tag{S.8}$$

$$= \underset{\boldsymbol{\theta} \in \mathbb{R}_+^K}{\operatorname{argmin}} \frac{1}{2mn} \|\boldsymbol{y}^\top - \boldsymbol{P}^\top \operatorname{diag}(\boldsymbol{s})\boldsymbol{\theta}\|_2^2 + \frac{\alpha_1}{2}\|\boldsymbol{\delta} \odot (\boldsymbol{\theta} - \boldsymbol{\theta}_0)\|_2^2 + \frac{\alpha_2}{2}\|\boldsymbol{\delta}^c \odot \boldsymbol{\theta}\|_2^2 + C_1(\boldsymbol{s}, \boldsymbol{P}), \tag{S.9}$$

where $\operatorname{diag}(\boldsymbol{s})$ and $\boldsymbol{P}$ are treated as fixed and $C_1(\boldsymbol{s}, \boldsymbol{P})$ is a constant with respect to $\boldsymbol{\theta}$. We call the function in (S.9) $f_\theta(\boldsymbol{\theta})$. (S.9) is derived from (S.8) by transposing. Now, to find the auxiliary function $h_\theta(\boldsymbol{\theta}|\boldsymbol{\theta}^t)$, by performing a second-order Taylor series expansion on $f_\theta(\boldsymbol{\theta})$ at $\boldsymbol{\theta} = \boldsymbol{\theta}^t$, we have

$$f_\theta(\boldsymbol{\theta}) = f_\theta(\boldsymbol{\theta}^t) + \nabla_{\boldsymbol{\theta}} f_\theta(\boldsymbol{\theta}^t)^\top(\boldsymbol{\theta} - \boldsymbol{\theta}^t) + \frac{1}{2}(\boldsymbol{\theta} - \boldsymbol{\theta}^t)^\top \nabla_{\boldsymbol{\theta}}^2 f_\theta(\boldsymbol{\theta}^t)(\boldsymbol{\theta} - \boldsymbol{\theta}^t).$$

Next, letting $\boldsymbol{V}(\boldsymbol{\delta}) = \operatorname{diag}(\boldsymbol{\delta})$ and $\boldsymbol{V}^c(\boldsymbol{\delta}^c) = \operatorname{diag}(\boldsymbol{\delta}^c)$, it can be observed that $\boldsymbol{\delta} \odot \boldsymbol{\theta} = \boldsymbol{V}\boldsymbol{\theta}$ and $\boldsymbol{\delta}^c \odot \boldsymbol{\theta} = \boldsymbol{V}^c\boldsymbol{\theta}$. Also, $\boldsymbol{V} + \boldsymbol{V}^c = \boldsymbol{I}$. The first and second derivatives of $f_\theta(\boldsymbol{\theta})$ are thus:

$$\nabla_{\boldsymbol{\theta}} f_\theta(\boldsymbol{\theta}) = \frac{1}{mn}\operatorname{diag}(\boldsymbol{s})\boldsymbol{P}(\boldsymbol{P}^\top \operatorname{diag}(\boldsymbol{s})\boldsymbol{\theta} - \boldsymbol{y}^\top) + \alpha_1 \boldsymbol{V}(\boldsymbol{\theta} - \boldsymbol{\theta}_0) + \alpha_2 \boldsymbol{V}^c\boldsymbol{\theta},$$

and

$$\nabla_{\boldsymbol{\theta}}^2 f_\theta(\boldsymbol{\theta}) = \frac{1}{mn}\operatorname{diag}(\boldsymbol{s})\boldsymbol{P}\boldsymbol{P}^\top \operatorname{diag}(\boldsymbol{s}) + \alpha_1 \boldsymbol{V} + \alpha_2 \boldsymbol{V}^c.$$

It can be guaranteed $\nabla_{\boldsymbol{\theta}}^2 f_\theta(\boldsymbol{\theta}) \succ \boldsymbol{0}$ given $\alpha_1, \alpha_2 > 0$, as $\nabla_{\boldsymbol{\theta}}^2 f_\theta(\boldsymbol{\theta}) \succeq \alpha_1 \boldsymbol{V} + \alpha_2 \boldsymbol{V}^c \succeq \min(\alpha_1, \alpha_2)\boldsymbol{I} \succ \boldsymbol{0}$. Then, define $\boldsymbol{d}(\boldsymbol{\theta}^t) = \operatorname{diag}(\boldsymbol{\theta}^t)^{-1}\nabla_{\boldsymbol{\theta}}^2 f_\theta(\boldsymbol{\theta}^t)\boldsymbol{\theta}^t$, and $\boldsymbol{D}(\boldsymbol{\theta}^t) = \operatorname{diag}(\boldsymbol{d}(\boldsymbol{\theta}^t))$. Lemma 1 implies that, on coordinates that correspond to the positive elements of $\boldsymbol{\theta}$,

$\boldsymbol{D}(\boldsymbol{\theta}^t) - \nabla_{\boldsymbol{\theta}}^2 f_\theta(\boldsymbol{\theta}^t) \succeq 0$. Define $\mathbb{R}_{++}^{K_1}$, where $K_1 \leq K$, as the open half space that contains this positive subset of $\boldsymbol{\theta}$. For simplification, we use the same notations for the full vectors and matrices on those positive coordinates. As mentioned in Section 2.2, we shall see that once a coordinate of $\boldsymbol{\theta}$ reaches zero, it stays zero thereafter, which justifies the focus on these positive coordinates. Next, define

$$h_\theta(\boldsymbol{\theta}|\boldsymbol{\theta}^t) = f_\theta(\boldsymbol{\theta}^t) + \nabla_{\boldsymbol{\theta}} f_\theta(\boldsymbol{\theta}^t)^\top (\boldsymbol{\theta} - \boldsymbol{\theta}^t) + \frac{1}{2}(\boldsymbol{\theta} - \boldsymbol{\theta}^t)^\top \boldsymbol{D}(\boldsymbol{\theta}^t)(\boldsymbol{\theta} - \boldsymbol{\theta}^t), \qquad \text{(S.10)}$$

which is an auxiliary function of (S.9) (thus of $f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P})$ when we combine all sub-problems together for $j = 1, \ldots m$). To demonstrate that it satisfies Definition 1, simply see that

$$h_\theta(\boldsymbol{\theta}|\boldsymbol{\theta}^t) - f_\theta(\boldsymbol{\theta}) = \frac{1}{2}(\boldsymbol{\theta} - \boldsymbol{\theta}^t)^\top (\boldsymbol{D}(\boldsymbol{\theta}^t) - \nabla_{\boldsymbol{\theta}}^2 f_\theta(\boldsymbol{\theta}^t))(\boldsymbol{\theta} - \boldsymbol{\theta}^t) \geq 0, \ \forall \boldsymbol{\theta}, \boldsymbol{\theta}^t \in \mathbb{R}_{++}^{K_1}$$

and due to Lemma 1 and that $h_\theta(\boldsymbol{\theta}^t|\boldsymbol{\theta}^t) - f_\theta(\boldsymbol{\theta}^t) = 0$, $\forall \boldsymbol{\theta}^t \in \mathbb{R}_{++}^{K_1}$. To derive the MU step from this auxiliary function, we find $\boldsymbol{\theta}^{t+1}$ such that

$$\boldsymbol{\theta}^{t+1} = \operatorname*{argmin}_{\boldsymbol{\theta} \in \mathbb{R}_+^{K_1}} h_\theta(\boldsymbol{\theta}|\boldsymbol{\theta}^t). \qquad \text{(S.11)}$$

To guarantee the existence of a solution for (S.11), the feasible set of $\boldsymbol{\theta}$ is a closed space that includes 0. Since $\boldsymbol{D}(\boldsymbol{\theta}^t) - \nabla_{\boldsymbol{\theta}}^2 f_\theta(\boldsymbol{\theta}^t) \succeq 0$ and $\nabla_{\boldsymbol{\theta}}^2 f_\theta(\boldsymbol{\theta}^t) \succ 0$ by the assumptions, $\boldsymbol{D}(\boldsymbol{\theta}^t) \succ 0$, which implies that (S.10) is strictly convex. Thus, (S.10) has a global minimizer in the closed half space $\mathbb{R}_+^{K_1}$. By the optimality condition, if a feasible $\tilde{\boldsymbol{\theta}} \in \mathbb{R}_+^{K_1}$ satisfies $\nabla_{\boldsymbol{\theta}} h_\theta(\tilde{\boldsymbol{\theta}}|\boldsymbol{\theta}^t) = 0$, then $\tilde{\boldsymbol{\theta}}$ minimizes (S.10) globally and $\boldsymbol{\theta}^{t+1} = \tilde{\boldsymbol{\theta}}$ becasue of (S.11). Thus, we solve

$$\nabla_{\boldsymbol{\theta}} h_\theta(\boldsymbol{\theta}^{t+1}|\boldsymbol{\theta}^t) = \nabla_{\boldsymbol{\theta}} f_\theta(\boldsymbol{\theta}^t) + \boldsymbol{D}(\boldsymbol{\theta}^t)(\boldsymbol{\theta}^{t+1} - \boldsymbol{\theta}^t) = 0. \qquad \text{(S.12)}$$

Because $\boldsymbol{D}(\boldsymbol{\theta}^t)$ is diagonal, for each $k \in 1, \ldots, K_1$, (S.12) implies

$$\theta_k^{t+1} = \theta_k^t - \frac{\nabla_{\boldsymbol{\theta}} f_\theta(\boldsymbol{\theta}^t)_k}{D(\boldsymbol{\theta}^t)_k}. \qquad \text{(S.13)}$$

We further simplify the notations by letting

$$\begin{aligned}
\nabla_{\boldsymbol{\theta}} f_\theta(\boldsymbol{\theta}) &= \frac{1}{mn} \operatorname{diag}(\boldsymbol{s}) \boldsymbol{P} (\boldsymbol{P}^\top \operatorname{diag}(\boldsymbol{s}) \boldsymbol{\theta} - \boldsymbol{y}^\top) + \alpha_1 \boldsymbol{V}(\boldsymbol{\theta} - \boldsymbol{\theta}_0) + \alpha_2 \boldsymbol{V}^c \boldsymbol{\theta} \\
&= \left(\frac{1}{mn} \operatorname{diag}(\boldsymbol{s}) \boldsymbol{P} \boldsymbol{P}^\top \operatorname{diag}(\boldsymbol{s}) + \alpha_1 \boldsymbol{V} + \alpha_2 \boldsymbol{V}^c \right) \boldsymbol{\theta} \\
&\quad - \left(\frac{1}{mn} \operatorname{diag}(\boldsymbol{s}) \boldsymbol{P} \boldsymbol{P}^\top \operatorname{diag}(\boldsymbol{s}) \boldsymbol{y}^\top + \alpha_1 \boldsymbol{V} \boldsymbol{\theta}_0 \right) \\
&= \boldsymbol{A} \boldsymbol{\theta} - \boldsymbol{b}.
\end{aligned}$$

It follows immediately that $\nabla_{\boldsymbol{\theta}}^2 f_\theta(\boldsymbol{\theta}) = \boldsymbol{A}$. Plugging in the values of $\nabla_{\boldsymbol{\theta}} f_\theta(\boldsymbol{\theta}^t)_k$ and $D(\boldsymbol{\theta}^t)_k$

to (S.13), we have

$$
\begin{aligned}
\theta_k^{t+1} &= \theta_k^t - \frac{(\boldsymbol{A}\boldsymbol{\theta}^t - \boldsymbol{b})_k}{\frac{1}{\theta_k^t}(\boldsymbol{A}\boldsymbol{\theta}^t)_k} \\
&= \theta_k^t - \frac{(\boldsymbol{A}\boldsymbol{\theta}^t)_k}{(\boldsymbol{A}\boldsymbol{\theta}^t)_k}\theta_k^t + \frac{b_k}{(\boldsymbol{A}\boldsymbol{\theta}^t)_k}\theta_k^t \\
&= \frac{b_k}{(\boldsymbol{A}\boldsymbol{\theta}^t)_k}\theta_k^t \\
&= \frac{\left(\frac{1}{mn}\mathrm{diag}(\boldsymbol{s})\boldsymbol{P}\boldsymbol{y}^\top + \alpha_1\boldsymbol{V}\boldsymbol{\theta}_0\right)_k}{\left[\left(\frac{1}{mn}\mathrm{diag}(\boldsymbol{s})\boldsymbol{P}\boldsymbol{P}^\top\mathrm{diag}(\boldsymbol{s}) + \alpha_1\boldsymbol{V} + \alpha_2\boldsymbol{V}^c\right)\boldsymbol{\theta}^t\right]_k}\theta_k^t \\
&= \frac{\left(\mathrm{diag}(\boldsymbol{s})\boldsymbol{P}\boldsymbol{y}^\top + mn\alpha_1\boldsymbol{V}\boldsymbol{\theta}_0\right)_k}{\left[(\mathrm{diag}(\boldsymbol{s})\boldsymbol{P}\boldsymbol{P}^\top\mathrm{diag}(\boldsymbol{s}) + mn(\alpha_1\boldsymbol{V} + \alpha_2\boldsymbol{V}^c))\boldsymbol{\theta}^t\right]_k}\theta_k^t. \quad\text{(S.14)}
\end{aligned}
$$

The multiplicative form of (S.14) guarantees that $\boldsymbol{\theta}^{t+1}$ is feasible. Therefore, $\boldsymbol{\theta}^{t+1}$ satisfies (S.11).

It is straightforward to check that (S.14) also applies to zero coordinates of $\boldsymbol{\theta}^t$ ($\{\theta_k^t : \theta_k^t \in \mathbb{R}_+^K \setminus \mathbb{R}_{++}^{K_1}\}$), as zero coordinates remain zero after any multiplication. Therefore, it is the unified update step for the entire $\boldsymbol{\theta}^t \in \mathbb{R}_+^K$. Since it corresponds to the update for the (j, k)-th element in $\boldsymbol{\Theta}$, combining all elements gives (4).

We remark that (S.13) reflects the block coordinate descent nature of the multiplicative update in $\boldsymbol{\Theta}$, as observed by Kim et al. (2014). In fact, the update steps for $\boldsymbol{P}$ and $\mathrm{diag}(\boldsymbol{s})$ can be written in similar fashions.

The update step for $\boldsymbol{P}$ in (5) can be similarly derived as that for $\boldsymbol{\Theta}$. We begin with specifying the sub-problem for $\boldsymbol{p}_i$, $i = 1, \ldots, n$, treating $\boldsymbol{\Theta}$ and $\mathrm{diag}(\boldsymbol{s})$ as fixed and omitting the subscripts and the iteration number (t for $\boldsymbol{s}$ and $t+1$ for $\boldsymbol{\Theta}$) for convenience in notations:

$$
\underset{\boldsymbol{p}\in\mathbb{R}_+^K}{\mathrm{argmin}}\ f_p(\boldsymbol{p}) = \frac{1}{2mn}\|\boldsymbol{y} - \boldsymbol{\Theta}\mathrm{diag}(\boldsymbol{s})\boldsymbol{p}\|_2^2 + \frac{\beta}{2}\boldsymbol{\rho}^{-1}\|\boldsymbol{p} - \boldsymbol{m}\|_2^2 + K_2(\boldsymbol{\Theta}, \boldsymbol{s}), \quad\text{(S.15)}
$$

where $K_2(\boldsymbol{\Theta}, \boldsymbol{s})$ only depends on $\boldsymbol{\Theta}$ and $\boldsymbol{s}$ and is a constant with respect to $\boldsymbol{P}$. Then, its first and second derivatives with respect to $\boldsymbol{p}$ are

$$
\begin{aligned}
\nabla_{\boldsymbol{p}}f_p(\boldsymbol{p}) &= \left(\frac{1}{mn}\mathrm{diag}(\boldsymbol{s})\boldsymbol{\Theta}^\top\boldsymbol{\Theta}\mathrm{diag}(\boldsymbol{s}) + \beta\boldsymbol{\rho}^{-1}\right)\boldsymbol{p} - \left(\frac{1}{mn}\mathrm{diag}(\boldsymbol{s})\boldsymbol{\Theta}^\top\boldsymbol{y} + \beta\boldsymbol{\rho}^{-1}\boldsymbol{m}\right) \\
&= \boldsymbol{C}\boldsymbol{p} - \boldsymbol{q}, \quad\text{(S.16)}
\end{aligned}
$$

and

$$
\nabla_{\boldsymbol{p}}^2 f_p(\boldsymbol{p}) = \frac{1}{mn}\mathrm{diag}(\boldsymbol{s})\boldsymbol{\Theta}^\top\boldsymbol{\Theta}\mathrm{diag}(\boldsymbol{s}) + \beta\boldsymbol{\rho}^{-1} = \boldsymbol{C}. \quad\text{(S.17)}
$$

To find the auxiliary function of (S.15), define $\boldsymbol{d}(\boldsymbol{p}^t) = \mathrm{diag}(\boldsymbol{p}^t)^{-1}\nabla_{\boldsymbol{p}}^2 f_p(\boldsymbol{p}^t)\boldsymbol{p}^t$, and $\boldsymbol{D}(\boldsymbol{p}^t) = \mathrm{diag}(\boldsymbol{d}(\boldsymbol{p}^t))$, and

$$
h_p(\boldsymbol{p}|\boldsymbol{p}^t) = f_p(\boldsymbol{p}^t) + \nabla_{\boldsymbol{p}}f_p(\boldsymbol{p}^t)^\top(\boldsymbol{p} - \boldsymbol{p}^t) + \frac{1}{2}(\boldsymbol{p} - \boldsymbol{p}^t)^\top\boldsymbol{D}(\boldsymbol{p}^t)(\boldsymbol{p} - \boldsymbol{p}^t). \quad\text{(S.18)}
$$

On the $K_2 \leq K$ positive coordinates in $\boldsymbol{p}^t$, (S.18) is the auxiliary function of (S.15) due to Lemma 1. The verification is the same to that for $h_\theta(\boldsymbol{\theta}|\boldsymbol{\theta}^t)$ and omitted here. Next, we find $\boldsymbol{p}^{t+1}$ such that

$$
\boldsymbol{p}^{t+1} = \underset{\boldsymbol{p}\in\mathbb{R}_+^{K_2}}{\mathrm{argmin}}\ h_p(\boldsymbol{p}|\boldsymbol{p}^t).
$$

This in turn corresponds to finding $\boldsymbol{p}^{t+1}$ such that

$$\nabla_{\boldsymbol{p}} h_p(\boldsymbol{p}^{t+1}|\boldsymbol{p}^t) = \nabla_{\boldsymbol{p}} f_p(\boldsymbol{p}^t) + \boldsymbol{D}(\boldsymbol{p}^t)(\boldsymbol{p}^{t+1} - \boldsymbol{p}^t) = 0,$$

given that such a $\boldsymbol{p}^{t+1}$ is feasible. After some algebra, for any $k = 1, \dots, K_2$,

$$
\begin{aligned}
p_k^{t+1} &= \frac{q_k}{(\boldsymbol{C}\boldsymbol{p}^t)_k} p_k^t \\
&= \frac{\left(\frac{1}{mn}\mathrm{diag}(\boldsymbol{s})\boldsymbol{\Theta}^\top \boldsymbol{y} + \beta \boldsymbol{\rho}^{-1}\boldsymbol{m}\right)_k}{\left\{\left(\frac{1}{mn}\mathrm{diag}(\boldsymbol{s})\boldsymbol{\Theta}^\top \boldsymbol{\Theta}\mathrm{diag}(\boldsymbol{s}) + \beta \boldsymbol{\rho}^{-1}\right)\boldsymbol{p}\right\}_k} p_k^t \\
&= \frac{\left(\mathrm{diag}(\boldsymbol{s})\boldsymbol{\Theta}^\top \boldsymbol{y} + mn\beta \boldsymbol{\rho}^{-1}\boldsymbol{m}\right)_k}{\left\{\left(\mathrm{diag}(\boldsymbol{s})\boldsymbol{\Theta}^\top \boldsymbol{\Theta}\mathrm{diag}(\boldsymbol{s}) + mn\beta \boldsymbol{\rho}^{-1}\right)\boldsymbol{p}\right\}_k} p_k^t,
\end{aligned}
\tag{S.19}
$$

which produces a feasible $\boldsymbol{p}^{t+1}$. Moreover, (S.19) also applies to the zero coordinates in $\boldsymbol{p}^t$. Since (S.19) corresponds to the update step for the (k, i)-th element for the $\boldsymbol{P}$ matrix, combining all of these elements yields (5).

Deriving the updates step for $\mathrm{diag}(\boldsymbol{s})$ in (6) largely follows the same footsteps as for $\boldsymbol{\Theta}$ and $\boldsymbol{P}$. To find a suitable sub-problem, we start from the following identity

$$\boldsymbol{\Theta}\mathrm{diag}(\boldsymbol{s})\boldsymbol{P} = \boldsymbol{\Theta}\mathrm{diag}(\boldsymbol{s})\boldsymbol{P} = \sum_{k=1}^{K} s_k \boldsymbol{\theta}_k \boldsymbol{p}_k^\top. \tag{S.20}$$

Denoting $\boldsymbol{G}_k = \boldsymbol{\theta}_k \boldsymbol{p}_k^\top$ and only assuming $\boldsymbol{s}$ a variable, we transform the objective function in $f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P})$ with respect to $\boldsymbol{s}$ as follows

$$
\begin{aligned}
f_s(\boldsymbol{s}) &= \frac{1}{2mn}\|\boldsymbol{Y} - \boldsymbol{\Theta}\mathrm{diag}(\boldsymbol{s})\boldsymbol{P}\|_F^2 + C_3(\boldsymbol{\Theta}, \boldsymbol{P}) && \text{(by rewriting } f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P})) \\
&= \frac{1}{2mn}\|\boldsymbol{Y} - \sum_{k=1}^{K} s_k \boldsymbol{G}_k\|_F^2 + C_3(\boldsymbol{\Theta}, \boldsymbol{P}) && \text{(by (S.20))} \\
&\propto \frac{1}{2}\left\{\sum_{k=1}^{K} s_k^2 \mathrm{tr}(\boldsymbol{G}_k^\top \boldsymbol{G}_k) - 2\sum_{k=1}^{K} s_k \mathrm{tr}(\boldsymbol{G}_k^\top \boldsymbol{Y}) + \sum_{k \neq l} s_k s_l \mathrm{tr}(\boldsymbol{G}_k^\top \boldsymbol{G}_l)\right\} + C_3'(\boldsymbol{\Theta}, \boldsymbol{P}) \\
& && \text{(by expanding the squared Frobenius norm and dropping } m, n) \\
&= \frac{1}{2}\left\{\sum_{k=1}^{K} s_k^2 \mathrm{tr}(\boldsymbol{G}_k^\top \boldsymbol{G}_k) + \sum_{k \neq l} s_k s_l \mathrm{tr}(\boldsymbol{G}_k^\top \boldsymbol{G}_l)\right\} - \sum_{k=1}^{K} s_k \mathrm{tr}(\boldsymbol{G}_k^\top \boldsymbol{Y}) + C_3'(\boldsymbol{\Theta}, \boldsymbol{P}) \\
& && \text{(by rearranging the last step)} \\
&= \frac{1}{2}\boldsymbol{s}^\top \boldsymbol{Z}\boldsymbol{s} - \boldsymbol{u}^\top \boldsymbol{s} + C_3'(\boldsymbol{\Theta}, \boldsymbol{P}), && \text{(S.21)}
\end{aligned}
$$

where $\boldsymbol{Z} \in \mathbb{R}_+^{K \times K}$, $Z_{ij} = \mathrm{tr}(\boldsymbol{G}_i^\top \boldsymbol{G}_j) = \mathrm{tr}(\boldsymbol{p}_i \boldsymbol{\theta}_i^\top \boldsymbol{\theta}_j \boldsymbol{p}_j^\top)$, $\boldsymbol{u} \in \mathbb{R}_+^K$, and $u_k = \mathrm{tr}(\boldsymbol{Y}^\top \boldsymbol{G}_k) = \mathrm{tr}(\boldsymbol{Y}^\top \boldsymbol{\theta}_k \boldsymbol{p}_k^\top)$. Both C_3 and C_3' only depend on $\boldsymbol{\Theta}$ and $\boldsymbol{P}$, hence are constant with respect to $\boldsymbol{s}$. (S.21) is then our sub-problem for $\boldsymbol{s}$. From (S.21), the first and second derivatives for $\boldsymbol{s}$ are $\nabla_{\boldsymbol{s}} f_s(\boldsymbol{s}) = \boldsymbol{Z}\boldsymbol{s} - \boldsymbol{u}$, and $\nabla_{\boldsymbol{s}}^2 f_s(\boldsymbol{s}) = \boldsymbol{Z}$. Again, define $\boldsymbol{d}(\boldsymbol{s}^t) = \mathrm{diag}(\boldsymbol{s}^t)^{-1}\nabla_{\boldsymbol{s}}^2 f_s(\boldsymbol{s}^t)\boldsymbol{s}^t$, $\boldsymbol{D}(\boldsymbol{s}^t) = \mathrm{diag}(\boldsymbol{d}(\boldsymbol{s}^t))$, and

$$h_s(\boldsymbol{s}|\boldsymbol{s}^t) = f_s(\boldsymbol{s}^t) + \nabla_{\boldsymbol{s}} f_s(\boldsymbol{s}^t)^\top(\boldsymbol{s} - \boldsymbol{s}^t) + \frac{1}{2}(\boldsymbol{s} - \boldsymbol{s}^t)^\top \boldsymbol{D}(\boldsymbol{s}^t)(\boldsymbol{s} - \boldsymbol{s}^t). \tag{S.22}$$

(S.22) is an auxiliary function of $f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P})$ for $\boldsymbol{s}$ according to Lemma 1. Since $\boldsymbol{s} \in \mathbb{R}_{++}^K$, there is no need for sub-setting to positive coordinates. Thus, we obtain $\boldsymbol{s}^{t+1}$ by finding a feasible

$$\boldsymbol{s}^{t+1} = \operatorname*{argmin}_{\boldsymbol{s} \in \mathbb{R}_+^K} h_s(\boldsymbol{s}|\boldsymbol{s}^t). \tag{S.23}$$

We remark that this is a slight relaxation of the problem to update $\boldsymbol{s}$, since it is originally assumed that $\boldsymbol{s} \in \mathbb{R}_{++}^K$. However, this allows us to obtain a guaranteed closed form global minimizer for (S.23) in the closed half space $\mathbb{R}_+^K$, and the resulting update satisfies is feasible in $\mathbb{R}_{++}^K$ as long as not all values of $\boldsymbol{\theta}_k$ or $\boldsymbol{p}_k^\top$ are 0 for any $k = 1, \ldots, K$, which is stated as Assumption 1(i) and 1(ii). (S.23) leads to solving

$$\nabla_{\boldsymbol{s}} h_s(\boldsymbol{s}^{t+1}|\boldsymbol{s}^t) = \nabla_{\boldsymbol{s}} f_s(\boldsymbol{s}^t) + \boldsymbol{D}(\boldsymbol{s}^t)(\boldsymbol{s}^{t+1} - \boldsymbol{s}^t) = 0,$$

from which we have

$$s_k^{t+1} = \frac{u_k}{(\boldsymbol{Z}\boldsymbol{s}^t)_k} s_k^t. \tag{S.24}$$

(S.24) corresponds to the update for the k-th coordinate in (6), which yields feasible s_k^{t+1} under the assumptions, since $Z_{ij} = \operatorname{tr}(\boldsymbol{G}_i^\top \boldsymbol{G}_j) = (\boldsymbol{\theta}_i^\top \boldsymbol{\theta}_j)\operatorname{tr}(\boldsymbol{p}_i \boldsymbol{p}_j^\top)$, and $u_k = \operatorname{tr}(\boldsymbol{Y}^\top \boldsymbol{G}_k) = \operatorname{tr}(\boldsymbol{Y}^\top \boldsymbol{\theta}_k \boldsymbol{p}_k^\top)$. Moreover,

$$Z_{ij} = \operatorname{tr}(\boldsymbol{p}_i \boldsymbol{\theta}_i^\top \boldsymbol{\theta}_j \boldsymbol{p}_j^\top) = (\boldsymbol{\theta}_i^\top \boldsymbol{\theta}_j)(\boldsymbol{p}_j^\top \boldsymbol{p}_i), \tag{S.25}$$

and that

$$u_k = \operatorname{tr}(\boldsymbol{Y}^\top \boldsymbol{\theta}_k \boldsymbol{p}_k^\top) = \sum_{j=1}^m \sum_{i=1}^n y_{ji} \theta_{jk} p_{ki}. \tag{S.26}$$

For the denominator in (S.24), we see that

$$(\boldsymbol{Z}\boldsymbol{s}^t)_k = \sum_{l=1}^K Z_{kl} s_l^t \geq Z_{kk} s_k^t = \|\boldsymbol{\theta}_k\|_2^2 \|\boldsymbol{p}_k\|_2^2 s_k^t > 0, \tag{S.27}$$

where the last step is a result of (S.25) and the stability of updates based Assumption 1(i) and 1(ii), which will be discussed in details below. As for the numerator of (S.24), taking from the stability of updates due to Assumption 1(i) and 1(ii) and assuming $\theta_{j_k k} > 0$ and $p_{k i_k} > 0$, by (S.26), we have

$$u_k = \sum_{j=1}^m \sum_{i=1}^n y_{ji} \theta_{jk} p_{ki} \geq y_{j_k i_k} \theta_{j_k k} p_{k i_k} > 0,$$

as long as $y_{j_k i_k} > 0$. Given Assumption 1(iii), we conclude that $\boldsymbol{s}^{t+1} \in \mathbb{R}_{++}^K$.

B.3. Stability of The Update Steps

To justify the claim that for any j and k, $\theta_{jk}^{t+1} > 0$ whenever $\theta_{jk}^t > 0$ for $t = 0, 1, 2, \ldots$, notice that the denominator of (S.14) is automatically positive whenever $\alpha_1, \alpha_2 > 0$. For guaranteeing that its numerator is also positive, we want to show

$$\left(\operatorname{diag}(\boldsymbol{s}^t)\boldsymbol{P}^t \boldsymbol{y}_j^\top\right)_k = s_k^t \sum_{i=1}^n p_{ki}^t y_{ji} \geq s_k^t p_{k i_k}^t y_{j i_k} > 0, \tag{S.28}$$

where the index i_k is defined as in Assumption 1(*ii*). By Assumption 1(*iii*), $y_{ji_k} > 0$, and $s_k^t > 0$ as discussed above. Thus, (S.28) holds whenever $p_{ki_k}^t > 0$. Similarly, for proving the claim that for any k and i, $p_{ki}^{t+1} > 0$ whenever $p_{ki}^t > 0$, notice that the denominator for (S.19) is also positive whenever $\beta > 0$. To prove that its numerator is positive, we need to show

$$\left(\mathrm{diag}(\boldsymbol{s}^t)(\boldsymbol{\Theta}^{t+1})^\top \boldsymbol{y}_i\right)_k = s_k^t \sum_{j=1}^m \theta_{jk}^{t+1} y_{ji} \geq s_k^t \theta_{j_kk}^{t+1} y_{j_ki} > 0, \tag{S.29}$$

where j_k is defined as in Assumption 1(*i*). (S.29) holds whenever $\theta_{j_kk}^{t+1} > 0$.

Since by Assumption 1(*ii*) and Assumption 1(*i*), $p_{ki_k}^0 > 0$ and $\theta_{j_kk}^0 > 0$. Therefore, we can show that $p_{ki_k}^t > 0$ and $\theta_{j_kk}^{t+1} > 0$ for any $t = 0, 1, 2, \ldots$ by induction through (S.28) and (S.29). Thus, the claims on the stability of updates hold.

C. Proof of Theorem 1

To prove Theorem 1, we leverage on the convergence properties of the block successive upper-bound minimization (BSUM) algorithms proved by Razaviyayn et al. (2013). Namely, we can view Algorithm 1 as a BSUM algorithm by counting a total of $m + n + 1$ separate blocks which the algorithm minimizes successively: m for updating $\boldsymbol{\Theta}^t$, n for $\boldsymbol{P}^t$, and one for $\boldsymbol{s}^t$. To utilize such properties, we need to prove the following proposition, which states the conditions on the update steps' auxiliary functions for Algorithm 1 to formally qualify as BSUM, which is translated from Assumption 2 of Razaviyayn et al. (2013):

Proposition 1 (Algorithm 1 is BSUM). *$h_\theta(\boldsymbol{\theta}_j|\boldsymbol{\theta}_j^t)$, $h_p(\boldsymbol{p}_i|\boldsymbol{p}_i^t)$, and $h_s(\boldsymbol{s}|\boldsymbol{s}^t)$ satisfy the following conditions for any $j = 1, 2, \ldots, m$ and $i = 1, 2, \ldots, n$: (1) $h_{\theta_j}(\boldsymbol{\theta}_j|\boldsymbol{\theta}_j^t) \geq f_\theta(\boldsymbol{\theta}_j^t)$; $h_p(\boldsymbol{p}_i|\boldsymbol{p}_i^t) \geq f_p(\boldsymbol{p}_i^t)$; $h_s(\boldsymbol{s}|\boldsymbol{s}^t) \geq f_s(\boldsymbol{s}^t)$; (2) $h_{\theta_j}(\boldsymbol{\theta}_j|\boldsymbol{\theta}_j^t)$, $h_p(\boldsymbol{p}_i|\boldsymbol{p}_i^t)$, and $h_s(\boldsymbol{s}|\boldsymbol{s}^t)$ are continous on their domains; (3) Any directional derivative of $h_{\theta_j}(\boldsymbol{\theta}_j|\boldsymbol{\theta}_j^t)$, $h_p(\boldsymbol{p}_i|\boldsymbol{p}_i^t)$, and $h_s(\boldsymbol{s}|\boldsymbol{s}^t)$ within their domains are equal to the direactional derivative of $f_\theta(\boldsymbol{\theta}_j^t)$, $f_p(\boldsymbol{p}_i^t)$, and $f_s(\boldsymbol{s}^t)$ respectively.*

Proof. The proof is quite straightforward. (1) and (2) are direct from the construction of the surrogate functions and Lemma 1, which also manifests the majorization-minimization (MM) property (Lange, 2016) of each block's update. (3) is seen by differentiating (S.10), (S.18), and (S.22), which leads to $\nabla_{\boldsymbol{\theta}_j} h_\theta(\boldsymbol{\theta}_j^t|\boldsymbol{\theta}_j^t) = \nabla_{\boldsymbol{\theta}_j} f_\theta(\boldsymbol{\theta}_j^t), \forall j$, $\nabla_{\boldsymbol{p}_i} h_p(\boldsymbol{p}_i^t|\boldsymbol{p}_i^t) = \nabla_{\boldsymbol{p}_i} f_p(\boldsymbol{p}_i^t), \forall i$, and $\nabla_{\boldsymbol{s}} h_s(\boldsymbol{s}^t|\boldsymbol{s}^t) = \nabla_{\boldsymbol{s}} f_s(\boldsymbol{s}^t)$. Equal gradients automatically guarantee equal directional derivatives, proving the proposition. $\qquad\square$

Now that Algorithm 1 is found to be BSUM, Theorem 1 can be proved directly by the results of Theorem 2(b) in (Razaviyayn et al., 2013). But a few additional conditions presented in the following propositions need to be verified, both of which are proved thereafter.

Proposition 2 (Compact Sub-level Sets). *The sub-level set $\mathcal{X}^0 : \{(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P}) : f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P}) \leq f(\boldsymbol{\Theta}^0, \boldsymbol{s}^0, \boldsymbol{P}^0)\}$ of $f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P})$ given the update steps in Algorithm 1 is compact.*

Proposition 3 (Unique Global Minimizer of Auxiliary Functions). *Given Assumption 1, $\forall t \in \mathbb{R}_+$ and for all i and j, the auxiliary functions $h_\theta(\boldsymbol{\theta}_j|\boldsymbol{\theta}_j^t)$, $h_p(\boldsymbol{p}_i|\boldsymbol{p}_i^t)$, and $h_s(\boldsymbol{s}|\boldsymbol{s}^t)$ each have a unique global minimizer in $\mathbb{R}_+^K$.*

C.1. Proof of Proposition 2

We first introduce the definition of coercive functions through:

Definition 2 (Coercivity). *A function $f(\boldsymbol{X}_1, \ldots, \boldsymbol{X}_N) : V_1 \times \cdots \times V_N \to \mathbb{R}$, where V_i is a Euclidean vector space endorsed with the L_2 norm if its elements are vectors or with the Frobenius norm if the elements are matrices, is called coercive if $\forall i$, $\|\boldsymbol{X}_i\| \to \infty$ implies $f(\boldsymbol{X}_1, \ldots, \boldsymbol{X}_N) \to \infty$.*

In fact, continuous coercive functions always have compact sub-level sets, as shown by the following lemma:

Lemma 2. *If a continuous function $f(\boldsymbol{X}_1, \ldots, \boldsymbol{X}_N) : V_1 \times \cdots \times V_N \to \mathbb{R}$ is jointly coercive, then its sublevel set $\mathcal{X}^r := \{(\boldsymbol{X}_1, \ldots, \boldsymbol{X}_N) : f(\boldsymbol{X}_1, \ldots, \boldsymbol{X}_N) \leq r\}$ is compact for any $r \in \mathbb{R}$.*

Proof. Because f is continuous and the space $f(\boldsymbol{X}_1, \ldots, \boldsymbol{X}_n) \leq r$ is closed, $\mathcal{X}^r$ is closed. We prove the boundedness of $\mathcal{X}^r$ by contrapositive. We let $\mathcal{X}^r = \mathcal{X}_1^r \times \cdots \times \mathcal{X}_N^r$, where $\mathcal{X}_i^r \subseteq V_i$. If $\mathcal{X}^r$ is unbounded, then one of $\mathcal{X}_i^r$ is unbounded. If $\mathcal{X}_i^r$ is unbounded, we can find a sequence of $\{\boldsymbol{X}_k\}_{k=1}^\infty$ such that $\|\boldsymbol{X}_k\| \to \infty$ as $k \to \infty$, which implies f is not jointly coercive. Therefore, whenever f is jointly coercive, $\mathcal{X}^r = \mathcal{X}_1^r \times \cdots \times \mathcal{X}_N^r$ is bounded. Since Euclidean spaces admit the Heine-Borel property, $\mathcal{X}^r$ is compact. $\square$

Coming back to the proof of Proposition 2, it is easy to check that $f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P})$ is continuously differentiable. To prove that it is also coercive, we need to show that $f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P})$ tends to ∞ whenever $\|\boldsymbol{\Theta}\|_F$, $\|\boldsymbol{s}\|_2$, or $\|\boldsymbol{P}\|_F$ does so.

Beginning with the case for $\boldsymbol{\Theta}$, we have, by $f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P})$,

$$
\begin{aligned}
f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P}) &\geq \frac{\alpha_1}{2}\|\boldsymbol{\Delta} \odot \boldsymbol{\Theta} - \boldsymbol{\Theta}_0\|_F^2 + \frac{\alpha_2}{2}\|\boldsymbol{\Delta}^c \odot \boldsymbol{\Theta}\|_F^2 \\
&\geq \min\left(\frac{\alpha_1}{2}, \frac{\alpha_2}{2}\right)\left(\|\boldsymbol{\Delta} \odot \boldsymbol{\Theta} - \boldsymbol{\Theta}_0\|_F^2 + \|\boldsymbol{\Delta}^c \odot \boldsymbol{\Theta}\|_F^2\right) \\
&\geq \min\left(\frac{\alpha_1}{2}, \frac{\alpha_2}{2}\right)\left(\|\boldsymbol{\Delta} \odot \boldsymbol{\Theta}\|_F^2 - \|\boldsymbol{\Theta}_0\|_F^2 + \|\boldsymbol{\Delta}^c \odot \boldsymbol{\Theta}\|_F^2\right) \\
&= \min\left(\frac{\alpha_1}{2}, \frac{\alpha_2}{2}\right)\left(\|\boldsymbol{\Theta}\|_F^2 - \|\boldsymbol{\Theta}_0\|_F^2\right) \\
&= \min\left(\frac{\alpha_1}{2}, \frac{\alpha_2}{2}\right)\|\boldsymbol{\Theta}\|_F^2\left(1 - \frac{\|\boldsymbol{\Theta}_0\|_F^2}{\|\boldsymbol{\Theta}\|_F^2}\right). \tag{S.30}
\end{aligned}
$$

Since $\min\left(\frac{\alpha_1}{2}, \frac{\alpha_2}{2}\right) > 0$ and $\frac{\|\boldsymbol{\Theta}_0\|_F^2}{\|\boldsymbol{\Theta}\|_F^2} \leq \frac{mK_0 M_\Theta}{\|\boldsymbol{\Theta}\|_F^2} \to 0$ as $\|\boldsymbol{\Theta}\|_F \to \infty$ by Assumption 2, (S.30) then guarantees that $f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P}) \to \infty$ when $\|\boldsymbol{\Theta}\|_F \to \infty$.

In the case for $\boldsymbol{P}$, whenever $\|\boldsymbol{P}\|_F \to \infty$, there is a k such that $\boldsymbol{p}_k^\top$ satisfies $\|\boldsymbol{p}_k^\top\|_2 \to \infty$. Also, since $\boldsymbol{p}_k \in \mathbb{R}_+^n$, there exists a column indexed i_k as defined in Assumption 1(*ii*) such that $p_{ki_k} \to \infty$. Then,

$$
\begin{aligned}
f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P}) &\geq \frac{\beta}{2}\sum_{l=1}^K r_l^{-1}\|\boldsymbol{p}_l^\top - \boldsymbol{m}_l\|_2^2 \geq \frac{\beta}{2}r_k^{-1}\|\boldsymbol{p}_k^\top - \boldsymbol{m}_k\|_2^2 \\
&= \frac{\beta}{2}r_k^{-1}\sum_{i=1}^n (p_{ki} - \nu_k)^2 \geq \frac{\beta}{2}r_k^{-1}(p_{ki_k} - \nu_k)^2 \to \infty,
\end{aligned}
$$

as ν_k is assumed fixed and $\beta > 0$.

Lastly for $\boldsymbol{s}$, we can similarly argue that, because $\boldsymbol{s} \in \mathbb{R}_{++}^K$, $\|\boldsymbol{s}\|_2 \to \infty$ implies there is a k such that $s_k \to \infty$. From the steps leading to (S.21), we have

$$
\begin{aligned}
f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P}) &\geq \frac{1}{2mn}\|\boldsymbol{Y} - \boldsymbol{\Theta}\mathrm{diag}(\boldsymbol{s})\boldsymbol{P}\|_F^2 = \frac{1}{2mn}\|\boldsymbol{Y} - \sum_{k=1}^K s_k \boldsymbol{G}_k\|_F^2 \\
&\geq \frac{1}{2mn}\left(\|\sum_{l=1}^K s_l \boldsymbol{G}_l\|_F^2 - \|\boldsymbol{Y}\|_F^2\right) \\
&= \frac{1}{2mn}\left(s_k\|\boldsymbol{G}_k\|_F^2 - \|\boldsymbol{Y}\|_F^2\right) \to \infty.
\end{aligned}
$$

The last line is because $\|\boldsymbol{G}_k\| > 0$ due to Assumption 1(i) and Assumption 1(ii) and that $\|\boldsymbol{Y}\|_F \leq mnM_Y < \infty$ by Assumption 2. Combining the cases for $\boldsymbol{\Theta}$, $\boldsymbol{P}$, and $\boldsymbol{s}$ completes the proof that $f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P})$ is coercive.

Since $f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P})$ is continuous and coercive, the set $\mathcal{X}^0 : \{(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P}) : f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P}) \leq f(\boldsymbol{\Theta}^0, \boldsymbol{s}^0, \boldsymbol{P}^0)\}$ is compact according to Lemma 2. We have concluded the proof of Proposition 2.

C.2. Proof of Proposition 3

Since $h_\theta(\boldsymbol{\theta}_j|\boldsymbol{\theta}_j^t)$, $h_p(\boldsymbol{p}_i|\boldsymbol{p}_i^t)$, and $h_s(\boldsymbol{s}|\boldsymbol{s}^t)$ are quadratic functions, proving Proposition 3 is equivalent to showing their Hessians are positive definite. Also, we only focus on the positive coordinates of $\boldsymbol{\theta}_j^t$ and $\boldsymbol{p}_i^t$ since other coordinates will remain zero through updates, and the proposition is vacuously true. By (S.10), (S.18), and (S.22), these Hessians are $\boldsymbol{D}(\boldsymbol{\theta}_j^t)$, $\boldsymbol{D}(\boldsymbol{p}_i^t)$, and $\boldsymbol{D}(\boldsymbol{s}^t)$ respectively. Since they are all diagonal matrices, we want to show that their diagonal values are all positive.

For $\boldsymbol{D}(\boldsymbol{\theta}_j^t)$, we have $D(\boldsymbol{\theta}_j^t)_{kk} = \boldsymbol{d}(\boldsymbol{\theta}_j^t)_k = \frac{(\boldsymbol{A}\boldsymbol{\theta}_j^t)_k}{\theta_{jk}^t}$. The denominator of $D(\boldsymbol{\theta}_j^t)_{kk}$ is positive. Dropping the index j, its numerator is

$$
\begin{aligned}
\left\{\left(\mathrm{diag}(\boldsymbol{s})\boldsymbol{P}\boldsymbol{P}^\top\mathrm{diag}(\boldsymbol{s}) + mn(\alpha_1\boldsymbol{V} + \alpha_2\boldsymbol{V}^c)\right)\boldsymbol{\theta}_j^t\right\}_k &\geq \left\{mn(\alpha_1\boldsymbol{V} + \alpha_2\boldsymbol{V}^c)\boldsymbol{\theta}_j^t\right\}_k \\
&\geq mn\min(\alpha_1, \alpha_2)\theta_{jk}^t > 0,
\end{aligned}
$$

since $\boldsymbol{V} + \boldsymbol{V}^c = \boldsymbol{I}$. Hence, $D(\boldsymbol{\theta}_j^t)_{kk}$ is positive.

For $\boldsymbol{D}(\boldsymbol{p}_i^t)$, from (S.16) and (S.17), we also have $D(\boldsymbol{p}_i^t)_{kk} = \boldsymbol{d}(\boldsymbol{p}_i^t)_k = \frac{(\boldsymbol{C}\boldsymbol{p}_i^t)_k}{p_{ki}^t}$. Likewise, the denominator is positive, and the numerator after dropping the index i is

$$
\begin{aligned}
\left\{\left(\mathrm{diag}(\boldsymbol{s})\boldsymbol{\Theta}^\top\boldsymbol{\Theta}\mathrm{diag}(\boldsymbol{s}) + mn\beta\boldsymbol{\rho}^{-1}\right)\boldsymbol{p}_i^t\right\}_k &\geq \left(mn\beta\boldsymbol{\rho}^{-1}\boldsymbol{p}_i^t\right)_k \\
&= mn\beta r_k^{-1}p_{ki}^t > 0.
\end{aligned}
$$

Lastly, $D(\boldsymbol{s}^t)_{kk}$ is already shown to be positive from the conclusion of (S.27). This completes the proof of Proposition 3.

To finally prove Theorem 1, we combine the results of Propositions 1-3, and apply these conclusions to Theorem 2(b) in Razaviyayn et al. (2013). This way, we have shown that Algorithm 1 converges to a set of stationary points of $f(\boldsymbol{\Theta}, \boldsymbol{s}, \boldsymbol{P})$.

D.   Supplementary Information for The Simulated Pseudo-Bulk Data

D.1.   Data Generation

For $k = 1, 2, \ldots, 5$, s_k^* was generated independently from a χ^2 distribution with 5 degrees of freedom. For the rare class, $\boldsymbol{p}_i^* \overset{i.i.d.}{\sim} \mathrm{Dir}(2^5, 2^4, 2^3, 2^2, 2)$, where Dir represents the Dirichlet distribution. For the uniform class, we first drew $\boldsymbol{\alpha}_i$ from a Dirichlet distribution with $\alpha_{ik} \overset{i.i.d.}{\sim} U(0.3, 0.7), k = 1, \ldots, 5$, and then generated $\boldsymbol{p}_i^*$ from $\mathrm{Dir}(\boldsymbol{\alpha}_i)$. For the extra class, $\boldsymbol{p}_i^* \overset{i.i.d.}{\sim} \mathrm{dir}(2, 2^2, 2^3, 2^4, 2^5)$, in reverse of the rare class. To generate values in $\boldsymbol{\Theta}^*$, for each gene $j = 1, 2, \ldots, M$, a 5×1 vector of expression was generated by

$$\boldsymbol{\theta}_j^{*\top} \sim \Pi \left[\chi_{200}^2 \cdot \{(1 - \gamma)\mathrm{dir}(1, 1, \ldots, 1) + \gamma \mathrm{dir}(1, \ldots, 1, 5)\} \right],$$

where $\Pi(\cdot)$ is the sampling without replacement operator. $\Pi(\cdot)$ allowed all of CT1-CT5 have a number of gene signatures.

The random errors $\boldsymbol{\epsilon}$ were generated based on the principle of mean-variance dependency in gene expression data: the higher the mean of a gene's expression, the higher their variation (Anders and Huber, 2010). For each gene j, its error-free mean of bulk expression $\bar{y}_j$ was calculated. Then its associated error for each sample i was generated by $e_{ji} \overset{i.i.d.}{\sim} N(0, \sigma^2 \bar{y}_j^2)$, $i = 1, 2, \ldots, n,$.

D.2.   Method For Selecting Highly Cell-Type-Specific Genes

First, for each gene, the cell type for which it had the highest expression was picked as the target cell type. This gene then became a candidate for a marker of this target cell type. Second, for each candidate marker, the ratio of expression in its target cell type to those in the cell type with the second highest expression was calculated. Third, for each cell type, its candidate marker genes were ranked by the ratio calculated in the second step from highest to lowest. Finally, the top 100 candidate marker genes for each cell type were selected.

D.3.   Tuning Grid And Algorithm Parameters

Cross-validation (CV) was performed in each simulation to find the best values for α_1 and β according to Section 2.4. A tuning grid $\mathfrak{A}_1 = \mathfrak{B} = (10^{-2}, 10^{-1}, \cdots, 10^3, 10^4)$ was set up with the tolerance parameter chosen as $\delta = 10^{-5}$.

D.4.   Matching The Estimated Proportions to Cell Types For Reference-Free Methods

We utilized $\hat{\boldsymbol{\Theta}}$ and the partial signature matrix $\boldsymbol{\Theta}_{K_0}^*$. Starting from the first column of $\boldsymbol{\Theta}_{K_0}^*$, $\boldsymbol{\theta}_{K_{01}}^*$, we found the column of $\hat{\boldsymbol{\Theta}}$, say column k, that had the highest Pearson's correlation with $\boldsymbol{\theta}_{K_{01}}^*$. Then, the k-th row of $\hat{\boldsymbol{P}}$ was determined to be estimated proportions for CT1. We did this for the rest of the columns of $\boldsymbol{\Theta}_{K_0}^*$ and those of $\hat{\boldsymbol{\Theta}}$, until there was one last unmatched column in $\hat{\boldsymbol{\Theta}}$. That unmatched column, and its corresponding row in $\hat{\boldsymbol{P}}$, were then matched to CT5.

E. Supplementary Method And Results For The Deconvolution of Bulk PBMC Samples from The Human Influenza Vaccine Study

E.1. Data Processing

FASTQ files containing 50bp (base pairs) pair-end raw nucleotide sequence reads of transcripts from Illumina Hiseq 2000 sequencers were downloaded from the Sequence Read Archive (SRA) BioProject PRJNA271578. The raw reads were pre-processed and filtered where reads with length < 50bp, with $> 30\%$ bases with quality scores < 30, having an average quality score < 25 in any 10bp interval, or corresponding to special adapter sequences were removed using the software `fastp` (, Chen et al., 2018). Then, reads passing the filters were mapped to reference human genome hg38 GRCh38 Release 43 (downloaded from the GENCODE website) and quantified using `salmon` (, Patro et al., 2017). The output files were then processed using the R package `tximport` (, Soneson et al., 2015): only protein-coding genes with an official gene symbol were selected and their transcripts per million (TPM) values were gathered into a gene-by-sample matrix of gene expression. Additionally, genes with a total TPM of less than 10 across the four cell types were excluded from consideration.

Distinct marker genes for the four cell types were individually selected for each subject using sorted bulk profiles of individual cell types. The selection process identified the top 10 genes exhibiting the highest cell-type specificity for each cell type from each subject according the ratio of expression method outlined in Section 3.1. The union of the two sets of marker genes, totalling 73 genes, were used in the deconvolution.

E.2. Tuning Grid And Algorithm Parameters

A random seed of 1000 was set in the R programming environment. The tuning grid for α_1 was set as the interval from 2^{-4} to 2^0 with a step size of 0.2 in the power, and that for β was set as the interval from 2^0 to 2^4 also with a step size of 0.2 in the power. α_2 was fixed at 10^{-12}. The difference in magnitudes between α_1 and β in the tuning grids ensured proper regularization, for there was a scale difference between gene expression and the proportions. A 4-fold cross-validation was invoked to choose the optimal tuning parameters. The eventual selected tuning parameter values were within the boundaries of the grids. The tolerance parameter was set as $\delta = 10^{-4}$.

E.3. Fixing mRNA Amounts as One For All Cell Types using ARTdeConv Results in Biased Deconvolution Results

In Section 3.2, it was posited that bulk and gene signature matrices whose expression values were measured in TPM would lose the information on cell-type mRNA amounts, necessitating the inclusion of $\boldsymbol{s}$ described in the underlying model (1). To verify this claim, we re-ran the data analysis on the same data in TPM as in Section 3.2 using the same empirical estimates $\boldsymbol{M}$ and $\boldsymbol{\rho}$, the same tolerance parameter, and the same tuning grid for the hyperparaters, and the same random seed, but coerced $\text{diag}(\boldsymbol{s})$ to be the identity matrix (i.e. coerced the cell-type mRNA amounts to 1) throughout updates in Algorithm 1. We observed that the estimated proportions for PBMC cell types for both

Figure 1: Estimated vs. true proportions of PBMC cell types of two subjects on Day 0 using ARTdeConv with the cell-type mRNA amounts coerced to 1s.

samples on Day 0 deviates more from those measured by flow cytometry when compared to when diag(s) is not coerced, as shown in the following Supplementary Figure 1.

Table 1: Means and ranges of PBMC cell type proportions calculated from the reference values.

	T cells	B cells	NK cells	Monocytes	Other cells
Means	0.6275	0.0625	0.1075	0.15	0.0525
Ranges	0.275	0.055	0.145	0.1	0.2

F. Supplementary Method And Results For The Deconvolution of Bulk PBMC Samples from The COVID-19 Study

F.1. Study Design And Data Processing

The dataset generated from the study of Arunachalam et al. (2020) and downloaded using NCBI GEO accession number GSE152418 contained 34 subjects recruited from Atlanta, GA, USA. It included gene expression from human blood PBMC bulks samples collected from 17 healthy control subjects, one convalescent subject, and 16 subjects

diagnosed with COVID-19. Healthy controls were asymptomatic adults whose samples were collected before the widespread circulation of SARS-COV-2 virus in the community. Subjects with COVID-19 diagnosis were further classified into three levels of disease severity based on the based on the adaptation of the Sixth Revised Trial Version of the Novel Coronavirus Pneumonia Diagnosis and Treatment Guidance. Moderate cases were defined as respiratory symptoms with radiological findings of pneumonia. Severe cases were defined as requiring supplemental oxygen, and ICU-hospitalized cases were those in critical conditions who needed ICU care due to organ failures.

The gene symbols were annotated to 24,259 genes using the gconvert method of the `gProfileR` package in R software. The raw counts were then converted to counts per million (CPM) using those annotated genes. Among the 16 samples from COVID-19 subjects, two (labeled S155 and S179) contained abnormally high levels of *HBB* expression (namely, *HBB* was the most highly expressed gene in those two samples; results are not shown), which could only been found in red blood cells and suggested sample contamination. Therefore, these two samples were removed from the deconvolution procedures. This also led to S155, which also provided the study with single cell samples, being excluded from the comparison in Figure 10. The one convalescent sample was also excluded due to its unique designation.

The scRNA-seq data were generated by Arunachalam et al. (2020) using CITE-seq of >63,000 cell samples from five healthy controls and seven COVID-19 diagnosed patients. Of the seven patients, three were labeled as moderate cases, three severe cases, and one requiring ICU care. Notably, dendritic cells were enriched by the experimenters and mixed back into the samples for CITE-seq. All five healthy subjects and six of the seven (except S155, which were previously excluded) had measured bulk gene expression from independent PBMC samples as well for the analysis. To process the raw scRNA-seq data downloaded from GEO accession GSE155673, we first merged the single cell gene expression from all 12 subjects together in R package `Seurat` V4 (Hao et al., 2021). We then performed a quality control procedure for the cells, retaining those with a detected number of genes per cell between 200 and 5000 and filtering out cells with >15% mitochondrial counts. Genes with total expression less or equal to 2 across all cell samples were removed as well. We applied the global-scaling normalization method LogNormalize to normalize the feature expression measurements for each cell by the total expression, multiplied this by a scale factor of 10^6, and log-transformed the result. We employed the UMAP dimensional reduction technique to determine the cell type notation. The R package `SingleR` (Aran et al., 2019) was used for the final determination of cell types, using gene signature data from Monaco et al. (2019) as the cell type reference. The single-cell data input now has genes as rows and estimated cell type notations with subjects as columns. In the end, 41,146 cells and 26,531 genes were retained in the gene-by-cell expression matrix of all 12 samples.

We then separated the gene-by-cell matrix into two matrices, one containing only cells from healthy controls (23,531 cells) and the other only those from COVID-19 subjects (17,615 cells). Due to the limits of memory allocations to CIBERSORTx users, 11,765 cells from healthy controls were randomly sampled for each cell type proportionally to its original relative abundance. The cell types were re-grouped by merging all cell subsets of the four major cell types of interest together. The cellular gene expression from cells not belonging to any of the four cell types were discarded for gene signature generation. The processed sub-matrices were then separately sent to CIBERSORTx software's Create Signature Matrix module (, Newman et al., 2019) to obtain gene signature matrices for both

control and COVID-19 cell samples. The signature matrix for healthy controls contains 1,368 genes for the four cell types, while that for COVID-19 samples containts 1,388 genes. After matching the common genes from the bulk expression matrices and the signature matrices returned from CIBERSORTx, there were 1,280 genes for the deconvolution of healthy control samples and 1,297 genes for the deconvolution of COVID-19 patient samples.

F.2.   Tuning Grid And Algorithm Parameters

We adopted a tuning grid and a set of parameters for ARTdeConv to the settings in Supplementary Material Section E.2 for the analysis in Section 3.2. A random seed of 1000 was set in the R programming environment. For the deconvolution of healthy control samples, the tuning grid for α_1 was set as the interval from 2^{-3} to 2^1 with a step size of 0.2 in the power, and that for β was set as the interval from 2^2 to 2^6 also with a step size of 0.2 in the power. α_2 was fixed at 10^{-12}. For that of COVID-19 infected samples, the tuning grid for α_1 was set as the interval from 2^{-4} to 2^0 with a step size of 0.2 in the power, and that for β was also set as the interval from 2^2 to 2^6 also with a step size of 0.2 in the power. α_2 was again fixed at 10^{-12}. A 4-fold cross-validation was invoked in both deconvolution analyses to choose the optimal tuning parameters. The eventual selected tuning parameter values were within the boundaries of the grids in each case. The tolerance parameter was set as $\delta = 10^{-4}$.

Figure 2: Bar graphs of the deconvolved cell type proportions of all COVID-19 infected samples, grouped by their classified disease severity

Figure 3: Bar graphs of the deconvolved cell type proportions of all healthy control samples.

Figure 4: Expression of *CD3* genes, the translation of which produces T cells' CD3 markers, among healthy samples with exceptionally low expression observed in sample S066.

Figure 5: Expression of *CD14* and *FCGR3A* genes among healthy samples with high *CD14* expression observed in sample S066. *CD14* is responsible for the production of CD14 surface markers on classical/intermediate monocytes and *FCGR3A* responsible for that of CD16 surface markers on non-classical/intermediate monocytes.

References

Razaviyayn, M., Hong, M., & Luo, Z.-Q. (2013). *A unified convergence analysis of block successive minimization methods for nonsmooth optimization. SIAM Journal on Optimization*, 23(2), 1126–1153.

Lin, C.-J. (2007). *On the convergence of multiplicative update algorithms for nonnegative matrix factorization. IEEE Transactions on Neural Networks*, 18(6), 1589–1596.

Lee, D., & Seung, H. S. (2000). *Algorithms for non-negative matrix factorization.* In *Advances in neural information processing systems*, 13.

Kim, J., He, Y., & Park, H. (2014). *Algorithms for nonnegative matrix and tensor factorizations: A unified view based on block coordinate descent framework. Journal of Global Optimization*, 58(2), 285–319.

Lange, K. (2016). *MM optimization algorithms.*

Anders S. & Huber W (2010). *Differential expression analysis for sequence count data. Nature Precedings*, pages 1–1.

Chen S., Zhou Y., Chen Y., & Gu J. (2018). *fastp: an ultra-fast all-in-one fastq preprocessor. Bioinformatics*, 34(17):i884–i890.

Patro R., Duggal G., Love M. I., Irizarry R. A., & Kingsford C. (2017). *Salmon provides fast and bias-aware quantification of transcript expression. Nature methods*, 14(4):417–419.

Soneson C., Love M. I., & Robinson M. D. (2015). *Differential analyses for rna-seq: transcript-level estimates improve gene-level inferences. F1000Research*, 4.

Arunachalam, P. S., Wimmers F., Mok C. K. P., Perera R. A. P. M., Scott M., Hagan T., Sigal N., et al. (2020). *Systems Biological Assessment of Immunity to Mild versus Severe COVID-19 Infection in Humans. Science* 369 (6508): 1210–20.

Monaco, G., Lee, B., Xu, W., Mustafah, S., Hwang, Y. Y., Carré, C., Burdin, N., Visan, L., Ceccarelli, M., Poidinger, M., Zippelius, A., Pedro de Magalhaes, J., & Larbi, A. (2019). *RNA-Seq Signatures Normalized by mRNA Abundance Allow Absolute Deconvolution of Human Immune Cell Types. Cell Reports*, 26(6), 1627–1640.e7.

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W. M., 3rd, Zheng, S., Butler, A., Lee, M. J., Wilk, A. J., Darby, C., Zager, M., Hoffman, P., Stoeckius, M., Papalexi, E., Mimitou, E. P., Jain, J., Srivastava, A., Stuart, T., Fleming, L. M., Yeung, B., ... Satija, R. (2021). *Integrated analysis of multimodal single-cell data. Cell*, 184(13), 3573–3587.e29.

Aran, D., Looney, A. P., Liu, L., Wu, E., Fong, V., Hsu, A., Chak, S., Naikawadi, R. P., Wolters, P. J., Abate, A. R., Butte, A. J., & Bhattacharya, M. (2019). *Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. Nature Immunology*, 20(2), 163–172.

Newman, A. M., Steen, C. B., Liu, C. L., Gentles, A. J., Chaudhuri, A. A., Scherer, F., Khodadoust, M. S., Esfahani, M. S., Luca, B. A., Steiner, D., Diehn, M., & Alizadeh, A. A. (2019). *Determining cell type abundance and expression from bulk tissues with digital cytometry. Nature Biotechnology*, 37(7), 773–782.