1

Functional annotation hypothetical proteins: a world to be explored in drug development in Trypanosomatids

Raissa Santos de Lima[1], Ana Carolina Silva Bulla[1], Manuela Leal da Silva[1,2*]

[1]Programa de Pós-Graduação em Biologia Computacional e Sistemas, Instituto Oswaldo Cruz, Manguinhos, Rio de Janeiro, RJ, Brazil.

[2]Instituto de Biodiversidade e Sustentabilidade NUPEM, Universidade Federal do Rio de Janeiro, Macaé, RJ, Brazil.

*Corresponding author:

E-mail: manuela@macae.ufrj.br

Author contributions: Conceived and designed the experiments: RSLR MLS. Performed the experiments: RSLR ACSB. Analyzed the data: RSLR ACSB. Wrote the paper: RSLR ACSB MLS.

Abstract (300 words)

Hypothetical proteins can provide an alternative pathway for finding potential targets in

the development of new drugs due to the fact that many Neglected tropical diseases are

caused by Trypanosomatids (Chagas Disease, Leishmaniasis, and Human African

Trypanosomiasis). In this work, we focus on applying functional prediction methods

based on both sequence and structure to analyze the hypothetical proteins of the

pathogenic agents that cause these diseases: *T. cruzi* (Tcr), *T. brucei brucei* (Tbr), *T.*

brucei gambiense (Tbg), *L. infantum* (Lif), *L. donovani* (Ldo), and *L. braziliensis* (Lbz).

By consulting databases and servers, we have predicted functional domains for twenty-

six proteins in Tcr, thirteen in Tbr, fifteen in Tbg, ten in Lif, and one in both Ldo and

Lbz. With the goal of developing multi-target therapies, we grouped the domains

according to how they are shared among the organisms and investigated those that are

shared among more species. By examining the existing literature using specific search

strategies, we described what has already been reported for these domains and also

analyzed protein structures and sequences, describing mutations among the species and

potential drug sites. The published works have unveiled that some of these domains are

non-essential for trypanosomatids, like the TRX domain, while others demand further

investigation due to a lack of information about metabolic processes (UFC1, Ufm1,

ACBP, AAA 18, and Fe-S). Although, we have identified three noteworthy domains

that hold promise as targets: TPR, which plays a crucial role in the ciliogenesis process;

Nuc deoxyrib tr, essential in purine recycling and recovery mechanisms; and MIX,

important for protein targeting and the assembly of complexes such as COX. These

three domains are promising targets for drug development due to their conservation,

their potential to affect multiple species and their exclusivity.

Introduction

Trypanosoma parasite species can infect several animals causing neglected

diseases such as Chagas disease (CD) (*Trypanosoma cruzi*), Human African

Trypanosomiasis (HAT) (*Trypanosoma brucei*) and Leishmaniasis (genus Leishmania)

[1,2]. From an epidemiological standpoint, these diseases' incidence rates are

alarmingly high, with millions of cases reported annually [3–5]. Furthermore, HAT was

close to eradication in the 60s but the disease has since experienced a resurgence in

reason of political problems, and a decline in control programs [5,6]. Besides that, many

of these diseases have a wide geographical distribution due to infected individuals

immigrating to non-endemic countries, and leishmaniasis in particular has three

endemic foci: India, East Africa, and Brazil [3,7,8].

Aside from the challenge of dissemination, the treatment has several problems,

mainly regarding efficacy and therapeutic adherence due to the high toxicity of the

drugs used [9–11]. Furthermore, there are already cases of resistance that demonstrate

the promiscuity not only of drugs but also of current targets for diseases caused by

trypanosomatids [12,13]. As a result, there is a pressing need to take the initiative to

develop new drugs, and one promising approach is the drug repositioning strategy. This

process is highly cost-effective since it reduces drug development time, optimizing pre-

clinical trials by utilizing drugs with known possible adverse effects [14,15]. Thus,

efforts are required to identify new molecular targets for new drugs and their implementation as effective treatments.

Drugs can have different molecular targets, one of the most common being proteins, and with the increasing popularity of sequencing techniques, there has been a rise in the use of bioinformatics tools, including those for protein analysis [16]. Despite the availability of predicted proteins in genomes, there is a limited amount of functional details data for these molecules. The reason for that is the sheer volume of data that experimental functional characterization produces, which is difficult to manage. So the functional prediction approach comes in as a solution, and it has been constantly improved [17].

There are two main approaches to protein function prediction and the first one is the SEquence-Based Functional Prediction (SEBFP), which assumes that the sequence contains a conserved "functional signature" that can be identified even in a small part of the sequence [18]. Consequently, if two proteins have this conserved "signature" it is possible to infer the protein function [19]. The second approach is Structure-Based Functional Prediction (STBFP) which has two premises, the first one is that all the essential information about a protein's structure is contained in its amino acid sequence [20]. The second premise is that proteins have a specific native folding state or a set of native states, and this structure is closely related to the catalytic protein function [21].

The primary aim of this study is to employ a hybrid approach, merging SEBFP and STBFP strategies, to predict the functional domains of hypothetical proteins (HP) as depicted in Fig 1. Our goal is to achieve more robust results compared to conventional

linear methodologies and to identify new targets for drug development that are

applicable across multiple neglected diseases caused by these parasites.

Fig 1. Scheme of the functional prediction methodology. Through the utilization of a

diverse range of computational tools and databases, we conducted an extensive analysis

of both sequence and structural data to identify functional domains within the

hypothetical proteins (HP). As illustrated in the sequence and structure, an exemplary

functional domain is highlighted in orange, showcasing its conservation across various

protein data levels. For the 3D protein representation, the PDBids were used: 3UM8,

4TU5, 6WEP, and 3Q1H.

To demonstrate our approach, we applied SEBFP and STBFP using various

databases, identifying a total of sixty-six protein domains for the organisms *T. cruzi*

(Tcr), *T. brucei brucei* (Tbr), *T. brucei gambiense* (Tbg), *L. infantum* (Lif), *L. donovani*

(Ldo), and *L. braziliensis* (Lbz). Notably, nine of these domains were shared by four

species (Tcr, Tbr, Tbg, and Lif), which we have provided detailed descriptions for. With

a focus on identifying promising multi-targets, we conducted sequence and structural

analyses in our pursuit of potential druggable sites for the three domains (TPR, Nuc

deoxyrib tr, and MIX) that hold promising functions.

Materials and methods

Proteomes selection

According to the disease, we selected the proteomes of the following species

Trypanosoma cruzi for CD, *Trypanosoma brucei* for HAT and *Leishmania* spp for

Leishmaniasis. To determine the strain among several deposited in the National Center

for Biotechnology Information (NCBI) databases [22], we first selected the most widely

reported strain based on literature data. Additionally, we considered the strains with the

highest assembly level available in the NCBI. After the selection process, using a script

that analyzes the identifiers, we separate the proteins annotated as HP from the others.

Large-scale comparative modeling

The multiFASTA file containing only the HP sequences was submitted to the

MHOLline web server for structural modeling [23]. MHOLline is a biological workflow

capable of performing three-dimensional modeling for different proteins at the same

time. The structure prediction process starts with searching for templates by aligning

protein sequences with structures in the Protein Data Bank (PDB) using the Basic Local

Alignment Search Tool (BLAST) [24]. Monomeric 3D models are generated with

MODELLER [25] for only results with e-value $\leq$ 10e-5, identity $\geq$ 0.25, and Length

Variation Index (LVI) $\leq$ 0. Subsequently, MHOLline classifies these models into five

groups among Very High and Very Low using the module called Filters, which is based

on the identity and LVI values. In this study, we selected those that were classified

between Very High and Good (Very High: ID $\geq$ 75% and LVI $\leq$ 0.1%; High: ID $\geq$ 50%

and <75% and LVI $\leq$ 0.1%; Good: ID $\geq$ 35% and < 50% and LVI $\leq$ 0.3%). Hence, only

the protein sequences that met the evaluation criteria were deemed suitable for the

functional prediction stage.

Functional prediction and description

As previously mentioned, we employed a combined SEBFP and STBFP approach to predict the selected HP sequence's functional domains. In total, nine databases and six web servers were utilized, as shown in Fig 2. To ensure the reliability of the functional prediction, we defined that the same domain must be predicted in at least eight of the SEBFP tools and two of the STBFP for functional inference. Similar strategies have been previously adopted by Shahbaaz, M., Hassan, M. I., & Ahmad, F. (2013) and Mazandu, G. K., & Mulder (2008).

Fig 2. Workflow containing databases and servers used in SEBFP and STBFP. Databases are represented as cylinders and the web servers as rectangles.

Consulting the literature, the functional domains were detailed, particularly through the use of bibliographic databases such as Web of Science, Google Scholar, The Lens, and PubMed. The search strategy employed was using keywords such as the functional domain acronym, disease's name, and the parasite species, for example, "TRX AND Chagas Disease AND Trypanosoma cruzi". Additionally, to find targets that are crucial for the parasite's metabolic pathways, we consulted the Kyoto Encyclopedia of Genes and Genom3es (KEGG) database [26] using the EC number or the functional domains acronym in the.

Sequence analysis

Sequence analysis began using tools that contain details about domains and families, which included: Pfam [27], Simple Modular Architecture Research Tool (SMART) [28], PROSITE [29], Conserved Domains Database (CDD) [30], Search Tool

for Retrieval of Interacting Genes/Proteins (STRING) [31] and the CATH [32] and

HMMER [33] web servers. These tools were used with default settings and allow the

identification of homologous sequences by aligning them with information in various

databases. This type of approach, the homologs identification, or evolutionary

relationships, is widely used for the structural and functional annotation of proteins [32–

34].

The proper characterization of a protein involves determining not only its

function but also its cellular localization. This information is crucial for understanding

its role in cellular processes and for the development of targeted therapies. To determine

the subcellular localization we used the Psort (Prediction of Protein Sorting Signals and

Localization Sites in Amino Acid Sequences) web server [35], more specifically WoLF

PSORT with default parameters. This tool predicts protein sorting signals and

localization sites based on amino acid sequences. Lastly, we consulted UniProtKB,

where is possible to search for similar sequences with predicted functions using the

integrated BLAST tool and we consulted the curated Swiss-Prot sequence database with

default parameters [36,37].

To further explore sequence similarity with other sequences with other already

known sequences deposited in data banks, we used the Basic Local Alignment Search

Tool (BLAST) tool with the Non-Redundant (NR) [24] database and the TriTrypDB

database [38], both with the default search parameters. The TriTrypDB database

contains sequence data for the Trypanosomatidae family species and sources its

information from banks such as the DNA Data Bank of Japan (DDBJ) and the European

Molecular Biology Laboratory (EMBL) [39].

Among species, the domain conservation, as well as mutations were analyzed

using the UGENE alignment tool, specifically with the MUtiple Sequence Comparison

by Log-Expectation (MUSCLE) algorithm [40]. MUSCLE realigns the sequences

repeatedly, considering a guiding phylogenetic tree to generate the distance matrix [40–

42]. Given that we were analyzing organisms from the same family, considering the

phylogenetic relationships when assembling the matrix was crucial for the alignment

quality [43].

Finally, in addition to the aforementioned analyses, we identified the signal

peptide and transmembrane regions. First, for signal peptide detection, we used the

SignalP web server, which can predict not only the signal peptide's presence but also

their cleavage sites [44]. For the transmembrane regions encounter, the DeepTMHMM

web server was used, which scores the protein region probability being inside, in the

transmembrane region, or outside the membrane [45]. We used both servers with the

default search parameters.

Structure analysis

STBFP analyses started by utilizing the structures generated by MHOLline to

search the SCOP server. This server classifies the protein based on its structure and

provides a structural relationship description among known proteins [46]. Besides that,

we used the VAST server, which compares the spatial coordinates of an input 3D

structure with other structures deposited in the Molecular Modeling Database (MMDB)

of NCBI [47]. Furthermore, VAST can identify distant homologs that may not be

recognized through a simple sequential comparison [48].

Another server utilized was the Flexible structure AlignmentT by Chaining

Aligned fragment pairs allowing Twists (FATCAT), which searches for similar

structures in databases such as PDB and SCOPe through a structure alignment

algorithm [49]. Lastly, the Distance matrix ALIgnment server (Dali) was used also for

comparison structures in the PDB database but it utilizes three different ways: the

heuristic search (all structures in PDB), the exhaustive search (a PDB subset), and an

AF-DB hierarchical lookup (AlphaFold subset) [50]. Both FATCAT, Dali, and VAST

servers perform alignment based on calculating of Root Mean Square deviation

(RMSD) using α carbons.

Regarding the RMSD, the Pymol (version 1.9) program [51] was utilized to

calculate it for the models generated by MHOLline with the templates used. In addition,

we examined the conservation of amino acid residues that compose the functional

domain, as well as the presence of mutations among the species in the structures. To

evaluate regions without clearly defined secondary structures, we referred to the

AlphaFold structure database [52], which provides information on models generated by

artificial intelligence [53], including details on species. Moreover, we utilized the

Quick2D web server, a tool that includes nine predictors of secondary structures, even

regions of disorder, to facilitate reaching a consensus on the structure in the region of

interest [54].

Modeling homo-oligomeric structures and validation and binding site identification

The final structures were remodeled considering the oligomeric state through AlphaFold2 on ColabFold [55] pipeline (num_relax: 5, template mode: none, msa_mode: MMSeq2 (uniRef□+□environmental), num_recycle: 3), relaxing the structure with an amber force field, considering the sequences in table S4. To verify the structural quality of the built models, the models were submitted to MolProbity [56] and ProSA – web [57]. To access the protein druggability, the structures were submitted to DogSiteScorer [58] server for prediction of potential pockets on the proteins (score between 0 and 1).

Results and Discussion

Selection and large-scale 3D modeling

Using the NCBI database, we conducted a search of organism's proteomes, specifically: *Trypanosoma cruzi* (Tcr) for CD; *Trypanosoma brucei brucei* (Tbr), and *Trypanosoma brucei gambiensi* (Tbg) for HAT; *Leishmania infantum* (Lif), *Leishmania donovani* (Ldo) *Leishmania braziliensis* (Lbz) for Leishmaniasis. Lif and Ldo species were selected due to their association with the most severe clinical form of the disease, visceral leishmaniasis (VL), which is more prevalent in Brazil [3,7].

The organisms and their respective strains were selected considering literature data and the proteomes with the highest assembly level deposited in the NCBI database (Table 1). Once the strains were chosen, the hypothetical proteins (HP) were separated

from the others and submitted to the MHOLline server. This resulted in the generation

of 3D models for 2928 proteins, ouch of which 111 models qualified for the functional

prediction stage due to their classification between Very High and Good (Very High: ID

$\geq$ 75% and LVI $\leq$ 0.1%; High: ID $\geq$ 50% and <75% and LVI $\leq$ 0.1%; Good: ID $\geq$ 35%

and <50% and LVI $\leq$ 0.3%). Lastly, we consulted sequential and structural databases to

infer functional domains for these proteins.

Table 1. Correlation of species, numbers of proteins, and 3D models

Species	Strains	NCBI Code	TP	HP	Gen 3D	Models selected
T. cruzi	CLBrener	GCA_000 209065.1	19.608	11.171	1.421	40
T. brucei brucei	TREU927	GCA_000 002445.1	8.758	5.711	658	20
T. brucei gambiensi	DAL972	GCA_000 210295.1	9.822	6.611	661	23
L. infantum	Jpcm5	GCA_000 002875.2	8.15	5.245	648	21
L. donovani	LdCl	GCA_003 719575.1	8.632	3.527	90	3
L. braziliensis	MHOM	GCA_900 537975.1	8.244	3.373	98	4

[a]TP - Total of proteins

[b]HP - Hypothetical protein

[c]Gen 3D - 3D models generated

Annotation of HPs

The nomenclature used to describe the results was based on the user in the

following databases: InterPro [59], Pfam [27], SMART [28], PROSITE [29], and CDD

[60] (S1 Table). To predict the functional domains, we established the following

criterion: the same function must be predicted in at least eight SEBFP tools and two

STBFP tools. Using sequence and structure analysis tools, we inferred 26 functional

domains for Tcr, 13 for Tbr, 16 for Tbg, 10 for, Lif, 1 for Ldo, and 1 for Lbz, totaling

67 proteins from the initial 111 (S2 Table).

By analyzing SEBFP and STBFP results together, it was observed that some of

the annotated domains were repeated within the same species and also interspecies (Fig

3). Specifically, we found that nine predicted domains were shared among four species

of trypanosomatids, namely Tcr, Tbr, Tbg, and Lif. Therefore, we narrowed our focus to

these shared domains, as our goal is to identify multi-targets for trypanosomatids.

Fig 3. Venn diagram illustrating the shared functional domains among

trypanosomatid species. Each circle in the diagram represents a distinct

trypanosomatid species, and the predicted functional domains are denoted by bold

abbreviations. Our analysis primarily concentrates on the centrally located shared

domains, which have the highest number of species located in the middle of the Venn

plot.

Multi-target functional domains identified

Ubiquitination: UFC1 and Ufm1

In our research, we employed a combination of SEBFP and STBFP

methodologies to uncover various functions. However, as we delved into the specifics

of each function, it became evident that there were significant gaps in our

understanding. For instance, we identified the UFC1 and Ufm1 domains, both of which

play roles in the ubiquitination process—a post-translational protein modification

system utilizing multiple ubiquitin molecules to regulate cellular functions [61].

The UFC1 domain is a member of the E2-type proteins (conjugators) in the

ubiquitination cascade. It plays a vital role in receiving a specific type of ubiquitin (Ub),

known as Ubiquitin fold modifier 1 (Ufm1), from Uba5 (E1-type activators) through a

thiolester bond mediated by a cysteine [62]. This process is crucial for central nervous

system (CNS) development in humans, and bi-allelic mutations in UFC1 have been

linked to compromised neurological phenotypes, leading to severe encephalopathy and

progressive microencephaly cases [63]. Mutations like Arg23Gln can affect binding to

E1 or E3 proteins (ligands), while Thr106Ile impairs the UFC1-Ufm1 complex

formation [63].

In trypanosomatids, specifically *T. brucei*, research has shown that the

glycoproteins on their surface rely on the ubiquitination process, particularly Invariant

Surface Glycoproteins (ISG). After undergoing ubiquitination, these proteins are

internalized and degraded, promoting cell surface renewal. This implies that

ubiquitination serves as a general mechanism for regulating surface proteins with

transmembrane domains [64]. However, research also reveals variations in the

machinery involved in this process, such as the absence of orthologs for E3 proteins in

the *T. brucei* genome, suggesting alternative mechanisms or factors may be in play in

these eukaryotes [65].

Regarding the Ufm1 domain, a study conducted by Yoo et al. in 2014 found that

certain ubiquitin ligases are overexpressed in tumors, promoting cell proliferation. This

study also identified interactions between estrogen receptors (ERα) and Activating

signal cointegrator 1 (ASC1) molecules through the Ufm1 domain, leading to the

activation of ERα target genes associated with excessive cell proliferation and breast

cancer [66]. The significance of ubiquitylation in the human body underscores the need

for further research into UFC1 and Ufm1, especially in the development of targeted

inhibitors for specific members of the E3 family [67]. Meanwhile, the proteasome

pathway, which is essential for the degradation of cytoskeletal proteins in *T. cruzi*,

shows promise as a drug development target. Studies suggest that inhibiting the

proteasome can disrupt the evolutionary cycle of *T. cruzi*, making it vital to explore this

pathway for new drug targets [68–70]. Hence, it is crucial to persist in the investigation

of this pathway to unearth novel drug targets.

Electron transfer: Fe-S cluster

The Fe-S domain assumes a pivotal role in modulating transcriptional and

translational processes within genes [71]. Originating during the prebiotic era on Earth,

this domain finds its basis in the abundance of iron and sulfur, coupled with their

structural versatility and chemical reactivity, facilitating the formation of Fe-S clusters

(iron and sulfur). These clusters function as "molecular switches," capable of

influencing gene regulation at multiple levels [72]. Their intrinsic chemical properties

make them active participants in various biological processes, encompassing

photosynthesis, DNA replication and repair, respiration, nitrogen fixation, and more.

Consequently, proteins containing Fe-S clusters are ubiquitous across all organisms

[73,74].

In humans, the cytosolic Fe-S cluster assembly (Cia) targeting complex is

constituted by Cia1, MMS19, Cia2A, and Cia2B (also known as Fam96b) [75].

Notably, Cia2A and Cia2B hold significance in iron homeostasis and participate in the

maturation of the iron regulatory protein (IRP1) [76]. Fam96a, also recognized as

Cia2A in mammals, has demonstrated its role in reducing gastrointestinal stromal

tumors, where its interaction with apoptotic peptidase activation factor 1 heightened

tumor sensitivity to apoptosis processes [77,78]. Additionally, Fam96a depletion has

been linked to organ damage protection and increased survival rates for sepsis patients

by reducing the secretion of pro-inflammatory cytokines by macrophages [79].

In trypanosomatids, like *T. brucei*, the presence of the Cia complex has been

acknowledged; however, the precise components for Cia in *T. brucei* remain uncertain,

and it's been suggested that these components may have influenced the divergence of

the Kinetoplastea class from other eukaryotes [80]. The ISC machinery, responsible for

the maturation of cytosolic and nuclear Fe-S proteins, has been proposed to directly

impact the formation of flagella in these parasites, particularly under conditions

affecting glycolysis and mitochondrial function [81,82]. *Leishmania* spp. also possess

Cia components with variations in their mechanisms. Given the conservation and

essential role of the Cia pathway in Fe-S cluster assembly within trypanosomatids, it

presents a promising source of potential pharmaceutical targets, with Fam96a being one

of the identified candidates in *T. brucei* [80] and detected via STBFP analysis (S2

Table).

Lipid metabolism: ACBP

ACBPs, in general, function by binding to acyl-CoAs, thereby preventing their

hydrolysis by hydrolases [83]. This action facilitates the efficient transport of acyl-

CoAs to cellular acylation machinery. In humans, ACBPs, also known as ACBDs, have

been linked to various cellular processes, including lipid metabolism and cell signaling

[84].

In trypanosomatids, ACBPs are believed to protect against inhibition by acyl-

CoA esters of acetyl-CoA carboxylase and long-chain mitochondrial adenine nucleotide

translocase [85]. These proteins also play a role in the transport and donation of acyl-

CoAs for β-oxidation in mitochondria and the synthesis of glycerolipids in microsomes

[86].

Experimental studies on trypanosomatids, particularly *T. brucei*, have revealed

the significant role of ACBP in the biosynthesis of glycosylphosphatidylinositol (GPI).

Specifically, ACBP donates myristoyl-CoA for fatty acid remodeling reactions, causing

a conformational change in fatty acids [87]. This alteration allows them to exit the

endoplasmic reticulum and anchor to Variant Surface Glycoproteins (VSG), which is

crucial for the formation of the VSGs [88]. Once anchored, VSGs undergo further

processing in the Golgi complex, including the addition of galactoses, before being

transported to the plasma membrane [85].

Furthermore, trypanosomatids rely on GPI anchors for various essential

functions, such as evading the host complement system in *T. brucei*, facilitating host

cell interaction and sialic acid uptake in *T. cruzi*, and protecting against extracellular

hydrolases while interacting with the insect vector's midgut in *Leishmani*a spp. These

membrane components are pivotal for various critical processes, highlighting the

importance of understanding the functionality of domains like ACBP in these parasites

[89].

Energy maintenance: AAA18

AAA 18 domain, also known as adenylate kinase (AK), plays a role in

regulating cellular energy management by maintaining adenine nucleotide homeostasis.

AAA 18 catalyzes the reversible conversion of ATP and AMP molecules into two ADP

molecules, contributing to the balance of adenine nucleotides and cellular energy

charge, thereby supporting optimal cellular functions and metabolism [84].

Studies have revealed the presence of multiple isoforms of AAA 18 with diverse

subcellular localizations and functions. In *T. cruzi*, a cytosolic isoform of AAA 18 has

been identified, and it plays a crucial role in the energy metabolism of the parasite,

particularly during the growth of the epimastigote form. This suggests a potential

correlation between the activity of AAA 18 and the energy demands of the parasite.

Interestingly, the expression of the cytosolic AAA 18 isoform decreases when another

phosphotransferase, arginine kinase, is overexpressed, indicating a potential functional

connection between these two enzymes and a compensatory mechanism for maintaining

the parasite's energy homeostasis [90].

Furthermore, the number of AAA 18 isoforms varies among different species. In

the case of *T. brucei*, Ginger et al. (2005) conducted a study that revealed seven

isoforms of AAA 18 in this organism. The existence of multiple isoforms suggests a

form of metabolic adaptation, leading to increased complexity in energy homeostasis

reactions. These isoforms are expressed in both procyclic trypanosomes found in the

midgut of the tsetse fly and metacyclic trypanosomes located in the bloodstream,

making them particularly relevant in the context of parasite infections [90].

Additionally, in mammals, several AAA 18 isoforms have been associated with cancer,

such as the overexpression of AK2 in lung cancer patients, which is linked to the
reprogramming of energy metabolism—an essential aspect of cancer progression [91].

Despite our understanding of the functions of ACBP and AAA 18 domains,
many mechanisms related to these domains remain enigmatic. One intriguing area of
research is the relationship between ACBPs and peroxisomes, given the critical role of
peroxisomes in various neurological disorders like Alzheimer's, autism, and
amyotrophic lateral sclerosis [92]. With these proteins playing a significant role in
humans, concerns arise when targeting them with drugs due to potential systemic
consequences. Therefore, further research is essential to enhance our understanding of
the selectivity and implications of targeting ACBPs and AAA 18 in the context of
human health.

Purine metabolism: TRX

TRX domain is a pivotal component of the purine metabolism pathway. Purines
play a crucial role in the synthesis of signaling molecules like cyclic adenosine 3,5-
monophosphate (cAMP) and are integral to the cellular energy system, including
adenosine triphosphate (ATP). Furthermore, they participate in the production of RNA
and DNA, working in tandem with pyrimidines [93,94]. However, experiments
conducted on *T. brucei* cultures have raised questions about the indispensability of this
protein due to the presence of a second TRX enzyme, TRX2 [95].

Significant differences in conservation between TRX2 and TRX1 exist among

various trypanosomatid species. TRX1 exhibits a lower degree of conservation, with

approximately 32% identity between *T. cruzi* and *T. brucei* [95]. However, our

alignment results indicate a higher identity of 50% between these species, with the

lowest being 44% between Tcr and Lif. This supports the hypothesis that the domain

we've identified is TRX1. Studies have revealed that the absence of TRX2 leads to

compromised parasite proliferation in *T. brucei*, while the deficiency of TRX1 doesn't

impact parasite proliferation [95]. Consequently, it appears that TRX1 may not be a

promising target for drug development.

In the upcoming sections, we will explore several protein domains that we have

identified as promising candidates for multi-target studies in the development of novel

drugs. These domains play critical roles in Trypanosomatids, making them valuable

subjects for therapeutic investigation. To gain a comprehensive understanding of these

domains, we conducted extensive analyses, including the generation of their oligomeric

structures using AlphaFold2, rigorous evaluation of quality parameters, and the

prediction of druggable sites. Detailed sequences of these domains are available in the

S3 table, which will aid in further elucidating their potential as therapeutic targets.

422   Cyclogenesis: IFT - TPR

423   The tetra-trico peptide repeat domain (TPR) is a highly conserved structure

424   comprised of multiple repeats of 34 hydrophobic amino acids, found in various

425   organisms [96]. Its involvement in critical processes, such as transcriptional inhibition,

426   stress response, and mitochondrial protein transport, positions it as a promising target

427   for drug development [97–99].

428    Our evaluation of the TPR's conservation across trypanosomatid species

429  revealed high levels of identity, ranging from 73% (between Lif and Tbg/Tbr) to 98%

430  (between Tbg and Tcr). Furthermore, the human isoforms of IFT70 (A and B) showed

431  an identity range of 51% to 54% (Fig 4A). While the isoforms share a considerable

432  degree of similarity, it's essential to emphasize the significance of the IFT train's

433  involvement in various processes, which we will describe in detail below.

434  **Fig 4. TPR - Sequential and structural analysis.** (A) Shows the alignment of

435  trypanosomatid and human (Hsa) sequences, accompanied by a graphical summary

436  revealing the percentage of identity. Notably, the proline-rich region is highlighted,

437  demonstrating its conservation across species. The alignment matrix below employs

438  color-coding within the UGENE software, indicating dissimilarity percentages: white

439  (0%), red (10%), orange (25%), green (50%), blue (70%), and gray (90%). (B)

440  Illustrates the alignment between the 3D model generated by MHOLline for the Lif

441  sequence (dark blue) and the 4UZY template (gray). The TPR domain is depicted in

442  pink, while the IFT52 protein, resolved with the PDBid: 4UZY structure, is represented

443  in orange. (C) Results for the druggability prediction where in a blue cartoon is IFT70

444  of *L. infantum* (Lif) and the druggability score and corresponding surface area values

445  are color-coded: green (0.84, 888.13 Å²), orange (0.81, 3311.56 Å²), purple (0.78,

446  878.57 Å²), pink (0.76, 746.52 Å²), and blue (0.76, 605.84 Å²).

447    The TPR domain's versatility in cellular location, being found in various

448  compartments such as the nucleus, cytoplasm, and mitochondria [96], was confirmed in

449  our Psort prediction for trypanosomatid species. Tcr, Tbr, and Lif were predicted to be

450  mitochondrial, while Tbg was cytoplasmic. The TPR domain's role in mediating

protein-protein interactions and assembly of protein complexes was underscored by our

Structure-Based Functional Prediction (STBFP) results, which identified the IFT70

protein (PDBid: 4UZY) [100]. IFT70 is present in the IFT train which is a crucial

component of the intraflagellar transport protein complex responsible for cyclogenesis,

an evolutionarily conserved transport process involving the bidirectional movement of

particles within cilia [101].

As noted in the structural analyses, the 4UZY template corresponds to the IFT70

protein complexed with part of IFT52 from the organism *Chlamydomonas reinhardtii*

(PDBid: 4UZY). The interaction between IFT70 and the IFT52 protein occurs through a

proline-rich region [102], and we observed that most of these regions are conserved,

even in humans (Fig 4A). Therefore, the tertiary and primary structures exhibited

significant similarity to a functionally characterized protein (Fig 4 A and B), reinforcing

the connection between structure and functionality in proteins.

To enhance the accuracy of our structure prediction, we employed AlphaFold2

to generate new models for each species, taking into account the oligomeric states of the

proteins. The results for IFT70 can be found in S1 Fig and to ensure the quality of these

models, conducted evaluations of the models using MolProbity [56] (Clash score,

MolProbity score, and Ramachandran favored regions), as well as ProSa [57] (as shown

in S4 Table). Notably, the best IFT70 model was identified for the *L. infantum* (Lif)

organism (accessible via this ZENODO link: https://doi.org/10.5281/zenodo.10072065).

Consequently, we employed this model to predict potential binding sites using

DoGSiteScorer [58], which assigns druggability scores within the range of 0 to 1.

Higher scores indicate more favorable druggable pockets, guiding our search for

potential drug-binding sites. The results for the Lif model indicate the presence of five

promising regions (Fig 4 C), with the majority of them located within the IFT52 binding

site, reaffirming the crucial role of this region in the functionality of IFT70.

Furthermore, the significance of the IFT52 protein in stabilizing the IFT-B1

complex and its direct involvement in cilia formation has been well-documented in

studies across various organisms [103,104]. For instance, the study by HUET et al. in

2019 demonstrated that other proteins within the IFT train, such as IFT172 (IFT-B2),

were pivotal for proper cyclogenesis and flagellar organization in *T. brucei*. These

findings underscore the indispensable role of the IFT train in cyclogenesis and

emphasize the critical contributions of individual proteins, like IFT52, in maintaining

the complex's stability and functionality.

Despite the significant progress in understanding the IFT train, questions remain

regarding the interactions and subunits involved in complex stabilization. However, the

potential of the IFT70 protein and other IFT train proteins as targets for drug

development in trypanosomatids is promising, given their pivotal role in critical cellular

processes and preliminary in vitro study results. Further investigations may reveal new

drug development opportunities.

Nucleosides transferases: Nuc deoxyrib tr

Nucleoside 2'-deoxyribosyltransferases (NDTs), also known as Nuc deoxyrib tr

according to domain databases' nomenclature, constitute a protein family that mediates

the transfer of the 2'-deoxyribosyl group between 2'-deoxyribonucleosides (donor) and

nucleobases (acceptor) [105]. These enzymes belong to the class of Nucleoside

Phosphorylases (NPs), which are further divided into ribo- and 2'-deoxyribonucleosides.

The class encompasses various subclasses, including type I NDTs (PDT), responsible

for deoxyribotransfer between purines only, and type II NDTs (NDT), which enable

transfer between purines and/or pyrimidines [106]. NDTs have been identified in

various species, including parasitic ones like *T. brucei*, *L. major*, and *L. mexicana*, as

well as in lactobacillus species like *Lactobacillus helveticus* [107,108].

Bosch et al., 2006 conducted a comparative analysis of NDTs among different

trypanosomatid species, including *T. cruzi*, *L. major*, *L. infantum*, and *T. brucei*, along

with a sequence from Lactobacillus helveticus. The study revealed that the percentage

of identity between *T. brucei* and other species ranged from 84.2% (*T. cruzi*) to 89.5%

(*L. major* and *L. infantum*), and only 21% with *L. helveticus*. Our analysis of predicted

NDT domain sequences showed identity ranging from 65% between Tcr and Lif to

100% between Tbg and Tbr (Fig 5A).

Bosch's work also highlighted conserved regions among trypanosomatids,

which included key residues within the 2'-deoxyribose binding site, such as Ala15,

Gly16, Val19, Phe20, Pro46, Thr47, Glu50, Ile57, Arg58, Asn61, Arg125, Glu127,

Phe129, Asn134, Leu135, and Met136 [108]. Comparing these regions among the

generated structures, it was evident that most of them were conserved across species,

with only a few mutations observed, such as Thr47Val in Tcr and Lif; Glu50Ile and

Arg125Ser in Lif; Glu50Met and Arg125Thr in Tcr (Fig 5B). The RMSD values

between the mold and the models ranged from 0.467 Å (Tcr) to 0.611 Å (Lif).

Fig 5. NDT - Sequential and structural analysis. (A) The alignment reveals the

comparison between trypanosomatid and human sequences. Notably, the highlighted

region contains three amino acid mutations. The right side of the figure displays the

aligned residues: Ala15, Gly16, Val19, Phe20, Pro46, Thr47, Glu50, Ile57, Arg58,

Asn61, Arg125, Glu127, Phe129, Asn134, Leu135, and Met136. Among these residues,

three mutations were observed: Thr47Val, Glu50Met, and Arg125Thr, as indicated in

bold. The numbering of *T. cruzi* amino acids was adjusted to account for the eight fewer

amino acids compared to the template. (B) The alignment between the predicted model

for the Lif protein (light blue) and the template with PDBid: 2F62 (chain A in pink and

chain B in gray). This template was selected due to having more mutations and the

lowest RMSD. The aligned residues match those presented in part (A) and highlight the

conserved regions among the sequences. (C) Results for the druggability prediction

where Nuc deoxyrib tr Lif is in red (chain A) and blue (chain B) in a blue and the

druggability score and corresponding surface area values are color-coded: orange 0.81

(1649.44 Å²); purple 0.57 (354.17 Å²); green 0.43 (231.66 Å²); blue 0.4 (264.39 Å²);

pink 0.35 (213.49).

Following the utilization of AlphaFold2 and the subsequent evaluation of model

quality (refer to S1 Fig and S4 table for details), we proceeded to select the Lif structure

for predicting druggable sites. Among the identified sites, the orange region (Fig 5C)

scored the highest, signifying its potential druggability. Notably, these druggable sites

are concentrated within a region that contains key amino acids critical for the proper

functioning of this protein.

Analysis in parasites such as *Trypanosoma brucei*, *Trypanosoma cruzi*, and

Leishmania major, showed that Nucleoside 2'-deoxyribosyltransferases (NDTs) play a

crucial role in the processes of recycling and recovery of purines, as well as in the de

novo biosynthesis of purines [108]. These parasites employ various enzymes to remove

ribose or deoxyribose from purine nucleosides, making it challenging to determine

which enzymes are essential in the purine turnover pathway.

In contrast, humans rely on Deoxynucleoside 5-monophosphate N-glycosidase

(DNPH1 – EC 3.2.2.) for similar functions, but differences in substrate specificity and

essentiality differentiate it from NDTs [109]. Furthermore, it has been suggested that

DNPH1 is not an essential gene, as many organisms can survive without it. These

findings support the idea that NDT and DNPH1 enzymes are distinct. To further

compare the two, we aligned the predicted NDT domain sequences of various species

with the DNPH1 sequence of Homo sapiens (UniProtKB code: O43598). The identities

between these sequences ranged from 16% (between Lif and Hsa) to 22% (between Tcr

and Hsa), confirming their significant differences (Fig 5A).

Given the precariousness of enzymes involved in de novo purine biosynthesis in

T. brucei and their reliance on the host for purine uptake, inhibiting NDT, an enzyme

crucial for their processes, could significantly impact trypanosome replication [108]. In

light of the similarity among trypanosomatid sequences and the absence of NDT in

humans, NDT presents a promising therapeutic target for further exploration.

Mitochondrial membrane: MIX

The MIX domain is a unique protein structure anchored to the mitochondrial

membrane and found exclusively in kinetoplastids [110], a group of organisms that

includes Trypanosoma and Leishmania species. In these kinetoplastids, the precise

mechanisms of mitochondrial protein import and processing are not fully understood.

One approach to gaining insights into the function of the MIX domain is to study its

association with the Cytochrome C Oxidase (COX) multiprotein complex, as proposed
by Gorman et al. in 2011 [111].

Cytochrome C Oxidase (COX) plays a crucial role in mitochondrial cellular
respiration, driving oxidative phosphorylation to generate ATP molecules. This energy
production process involves various protein complexes (I, II, III, IV, and V) [112], with
COX representing complex IV in trypanosomatids [111]. Notably, the COX complex in
these organisms exhibits distinctive features in 8 of the 10 smaller subunits, for which
no counterparts are identified outside the Kinetoplastida group [113]. Early hypotheses
suggested that the MIX domain may be instrumental in directing or assembling COX
components [111].

The MIX domain comprises small sequences characterized by a conserved
primary structure, facilitating their targeting of mitochondria in parasitic and yeast cells
[110]. In their 2011 study, Gorman et al. compared the MIX sequence from
Trypanosoma brucei with that of other trypanosomatids, specifically *Leishmania major*
and *Trypanosoma cruzi*, revealing identity percentages ranging from 75% to 66%,
respectively. In the present study, comparisons of sequences predicted to contain this
domain show identity percentages ranging from 67% to 99% (Fig 6A).

Fig 6. MIX - Sequential and structural analysis. (A) This section features the
alignment of trypanosomatid sequences, with a specific emphasis on critical amino acid
residues (Arg86, Glu87, and Glu115) that play a pivotal role in protein dimerization. (B)
The structural alignment showcases the model sequence CAM65809 from *L. infantum*
in light blue, contrasted with the template PDBid: 3HA4 (F chain in pink and H chain in
gray). Notably, this template contains the sole mutation found in Asn158Lys,

represented in bold. Additionally, predictions from the DeepTMHMM web server are

illustrated, highlighting the transmembrane region in vibrant yellow and the

extracellular region in regal purple. A display of amino acid residues is provided on the

right side of the image, featuring elements like salt bridge formers (Arg86, Glu87, and

Glu115), as well as residues that engage in hydrogen bonds without mediation by water

molecules (Gln80, Glu160, Pro157, Gly159, and Asn163). (C) The results of

druggability assessments for Mix Lif are presented in red (chain A) and blue (chain B),

utilizing a cartoon representation (excluding the initial 70 amino acids). Druggability

scores and their corresponding surface area values are color-coded: green 0.87 (602.82

Å²); orange 0.81 (1446.28 Å²); purple 0.8 (772.76 Å²); pink 0.64 (404.49 Å²); blue 0.6

(369.83 Å²).

Moreover, Gorman et al. in 2011 identified three critical regions for the MIX

domain's structural integrity in various trypanosomatid species, using *Leishmania major*

(Lma) as the model organism. The first region involves three amino acid residues

(Arg86, Glu87, and Glu115) essential for dimerization. The second region consists of

amino acid residues that form salt bridges (Arg89, Glu90, and Glu118), while the third

region comprises amino acid residues that engage in hydrogen bonding without water

molecule mediation, including Gln80, Glu160, Pro157, Asn158, Gly159, and Asn163.

In our own work, we compared the same Lma sequence used by Gorman et al. in

2011 and found identity percentages ranging from 56% (between Lma and Tbg/Tbr) to

77% (between Lma and Lif). The first two regions remained conserved, while we

identified a single mutation in the third region, specifically in Lif: Asn158Lys (Fig 6B).

This mutation may impact the formation of chemical bonds due to the distinct

physicochemical characteristics of the lysine radical, suggesting the need for further

investigation to understand its effects on hydrogen bond formation.

In our structural analyses, the STBFP results and the template used led to the

structure PDBid:3HA4, representing a MIX protein from the *L. major* organism [114].

However, we observed that the generated models exhibited a significant region without

a clearly defined secondary structure, encompassing residues 1 to 70. To address this

issue, we employed the transmembrane region predictor, DeepTMHMM, based on prior

descriptions of the MIX domain. The results indicated a consensus for the Lif and

EAN89737 species of Tcr, while transmembrane regions were not predicted for Tbg and

Tbr (Table 2). This provides valuable insights into the structural characteristics of the

MIX domain in different trypanosomatid species.

Table 2. Results DeepTMHMM

Organism	Regions	Prediction
Lif	1 – 22	I
	23 – 32	Tmhelix
	33	O
	34 – 43	Tmhelix
	44 –T5	I
Tbg	1 – 198	I
Tbr	1 – 198	I
Tcr	1 – 27	I
	28 – 37	Tmhelix

	38	O
	39 – 48	Tmhelix
	49 – 198	I

[a]i: Intracelular region.

[b]TMhelix: Transmembrane region.

[c]o: Extracellular region.

Uboldi et al. (2006) postulated the existence of an N-terminal amphipathic α-

helix within the MIX domain, followed by a subsequent transmembrane region, with the

transmembrane portion typically being laterally inserted into the mitochondrial

membrane. In the case of *L. major*, this transmembrane region consists of the initial 45

residues [110]. The DeepTMHMM results for proteins that possess the MIX domain,

even those with unknown functions, align with this characteristic.

In terms of the structural attributes of MIX, it comprises seven α-helices that

collectively form the coiled domain [114]. This pattern of folding is consistent with our

generated models, and the root mean square deviation (RMSD) values derived from the

comparison of these models with the template (PDBid: 3HA4) ranged from 2.069 Å

(Tbr) to 2.493 Å (Lif). It is worth noting that the variability in RMSD values is likely

associated with the N-terminal segment (residues 1-45), for which the template lacks a

well-defined structure (Fig 6B).

In summary, the MIX domain is universally expressed across all evolutionary

stages of Leishmania, including the amastigote form residing in the mammalian host.

Intriguingly, both in vitro and in vivo experiments have demonstrated that the deletion

of two MIX gene alleles leads to a reduction in infectivity and virulence [110]. While

initial assessments in *T. brucei* suggested that COX activity was solely required in the

epimastigote form, indicating that the interaction between MIX and COX was specific

to this developmental stage [110,114], subsequent research has unveiled the presence of

MIX in the blood stage. This suggests that MIX could engage with other proteins and

catalyze additional reactions [111]. Given the unique characteristics of MIX within the

Kinetoplastida group and its pivotal roles in diverse processes, it emerges as a

promising molecular target for the development of novel drugs.

Conclusion

Based on the SEquence-Based Functional Prediction (SEBFP) and Structure-

Based Functional Prediction (STBFP) results, we were able to predict functional

domains for 67 trypanosomatid proteins, of which nine domains were shared among the

majority of trypanosomatid species. These domains were analyzed, by consulting

bibliographic resources, and we were able to clarify their functions and their existing

correlations with trypanosomatids.

Among the domains analyzed, we found those that had several isoforms, not

being crucial for the parasites under study, TRX, ACBP, and AAA 18 [95,115][115].

Additionally, the complexity of the pathways in which they are inserted, along with the

lack of detailed information on all the inherent metabolic processes, currently hinders

the development of inhibitors for these targets.

In addition to the aforementioned domains, we also identified two domains that

are integral to the ubiquitination process, a crucial mechanism for protein degradation.

Research has shown that mutations in UFC1, for instance, can result in neurological

impairment [63]. Similarly, Ufm1 can recruit transcriptional coactivators in humans,

contributing to excessive cell proliferation and ultimately leading to conditions such as

breast cancer [66]. Both domains possess complex intricacies that need to be addressed

before considering them as therapeutic targets. Nonetheless, the proteasome, another

component of the ubiquitination pathway, has been studied extensively due to its

significant role in *T. cruzi* [116]. Inhibitors of the proteasome have already been tested,

and the results demonstrate impairment of the infective evolutionary form, indicating

that a study considering the proteasome as a therapeutic target could yield promising

results.

Lastly, we identified domains that can be essential for the survival and virulence

of the trypanosomatids under study, namely Nuc deoxyrib tr, MIX, and TPR. Nuc

deoxyrib tr is critical in de novo purine biosynthesis, and studies on *T. brucei* have

shown that these parasites do not produce an abundance of such proteins, making them

heavily reliant on the catalytic activity of this domain for survival [108]. Regarding the

MIX domain, this is exclusive to kinetoplastids and *in vitro* and *in vivo* tests

demonstrate that its deletion led to a reduction in infectivity and virulence [111].

Furthermore, we also discovered the TPR domain, specifically the IFT70 protein, which

is involved in a protein complex that facilitates intraflagellar transport. IFT70 is a vital

member of the IFT train, and studies on *T. brucei* have confirmed that the train is

necessary for the transportation of ciliary assembly precursors from the cytoplasm into

the cell body [103]. Therefore, we believe these targets hold promise for multi-target

studies aimed at developing innovative therapeutic approaches.

Acknowledgments

References

[1]   Rasoanoro M, Ramasindrazana B, Goodman SM, Rajerison M, Randrianarivelojosia M. A review of Trypanosoma species known from Malagasy vertebrates. Malagasy Natiora 2019;13:65–75.

[2]   Thompson CK, Godfrey SS, Thompson RCA. Trypanosomes of Australian mammals: A review. Int J Parasitol Parasites Wildl 2014;3:57–66. https://doi.org/10.1016/j.ijppaw.2014.02.002.

[3]   Fahrion A, Gasimov E, Joseph S, Grout L, Allan M, Postigo JR. Surveillance of leishmaniasis in the WHO European Region. Rev Epidemiol Sante Publique 2018;66:S394. https://doi.org/10.1016/j.respe.2018.05.429.

[4]   WHO. Neglected tropical diseases 2017.

[5]   Imran M, Khan SA, Alshammari MK, Alqahtani AM, Alanazi TA, Kamal M, et al. Discovery, Development, Inventions and Patent Review of Fexinidazole: The First All-Oral Therapy for Human African Trypanosomiasis. Pharmaceuticals 2022;15. https://doi.org/10.3390/ph15020128.

[6]   Capewell P, Atkins K, Weir W, Jamonneau V, Camara M, Clucas C, et al. Resolving the apparent transmission paradox of African sleeping sickness. PLoS Biol 2019;17:1–6. https://doi.org/10.1371/journal.pbio.3000105.

[7]   Abdellahi L, Iraji F, Mahmoudabadi A, Hejazi SH. Vaccination in

leishmaniasis: A   review   article.   Iran   Biomed   J   2022;26:1–35.

https://doi.org/10.52547/ibj.26.1.35.

[8]   Pires M, Wright B, Kaye PM, da Conceição V, Churchill RC. The impact of

leishmaniasis on mental health and psychosocial well-being: A systematic review. PLoS

One 2019;14:e0223313. https://doi.org/10.1371/journal.pone.0223313.

[9]   Franco J, Scarone L, Comini MA. Drugs and Drug Resistance in African and

American   Trypanosomiasis.   vol.   51.   1st   ed.   Elsevier   Inc.;   2018.

https://doi.org/10.1016/bs.armc.2018.08.003.

[10]   De Fuentes-Vicente JA, Vidal-López DG, Flores-Villegas AL, Moreno-

Rodríguez A, De Alba-Alvarado MC, Salazar-Schettino PM, et al. Trypanosoma cruzi:

A review of biological and methodological factors in Mexican strains. Acta Trop

2019;195:51–7. https://doi.org/10.1016/j.actatropica.2019.04.024.

[11]   Field MC, Horn D, Fairlamb AH, Ferguson MAJ, Gray DW, Read KD, et al.

Anti-trypanosomatid drug discovery: An ongoing challenge and a continuing need. Nat

Rev Microbiol 2017;15:217–31. https://doi.org/10.1038/nrmicro.2016.193.

[12]   Pradhan S, Schwartz RA, Patil A, Grabbe S, Goldust M. Treatment options for

leishmaniasis. Clin Exp Dermatol 2022;47:516–21. https://doi.org/10.1111/ced.14919.

[13]   Zingales B. Trypanosoma cruzi genetic diversity: Something new for something

known about Chagas disease manifestations, serodiagnosis and drug sensitivity. Acta

Trop 2018;184:38–52. https://doi.org/10.1016/j.actatropica.2017.09.017.

[14]   Deotarse PP, Jain AS, Baile MB, Kolhe NS, Kulkarni AA. Drug

Repositioning : A Review. Int J Pharma Res Rev 2018;4:51–8.

[15]   Luo H, Li M, Yang M, Wu FX, Li Y, Wang J. Biomedical data and

736 computational models for drug repositioning: A comprehensive review. Brief Bioinform

737 2021;22:1604–19. https://doi.org/10.1093/bib/bbz176.

738 [16] Sadowski MI, Jones DT. The sequence-structure relationship and protein

739 function prediction. Curr Opin Struct Biol 2009;19:357–62.

740 https://doi.org/10.1016/j.sbi.2009.03.008.

741 [17] Radivojac P, Clark W, Oron T, Schnoes A. A large-scale evaluation of

742 computational protein function prediction. Nature 2013;10:221–7.

743 https://doi.org/10.1038/nmeth.2340.A.

744 [18] Araújo CL, Blanco I, Souza L, Tiwari S, Pereira LC, Ghosh P, et al. In silico

745 functional prediction of hypothetical proteins from the core genome of Corynebacterium

746 pseudotuberculosis biovar ovis. PeerJ 2020;8:1–24. https://doi.org/10.7717/peerj.9643.

747 [19] Shahbaaz M, Hassan MI, Ahmad F. Functional annotation of conserved

748 hypothetical proteins from Haemophilus influenzae Rd KW20. PLoS One 2013;8.

749 https://doi.org/10.1371/journal.pone.0084263.

750 [20] VERLI H. Bioinformática da Biologia à Flexibilidade Molecular. vol. 53. 2014.

751 https://doi.org/10.1017/CBO9781107415324.004.

752 [21] Punta M, Ofran Y. The rough guide to in silico function prediction, or how to

753 use sequence and structure information to predict protein function. PLoS Comput Biol

754 2008;4. https://doi.org/10.1371/journal.pcbi.1000160.

755 [22] Agarwala R, Barrett T, Beck J, Benson DA, Bollin C, Bolton E, et al. Database

756 resources of the National Center for Biotechnology Information. Nucleic Acids Res

757 2016;44:D7–19. https://doi.org/10.1093/nar/gkv1290.

758 [23] Capriles PVSZ, Guimarães ACR, Otto TD, Miranda AB, Dardenne LE, Degrave

759 WM. Structural modelling and comparative analysis of homologous, analogous and

specific proteins from Trypanosoma cruzi versus Homo sapiens: Putative drug targets for chagas' disease treatment. BMC Genomics 2010;11. https://doi.org/10.1186/1471-2164-11-610.

[24]   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol 1990;215:403–10. https://doi.org/10.1016/S0022-2836(05)80360-2.

[25]   Eswar N, Webb B, Marti-Renom MA, Madhusudhan MS, Eramian D, Shen M, et al. Comparative protein structure modeling using Modeller. vol. Chapter 5. 2006. https://doi.org/10.1002/0471250953.bi0506s15.Comparative.

[26]   Kanehisa M, Sato Y, Furumichi M, Morishima K, Tanabe M. New approach for understanding genome variations in KEGG. OXFORD Acad Nucleic Acids Res 2018.

[27]   El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, et al. The Pfam protein families database in 2019. Nucleic Acids Res 2019;47:D427–32. https://doi.org/10.1093/nar/gky995.

[28]   Letunic I, Khedkar S, Bork P. SMART: Recent updates, new developments and status in 2020. Nucleic Acids Res 2021;49:D458–60. https://doi.org/10.1093/nar/gkaa937.

[29]   Sigrist CJA, De Castro E, Cerutti L, Cuche BA, Hulo N, Bridge A, et al. New and continuing developments at PROSITE. Nucleic Acids Res 2013;41:344–7. https://doi.org/10.1093/nar/gks1067.

[30]   Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, et al. CDD/SPARCLE: The conserved domain database in 2020. Nucleic Acids Res 2020;48:D265–8. https://doi.org/10.1093/nar/gkz991.

[31]   Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, et al.

STRING v11: Protein-protein association networks with increased coverage, supporting

functional discovery in genome-wide experimental datasets. Nucleic Acids Res

2019;47:D607–13. https://doi.org/10.1093/nar/gky1131.

[32] Sillitoe I, Bordin N, Dawson N, Waman VP, Ashford P, Scholes HM, et al.

CATH: Increased structural coverage of functional space. Nucleic Acids Res

2021;49:D266–73. https://doi.org/10.1093/nar/gkaa1079.

[33] Potter SC, Luciani A, Eddy SR, Park Y, Lopez R, Finn RD. HMMER web

server: 2018 update. Nucleic Acids Res 2018;46:W200–4.

https://doi.org/10.1093/nar/gky448.

[34] Mistry J, Finn RD, Eddy SR, Bateman A, Punta M. Challenges in homology

search: HMMER3 and convergent evolution of coiled-coil regions. Nucleic Acids Res

2013;41. https://doi.org/10.1093/nar/gkt263.

[35] Peabody MA, Laird MR, Vlasschaert C, Lo R, Brinkman FSL. PSORTdb:

Expanding the bacteria and archaea protein subcellular localization database to better

reflect diversity in cell envelope structures. Nucleic Acids Res 2016;44:D663–8.

https://doi.org/10.1093/nar/gkv1271.

[36] Bateman A, Martin MJ, Orchard S, Magrane M, Agivetova R, Ahmad S, et al.

UniProt: the universal protein knowledgebase in 2021. Nucleic Acids Res

2021;49:D480–9. https://doi.org/10.1093/nar/gkaa1100.

[37] Bateman A. UniProt: A worldwide hub of protein knowledge. Nucleic Acids

Res 2019;47:D506–15. https://doi.org/10.1093/nar/gky1049.

[38] Aslett M, Aurrecoechea C, Berriman M, Brestelli J, Brunk BP, Carrington M, et

al. TriTrypDB: A functional genomic resource for the Trypanosomatidae. Nucleic Acids

Res 2009;38:457–62. https://doi.org/10.1093/nar/gkp851.

[39]   Amos B, Aurrecoechea C, Barba M, Barreto A, Basenko EY, Bażant W, et al.

VEuPathDB: The eukaryotic pathogen, vector and host bioinformatics resource center.

Nucleic Acids Res 2022;50:D898–911. https://doi.org/10.1093/nar/gkab929.

[40]   Okonechnikov K, Golosova O, Fursov M, Varlamov A, Vaskin Y, Efremov I, et

al. Unipro UGENE: A unified bioinformatics toolkit. Bioinformatics 2012;28:1166–7.

https://doi.org/10.1093/bioinformatics/bts091.

[41]   Löytynoja A, Goldman N. An algorithm for progressive multiple alignment of

sequences with insertions. Proc Natl Acad Sci U S A 2005;102:10557–62.

https://doi.org/10.1073/pnas.0409137102.

[42]   Edgar RC. MUSCLE: Multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res 2004;32:1792–7. https://doi.org/10.1093/nar/gkh340.

[43]   Naser-Khdour S, Minh BQ, Zhang W, Stone EA, Lanfear R, Bryant D. The

Prevalence and Impact of Model Violations in Phylogenetic Analysis. Genome Biol

Evol 2019;11:3341–52. https://doi.org/10.1093/gbe/evz193.

[44]   Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O,

Brunak S, et al. SignalP 5.0 improves signal peptide predictions using deep neural

networks. Nat Biotechnol 2019;37:420–3. https://doi.org/10.1038/s41587-019-0036-z.

[45]   Hallgren J, Tsirigos KD, Damgaard Pedersen M, Juan J, Armenteros A,

Marcatili P, et al. DeepTMHMM predicts alpha and beta transmembrane proteins using

deep neural networks. BioRxiv 2022:2022.04.08.487609.

[46]   Hubbard TJP, Murzin AG, Brenner SE, Chothia C. SCOP: A Structural

Classification of Proteins database. Nucleic Acids Res 1997;25:236–9.

https://doi.org/10.1093/nar/25.1.236.

[47]   Madej T, Lanczycki CJ, Zhang D, Thiessen PA, Geer RC, Marchler-Bauer A, et

al. MMDB and VAST+: Tracking structural similarities between macromolecular

complexes. Nucleic Acids Res 2014;42:297–303. https://doi.org/10.1093/nar/gkt1208.

[48]   Madej T, Marchler-Bauer A, Lanczycki C, Zhang D, Bryant SH. Biological

Assembly Comparison with VAST+. Struct Bioinforma 2020;2112.

[49]   Li Z, Jaroszewski L, Iyer M, Sedova M, Godzik A. FATCAT 2.0: Towards a

better understanding of the structural diversity of proteins. Nucleic Acids Res

2020;48:W60–4. https://doi.org/10.1093/NAR/GKAA443.

[50]   Holm L. Using Dali for protein structure comparison. Struct Bioinforma

Methods Protoc Methods Mol Biol 2020;2112:29–42.

[51]   DeLano WL. The PyMol molecular graphics system 2002.

[52]   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al.

Highly accurate protein structure prediction with AlphaFold. Nature 2021;596:583–9.

https://doi.org/10.1038/s41586-021-03819-2.

[53]   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al.

AlphaFold Protein Structure Database: Massively expanding the structural coverage of

protein-sequence space with high-accuracy models. Nucleic Acids Res 2022;50:D439–

44. https://doi.org/10.1093/nar/gkab1061.

[54]   Gabler F, Nam SZ, Till S, Mirdita M, Steinegger M, Söding J, et al. Protein

Sequence Analysis Using the MPI Bioinformatics Toolkit. Curr Protoc Bioinforma

2020;72. https://doi.org/10.1002/cpbi.108.

[55]   Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M.

ColabFold: making protein folding accessible to all. Nat Methods 2022;19:679–82.

https://doi.org/10.1038/s41592-022-01488-1.

[56]   Williams CJ, Headd JJ, Moriarty NW, Prisant MG, Videau LL, Deis LN, et al.

MolProbity: More and better reference data for improved all-atom structure validation.

Protein Sci 2018;27:293–315. https://doi.org/10.1002/pro.3330.

[57]   Wiederstein M, Sippl MJ. ProSA-web: Interactive web service for the

recognition of errors in three-dimensional structures of proteins. Nucleic Acids Res

2007;35:407–10. https://doi.org/10.1093/nar/gkm290.

[58]   Volkamer A, Kuhn D, Rippmann F, Rarey M. Dogsitescorer: A web server for

automatic binding site prediction, analysis and druggability assessment. Bioinformatics

2012;28:2074–5. https://doi.org/10.1093/bioinformatics/bts310.

[59]   Blum M, Chang HY, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, et

al. The InterPro protein families and domains database: 20 years on. Nucleic Acids Res

2021;49:D344–54. https://doi.org/10.1093/nar/gkaa977.

[60]   Marchler-Bauer A, Zheng C, Chitsaz F, Derbyshire MK, Geer LY, Geer RC, et

al. CDD: Conserved domains and protein three-dimensional structure. Nucleic Acids

Res 2013;41:348–52. https://doi.org/10.1093/nar/gks1243.

[61]   Kimura Y, Tanaka K. Regulatory mechanisms involved in the control of

ubiquitin homeostasis. J Biochem 2010;147:793–8.

[62]   Komatsu M, Chiba T, Tatsumi K, Lemura SI, Tanida I, Okazaki N, et al. A

novel protein-conjugating system for Ufm1, a ubiquitin-fold modifier. EMBO J

2004;23:1977–86. https://doi.org/10.1038/sj.emboj.7600205.

[63]   Nahorski MS, Maddirevula S, Ishimura R, Alsahli S, Brady AF, Begemann A, et

al. Biallelic UFM1 and UFC1 mutations expand the essential role of ufmylation in brain

development. Brain 2018;141:1934–45. https://doi.org/10.1093/brain/awy135.

[64]   Chung WL, Leung KF, Carrington M, Field MC. Ubiquitylation is required for

degradation of transmembrane surface proteins in Trypanosomes. Traffic 2008;9:1681–

97. https://doi.org/10.1111/j.1600-0854.2008.00785.x.

[65]   Leung KF, Riley FS, Carrington M, Field MC. Ubiquitylation and

developmental regulation of invariant surface protein expression in trypanosomes.

Eukaryot Cell 2011;10:916–31. https://doi.org/10.1128/EC.05012-11.

[66]   Yoo HM, Kang SH, Kim JY, Lee JE, Seong MW, Lee SW, et al. Modification

of asc1 by ufm1 is crucial for erα transactivation and breast cancer development. Mol

Cell 2014;56:261–74. https://doi.org/10.1016/j.molcel.2014.08.007.

[67]   Huang X, Dixit VM. Drugging the undruggables: Exploring the ubiquitin system

for drug development. Cell Res 2016;26:484–98. https://doi.org/10.1038/cr.2016.31.

[68]   Hochstrasser M. Evolution and function of ubiquitin-like protein-conjugation

systems 2000;2:153–7.

[69]   González J, Ramalho-Pinto FJ, Frevert U, Ghiso J, Tomlinson S, Scharfstein J

ulio, et al. Proteasome activity is required for the stage-specific transformation of a

protozoan parasite. J Exp Med 1996;184:1909–18.

[70]   Kurup SP, Tarleton RL. The trypanosoma cruzi flagellum is discarded via

asymmetric cell division following invasion and provides early targets for protective

CD8+   T   cells.   Cell   Host   Microbe   2014;16:439–49.

https://doi.org/10.1016/j.chom.2014.09.003.

[71]   Ayala-Castro C, Saini A, Outten FW. Fe-S Cluster Assembly Pathways in

Bacteria.   Microbiol   Mol   Biol   Rev   2008;72:110–25.

https://doi.org/10.1128/mmbr.00034-07.

[72]   Kispal G, Csere P, Prohl C, Lill R. The mitochondrial proteins Atm1p and Nfs1p

are essential for biogenesis of cytosolic Fe/S proteins. EMBO J 1999;18:3981–9.

https://doi.org/10.1093/emboj/18.14.3981.

[73]   Mettert EL, Kiley PJ. How Is Fe-S Cluster Formation Regulated? Annu Rev

Microbiol   2015;69:505–26.   https://doi.org/10.1146/annurev-micro-091014-

104457.How.

[74]   Py B, Barras F. Building Fe-S proteins: Bacterial strategies. Nat Rev Microbiol

2010;8:436–46. https://doi.org/10.1038/nrmicro2356.

[75]   Basu S, Netz DJ, Haindrich AC, Herlerth N, Lagny TJ, Pierik AJ, et al.

Cytosolic iron-sulphur protein assembly is functionally conserved and essential in

procyclic and bloodstream Trypanosoma brucei. Mol Microbiol 2014;93:897–910.

https://doi.org/10.1111/mmi.12706.

[76]   Stehling O, Mascarenhas J, Vashisht AA, Sheftel AD, Niggemeyer B, Rösser R,

et al. Human CIA2A-FAM96A and CIA2B-FAM96B integrate iron homeostasis and

maturation of different subsets of cytosolic-nuclear iron-sulfur proteins. Cell Metab

2013;18:187–98. https://doi.org/10.1016/j.cmet.2013.06.015.

[77]   Schwamb B, Pick R, Fernández SBM, Völp K, Heering J, Dötsch V, et al.

FAM96A is a novel pro-apoptotic tumor suppressor in gastrointestinal stromal tumors.

Int J Cancer 2015;137:1318–29. https://doi.org/10.1002/ijc.29498.

[78]   Yin A, Luo Y, Chen W, He M, Deng JH, Zhao N, et al. FAM96A Protects Mice

From Dextran Sulfate Sodium (DSS)-Induced Colitis by Preventing Microbial

Dysbiosis.   Front   Cell   Infect   Microbiol   2019;9:1–14.

https://doi.org/10.3389/fcimb.2019.00381.

[79]   Yin A, Chen W, Cao L, Li Q, Zhu X, Wang L. FAM96A knock-out promotes

alternative macrophage polarization and protects mice against sepsis. Clin Exp Immunol

2021;203:433–47. https://doi.org/10.1111/cei.13555.

[80]   Lukeš J, Basu S. Fe/S protein biogenesis in trypanosomes - A review. Biochim

Biophys Acta - Mol Cell Res 2015;1853:1481–92.

https://doi.org/10.1016/j.bbamcr.2014.08.015.

[81] Sipos K, Lange H, Fekete Z, Ullmann P, Lill R, Kispal G. Maturation of

cytosolic iron-sulfur proteins requires glutathione. J Biol Chem 2002;277:26944–9.

https://doi.org/10.1074/jbc.M200677200.

[82] Long S, Changmai P, Tsaousis AD, Skalický T, Verner Z, Wen YZ, et al. Stage-

specific requirement for Isa1 and Isa2 proteins in the mitochondrion of Trypanosoma

brucei and heterologous rescue by human and Blastocystis orthologues. Mol Microbiol

2011;81:1403–18. https://doi.org/10.1111/j.1365-2958.2011.07769.x.

[83] Milne KG, Güther MLS, Ferguson MAJ. Acyl-CoA binding protein is essential

in bloodstream form Trypanosoma brucei. Mol Biochem Parasitol 2001;112:301–4.

https://doi.org/10.1016/S0166-6851(00)00369-8.

[84] Konrad M. Molecular analysis of the essential gene for adenylate kinase from

the fission yeast Schizosaccharomyces pombe. J Biol Chem 1993;268:11326–34.

https://doi.org/10.1016/s0021-9258(18)82128-7.

[85] Milne KG, Ferguson MAJ. Cloning, expression, and characterization of the

acyl-coA-binding protein in African trypanosomes. J Biol Chem 2000;275:12503–8.

https://doi.org/10.1074/jbc.275.17.12503.

[86] Lee SH, Stephens JL, Englund PT. A fatty-acid synthesis mechanism specialized

for parasitism. Nat Rev Microbiol 2007;5:287–97. https://doi.org/10.1038/nrmicro1617.

[87] Smith TK, Bütikofer P. Lipid metabolism in Trypanosoma brucei. Mol Biochem

Parasitol 2010;172:66–79. https://doi.org/10.1016/j.molbiopara.2010.04.001.

[88] Mehlert A, Bond CS, Ferguson MAJ. The glycoforms of a Trypanosoma brucei

variant surface glycoprotein and molecular modeling of a glycosylated surface coat.

Glycobiology 2002;12:607–12. https://doi.org/10.1093/glycob/cwf079.

[89]   McConville MJ, Mullin KA, Ilgoutz SC, Teasdale RD. Secretory Pathway of Trypanosomatid Parasites. Microbiol Mol Biol Rev 2002;66:122–54. https://doi.org/10.1128/mmbr.66.1.122-154.2002.

[90]   Bouvier LA, Miranda MR, Canepa GE, Alves MJM, Pereira CA. An expanded adenylate kinase gene family in the protozoan parasite Trypanosoma cruzi. Biochim Biophys Acta - Gen Subj 2006;1760:913–21. https://doi.org/10.1016/j.bbagen.2006.02.013.

[91]   Klepinin A, Zhang S, Klepinina L, Rebane-Klemm E, Terzic A, Kaambre T, et al. Adenylate Kinase and Metabolic Signaling in Cancer Cells. Front Oncol 2020;10:1–9. https://doi.org/10.3389/fonc.2020.00660.

[92]   Berger J, Dorninger F, Forss-Petter S, Kunze M. Peroxisomes in brain development and function. Biochim Biophys Acta - Mol Cell Res 2016;1863:934–55. https://doi.org/10.1016/j.bbamcr.2015.12.005.

[93]   Hubert L, Sutton VR. Disorders of purine and pyrimidine metabolism. Elsevier Inc.; 2017. https://doi.org/10.1016/b978-0-12-802896-4.00009-2.

[94]   Yin J, Ren W, Huang X, Deng J, Li T, Yin Y. Potential Mechanisms Connecting Purine Metabolism and Cancer Therapy. Front Immunol 2018;9:1–8. https://doi.org/10.3389/fimmu.2018.01697.

[95]   Currier RB, Ulrich K, Leroux AE, Dirdjaja N, Deambrosi M, Bonilla M, et al. An essential thioredoxin-type protein of Trypanosoma brucei acts as redox-regulated mitochondrial chaperone. vol. 15. 2019. https://doi.org/10.1371/journal.ppat.1008065.

[96]   Zeytuni N, Zarivach R. Structural and functional discussion of the tetra-trico-peptide repeat, a protein interaction module. Structure 2012;20:397–405.

https://doi.org/10.1016/j.str.2012.01.006.

[97]   Young JC, Barral JM, Hartl FU. More than folding: localized functions of

cytosolic chaperones. Trends Biochem Sci 2003;28.

[98]   Baker MJ, Frazier AE, Gulbis JM, Ryan MT. Mitochondrial protein-import

machinery: correlating structure with function. Trends Cell Biol 2007;17.

[99]   Brocard C, Hartig A. Peroxisome targeting signal 1: Is it really a simple

tripeptide? Biochim Biophys Acta 2006;1763:1565–73.

[100] Takei R, Katoh Y, Nakayama K. Robust interaction of IFT70 with IFT52–IFT88

in   the   IFT-B   complex   is   required   for   ciliogenesis.   Biol   Open   2018;7.

https://doi.org/10.1242/bio.033241.

[101] Taschner M, Kotsis F, Braeuer P, Wolfgang Kuehn E, Lorentzen E. Crystal

structures of IFT70/52 and IFT52/46 provide insight into intraflagellar transport B core

complex   assembly.   J   Cell   Biol   2014;207:269–82.

https://doi.org/10.1083/jcb.201408002.

[102] Guido RVC, Andricopulo AD, Oliva G. Drug design, biotechnology and

medicinal chemistry: applications to infectious diseases. Estud Avançados 2010;24:81–

98. https://doi.org/10.1016/j.neuroscience.2012.06.055.

[103] Wachter S, Jung J, Shafiq S, Basquin J, Fort C, Bastin P, et al. Binding of IFT22

to the intraflagellar transport complex is essential for flagellum assembly. EMBO J

2019;38:1–18. https://doi.org/10.15252/embj.2018101251.

[104] Bertiaux E, Mallet A, Rotureau B, Bastin P. Intraflagellar transport during

assembly of flagella of different length in Trypanosoma brucei isolated from tsetse flies.

J Cell Sci 2020;133. https://doi.org/10.1242/jcs.248989.

[105] Cardinaud   R,   Holguin   J.   Nucleoside   deoxyribosyltransferase-II   from

Lactobacillus helveticus. Substrate specificity studies. Pyrimidine bases as acceptors.

Biochim Biophys Acta - Enzymol 1979;568:339–47.

[106] Fresco-Taboada A, De La Mata I, Arroyo M, Fernández-Lucas J. New insights

on nucleoside 2′-deoxyribosyltransferases: A versatile biocatalyst for one-pot one-step

synthesis of nucleoside analogs. Appl Microbiol Biotechnol 2013;97:3773–85.

https://doi.org/10.1007/s00253-013-4816-y.

[107] Fernández-Lucas J, Fresco-Taboada A, Acebal C, De La Mata I, Arroyo M.

Enzymatic   synthesis   of   nucleoside   analogues   using   immobilized   2′-

deoxyribosyltransferase   from   Lactobacillus   reuteri.   Appl   Microbiol   Biotechnol

2011;91:317–27. https://doi.org/10.1007/s00253-011-3221-7.

[108] Bosch J, Robien MA, Mehlin C, Boni E, Riechers A, Buckner FS, et al. Using

fragment cocktail crystallography to assist inhibitor design of Trypanosoma brucei

nucleoside   2-deoxyribosyltransferase.   J   Med   Chem   2006;49:5939–46.

https://doi.org/10.1021/jm060429m.

[109] Reintamm T, Aas-Valleriani N, Kelve M. Evolutionary distribution of

deoxynucleoside 5-monophosphate N-glycosidase, DNPH1. Gene 2019;683:1–11.

https://doi.org/10.1016/j.gene.2018.10.005.

[110] Uboldi AD, Lueder FB, Walsh P, Spurck T, McFadden GI, Curtis J, et al. A

mitochondrial protein affects cell morphology, mitochondrial segregation and virulence

in   Leishmania.   Int   J   Parasitol   2006;36:1499–514.

https://doi.org/10.1016/j.ijpara.2006.08.006.

[111] Zíková A, Panigrahi AK, Uboldi AD, Dalley RA, Handman E, Stuart K.

Structural and functional association of Trypanosoma brucei MIX protein with

cytochrome   c   oxidase   complex.   Eukaryot   Cell   2008;7:1994–2003.

https://doi.org/10.1128/EC.00204-08.

[112] Fitzpatrick F. Cyclooxygenase Enzymes: Regulation and Function. Curr Pharm

Des 2005;10:577–88. https://doi.org/10.2174/1381612043453144.

[113] Speijer D, Breek CKD, Muijsers AO, Groenevelt PX, Dekker H, De Haan A, et

al. The sequence of a small subunit of cytochrome c oxidase from Crithidia fasciculata

which is homologous to mammalian subunit IV. FEBS Lett 1996;381:123–6.

https://doi.org/10.1016/0014-5793(96)00095-6.

[114] Gorman MA, Uboldi AD, Walsh PJ, Tan KS, Hansen G, Huyton T, et al. Crystal

structure of the Leishmania major MIX protein: A scaffold protein that mediates

protein-protein interactions. Protein Sci 2011;20:1060–8.

https://doi.org/10.1002/pro.631.

[115] Burton M, Rose TM, Færgeman NJ, Knudsen J. Evolution of the acyl-CoA

binding protein (ACBP). Biochem J 2005;392:299–307.

https://doi.org/10.1042/BJ20050664.

[116] Bijlmakers MJ. Ubiquitination and the Proteasome as Drug Targets in

Trypanosomatid Diseases. Front Chem 2021;8:1–18.

https://doi.org/10.3389/fchem.2020.630888.

1041

Supporting Files

S1 Fig. AlphaFold2 prediction. Presented in both cartoon and surface views. These

models pertain to the TPR, Nuc deoxyrib tr, and MIX domains, covering various

species, namely *L. infantum* (Lif), *T. brucei gambiensi* (Tbg), *T. brucei brucei* (Tbr),

and *T. cruzi* (Tcr). In this representation, monomeric TPR models are depicted in

vibrant purple, offering insights into the individual subunits. Dimeric structures are

presented with chain A in a distinct orange shade and chain B in a contrasting blue hue,

allowing for clear differentiation between the two subunits. Arrows within the figures

highlight regions where MIX transmembrane segments are predicted, indicating

potential areas of interest for further study and analysis.

S1 Table. Domain index.

S2 Table. Functional prediction results. The abbreviations in the table mean WR –

Without Results; UP - Uncharacterized Protein; UCP - Uncharacterized Conserved

Protein.

S3 Table. Proteins/Domains sequences.

S4 Table. Oligomeric AlphaFold Models - Validation.

Functional prediction

SEBFP

STBFP

Functional Domain

*SEBFP = SEquence-Based Functional Prediction
*STBFP = STructure-Based Functional Prediction

A

	Hsa_IFT70A	Hsa_IFT70B	Lif_IFT70_CAM69560	Tbg_IFT70_CBH10276	Tbr_IFT70_AAZ10580	Tcr_IFT70_EAN96986
Hsa_IFT70A	100%	96%	51%	52%	54%	51%
Hsa_IFT70B	96%	100%	52%	52%	54%	51%
Lif_IFT70_CAM69560	51%	52%	100%	74%	73%	75%
Tbg_IFT70_CBH10276	52%	52%	74%	100%	98%	90%
Tbr_IFT70_AAZ10580	54%	54%	73%	98%	100%	88%
Tcr_IFT70_EAN96986	51%	51%	75%	90%	88%	100%

B

Model
4UZY
TPR
IFT52

C

0.81
0.76
0.84
0.78
0.76

A

	Hsa_DNPH1	Lif_Nucdeoxyrib_CBZ08689.1	Tbg_Nucdeoxyrib_CBH11048.1	Tbr_Nucdeoxyrib_AAZ11251.1	Tcr_Nucdeoxyrib_EAN91728.1
Hsa_DNPH1	100%	16%	20%	20%	22%
Lif_Nucdeoxyrib_CBZ08689.1	16%	100%	72%	72%	65%
Tbg_Nucdeoxyrib_CBH11048.1	20%	72%	100%	100%	68%
Tbr_Nucdeoxyrib_AAZ11251.1	20%	72%	100%	100%	68%
Tcr_Nucdeoxyrib_EAN91728.1	22%	65%	68%	68%	100%

B

C

A

	Lma_MIX_3HA4	Lif_MIX_CAM65809.1	Tbg_MIX_CBH11287.1	Tbr_MIX_AAZ11417.1	Tcr_MIX_EAN86385.1	Tcr_MIX_EAN89737.1
Lma_MIX_3HA4	100%	77%	56%	56%	61%	60%
Lif_MIX_CAM65809.1	77%	100%	67%	67%	74%	73%
Tbg_MIX_CBH11287.1	56%	67%	100%	99%	83%	83%
Tbr_MIX_AAZ11417.1	56%	67%	99%	100%	83%	83%
Tcr_MIX_EAN86385.1	61%	74%	83%	83%	100%	99%
Tcr_MIX_EAN89737.1	60%	73%	83%	83%	99%	100%

B

C