1   **High sugar diets can increase susceptibility to bacterial infection in *Drosophila***

2   ***melanogaster***

4   Andrea M. Darby[1,2], Destiny O. Okoro[1,2], Sophia Aredas,[2,3,4], Ashley M. Frank[1,5], William H.

5   Pearson[6,7], Marc S. Dionne[6,7], Brian P. Lazzaro[1,2]

6   1. Department of Entomology, Cornell University, Ithaca, NY, United States of

7   America

8   2. Cornell Institute of Host-Microbe Interactions and Disease, Cornell University,

9   Ithaca, NY, United States of America

10   3. University of California, Irvine, Irvine, CA, United States of America

11   4. Department of Microbiology, Cornell University, Ithaca, NY, United States of

12   America

13   5. Battelle, Columbus, Ohio, United States of America

14   6. Department of Life Sciences, Imperial College London, London SW7 2AZ, United

15   Kingdom

16   7. Centre for Bacterial Resistance Biology, Imperial College London, London SW7

17   2AZ, United Kingdom

Short title: Effects of high-sugar diets on innate immunity

Keywords: *Drosophila,* high-sugar, innate immunity, metabolism, bacterial infection

Abstract

Overnutrition with dietary sugar can worsen infection outcomes in diverse organisms including

insects and humans, generally through unknown mechanisms. In the present study, we show

that adult *Drosophila melanogaster* fed high-sugar diets became more susceptible to infection

by the Gram-negative bacteria *Providencia rettgeri* and *Serratia marcescens,* although diet had

no significant effect on infection by Gram-positive bacteria *Enterococcus faecalis* or

Lactococcus lactis. We found that *P. rettgeri* and *S. marcescens* proliferate more rapidly in *D.*

melanogaster fed a high-sugar diet, resulting in increased probability of host death. *D.*

melanogaster become hyperglycemic on the high-sugar diet, and we find evidence that the

extra carbon availability may promote *S. marcescens* growth within the host. However, we found

no evidence that increased carbon availability directly supports greater *P. rettgeri* growth. *D.*

melanogaster on both diets fully induce transcription of antimicrobial peptide (AMP) genes in

response to infection, but *D. melanogaster* provided with high-sugar diets show reduced

production of AMP protein. Thus, overnutrition with dietary sugar may impair host immunity at

the level of AMP translation. Our results demonstrate that dietary sugar can shape infection

dynamics by impacting both host and pathogen, depending on the nutritional requirements of

the pathogen and by altering the physiological capacity of the host to sustain an immune

response.

Author Summary

Diet has critical impact on the quality of immune defense, and high-sugar diets increase

susceptibility to bacterial infection in many animals. Yet it is unknown which aspects of host and

pathogen physiology are impacted by diet to influence infection dynamics. Here we show that

high-sugar diets increase susceptibility to some, but not all, bacterial infections in *Drosophila*.

We find that feeding on high sugar diet impairs the host immune response by reducing the level

of antimicrobial peptides produced. The expression of genes encoding these peptides is not

affected, so we infer that protein translation is impaired. We further show that flies on high-sugar

diets are hyperglycemic, and that some pathogens may use the excess sugar in the host to

promote growth during the infection. Thus, our study demonstrates that dietary impacts on

infection outcome arise through physiological effects on both the host and pathogen.

Introduction

Nutritional status and diet are important factors for immune defense in both mammals and

insects (1–5). Hyperglycemia and diets excessively high in sugar have adverse effects on

metabolic homeostasis and infection outcome (6–10). For example, human patients admitted to

the intensive care unit with sepsis have a higher probability of death if they have elevated blood

sugar (11) and non-diabetic hyperglycemia increases the severity of *Mycobacterium*

tuberculosis infections in guinea pigs (1). A similar phenomenon is observed in the fruit fly

Drosophila melanogaster, where adults reared on high sugar diets experience higher pathogen

burden (3) and increased mortality from systemic bacterial infection (2,12). Despite these clear

effects of diet and nutritional state on infection outcome, the molecular and physiological

mechanisms by which high sugar impacts infections is often unclear. By uncovering these

mechanisms, we can clarify the role that diet has on shaping organismal physiology during an

active infection and identify strategies to alleviate consequences of infection that are

exacerbated by obesogenic diets.

Drosophila is an excellent model to understand the mechanisms by which high-sugar diets

affect infection dynamics, with extensive prior study on infection, diet, and metabolism (e.g., 13–

16). Lacking adaptive immune systems, *Drosophila* rely on innate immune defenses that include

humoral production and secretion into circulation of antimicrobial peptides (AMP) and cytokines

secreted that share homologous function with innate immune defenses in mammals and other

invertebrate species (17,18). The Toll and the immune deficiency (IMD) pathways are the two

major regulators of the *Drosophila* humoral immune response to infection (14). Gram-positive

bacterial and fungal infections predominantly stimulate Toll activity (19,20) while Gram-negative

bacterial infections activate the IMD pathway (19,21). Once activated, both pathways lead to the

nuclear translocation of nuclear factor-κB (NF-κB) family transcription factors to drive expression

of hundreds of infection-responsive genes, including those encoding antimicrobial peptides

(22,23). Infection and/or constitutive activation of these pathways stimulates major shifts in

metabolic processes to support the immune response, including suppressed insulin signaling,

reduction in energetic stores like glycogen and triglycerides, and alterations in carbohydrate and

lipid metabolism (24–28).

Both high-sugar diets and infection alter metabolism and energetic usage in *Drosophila* (29).

In the absence of infection, high-sugar diets can also upregulate genes involved in the immune

response, including antimicrobial peptides (12,30), which is similar to high levels of dietary

sugar stimulating low-grade inflammation responses in humans (31). Additionally, uninfected

Drosophila larvae reared on high-sugar diets exhibit impaired melanization and reduced

phagocytic capacity of fungal spores (12,32). While there is evidence that high-sugar diets affect

the *Drosophila* immune response in uninfected states, it is less clear whether high-sugar diets

impact immune activity during a live infection. Resistance to infection can be measured as

immune system activity or control of pathogen burden (33). Feeding *Drosophila* high-glucose

diets results in higher pathogen burden after systemic infection with the Gram-negative bacteria

Providencia rettgeri (2,3) and *Pseudomonas aeruginosa* (6). This could suggest that high-sugar

diets impair host immune system activity, but *Pseudomonas aeruginosa* is additionally able to

use glucose in hyperglycemic hosts to establish higher pathogen burdens (6). Thus, it has been

difficult to discern whether elevated pathogen burden in *Drosophila* provided with high-sugar

diets is due to impaired host immune responses or to bolstered pathogen growth capacity.

These are not mutually exclusive alternatives. Furthermore, prior *D. melanogaster* studies that

investigated the effects of high dietary sugar on systemic infection outcome were performed

using flies that were reared throughout their entire development on high-sugar diets (2,3,12,32).

Rearing on elevated dietary sugar diets causes hyperglycemia, lipidemia, and reduced body

weight in both larvae and as adults, and prolongs larval developmental time (12,34). Thus, it is

difficult to determine in those experiments whether effects of diet on adult infection outcome

result from altered metabolism and immunity or from indirect consequences of altered

development.

In the present study, we specifically focus on the immediate effects of high dietary sugar on

infection dynamics in adult *D. melanogaster*. We do this by rearing *D. melanogaster* larvae on a

common base diet, then switching adults to diets that vary in sugar content prior to delivering

systemic bacterial infection. To determine whether adverse effects from high-sugar diets are

universal across pathogenic infections, we test infection with four different bacterial pathogens

representing Gram-positive and Gram-negative bacteria. We find that flies provided with high-

sugar diet as adults are particularly susceptible to Gram-negative bacterial infection, suffering

increased mortality and more rapid pathogen proliferation during the early hours of infection. We

find evidence that increased carbon availability in high-sugar hosts can accelerate the growth of

certain pathogens, and that high-sugar diets impaired immune system function at the level of

antimicrobial peptide translation without affecting transcription levels.

Results

High-sucrose diets increase mortality from infection with some bacterial pathogens

We first wanted to test whether high-sugar diets reduce immune defense in adult *D.*

melanogaster independent of any developmental consequences. We therefore reared larvae on

a yeast-cornmeal diet containing 4% sucrose, then split the population across the experimental

diets during the non-feeding pupal stage such that eclosing adults emerged onto experimental

diets that were 0%, 2%, 4%, 8%, 16%, or 24% (w/v) sucrose (Fig. 1A), which covers the range

used in prior studies (3,12,32). After 3-5 days on the experimental diets, adult female flies were

given a systemic infection with one of four bacteria: *Providencia rettgeri*, *Serratia marcescens,*

Enterococcus faecalis, Lactococcus lactis. These bacteria were chosen as natural pathogens of

D. melanogaster (35,36) and include both Gram-positive and Gram-negative bacteria, which are

respectively expected to predominantly activate the Toll and IMD immune response pathways

(19).

We found that flies provided with high-sucrose diets experienced higher mortality after

infection with the Gram-negative bacteria *S. marcescens* and *P. rettgeri* than flies fed low sugar

(Figs. 1B and 1C). This increased mortality occurred in a concentration-dependent manner

depending on the pathogen. Flies provided with 0%, 2%, or 4% sucrose experienced low

mortality and flies provided with 16% or 24% sucrose experienced high mortality after infection

with both pathogens. However, on the 8% sucrose diet, flies infected with *S. marcescens*

exhibited high survival (Fig. 1B), but flies infected with *P. rettgeri* exhibited low survival (Fig.

1C). Overall, there was no effect of diet on survivorship of infection with the Gram-positive

bacteria *L. lactis* and *E. faecalis* (Figs. 1D and 1E). Although flies provided with the 24%

sucrose diet experienced the highest mortality after *L. lactis* infection (Fig. 1D), we attribute this

to the diet being generally poor as sham-infected control flies fed on the 24% sucrose diet also

experienced elevated mortality (Fig. S1A). Varying the diet of the host can alter the composition

of the gut microbiota (37–39), which can yield effects on the host's metabolic status (40–42).

Flies provided with 16% sucrose in our experiments showed a reduced abundance of gut flora

compared to flies provided with 2% sucrose diet (Fig S2A). However, we found that high-sugar

diets increased the probability of death from systemic *P. rettgeri* infection even in axenic flies

with no microbiota (Fig S2B), suggesting that our observed effect of diet on immunity is not

mediated by effects on gut flora.

Fig 1. High-sugar diet increases mortality after infection with some pathogens. (A) Experimental setup for
assaying effects of high-sugar diets on adult survival of infection. The six experimental diets used have
the same yeast and cornmeal content, but they varied in sucrose (w/v). Diets with 0%, 2%, and 4% are
represented in shades of blue while diets with 8%, 16%, 24% sucrose are represented in shades of
orange. The rearing diet contains 4% sucrose. (B) Flies fed 8%, 16%, and 24% sucrose diets exhibit

significantly higher mortality after *Providencia rettgeri* infection than flies given 0%, 2%, and 4% sucrose
diets (p<0.0001; Cox Mixed Effects Model). There was no difference in survival among flies provided with
the three high-sugar diets, while flies provided with the 4% sucrose diet exhibited significantly lower
survivorship than flies provided with 0% added sucrose (p = 0.035; Cox Mixed Effects Model). (C) Flies
fed 16% and 24% sucrose diets exhibit significantly higher mortality after *Serratia marcescens* than flies
given diets with 8% sucrose or lower (p<0.0001; Cox Mixed Effects Model). (D) Flies fed the 24% sucrose
diet have the highest mortality after *Lactococcus lactis* infection compared to all other diets (p<0.001, Cox
Mixed Effects Model). Flies fed 16% sucrose died significantly faster than flies fed 0% sucrose diets (p =
0.047; Cox Mixed Effects Model) but there were no other significant pairwise differences in survivorship
among flies provided the 16% sucrose diet or lower. (E) There was no effect of diet on survivorship after
infection with *Enterococcus faecalis*. Letters denote significant pairwise differences between diets
(p<0.05).

Fig. S1. A) Survivorship of PBS injury controls across all infection experiments pooled together.
Flies fed 24% (w/v) sucrose diet have higher proportion of death after sham-infection with sterile
PBS compared to all other diets (p<0.05; Cox Proportional Hazards Mixed effects model).

169
Fig. S2. A) To measure the abundance of the endogenous gut microbiota in uninfected flies fed on
the 2% (w/v) sucrose or 16% sucrose diets, 10 whole flies were pooled and homogenized in 250 µl
of PBS, and 50 µl of the homogenate was plated on lysogeny broth (LB) or brain heart infusion (BHI)
agar. Flies fed on the 2% diets exhibit higher microbiota loads on LB (p<0.001, n = 15 pools of 10
flies per diet) and BHI (p<0.001, n = 15 pools of 10 flies per diet) plates than flies fed 16% sucrose.
C) Axenic flies were infected with *P. rettgeri*, and flies fed 16% sucrose diets exhibited higher
mortality than flies given 2% sucrose (p<0.001; Cox Proportional Hazards model).

Flies fed on high-sucrose diets become hyperglycemic despite eating less

D. melanogaster will modulate feeding behavior depending on the content of their diet (43–

45), and the nutritional composition of the diet influences metabolic state (3,34,46). We

therefore first assessed whether varying sucrose levels in the diet impacts feeding rate. We

performed an excreta quantification assay (ExQ) (47) on uninfected females to measure feeding

across the six experimental diets shown in Figure 1A. Flies were housed in ExQ chambers over

a 2-day period with experimental diets containing the dye erioglaucine (1% w/v) and their

excreta was collected every 24-hours. We found that flies ate significantly more of the three

diets with the lowest sucrose (0%, 2%, and 4%) than the three diets with the highest sucrose

(8%, 16%, 24%; $p < 0.0001$, Fig. 2A). There was no significant difference in feeding rate among

the three lowest-sucrose diets or among the three highest-sucrose diets. In subsequent

experiments, we focus on the 2% and 16% sucrose diets as representative the low- and high-

sugar diets.

To assess the effects of diet on the metabolic status of adults, we measured glucose,

trehalose, glycogen, and soluble protein in the whole body of uninfected flies provided with

either 2% or 16% sucrose diets. We found that uninfected flies fed the high-sucrose diet exhibit

higher levels of glucose ($p < 0.001$, Fig. 2B), trehalose ($p = 0.001$, Fig. 2C), and glycogen (Fig.

2D, $p = 0.0479$) than flies fed the low-sugar diet (Fig. 2B-D). Given that infection can stimulate

shifts in carbohydrate levels (26,48), we also tested whether flies fed high-sugar continue to

exhibit elevated carbohydrate levels after infection. We indeed found that flies fed the high-

sucrose diet exhibited higher levels of glucose ($p<0.01$, Fig. 2B), trehalose ($p < 0.01$, Fig. 2C),

and glycogen ($p = 0.019$, Fig. 2C) than flies fed the low-sucrose diet at 24 hours after *P. rettgeri*

infection. Interestingly, there was no difference in glucose content (Fig. 2B) or trehalose content

(Fig. 2C) between flies fed low-sucrose or high-sucrose at 24 hours after infection with *S.*

marcescens. However, flies given a high-sucrose diet exhibited higher glycogen stores after *S.*

marcescens infection than flies given low sucrose (p = 0.035 Fig. 2D). Despite reduced feeding

on high sugar, flies on the high-sucrose diet still sustain higher levels of carbohydrate stores in

both uninfected and infected states.

205
Fig. 2. Flies fed high sugar sustain higher carbohydrate stores despite reduced feeding. (A) Excreta
quantification of uninfected female flies fed on diets ranging from 0%-24% sucrose was measured
over two days. Flies feeding on the 8%, 16%, and 24% diets consumed significantly lower dye than
those fed on the 0%, 2%, 4% sucrose diets (p < 0.001). (B) Uninfected flies fed on the 16% diet had
higher glucose levels compared to flies fed on 2% sucrose (p < 0.001; ANOVA with Tukey HSD
post-hoc). After 24 hours of *P. rettgeri* infection, flies fed on high sugar continued to exhibit higher
glucose levels compared to flies fed 2% sucrose (p < 0.01; ANOVA with Tukey HSD post-hoc).
However, there was no difference in glucose levels between flies provided with 2% and 16% diets
24-hours after *S. marcescens* infection (p=0.658). Flies fed 16% sucrose exhibit significantly
reduced glucose levels after *S. marcescens* infection (p<0.001), and there was a nearly significant
decrease in glucose levels of flies fed 2% sucrose after *S. marcescens* infection (p = 0.0877) (C)
Trehalose levels were significantly higher in flies fed 16% sucrose than in flies fed 2% sucrose when

the flies were uninfected (p<0.001) or when they were infected with *P. rettgeri* (p<0.01). However, *S.*
marcescens infection had no effect on trehalose levels between flies given the 2% and 16% diets (p
= 0.38). *P. rettgeri* infection led to higher trehalose levels in flies on both 2% (p = 0.0554) and 16%
(p = 0.0446) sucrose diets compared to uninfected flies, while *S. marcescens* had no effect on
trehalose levels on either diet. (D) Glycogen levels were significantly higher in flies on 16% versus
2% diets when flies were uninfected (p = 0.0479), infected with *P. rettgeri* (p = 0.019), or infected
with *S. marcescens* (p = 0.035). On both diets, infection with both pathogens led to reduced
glycogen levels compared to uninfected (2% *Providencia*, p<0.01; 16% *Providencia,* p = 0.036; 2%
Serratia, p<0.01; 16% *Serratia*, p < 0.01). Legend for panel figure: * = p<0.05, ** = p<0.01, *** = p
<0.001.

Feeding on high sucrose increases pathogen load in the early stages of infection

High pathogen load is associated with higher mortality from infection (49,50). Since we

observed that high-sugar diets lead to higher mortality during infection with the Gram-negative

bacteria *S. marcescens* and *P. rettgeri*, we wanted to establish whether this high mortality arises

from higher bacterial load. To test whether flies fed on high sugar have higher pathogen loads,

we performed an *in vivo* bacterial growth assay where we sampled infected flies at two-hour

intervals from 0-16 hours post-infection and at 24, 36, and 48 hours post-infection. At each time

of sampling, individual flies were homogenized and plated on agar plates to estimate the

number of live bacterial cells in the fly.

Overall, the bacteria proliferated more quickly in flies fed on the high-sugar diet after both *P.*

rettgeri (Fig. 3A) and *S. marcescens* (Fig. 3B) infection. However, the divergence in load

between the two diets occurs at different times for the two pathogens. *S. marcescens* infections

start to exhibit a higher pathogen burden as soon as 6 hours into the infection, while *P. rettgeri*

burden becomes higher by 12 hours into the infection. Flies that died during the sampling period

were not included in this assay, which may mean that pathogen burdens are underestimated if

the surviving flies at any given timepoint tend to be the ones with lower pathogen burdens. At 10

hours and later post-infection with *S. marcescens*, we observed a lower proportion of surviving

flies on the high-sugar diet than on the low-sugar diet (Fig. 3B), which may explain the apparent

absence of difference in pathogen burden 10-16 hours into the infection. We similarly observe a

248 lower proportion of flies provided with high-sugar diet surviving *P. rettgeri* infection 10 hours and

249 later, although the difference in pathogen burden between flies on the two diets remains

250 significant (Fig. 3A).

A *Providencia rettgeri*

Sucrose (w/v)
■ 2%
■ 16%

Log2 CFU per Fly

Time (hour)

Proportion of flies alive at each time point

Hour	0	2	4	6	8	10	12	14	16		24	36	48
2%	1.0	0.97	0.93	0.93	1.0	0.93	0.90	0.90	0.97		0.76	0.76	0.71
16%	1.0	0.97	1.0	0.89	1.0	0.83	0.85	0.81	0.59		0.47	0.29	0.29

B *Serratia marcescens*

Sucrose (w/v)
■ 2%
■ 16%

Log2 CFU per Fly

Time (hour)

Proportion of flies alive at each time point

Hour	0	2	4	6	8	10	12	14	16		24	36	48
2%	1.0	1.0	1.0	0.96	0.90	0.93	0.85	0.90	0.83		0.80	0.74	1.0
16%	1.0	1.0	0.93	1.0	0.83	0.69	0.76	0.67	0.63		0.62	0.42	0.44

Fig 3. Pathogens proliferate faster in flies fed high sugar. (A) Pathogen load was assayed at 2-hour intervals from 0-16 hours post-infection and at 24, 36, and 48 hr post-infection in flies fed 2% and 16% sucrose diets. After *P. rettgeri* infection, flies provided with high-sugar diets have higher pathogen loads at 12, 14, 16, 36, and 48 hr than flies given low-sugar diet (12 hr, p =0.031; 14 hr, p <0.001, 16 hr, p< 0.001; 36 hr, p < 0.01; 48 hr, p = 0.022; Generalized Least Square Means (GLS) with Tukey HSD post-hoc for pairwise comparisons). Flies given high-sugar diets have consistently lower survival to time of sampling at hour 10 and onward. (B) After *S. marcescens* infection, flies given high-sugar diets have higher pathogen loads than flies given low-sugar diet at 4 hr, 6 hr, 24 hr, and 36 hr post-infection (4 hr, p = 0.011; 6 hr, p = 0.044, 24 hr, p< 0.001; 36 hr, p < 0.01; Generalized Least Square Means (GLS) with Tukey HSD post-hoc for pairwise comparisons). Flies given high-sugar diets have consistently lower survival to time of sampling at hour 8 and beyond. At 2 hours post-infection, flies given low-sugar diets have higher pathogen load than flies given high-sugar diets (p < 0.01). Each dot represents the pathogen burden of an individual fly. Legend for panel figure: * = p<0.05, ** = p<0.01, *** = p <0.001.

Nutrient availability promotes *S. marcescens* proliferation within hosts on high-sugar diets

Having determined that *D. melanogaster* provided with high-sucrose diets are both more

susceptible to infection and sustain higher internal carbohydrate stores, we next sought to

understand whether the increased susceptibility to infection on the high-sugar diet is due to

impaired immune system activity or to an altered nutritional environment experienced by the

infecting pathogen. To test this, we measured pathogen proliferation during *in vivo* infection of

immune compromised *D. melanogaster* on the low- and high-sugar diets. We predicted that the

effect of diet on sensitivity to infection would be eliminated in immunocompromised flies if the

dietary effect is due to impaired immunity, but it would remain if the dietary effect were due to

altered nutritional environment for the pathogen. We used the *D. melanogaster* strain ΔAMP10

(51), which has intact Toll and IMD signaling but is missing 10 genes encoding major infection-

inducible antimicrobial peptides. Antimicrobial peptides are small, cationic peptides that are

secreted from the insect fat body in response to a systemic infection, and they are critical for

killing pathogens by disrupting function of specific microbial processes (52–55). The ΔAMP10

flies have mutations that eliminate *Diptericin A*, which is critical for controlling *P. rettgeri*

infection (51,56), and the four *Attacin* genes, three of which are critical for the control of *S.*

marcescens infection (P. Nagy, N. Buchon, and B.P. Lazzaro, unpublished). We can use these

flies to monitor pathogen growth in response to diet without any interference from the humoral

immune response. There was no difference in the rate of *P. rettgeri* proliferation in ΔAMP10

hosts provided with either low or high sucrose (Fig. 4A), indicating that the typically observed

effect of diet on sensitivity to this infection may be a consequence of diet-dependent immune

impairment. In contrast, *Serratia marcescens* continued to proliferate faster in ΔAMP10 flies fed

high sucrose than in those fed low sucrose (Fig. 4B), indicating an effect of diet beyond direct

mediation of the immune system.

As an independent, albeit indirect, test of whether *S. marcescens* might utilize excess

available carbon to grow faster in hosts provided with high-sugar diets, we measured *in vitro*

growth curves of *P. rettgeri* and *S. marcescens* in minimal medium supplemented with either a

carbon source or a nitrogen source. We performed an 18-hour growth curve assay in M9

minimal media supplemented with different combinations of casamino acids, sucrose, and

glucose. We found that supplementation with 1% casamino acids enabled *P. rettgeri* growth, but

there was no further change in the growth trajectory or final OD with supplementation with 1%

glucose, 1% sucrose, or no additional sugar (p >0.05, Fig. 4C). These data indicate that *P.*

rettgeri is nitrogen-limited but not carbon-limited in minimal media. The addition of either

glucose or sucrose dramatically increased the growth of *S. marcescens* in minimal media and

resulted in a significantly higher OD at the end of the assay (p<0.01, Fig. 4D), indicating a

greater dependence on environmental carbon under these growth conditions. While we do not

know precise bioavailability of carbon to bacteria infecting *D. melanogaster*, the data in Figure 2

show that flies on high-sugar diets have higher levels of circulating and stored sugars, and in

combination these data offer conceptual support to the hypothesis that the increased

proliferation of *S. marcescens* within hosts provided with a high-sugar diet may arise in part

from increased availability of carbon to the infecting bacteria.

We had observed that flies provided with high-sucrose diets show elevated glucose and

trehalose levels after infection with *S. marcescens* (Fig. 2B, 2C) despite reduction in glycogen

stores in both diets (Fig. 2D), leading us to hypothesize that *S. marcescens* might consume

excess circulating sugar within the host. We measured the change in carbohydrate levels after

infection with either *P. rettgeri* or *S. marcescens* relative to uninfected flies on both diets using

the data in Fig. 2B-D. We found that flies infected with *S. marcescens* have a significantly higher

loss in glucose levels compared to flies infected with *P. rettgeri* on both the low-sugar ($p < 0.01$)

and high-sugar ($p<0.001$, Fig. 4E) diets, although they exhibit similar relative loss in glycogen (p

> 0.05, Fig. 4F). Interestingly, we observed significantly higher loss of glycogen in flies provided

with low-sugar compared to high-sugar diets after both *P. rettgeri* ($p = 0.0178$) and *S.*

marcescens infection ($p = 0.029$, Fig. 4F). There was no significant difference between the

infections in proportional change in trehalose levels on either diet (Fig. 4G). Our observation

that both infections stimulate loss in glycogen stores but only *S. marcescens* infection results in

a significant reduction in glucose levels is consistent with *S. marcescens* consumption of free

sugars during infection.

Fig 4. Nutrient availability promotes *Serratia marcescens* proliferation within hosts on high-sugar
diets. (A) Diet did not affect the proliferation of *P. rettgeri* in flies lacking 10 major inducible
antimicrobial peptides (p> 0.05 time*diet; GLS model). (B) *S. marcescens* proliferate significantly
more rapidly in ΔAMP10 flies fed high sugar (p<0.001 time*diet; GLS model). Two hours into the
infection, flies fed high sugar exhibit a lower *Serratia* burden (p < 0.01; GLS model followed by Tukey
HSD post-hoc), but flies fed high sugar exhibit significantly higher loads by hours by hours 4
(p<0.001) and 6 (p<0.0001; GLS model followed by Tukey HSD-post-hoc). (C) There was no effect of

sugar supplementation on *P. rettgeri* growth over 1080 mins in M9 minimal media supplemented with
1% casamino acids (CAS; p> 0.05; ANOVA followed by Tukey HSD post-hoc). CAS supplementation
significantly increased OD at 1080 minutes compared to media without casamino acids (p<0.001;
ANOVA). (D) *S. marcescens* grew significantly more in minimal media supplemented with the
combination of casamino acids and glucose (p<0.001; ANOVA with Tukey HSD post-hoc) or sucrose
(p < 0.01, ANOVA with Tukey HSD post-hoc) than in media supplemented with casamino acids alone.
S. marcescens OD at 1080 mins was significantly lower in media supplemented only with glucose (p
< 0.001; ANOVA with Tukey HSD). Each curve represents 6-8 replicates per treatment, pooled.
Letters in figures denote significant pairwise differences with p <0.05. Legend: G = glucose, S =
sucrose, C = casamino acids. Colors for *in vitro* curves: 1% glucose (light orange), 1% sucrose (light
blue), 1% casamino acids (black), or 1% glucose & casamino acids (dark orange), and 1% sucrose &
casamino acids (dark blue). (E) Glucose levels declined significantly more after infection with *S.
marcescens* but increased in flies infected with *P. rettgeri* on both 2% (p < 0.01; ANOVA followed by
Tukey HSD post-hoc) and 16% (p< 0.0001; ANOVA followed by Tukey HSD post-hoc) sucrose diets.
There was no significant difference in change in glucose levels after *P. rettgeri* infection between
diets (p = 0.24; ANOVA followed by Tukey HSD post-hoc). There was a trend for increased relative
loss of glucose after *S. marcescens* infections in flies fed the 16% sucrose diet compared to 2%,
however this was not statistically significant (p = 0.0601, ANOVA followed by Tukey HSD post-hoc).
(F) Glycogen levels were reduced after infection with both *P. rettgeri* and *S. marcescens*. Flies fed
2% sucrose had significantly higher loss of glycogen stores compared to flies fed 16% sucrose after
infection with both *P. rettgeri* (p = 0.0178, ANOVA with Tukey HSD post-hoc) and *S. marcescens* (p =
0.0291, ANOVA with Tukey HSD post-hoc). (G) There was no significant difference in change upon
infection in trehalose levels between *P. rettgeri* and *S. marcescens* on either the 2% diet (p = 0.197)
or the 16% sucrose diet (p = 0.215). There was a significant reduction in trehalose levels of flies
infected with *S. marcescens* on 16% sucrose diets compared to flies fed 2% sucrose (p= 0.0372,
ANOVA with Tukey HSD post-hoc). There was a trend toward reduced trehalose in flies provided with
16% sucrose diet after *P. rettgeri* infection, but it was not statistically significant (p = 0.0816, ANOVA
with Tukey HSD post-hoc). Legend for panel figure: * = p<0.05, ** = p<0.01, *** = p <0.001.

Drosophila on high sugar diets produce less AMP peptide after infection

We inferred that high-sugar diets increase sensitivity to *P. rettgeri* infection by impairing the

immune system because the effect of diet was eliminated in immunocompromised flies.

However, the inference that *S. marcescens* may consume excess carbohydrates present in flies

on high-sugar diets does not preclude the possibility that those flies might also have reduced

immune responses. To directly test whether high-sugar diets impair the immune response, we

first measured mRNA transcripts of a panel of genes encoding six *D. melanogaster*

antimicrobial peptides (AMPs). Gene expression was measured in flies fed low or high-sugar

diets at 8-hours after infection with either *S. marcescens* or *P. rettgeri*. We found that diet had

no effect on the infection-induced expression levels of *Diptericin A, Attacin A, Cecropin A1,*

Drosocin, Drosomycin, or *Defensin* after either *P. rettgeri* and *S. marcescens* infection (Figs 5A

and 5B).

Measuring the amount of mRNA transcript does not directly demonstrate the abundance of

translated peptide (57), and translation can be impaired by cellular stress during the *D.*

melanogaster response to infection (58). To test whether high-sugar diets impact AMP protein

levels in response to infection, we used sandwich ELISA quantify Drosocin and Cecropin A1

from flies that have hemagglutinin (HA) and FLAG epitopes tagged to the endogenous AMPs.

Infection-induced production of tagged AMP peptides is detectable as early as 3 hours post-

infection (Fig. S3A, S3B). We quantified tagged AMPs from individual flies at 6-hours after

infection with either *P. rettgeri* or *S. marcescens.* Flies fed on the high-sugar diet produced

significantly lower levels of tagged Drosocin than flies fed on the low-sugar diet after *P. rettgeri*

infection (p = 0.008, Fig. 5C). However, we observed no effect of diet on Drosocin levels in flies

infected with *S. marcescens* (Fig. 5D). An unexpectedly high number of flies did not have

detectable levels of tagged Cecropin peptide after infection with either *P. rettgeri* or *S.*

marcescens (Fig. S4A, S4B). We performed a hurdle model to statistically test for differences in

Cecropin abundance. The hurdle model first tests the probability that detectable levels of tagged

Cecropin are present using a binomial logistic regression, and subsequently tests whether there

are quantitative differences in peptide levels conditional on having observed a detectable level

of peptide. While the probability of detecting Cecropin was equivalent flies infected with *P.*

rettgeri on both diets (Fig. 5E), flies infected with *S. marcescens* had lower probability of

expressing detectable Cecropin when fed the high-sugar diet (p < 0.01, Fig. 5E). In the flies for

which Cecropin was detectable, peptide abundance was significantly lower in the flies on the

high-sugar sucrose diet than in the flies on the low-sugar diet after infection with *P. rettgeri* (p =

0.023, Fig. 5F). Diet had no effect on the abundance of detectable Cecropin after *S.*

marcescens infection (p = 0.21, Fig 5F). Overall, flies provided with 16% sucrose diets produce

less Drosocin and Cecropin after infection with *P. rettgeri* and have a lower probability of

detectable Cecropin 6 hours post-infection with *S. marcescens*. Thus, the translational

production of AMPs seems to be impaired in flies on high-sucrose diets even when there is no

decrease in transcription level.

Fig 5. High sugar diets reduce production of AMP peptides in response to infection without affecting
AMP gene expression. (A) There was no effect of diet on expression of genes encoding the AMPs
Diptericin, Attacin A, Cecropin A1, Drosocin, Drosomycin, and Defensin at 8-hours post-infection with
P. rettgeri or (B) *S. marcescens* (p > 0.05). (C) *P. rettgeri* infection resulted in significantly higher
levels of tagged Drosocin than in uninfected flies on both 2% (p < 0.0001) and 16% (p = 0.0002)
diets. Flies on 16% sucrose diet had significantly lower amounts of tagged Drosocin six hours after *P.*
rettgeri infection than flies on 2% sucrose diet (p = 0.008). (D) Tagged Drosocin levels were higher at
6 hours after *S. marcescens* infection than in uninfected flies on both 2% (p < 0.001) and 16% (p <
0.001) diets, but there was no difference between the diets after *S. marcescens* infection (p = 0.71).
(E) The probability of detectable Cecropin peptide was similar after *P. rettgeri* infection on both diets
(p = 0.45), while flies fed the 2% diet were significantly more likely to have detectable peptide present
after *S. marcescens* infection than those fed the 16% diet (p = 0.003). (F) Of the flies with detectable
Cecropin levels, there were significantly higher peptide levels in flies fed the 2% diet compared to
16% diet after both *P. rettgeri* (p = 0.0229) and *S. marcescens* (p = 0.211) infections.

Fig S3. Demonstration that HA- and FLAG-tagged Drosocin and Cecropin can be detected by
sandwich ELISA following bacterial infection. (A) Tagged Drosocin levels are significantly higher 3-
hours after injection with PBS (p = 0.0455) or ~6.0x10^4 cells of the Gram-negative bacterium
Enterobacter cloacae (p<0.001) than they are in uninjured flies. *E. cloacae* infection results in
significantly higher Drosocin levels than PBS injection (p<0.001; linear model with Tukey post-hoc).
(B) Tagged Cecropin levels are significantly higher 3-hours after infection with *E. cloacae* than after
PBS injection (p<0.001) or in uninjured flies (p<0.001). There was no detectable difference in
Cecropin levels between uninjured and PBS-injected flies (p = 0.32). Letters denote pairwise
difference p<0.05.

A

	Providencia detected	Providencia undetected	Proportion in linear range
2%	13	16	0.45
16%	11	21	0.34

B

	Serratia detected	Serratia undetected	Proportion in linear range
2%	21	9	0.70
16%	8	19	0.30

Fig. S4. (A) The number of flies fed the 2% and 16% diets that have, or do not have, detectable
tagged Cecropin six hours after *P. rettgeri* infection. (B) The number of flies fed the 2% and 16%
sucrose diets that have, or do not have, detectable tagged Cecropin six hours after *S. marcescens*
infection.

Discussion

In this study, we show that *D. melanogaster* provided with high-sugar diets exhibit higher

mortality and elevated pathogen load after infection with the Gram-negative bacteria

Providencia rettgeri and *Serratia marcescens*, even when the high-sugar diet is provided only to

the adult stage after development is complete. *S. marcescens* may be able to take a growth

advantage from the excess carbohydrates available in hosts provided with high-sugar diets, and

the effect of diet on sensitivity to both infections seems to be mediated by impaired host immune

response on high-sugar diet. Interestingly, we saw no effect of diet on susceptibility to infection

with the Gram-positive bacteria *Enterococcus faecalis* and *Lactococcus lactis*. As Gram-

negative (*S. marcescens* and *P. rettgeri*) infections should primarily activate the IMD signaling

pathway whereas Gram-positive (*E. faecalis* and *L. lactis*) infections should primarily activate

the Toll pathway (50,59), the much larger effect of diet on sensitivity to Gram-negative infections

may implicate an interaction with the IMD pathway.

Our approach of feeding flies on diets that range from 0%-24% sucrose was based on

previous studies of systemic infection outcome that defined "high" dietary sugar as ranging from

10% glucose (3) to 24% (w/v) sucrose (12). We predicted that there would be a concentration-

responsive decrease in host survival post-infection as dietary sugar increased. However, what

we observed was a threshold response where quantitatively variable low-sucrose diets (0%, 2%

and 4%) were statistically equivalent to each other but different than high-sucrose diets (16%

and 24%) with respect to host survivorship of infection. However, the precise threshold

depended on the pathogen, with 8% sucrose being a functionally high-sugar diet for a *P. rettgeri*

infection but low-sugar for a *S. marcescens* infection.

Our observation of immunological effects of diet that vary among pathogens is consistent

with previous reports in the literature. For example, Ayres and Schneider (2009) (60)

demonstrate that dietary restriction at 24 hours prior to infection of adult *D. melanogaster* is

protective against a *Salmonella typhimurium* infection, decreases resistance to *Listeria*

monocytogenes, and has no effect on *E. faecalis* infection (60). Similarly, Singh et al. (2022)

(61) found that *D. melanogaster* reared on a calorie-restricted diet were more susceptible to

Pseudomonas entomophila infection while diet had no effect on mortality after *E. faecalis*

infection. Our finding that diet had no effect on *E. faecalis* infection outcome is corroborated by

several studies (60–62) that suggest negligible impact of diet or relevant host physiology on

defense against this pathogen. Our data in combination with these previous studies suggest that

the effects of diet on immune outcome are in part dependent on the unique physiology of the

infecting pathogen.

We observed that *S. marcescens* and *P. rettgeri* pathogen burdens are both higher in *D.*

melanogaster fed high sugar. However, the increased pathogen load arises differently for the

two pathogens. The effect of diet on defense against *P. rettgeri* is mediated by the host immune

system, as flies provided with 16% sucrose diet produce less Drosocin after infection with *P.*

rettgeri and the effect of dietary sugar on *P. rettgeri* proliferation is eliminated in *D.*

melanogaster lacking a functional immune response. However, while flies infected with *S.*

marcescens produced less Cecropin when provided with 16% sucrose diet, we also observed

that *S. marcescens* continued to proliferate more rapidly in flies provided with 16% sucrose diets

even when those flies lack most inducible AMPs. We found that high-sugar diet increases the

level of circulating sugar within the fly, and that *S. marcescens* benefits more than *P. rettgeri*

from carbon supplementation when growing *in vitro.* These data suggest *S. marcescens* may

perhaps gain a growth benefit from the hyperglycemia of *D. melanogaster* feeding on high-sugar

diets. A similar phenomenon has been observed with the Gram-negative opportunistic pathogen

Pseudomonas aeruginosa infecting hyperglycemic mice. Wild-type *P. aeruginosa* proliferates

more quickly in hyperglycemic mice, but mutants for glucose uptake and metabolism genes

(*oprB, gltK, gtrS*, *glk)* exhibit similar growth in control and hyperglycemic mice (6). More detailed

future study of *S. marcescens* mutants deficient for glucose uptake or metabolism infecting *D.*

melanogaster mutant for key factors in metabolism and immune response could probe the

relationship between diet, host physiology, and pathogen physiology as determining infection

outcome.

Interestingly, we noticed that both wild-type and ΔAMP10 *D. melanogaster* genotypes had

lower *S. marcescens* burdens at 2 hours after infection on high-sugar diet than on low-sugar

diet (Figs. 3B and 4B). Phagocytosis is known to play a role in controlling *S. marcescens*

infection in some contexts (63) and infection has previously been shown to increase aerobic

glycolysis in *Drosophila* phagocytes (64). Since ΔAMP10 flies still have intact cellular immune

systems, it is possible that the elevated dietary sugar provides energy for phagocytes to use in

clearing pathogens sensitive to phagocytosis, although this process alone is not sufficient to

manage the infection delivered in our experimental context.

We found that elevated dietary sugar does not impact the transcription of antimicrobial

peptides genes (Fig. 5A, 5B), but it does reduce translation of the encoded peptides Drosocin

and Cecropin A1 (Fig. 5C-5E). A similar transcriptional effect was observed in larvae of the beet

army worm (*Spodoptera exigua*) and fall army worm (*Spodoptera frugiperda*), where infection-

induced expression of the AMPs *attacin* and *gloverin* transcripts were equivalent between a low

protein and control diet despite reduced resistance to infection on the low-quality diet (65). This

suggests that dietary impacts on the immune system may not occur at the transcriptional level

but at the level of translation. A recent proteomics study on *D. melanogaster* found that naive

flies subjected to dietary restriction exhibit lower peptide levels of the AMPs Attacin A and

Cecropin C (66), although those authors did not measure the effect of diet on production of

antimicrobial peptides in the context of a systemic infection.

Antimicrobial peptides are critical for controlling Gram-negative infections in *D.*

melanogaster. At the transcriptional level, there is a high degree of co-regulation of genes

encoding AMPs (19,67), and we would anticipate corresponding similarity in translational

efficiency (68). Given that high dietary sugar reduced production of Drosocin and Cecropin, we

anticipate it would also reduce production of other AMPs. It should also be noted that the IMD

and Toll immune regulatory pathways do more than simply stimulate production of AMPs.

Activation of these pathways alters host metabolism and is influenced by host metabolic status

(13), and even stimulates host sequestration of micronutrients to limit pathogens from accessing

them (69). Future experiments will be necessary to establish the mechanistic details of how

high-sugar diets impact the IMD and Toll pathways to shape resistance to infection.

Our present work establishes that adverse effects of high-sugar diets on infection outcome

are due to the combination of impaired host immune function as well as, in some cases, faster

proliferation of the infecting pathogen in a hyperglycemic environment. Notably, the immune

impairment is not necessarily at the level of transcriptional activation of the immune response

but appears to be due to reduced translation of antimicrobial peptides. It is also notable that

dietary effects on immune defense are pathogen-specific, and there is not one universal

immunological consequence of high-sugar diets. Our findings illustrate the importance of

understanding the nutritional requirements of the infecting pathogens as well as the

physiological intertwining of host metabolism and immune responses when interpreting effects

of diet on infection outcome. We expect this principle to be general across host-pathogen

infection systems.

Methods

Fly stocks and husbandry

All flies were reared and maintained at 25 °C on a 12-hour light/dark cycle. Mated females of

the wild-type strain Canton S (CS) were used for all experiments unless otherwise indicated.

Females were cohoused with males in culture chambers and were aged to 3-5 days post-

eclosion prior to infection for all experiments. Flies were anesthetized on CO_2 pads for collection

and during infection.

Generation of Tagged AMP Flies

Flies carrying tagged Cecropin A1 or Drosocin were generated using CRISPR-facilitated

homology-directed repair. In brief, guide sequences for *CecA1* or *Dro* were cloned into pCFD5

(*CecA1* guides: guide 1: CCATTGGACAATCGGAAGCT; guide 2:

ATAATTATAAATAATCATCG; *Dro* guides: guide 1: CAAAAACGCAAGCAAGCAGC; guide 2:

CAATCAATTGTGACACAATG) (70). Homology-directed repair cassettes were cloned into pHD-

Scarless-dsRed, encoding the replacement of endogenous *CecA1* and *Dro* with peptides that

included FLAG and HA tags (predicted sequences of mature peptides: CECA1[HF],

GWLDYKDDDDKKIGKKIERVGQHTRDATIQGLGIAQQAANVAATYPYDVPDYAR; DRO[HF],

GDYKDDDDKPRPYSPRPTSHPRPIYPYDVPDYARV) (71). These two plasmids were then

injected together into embryos of the genotype *y¹ P(nos-cas9, w-) M(RFP-.attP)ZH-2A w**. Male

flies emerging from this injection were crossed with females of the genotype *w¹¹¹⁸*; *If / SM6a* or

w¹¹¹⁸; *TM2 / TM6c, Sb¹*, as appropriate. Males expressing dsRed in the eye were then selected

and crossed to flies carrying *CyO, Tub-PBac* to enable excision of the 3xP3-dsRed cassette.

The final structure of each locus was verified by PCR.

Experimental Diets

All flies were reared from egg to pupation in rearing chambers with interchangeable food plates

on a standard base sucrose-yeast-cornmeal diet (per liter: 60 g yeast, 60 g cornmeal, 40 g

sucrose, 7 g *Drosophila* Agar, 0.5 mL phosphoric acid, 5 mL propionic acid, and 6 g

methylparaben dissolved in 26.5 mL 100% ethanol). *D. melanogaster* pupated on the chamber

walls and, prior to eclosion, the food plates containing the standard rearing media were

swapped with one of the experimental diets. Emerging adult flies were aged on the experimental

media for 3-5 days post-eclosion prior to infection for all experiments. Experimental diets were

identical to the rearing diet aside from the concentration of sucrose. Experimental diets

contained either 0%, 2%, 4%, 8%, 16%, and 24% (w/v) of sucrose.

Infection Survival Assays

Flies were anesthetized on CO_2 and injected in the thorax with either a bacterial suspension or

sterile PBS (as injury control) using a pulled glass capillary needle and nano-injector (72). The

following bacteria were injected in at least four experimental blocks: *Providencia rettgeri* strain

Dmel, *Serratia marcescens* strain 2698B, *Enterococcus faecalis* strain Dmel*, Lactococcus lactis*

strain Dmel*.* All bacteria were originally isolated as natural infections of *D. melanogaster* (36,73)

except *S. marcescens* 2698B, which is a clinical isolate (74). Cultures for *P. rettgeri* and *S.*

marcescens were started from a frozen glycerol stock then cultured overnight in liquid lysogeny

broth (LB) at 37 °C on a shaker, then subcultured the next morning in fresh LB to achieve

growth phase. *L. lactis* was cultured overnight from a frozen glycerol stock in liquid Brain Heart

Infusion (BHI) at 37 °C with shaking, then subcultured the next morning in fresh BHI to achieve

log-phase. *E. faecalis* was cultured from a frozen glycerol stock at room temperature (~24 °C) in

BHI. All bacteria were resuspended in 1x phosphate-buffered saline (PBS), diluted to $A_{600} = 0.1$,

and injected at a volume of 23 nL per fly. There were ~4000 *P. rettgeri* cells, ~3000 *S.*

marcescens cells, ~1000 cells *E. faecalis* cells injected, or ~ 1000 *L. lactis* cells injected per fly.

Flies were housed in groups of ten and each experimental block contained 3-4 vials of flies per

experimental diet. Survival curves were assessed using a Cox proportional hazards mixed effect

model (coxme) with the "CoxMe" package in R.

Model A: coxme(Surv(status, time) ~ diet + (1|block)

Diet was treated as a main effect and experimental block was treated as a random factor in our

model. This model was used to perform all pairwise comparisons using the "emmeans" package

in R with a Tukey p-value correction ($p < 0.05$).

Bacterial Load Trajectory Assay

To assay bacterial proliferation during *in vivo* infection, flies were infected with *P. rettgeri* or

S. marcescens as described above, then collected for measurement of bacterial load at 0, 2, 4,

6, 8, 10, 12, 14, 16, 24, 36, and 48 hours post-infection. Single, live flies were homogenized in

500 µL sterile 1x-PBS with a metal bead and 50 µL of the homogenate was plated onto LB agar

using a spiral plater (Whitley Automated Spiral Plater-2). To ensure there was no overgrowth of

bacteria on the plates, fly homogenates were diluted with PBS at time points where we

anticipated higher bacterial growth. For wild-type CS flies, fly homogenates from timepoints 0, 2,

and 4 hours were directly plated, homogenates from timepoints 6-12 were diluted 1:10, and

homogenates from timepoints 14-48 were diluted 1:100. For ΔAMP10 mutants, fly homogenates

were directly plated for time points 0, 2, and 4, diluted 1:10 at hour 6, and 1:100 at time points 8,

10 and 12. Agar plates were incubated at 37°C overnight. The resulting colonies were counted

using PROTOCOL3 software to estimate the number of colony forming units (CFU) per mL of fly

homogenate. We then used the CFU/mL value to approximate the number of bacteria in the

individual fly at the time of sampling by using the following formula:

CFU/fly = CFU/mL * (dilution factor) * (0.5 mL/fly of original homogenate)

Two experimental blocks were completed for *P. rettgeri* and *S. marcescens* in CS flies, and

three experimental blocks were completed for ΔAMP10 mutants with each pathogen. A

generalized least squares (GLS) model from the "nlme" package in R was used to determine the

effect of diet on pathogen loads over time. Fixed effects in the model include time point sampled

(T) and diet (D). The function weights = varIdent(form = ~1|time point) from the "nlme" package

was incorporated into the model to account for unequal variances across time points, which

arises as the pathogen grows from the initial injection time point. The emmeans function was

used for pairwise comparisons between the experimental diets within each time point using the

Tukey method to determine p-values ($p<0.05$)

Model B: gls($\log_2$(pathogen load) ~ T * D, weights = varIdent(form = ~ 1|T))

Excreta Quantification (ExQ) to measure feeding

Methods for the ExQ assay are further detailed in Wu (2020) (47). Briefly, food was prepared by

melting down experimental diets and adding 1% (w/v) of dry erioglaucine powder (Sigma #

861146). The dyed food mixture was then dispensed into microcentrifuge tubes caps filled up to

50 µL volume. ExQ chambers were prepared from 50 mL conical vials (Cell Treat # 229421). Air

holes were poked along the sides of tubes and the lid using a pushpin (~0.5 mm diameter). A

precision knife was used to cut a hole in the lid of the conical vial to fit the diameter of the food

caps. Naïve flies were anesthetized with CO_2 and sorted into ExQ chambers in groups of 8-10.

Every 24 hours, the flies were transferred to new chambers with fresh food. After 48 hours, flies

were emptied from chambers and 3 mL of 1x-PBS was used to collect the excreta. Total dye

concentration was determined by using a spectrophotometer to measure absorbance of the dye

excreted from flies. A standard curve was prepared from a stock solution of erioglaucine by

dissolving 10 mg of powder dye in 10 mL of PBS. A serial 2-fold dilution was then created from

this initial stock, and the final range of standards used to create the standard curve was 312.5

µg/mL to 5 µg/mL. Absorbance was measured at 630 nm in a 96-well plate on a spectrometer

(Molecular Biosciences Spectra Max Series Plus 384). Standards and samples were run in

duplicates with 100 µL per well and then averaged for data analysis. Dye consumed per fly

within a single ExQ chamber was determined using the following equation:

Dye (µg) per fly = [(Absorbance – Slope)]/(Intercept * # flies)

A two-way Analysis of Variance (ANOVA) in R was used to test for differences in dye

consumed by flies on each diet on Day 1 and Day 2 of feeding. Then the emmeans function in R

was used for pairwise comparisons between diets within each day using the Tukey test to

determine p-values (p<0.05).

Model C: aov(dye ~ diet + day)

Quantitative real-time PCR (RT-qPCR) for Antimicrobial Peptide Gene Expression

Female flies provided with 2% and 16% sucrose diets were injected with ~3000 CFU of log-

growth *S. marcescens* or *P. rettgeri* then sampled at 8 hours after injection for quantification of

gene expression. Flies were pooled in groups of 5 and frozen at -80 °C for subsequent RNA

extraction. In 1.5 mL microcentrifuge tubes, RNA was isolated and purified using a TRIzol

extraction procedure according to the manufacturer's instructions (Zymo Kit direct-Zol RNA kit

#R2050). Briefly, flies were homogenized in TRIZoL (Invitrogen #15596018) using a motorized

pestle in a microcentrifuge tube and RNA was then extracted using the Zymo Spin Column

(Zymo Research #C1078). Samples were then treated with the kit's DNAse and RNA was

dissolved 50 µL of molecular grade water. RNA concentration was determined using a Qubit4

Broad Sense RNA kit (Thermo #Q10210). cDNA was generated using 500 ng of total RNA with

the iSynthesis cDNA kit (Bio-Rad #1708891). RT-qPCR was performed on a CFX Connect

Real-Time Detection System (Bio-Rad) using PerfeCTa SYBER Green fast mix (VWR #101414-

270). Expression of the AMPs *Drosocin, Defensin, CecropinA, Dipterin, Attacin A, Drosomycin*

was normalized to the expression of the housekeeping gene *Actin 5C*. Uninfected controls were

not subject to injury. Primer sequences used are listed in Supplementary Table 1. A linear

model was used to test the contributions of diet and infection status to expression of each AMP

gene, standardized to the housekeeping gene (75).

Model D: lm(AMP_CT ~ *Actin5C*_CT + Diet + Infection Status * Diet)

Diet and Infection Status were fixed effects in the model. Estimated marginal mean CT values

for the Diet x Infection interaction terms were extracted using the emmeans package in R. Then

we subtracted the estimated marginal mean CT value for uninfected flies from the estimated

marginal means from the infected flies for each diet. This difference in estimated marginal mean

CT between uninfected and infected values indicates infection-induced change in AMP gene

expression on a $\log_2$ scale. Using the emmeans function, we compared infection-induced

expression values between experimental diets with Tukey's corrections test for p-values

($p<0.05$).

Gene	Forward Primer	Reverse Primer
Actin-5C	AGCGCGGTTACTCTTTCACCAC	GTGGCCATCTCCTGCTCA AAGT
Attacin A	CGTTTGGATCTGACCAACG	AAAGTTCCGCCAGGTGTGAC
CecropinA1	CTCTCATTCTGGCCATCACC	TCTTGAGCGATTCCCAGTC
Defensin	GCGGATCATGTCCTGGTGCAT	TCGCTTTGGCGGCTATG
Diptericin A	GCGGCGATGGTTTTGG	CGCTGGTCCACACCTTCTG
Drosocin	TTTTCCTGCTGCTTGCTTGC	GGCAGCTTGAGTCAGGTGAT
Drosomysin	CTGCCTGTCCGGAAGATACAA	TCCCTCCTCCTTGCACACA

Supplementary Table 1: Primer sequences for AMP and housekeeping genes ('5-3').

Sandwich ELISA to quantify tagged AMPs

HA/FLAG tagged AMPs were quantified 6 hr post-infection. The fly line w[1118] ; Dro[HF]/Dro[Hf]

was used to quantify tagged Drosocin and w[1118] ; Cec[HF]/+ was used to quantify tagged

Cecropin. Maxisorp Nunc 96-well ELISA plates (Thermo Scientific 232698) were coated with 2.5

µg/mL of anti-FLAG antibody (Sigma-Aldrich F1804) in coating buffer (0.2 M sodium

carbonate/bicarbonate buffer) overnight at 4 °C, then the plates were blocked for 2 hours at

room temperature with 2% bovine serum albumin (Sigma-Aldrich A2153) prior to use. At the

time of sampling, flies were homogenized in 200 µL of 0.2% Triton-X buffer and centrifuged at

10,000 RPM for 5 mins to pellet fly debris. Drosocin samples were diluted 1:20 and Cecropin

samples were diluted 1:10 so that they fell within linear range of a HA/FLAG peptide standard

curve (sequence NH2-DYKDD DDKGG GGGSY PYDVP DYA-NH2 made from AAPTec). The

standard curve was prepared by performing a 2-fold serial dilution starting from a stock solution

of 2 ng/mL of HA/FLAG peptide to yield 14 standard concentrations. For sample incubation, 50

µL of each standard and sample were added to the ELISA plate and then incubated overnight.

Any unused wells were filled with 50 µL of 0.2% Triton-X buffer to serve as blanks. The next

day, anti-HA diluted 1:5000 in 0.2% Triton-X was added to each well for another overnight

incubation at 4 °C. The following morning, 100 µL of room temperature 1-Step Ultra TMB-ELISA

Substrate Solution (ThermoFisher #34028) was added to each well and the plate was incubated

at 25 °C for 30 min. After incubation, 100 µL of 2M sulfuric acid was added to each well to stop

the reaction. Plates were then read on a spectrometer at 450 nm to measure absorbance. The

amount of peptide per fly was determined from the HA/FLAG standard curve with the following

equation:

Peptide per fly = [(Absorbance-Slope) * Dilution Factor]/ (Intercept * Fly Homogenate)

To determine whether diet affects Drosocin peptide levels, a linear mixed effects model was

applied, with diet and infection status as main factors and experimental block as a random

factor. The emmeans function in R was used to compare peptide yield from flies on each diet

within infection status.

Model E: lmer(Dro_levels ~ diet * infection + diet + (1|block))

To determine effects of diet on Cecropin peptide levels, a hurdle model was applied. In the first

step, a binomial generalized linear model (GLM) was used to test whether diet and infection

status (*Serratia* or *Providencia* infection versus uninjured) affected the probability of detecting

Cecropin peptide in the samples.

Model F: glm(Cec_present ~ diet * infection + diet, family = binomial)

In the second step, a GLM was performed on samples with detectable peptide to test

whether diet and infection resulted in quantitative differences in peptide abundance.

Model G: glm(Cec_level ~ diet *infection + diet)

The significance of the contrast between diets within each infection was determined with a

post-hoc Tukey test (p<0.05) using the emmeans package in R.

Measurement of nutritional indices

Nutritional indices were measured in naïve flies and in flies 24 hours post-infection with

either *P. rettgeri* or *S. marcescens.* Flies were homogenized in pools of three in 150 µL of cold

lysis buffer (10 mM Tris, 1 mM EDTA, pH 8.0 with 0.1% (v/v) Triton-X-100) with a motorized

pestle in 1.5 mL microcentrifuge tubes. Homogenates were centrifuged for 1 minute at 13,000

rpm at 4°C to pellet fly debris. The supernatants were transferred to fresh 1.5ml microfuge

tubes, and 10 µL of homogenized sample was removed for protein quantification. The remaining

samples were placed in a 72°C water bath for 20 minutes to denature endogenous enzymes.

The tubes were stored at -80 °C for later use.

Protein measurements: Protein was measured to normalize all carbohydrate measures. Non-

heated fly homogenates were diluted to 1:8 in lysis buffer. Protein was quantified according to

Bio-Rad Assay DC Protein Assay Kit instructions. Briefly, 5 µL of standard and sample were

incubated with 25 µL reagent A and 200 µL of reagent B. Bovine serum albumin was serially

diluted to generate a standard curve that was then used to determine the quantity of protein in

each sample. Samples and standards were all run in duplicates. Samples were incubated at

room temperature for 15 minutes and then read on a spectrometer plate reader at 750 nm.

Glucose, trehalose, and glycogen level measurements: Glucose, trehalose, and glycogen

levels were measured using reagents from the Sigma Aldrich Glucose (GO) Assay Kit

(GAGO20-1KT) and processed according to manufacturer's instructions. The same fly

homogenates were used across all nutrients assayed. Glucose was measured by mixing 5 µL of

standard or 5 µL of sample with 150 µL of glucose assay reagent. A glucose standard curve

was generated from 2-fold serial dilution from a 1 mg/mL glucose stock concentration. Each

sample and standard were run in duplicate. The plate was incubated for 30 minutes at 37°C,

then 150 µL of 12M sulfuric acid was added to each well to stop the reaction. Plates were then

measured for absorbance at 544 nm on a spectrometer. For trehalose measurements, 5 µL of

sample and standards were incubated with 2.5 µL of trehalase overnight at 37 °C. For each

sample, two replicates were treated with trehalase enzyme and two were not. Trehalose

standard curves were generated from a glucose standard curve, trehalose standards treated

with trehalase, and trehalose standards without enzyme. Glucose liberated from trehalose

digestion was quantified the next day using the procedure described above, and the abundance

of trehalose in the standard was inferred from the difference between the samples that were

treated with trehalase and those that were not. Glycogen measurements were conducted

similarly, except 5 µL of standard and 5 µL of sample were incubated overnight with

amylogucosidase instead of trehalase. The glycogen standard curve was prepared from a

glycogen stock solution over the range 1 mg/mL to 0.1 mg/mL. All estimates of carbohydrate

levels were normalized to measured protein level for each sample to correct for potential

differences in the size of the fly. To calculate change in metabolite level after infection, we

subtracted the average post-infection value of the metabolite level from the average level in

uninfected samples within an experimental block. A linear mixed-effects model was performed

to test for the main effects of diet and infection on carbohydrate/protein ratios with experimental

block as a random factor. The emmeans package was used to perform a post-hoc Tukey's test

to determine the significance of pairwise comparisons between diet and infection ($p<0.05$).

Bacteria growth curves *in vitro*

Serratia marcescens and *Providencia rettgeri* were measured for *in vitro* growth using the

kinetic cycle in a spectrometer (Molecular Biosciences Spectra Max Series Plus 384) at room

temperature. Bacteria were first cultured overnight with shaking in liquid LB at 37°C. The next

morning, bacterial cells were resuspended in 1X M9 minimal media (M9) to achieve $A_{600} = 1.0$,

then further diluted to $A_{600} = 0.20$ in one of six different experimental M9 media conditions. The

combinations of M9 media used were 1% glucose, 1% sucrose, 1% casamino acids (CAS), 1%

casamino acids and 1% glucose, 1% casamino acids and 1% sucrose, and a M9 blank. CAS

was added to the media to provide a nitrogen source while glucose and sucrose provide a

carbon source. In a sterile 96-well plate with a clear flat bottom, 100 µL of bacteria resuspended

in the experimental media were pipetted into individual wells in 3-fold replicate. Two wells of

uninoculated experimental media were included to monitor for contamination. The plate was

covered with a lid and inserted into the spectrometer to measure A_{600} at 15-minute intervals over

18 hours at 37°C. We used a linear mixed effect model to test for the main effects of CAS and

sugar with block as a random factor to test whether the addition of CAS and/or sugar led to

increased bacterial growth, measured as a higher final A_{600}. A post-hoc Tukey HSD was

performed using the emmeans package in R to test pairwise comparisons ($p<0.05$) between

media containing sugars and CAS.

Model H: lmer(OD600 at 18 hours ~ sugar *CAS + sugar + (1|block))

Preparation of Axenic Flies

Axenic flies were prepared by embryo bleaching. Conventionally reared (microorganism

associated) flies were fed on lab standard yeast cornmeal diet (4% sucrose w/v), and

approximately 100-150 flies were placed into an egg laying chamber and allowed to lay eggs on

grape juice agar plates. Embryos were collected after 20 hours and removed from agar plates

with a clean paint brush into a cell strainer. In a six-well tissue culture plate, embryos were

washed within the cell strainer first in 10% bleach twice for 1.5 minutes each, followed by two

30-second washes in 70% ethanol. Lastly, embryos were washed twice for 10 seconds with

autoclaved 1X PBS. Embryos were continuously mixed throughout the process to prevent them

from sticking to each other or the cell strainer. Once sterilized, embryos were transferred with a

sterile brush onto autoclaved yeast-cornmeal rearing media in groups of 50. During pupation,

the rearing media was replaced with autoclaved 2% and 16% sucrose diets for adult flies to

eclose onto.

Acknowledgements

We thank all members of the Lazzaro lab for feedback on the experimental design for this

project and for comments on the manuscript. Figure 1A was designed using biorender.com.

References

1. Podell BK, Ackart DF, Kirk NM, Eck SP, Bell C, Basaraba RJ. Non-diabetic hyperglycemia exacerbates eisease severity in *Mycobacterium tuberculosis* infected guinea pigs. PLOS ONE. 2012 Oct 4;7(10):e46824.

2. Howick VM, Lazzaro BP. Genotype and diet shape resistance and tolerance across distinct phases of bacterial infection. BMC Evol Biol. 2014 Mar 22;14(1):56.

3. Unckless RL, Rottschaefer SM, Lazzaro BP. The complex contributions of genetics and nutrition to immunity in *Drosophila melanogaster.* PLoS Genet. 2015;11(3):1005030.

4. Kutzer MAM, Armitage SAO. The effect of diet and time after bacterial infection
on fecundity, resistance, and tolerance in *Drosophila melanogaster.* Ecology and Evolution.
2016 Jul 1;6(13):4229–42.

5. Cotter SC, Reavey CE, Tummala Y, Randall JL, Holdbrook R, Ponton F, et al. Diet
modulates the relationship between immune gene expression and functional immune
responses. Insect Biochem Mol Biol. 2019 Jun;109:128–41.

6. SK, Hui K, Farne H, Garnett JP, Baines DL, Moore LS, Holmes AH, Filloux A, Tregoning JS.
Increased airway glucose increases airway bacterial load in hyperglycaemia. Sci Rep. 2016
Jun 8;6:27636. doi: 10.1038/srep27636.

7. Thaiss CA, Levy M, Grosheva I, Zheng D, Soffer E, Blacher E, et al. Hyperglycemia drives
intestinal barrier dysfunction and risk for enteric infection. Science. 2018 Mar
23;359(6382):1376–83.

8. Gupta S, Koirala J, Khardori R, Khardori N. Infections in diabetes mellitus and
hyperglycemia. Infectious Disease Clinics of North America. 2007 Sep 1;21(3):617–38 vii.
doi: 10.1016/j.idc.2007.07.003.

9. Lecube A, Pachón G, Petriz J, Hernández C, Simó R. Phagocytic activity is impaired in
Type 2 diabetes mellitus and increases after metabolic improvement. PLoS ONE. 2011;6(8).

10. Ilias I, Zabuliene L. Hyperglycemia and the novel Covid-19 infection: Possible
pathophysiologic mechanisms. Med Hypotheses. 2020 Jun;139:109699.

11. van Vught LA, Wiewel MA, Klein Klouwenberg PMC, Hoogendijk AJ, Scicluna BP, Ong
DSY, et al. Admission hyperglycemia in critically ill sepsis patients: Association With
Outcome and Host Response*. Critical Care Medicine. 2016 Jul;44(7):1338–46.

12. Musselman LP, Fink JL, Grant AR, Gatto JA, Tuthill BF, Baranski TJ. A complex relationship
between immunity and metabolism in *Drosophila* diet-induced insulin resistance. Molecular
and Cellular Biology. 2017 Oct 30;38(2).

13. Darby AM, Lazzaro BP. Interactions between innate immunity and insulin signaling affect
resistance to infection in insects. Frontiers in Immunology. 2023;14. Available from:
https://www.frontiersin.org/articles/10.3389/fimmu.2023.1276357

14. Buchon N, Silverman N, Cherry S. Immunity in *Drosophila melanogaster* — from microbial
recognition to whole-organism physiology. Nature Reviews Immunology. 2014
Dec;14(12):796–810.

15. Bland ML. Regulating metabolism to shape immune function: Lessons from *Drosophila.*
Seminars in Cell & Developmental Biology. 2023 Mar 30;138:128–41.

16. Piper MD. Using artificial diets to understand the nutritional physiology of *Drosophila
melanogaster.* Current Opinion in Insect Science. 2017 Oct 1;23:104–11.

17. Imler JL, Zheng L. Biology of Toll receptors: lessons from insects and mammals. Journal of
Leukocyte Biology. 2004 Jan 1;75(1):18–26.

18. Paquette N, Broemer M, Aggarwal K, Chen L, Husson M, Ertürk-Hasdemir D, et al. Caspase
mediated cleavage, IAP binding and ubiquitination: Linking three mechanisms crucial for
Drosophila NF-κB signaling. Mol Cell. 2010 Jan 29;37(2):172.

19. Troha K, Im JH, Revah J, Lazzaro BP, Buchon N. Comparative transcriptomics reveals
CrebA as a novel regulator of infection tolerance in *D. melanogaster.* PLoS pathogens.
2018;14(2):e1006847.

20. Valanne S, Wang JH, Rämet M. The *Drosophila* Toll signaling pathway. The Journal of
Immunology. 2011 Jan 15;186(2):649–56.

21. Myllymäki H, Valanne S, Rämet M. The *Drosophila* Imd signaling pathway. The Journal of
Immunology. 2014 Apr 15;192(8):3455–62.

22. Tanji T, Hu X, Weber ANR, Ip YT. Toll and IMD pathways synergistically activate an innate
immune response in *Drosophila melanogaster.* Mol Cell Biol. 2007 Jun;27(12):4578–88.

23. De Gregorio E. The Toll and Imd pathways are the major regulators of the immune
response in *Drosophila.* The EMBO Journal. 2002 Jun 3;21(11):2568–79.

24. Martínez BA, Hoyle RG, Yeudall S, Granade ME, Harris TE, Castle JD, et al. Innate immune
signaling in *Drosophila* shifts anabolic lipid metabolism from triglyceride storage to
phospholipid synthesis to support immune function. PLOS Genetics. 2020 Nov
23;16(11):e1009192.

25. Dionne MS, Pham LN, Shirasu-Hiza M, Schneider DS. Akt and foxo dysregulation contribute
to infection-induced wasting in *Drosophila.* Current Biology. 2006 Oct 24;16(20):1977–85.

26. Chambers MC, Song KH, Schneider DS. *Listeria monocytogenes* infection causes
metabolic shifts in *Drosophila melanogaster.* PLOS ONE. 2012 Dec 13;7(12):e50679.

27. Davoodi S, Galenza A, Panteluk A, Deshpande R, Ferguson M, Grewal S, et al. The
immune deficiency pathway regulates metabolic homeostasis in *Drosophila.* The Journal of
Immunology. 2019 May 1;202(9):2747–59.

28. DiAngelo JR, Bland ML, Bambina S, Cherry S, Birnbaum MJ. The immune response
attenuates growth and nutrient storage in *Drosophila* by reducing insulin signaling.
Proceedings of the National Academy of Sciences of the United States of America. 2009
Dec 8;106(49):20853–8.

29. Nunes RD, Drummond-Barbosa D. A high-sugar diet, but not obesity, reduces female
fertility in *Drosophila melanogaster.* Development. 2023 Oct 16;150(20):dev201769.

30. Hemphill W, Rivera O, Talbert M. RNA-sequencing of *Drosophila melanogaster* head tissue
on high-sugar and high-fat diets. G3: Genes, Genomes, Genetics. 2018 Jan 1;8(1):279–90.

31. Ma X, Nan F, Liang H, Shu P, Fan X, Song X, et al. Excessive intake of sugar: An
accomplice of inflammation. Frontiers in Immunology [Internet]. 2022 [cited 2023 Apr 25];13.
Available from: https://www.frontiersin.org/articles/10.3389/fimmu.2022.988481

32. Yu S, Zhang G, Jin LH. A high-sugar diet affects cellular and humoral immune responses in
Drosophila. Experimental Cell Research. 2018 Jul;368(2):215–24.

33. Baucom RS, de Roode JC. Ecological immunology and tolerance in plants and animals.
Functional Ecology. 2011;25(1):18–28.

34. Musselman LP, Fink JL, Narzinski K, Ramachandran PV, Hathiramani SS, Cagan RL, et al.
A high-sugar diet produces obesity and insulin resistance in wild-type *Drosophila.* DMM
Disease Models and Mechanisms. 2011 Nov;4(6):842–9.

35. Lazzaro BP, Sackton TB, Clark AG. Genetic variation in *Drosophila melanogaster*
resistance to infection: A comparison across bacteria. Genetics. 2006 Nov 1;174(3):1539–
54.

36. Juneja P, Lazzaro BP. *Providencia sneebia* sp. nov. and *Providencia burhodogranariea* sp.
nov., isolated from wild *Drosophila melanogaster.* International Journal of Systematic and
Evolutionary Microbiology. 2009;59(5):1108–11.

37. Vandehoef C, Molaei M, Karpac J. Dietary adaptation of microbiota in *Drosophila* requires
NF-κB-dependent control of the translational regulator 4E-BP. Cell Reports. 2020 Jun
9;31(10):107736.

38. Sharon G, Segal D, Ringo JM, Hefetz A, Zilber-Rosenberg I, Rosenberg E. Commensal
bacteria play a role in mating preference of *Drosophila melanogaster.* Proceedings of the
National Academy of Sciences. 2010 Nov 16;107(46):20051–6.

39. Fink C, Staubach F, Kuenzel S, Baines JF, Roeder T. Noninvasive analysis of microbiome
dynamics in the fruit fly *Drosophila melanogaster.* Applied and Environmental Microbiology.
2013;79(22):6984–8.

40. Shin SC, Kim SH, You H, Kim B, Kim AC, Lee KA, et al. *Drosophila* microbiome modulates
host developmental and metabolic homeostasis via insulin signaling. Science.
2011;334(6056):670–4.

41. Wong ACN, Dobson AJ, Douglas AE. Gut microbiota dictates the metabolic response of
Drosophila to diet. Journal of Experimental Biology. 2014 Jun 1;217(11):1894–901.

42. Fischer CN, Trautman EP, Crawford JM, Stabb EV, Handelsman J, Broderick NA.
Metabolite exchange between microbiome members produces compounds that influence
Drosophila behavior. Scott K, editor. eLife. 2017 Jan 9;6:e18855.

43. Edgecomb RS, Harth CE, Schneiderman AM. Regulation of feeding behavior in adult
Drosophila melanogaster varies with feeding regime and nutritional state. Journal of
Experimental Biology. 1994 Dec 1;197(1):215–35.

44. Yapici N, Cohn R, Schusterreiter C, Ruta V, Correspondence LBV, Vosshall LB. A taste
circuit that regulates ingestion by integrating food and hunger signals in brief article a taste
circuit that regulates ingestion by integrating food and hunger signals. 2016 April 21; 165(3)
715-729

45. Ro J, Harvanek ZM, Pletcher SD. FLIC: High-Throughput, continuous analysis of feeding
behaviors in *Drosophila.* PLOS ONE. 2014 Jun 30;9(6):e101107.

46. Klepsatel P, Knoblochová D, Girish TN, Dircksen H, Gáliková M. The influence of
developmental diet on reproduction and metabolism in *Drosophila.* BMC Evol Biol. 2020 Jul
29;20(1):93.

47. Wu Q, Yu G, Park SJ, Gao Y, Ja WW, Yang M. Excreta quantification (EX-Q) for
longitudinal measurements of food intake in *Drosophila.* iScience. 2020 Jan
24;23(1):100776.

48. Bajgar A, Dolezal T. Extracellular adenosine modulates host-pathogen interactions through
regulation of systemic metabolism during immune response in *Drosophila.* PLOS
Pathogens. 2018 Apr 27;14(4):e1007022.

49. Schwenke RA, Lazzaro BP. Juvenile hormone suppresses resistance to infection in mated
female *Drosophila melanogaster.* Current Biology. 2017;27(4).

50. Chambers MC, Jacobson E, Khalil S, Lazzaro BP. Consequences of chronic bacterial
infection in *Drosophila melanogaster.* PLOS ONE. 2019 Oct 24;14(10):e0224440.

51. Hanson MA, Dostálová A, Ceroni C, Poidevin M, Kondo S, Lemaitre B. Synergy and
remarkable specificity of antimicrobial peptides in vivo using a systematic knockout
approach. MacPherson AJ, Garrett WS, Hornef M, Hooper LV, editors. eLife. 2019 Feb
26;8:e44341.

52. Rolff J, Schmid-Hempel P. Perspectives on the evolutionary ecology of arthropod
antimicrobial peptides. Philosophical Transactions of the Royal Society B: Biological
Sciences. 2016 May 26;371(1695):20150297.

53. Bulet P, Hetru C, Dimarcq JL, Hoffmann D. Antimicrobial peptides in insects; structure and
function. Developmental & Comparative Immunology. 1999 Jun 1;23(4–5):329–44.

54. Lazzaro BP, Zasloff M, Rolff J. Antimicrobial peptides: Application informed by evolution.
Science. 2020 May 1;368(6490). Available from:
http://science.sciencemag.org/content/368/6490/eaau5480

55. Kuroda K, Okumura K, Isogai H, Isogai E. The human cathelicidin antimicrobial peptide LL-
37 and mimics are potential anticancer drugs. Frontiers in Oncology. 2015;5. Available from:
https://www.frontiersin.org/articles/10.3389/fonc.2015.00144

56. Unckless RL, Howick VM, Lazzaro BP. Convergent balancing selection on an antimicrobial
peptide in *Drosophila.* Current Biology. 2016;26(2):257–62.

57. Buccitelli C, Selbach M. mRNAs, proteins and the emerging principles of gene expression
control. Nat Rev Genet. 2020 Oct;21(10):630–44.

58. Gupta V, Frank AM, Matolka N, Lazzaro BP. Inherent constraints on a polyfunctional tissue
lead to a reproduction-immunity tradeoff. BMC Biology. 2022 Jun 2;20(1):127.

59. Sackton TB, Lazzaro BP, Clark AG. Genotype and gene expression associations with
immune function in *Drosophila.* PLOS Genetics. 2010 Jan 8;6(1):e1000797.

60. Ayres JS, Schneider DS. The role of anorexia in resistance and tolerance to infections in
Drosophila. PLoS biology. 2009 Jul;7(7):e1000150.

61. Singh A, Basu AK, Bansal N, Shit B, Hegde T, Prasad NG. Effect of larval diet on adult
immune function is contingent upon selection history and host sex in *Drosophila*
melanogaster. bioRxiv; [Preprint] 2022 [cited 2023 Aug 1]. p. 2022.03.03.482770. Available
from: https://www.biorxiv.org/content/10.1101/2022.03.03.482770v1

62. Libert S, Chao Y, Zwiener J, Pletcher SD. Realized immune response is enhanced in long-
lived puc and chico mutants but is unaffected by dietary restriction. Molecular Immunology.
2008;8.

63. Nehme NT, Liégeois S, Kele B, Giammarinaro P, Pradel E, Hoffmann JA, et al. A model of
bacterial intestinal infections in *Drosophila melanogaster.* PLOS Pathogens. 2007 Nov
23;3(11):e173.

64. Krejčová G, Danielová A, Nedbalová P, Kazek M, Strych L, Chawla G, et al. *Drosophila*
macrophages switch to aerobic glycolysis to mount effective antibacterial defense. eLife.
2019;8. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6867711/

65. Mason CJ, Peiffer M, Felton GW, Hoover K. Host-specific larval lepidopteran mortality to
pathogenic *Serratia* mediated by poor diet. Journal of Invertebrate Pathology. 2022 Aug
13;107818.

66. Gao Y, Zhu C, Li K, Cheng X, Du Y, Yang D, et al. Comparative proteomics analysis of
dietary restriction in *Drosophila.* PLOS ONE. 2020 Oct 16;15(10):e0240596.

67. Lemaitre B, Reichhart JM, Hoffmann JA. *Drosophila* host defense: Differential induction of
antimicrobial peptide genes after infection by various classes of microorganisms. PNAS.
1997 Dec 23;94(26):14614–9.

68. Vasudevan D, Clark NK, Sam J, Cotham VC, Ueberheide B, Marr MT, et al. The GCN2-
ATF4 signaling pathway activates 4E-BP to bias mRNA translation and boost antimicrobial
peptide synthesis in response to bacterial infection. Cell Rep. 2017 Nov 21;21(8):2039–47.

69. Iatsenko I, Marra A, Boquete JP, Peña J, Lemaitre B. Iron sequestration by transferrin 1
mediates nutritional immunity in *Drosophila melanogaster.* PNAS. 2020 Mar
31;117(13):7317–25.

70. Port F, Bullock SL. Augmenting CRISPR applications in *Drosophila* with tRNA-flanked
sgRNAs. Nat Methods. 2016 Oct;13(10):852–4.

71. Bruckner JJ, Zhan H, Gratz SJ, Rao M, Ukken F, Zilberg G, et al. Fife organizes synaptic
vesicles and calcium channels for high-probability neurotransmitter release. Journal of Cell
Biology. 2016 Dec 20;216(1):231–46.

72. Khalil S, Jacobson E, Chambers MC, Lazzaro BP. Systemic bacterial infection and immune
defense phenotypes in *Drosophila melanogaster.* Journal of Visualized Experiments. 2015
May 13;2015(99).

73. Lazzaro BP. A population and quantitative genetic analysis of the Drosophila melanogaster
antibacterial immune response. PhD. Thesis, The Pennsylvania State University; 2002.
Available from:
https://www.proquest.com/docview/305486474/abstract/CF5BBC5BAB284669PQ/1

74. Jensen EC, Amburn DSK, Schlegel AH, Nickerson KW. A collection of *Serratia marcescens*
differing in their insect pathogenicity towards *Manduca sexta* larvae. bioRxiv. [Preprint] 2020
Jul 29[cited Nov 2023] ;2020.07.29.226613. Available from:
https://www.biorxiv.org/content/10.1101/2020.07.29.226613v1

75. Gordon KE, Wolfner MF, Lazzaro BP. A single mating is sufficient to induce persistent
reduction of immune defense in mated female *Drosophila melanogaster.* Journal of Insect
Physiology. 2022 Jul 1;140:104414.

A Sucrose Dose Infection Survival Experimental Design

Rear Canton-S flies on yeast-cornmeal rearing diet. → When larvae pupate on sides of the chamber, swap rearing medium with on of six (w/v) sucrose diets. → Allow adults to feed for 3-5 days prior to systemic infection. Score survivorship daily for 5-days.

B *Providencia rettgeri* (G-)

Sucrose (w/v)
- 0% a n = 108
- 2% | n = 120
- 4% |b n = 124
- 8% | n = 120
- 16% | n = 120
- 24% |c n = 120

C *Serratia marcescens* (G-)

Sucrose (w/v)
- 0% | n = 167
- 2% |a n = 152
- 4% | n = 171
- 8% | n = 172
- 16% |b n = 167
- 24% | n = 157

D *Lactococcus lactis* (G+)

Sucrose (w/v)
- 0% a n = 123
- 2% | n = 119
- 4% | n = 121
- 8% |b n = 121
- 16% | n = 120
- 24% |c n = 119

E *Enterococcus faecalis* (G+)

Sucrose (w/v)
- 0% | n = 120
- 2% | n = 120
- 4% | n = 120
- 8% |a n = 120
- 16% | n = 121
- 24% | n = 120

A — Quantified Excreta

B — Glucose Levels

C — Trehalose Levels

D — Glycogen Levels

A *Providencia rettgeri*

Proportion of flies alive at each time point

Hour	0	2	4	6	8	10	12	14	16		24	36	48
2%	1.0	0.97	0.93	0.93	1.0	0.93	0.90	0.90	0.97		0.76	0.76	0.71
16%	1.0	0.97	1.0	0.89	1.0	0.83	0.85	0.81	0.59		0.47	0.29	0.29

B *Serratia marcescens*

Proportion of flies alive at each time point

Hour	0	2	4	6	8	10	12	14	16		24	36	48
2%	1.0	1.0	1.0	0.96	0.90	0.93	0.85	0.90	0.83		0.80	0.74	1.0
16%	1.0	1.0	0.93	1.0	0.83	0.69	0.76	0.67	0.63		0.62	0.42	0.44

A *Providencia rettgeri:* AMP expression 8-hpi

Infection-Induced Expression (log_{10})

Sucrose (w/v): 2% (blue), 16% (orange)

Diptericin, Attacin A, Cecropin, Drosocin, Drosomycin, Defensin

B *Serratia marcescens:* AMP expression 8-hpi

Infection-Induced Expression (log_{10})

Sucrose (w/v): 2% (blue), 16% (orange)

Diptericin, Attacin A, Cecropin, Drosocin, Drosomycin, Defensin

C *P. rettgeri:* Drosocin levels 6-hpi

Tagged Drosocin (*Mr*/fly)

Sucrose (w/v): 2% (blue), 16% (orange)

Lorem ipsum

*** , ** , ***

uninfected, *Providencia*

D *S. marcescens:* Drosocin levels 6-hpi

Tagged Drosocin (*Mr*/fly)

Sucrose (w/v): 2% (blue), 16% (orange)

*** , p = 0.71 , ***

uninfected, *Serratia*

E Probability of Cecropin Present

Probability of peptide present

p = 0.35 , **

Providencia, *Serratia*

Sucrose (w/v): 2% (blue), 16% (orange)

F Cecropin levels 6-hpi

Tagged Cecropin (*Mr*/fly)

* , p = 0.21

Providencia, *Serratia*

Sucrose (w/v): 2% (blue), 16% (orange)

A w^{1118}; $Dro^{\Delta HF}$/ c564Gal4

B w^{1118}; c564Gal4 / + ; $CecA1^{\Delta HF}$/ +

Gene	Forward Primer	Reverse Primer
Actin-5C	GGCGCCCCTTGAACCCCAAAG	AAGTTGCCCGGAGGCGTACAG
Attacin A	CTCTCATTCTGGCCACACC	TGCTTTGGCCGGACATTCC
Cecropin I	GTCGCTGTGCTTGTTGA	TCTTGGCAACATACTGGG
Defensin A	GCGGCCATGGTTCTTCG	TCCTGGTGTCCCACAGA
Diptericin	GCTGCGCAATCGCTTCTACT	TGGTGGAGTGGGCTTCAT
Drosocin	CGGTTGCGCTTGCGTGC	TTGGACAGCTCCTGCTT
Drosomycin	CGTGAGAACCTTTTCCAATATGATG	TCCCAGGACCACCAGCAT