BBB recovery after stroke by self-replenishing E-pericytes transdifferentiated from local endothelial cells.

Tingbo Li[1], Jiaqi Tu[2], Qingzhu Gao[2], Xuzhao Li[2], Zhu Zhu[2], Yuxiao Jin[2], Yiyi Zhang[2], Jiongfang Xie[3], Peiran Zhu[4], Bingrui, Zhao[2], Jie-Min Jia*[2, 5, 6, 7]

[1]College of Life Sciences, Zhejiang University, Hangzhou, Zhejiang, 310058, China.

[2]Laboratory of Neurovascular Biology, School of Life Sciences, Westlake University, Hangzhou, 310024, China.

[3]Key Laboratory of Structural Biology of Zhejiang Province, School of Life Sciences, Westlake University, Hangzhou, Zhejiang, 310024, China.

[4]School of Materials Science and Engineering, Zhejiang University, Hangzhou, China.

[5] Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou, 310024, China

[6]Key Laboratory of Growth Regulation and Translational Research of Zhejiang Province, School of Life Sciences, Westlake University, Hangzhou, 310024, China.

[7]Institute of Basic Medical Sciences, Westlake Institute for Advanced Study, Hangzhou, 310024, China.

*Correspondence: jiajiemin@westlake.edu.cn

Ischemic stroke causes a significant amount of cell death, including pericytes. Chronic perfused phase of ischemic stroke, the number of pericytes increases, and limited research has been explored on the reasons for the recovery of pericytes. By genetic tracing, we identified ECs undergoing cell-fate changing, through TGFb-pSMAD3-EndoMT pathway transdifferentiation, into pericytes (E-pericytes). Some of the migrating E-pericytes induced by stroke are derived from detached ECs. The E-pericytes can integrate into vasculature again, suggesting ischemic brains can recycle ECs from unfunctional vessels to replenish lost pericytes as an innate self-maintenance program sustaining repair phenomena. We also found E-pericytes promotes BBB restoration, increases angiogenesis and restores cerebral blood flow(CBF), resulting more neurons survive and accelerating behavioral rehabilitation.

Stroke kills millions each year and no effective treatment is available so far[1]. Brains possess an intrinsic, limited self-repair capacity[2, 3], e.g. spontaneous angiogenesis after stroke. However, understanding the definitive spontaneous repairing processes is largely incomplete. ECs form vascular tubes, while pericytes encircle the endothelial tube. The observation that more pericytes die faster than ECs may explain neurodegeneration exacerbation and brain self-recovery hindering after stroke because pericytes are a type of pericytess known to be critical for the blood-brain barrier and cerebral blood flow[4-6]. However, little is known about cellular sources for pericyte regeneration in the adult ischemic brain while it has been well documented that the cephalic neural crest gives rise to pericytes in all blood vessels of the face and forebrain during embryonic development[7, 8]. Many capillaries collapse after stroke, however, the

fate of endothelial cells from these unfunctional vessels to be recycled or to die is

unclear. Recently, Yu et al. have demonstrated that pericytes can be differentiated

locally from a subgroup of ECs in multiple tissues under homeostasis[9]. However, the

case in brains with or without ischemia remains unknown. Identification of origin cells for

regenerated pericytes and EC's fate in the ischemic brain is of importance for

enhancing endogenous capacity to repair a stroke-damaged brain.

First, we found that using flow cytometric analysis revealed that after 2 days of

reperfusion, close to 50% of pericytes were lost, while endothelial cells accounted for

less than 20% (Fig. 1a). With the prolonged reperfusion time, the numbers of pericytes

and endothelial cells both show recovery, with the percentage of pericyte recovery

exceeding 40% and the percentage of endothelial cell recovery approaching 10% at

RP34D, comparion to RP2D(Fig. 1b-d). This indicates that in the chronic phase of

ischemic reperfusion, there are ways for both endothelial cells and pericytes to increase

the cell number. In ischemic stroke mice continuously treated with EdU by

intraperitoneal injection every day for 34 days, we observed the presence of EdU$^+$

pericytes(Fig. 1e-f). After statistical analysis(Fig. 1g), it was found that EdU$^+$ pericytes

accounted for less 20% of the total pericytes, suggesting that over 20% of pericytes

may not be generated through self-proliferation, comparison with flow analysis results.

Could these non-EdU$^+$ pericytes originate from the transformation of endothelial cells?

To test if cerebral ECs differentiate into pericytes during adult angiogenesis as do

those in the mammary gland, skin, and retina[9], we employed a cell fate tracing system

by crossing EC-specific DNA recombinase driver line *Cdh5-CreERT2*[10] with triple EGFP

reporter (Ai47)[11] or tdTomato reporter (Ai14) (The Jackson Laboratory) (Fig. 2a). ECs

and their potential progeny cells are permanently labeled by EGFP or tdTomato

fluorescent proteins once tamoxifen (TAM) is provided(Extended Data Fig.2a). At the

same time, When tamoxifen was not given, there was no EGFP signal(Extended Data

Fig.2a-b). Upon long-term TAM exposure (Fig. 2b), although some of these EGFP[+] ECs

in skin tissue change their fates into EGFP[+] pericytes[9], brain EGFP[+] ECs, however,

seemed never (Fig. 2c-e). all EGFP[+] cells remained expressing multiple classical

endothelial markers, such as ERG, GluT1 and VE-cadherin, despite up to a two-year

tracing (Extended Data Fig.2c-f). Consistent with previous reports[9, 10, 12], these

fluorescent proteins were restricted in ECs but not in any other types of brain cells

short-term after TAM exposure (4 - 7 days, Extended Data Fig. 2g-m). At the same time，

ECs and their potential progeny cells are never labeled by pericyte markers: PDGFRβ,

NG2, CD13, Caldesmon and α-SMA (Fig.2f-h;Extended Data Fig.2n-q). Together, these

results demonstrated unlike in the peripheral system and retina, cerebral ECs are not

the cellular source for pericytes in adult brains under homeostasis.

Ischemic stroke, modeled by middle cerebral artery occlusion (MCAO, Extended

Data Fig.2r-t), induced cerebral ECs detaching from vessels, gradually losing EC

markers, basal membrane protein, and undergoing drastic morphological changes such

as losing hollow structure and growing multiple processes from cell soma (Fig. 2i;

Extended Data Fig.2u-v). This finding suggested that ECs might undergo cell fate

changing in response to ischemia attack. Thus, we referred to this subgroup of EGFP[+]

cells as 'EGFP[+] non-EC cells'. TAM was delivered to 1.5-month-old mice and then at

least two weeks later we performed MCAO surgery (Fig. 2b), aiming to ensure no extra

residual TAM. After 2-hour MCAO, sutures were withdrawn, permitting various periods

of reperfusion, ranging from 8 all the way through 514 days (RP2D to RP514D)

(Extended Data Fig.2w-z). These EGFP⁺ non-EC cells can be found anywhere possible

in the ischemic region, showing no regional specificity (Extended Data Fig. 2a1).

According to whether having secondary processes, we classified EGFP⁺ non-EC cells

into two types: type I, short, and fewer primary processes, no secondary processes

(Extended Data Fig. b1,d1-e1); type II, longer and more primary branches with

secondary ones (Extended Data Fig. b1,d1-e1), while their cell bodies were comparable

in shape and size (Extended Data Fig. f1).

We found that the prevalence of these two types of cells switched when reperfusion

period prolonged. (Extended Data Fig. c1). This morphological switching suggests type I

becoming type II gradually. The prevalence of EGPF⁺ non-EC cells started to rise at

RP8D, peaked by RP52D, and finally declined back to a low level (Extended Data Fig.z).

This curve suggested that these transformed avascular EGFP⁺ non-EC cells may die or

be recruited back to vessels, becoming vascular cells again. We termed this possibility

a recycling process. Furthermore, FACS analysis (RP34D) revealed an EGFP⁺ but

CD31 negative population (EGFP⁺ non-EC cells, Fig. 2j). These EGFP⁺ transformed

cells accounted for ~1.2% of total EGFP⁺ cells which included ECs and non-EC cells

(Fig. 2k).

Cdh5-CreERT2 may label a small portion of neutrophils and monocytes in the bone

marrow and circulation[12] (Extended Data Fig. 3 a-b), which are known to enter brain

parenchyma shortly after stroke onset but dies out[13]. To discriminate whether

parenchyma EGFP⁺ non-EC cells are derived from EGFP⁺ bone marrow cells in

Cdh5CreERT2;Ai47 mice, we introduced a viral expression system restricted to ECs

preventing EGFP expression in hematopoietic cells (Extended Data Fig. 3c-e).

Specifically, we injected cranially the Cre-dependent *AAV-CAG-DIO-EGFP* virus to

Cdh5-CreERT2 single-positive mice that do not carry *Ai47* or *Ai14* reporter (Extended

Data Fig. 3c), resulting in high labeling specificity to ECs (Extended Data Fig. 3f-g).

Once again, we found EGFP⁺ ECs transforming into EGFP⁺ non-EC cells after stroke

(Extended Data Fig. 3h). Notably, some of these cells extended their long processes

and enwrapped capillaries, reminiscent of pericyte characteristics (Extended Data Fig.

3h). Independent radiation experiment was implemented to exclude the same possibility,

we replaced *Cdh5CreERT2;Ai47* mice's bone marrow with wild type ones (Extended

Data Fig. 3i-j). Despite the depletion of EGFP⁺ bone marrow cells under this

circumstance (Extended Data Fig. 3i-j), we still found comparable EGFP⁺ non-EC cells

in ischemic regions (Extended Data Fig. 3k-n), demonstrating that EGFP⁺ immune cells

were not the cell source of EGFP⁺ non-EC cells in parenchyma. These results together

indicate that a small proportion (1.2%) of ECs detached from vessels in the parenchyma,

lost EC features, and changed their cellular identity after subjecting them to ischemia

insult.

To assess whether these EGFP⁺ non-EC cells are derived from vascular stem

endothelial cells as reported previously[9] which involves cell proliferation, we traced cell

division events by injecting EdU for consecutive 34 days. No single EGFP⁺ non-ECs

were found to be EdU positive after examining 15 brain slices from two mice (Fig. 2l-m).

similarly, neither Ki67 signals were found (Extended Data Fig. 4a-b). Thus, we

concluded these EGFP⁺ non-EC cells may not be progenies of stem cells, suggesting

that a transdifferentiation process may be required. Fortunately, EGFP+ non-EC cells

appear at the classical pericyte location (Fig. 2n-o).

By employing a list of cell type markers, we identified that 90% of EGPF[+] non-EC

cells were pericytes (namely, E-pericytes) as they were positive for typical pericyte

markers of CD13(90%), PDGFRβ(75%), and NG2 (Fig. 2p-s;Extended Data Fig. 4c), as

well as that they were negative for all of the markers of immature and mature neurons

(doublecortin and NeuN), astrocytes (GFAP), CNS macrophages (Iba1, CD68, and

F4/80), immuce cells (CD45), and smooth muscle cells (α-SMA) (Extended Data Fig.

4d-f). The cell identity of the rest 10% of EGFP+ non-EC cells remained unknown (Fig.

3g) though as these cells were not labeled by any cell markers that we used.

Furthermore, more than a quarter of EGFP[+] pericytes were recruited back to vessels

(Fig. 2p-q,v), suggesting ECs were fully recycled not only by expressing pericyte

markers but also by locating the right anatomical position. A transformed tdTomato[+]

non-ECs (here, Ai14 reporter was used instead of Ai47) that were also tracked by *AAV-*

DIO-EGFP virus exhibited one process attaching a capillary while the other did not

(Extended Data Fig. 4g), implying capturing a recruiting moment. It has been repeatedly

reported that pericytes detach from vessels after stroke[14]. We found actually a third of

these avascular, migrating pericytes were derived from ECs (Fig. 2v). Further analysis

revealed that 23.3% of total vascular pericytes were EGFP[+] (Fig. 2t), suggesting ECs

are a fair contributor cell source for pericyte regeneration. The virus-tracked ECs also

can generate pericytes, coinciding with results from genetic tracing (Extended Data Fig.

4h-i). Meanwhile, we also employed other methods to specifically label endothelial cells

to validate the E-pericytes phenomenon. Tie2Dre;Msfd2aXER mice were used for

specific tracing of endothelial cells in the brain[15]. Firstly, we verified their specificity

using endothelial cell markers(Extended Data Fig. 4j-k). After MCAO/RP34D, we still

observed the presence of CD13$^+$E-periytes (Extended Data Fig. 4n). We also use AAV-

Bi30-Cre virus which can be traced ECs in mouse brain[16]. AAV-Bi30-Cre virus was

injected via the eye socket in Ai47 mice and we still observed the presence of CD13$^+$E-

periytes (Extended Data Fig. 4i-m,o).These results together demonstrate that after

ischemic stroke, ECs transdifferentiated into E-pericytes, likely capable of integrating

back into the cerebral vasculature.

To further verify the identified EGFP$^+$&CD31$^-$ cells as pericyte, we analyzed the

transcriptomic characteristics of EGFP$^+$&CD31$^-$ cells. First, we obtained EGFP$^+$&CD31$^-$

cells (MCAO:2H/RP34D) using flow cytometric sorting. The flow cytometric gating

strategy was CD45$^-$&Viability$^+$EGFP$^+$&CD31$^-$ cells(Fig. 3a). The sorted cells from each

group were as follows: non-stroke endothelial cells (Con-ECs), ipsilateral stroke

endothelial cells (Ipsi-ECs), and EGFP$^+$&CD31$^-$ cells derived from the ipsilateral side

(Ipsi-ETCs). After library construction, sequencing was performed. PCA analysis

revealed that Con-ECs and Ipsi-ECs had similar principal component relationships

because they were both endothelial cells, although they were derived from different

environments of the endothelial cells (non-ischemia and ischemia brain tissue,

respectively)(Fig. 3b). On the other hand, Ipsi-ETCs showed greater dissimilarity to the

principal component relationships of Con-ECs and Ipsi-ECs(Fig. 3b). This result

remained consistent different batches, suggesting significant differences in the

transcriptomes between Ipsi-ETCs and the endothelial cells represented by Con-ECs

and Ipsi-ECs, even though Ipsi-ETCs were derived from EGPF$^+$ normal endothelial cells.

Heatmap analysis of all genes indicated that the transcriptomes of Con-ECs and Ipsi-ECs were more similar and clustered closely, despite some differences caused by the ischemic environment(Fig. 3c). In contrast, the transcriptome differences between Ipsi-ETCs and Con-ECs/Ipsi-ECs were more pronounced, and Ipsi-ETCs from different batches tended to cluster together(Fig. 3c). These findings indicated significant differences between the transcriptomes of CD45$^-$&Viability$^+$EGFP$^+$&CD31$^-$ cells and endothelial cells. Subsequently, we analyzed the expression of endothelial cell marker proteins among different groups. We found that the expression of endothelial cell markers genes in Ipsi-ETCs significantly decreased compared to Con-ECs and Ipsi-ECs, and this decrease was observed across most endothelial cell marker genes(Fig. 3d). Con-ECs and Ipsi-ECs still expressed numerous endothelial cell marker genes, although differences existed between them(Fig. 3d). Additionally, we analyzed the expression of pericyte-related transcriptomes genes among different groups. We observed a significant increase in the expression of pericyte marker genes in Ipsi-ETCs, particularly in specific transcription factors(Fig. 3e), membrane proteins(Fig. 3f), and ligand proteins associated with pericytes(Fig. 3g). Conversely, the expression of pericyte-related marker genes was lower in Con-ECs and Ipsi-ECs, suggesting that the transcriptome of Ipsi-ETCs exhibited characteristics of pericyte cell transcriptomes genes. Furthermore, volcano plot analysis of ETCs/Ipsi-ECs identified differentially expressed genes with statistical significance(Fig. 3h). GO enrichment analysis revealed that the Ipsi-ETCs group was more enriched in functionally related genes associated with fate conversion, such as high expression of genes involved in cell differentiation, cell movement, cell migration, and endothelial cell migration, as well as low expression

of genes related to endothelial cell junction genes(Fig. 3i). These findings were consistent with the observations in ischemic brain slices, where 2/3 of Ipsi-ETCs detached from the blood vessels. Additionally, KEGG pathway enrichment analysis revealed that genes related to metabolic pathways were upregulated, while genes related to endothelial cell tight junctions were downregulated(Fig. 3j). GSEA analysis showed that genes associated with endothelial cell features, such as endothelial cell differentiation, cell junctions, and cell apoptosis, were downregulated(Fig. 3k-m), while genes associated with glucose metabolism pathways and ATP production through aerobic respiration were upregulated(Fig. 3n-p). It is known that endothelial cells have fewer mitochondria and primarily produce ATP through glycolysis, whereas Ipsi-ETCs showed a significant increase in genes related to glucose metabolism, as well as genes involved in ATP production. This suggests that important genes in the metabolic pathway of glucose metabolism are altered in Ipsi-ETCs. These results collectively indicate that Ipsi-ETCs exhibit significant differences in the genomic profile compared to endothelial cells and show transcriptional characteristics of pericyte cells, providing further evidence from the transcriptome analysis that EGFP[+] non-EC cells are E-pericytes.

The above experiments have demonstrated that the ischemic microenvironment after stroke promotes the transdifferentiation of endothelial cells into E-pericytes. Next, we attempted to explore the mechanism by which endothelial cells transform into E-pericyte. In 2016, Yu QC et al. reported the identification of endothelial cells expressing Protein C Receptor (Procr) in breast tissue, which possess characteristics of vascular endothelial stem cells (VESCs). Procr+ VESCs have strong proliferative abilities in vitro and also

participate in vascular remodeling in live animals. Lineage tracing studies have found

that this group of cells can transform into pericytes during breast development, and the

transcriptome of Procr+ VESCs exhibits characteristics of Endothelial-to-Mesenchymal

Transition (EndoMT) and Epithelial-to-Mesenchymal Transition (EMTs), suggesting that

the transformation of endothelial cells into pericytes in breast development may occur

through EndoMT[9]. We found that after ischemic stroke, the expression of EndoMT

marker proteins in ischemic regions of endothelial cells was significantly increased.

Specifically, at MCAO:2H/RP2D, the expression of phosphorylated SMAD3 (an impor-

tant kinase in the TGF β pathway-activated by phosphorylation) as an EndoMT marker

protein increased in the ischemic region(Fig. 4a-c). Furthermore, MCAO:2H/RP8D, the

expression of p-SMAD3 in ECs also increased in the ischemic region(Fig. 4a-c).

However, MCAO:2H/RP34D, endothelial cells almost did not express p-SMAD3 as an

EndoMT marker protein, and E-pericytes cells also almost did not express p-

SMAD3(Fig. 4a-c). At 8 days post ischemia, we also observed two adjacent EFGP[+]

cells on the same blood vessel, one cell exhibiting CD31[+]&p-SMAD3[-], still maintaining

endothelial cell characteristics, while the other cell exhibiting CD31[-]& p-SMAD3[+],

indicating that after p-SMAD3 is expressed in some endothelial cells, the characteristic

molecule CD31 is reduced or not expressed(Fig. 4d), indicating successfully catching

cells undergoing transition on time. At 2 days post ischemia, the expression of KLF4 (an

important transcription factor in the EndoMT pathway) as an EndoMT marker protein in

the ischemic region of endothelial cells increased(Extended Data Fig.5a-c). At 8 days

post ischemia, the expression of KLF4 in the ischemic region of endothelial cells

decreased, and at 34 days post ischemia, the expression of KLF4 as an EndoMT

marker protein in endothelial cells further decreased, and E-pericytes cells almost did

not express KLF4(Extended Data Fig.5a-c). At 2 days post ischemia, the expression of

α-SMA (an important marker protein in EndoMT) in the ischemic region of endothelial

cells increased(Extended Data Fig.5d-f). At 8 days post ischemia, the expression of α-

SMA as an EndoMT marker protein in the ischemic region of endothelial cells

decreased, and at 34 days post ischemia, the expression of α-SMA in endothelial cells

also decreased, and E-pericytes cells almost did not express α-SMA (Extended Data

Fig.5d-f). Furthermore, cytoplasmic flow cytometry demonstrated that at 2 days post

ischemia, the expression of α-SMA in endothelial cells increased, reaching 1.6% of the

proportion of endothelial cells in the ischemic region, which is close to the proportion of

E-pericytes cells (1.2%) in the ischemic region(Extended Data Fig.5g-h). Next, we

investigate the potential molecular signaling pathway that might mediate this E-pericytes.

Since TGFβ signaling is well documented for cell fate changing including the endothelial

mesenchymal transition (EndMT) process, a newly recognized type of

transdifferentiation[17,18]. Therefore, we injected LY364947 and SB431542, TGFβRI

inhibitors when MCAO was performed, followed by additional daily injections for

consecutive 2 days of reperfusion. The expression of p-SMA3, KLF4 and α-SMA

protein in the ischemic region of endothelial cells decreased in TGFβRI inhibitors

group(Extended Data Fig.5i-j), and cytoplasmic flow cytometry demonstrated that at 2

days post ischemia, the expression of α-SMA in endothelial cells decrease(Extended

Data Fig.5k-i). For consecutive 34 days of reperfusion, We found significantly fewer

EGFP[+] non-EC cells appeared overall(Extended Data Fig.5m-n). FACS analysis further

confirmed this conclusion quantitively (Extended Data Fig.5o-p). The specific knockout

of the *Tgfbr2* gene in endothelial cells also resulted in a decrease of EndoMT markers

(p-SMAD3, KLF4, and α-SMA) at the RP2D(Fig. 4e-h), along with a decrease in the

number of E-pericytes(Fig. 4i-l). These results strongly suggested that TGFβ-pSMAD3-

EndoMT signaling involved in E-pericytes is centrally important. Together, under

ischemic conditions, ECs exploited TGFβ signaling to transform ECs into pericytes

during reperfusion to replenish the shrunk pericyte pool occurring in the acute ischemia

period.

Pericyte-deficient mouse mutants shows that pericyte deficiency increases the

permeability of the BBB to water and a range of low-molecular-mass and high-

molecular-mass tracers[19,20]. Therefore, we speculate that E-pericytes participate in the

recovery of the blood-brain barrier (BBB) during the chronic recovery period after

ischemic stroke. E-pericytes can return to the blood vessels and perform functions such

as maintaining BBB integrity, reducing the entry of harmful substances into the brain,

promoting angiogenesis, increasing capillary density, and improving blood supply to

facilitate the recovery of neurons and other cells, ultimately promoting self-repair.

Studies in mouse models showed that, in the absence of key TGFβ receptors,

angiogenesis stalls in the yolk sac at an early stage with fatal consequences[21]. The

effect of endothelial cell-specific knockout of Tgfbr2 on blood vessel development and

endothelial function in adult mice is unknown. Firstly, we found that specific knockout of

the Tgfbr2 gene in endothelial cells does not affect the function of BBB (Extended Data

Fig.6a). At the same time, it also did not affect the expression of the BBB-related protein

ZO-1(Extended Data Fig.6b-c). Furthermore, under electron microscopy, the basement

membrane remains intact(Extended Data Fig.6d-e), and the tight connections between

endothelial cells are normal(Extended Data Fig.6f). There is also no obvisous

occurrence of transcytosis phenomena caused by BBB disruption in the endothelial

cells(Extended Data Fig.6g). This indicates that knocking out Tgfbr2 in adult mice does

not affect the BBB function of vascular endothelial cells.

At different reperfusion time points, reducing E-pericytes by specifically knocking out

Tgfbr2 in endothelial cells can aggravate the chronic BBB leakage. In the group with

Tgfbr2 knocked out specifically in endothelial cells, the BBB function was significantly

impaired when the reperfusion time exceeded 7 days(Fig.5a-b). When reperfused timely

for 118 days, it still showed low Evan blue leakage(Fig.5a-b). This suggests that

reducing E-pericytes through specific knockout of Tgfbr2 in endothelial cells can affect

the recovery of chronic BBB during reperfusion.

We observed a decrease in NeuN$^+$ neurons after ischemic stroke, with a more

significant decrease in the reducing E-pericytes mouse group(Fig.5c-d). Additionally, we

quantified the number of CD13$^+$ pericytes in the ischemic area and found a significant

decrease in pericyte density(Fig.5e-f), which is consistent with previous studies. In the

reducing E-pericytes group, the reduction in pericyte density was more significant

(Fig.5e-f). The ratio of endothelial cells to pericytes on the blood vessels was also

significantly lower than the normal group(Fig.5g). We measured the length of CD31$^+$

endothelial cells in the ischemic area and found a decrease in length(Fig.5h-i), and flow

cytometry analysis showed a more pronounced decrease in CD31$^+$ endothelial cells in

the reducing E-pericytes mouse group, indicating decreased angiogenesis(Fig.5j-k).

Furthermore, laser speckle contrast analysis revealed poorer blood flow recovery in the

reducing E-pericytes mouse group(Fig.5i-m), as well as more pronounced brain atrophy

(Fig.5n-o). This suggests that in the E-pericytes mouse group, reduced E-periytes lead

to impaired blood flow recovery during reperfusion, resulting in more cell death and

more significant brain atrophy. To test animal locomotion function, Tgfbr2$^{fl/fl}$, EC$^{Tgfbr2fl/+}$,

and EC$^{Tgfbr2fl/fl}$ mice groups were subjected to 2 hours of ischemia followed by 34 days of

reperfusion, and behavioral tests were conducted at different time points (0D, 1 day, 3

days, 7 days, 14 days, 21 days, and 34 days of reperfusion). Animal locomotion function

tests related to motor function, including pole test, beam walking test, turning test, and

adhesive removal test, were performed. Compared to the Tgfbr2$^{+/+}$ group, mice in

EC$^{Tgfbr2fl/fl}$ group showed shorter pole test duration and had statistically significant

differences starting from 21 days of reperfusion(Fig.5p); mice in EC$^{Tgfbr2fl/fl}$ group spent

less time on the beam and had statistically significant differences starting from 21 days

of reperfusion(Fig.5q); mice in EC$^{Tgfbr2fl/fl}$ group exhibited more pronounced differences

in the turning test compared to the normal values, with statistically significant

differences starting from 14 days of reperfusion(Fig.5r); mice in EC$^{Tgfbr2fl/fl}$ group were

less likely to remove the adhesive tape from their forepaws in the adhesive removal test,

with statistically significant differences starting from 14 days of reperfusion(Fig.5s).

These results indicate that the motor recovery ability of EC$^{Tgfbr2fl/fl}$ group is poorer, which

may be associated with reducng E-periytes.

Overall, using mouse genetic tracing tools combined with virus infection,

pharmacology and genetically modified mice, we demonstrated adult ischemic brains

spontaneously repair microvasculatures by regenerating new pericytes that are

transformed from ECs after detaching from vessels. Distinct from the previous findings

in peripheral organs in which a small group of endothelial stem cells Procr$^+$ ECs

differentiate into pericytes during adult angiogenesis under homeostasis[9], ECs do not

become pericytes in normal brains but transdifferentiate into pericytes without cell

division. The death rate is dozens of times higher in pericytes than in ECs within 24

hours after a transient focal ischemia[4], suggesting abundant capillaries become

damaged by being naked. What is the fate of these naked capillaries made up by ECs?

The present study suggests that these naked vessels may release ECs, some of which

may not die but rather be recycled and repurposed by the ischemic brain. By observing

type I simpler form switching to type II more complex EGFP$^+$ E-pericytes (Extended

Data Fig.2c1) and the medium stage of the transformed EGFP$^+$ E-pericytes (Extended

Data Fig. 4g), we extrapolate them likely being recruited and serve as building blocks to

repair the existing damaged vasculatures. This could be an efficient system to reuse

brain cells that have lost their original functions. Further studies are required to live

imaging the whole dynamic process during reperfusion periods.

Identifying new cellular sources for pericyte replenishment is important not only for

stroke but also for many other either acute or chronic brain injuries. This study may

suggest a novel therapy for generating EC-derived pericyte without involving stemness

which has long been a concern for tumorigenesis.

Methods

Experimental design. Transgenic Cdh5CreERT2 transgenic mice[10] cross with reporter

mice *Ai47 (EGFP$^{f/f}$) reporter* or *Ai14 (tdTomato$^{f/f}$) reporter mice*. Thus, ECs are labeled

in green or red initially. TAM was delivered when mice were 1.5 months old (6 weeks), i.

e. genetic tracing started. After waiting for 2 weeks, MCAO was performed. After a 2-

hour occlusion, reperfusion began. Contralateral sides of the ischemic brain were used

as objects for homeostasis conditions, except for the time point of tracing 4 days. Thus, it was traced for 22 days in the PR8D mice (N = 3 mice), 1.6 months in RP34D (N = 27 mice), 2.8 months in RP71D (N = 3 mice), 14 months in RP408D (N = X mice), 8.2 months in RP232D (N = 3 mice), 14 months in RP408D (N = 2 mice),17.6 months in RP51D mice (N = 2 mice). The tracing time point of 1.6 months (RP34D) was the most frequently examined. FACS and immunohistochemistry were the major two techniques used for cell identity confirmation.

Animal care and tissue dissection.

All animal experiments were carried out in accordance with protocols approved by the Institutional Animal Care and Use Committee (IACUC) at the School of Life Sciences, Westlake University. Wild-type C57BL6/J mice were purchased from the Laboratory Animal Resources Center of Westlake University. Cdh5CreERT2 mice were a gift from Le-Ming Zheng (Peking University). Gt(ROSA)26Sor$^{tm47(CAG-EGFP*)Hze}$ (Ai47) was share by Zilong Qiu (ION). We crossed Cdh5Cre-ERT2 mice to Ai47 mice to generate Cdh5CreERT2:Ai47 mice. Standard chow and water were provided to mice *ad libitum*. Four mice were housed in each cage. All animals were housed in a standard animal room with a 12/12-hour light/dark cycle at 25 °C. Both male and female mice were used in this study. Mice were anaesthetized by intraperitoneal injection pentobarbital sodium (100 mg/kg), and brain dissection was performed immediately for FACS experiments or cardiac PFA perfusion was performed for immunohistochemistry experiment. After fixation, brains were dissected and kept in 4% PFA for additional 4 hours and then transferred to PBS until further processing.

MCAO

Adult mice were anesthetized with pentobarbital sodium (100 mg/kg) and body
temperature was maintained during surgery with a heating pad. A midline neck incision
was made, the right common carotid artery was carefully separated from the vagus
nerve, and the artery was ligated using a 5.0-string. A second knot was made on the left
external carotid artery. The right internal carotid artery (ICA) was isolated, and a knot
was left loose with a 5.0-string. This knot was not tightened until the intraluminal
insertion was done. A small hole was cut on the common carotid artery before it was
bifurcated to the external carotid artery and ICA. A silicon-coated monofilament (tip
diameter = 230 μm, Doccol Corporation, 602256PK5Re) was gently advanced into the
ICA until it stopped at the origin of the MCA in the circle of Willis. The third knot on the
ICA was closed to fix the filament in position. During MCA occlusion (2 h), mice were
kept in a warm cage in which the temperature was maintained at 35 °C.

Laser speckle contrast imaging (LSCI)

LSCI provides a measure of blood flow velocity by quantifying the extent of blurring of
dynamic speckles caused by the motion of red blood cells through the vessels. Briefly,
mice were placed under an RFLSI III device (RWD Life Sciences) before and after the
suture was successfully inserted through CCA. The skull over both hemispheres was
exposed by making an incision along the midline of the scalp. When a 785 nm laser is
used to illuminate the brain, it produces a random interference effect that represented
blood flow in the form of a speckle pattern. Scattering light was detected by a charge-
coupled device (CCD) camera, and the images were acquired by custom software from
RWD Life Sciences Company. For each acquisition, a total of 160 images, each of
which measured 2048 ×2048 pixels, were collected at 16 Hz.

Tamoxifen

Tamoxifen, 10 mg/ml dissolved in 200 μl corn oil, was administrated intragastrically for 4 consecutive days. Prior to further experiments, fluorescent protein expressions in mouse ear vasculatures were first examined. Tamoxifen was administrated at least 2 weeks before MCAO surgery.

Immunohistochemistry

Cryosections of fixed mouse brains (50 μm) were handled free-floating and washed for 5 minutes in PBS prior to further procedures. Tissue sections were permeablized and blocked in the PBS containing 0.5% Triton and 5% BSA at room temperature for 1 hour. Then, brain sections were incubated with a different primary antibodies, including CD31(1:400, BD, Cat# 557355), VE-Cadherin(1:200, BiCellScientific, Cat# 00105), ERG(1:300, Sigma, Cat# ab110639), GLUT1 (1:300, Sigma, Cat# 07-1401), NeuN (1:300, Merck, Cat# ABN90P), Doublefortin (1:300, Santa clus, Cat# sc-271390), Iba1 (1:300, HUABIO, Cat# ET1705-78), GFAP (1:300, Proteintech, Cat# CL488-60190), PDGFR β (1:300, ThermoFisher, Cat# 14-1402-82), NG2 (1:300, Merck, Cat# AB5320) ,CD68 (1:300, BIO-RAD, Cat# MCA1957T), CD45-APC (1:200, Biogend, Cat# 103112), Ki67 (1:300, thermoscientific, Cat# RM-9106-S0), F4/80 (1:300, Cell Signaling, Cat# 30325), CD13 (1:300, R&D system, Cat# AF2335), α-SMA-CY3 (1:300, Sigma, Cat# C6198), which were all diluted in the blocking solution (1% percelin, 5% BSA, and 0.1% Triton in PBS) at 4 °C overnight. After rinsing in PBS 3 times for 5 minutes each, brain slices were incubated with secondary antibodies accordingly, including donkey anti-mouse IgG (A21202, Thermo Fisher Scientific, 1:1,000) and donkey anti-rabbit IgG

(A10040, Thermo Fisher Scientific, 1:1,000). They were diluted in the blocking solution.

After a 2-hour incubation, brain slices were washed with PBS three times for 15 minutes

each. Finally, Hoechst counterstaining was performed for each specimen. After

mounting in an anti-fade mounting medium, images were acquired by using a Zeiss

LSM800 laser-scanning confocal microscope with a 20X or 63X objective, with or

without Airyscan mode.

Flow cytometry

Primary cell preparation was conducted as described below. Mouse brains were

dissected into cold HBSS with glucose and BSA (HBSS^{++}, Vendor info) were cut into

small pieces by sterile scissors, followed by a centrifuge at 1600 rpm/min for 5 mins to

get rid of supernatant. The minced tissues after the centrifuge were incubated with

collagenase IV (2.5 mg/ml, ROCHE, Cat# 11088858001) at 37 °C with gentle rotation

for 30 min in 2 ml of HBSS^{++}. Next, samples were passed through a 70-µm filter and

washed with HBSS^{++}. To obtain higher cell yield, grind the tissue on the 70-µm filter

using a syringe piston and guarantee all tissue pass the filter. The following antibodies

in 1:200 dilutions were used: CD31-PE-594 (1:200, Biolegend, Cat# 102520), CD45-

APC (1:200, Biogend, Cat# 103112), Ly6C-APC (1:200, Biolegend, Cat# 128016),

CD11b-PE (1:200, Biolegend, Cat#101212), Viability dye (1:1000, ThermoFisher,

Cat#65-0863-14). Antibody incubation was performed on ice for 30min in HBSS with

glucose and BSA. All analyses were performed with CytoFLEX LX-5L1(Beckman).

FlowJo x software was used for data analyses.

EdU injection

EdU (200 mg/kg, Alfa Aesar Chemical) was injected intraperitoneally for consecutive 34

days since when MCAO was performed. EdU was initially dissolved in DMSO at the

stock concentration of XX mg/ml and was diluted with saline prior to injection.

Virus injection

The following adenovirus vector *AAV-CAG-DIO-EGFP* was purchased for ShuMi,

Wuhan, China (PT-0168). Half of the one-micron virus was delivered to postnatal day 2

mouse pups by intracerebroventricular injection to both sides. After 1.5 months,

tamoxifen was administrated. One month later, these mice were subjected to MCAO to

induce ischemic stroke.

Radiation and bone marrow transplantation

At two weeks post tamoxifen injection, adult *Cdh5CreERT2:Ai47* mice received

radiation (total: 11Gy, 2 times, 5.5Gy per time, interval 2 hours) and bone marrow cell

transplantation from wild-type mice with the same genetic background. RBCs were

removed from the bone marrow cells by using ACK Lysis Buffer (4 Million cells/mouse).

These mice were subjected to MCAO 2 months after bone marrow reconstruction.

Mouse cranial imaging by two-photon microscopy

Adult mice were anesthetized with pentobarbital sodium, and analgesia was provided by

subcutaneous injection of 0.2% meloxicam. The scalp was removed, and the skull was

exposed and cleaned. A dental drill with a 0.6-mm-diameter bit was used to engrave

and thin the bone around the circular craniotomy area at a size of 3 mm. The piece of

skull was carefully peeled off with fine-pointed forceps, and a cranial window was

generated. A coverslip with a 3-mm diameter was placed on the cranial window, and its

perimeter was completely sealed with a 1% agarose gel. The metal headplate was

glued onto skull with dental acrylic, through which the mouse brain was fixed on a

headplate holder. The headplate holder together with the mouse was placed under a

two-photon microscope (FVMPE-RS, Olympus). The cerebral vasculature

Cdh5CreERT2:Ai47 mice was visualized and imaged once a day before and after stroke.

All surgeries were performed with sterilized instruments and an environment.

Statistics.

The quantified data in all figures were analyzed with GraphPad Prism 7.0 (La Jolla, CA,

USA) and presented as the mean ± SEM with individual data points shown. Unpaired

two-tailed Student's t-test was used for assessing the statistical significance between

two groups. Statistical significance was determined by calculation of p-value (*$p < 0.05$,

**$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$, ns: not significant). The repetition of our

data is independent biological replicates and the number of replicates for each

experiment is noted in the corresponding figure legend.

Data availability

The source data underlying the graphs and charts shown in the figures and tables are

provided in Supplementary Data 1. All data generated or analyzed during this study are

included in this published article (and its supplementary information file).

Acknowlegements

We thank Xuzhao Li, Zhu Zhu, Dongdong Zhang, Jinze Li, Jiayu Ruan, Lili Zhou for

their constructive discussions. We thank Danping Lu for her responsive and timely

purchasing support, Jiongfang Xie for sharing AAV virus. We thank the animal facility for

its technical assistance with rodent housing, Biomedical Research center platform for

technique support. This work is supported by the National Natural Science Foundation

of China (31800864 and 31970969 to J.-M.J, 82101475 to Z.Z.), Westlake startup

funds, Westlake Education Foundation and MRIC funds (103536022011) to J.-M.J.;

Westlake Education Foundation to B.D.

Author contributions

J.-M.J. conceived the research. T.L. and J.-M.J. designed experiments. T.L. performed

all the experiments and data quantification, except for the virus cranial injection. Y.W.

performed FACS together with T.L.. Z.Z. conducted some stroke models. Q.G. and B.Z.

conducted mouse genotyping.

Competing interests

All authors declare no competing interests.

Fig.1

a **RP2D**
b **RP7D**
c **RP34D**

Contralateral
Contralateral
Contralateral
Contralateral
Contralateral
Contralateral

Ipsilateral
Ipsilateral
Ipsilateral
Ipsilateral
Ipsilateral
Ipsilateral

SSC-A
CD31
PDGFRβ

d

The percentage of cell
(Ipsilateral/Contralateral)

● ECs
■ Pericytes

0.9
0.676
0.473

e
EGFP
CD13
CD13
EGFP

EdU
CD13
EdU
CD13
EGFP
EdU
Hoechst

10 μm

f
EGFP
CD13
EGFP
CD13

EdU
CD13
EdU
EGFP
CD13
EdU
Hoechst

10 μm

g

Percentage(%)

****

N=3

EdU⁺
EdU⁻

Figure legends

Fig. 1 Ischemic stroke leads to a large number of pericyte death in the acute phase, while the number of pericytes gradually increases in the chronic phase of ischemia. a, FACS analysis the number of CD31$^+$ECs and PDGFRβ$^+$ pericytes in contralateral control hemisphere (up plot) and ischemic side (down plot) of the same mouse at RP2D. b, FACS analysis the number of CD31$^+$ECs and PDGFRβ$^+$ pericytes in contralateral control hemisphere (up plot) and ischemic side (down plot) of the same mouse at RP7D. c, FACS analysis the number of CD31$^+$ECs and PDGFRβ$^+$ pericytes in contralateral control hemisphere (up plot) and ischemic side (down plot) of the same mouse at RP34D. d, Quantification of percentages of CD31$^+$ECs and PDGFRβ$^+$ pericytes at RP2D ,RP7D and RP34D(N=3). e, Illustration of the EdU$^{+\&}$CD13$^+$ pericytes on blood vessel at RP34D. f, Illustration of the EdU$^{+\&}$CD13$^+$ pericytes in brain parenchyma at RP34D. g, Quantification of percentages of EdU$^{+\&}$CD13$^+$ pericytes at R34D(N=3). Data presented as mean ± SEM, ** P < 0.01, *** P < 0.001, **** P < 0.0001.

Fig.2

Fig. 2 Genetic tracing of cerebral ECs demonstrates that they can give rise to pericyte-like cells via transdifferentiation after ischemic stroke. a, Illustration of genetic tracing strategy using *Cdh5CreERT2;Ai47* or *Ai14* line. b, Experimental design for tracing periods and reperfusion periods following 2-hour ischemia. c, Confocal images of brain slice vasculatures traced for 1.6 months, stained with EC markers (CD31). d, FACS analysis the specificity of EGFP$^+$ cells using EC marker (CD31). e, Quantification of the specificity of EGFP$^+$ cells using EC marker (CD31)(N=4). f, Confocal images of brain slice EGFP$^+$ cells stained with pericytes markers (PDGFRβand NG2). h, FACS analysis the specificity of EGFP$^+$ cells using pericytes marker (PDGFRβ). i, Confocal images of brain slice EGFP$^+$ cells stained with EC marker (CD31). j, FACS analysis the percentage of EGFP$^+$&CD31$^-$cells using EC marker (CD31). k, Quantification of FACS analysis the percentage of EGFP$^+$&CD31$^-$ cells using EC markers (CD31)(N=4). l, Confocal images of brain slice EGFP$^+$ cells stained with proliferation marker (EdU). m, Quantification of confocal images of brain slice EGFP$^+$ cells stained with proliferation markers (EdU).(N=2, 15 brain slices*2). n, Confocal images of brain slice EGFP$^+$ cells stained with ECs marker (CD31). o, Confocal images of brain slice EGFP$^+$ cells stained with ECs marker (CD31)in classic pericyte location. p, Confocal images of brain slice EGFP$^+$ cells stained with pericytes marker (CD13)in classic pericyte location. q, Confocal images of brain slice EGFP$^+$ cells stained with pericytes marker (PDGFRβ)in classic pericyte location. r, Quantification of confocal images of brain slice EGFP$^+$ cells stained with pericytes marker (CD13) in classic pericyte location..(N=3). s, Quantification of confocal images of brain slice EGFP$^+$ cells stained with pericytes marker (PDGFRβ) in classic pericyte location..(N=3).

562   **t**, The transforming efficiency of EGFP$^+$ non-EC cell to pericytes. **u**, Quantification of the

563   percentages of vascular and avascular EGFP$^+$ pericytes. N = 3 mice. **v**, The percentage

564   of avascular EGFP$^+$ pericytes in the total migrating pericyte population after stroke. **w**,

565   The contribution of vascular EGFP$^+$ pericytes to the total vascular pericyte population.

566   Data presented as mean ± SEM, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

Fig.3

Fig.3 The transcriptome demonstrates that EGFP⁺&CD31⁻ cells exhibit

characteristics of pericytes. a, Illustration the gate of sorting EGFP⁺&CD31⁻ cells. b.

PCA analysis each group similar principal component relationships. c, Heatmap

analysis of all genes expression. d, Illustration the expression of endothelial cell

markers genes in each group. e, Illustration the expression of pericytes specific

transcription factors genes in each group. f, Illustration the expression of pericytes

specific membrane proteins genes in each group. g, Illustration the expression of

pericytes specific ligand proteins genes in each group. h, Illustration the different genes

expression of ETCs group and Ipsi-ECs group. i, Illustration the GO enrichment analysis

of ETCs group and Ipsi-ECs group. j, Illustration the KEGG pathway enrichment

analysis of ETCs group and Ipsi-ECs group. j-m, Illustration the GSEA analysis

analysis of ETCs group and Ipsi-ECs group.

Fig.4

Fig.4 TGFβ-pSMAD3-EndoMT signaling involved in E-pericytes formation after storke. a, Confocal images of brain slice EGFP$^+$ cells stained with EndoMT marker (p-SMAD3) at RP2D,RP8D and RP34D. b, Quantification the number of confocal images of brain slice EGFP$^+$ cells stained with EndoMT marker (p-SMAD3) at RP2D,RP8D and RP34D.(N=3). c, Quantification the intensity of confocal images of brain slice EGFP$^+$ cells stained with EndoMT marker (p-SMAD3) at RP2D,RP8D and RP34D(N=3). d, Confocal images of brain slice EGFP$^+$ cells stained with EndoMT marker (p-SMAD3) at RP8D. e, Confocal images of brain slice EGFP$^+$ cells stained with EndoMT marker (p-SMAD3,KLF4 and α-SMA) at RP2D after specific knockout of Tgfbr2 in endothelial cells. f, Quantification the number of confocal images of brain slice EGFP$^+$ cells stained with EndoMT marker (p-SMAD3,KLF4 and α-SMA) at RP2D after specific knockout of Tgfbr2 in endothelial cells. g, FACS analysis the percentage of α-SMA$^+$ ECs using EC marker (CD31) at RP2D after specific knockout of Tgfbr2 in endothelial cells. h,Quantification the FACS analysis the percentage of α-SMA$^+$ ECs using EC marker (CD31)at RP2D after specific knockout of Tgfbr2 in endothelial cells. i, Confocal images of brain slice EGFP$^+$&CD31$^-$ cells stained with ECs marker (CD31) at RP34D after specific knockout of Tgfbr2 in endothelial cells. j, Quantification the confocal images of brain slice EGFP$^+$&CD31$^-$ cells stained with ECs marker (CD31) at RP34D after specific knockout of Tgfbr2 in endothelial cells. k, FACS analysis EGFP$^+$&CD31$^-$ cells stained with ECs marker (CD31) at RP34D after specific knockout of Tgfbr2 in endothelial cells. Data presented as mean ± SEM, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

Fig.5

Fig.5 Reducing E-pericytes through specific knockout of Tgfbr2 in endothelial cells can impede the recovery of stroke during chronic reperfusion. a**,** Images of whole brain slice injection with Evans blue after stroke at RP7D, RP21D, RP34D and 118D. b, Quantification the whole brain slice injection with Evans blue after stroke at RP2D, RP7D, RP21D and RP34D. c,Confocal images of brain slice neuron stained with marker (Neun) at RP34D. d, Quantification the confocal images of brain slice neuron stained with marker (Neun) at RP34D. e, Confocal images of brain slice pericyte stained with marker (CD13) at RP34D. f, Quantification the confocal images of brain slice pericyte stained with marker (CD13) at RP34D. g, Quantification the confocal images of brain slice the ration of ECs/pericyte (CD13) at RP34D. h, Confocal images of brain slice blood vessel stained with ECs marker (CD31) at RP34D. i, Quantification the confocal images of brain slice blood vessel stained with ECs marker (CD31) at RP34D. j, FACS analysis the percentage of CD31$^+$ ECs at RP34D. k, Quantification the FACS analysis the percentage of CD31$^+$ ECs at RP34D. i, Images of whole brain CBF after stroke at 34D. m, Quantification the images of whole brain CBF after stroke at 34D. n, Images of whole brain volume after stroke at 34D. o, Quantification the images of whole brain volume after stroke at 34D. p, Quantification the time of pole test after stroke at RP0D, RP2D, RP7D, RP14D, RP21D and RP34D. q, Quantification the time of beam walking test after stroke at RP0D, RP2D, RP7D, RP14D, RP21D and RP34D. r, Quantification the time of turning test after stroke at RP0D, RP2D, RP7D, RP14D, RP21D and RP34D. s, Quantification the time of adhesive removal test after stroke at RP0D, RP2D, RP7D, RP14D, RP21D and RP34D. Data presented as mean ± SEM, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

Extended Data Fig.1

Extended Data Fig. 1 The gate of FACS for the next of experiments. a, The flow of FACS using CD45+&Viability+ cells for the next analysis and soritng.

Extended Data Fig.2

Extended Data Fig. 2 The endothelial cells are capable of undergoing cell fate transition after stroke. a, Confocal images of genetic tracing strategy using *Cdh5CreERT2;Ai47* or *Ai14* line without Tamoxifen. b, FACS analysis genetic tracing strategy using *Cdh5CreERT2;Ai47* or *Ai14* line without Tamoxifen(N=4). c-e, Confocal images of brain slice vasculatures traced for 1.6 months, stained with EC markers (GLUT1, ERG and VE-Cadherin). f, Quantification of confocal images of brain slice EGFP cells stained with EC markers (GLUT1, ERG and VE-Cadherin). g-j, Confocal images of brain slice EGFP cells with other cells markers (NeuN:neuron; Doublecortin:immature neuron; Iba1: microglia; GFAP;astrocyte; CD68:microglia and monocyte-macrophage; CD45:immuce cell). m, Quantification of confocal images of brain slice EGFP cells stained with other cells markers. n-p, Confocal images of brain slice EGFP cells stained with pericytes markers (CD13, Caldesmon and α-SMA). q, Quantification of confocal images of brain slice EGFP cells stained with pericytes markers. r, Images of the CBF of mice after MCAO:2H with LSCI. s, Quantification of the CBF of mice after MCAO:2H with LSCI. t, Images of the whole brain after MCAO:2H. u-v, Confocal images of brain slice EGFP⁺ cells stained with EC marker (VE-Cadherin , ERG, GLUT1 and IB4). w-y, Confocal images of brain slice EGFP cells stained with ECs markers (CD31) at RP81D, RP232D and RP514D. z, Quantification of confocal images of brain slice EGFP cells stained with ECs markers (CD31) at RP81D, RP232D and RP514D. a1, Schematic diagram show of the location of EGFP⁺&CD31⁻ cells at different RP. b1, Confocal images of brain slice EGFP⁺&CD31⁻ cells morphology. c1, Quantification of confocal images of brain slice EGFP⁺&CD31⁻ cells morphology. d1-f1, Quantification of confocal images of brain slice EGFP⁺&CD31⁻ cells morphology

689   (process length, primary process and diameter of soma). Data presented as mean ±

690   SEM, ** P < 0.01, *** P < 0.001, **** P < 0.0001.

a *Cdh5CreERT2:Ai47*

EGFP / Bone mallow cells without RBCs | Hoechst | Merge

b Bone marrow / Blood

monocyte / neutrophil

EGFP | CD11b | CD11b

c
P2 — Intraventricular injection of *AAV-DIO-EGFP* virus
1.5M — TAM
2.5M — MCAO
RP:34Days
3.5M — Analysis

d *Cdh5CreERT2:Ai47* | *Cdh5CreERT2* | *Cdh5CreERT2* + Virus

e % of EGFP+ cells in CD45+ cells
Cdh5CreERT2:Ai47 / *Cdh5CreERT2* / *Cdh5CreERT2* +virus

f *Cdh5CreERT2; AAV-DIO-EGFP* (Contralateral side)
EGFP | CD31 | EGPP CD31

g % of EGFP+ Cells
CD31 / VE-Cadherin / ERG / GLUT1

h *Cdh5CreERT2; AAV-DIO-EGFP* (Ischemic side)
EGFP | CD31 | EGFP CD31

i without Tamoxifen *WT* | Tamoxifen *Cdh5CreERT2:Ai47* | Tamoxifen+Radiation *Cdh5CreERT2:Ai47*

j % of EGFP+ cells in total bone marrow cells
WT / Tamoxifen / Tamoxifen+Radiation

k *Cdh5CreERT2:Ai47* Contralateral side | *Cdh5CreERT2:Ai47* Ischemic side
EGFP | CD31 | EGFP CD31 | EGFP CD31 Hoechst

l EGFP+ non-ECs cells / 10 mm²
Contralateral hemisphere / Ischemic hemisphere / Contralateral hemisphere / Ischemic hemisphere
Radiation

m Ischemic hemisphere without radiation | Ischemic hemisphere with radiation
CD31 / EGFP

n % of EGFP+ non-ECS in total EGFP+ cells
Ischemic hemisphere / Ischemic hemisphere + Radiation

Extended Data Fig. 3 EGFP$^+$&CD31$^-$ cells do not originate from peripheral blood non-specific cells labeled in Cdh5CreERT2 mouse. a, Confocal images of bone marrow cells without RBCs in Cdh5CreERT2 mouse. b, FACS analysis the EGFP+ cells from bone marrow cells without RBCs in Cdh5CreERT2 mouse. c, Illustration of virus strategy using *Cdh5CreERT2* line. d, FACS analysis the EGFP+ cells in bone marrow cells without RBCs form in Cdh5CreERT2 mouse with virus. e, Quantification of FACS analysis the EGFP+ cells in bone marrow cells without RBCs form in Cdh5CreERT2 mouse with virus. f, Confocal images of EGFP+ cells in brain from Cdh5CreERT2 mouse stained with EC marker CD31. g, Quantification of Confocal images of EGFP+ cells in brain from Cdh5CreERT2 mouse stained with EC markers(CD31, VE-Cadherin, ERG and GLUT1). h, Confocal images of EGFP+ cells in brain from Cdh5CreERT2 mouse stained with EC marker CD31. i, FACS analysis the percentage of EGFP$^+$ cells after rebuilding bone marrow using radiation. j, Quantification of FACS analysis the percentage of EGFP$^+$ cells after rebuilding bone marrow using radiation. k, Confocal images of EGFP$^+$&CD31$^-$cells in Cdh5CreERT2 with rebuilding bone marrow using radiation. l, Quantification of confocal images of EGFP$^+$&CD31$^-$cells in Cdh5CreERT2 with rebuilding bone marrow using radiation. m, FACS analysis EGFP$^+$&CD31$^-$cells in Cdh5CreERT2 with rebuilding bone marrow using radiation. n, Quantification of FACS analysis EGFP$^+$&CD31$^-$cells in Cdh5CreERT2 with rebuilding bone marrow using radiation. Data presented as mean ± SEM, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

Extended Data Fig.4

Extended Data Fig. 4 The identification of EGFP$^+$&CD31$^-$ cells is pericyte-like cells.

a, Confocal images of EGFP$^+$&CD31$^-$cells stained with Ki67. b, Quantification of confocal images of EGFP$^+$&CD31$^-$cells stained with Ki67(N=2, 15 brain slice*2). c, Confocal images of EGFP$^+$&CD31$^-$cells stained with pericyte marker(NG2). d-f, Confocal images of EGFP$^+$&CD31$^-$cells stained with other cells markers (NeuN:neuron; Doublecortin:immature neuron; Iba1: microglia; α-SMA:SMC; GFAP;astrocyte; CD68: microglia and monocyte-macrophage). g, Confocal images of EGFP$^+$&CD31$^-$cells in Cdh5CreERT2 mice injection virus. h, Confocal images of EGFP$^+$&CD31$^-$cells stained with pericyte marker(CD13) in Cdh5CreERT2 mice injection virus. i, Confocal images of EGFP$^+$&CD31$^-$cells stained with pericyte marker(PDGFRβ) in Cdh5CreERT2 mice injection virus. j, Confocal images of Tie2Dre;Msfd2a mice brain slice stained with EC marker(CD31). k, Quantification of confocal images of Tie2Dre;Msfd2a mice brain slice stained with EC marker(CD31). l, Confocal images of brain slice from Ai47 mice injection with AAV-Bi30-Cre virus , stained with EC marker(CD31). l, Quantification of confocal images of brain slice from Ai47 mice injection with AAV-Bi30-Cre virus , stained with EC marker(CD31). n, Confocal images of EGFP$^+$ cells in Tie2Dre;Msfd2a mice brain slice stained with EC marker(CD31) and pericyte marker (CD13). o, Confocal images of EGFP$^+$ cells in Ai47 mice injection with AAV-Bi30-Cre virus brain slice stained with EC marker(CD31) and pericyte marker (CD13). Data presented as mean ± SEM, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

Extended Data Fig. 5 TGFβ-pSMAD3-EndoMT signaling involved in E-pericytes

formation after storke. a, Confocal images of brain slice EGFP⁺ cells stained with

EndoMT marker (KLF4) at RP2D,RP8D and RP34D. b, Quantification the number of

confocal images of brain slice EGFP⁺ cells stained with EndoMT marker (KLF4) at

RP2D,RP8D and RP34D.(N=3). c, Quantification the intensity of confocal images of

brain slice EGFP⁺ cells stained with EndoMT marker (KLF4) at RP2D,RP8D and

RP34D(N=3). d, Confocal images of brain slice EGFP⁺ cells stained with EndoMT

marker (α-SMA) at RP2D,RP8D and RP34D. e, Quantification the number of confocal

images of brain slice EGFP⁺ cells stained with EndoMT marker (α-SMA) at at

RP2D,RP8D and RP34D. f, Quantification the indensity of confocal images of brain slice

EGFP⁺ cells stained with EndoMT marker (α-SMA) at at RP2D,RP8D and RP34D. g,

FACS analysis the percentage of α-SMA⁺ ECs using EC marker (CD31) at RP2D.

h,Quantification the FACS analysis the percentage of α-SMA⁺ ECs using EC marker

(CD31)at RP2D. i, Confocal images of brain slice EGFP⁺cells stained with EndoMT

markers (p-SMAD3, KLF4 and α-SMA) at RP2D after using TGFBR2 inhibitors. j,

Quantification the confocal images of brain slice EGFP⁺cells stained with EndoMT

markers (p-SMAD3, KLF4 and α-SMA) at RP2D after using TGFBR2 inhibitors. k, FACS

analysis the percentage of α-SMA⁺ ECs using EC marker (CD31) at RP2D after using

TGFBR2 inhibitors. l, Quantification the FACS analysis the percentage of α-SMA⁺ ECs

using EC marker (CD31) at RP2D after using TGFBR2 inhibitors. m, Confocal images of

brain slice EGFP⁺ cells stained with EC marker (CD31) at RP34D after using TGFBR2

inhibitors. n, Quantification the confocal images of brain slice EGFP⁺ cells stained with

EC marker (CD31) at RP34D after using TGFBR2 inhibitors. o, FACS analysis brain

slice EGFP$^+$ cells stained with EC marker (CD31) at RP34D after using TGFBR2

inhibitors. p, Quantification the FACS analysis brain slice EGFP$^+$ cells stained with EC

marker (CD31) at RP34D after using TGFBR2 inhibitors. Data presented as mean ±

SEM, ** P < 0.01, *** P < 0.001, **** P < 0.0001.

Extended Data Fig.6

Extended Data Fig. 6 Knocking out Tgfbr2 ECs from adult mouse does not affect the BBB function. a, Images of the whole brain injection with Evans blue in normal state. b, confocal images of brain slice CD31$^+$ ECs stained with BBB marker (ZO-1) in normal state. c, Quantification of confocal images of brain slice CD31$^+$ ECs stained with BBB marker (ZO-1) in normal state. d, Images of the thickness of basement membrane brain slice useing EM in normal state. e, Quantification of images of the thickness of basement membrane in brain slice using EM in normal state. f, Images of the endothelial tight junction in brain slice using EM in normal state. g, Images of the endothelial transcytosis in brain slice using EM in normal state. Data presented as mean ± SEM, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

1.   Kuriakose, D. & Xiao, Z. Pathophysiology and Treatment of Stroke: Present

Status and Future Perspectives. *Int J Mol Sci* **21** (2020).

2.   Martino, G. How the brain repairs itself: new therapeutic strategies in

inflammatory and degenerative CNS disorders. *Lancet Neurol* **3**, 372-378 (2004).

3.   Zhao, L.R. & Willing, A. Enhancing endogenous capacity to repair a stroke-

damaged brain: An evolving field for stroke research. *Prog Neurobiol* **163-164**, 5-

26 (2018).

4.   Hall, C.N. *et al.* Capillary pericytes regulate cerebral blood flow in health and

disease. *Nature* **508**, 55-60 (2014).

5.   Díaz-Flores, L., Gutiérrez, R., Varela, H., Rancel, N. & Valladares, F.

Microvascular pericytes: a review of their morphological and functional

characteristics. *Histol Histopathol* **6**, 269-286 (1991).

6.   Pallone, T.L. & Silldorff, E.P. Pericyte regulation of renal medullary blood flow.

Exp Nephrol **9**, 165-170 (2001).

7.   Simon, C., Lickert, H., Götz, M. & Dimou, L. Sox10-iCreERT2 : a mouse line to

inducibly trace the neural crest and oligodendrocyte lineage. *Genesis* **50**, 506-

515 (2012).

8.   Etchevers, H.C., Vincent, C., Le Douarin, N.M. & Couly, G.F. The cephalic neural

crest provides pericytes and smooth muscle cells to all blood vessels of the face

and forebrain. *Development* **128**, 1059-1068 (2001).

9.   Yu, Q.C., Song, W., Wang, D. & Zeng, Y.A. Identification of blood vascular

endothelial stem cells by the expression of protein C receptor. *Cell Res* **26**, 1079-

1098 (2016).

10. Wang, Y. *et al.* Ephrin-B2 controls VEGF-induced angiogenesis and

lymphangiogenesis. *Nature* **465**, 483-486 (2010).

11. Science, A.I.f.B. TECHNICAL WHITE PAPER: TRANSGENIC

CHARACTERIZATION. *Allen Brain Atlas* **9** (2015).

12. Kilani, B. *et al.* Comparison of endothelial promoter efficiency and specificity in

mice reveals a subset of Pdgfb-positive hematopoietic cells. *J Thromb Haemost*

17, 827-840 (2019).

13. Planas, A.M. Role of Immune Cells Migrating to the Ischemic Brain. *Stroke* **49**,

2261-2267 (2018).

14. Fernandez-Klett, F. *et al.* Early loss of pericytes and perivascular stromal cell-

induced scar formation after stroke. *J Cereb Blood Flow Metab* **33**, 428-439

(2013).

15. Pu W, et al. Genetic Targeting of Organ-Specific Blood Vessels. Circ Res. 2018

Jun 22;123(1):86-99.

16. Krolak T, et al. A High-Efficiency AAV for Endothelial Cell Transduction

Throughout the Central Nervous System. Nat Cardiovasc Res. 2022

Apr;1(4):389-400.

17. Piera-Velazquez, S. & Jimenez, S.A. Endothelial to Mesenchymal Transition:

Role in Physiology and in the Pathogenesis of Human Diseases. *Physiol Rev* **99**,

1281-1324 (2019).

18. Hulshoff, M.S., Xu, X., Krenning, G. & Zeisberg, E.M. Epigenetic Regulation of

Endothelial-to-Mesenchymal Transition in Chronic Heart Disease. *Arterioscler*

Thromb Vasc Biol **38**, 1986-1996 (2018).

19.   Armulik A, et al. Pericytes regulate the blood-brain barrier. Nature. 2010 Nov

25;468(7323):557-61. doi: 10.1038/nature09522. Epub 2010 Oct 13. PMID:

20944627.

20.   Daneman R, et al. Pericytes are required for blood-brain barrier integrity during

embryogenesis. Nature. 2010 Nov 25;468(7323):562-6.

21.   Goumans MJ, Mummery C. Functional analysis of the TGFbeta receptor/Smad

pathway through gene ablation in mice. Int J Dev Biol. 2000 Apr;44(3):253-65.

PMID: 10853822.