Ischemic brains regenerate vascular mural pericytes by recycling detached endothelial cells, depending on TGFβ signaling

Tingbo Li[1], Qingzhu Gao[2], Xuzhao Li[2], Zhu Zhu[2], Yuxiao Jin[2], Yiyi Zhang[2], Jiongfang Xie[3], Peiran Zhu[4], Bingrui, Zhao[2], Jie-Min Jia*[2, 5, 6, 7]

[1]College of Life Sciences, Zhejiang University, Hangzhou, Zhejiang, 310058, China.

[2]Laboratory of Neurovascular Biology, School of Life Sciences, Westlake University, Hangzhou, 310024, China.

[3]Key Laboratory of Structural Biology of Zhejiang Province, School of Life Sciences, Westlake University, Hangzhou, Zhejiang, 310024, China.

[4]School of Materials Science and Engineering, Zhejiang University, Hangzhou, China.

[5] Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou, 310024, China

[6]Key Laboratory of Growth Regulation and Translational Research of Zhejiang Province, School of Life Sciences, Westlake University, Hangzhou, 310024, China.

[7]Institute of Basic Medical Sciences, Westlake Institute for Advanced Study, Hangzhou, 310024, China.

*Correspondence: jiajiemin@westlake.edu.cn

Cerebral endothelial cells (ECs) fate and cellular sources for pericyte regeneration after stroke are unclear. by genetic tracing, we identified ECs undergoing cell-fate changing, likely through TGFβ-mediated transdifferentiation, into pericytes. Some of the migrating pericytes induced by stroke are derived from detached ECs. The EC-transformed pericytes can integrate into vasculature again, suggesting ischemic brains can recycle ECs from unfunctional vessels to replenish lost pericytes as an innate self-maintenance program sustaining repair phenomena.

Stroke kills millions each year and no effective treatment is available so far[1]. Brains possess an intrinsic, limited self-repair capacity[2, 3], e.g. spontaneous angiogenesis after stroke. However, understanding the definitive spontaneous repairing processes is largely incomplete. ECs form vascular tubes, while pericytes encircle the endothelial tube. The observation that more pericytes die faster than ECs may explain neurodegeneration exacerbation and brain self-recovery hindering after stroke because pericytes are a type of mural cells known to be critical for the blood-brain barrier and cerebral blood flow[4-6]. However, little is known about cellular sources for pericyte regeneration in the adult ischemic brain while it has been well documented that the cephalic neural crest gives rise to pericytes in all blood vessels of the face and forebrain during embryonic development[7, 8]. Many capillaries collapse after stroke, however, the fate of endothelial cells from these unfunctional vessels to be recycled or to die is unclear. Recently, Yu et al. have demonstrated that pericytes can be differentiated locally from a subgroup of ECs in multiple tissues under homeostasis[9]. However, the case in brains with or without ischemia remains unknown. Identification of origin cells for regenerated pericytes and EC's fate in

the ischemic brain is of importance for enhancing endogenous capacity to repair a stroke-

damaged brain.

To test if cerebral ECs differentiate into pericytes during adult angiogenesis as do those

in the mammary gland, skin, and retina[9], we employed a cell fate tracing system by

crossing EC-specific DNA recombinase driver line *Cdh5-CreERT2*[10] with triple EGFP

reporter (Ai47)[11] or tdTomato reporter (Ai14) (The Jackson Laboratory) (Fig. 1a). ECs and

their potential progeny cells are permanently labeled by EGFP or tdTomato fluorescent

proteins once tamoxifen (TAM) is provided (Extended Data Fig.1a). Consistent with

previous reports[9, 10, 12], these fluorescent proteins were restricted in ECs but not in any

other types of brain cells short-term after TAM exposure (4 - 7 days, Extended Data Fig.

1c-g). Upon long-term TAM exposure (Fig. 1b), although some of these EGFP+ ECs in skin

tissue change their fates into EGFP+ pericytes[9], brain EGFP+ ECs, however, seemed never

(Fig. 1c-i). By using immunofluorescent staining of typical pericyte markers (CD13 and

PDGFRβ), we did not observe any EGFP+ pericytes at various tracing periods, ranging

from 1.5 to 17.6 months (Fig. 1c-d, i). Whereas, all EGFP+ cells remained expressing

multiple classical endothelial markers, such as CD31, VE-cadherin, ERG, and GluT1,

despite up to a two-year tracing (Fig.1e-i). Together, these results demonstrated unlike in

the peripheral system and retina, cerebral ECs are not the cellular source for pericytes in

adult brains under homeostasis.

Ischemic stroke, modeled by middle cerebral artery occlusion (MCAO, Extended Data

Fig. 2), induced cerebral ECs detaching from vessels, gradually losing EC markers, basal

membrane protein, and undergoing drastic morphological changes such as losing hollow

structure and growing multiple processes from cell soma (Fig. 1j-k and Extended Data

Fig. 3a). This finding suggested that ECs might undergo cell fate changing in response to

ischemia attack. Thus, we referred to this subgroup of EGFP$^+$ cells as 'EGFP$^+$ non-EC

cells'. TAM was delivered to 1.5-month-old mice and then at least two weeks later we

performed MCAO surgery (Fig. 1b), aiming to ensure no extra residual TAM. After 2-hour

MCAO, sutures were withdrawn, permitting various periods of reperfusion, ranging from

8 all the way through 514 days (RP2D to RP514D) (Fig. 1b). These EGFP$^+$ non-EC cells

can be found anywhere possible in the ischemic region, showing no regional specificity

(Extended Data Fig. 3b). According to whether having secondary processes, we classified

EGFP$^+$ non-EC cells into two types: type I, short, and fewer primary processes, no

secondary processes (Fig. 1j, l, Extended Data Fig. 3c-e); type II, longer and more primary

branches with secondary ones (Fig. 1k, l, Extended Data Fig. 3c-e), while their cell bodies

were comparable in shape and size (Extended Data Fig. 3e). We found that the prevalence

of these two types of cells switched when reperfusion period prolonged. (Fig. 1l). This

morphological switching suggests type I becoming type II gradually. The prevalence of

EGPF$^+$ non-EC cells started to rise at RP8D, peaked by RP52D, and finally declined back

to a low level (Fig. 1m). This curve suggested that these transformed avascular EGFP$^+$

non-EC cells may die or be recruited back to vessels, becoming vascular cells again. We

termed this possibility a recycling process. Furthermore, FACS analysis performed on the

peak day (RP52D) revealed an EGFP$^+$ but CD31 negative population (EGFP$^+$ non-EC cells,

Fig. 1l). These EGFP$^+$ transformed cells accounted for ~0.4% of total EGFP$^+$ cells which

included ECs and non-EC cells (Fig. 1m). *Cdh5-CreERT2* may label a small portion of

neutrophils and monocytes in the bone marrow and circulation[12] (Extended Data Fig. 4

a-b), which are known to enter brain parenchyma shortly after stroke onset but dies out[13].

To discriminate whether parenchyma EGFP+ non-EC cells are derived from EGFP+ bone

marrow cells in *Cdh5CreERT2;Ai47* mice, we introduced a viral expression system

restricted to ECs preventing EGFP expression in hematopoietic cells (Extended Data Fig.

4c-e). Specifically, we injected cranially the Cre-dependent *AAV-DIO-EGFP* virus to

Cdh5-CreERT2 single-positive mice that do not carry *Ai47* or *Ai14* reporter (Extended

Data Fig. 4c), resulting in high labeling specificity to ECs (Extended Data Fig. 4f-g). Once

again, we found EGFP+ ECs transforming into non-EC cells after stroke (Extended Data

Fig. 4h). Notably, some of these cells extended their long processes and enwrapped

capillaries, reminiscent of pericyte characteristics (Extended Data Fig. 4h). Independent

radiation experiment was implemented to exclude the same possibility, we replaced

Cdh5CreERT2;Ai47 mice's bone marrow with wild type ones (Extended Data Fig. 5a-b).

Despite the depletion of EGFP+ bone marrow cells under this circumstance (Extended

Data Fig. 5a-b), we still found comparable EGFP+ non-EC cells in ischemic regions

(Extended Data Fig. 5c-f), demonstrating that EGFP+ immune cells were not the cell

source of EGFP+ non-EC cells in parenchyma. These results together indicate that a

small proportion (0.4%) of ECs detached from vessels in the parenchyma, lost EC

features, and changed their cellular identity after subjecting them to ischemia insult.

To assess whether these EGFP+ non-EC cells are derived from vascular stem

endothelial cells as reported previously[9] which involves cell proliferation, we traced cell

division events by injecting EdU for consecutive 34 days. No single EGFP+ non-ECs were

found to be EdU positive after examining 15 brain slices from two mice (Fig. 2a-b).

similarly, neither Ki67 signals were found (Fig. 2c-d). Thus, we concluded these EGFP[+]

non-EC cells may not be progenies of stem cells, suggesting that a transdifferentiation

process may be required.

By employing a list of cell type markers, we identified that 90% of EGPF[+] non-EC cells

were pericytes (namely, 'EGFP[+] pericytes') as they were positive for typical pericyte

markers of CD13, PDGFRβ, and NG2 (Fig. 3a-g), as well as that they were negative for all

of the markers of immature and mature neurons (doublecortin and NeuN), astrocytes

(GFAP), CNS macrophages (Iba1, CD68, and F4/80), myeloid cells (CD45), and smooth

muscle cells (α-SMA) (Extended Data Fig. 6a). The cell identity of the rest 10% of EGFP+

non-EC cells remained unknown (Fig. 3g) though as these cells were not labeled by any

cell markers that we used. Furthermore, more than a quarter of EGFP[+] pericytes were

recruited back to vessels (Fig. 3e-f and h, Movie1), suggesting ECs were fully recycled

not only by expressing pericyte markers but also by locating the right anatomical position.

A transformed tdTomato[+] non-ECs (here, Ai14 reporter was used instead of Ai47) that

were also tracked by *AAV-DIO-EGFP* virus exhibited one process attaching a capillary

while the other did not (Extended Data Fig. 6b), implying capturing a recruiting moment.

It has been repeatedly reported that pericytes detach from vessels after stroke[14]. We

found actually a third of these avascular, migrating pericytes were derived from ECs (Fig.

3i). Further analysis revealed that 23.3% of total vascular pericytes were EGFP[+] (Fig. 3h),

suggesting ECs are a fair contributor cell source for pericyte regeneration. The virus-

tracked ECs also can generate pericytes, coinciding with results from genetic tracing

(Extended Data Fig. 6c-d). These results together demonstrate that after ischemic stroke,

ECs transdifferentiated into pericytes, likely capable of integrating back into the cerebral

vasculature.

In addition, during EdU assay, we found that there were EdU positive pericytes that

were EGFP negative in both vascular and migrating pericytes (Fig. 3k), suggesting

alternate cell sources for pericyte regeneration except for EC transdifferentiation. Each

source contributed 20% of pericytes respectively (Fig. 3l). Therefore, we inferred that at

RP34D, 40% of total pericytes in ischemic regions were not pre-existing ones but

regenerated ones (Fig. 3l). Together, these findings indicated that ischemic brains have

more than one way to replenish pericytes when exerting self-repair after injury.

Next, we investigate the potential molecular signaling pathway that might mediate this

EC-to-pericyte transdifferentiation. Since TGFβ signaling is well documented for cell fate

changing including the endothelial mesenchymal transition (EndMT) process, a newly

recognized type of transdifferentiation[15, 16]. Therefore, we injected LY364947, a TGFβRI

inhibitor when MCAO was performed, followed by additional daily injections for

consecutive 34 days of reperfusion. We found significantly fewer EGFP+ non-EC cells

appeared overall (Extended Data Fig. 6e-f). FACS analysis further confirmed this

conclusion quantitively (Fig. 3m-n). These results strongly suggested that TGFβ signaling

in ECs is centrally important. Together, under ischemic conditions, ECs exploited TGFβ

signaling to transform ECs into pericytes during reperfusion to replenish the shrunk

pericyte pool occurring in the acute ischemia period.

Overall, using mouse genetic tracing tools combined with virus infection and

pharmacology, we demonstrated adult ischemic brains spontaneously repair

microvasculatures by regenerating new pericytes that are transformed from ECs after detaching from vessels. Distinct from the previous findings in peripheral organs in which a small group of endothelial stem cells Procr+ ECs differentiate into pericytes during adult angiogenesis under homeostasis[9], ECs do not become pericytes in normal brains but transdifferentiate into pericytes without cell division. The death rate is dozens of times higher in pericytes than in ECs within 24 hours after a transient focal ischemia[4], suggesting abundant capillaries become damaged by being naked. What is the fate of these naked capillaries made up by ECs? The present study suggests that these naked vessels may release ECs, some of which may not die but rather be recycled and repurposed by the ischemic brain. By observing type I simpler form switching to type II more complex EGFP+ non-EC cells (Fig. 1l) and the medium stage of the transformed EFGP+ pericyte (Extended Data Fig. 6b), we extrapolate them likely being recruited and serve as building blocks to repair the existing damaged vasculatures. This could be an efficient system to reuse brain cells that have lost their original functions. Further studies are required to live imaging the whole dynamic process during reperfusion periods.

Identifying new cellular sources for pericyte replenishment is important not only for stroke but also for many other either acute or chronic brain injuries. This study may suggest a novel therapy for generating EC-derived pericyte without involving stemness which has long been a concern for tumorigenesis.

Methods

Experimental design. Transgenic Cdh5CreERT2 transgenic mice[10] cross with reporter mice *Ai47 (EGFP^{f/f}) reporter* or *Ai14 (tdTomato^{f/f}) reporter mice*. Thus, ECs are labeled in green or red initially. TAM was delivered when mice were 1.5 months old (6 weeks), i. e.

genetic tracing started. After waiting for 2 weeks, MCAO was performed. After a 2-hour

occlusion, reperfusion began. Contralateral sides of the ischemic brain were used as

objects for homeostasis conditions, except for the time point of tracing 4 days. Thus, it

was traced for 22 days in the PR8D mice (N = 3 mice), 1.6 months in RP34D (N = 27

mice), 2.8 months in RP71D (N = 3 mice), 14 months in RP408D (N = X mice), 8.2 months

in RP232D (N = 3 mice), 14 months in RP408D (N = 2 mice),17.6 months in RP51D mice

(N = 2 mice). The tracing time point of 1.6 months (RP34D) was the most frequently

examined. FACS and immunohistochemistry were the major two techniques used for cell

identity confirmation.

Animal care and tissue dissection.

All animal experiments were carried out in accordance with protocols approved by the

Institutional Animal Care and Use Committee (IACUC) at the School of Life Sciences,

Westlake University. Wild-type C57BL6/J mice were purchased from the Laboratory

Animal Resources Center of Westlake University. Cdh5CreERT2 mice were a gift from

Le-Ming Zheng (Peking University). Gt(ROSA)26Sor$^{tm47(CAG-EGFP*)Hze}$ (Ai47) was share by

Zilong Qiu (ION). We crossed Cdh5Cre-ERT2 mice to Ai47 mice to generate

Cdh5CreERT2:Ai47 mice. Standard chow and water were provided to mice *ad libitum*.

Four mice were housed in each cage. All animals were housed in a standard animal room

with a 12/12-hour light/dark cycle at 25 °C. Both male and female mice were used in this

study. Mice were anaesthetized by intraperitoneal injection of 无巴比妥钠 (10 mg/kg?),

and brain dissection was performed immediately for FACS experiments or cardiac PFA

perfusion was performed for immunohistochemistry experiment. After fixation, brains

were dissected and kept in 4% PFA for additional 4 hours and then transferred to PBS

until further processing.

MCAO

Adult mice were anesthetized with pentobarbital sodium (100 mg/kg) and body

temperature was maintained during surgery with a heating pad. A midline neck incision

was made, the right common carotid artery was carefully separated from the vagus nerve,

and the artery was ligated using a 5.0-string. A second knot was made on the left external

carotid artery. The right internal carotid artery (ICA) was isolated, and a knot was left

loose with a 5.0-string. This knot was not tightened until the intraluminal insertion was

done. A small hole was cut on the common carotid artery before it was bifurcated to the

external carotid artery and ICA. A silicon-coated monofilament (tip diameter = 230 μm,

Doccol Corporation, 602256PK5Re) was gently advanced into the ICA until it stopped at

the origin of the MCA in the circle of Willis. The third knot on the ICA was closed to fix the

filament in position. During MCA occlusion (2 h), mice were kept in a warm cage in which

the temperature was maintained at 35 °C.

Laser speckle contrast imaging (LSCI)

LSCI provides a measure of blood flow velocity by quantifying the extent of blurring of

dynamic speckles caused by the motion of red blood cells through the vessels. Briefly,

mice were placed under an RFLSI III device (RWD Life Sciences) before and after the

suture was successfully inserted through CCA. The skull over both hemispheres was

exposed by making an incision along the midline of the scalp. When a 785 nm laser is

used to illuminate the brain, it produces a random interference effect that represented

blood flow in the form of a speckle pattern. Scattering light was detected by a charge-

coupled device (CCD) camera, and the images were acquired by custom software from

RWD Life Sciences Company. For each acquisition, a total of 160 images, each of which

measured 2048 ×2048 pixels, were collected at 16 Hz.

Tamoxifen

Tamoxifen, 10 mg/ml dissolved in 200 µl corn oil, was administrated intragastrically for 4

consecutive days. Prior to further experiments, fluorescent protein expressions in mouse

ear vasculatures were first examined. Tamoxifen was administrated at least 2 weeks

before MCAO surgery.

Immunohistochemistry

Cryosections of fixed mouse brains (50 µm) were handled free-floating and washed for

5 minutes in PBS prior to further procedures. Tissue sections were permeablized and

blocked in the PBS containing 0.5% Triton and 5% BSA at room temperature for 1 hour.

Then, brain sections were incubated with a different primary antibodies, including

CD31(1:400, BD, Cat# 557355), VE-Cadherin(1:200, BiCellScientific, Cat# 00105),

ERG(1:300, Sigma, Cat# ab110639), GLUT1 (1:300, Sigma, Cat# 07-1401), NeuN

(1:300, Merck, Cat# ABN90P), Doublefortin (1:300, Santa clus, Cat# sc-271390), Iba1

(1:300, HUABIO, Cat# ET1705-78), GFAP (1:300, Proteintech, Cat# CL488-60190),

PDGFRβ (1:300, ThermoFisher, Cat# 14-1402-82), NG2 (1:300, Merck, Cat# AB5320)

,CD68 (1:300, BIO-RAD, Cat# MCA1957T), CD45-APC (1:200, Biogend, Cat# 103112),

Ki67 (1:300, thermoscientific, Cat# RM-9106-S0), F4/80 (1:300, Cell Signaling, Cat#

30325), CD13 (1:300, R&D system, Cat# AF2335), α-SMA-CY3 (1:300, Sigma, Cat#

C6198), which were all diluted in the blocking solution (1% percelin, 5% BSA, and 0.1%

Triton in PBS) at 4 °C overnight. After rinsing in PBS 3 times for 5 minutes each, brain

slices were incubated with secondary antibodies accordingly, including donkey anti-

mouse IgG (A21202, Thermo Fisher Scientific, 1:1,000) and donkey anti-rabbit IgG

(A10040, Thermo Fisher Scientific, 1:1,000). They were diluted in the blocking solution.

After a 2-hour incubation, brain slices were washed with PBS three times for 15 minutes

each. Finally, Hoechst counterstaining was performed for each specimen. After mounting

in an anti-fade mounting medium, images were acquired by using a Zeiss LSM800 laser-

scanning confocal microscope with a 20X or 63X objective, with or without Airyscan

mode.

Flow cytometry

Primary cell preparation was conducted as described below. Mouse brains were

dissected into cold HBSS with glucose and BSA (HBSS^{++}, Vendor info) were cut into small

pieces by sterile scissors, followed by a centrifuge at 1600 rpm/min for 5 mins to get rid

of supernatant. The minced tissues after the centrifuge were incubated with collagenase

IV (2.5 mg/ml, ROCHE, Cat# 11088858001) at 37 °C with gentle rotation for 30 min in

2 ml of HBSS^{++}. Next, samples were passed through a 70-μm filter and washed with

HBSS^{++}. To obtain higher cell yield, grind the tissue on the 70-μm filter using a syringe

piston and guarantee all tissue pass the filter. The following antibodies in 1:200 dilutions

were used: CD31-PE-594 (1:200, Biolegend, Cat# 102520), CD45-APC (1:200, Biogend,

Cat# 103112), Ly6C-APC (1:200, Biolegend, Cat# 128016), CD11b-PE (1:200,

Biolegend, Cat#101212), Viability dye (1:1000, ThermoFisher, Cat#65-0863-14).

Antibody incubation was performed on ice for 30min in HBSS with glucose and BSA. All

analyses were performed with CytoFLEX LX-5L1(Beckman). FlowJo x software was used

for data analyses.

EdU injection

268 EdU (200 mg/kg, Alfa Aesar Chemical) was injected intraperitoneally for consecutive 34

269 days since when MCAO was performed. EdU was initially dissolved in DMSO at the stock

270 concentration of XX mg/ml and was diluted with saline prior to injection.

Virus injection

272 The following adenovirus vector *AAV-CAG-DIO-EGFP* was purchased for ShuMi, Wuhan,

273 China (PT-0168). Half of the one-micron virus was delivered to postnatal day 2 mouse

274 pups by intracerebroventricular injection to both sides. After 1.5 months, tamoxifen was

275 administrated. One month later, these mice were subjected to MCAO to induce ischemic

276 stroke.

Radiation and bone marrow transplantation

278 At two weeks post tamoxifen injection, adult *Cdh5CreERT2:Ai47* mice received radiation

279 (total: 11Gy, 2 times, 5.5Gy per time, interval 2 hours) and bone marrow cell

280 transplantation from wild-type mice with the same genetic background. RBCs were

281 removed from the bone marrow cells by using ACK Lysis Buffer (4 Million cells/mouse).

282 These mice were subjected to MCAO 2 months after bone marrow reconstruction.

Mouse cranial imaging by two-photon microscopy

Adult mice were anesthetized with pentobarbital sodium, and analgesia was provided by

subcutaneous injection of 0.2% meloxicam. The scalp was removed, and the skull was

exposed and cleaned. A dental drill with a 0.6-mm-diameter bit was used to engrave and

thin the bone around the circular craniotomy area at a size of 3 mm. The piece of skull

was carefully peeled off with fine-pointed forceps, and a cranial window was generated.

A coverslip with a 3-mm diameter was placed on the cranial window, and its perimeter

was completely sealed with a 1% agarose gel. The metal headplate was glued onto skull

with dental acrylic, through which the mouse brain was fixed on a headplate holder. The

headplate holder together with the mouse was placed under a two-photon microscope

(FVMPE-RS, Olympus). The cerebral vasculature *Cdh5CreERT2:Ai47* mice was

visualized and imaged once a day before and after stroke. All surgeries were performed

with sterilized instruments and an environment.

Statistics.

The quantified data in all figures were analyzed with GraphPad Prism 7.0 (La Jolla, CA,

USA) and presented as the mean ± SEM with individual data points shown. Unpaired two-

tailed Student's t-test was used for assessing the statistical significance between two

groups. Statistical significance was determined by calculation of p-value (*$p < 0.05$, **$p <$

0.01, ***$p < 0.001$, and ****$p < 0.0001$, ns: not significant). The repetition of our data is

independent biological replicates and the number of replicates for each experiment is

noted in the corresponding figure legend.

Data availability

The source data underlying the graphs and charts shown in the figures and tables are

provided in Supplementary Data 1. All data generated or analyzed during this study are

included in this published article (and its supplementary information file).

Acknowlegements

We thank Xuzhao Li, Zhu Zhu, Dongdong Zhang, Jinze Li, Jiayu Ruan, Lili Zhou for their

constructive discussions. We thank Danping Lu for her responsive and timely purchasing

support, Jiongfang Xie for sharing AAV virus. We thank the animal facility for its technical

assistance with rodent housing, Biomedical Research center platform for technique

support. This work is supported by the National Natural Science Foundation of China

(31800864 and 31970969 to J.-M.J, 82101475 to Z.Z.), Westlake startup funds, Westlake

Education Foundation and MRIC funds (103536022011) to J.-M.J.; Westlake Education

Foundation to B.D.

Author contributions

J.-M.J. conceived the research. T.L. and J.-M.J. designed experiments. T.L. performed

all the experiments and data quantification, except for the virus cranial injection. Y.W.

performed FACS together with T.L.. Z.Z. conducted some stroke models. Q.G. and B.Z.

conducted mouse genotyping.

Competing interests

All authors declare no competing interests.

Figure legends

Fig. 1 Genetic tracing of cerebral Cdh5$^+$ ECs demonstrates that they do not contribute to pericyte generation during adult angiogenesis but change cell fate to non-EC identity after ischemic stroke. a, Illustration of genetic tracing strategy using *Cdh5CreERT2;Ai47* or *Ai14* line. **b**, Experimental design for tracing periods and reperfusion periods following 2-hour ischemia. **c-h**, Confocal images of brain slice vasculatures traced for 1.6 months, stained with pericyte markers (CD13 and PDGFβR) and EC markers (CD31, VE-Cadherin, ERG, and GluT1). **i**, Summary of EGFP$^+$ cell's identity confirmation. CD13 antibody staining used brain slices from 6 mice, two mice for each time point of 1.6 months, 8.2 months, and 14 months; PDGFβR staining used two mouse brains traced for 1.6 months; Mice used for CD31 antibody staining was summarized after examining 27 mice that included all the time points shown in **b**; for VE-Cadherin, ERG, and GluT1 antibody staining, each of them used 3 mice that were traced for 1.6 months. **j-m**, Confocal images of ischemic brain region from mice at RP34D (correspondingly, trace period 1.6 months), showing type I (**j**) and type II(**k**) EGFP$^+$CD31$^-$ cells (named as EGFP$^+$ non-EC cells). **l**, Quantification of percentages of each type of EGFP$^+$ non-EC cells over different reperfusion periods. N = 3 mice for each time point, excepting only 2 mice for RP408D and RP514D. **m**, The time course of EGFP$^+$ non-EC cell densities in the ischemic regions. N = 3 mice for each time point, excepting only 2 mice for RP408D and RP514D. **n**, FACS analysis of EGFP$^+$ non-EC cells from contralateral control hemisphere (left plot) and ischemic side (right plot) of the same mouse at RP34D. **p**, Quantification of EGFP$^+$ non-EC cells for **o**. N = 4 mice. Data presented as mean ± SEM, ** P < 0.01, *** P < 0.001, **** P < 0.0001

Fig. 2 EGFP⁺ non-EC cell generation does not involve cell proliferation. **a**, Representative confocal images of a brain slice from the RP34D mouse that received EdU daily injection for 34 days. **b**, Quantification of EdU⁺ cells in total Hoechst⁺ cell population and EdU⁺ EGFP⁺ double positive and CD31⁻ cells in the total EGFP⁺ non-EC cell population. n = 15 brain slices from two mice. **c**, Representative confocal images of cell division marker Ki67 immunostaining from the RP34D mouse. **d**, Quantification of Ki67⁺ cells in total Hoechst⁺ cell population and Ki67⁺ EGFP⁺ double positive and CD31⁻ cells in the total EGFP⁺ non-EC cell population. n = 15 brain slices from three mice.

Fig. 3 Majority of EGFP⁺ non-EC cells transdifferentiate into pericytes in the

ischemic brain, which was attenuated by TGFβRI inhibitor. a-**f**, Representative

confocal images of avascular EGPF⁺ pericytes (**a**-**c**) and vascular EGFP⁺ pericytes (**d**-**f**).

g, The transforming efficiency of EGFP⁺ non-EC cell to pericytes. **h**, Quantification of the

percentages of vascular and avascular EGFP⁺ pericytes. N = 3 mice. **i**, The percentage

of avascular EGFP⁺ pericytes in the total migrating pericyte population after stroke. **j**, The

contribution of vascular EGFP⁺ pericytes to the total vascular pericyte population. **k**,

Confocal images of EGFP negative pericytes that were EdU positive after stroke. **l**, The

individual contributions of EGFP⁺ & EdU⁻ pericytes and EGFP⁻ & EdU⁺ pericytes to the

total pool of pericytes in ischemic regions. **m**, FACS plot comparison of EGFP⁺ non-EC

cell number in mice that received daily injection DMSO or TGFβRI inhibitors for 34 days

since when MCAO was performed. **n**, Quantification for **m**. N = 2 mice. All the mice used

were at RP34D, correspondingly to the trace period of 1.6 months.

Extended Data Fig. 1 The specificity of *Cdh5CreERT2* driver mouse line in ECs for a 4-day tracing period. a-b, Genotyping results of two *Cdh5CreER:Ai47* mice (**a**), one with TAM injection (right in **b**) and the other with no TAM (left in **b**). c. FACS identification of EGFP$^-$ and EGFP$^+$ cells population from individual two mice with TAM (right plot) or without TAM injection (left plot). **d**, FACS analysis of EC labeling efficiency of TAX-dependent EGFP expression. **e**. All EGFP$^+$ cells were CD31$^+$ at 1.6 months of the tracing period. N = 4 mice. **f**. Confocal images of brain sections from *Cdh5CreER:Ai47* mouse after staining with markers for a variety of cell types, NeuN for mature neurons, doublecortin for immature neurons, Iba1 and CD68 for CNS macrophages, GFAP for astrocytes, CD 45 for myeloid cells, and CD31 and GluT1 for ECs. **g**. Summary for **f**. N = 27 mice for CD31 and N = 3 mice for the rest of the markers.

a

b

c

383 **Extended Data Fig. 2 Demonstration of successful MCAO for modeling stroke. a**,

384 Laser speckle contrast imaging of cerebral blood flow before and after MCAO. **b**,

385 Quantification of CBF reductions from 10 mice that were subjected to MCAO. **c**,

386 Representative bright field image of an atrophy mouse brain at RP34D. Data presented

387 as mean ± SEM, **** P < 0.0001

a

b

Extended Data Fig. 3 General characterizations of EGFP⁺ non-EC cells. a, Confocal

images of brain slices from mice at RP34D, stained by EC markers (GluT1, Ve-Cadherin,

and ERG), and basal membrane component IB4. White arrows indicate EGFP⁺ non-EC

cells. **b**, An illustrative summary of anatomical locations of EGFP⁺ non-EC cells after

examining 19 mice that included all the 8 time-points as shown Fig. 1**b**. **c-e**. Quantification

of the total process lengths (**c**), the number of primary processes (**d**), and the diameter

soma (**e**) for both type I and type II EGFP⁺ non-EC cells. Data presented as mean ± SEM,

*** $P < 0.0001$, **** $P < 0.0001$

a *Cdh5CreERT2:Ai47*

EGFP
Bone mallow cells without RBCs

Hoechst

Merge

50 µm

b

Bone marrow

Blood

0.92

0.86

monocyte
neutrophil

monocyte
neutrophil

SSC-A

Ly6C

Ly6C

EGFP

CD11b

CD11b

c

RP:34Days

P2   1.5M   2.5M   3.5M

Intraventricular injection of *AAV-DIO-EGFP* virus   TAM   MCAO   Analysis

d

Cdh5CreERT2:Ai47

Cdh5CreERT2

Cdh5CreERT2 + Virus

SSC-A

EGFP

e

% of EGFP⁺ cells in CD45⁺ cells

Cdh5CreERT2:Ai47
Cdh5CreERT2
Cdh5CreERT2 +virus

f *Cdh5CreERT2; AAV-DIO-EGFP* (Contralateral side)

EGFP

CD31

EGPP
CD31

50 µm

g

% of EGFP⁺ Cells

CD31   VE-Cadherin   ERG   GLUT1

h *Cdh5CreERT2; AAV-DIO-EGFP* (Ischemic side)

50 µm

Extended Data Fig. 4 Exclusion of the possibility for Cdh5⁺ peripheral circulating cells to be the EGFP⁺ non-EC cells found in parenchyma after stroke. a, Wild-field fluorescent images of white blood cells, stained with Hoechst, from *Cdh5CreERT2:Ai47* mice. The white arrow indicates an EGFP⁺ white blood cell. **b**, FACS analysis of the EGFP⁺ white blood cell identity in the bone marrow and circulating blood by using neutrophil marker Ly6G and monocyte marker CD11b. Two independent experiments were performed, resulting in the same conclusion. **c**, Illustration of *AAV-DIO-EGFP* virus tracing strategy. **d**, FACS plots of EGFP+ cells in bone marrow cells from *Cdh5CreERT2:Ai47, Cdh5CreERT2 alone, and Cdh5CreERT2* with virus injection. **e**, Quantification of two independent experiments for **d**. **f**, EGFP expression in ECs after virus infection in the contralateral side of *Cdh5CreERT2* mouse brain. **g**, Quantification of EC markers in EGFP+ cells for f. **h**. EGFP⁺ cells in the ischemic region of mice that ECs were traced by virus labeling strategy. N = 7 brain slices from 1 mouse (trace period 1.6 months, RP34D) for **g**.

Extended Data Fig. 5 Radiation-depleted *Cdh5CreERT2;Ai47* mice exhibit EC cell fate change. **a**, FACS analysis of EGFP+ cells in three groups of *Cdh5CreERT2;Ai47* mice that receive corn oil, TAM, or TAM together with radiation. **b**, Quantification for **a**, N = 3 mice. **c**, Confocal images of vasculatures of the contralateral and ischemic side of the brain from *Cdh5CreERT2;Ai47* mouse that received radiation. **d**, Quantification for **c**. N = 2 mice. e-f, FACS analysis and quantification of EGFP+ non-EC cells in brains of mice that received the same treatment as **c**. N =. 2 mice (trace period 1.6 months, correspondingly RP34D). Data presented as mean ± SEM, * P < 0.05

Extended Data Fig. 6 EGFP⁺ non-EC cells transformed into pericytes, depending on TGFβ signaling cascade. a, EGFP⁺ non-EC cells were negative for a variety of cell markers such as neurons (NeuN and doublecortin), astrocytes (GFAP), CNS macrophage (Iba1), and SMC (α-SMA). **b**, A bipolar-shape EGFP⁺ non-EC cell traced by the virus in *Cdh5CreERT2:Ai14* mouse. **c-d**, EGFP⁺ non-EC cells traced by the virus were positive for pericyte markers (CD13 and PDGFβR). **e**. Brain sections from mice that were traced for 1.6 months meanwhile received a daily injection of vehicle DMSO or TGFβRI inhibitors since the day when MCAO was performed. **f**. Quantification of EGFP⁺ non-EC cell densities for **e**. N = 2 mice.

1.   Kuriakose, D. & Xiao, Z. Pathophysiology and Treatment of Stroke: Present Status and Future Perspectives. *Int J Mol Sci* **21** (2020).

2.   Martino, G. How the brain repairs itself: new therapeutic strategies in inflammatory and degenerative CNS disorders. *Lancet Neurol* **3**, 372-378 (2004).

3.   Zhao, L.R. & Willing, A. Enhancing endogenous capacity to repair a stroke-damaged brain: An evolving field for stroke research. *Prog Neurobiol* **163-164**, 5-26 (2018).

4.   Hall, C.N. *et al.* Capillary pericytes regulate cerebral blood flow in health and disease. *Nature* **508**, 55-60 (2014).

5.   Díaz-Flores, L., Gutiérrez, R., Varela, H., Rancel, N. & Valladares, F. Microvascular pericytes: a review of their morphological and functional characteristics. *Histol Histopathol* **6**, 269-286 (1991).

6.   Pallone, T.L. & Silldorff, E.P. Pericyte regulation of renal medullary blood flow. *Exp Nephrol* **9**, 165-170 (2001).

445   7.   Simon, C., Lickert, H., Götz, M. & Dimou, L. Sox10-iCreERT2 : a mouse line to

446   inducibly trace the neural crest and oligodendrocyte lineage. *Genesis* **50**, 506-515

447   (2012).

448   8.   Etchevers, H.C., Vincent, C., Le Douarin, N.M. & Couly, G.F. The cephalic neural

449   crest provides pericytes and smooth muscle cells to all blood vessels of the face

450   and forebrain. *Development* **128**, 1059-1068 (2001).

451   9.   Yu, Q.C., Song, W., Wang, D. & Zeng, Y.A. Identification of blood vascular

452   endothelial stem cells by the expression of protein C receptor. *Cell Res* **26**, 1079-

453   1098 (2016).

454   10.   Wang, Y. *et al.* Ephrin-B2 controls VEGF-induced angiogenesis and

455   lymphangiogenesis. *Nature* **465**, 483-486 (2010).

456   11.   Science, A.I.f.B. TECHNICAL WHITE PAPER: TRANSGENIC

457   CHARACTERIZATION. *Allen Brain Atlas* **9** (2015).

458   12.   Kilani, B. *et al.* Comparison of endothelial promoter efficiency and specificity in

459   mice reveals a subset of Pdgfb-positive hematopoietic cells. *J Thromb Haemost*

460   **17**, 827-840 (2019).

461   13.   Planas, A.M. Role of Immune Cells Migrating to the Ischemic Brain. *Stroke* **49**,

462   2261-2267 (2018).

463   14.   Fernandez-Klett, F. *et al.* Early loss of pericytes and perivascular stromal cell-

464   induced scar formation after stroke. *J Cereb Blood Flow Metab* **33**, 428-439 (2013).

465   15.   Piera-Velazquez, S. & Jimenez, S.A. Endothelial to Mesenchymal Transition: Role

466   in Physiology and in the Pathogenesis of Human Diseases. *Physiol Rev* **99**, 1281-

467   1324 (2019).

468   16.   Hulshoff, M.S., Xu, X., Krenning, G. & Zeisberg, E.M. Epigenetic Regulation of

469   Endothelial-to-Mesenchymal Transition in Chronic Heart Disease. *Arterioscler*

470   *Thromb Vasc Biol* **38**, 1986-1996 (2018).