Rhythmic sampling of multiple decision alternatives in the human brain

Marcus Siems[1]*, Yinan Cao[1], Maryam Tohidi-Moghaddam[1], Tobias H. Donner[1], & Konstantinos Tsetsos[1,2]

[1] Department of Neurophysiology and Pathophysiology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

[2] School of Psychological Science, University of Bristol, Bristol, United Kingdom

* Corresponding author

Contact Information

Correspondence: m.siems@uke.de & k.tsetsos@bristol.ac.uk

Highlights

- Humans navigate complex decisions by dynamically shifting their attention to sequentially sample different aspects of the decision problem.

- Using magnetoencephalography (MEG), we tracked the locus and strength of covert attention in humans performing a multialternative choice task.

- The strength of covert attention fluctuated rhythmically around 11 Hz, independent from oculomotor activity.

- At the peak of the 11 Hz attentional oscillation, the previously sampled alternative was resampled, while at the trough of the oscillation, sampling switched to a different alternative

Summary

Humans and other animals navigate real-world decisions by sequentially attending to (*sampling*) subsets of the available information. The neural mechanisms underlying the selective sampling of decision-relevant information remain unknown. Here we used magnetoencephalography (MEG) recordings and neural decoding to track the spontaneous dynamics of the locus and strength of covert attention as human participants performed a three-alternative perceptual choice task. The strength of covert attention fluctuated rhythmically around 11 Hz. At the peak of this oscillation, attention re-focused on the location of the previously sampled alternative, presumably promoting the in-depth evaluation of a single alternative. By contrast, at the trough, attention was oriented to a different alternative, that way facilitating the comparison between alternatives. Further, switching attention to a different alternative intrinsically reset the attentional rhythmicity. We propose rhythmic attentional sampling as a cognitive mechanism underpinning the flexible information processing necessary for key decision computations.

Introduction

Every day of our lives we make decisions based on varying amounts of external information. For example, deciding when to cross the street involves considering multiple external cues - i.e. the number of cars, the state of the traffic lights, the movement of other pedestrians, etc. - and translating these into actions guided by internal goals (e.g. the need to catch the bus). Unlike in simple two-alternative perceptual decision tasks where incoming sensory information can be directly integrated into motor plans[1–6], in many real-life decisions, the brain may not be able to parse and process all decision-relevant information in parallel[7,8]. Thus, subsets of the available information may need to be sampled serially as the decision unfolds[9–12], i.e. via attentional mechanisms.

With limited attentional resources[13–15] a trade-off during challenging decision computations emerges: The brain needs to process and evaluate single alternatives while also reallocate attention between alternatives to compare them[12,13,16]. Recent work on the neurophysiological basis of attention in simple detection tasks has provided a candidate mechanism that might help decision-makers navigate this trade-off: Even during periods of sustained focus on a single stimulus attention is not static but oscillates[17–23]. This oscillation is evident in behavioral reports[18,20,21,24–27], and neuronal activity in the theta[18,20,24,28], alpha[19,29,30] and gamma[22,24,25,30] frequency ranges. As a result, when attention is at an oscillatory peak, processing of a selected stimulus is enhanced[17,18,20,22,25,31–33]. By contrast, when at an oscillatory trough, attention is relatively disengaged and might be shifted from one stimulus to another[34]. Distinct attentional modes supporting different functions can be thus confined in one cycle of this oscillation[17]. Investigations in pupillometry and neuromodulatory systems further highlight the distinction between a focused versus a disengaged and exploratory attentional mode on longer timescales[35,36].

Here we hypothesized that the rhythmic alternation between focused and disengaged attentional modes may underlie the narrow and exploratory processing needed for valuation and comparison during multi-alternative decisions, respectively. Previous neurophysiological studies have established the role of rhythmic attention during simple detection tasks[37], tasks that could be solved through bottom-up strategies based on perceptual salience. However, it remains unknown whether equivalent rhythmic mechanisms are implicated during complex multi-alternative decisions that recruit top-down mechanisms and require protracted deliberation about alternatives.

Here we harnessed the high temporal resolution of MEG recordings to continuously track the locus and strength of covert spatial attention during protracted multialternative decisions. Focusing on covert attention enabled us to gauge rich sampling dynamics that cannot be perfectly captured by oculomotor activity alone[28,31,38–40], making them difficult to assess when only overt attention is tracked[10–12,16,41,42]. Using a multialternative choice task that requires valuing and comparing decision alternatives on intrinsically set timescales, enabled us to identify attention-driven serial processing of information while the decision unfolded. Importantly, we decoupled low-level stimulus features (contrast) from the decision value through randomized task-framing. We expected that the intrinsic dynamics of attentional allocation are rhythmic.

Indeed, in what follows we show that the strength of covert attention intrinsically oscillates around 11 Hz. Refocusing attention to the previously sampled alternative occurs at the peak of this oscillation while switching to a different alternative occurs at the trough and resets the ongoing rhythm. Thus, the conflict between gathering additional information about an alternative and exploring other alternatives is resolved through rhythmic sampling. We propose that rhythmic attentional sampling supports the flexible information processing needed to perform cognitive tasks with increased processing demands.

Results

We recorded non-invasive MEG from a total of 20 participants (9 female, age_{mean} = 28.05 years ± age_{STD} 4.6 years). The participants performed two interleaved tasks: A three-alternative perceptual choice task at threshold discriminability (main task) and a flicker-detection task (sensory localizer). Our experimental protocol for the main task followed a three-alternative forced choice design with fixed response windows (Fig. 1a). A trial started with a central fixation, followed by three static Gabor-patches of different contrast levels appeared at 2.2° eccentricity. To control for bottom-up sensory capture of attention induced by the presentation of the stimulus, we employed two framing conditions that were randomly assigned on each trial. In particular, in different framing conditions, participants were instructed to either choose the Gabor-patch with the highest or the lowest contrast (the highest value), with the framing instruction being revealed one second after the stimulus onset. Because the framing manipulation decoupled the low-level features (i.e., contrast) from the task-relevance of the choice alternatives, the task can probe goal-directed (top-down) attention independent from bottom-up sensory cortical responses. Participants performed the task with around 77% accuracy ($range_{min,max}$ = [66%, 86%]; Fig. 1b) and there was no accuracy difference between the two task framings ($|t_{19}|$ = 1.679; p = 0.11).

Fig. 1 | Experimental task & behavioral performance. a Trial structure of the three-alternative decision-making task with variable task framing. Stimulus size, contrast and eccentricity are indicative and have been changed for visualization purposes. Participants were asked to select the stimulus that had the highest (framing high: 'Hig') or the lowest (framing low: 'Low') contrast. **b** Choice probability across the three alternatives (H, M, L indicates high, medium, low decision value) for all trials and within each framing condition. The highest contrast stimulus has the highest value in high framing trials, while it has the lowest value in low framing trials. The dots indicate individual participant's choice probability.

Tracking the locus and strength of covert attention using MEG

We developed an approach to continuously track the locus and strength of covert attention during protracted decision processing. Specifically, we used the sensory localizer trials to train an inverted encoding model[43] on the angular positions of the stimuli, presented one at a time at different positions. This enabled us to estimate the focus of spatial attention during the decision-making task, when all stimulus locations were occupied (Fig. 2a and Supplementary S1). We followed the assumption that the neural representation, i.e. decodability, will be stronger for attended and weaker for unattended targets[13,14]. Inverting the model yielded a time-resolved estimate of the response to each of the three angular positions during the decision-making task (Fig. 2b). From the estimated responses we reconstructed spatial attention as the vector sum of the three channel responses (left, top, right). We focused on the resulting vector (in the following 'Attentional Vector') to precisely track internal attentional fluctuations. The vector length corresponded to the *strength* of attention. Thus, high attention strength might signal increased sensory processing, task engagement or arousal. The angular similarity to the three stimulus positions corresponded to the *locus* of attention. We

further tracked the eye-gaze position to ensure central fixation and disentangle covert attention from eye-movements.

Fig. 2 | Reconstructing the locus and strength of spatial attention with inverted encoding models. a Training the encoding model. The encoding model consists of a set of linear spatial filters defining the response given a certain stimulus. It comprises each information channel's tuning curve, which when discretized at presentation angles forms the Design Matrix with the normalized predicted response (from 0 to 1). In combination with MEG data from the sensory localizer (see Methods) the weighting of the information channels within each MEG sensor can be estimated by solving the general linear model (GLM). **b** Inverting the weights to estimate the channel responses during the decision-making task. We multiplied the inverted weighting matrix with the MEG data to estimate the response of the information channels at each time point (t). From the estimated responses we reconstructed the focus of spatial attention as the vector sum of the three spatial channel responses (left, top, right). The resulting Attentional Vector featured first, the length corresponding to the overall amount of attention and second, the angular similarity to the three stimulus targets.

Attention strength oscillates between 7 - 13 Hz

Building on recent work on the neurophysiological basis of attention in simple perceptual tasks[37], we hypothesized that attention strength (i.e., vector length) would vary throughout the trial as a result of distinct processing demands in the sensory- and decision-phases of the task. First, we quantified the trial-averaged attention strength (Fig. 3a, z-scored over stimulus presentation). After the stimulus onset, attention strength steeply ramps up and, relative to the framing cue onset, displays three peaks at

around -0.8 s, 0.2 s and between 0.7-1.2 s overlapping with key moments of the trial

(stimulus onset, framing and early decision formation). For external validation of this

metric we compared attention strength to the trial-averaged pupil diameter, a peripheral

proxy for arousal and task engagement[44]. The pupil diameter (Fig. 3b) resembled the

attentional vector distribution (Fig. 3a), albeit with more sluggish dynamics ($r_{pupil,veclen}$ =

0.54; p = 0.007).

Fig.3 | Attention strength fluctuates at around 11 Hz. a The distribution of the z-scored attention strength, i.e. vector length. The blue line indicates the average over trials and participants, the shaded area the standard error of the mean (SEM). Vertical dashed lines indicate the stimulus onset, framing cue onset and offset as well as the stimulus offset (from left to right). **b** The average normalized pupil diameter during task execution. The thick line and shaded area indicate the average and the SEM over participants. **c** Single trial example traces of the normalized attention strength. The depicted trials are from participant 1, session 1. **d** Windowed normalized amplitude spectra of attention strength fluctuations over the trial. Lines indicate the participant average, shaded areas display the SEM. The lines are color-coded according to the window onset (0.5 sec. window length) with regard to the framing cue, from early (blue) to late (pink) windows. **e-g** Frequency-specific change of attention strength amplitude relative to baseline. The frequency-wise amplitude strength values were normalized by dividing the average amplitude during the 0.5 seconds prior to stimulus onset and displayed in dB. **e** Time-frequency resolved (TFR) distribution of the amplitude of attention strength fluctuations. Red and blue colors indicate an increase or decrease relative to baseline, respectively. The values were statistically masked at p_{FDR}< 0.01 within each frequency. Vertical dashed lines as in **a**. **f** Average frequency-wise amplitude change (as in **e**) throughout the stimulus presentation period. Thick and thin lines display the average and single participant amplitude changes, respectively. Colored bars denote change from baseline (p_{FDR}< 0.05). **g** Attention strength fluctuation amplitude averaged for

frequencies between 7 - 13 Hz, relative to baseline in dB. Shaded area indicates the SEM. **h** Correlation of trial-averaged amplitude envelopes of the neuronal signal (compare Supplementary Fig. S2) and attention strength fluctuation. The correlation was computed for the broadband as well as frequency resolved signals. The values were statistically masked at p_{FDR}< 0.05 within each panel.

Importantly, there were substantial fluctuations of attention strength between trials (Fig. 3c), which we quantified using spectral analyses (Fig. 3d-g). Descriptively, attention strength fluctuated most strongly in the frequency range between 8 - 22 Hz (Fig. 3d), and significantly increased relative to the pre-stimulus baseline (-1.5 to -1 seconds prior to framing) between 4 - 16 Hz, and decreased between 25 - 50 Hz (Fig. 3e,f; $|t_{19}| > 2.38$, $p_{FDR} < 0.05$). The prolonged increase of the fluctuations between 7 – 13 Hz was most pronounced between 0.5 seconds prior to 1.5 seconds after the framing cue (Fig. 3g).

To examine the link between attentional and neural dynamics, we pairwise correlated the time-frequency resolved attention strength fluctuations with the frequency-equivalent neuronal signal amplitude envelope (Supplementary Fig. S2) over the entire stimulus presentation period. Following the spectral properties of attention strength (Fig. 3g) we focused our analysis on the alpha (7 – 13 Hz) and gamma (25 - 50 Hz) frequencies as well as on the broadband activity (0.2 – 200 Hz). We found strong positive correlations of attention strength with visual cortex activity in alpha, gamma, and broadband activity ($r > 0.25$; $p_{FDR} < 0.05$; Fig. 3h). Attention-neuronal correlations in the alpha-band highlighted lateral prefrontal cortical areas ($p_{FDR} < 0.05$).

The observed attentional fluctuations could be a consequence of stimulus-evoked cortical (and in particular visual) activity[45]. However, conversely to the 7 - 13 Hz increase in attention strength, we observed stimulus-evoked suppression of cortical activity in the alpha and neighboring frequency ranges throughout the entire trial duration (Supplementary Fig. S2). Similarly, we assessed the dynamics of the estimated

responses (Fig. 2b), the precursor to the attentional vector, and remapped each response to its option value class (High, Medium, Low). Here, the estimated responses displayed a broadband decrease between 8 – 45 Hz for all three value classes (Supplementary Fig. S3). Further, the difference in physical contrast between framing conditions appeared to not affect the estimated response within its value class apart from a ~0.25 second stimulus onset response (Supplementary Fig. S4). Thus, the increased attentional fluctuation around 11 Hz cannot be readily explained by stimulus induced cortical activity responses or low-level feature decodability dynamics (Supplementary Fig. S2,S3).

Attention strength fluctuations correlate with behavioral performance

As shown above, attention strength varies throughout the trial and displays prominent band-limited fluctuations but it remains to be shown whether these attention dynamics are relevant for decision-making performance. To evaluate the behavioral relevance of these dynamics we compared attentional fluctuations between correct and error trials. In a first analysis, broadband attention strength (Fig. 3a) was slightly larger (although not significantly) late in the decision phase for correct choices relative to incorrect ones (Fig. 4a; all $|t_{19}| < 1.5$, $p > 0.05$).

In a second analysis we repeated the comparison between correct and error trials by focusing on the frequency resolved attention strength fluctuations (compare Fig. 3e). We identified significantly increased fluctuations in attention strength associated with correct choices from 1.5 seconds post framing until the end of the trial, peaking in a frequency range from 8 – 16 Hz ($p_{FDR} < 0.05$). This indicates that band-limited attention strength fluctuations are predictive of choosing correctly (Fig. 4b). Importantly, we matched correct and error trials in difficulty and framing conditions (see Methods).

Fig. 4 | Attention strength fluctuations correlate with performance. a The distribution of the overall attention strength difference, i.e. vector length, between correct and error trials (condition matched, see Methods). The blue line and shaded area indicate the average over participants and the standard error of the mean (SEM). Vertical dashed lines denote the stimulus onset, framing cue onset and offset as well as the stimulus offset (from left to right). **b-c** Time-frequency resolved (TFR) attention strength fluctuation differences between correct and error trials as percent change to correct. **b** The distribution of TFR amplitude differences (correct-error) over the full spectrum and trial duration. The values were statistically masked at pFDR< 0.05 over the spectrum. **c** Frequency-wise amplitude change (as in **b**) averaged over the stimulus presentation period (-1 to 2.8 seconds). Thick and thin lines display the average and single participant amplitude changes, respectively.

Focusing and switching attention is nested within an 11 Hz rhythm

The above analyses highlight prominent and task-relevant modulations of attention strength around 7 - 13 Hz. However, these findings are unrelated to the locus of attention. Is there a relationship between attention strength and the dynamics of spatial attention? According to recent literature and the "rhythmic sampling" framework[17,19,22],

attention strength is expected to index the stability of the attention locus. In particular, it is expected that when attention strength peaks, the tendency to maintain the representation of one alternative is strong (locus stability). When attention is weaker, the tendency to switch to a different alternative increases (locus instability)[17,34,46], with reallocations occurring swiftly and analogously to covert 'attentional saccades'[21,31,47].

To test this hypothesis directly, we extracted brief events of maximal engagement with one decision alternative and assessed their temporal dynamics. This approach allows us to quantify several aspects of information sampling, such as identifying periods when participants are engaged with a given alternative, or quantifying how long it takes to switch attention between alternatives. Crucially, we can ask if these temporal regularities occur within the prominent 7 - 13 Hz attention strength rhythm.

Attentional events are defined as points in time when the Attentional Vector aligns with one of the three decision alternatives (top, left, right). Quantitatively, these are timepoints of maximum angular similarity (MAS) between the Attentional Vector and one of the decision alternatives (Fig. 5a). Hereby, MAS displays no systematic variations over the course of a trial. Next, we binarized MAS by applying a threshold (MAS 90% percentile, see Methods) to define brief events associated with selective attention to a single alternative (akin to an 'attentional saccade'). It is worth noting that the MAS metric is defined in a circular space and thus under completely random conditions (periods without visual stimulation) or when decodability of the alternatives is weak (inattentiveness) MAS will still exceed the angular threshold at random times. In these cases, we would falsely infer that attention is allocated to one of the three alternatives. We thus included a second threshold on the vector length, enabling us to exclude purely noise-driven events as much as possible (Fig. 5b and Supplementary S5). An attentional event is thus only recorded when both the attention locus is selective for one decision alternative and the attention strength is sufficiently high. Importantly, these events

appear to carry information about the decision behavior (Fig. 5c): During correct choices the high-value alternatives are sampled more often (logistic regression predicting choice from event counts within participants, weights t-test against zero: $t_{19,chooseH,stayH}$ = 2.35, p = 0.03; $t_{19,chooseH,switchH}$ = 2.42, p = 0.02). When choosing the second-highest value alternative the highest value target is sampled less ($t_{19,chooseM,stayH}$ = -2.52, p = 0.02; $t_{19,chooseM,switchH}$ = -2.71, p = 0.01). Additionally, the mid-value alternative was sampled less frequently when the low-value alternative was chosen ($t_{19,chooseL,stayM}$ = -3.83, p = 0.001; $t_{19,chooseL,switchM}$ = -3.91, p = 0.001). Further, we found pupillometric measures of neuromodulation to display a general link between arousal and focused information sampling (Supplementary Information & Supplementary Fig. S6).

Next, we characterized the dynamics of attentional events in a data-driven fashion. To assess their temporal structure, we timed the interval in-between successive attentional events (inter-event-intervals, IEI; Fig. 5d). Assuming that attentional allocation is governed by a random Poisson process (i.e., no temporal structure), the IEI histogram would display an exponential decay of probability density with interval duration (Fig. 5d,e, dashed lines). However, we identified clear deviations from this prediction, as the IEI histogram peaked at 0.13 seconds and at around 0.18 seconds (p_{FDR} < 0.05). These short peak latencies indicate that events tend to cluster together in a burst-like pattern over the course of the trial. These results confirm that the temporal structure of attention allocation is not random.

Fig. 5 | Attention allocation to distinct stimuli is nested within an 11 Hz rhythm. a Maximum angular similarity (MAS) averaged over trials. The blue line indicates the z-scored average over participants. The shaded area displays the SEM. Vertical dashed lines indicate the stimulus onset, framing cue onset and offset as well as the stimulus offset (from left to right). **b** Schematic definition of attentional events in an example trial. An attentional saccade (red mark) is recorded when vector orientation (90% MAS percentile) and attention strength are sufficiently high (90% vector length percentile). **c** Logistic regression predicting choice (H, M, L) from the event count per trial. Bars and vertical lines denote the mean and SEM over participants. Asterisks denote significant differences from zero (p < 0.05). **d** Inter-event-interval (IEI) probability histograms over all trials and attentional events. The black line indicates the average over participants, the shaded area the SEM. The black bars denote significant ($p_{FDR} < 0.05$) interval probabilities exceeding random conditions (dashed line). **e** Inter-event-interval histograms separately for stay (orange line, ± SEM) and switch (green line, ± SEM) attentional events. The bars denote significant ($p_{FDR} < 0.05$) interval probabilities exceeding random conditions. **f** Same data as in **e** plotted as t-score against random. Dashed lines indicate the uncorrected significance-levels and zero. **g** Box-plots of first-to-second peak latencies. Colored dots display single participants, color-coded as above. **h** First peak latency relative to the cycle length (see **f**). Asterisk denotes a significant difference between event types ($p_{FDR} < 10^{-4}$). N denotes the number of participants with significant peaks within the IEI histogram. **i** Schematic - Covert attention "spotlight" allocations nested within the underlying 11 Hz rhythm (gray) with two sets of attentional events offset by half a cycle: "staying" on the same stimulus and "switching" the focus of attention.

We next examined whether the dynamics of attentional allocation change as a function of the identity of the previously attended alternative. Attention can either re-focus on the previously attended alternative (staying) or shift to a different alternative (switching). As evident in the corresponding IEI histograms, staying and switching events exhibited distinct temporal profiles: staying predominantly occurred after 0.1 s and 0.18 s from the preceding event; and switching mostly after 0.05 s and 0.15 s (Fig. 5e,f and Supplementary S7), independent of the framing condition (Supplementary Fig. S8). Generally, we observed more stay than switch events (mean p_{Stay} = 0.63, $p_{Stay,25\%-75\%}$ = 0.60 – 0.64) and that switching was relatively increased right after stimulus onset and later in the decision phase (Supplementary Fig. S9).

The IEI histograms associated with stay and switch events can help assess the (potential) rhythmicity of these events: Assuming that events likely occur at distinct phases of an underlying rhythm, we can expect maxima of the IEI probability density reappearing in the first, second, third, etc. cycle of a putative oscillation at event-specific phases of each cycle. Thus, the latency between reappearing peaks in the IEI histogram (effectively the cycle length) would enable us to quantify rhythmicity. Both staying and switching events displayed approximately 0.09 s first-to-second peak latencies (median$_{stay}$ = 0.09 s, range$_{stay,25\%-75\%}$ = 0.088 s – 0.098 s; median$_{switch}$ = 0.088 s, range$_{switch,25\%-75\%}$ = 0.08 s – 0.095 s) – corresponding to rhythmicity at 11 Hz (Fig. 5g). Thus, the probability distribution of the events is consistent with the frequency of the dominant attention rhythm (Fig. 3) and the events appear to follow the attentional rhythm (Fig. 5i).

We further observed a prominent offset between switch and stay peak probabilities (Fig. 5e,f). While stay events displayed the first peak at around 0.085 s (median = 0.085 s, range$_{25\%-75\%}$ = 0.076 s – 0.094 s), switch events peaked earlier at around 0.045 s (median = 0.046 s, range$_{25\%-75\%}$ = 0.041 s – 0.054 s; paired $t_{18,stay-switch}$ =

12.35, p < 10^{-9}). With respect to each participant's event-probability cycle length (Fig. 5g) we identified the first 'switching' peak at around 0.5 (median = 0.51 cycle length, range$_{25\%-75\%}$ = 0.46 - 0.64) and the first 'staying' peak at a full cycle length (median = 0.95 cycle length, range$_{25\%-75\%}$ = 0.79 - 1.01) relative to the previous event, with a consistent switch-to-stay phase-shift slightly below half a cycle (median difference = 0.45 cycle length, range$_{25\%-75\%}$ = 0.31 - 0.48; Fig. 5h, grey box).

In sum, the temporal regularities characterizing attentional events provided a window into the dynamics of covert attentional sampling during decision formation. These results show that both event types rhythmically reappear over time, offset by half a cycle (Fig. 5h,i)[23] representing two distinct attentional modes: focus and reallocation.

Brief attentional saccades impact the ongoing attentional rhythm

The offset between switch and stay events introduces an insightful caveat. If attentional modes simply follow the attentional oscillation (hereafter, the "follow" interpretation), with stay events occurring at the peak and switch events occurring at the trough, then a switch (stay) event should follow half a cycle after a previous stay (switch) event or a full cycle after a switch (stay) event (Fig. 6a). This would result in IEI peaks at double the attentional strength oscillation frequency and no offset between switch- and stay events (Fig. 6a, lower panel). However, this runs against our reported findings (Fig. 5).

Further, the 'follow interpretation' does not hold when we assess second order dependencies in longer event sequences. We defined IEI histograms for four sequence types: Switch after stay, switch after switch, stay after stay, stay after switch. Under the "follow" interpretation, we would expect repetition (switch-after-switch, stay-after-stay) histograms to be offset by half a cycle relative to alternations (Fig. 6a). However, we found that the IEI histograms within each event type were invariant to the preceding

event (Fig. 6b,c). In other words, the temporal dynamics patterns expected under the "follow" interpretation do not hold.

Fig. 6 | Schematic and empirical second-order sequence inter-event-interval probabilities. a
Schematic of IEI probability depending on previous event identity. If the attention oscillation persists irrespective of events ("follow" interpretation), we would expect **1)** no offset between stay and switch peaks and **2)** peak event probabilities to fluctuate at double the attention modulation frequency (Fig. 3e vs. 5g). **b,c** Second-order sequence effects on inter-event-interval probabilities separately for **b** stay (orange line, ± SEM) and **c** switch (green line, ± SEM) attentional events. Solid and dashed colored lines indicate the mean interval probability following a stay or a switching event, respectively. The bars denote significant ($p_{FDR} <$ 0.05) interval probabilities differences between respective 2nd-order sequence types, color-coded as above. The empirical data is not in line with the follow interpretation (compare **a**).

Taken together, this finding challenges the concept that the attentional rhythm is unaltered in amplitude and frequency over time. Attentional events do not merely follow the attentional strength oscillation. In the next section we present evidence for a more nuanced "interaction" interpretation, in which attentional saccades can interfere with the rhythm, starting it anew.

Attentional switching resets rhythmic attention

To better understand how attentional events might alter the attentional oscillation we focused on attention strength around event onset. We averaged attention strength in a time-window from -0.2 s prior to 0.2 s after each event (Fig. 7a,b; MAS Supplementary Fig. S10). Prior to events, the traces of attention strength were relatively consistent over participants but largely unaligned between switch and stay (mean r_{pre} = 0.13, $r_{pre,25\%-75\%}$ = -0.05 - 0.34; t_{19} = 0.49, p = 0.31; correlations on detrended signals, Fig. 7b). By contrast, right after the event, the attention strength traces indicated a strong overlap between both event types (mean r_{post} = 0.60, $r_{post,25\%-75\%}$ = 0.54 – 0.76; t_{19} = 2.59, p = 0.009; correlations on detrended signals). Moreover, attention strength was weaker around switching- compared to stay events (Fig. 7c) both pre- (mean $ratio_{Switch/Stay}$ = 0.90, $range_{25\%-75\%}$ = 0.85 – 0.94; paired t_{19} = -8.09, p < 10^{-6}) and post-event (mean $ratio_{Switch/Stay}$ = 0.92, $range_{25\%-75\%}$ = 0.88 – 0.95; paired t_{19} = -7.40, p < 10^{-6}) but increased post-event by almost 2% (mean $\Delta ratio_{Switch/Stay}$ = 0.019, $range_{25\%-75\%}$ = 0.011 – 0.029; paired t_{19} = 7.21, p < 10^{-6}).

Fig. 7 | Peri-event fluctuations of attention strength. a Attentional event-triggered average of the z-scored attention strength, i.e. vector length. The colored lines and shaded area indicate the average and SEM over events and participants separately for stay (orange) and switch events (green). Pre- and post-event epochs are highlighted in blue and purple. Gray lines display full 11 Hz cycles from the event. **b** The same data as in **a** detrended individually pre- (-0.2 s to -0.05 s) and post-event (0.05 s to 0.2 s). **c** Peri-event attention strength difference between switch and stay, displayed as the ratio of average activity over a time-window from -0.15 to -0.05 (Pre, blue) and 0.05 to 0.15 (Post, purple). Colored dots and gray lines indicate the single-participant ratio and the difference from pre- to post-event, respectively. Asterisks indicate significant differences ($p < 10^{-6}$). **d** Dominant peri-event fluctuation frequencies based on **b**. *N* denotes the number of participants with two peaks per interval (see Methods). **e** Correlation between the attentional event probability (Supplementary Fig. S9) and the amplitude of the attentional fluctuations (Fig. 3e) over the stimulus presentation period. Lines depict the average, shaded areas the SEM over participants. Thick lines indicate significant differences ($p_{FDR} < 0.05$) and crosses a statistical trend for significance ($p_{FDR} < 0.1$). **f** Schematic - pre-event activity dissociation and the reset of attentional fluctuations at each event.

Focusing on pre-event attention strength, we observed a faster fluctuation for switch events (Fig. 7a,b). We thus quantified the dominant fluctuation frequencies pre- and post-event, and for both event types. Attention strength fluctuated at around 9-12 Hz before and after staying events, as well as after switching events (Fig. 7d). However, prior to switching, there was an increase (paired t_{16} = 5.82, p_{FDR} = 10^{-4}) of the fluctuation frequency to around 20-26 Hz, dissociating switch and stay events. Correlations between event distributions over the trial and band-limited neuronal activity further underline this qualitative dissociation between switch and stay events in an extended visual and lateral prefrontal cortical network (Supplementary Fig. S11). Overall, our findings suggest that switching events in particular are preceded by a higher frequency profile (Fig. 7d) and they 'reset' the ongoing attention oscillation to the dominant frequency around 11 Hz. Following both event types, the 11 Hz attentional rhythm starts from the local maximum (Fig. 7a,b) and attention strength peaks at around 0.09 s and 0.18 s post-event, i.e. approximately after one and two full cycles. Further, this reset can explain the lack of second-order sequence effects (Fig. 6).

Given the presence of a higher frequency attentional rhythm prior to switching, we next asked whether faster or slower attentional fluctuations predict the occurrence of event types[15,17]. We thus correlated the attention strength fluctuation amplitude in each frequency range (Fig. 3e) with the trial-averaged event probability (Supplementary Fig. S9). Stronger 9 - 11 Hz attention modulation trended towards more stay events (Fig. 7e, paired t_{19} > 1.92, p_{FDR} < 0.08) and stronger 19 - 40 Hz modulations significantly predicted more switch events (Fig. 7e, paired t_{19} < -3.42, p_{FDR} < 0.05). Hence, faster attention rhythms corresponded to an increased probability to re-allocate attention (switching).

Taken together, our results suggest that switching the attention locus exhibited a unique relation to attentional fluctuations (Fig. 7f), resetting the rhythm frequency, as

well as distinct spectral fingerprints particularly highlighting higher frequencies (> 11 Hz). Switch events appear to represent a neurophysiological mechanism of reallocating attention between options dissociable from the (re-)focus on single alternatives (stay). These two distinct modes of attention appear to be organized through a common oscillation: With attentional switches embedded in down-phases of the attention oscillation the conflict between prolonged focusing on one decision alternative and the need to shift attention between alternatives to compare them can be resolved temporally.

Dynamics of covert attention can be dissociated from (micro)saccadic eye movements

Previous research has shown that overt, i.e. eye-movements, and covert attention are linked. Particularly, it has been proposed that (micro)saccades are overt manifestations of covert attentional processing[48–50]. Similarly, the recent literature on the interplay between attention and decision-making has predominantly focused on overt attention via saccadic eye movements[12,16,42]. However, new findings indicate that overt and covert attention allocation is dissociable and may offer distinct insights into information sampling and intrinsic processing[28,31,38,39]. We thus assessed if and how the rhythmic fluctuations of covert attention relate to, or potentially originate from, the activity of the oculomotor system. In other words, do eye-movements show compatible dynamics as covert shifts of attention?

We applied eye-tracking and extracted (micro)saccadic eye-movements (MS), to evaluate the relationship of overt gaze to our measures of covert attention. The trial-averaged MS rate displayed pronounced variation with the trial structure (Fig. 8a), increasing after stimulus- and framing cue onset and following a ~0.2 s suppression period. To characterize the temporal structure of (micro)saccades, we computed the

inter-saccade-intervals (ISI) over the stimulus period. ISI histograms displayed a single

peak at around 0.25 s (Fig. 8b). The oculomotor first-peak latencies were thus

considerably longer in comparison to those of covert attention (Fig. 5e; $p < 10^{-10}$).

Moreover, due to the lack of additional local maxima, there is no evidence of persistent

rhythmicity in the (micro)saccade distribution beyond isolated serial saccades (i.e.

absence of first-to-second peak latencies).

Fig. 8 | Relationship between covert attentional fluctuations and oculomotor activity. a Average
(micro)saccade (MS) rate during task execution. The thick line indicates the average over participants and
the shaded area the SEM. Vertical dashed lines indicate the stimulus onset, framing cue onset and offset as
well as the stimulus offset. The horizontal line indicates the average MS rate over the stimulus on period
(1.14 Hz). **b** Inter-saccade-interval histograms for micro(saccades). Thick and thin lines display the group-
average and single-participant ISI histograms, respectively. **c** Oculomotor event, i.e. (micro)saccade, rates
pre- and post-attentional saccades (switch in green, stay orange) tested against shuffled events (t-test). The
thick colored bar denotes statistical significance ($p_{FDR} < 0.05$). **d** Attention strength around (micro)saccade
onset. The data is z-scored based on the 0.4 seconds prior to MS. The black line indicates the average over
participants and the shaded area the SEM. The thick bar denotes statistical significance from zero ($p_{FDR} <$
0.05). **e** The correlation of the attentional event probability (Supplementary Fig. S9) with the trial-averaged
(micro)saccade rate **a** for each participant. The bars and vertical lines display the average and SEM over
participants, respectively. The colored dots represent the single participant Pearson correlation coefficients
(color-coded as in **c**). Asterisk denotes significant correlations ($|r| > 0$, $p = 0.02$). **f** Proportion of attentional
events preceded by (micro)saccades. Color-coded as above. Bars and vertical black lines display the mean

and 25%-75% range. Asterisk denotes significant difference between switch and stay ($p < 0.001$). **g,h** Inter-event-interval histograms for stay **f** and switch **h** attentional events split into events with (dashed) and without a (micro)saccade (solid line) between successive events. Color-coded as above. The thick colored bar denotes statistical significance (No MS > post-MS; $p_{FDR} < 0.025$).

To directly assess the relationship between oculomotor activity and covert attentional events, we computed the (attentional) event-triggered MS rate. We identified a prominent increase around -0.15 s, exclusively before switching events (Fig. 8c; $p_{FDR} < 0.05$). Conversely, attention strength was strongly increased following a (micro)saccade at around 0.17 s post-saccade (Fig. 8d). Correlating the average MS rate with the corresponding attentional event probability within each participant (Fig. 8e), we found that attention switches are positively correlated with the MS rate ($r_{mean} = 0.06$; $t_{19} = 3.23$; $p = 0.005$). Further, we found that (micro)saccades preceded switching events (mean = 35%, range = 25 – 50%) more often than staying events (mean = 26%, range = 17 – 38%; $t_{19} = -13.2$, $p < 10^{-9}$). Taken together, oculomotor activity appears to play a prominent role in switching attention between decision alternatives but less when staying focused on the same alternative.

We next evaluated the effect of oculomotor activity on the ongoing attention oscillation. We examined if a (micro)saccade occurring between two attentional events affects the respective interval lengths. We found that on average, (micro)saccades decrease the probability for short-latency intervals up to 0.2 s both for staying and switching events (Fig. 8g,h). Thus, overt gaze displacements appear to delay the upcoming attentional event but the rhythmicity persisted (Fig. 8g,h): We identified post-saccadic rhythmicity (first-to-second peak latency) in 15 out of 20 participants for both event types with no detectable change in oscillation frequency relative to events without (micro)saccades (paired $t_{14,stay} < 1.8$, $p_{stay} = 0.09$; paired $t_{14,switch} < 1.1$, $p_{switch} = 0.29$).

In conclusion, oculomotor activity interacted with the way attention switched between decision alternatives. However, overt gaze cannot explain the dynamic aspects of attentional allocation. Thus, the analyses reported in this section indicate that the presented insights on covert attention dynamics cannot readily be inferred from monitoring eye position.

Discussion

Attention likely plays a pivotal role in complex, naturalistic decision-making, steering information sampling in order to assess and compare alternatives. However, little has been known about the role of internal attentional dynamics in decision formation. By means of high-precision, continuous tracking of the locus and strength of covert attention in the human brain, we here uncovered rich spontaneous attentional dynamics that govern information sampling during multialternative decisions. We established that the strength of attention fluctuated around 11 Hz frequency and that peaks of this attentional oscillation were associated with an increased tendency to maintain focus on a single decision alternative indicative of focal processing. By contrast, troughs of the attentional oscillation were associated with a tendency to switch attention between decision alternatives. Redirecting attention to another alternative was further preceded by faster attentional fluctuations, and resetting the ongoing attentional oscillation.

Recent literature has quantified generic temporal fluctuations in behavioral performance[20,22,25,32,46] and/or in cortical population activity[18,20,21,26,30], without explicating the information content of the latter. By contrast, we here constructed content-specific decoders from patterns of cortical population activity (trained on independent localizer data). This enabled us to directly track the strength and locus of attention during decision formation. Importantly, this approach enabled us to discover attentional dynamics in a data-driven fashion, without relying on prior assumptions about rhythmicity or temporal structure. The framing manipulation in our decision task further decoupled the (bottom-up) stimulus intensity from the (top-down) task-relevance of the choice alternatives. Further, apart from a ~0.25 second stimulus onset response we did not find evidence that the physical contrast was affecting the decoding between framing conditions (Supplementary Fig. S4). We can thus conclude that our findings, and particularly the

dominant attention oscillation frequency around 11 Hz, were not driven by bottom-up sensory cortical responses (Supplementary Fig. S2,S3). The exact frequency channel for rhythmic sampling, here 11 Hz, might further depend on both the cognitive domain and also the task and stimulus complexity[19,31,51–53]. However, further research is needed to explicate the influence of external (e.g. binary vs. ternary decisions) and internal factors (e.g. neuromodulation) on the sampling speed.

These findings provide a window into the computational and neural basis of decision-making by alluding to the mechanisms underlying key decision computations. In decision-theoretic models, central computations such as valuation and comparison are conceptualized in idealized ways and assumed to occur at distinct stages of the decision process[54]. Within our proposed framework, re-focusing attention on a previously attended alternative corresponds to a processing mode that could implement valuation. Similarly, switching attention between alternatives may underpin comparative processing. We show that although tendencies to re-focus (valuation) and switch (comparison) attention predominantly occur at different times (Supplementary Fig. S9)[23], effectively they are temporally coordinated within a single cycle of attentional oscillations. Oscillating attentional strength, a feature potentially established through cortico-thalamic connections[30,55,56] or center-surround properties of the visual cortex[28], resolves the competition between sampling information about each decision alternative and comparing between them[15,17,22]. Thus, rhythmicity (identified at the level of single participants) is the mechanism that allows key computations to be performed without competing for the same cognitive resources or substrates[15,17,23].

One central finding is that stronger attentional rhythmicity right before the response is associated with more accurate decisions (Fig. 4), potentially due to enhanced information sampling for brief periods of time through attentional gain[22]. At times of strong attentiveness participants were more engaged[36,44] with the task

(Supplementary Fig. S6) and more likely to increase processing of the attended decision alternative (Fig. 5). Consequently, decision formation processes at the neuronal level might weigh incoming evidence in a similarly time dependent manner[57]. Attention fluctuates rhythmically, thus the attention-modulated input driving evidence accumulation is likely rhythmic as well. Integrating covert attention as a time-varying variable, both in locus as well as strength, into existing models[11,12] of evidence accumulation can improve our understanding of dynamic decision making as well as the suboptimalities characterizing behavior in complex, multiattribute contexts[10,12,58].

A further prominent aspect of our findings is that, unlike re-focusing, switching events are accompanied by marked changes in neuronal processing. Specifically, the frequency of the attentional oscillation is higher prior to switching (Fig. 7), indicating dissociable spectral features for the two attentional modes. 'In-depth evaluation' (stay, ~11 Hz) and the 'comparison of alternatives' (switch, ~22 Hz) might for example be coordinated through cross-frequency interactions of distinct attentional oscillations[15,17,22]. With a switch event, the frequency of the attentional oscillation is reset to the dominant ~11 Hz, signaling a shift from a disengaged to a focused mode of processing. This reset of the attentional rhythm, speaks for a transient interaction between the two modus operandi of attention, i.e. switching and orienting vs. staying and alertness[13], and is further supported by interactions between the corresponding neural substrates of the two modes (Supplementary Fig. S6,S11). The reset might serve two purposes. First, it could correspond to an internal go cue, indicating readiness of the system to change[14] and briefly enhancing processing power[59,60]. Second, it might temporally (re)align processing at various stages of the functional hierarchy[20,22,61] or multiplexed information channels[62]. In the context of decision theory, an attentional reset and its accompanying neural dynamics could be the cornerstone neurocomputational mechanism underlying binary

comparisons[1,10,41,60,63]. However, further research is needed to better understand the functional role that switching resets have during decision-making.

Finally, our findings suggest that the fast dynamics of sampling information from various decision alternatives are distinct from oculomotor activity. In principle, oculomotor responses, i.e. (micro)saccades, can guide subsequent neuronal processing[13,49,60] by triggering shifts of attention locus. Indeed, we showed that after fixational eye movements and microsaccades, the probability to switch attention between alternatives increased (Fig. 8). Overt gaze displacements yielded a brief post-saccadic suppression of attentional event probability followed by strengthened visual representations of the focused alternative. However, not every covert shift of attention is accompanied by compatible eye-movements[28,31,38,39,47]. Particularly, oculomotor activity itself did not display rhythmicity, inter-saccadic intervals peaked later than attentional event intervals and (micro)saccades displayed a relationship to switching but not to re-focusing attentional events. Lastly, the rhythmicity within the attentional cycle was not affected by overt gaze. Although recent experimental and modeling studies have emphasized the causal role of overt fixations on choice[16,41] and intrinsic value representations[12,42], our results paint a more nuanced picture. We show that the relationship between covert information sampling and eye-movements might be one-way: microsaccades can elicit covert reorienting, but covert reorienting is not necessarily associated with microsaccades. Thus, where one is fixating and what is intrinsically processed aren't interchangeable units of attention and neuronal processing. Our study offers a methodological framework that dissociates the role that latent and overt attentional components play on decision formation.

In conclusion, our findings highlight rhythmic sampling as a general neurophysiological mechanism that resolves conflicts between neuronal processing stages in visual search, target detection, perception, working memory[64] and, as shown

here for the first time, multialternative decision-making. Focused and comparative sampling of information display distinct spectral profiles but are coordinated within a common oscillatory cycle of about 0.1 seconds, and operate distinctly from overt oculomotor markers. Similarly, decision-relevant quantities such as the value representations of decision alternatives, likely fluctuate over time. These dynamics challenge extant sequential sampling models of choice, which assume stable and parallel throughput of value information to downstream accumulator representations. These insights open new theoretical avenues, highlighting the role of rhythmic information processing in shaping key decision computations.

Acknowledgements

This research was supported by the EU Horizon 2020 Research and Innovation Program (ERC starting grant no. 802905, K.T.) and the BMBF (01GQ1907, T.H.D.)

Author Contributions

All authors contributed to the conceptualization and experimental protocol. M.S., Y.C. & M.T.M. collected the data. M.S. & Y.C. did the formal data analysis and M.S. visualized the results. M.S., T.H.D. & K.T. wrote the initial draft, all authors contributed to editing the manuscript.

Declaration of Interests

The authors declare no competing interests.

Data and Code Availability

Data and Code for reproducing the main results will be deposited on osf.io (under CC

4.0 license) and made publicly available upon publication.

Methods

Participants and experimental procedure

Twenty right-handed participants (9 female, age_{mean} = 28.05 years ± age_{STD} 4.6 years) participated in a three-alternative decision-making task with variable task framing. The experiment was approved by the ethics committee of the Hamburg Medical Association and conducted in accordance with the Declaration of Helsinki. All participants gave their written informed consent and received monetary compensation for their participation.

Decision-making task. During central fixation three Gabor patches of varying contrast levels and orientations appeared on screen. The stimuli were presented peripherally (2.2° eccentricity, 1.6° stimulus diameter) at the top and 120° left and right of it. The three stimuli were drawn without replacement from a set of five linearly spaced contrast levels (10 possible combinations). We derived the distance between neighboring contrast levels via a staircase procedure individually within each participant. The patch orientations were drawn from a set of three equally distant orientations with either all three being identical or different and pooled our analyses over orientation conditions. After 1 second into the stimulus presentation a central framing cue (0.3 sec.) indicated the target contrast, either the highest ("Hig") or lowest ("Low"). The high contrast stimulus has the highest value in high framing trials, while it has the lowest value in low framing trials. We randomized the task framing between trials. After the framing cue offset the central fixation cross re-appeared and initiated a fixed decision phase (2.5 sec.). The three Gabor stimuli were visible and unaltered all throughout the 3.8 seconds stimulus period. After the stimuli turned off the fixation cross rotated 45° and cued the response. Participants had 2 seconds to indicate their decision with a left thumb, right thumb or foot button press for the left, right or top alternative, respectively. We randomized the foot (left or right) between participants. To visually guide the participants

the entire stimulus array was displayed with a 6° clockwise or counter-clockwise tilt when participants were using their right or left foot, respectively. The foot pedal was mounted on Styrofoam. We adjusted the height and location of the pedal for each participant such that the position was comfortable and a response could be triggered with a small flexion of the porcellus fori. With the individual adjustments to the foot pedal placement, we tried to make the motions between effectors as compatible as possible. Median response times (RT) were slightly but not significantly slower for foot responses (RT_{foot} = 0.46 sec $\pm$ 0.18 sec; RT_{left} = 0.40 sec $\pm$ 0.18 sec; RT_{right} = 0.39 sec $\pm$ 0.16 sec, each mean $\pm$ standard deviation; paired sign test $p_{foot,left}$ = 0.06; $p_{foot,right}$ = 0.08).

Task framing, contrast levels, Gabor orientation and stimulus locations were fully randomized, yielding a total of 1080 unique trials. The trials were blocked into random sets of 120 trials and each participants concluded two full randomization runs (18 blocks).

Sensory localizer. Prior to each block of the decision-making task the participants concluded a sensory localizer block. Each block consisted of 90 consecutive 0.35-second presentations of single Gabor stimuli at full contrast with 0.5 seconds inter-trial intervals. Each stimulus was presented with the same size and eccentricity as during the decision-making task. We instructed the participants to ignore the peripheral stimuli and fixate at the center. The participants' task was to detect a flickering of the central fixation cross that could occur randomly at each stimulus, in total 9 times per block, and indicate the detection with a finger button press. We excluded trials with flicker presentation from further analyses. In each block we fully randomized the orientation and the position over the three locations. Each unique stimulus combination was repeated ten times paired once with a central fixation flickering.

Functional data acquisition and preprocessing

Data acquisition. During the experiment the participants were sitting upright in a whole-head magnetoencephalography scanner (MEG; 275 axial gradiometer sensors, CTF Systems Inc.) in a magnetically shielded room. The data was acquired at a sampling rate of 1200 Hz. Each participant's head position was tracked online using three head localization coils (nasion, left/right preauricular points). Additionally, we recorded a 2-channel bipolar electro-oculogram (horizontal and vertical EOG) and a single channel bipolar electro-cardiogram (ECG) to control for cardiovascular and ocular signal artifacts. Eye-movements and pupil diameter were tracked with an MEG-compatible Eyelink 1000 system (SR Research). We further collected T1-weighted structural magnetic resonance images (MRI; sagittal MP-RAGE) to reconstruct individualized high-resolution head models.

Preprocessing. We band-pass filtered the continuous MEG data segments between 0.2 and 200 Hz (4th order Butterworth filter) and down-sampled it to 400 Hz. Next, we filtered the line noise with a notch filter at 50 Hz and its first 6 harmonics (1 Hz stop-band filter width). We implemented a two-stage procedure for artifact rejection[65,66]: First, we inspected all data segments for technical- (i.e. SQUID-jumps, passing cars), muscle- and eye-blink artifacts and excluded corresponding time intervals from further analyses. Second, we filtered the data into two distinct frequency ranges, a low- from 0.2 to 30 Hz and a high-range from 30 to 200 Hz, with a 4th order Butterworth filter. We computed independent component analyses (ICA) on both frequency ranges separately. This approach takes advantage of the intrinsic properties of different artifact types. On the one hand, eye-blinks, eye-movements, passing cars and cardiovascular artifacts display predominantly low-frequency features. On the other hand, muscle activity shows the strongest effects in higher frequencies. We visually inspected the independent components, rejected artifact components based on their topology, power-spectra and time-courses[65] and subsequently recombined both frequency bands.

Spectral analysis and source reconstruction

Spectral analysis. We estimated time-frequency representations of the time-domain

MEG signals using Morlet's wavelets[66]. We set the wavelet bandwidth at 0.5 octaves

(f/σ_f = 5.83, kernel width covered $\pm$ σ_t; σ_f and σ_t corresponds to standard deviation in

frequency and time domain) and derived complex spectral estimates for frequencies

between 2 to 90.5 Hz in quarter octave steps (2^1 to $2^{6.5}$ Hz).

Source projection. Based on each participant's individual structural MRI we generated

single-shell boundary-element method models[67] (BEM) and computed the physical

forward model (leadfields) for 457 equally spaced cortical sources[68] (~1.2 cm distance,

at 0.7 cm depth below pial surface). Sensor locations, source and leadfield positions

were co-registered to the common MNI-space using the position of the three head-

localization coils. We applied dynamic imaging of coherent sources[69] (DICS) linear

beamforming to estimate source-level neuronal activity. DICS is a spatial filtering

approach that reconstructs activity at the source of interest with unit gain while at the

same time maximally suppressing signal contributions from other sources.

Encoding the focus of covert attention

To reconstruct the focus of covert attention during free information sampling of multiple

stimuli we implemented an inverted encoding model[43]. The modeled activity hereby

displays momentary changes of the visual representations of the presented stimuli (Fig.

2). The model assumes that the sensor-level MEG activity can be described as a

weighted sum of distinct abstract neuronal populations, i.e. information channels, with

individual tuning properties to the angular positions plus noise *N*.

$$MEG_{Loc} = WC_{Loc} + N$$

We trained the encoding model on the functional localizer task *MEG_{Loc}*. *C_{Loc}* was a

design matrix describing a set of basis functions, i.e. the tuning curves, for each

information channel (Fig. 2a) and *W* described the weight matrix specifying the

relationship between the information channels and the neuronal data. Each information

channel was modelled as a half-wave rectified sinusoidal filter *f* with a fifth power

exponent ($\alpha = 5$). We derived the basis functions *f* for the design matrix as

$$f = |cos^{\alpha}[0.5(\theta - \mu)\pi/180]|$$

Here, θ is a continuous variable indicating the angular position relative to central fixation,

and μ the preferred angle of each filter (θ and μ are in degrees, with 0° pointing upward

along the vertical midline and angular value increasing counter-clockwise e.g., 90°

pointing leftward). α was the power exponent set to 5. The target stimuli were set to μ =

0°, 120° and 240° for top, left and right, respectively. During the functional localizer task

only one stimulus out of the full set has been visible on screen. Hence, we trained the

model during the localizer, as both *MEG_{Loc}* and *C_{Loc}* were known variables. We estimated

the weight matrix *W* (sensors x information channels) as follows

$$w_k = MEG_{Loc}c_{Loc,k}^T(c_{Loc,k}c_{Loc,k}^T)^{-1}$$

Where *w_k* denotes the weights of the *kth* information channel $c_{Loc,k}$, i.e. a single tuning

curve. As neighboring MEG channels are not independent estimates of neuronal activity

due to volume conduction we further included multivariate noise normalization to

improve the pattern reliability. Thus, we transformed each *w_k* into the pre-whitened

weighting vector *g_k*

$$g_k = \frac{E_k^{-1}w_k}{w_k^T E_k^{-1}w_k}$$

Where *E* is the noise covariance matrix for channel *k* using the channels noise term as

$\varepsilon_k = MEG_{Loc} - w_k c_{Loc,k}$. *E* was regularized by shrinking it towards the diagonal matrix,

using the optimal shrinkage factor, i.e., minimizing the expected squared loss of the

resultant covariance estimator. With the optimized weight-matrix *G* we can estimate the latent visual representation of the stimuli during the multialternative decision-making task (Fig. 2b). Therefore, we invert the encoding model

$$C_{MDM} = G^T MEG_{MDM}$$

C~MDM~ comprises the estimated response as the representational activity distribution over angular positions for each time point during the task. The strength of the representation indexes the covert visual attention towards each target on screen.

Lastly, we reconstructed the momentary spatial focus as the vector sum over the three estimated stimulus activities. The vector orientations were set to 0^o, 120^o and 240^o for the top, left and right stimulus, respectively and the corresponding length at each orientation from the channel response *C~MDM~*. This reconstructed attentional vector visualizes two main characteristics of spatial attention, namely the overall attention paid at each time point and from which target information is sampled. We considered the attentional vector length as a proxy of overall attention and the maximum of the angular similarity to the displayed alternatives as the most likely target of covert spatial attention. We trained the model for different time points during the localizer task with respect to stimulus onset. To achieve the best decoding performance, we selected the optimal training epoch using cross-validation (leave-one-block-out, 18 blocks per participant) within the sensory localizer task. We found decoding strength to peak between 140 to 170 msec post stimulus onset (Supplementary Fig. S1) and averaged the MEG signal over this period to construct the decoding model for the application on the *MEG~MDM~* data. Important for the interpretation of the IEM output is whether or not a stimulus is on screen. In the pre-stimulus phase the Attentional Vector might vary due to for example random fluctuations of ongoing neuronal activity. On the other hand, directly after the stimulus offset in the post-stimulus period participants might have still been

remembering the options or display extended stimulus processing from memory. We thus centered our analyses on the stimulus presentation period.

Spectral analysis of attention strength

We computed time-frequency resolved estimates of the attentional vector length to assess the spectral dynamics of attention strength. We band-pass filtered (4th order Butterworth filter) the trialed vector length (f_{sample} = 400 Hz, from -1.5 seconds relative to stimulus onset to 2 seconds after stimulus offset) to center frequencies between 2.8 Hz to 90.5 Hz in quarter octave steps ($2^{1.5}$ to $2^{6.5}$). The pass-band covered the center frequency plus/minus half an octave, e.g. for $2^{1.5}$ Hz center frequency we filtered the data between 2^1 and 2^2 Hz. We estimated the frequency dependent attention strength as the absolute value of the complex Hilbert transform from the time-domain vector length signal. To exclude edge artifacts, we cut two cycles of the center frequency from the beginning and end of each trial. We visualized the data between -1.5 to +3 seconds relative to the framing cue onset. We applied the same procedure for the reconstructed stimulus activities at each location (Supplementary Fig. S2,S3).

For the analysis of attention strength fluctuations as a function of correct versus error trials we stratified the included trials. Hereby, we matched the numbers of trials included from each of the 10 possible contrast combinations per framing condition. For all participants we identified more correct choices than error choices in all stimulus combinations. We thus randomly subsampled one hundred times from all correct trials according to the stratification assumptions and averaged the spectral amplitude and the vector length over repetitions before comparing against error trials (Fig. 4).

Defining attentional events

We estimated the momentary focus of spatial attention via the angular similarity of the attentional vector to the three stimuli on screen. At each point in time the attentional vector can display the focus to be close to one or in between targets. The maximum of

the angular similarity (MAS) between the Attentional Vector and the three alternatives is an estimate of how focused a participant is. If MAS equals 1 then the participant is focused solely on one alternative at a given time point. The minimum MAS is 0.5 for the case when the Attentional Vector points exactly in between two alternatives. To investigate the dynamics of information sampling, i.e. when one alternative is selectively focused, we thresholded the maximum angular similarity at the participant-specific 90^{th} percentile over the entire dataset and binarized the time-courses accordingly. However, as the vector orients in circular space and even under completely random conditions will point towards one of the three stimuli at times, we added a second threshold on the vector length (90^{th} percentile within a trial). Thus, we recorded attentional events when both thresholds were crossed at the same time. In other words, we traced points in time when participants were both attentive and locked on one stimulus. We recorded the first time point of passing both thresholds as an event onset and excluded events occurring within 0.025 sec after the previous onset. We further tested the behavioral relevance of event occurrences and event targets on the reported choice using logistic regression. Within each participant we predicted the categorical choice (H, M, L individually) from the number of events at each decision alternative (stay-H, stay-M, stay-L, switch-H, switch-M, switch-L). We computed t-tests against zero over participants and report the related t-scores and p-values.

Event dynamics. To assess the dynamics of these attentional saccades we investigated the intervals between subsequent events. Here, we recorded the time from event *e* onset to the onset of event *e+1* within a trial and generated inter-event-interval (IEI) histograms over all events and trials within a participant. We excluded intervals reaching over trials. The histograms were computed over a median of 17,696 events per participant (range 12,377 to 28,909; Supplementary Fig. S5). We further generated Null-models for inference statistical testing using randomization. Hereby, we randomized the

trial each event was belonging to while keeping its position within the trial. In other words, when we detected for example 10 events at time point t scattered over 2160 trials, we randomly reassigned trial identities to these events (without replacement) and proceeded in the same way with time point $t+1$. We computed the IEI histogram after every randomization run. With this approach we can randomize event relationships without tempering with the task-dependent dynamics. We repeated the randomization step 1000 times for each participant.

Event sequences. We further tracked first order event sequences. An event can occur either at the same stimulus location as the one before or a different location, i.e. a participant either refocuses (stay) or reorients (switch) their attention. We generated IEI histograms and the randomization for both event types separately and statistically tested interval probabilities against the mean of the randomized Null-models (one-sided paired t-test). Hereby, we obtained t-scores for each interval from 0.025 to 0.5 seconds (at f_{sample} = 400 Hz) for both stay and switch.

Interval-probability fluctuations. If we assume that attention is rhythmic then attentional events would occur probabilistically at preferential phases within an underlying oscillation and we would expect to observe distinct peaks in the IEI histograms: The first peak would display the preferential phase within the first oscillatory cycle, the second peak the second cycle, etc. Thus, we can estimate the prevalent attentional fluctuation frequency as the inverse of the latency between peaks. Therefore, we subtracted the participant-specific expected IEI histogram (Null-model) from the observed histogram. All non-significant values were set to zero according to random-effect statistics based on randomization (one-sided paired t-test against zero, $p < 0.01$; see also above). We smoothed the resulting truncated histograms twice with a 0.01 sec boxcar kernel, padding the edges to the first and last histogram value. Finally, we extracted the local maxima of the truncated and smoothed histograms. Overall, we identified one significant

peak in 17 and 18 participants and at least two peaks in 16 and 17 participants for the stay- and switch-IEI histograms, respectively. For further analyses we defined the cycle length of the attention rhythm as the latency from the first to the second peak.

Threshold assessment. We further controlled for effects of the applied threshold through a systematic assessment for a wide range of threshold choices. In theory, if the thresholds are too liberal the analysis will include several random events converging the IEI histogram to an exponential decay. If the threshold is too high there won't be enough events to have sufficient statistical power for the full IEI distribution. We varied the applied thresholds between the 50% and 95% percentile in 5% steps as well as for the 97.5% and 99% percentile separately for the maximum angular similarity and the vector length (Supplementary Fig. S12). This approach yielded a total of 144 (12 x 12) threshold combinations. We replicated our main findings (individual peaks in the IEI histogram) for a variety of threshold choices between the 80% and 95% percentile (Supplementary Fig. S12a,b). Importantly, the timing of histogram peaks appeared to be stable over thresholds. Further, we quantified the deviation of the histogram from an exponential decay, i.e. randomly distributed events, through the variability of interval probabilities further referred to as modulation intensity. Hereby, we first log-transformed the interval probabilities and linearly detrended the resulting values. Next, we smoothed these distributions with a 0.025 second (10 samples) box-car kernel twice and defined local maxima. Finally, we computed the modulation intensity as the standard deviation of the detrended histogram between the first and second maximum (Supplementary Fig. S12c,d). We found the strongest median modulation intensity for the highest vector length thresholds at over 95% with variably applicable thresholds for the MAS. Further increasing both thresholds beyond the 95% percentile yielded inconclusive modulation in several participants due to a lack of identifiable events (Supplementary Fig. S12e,f). We thus conclude that our main findings are threshold invariant within a sensible range

between the 80% to 97.5% percentile with an optimal range approximately between the 90% and 95% percentile for both parameters. At the 90% percentile threshold for both measures, i.e. the applied threshold during the main text, thus granted both reasonably reliable modulation intensity and further a sufficient number of attentional events to increase statistical power for event-triggered analyses (Supplementary Fig. S12g,h). Finally, the optimal threshold parameters depend on the research question, the number and the length of the trials. It is thus pivotal to control these parameters for each new analysis. However, we must stress that optimizing the parameters does not evoke rhythmicity or temporal structure but can only reveal existing structure.

Peri-event fluctuations of attention

To compare the dynamics leading up to attentional events we investigated the event-triggered fluctuations of attention from 0.2 sec before to 0.2 sec after each event. Here, we averaged the attention strength, i.e. attention vector length, and the attention focus, i.e. maximum angular similarity, over all events within each participant and z-scored the traces according to the distribution between -0.2 to -0.03 sec (pre-event period) and 0.03 to 0.2 sec (post-event period) before averaging over participants. We excluded the peri-event period (± 0.03 sec.) from normalization as the event itself, by definition the highest values, would skew the distribution. For further visualization we de-trended and smoothed the event-triggered traces with a 0.025 sec boxcar kernel separately for the pre- and post-event period. Further, we used the de-trended and smoothed traces to define prominent peri-event fluctuation frequencies via the inverse of the average peak-to-peak latencies. We used the two peaks closest to the event both in the pre- and post-event interval to define peri-event fluctuation frequencies. We could only identify a frequency when a participant would display at least two peaks within an interval.

Pupillometry and oculomotor activity analysis

We tracked the eye-gaze and pupil size of the right eye at 1000 Hz sampling rate with an EyeLink 1000 Long Range Mount (SR Research, Osgoode, Canada). We calibrated and validated the eye-tracker at the start of each block. The system's spatial resolution was <0.01° and the average position accuracy between 15 – 30 min arc. We down-sampled the sampling to 400 Hz compatible to the MEG data. Due to hardware issues we had to exclude one subject from the analysis.

Defining pupil responses. We detected blinks according to the MEG preprocessing pipeline. Within each participant we normalized the average pupil size between the 5% and 95% percentile. We derived the change in pupil diameter as the temporal derivative of the pupil size by subtracting adjacent frames.

Microsaccade detection. We converted the SR Research .edf files to Matlab through FieldTrip[67]. We normalized the gaze position through the data from the calibration. We defined (micro)saccades as brief events of high gaze velocity. We quantified gaze velocity as the absolute value of temporal derivate over x-y coordinates and smoothed with a 0.02 second Gaussian kernel. A saccade was recorded when the velocity exceeded 6 times the trial-based median[70]. To avoid counting the same event multiple times we imposed a minimum delay of 0.025 s between gaze shifts. This saccade definition is agnostic to the length of the gaze displacement and can thus include both saccades and microsaccades (microsaccade$_{threshold}$: gaze displacement <1 deg visual angle[33]). We thus denote all oculomotor events as (micro)saccades. The (micro)saccade rate was generated as the saccade probability over trials at each sample multiplied by the sampling rate (400 Hz).

Relating oculomotor events and attentional saccades. We assessed the attentional event-triggered (micro)saccade rate in time-windows between -0.3 to 0.3 s from attention event onset and averaged over events separately for switch and stay. We compared the average (micro)saccade rate within these windows against trial-shuffled attention-event

definitions (see also above), i.e. the change in (micro)saccade rate. We further analyzed the (micro)saccade triggered attention strength, i.e. vector length, in a time-window between -0.4 to 0.8 s from (micro)saccade onset. Finally, we investigated the effect of (micro)saccades on the rhythmicity of attentional events by re-computing the inter-event-interval histograms separately for intervals containing or not containing a (micro)saccade.

Neural correlates of covert attentional saccades

We computed the neuronal signal strength, i.e. the amplitude envelope, over the entire cortex to assess the relationship to attentional events (see also Spectral analysis and source reconstruction). According to our results (Fig. 3e and 5g) we focused the analysis on the alpha (11 Hz), beta (22 Hz) and gamma (40 Hz) frequency range. With respect to the literature we further included the theta frequency range (4 Hz)[17,19,22]. Within each participant and source, we correlated the frequency-specific signal amplitude envelope with the event probability throughout the stimulus on period. For each event type as well as the difference between them we tested the correlation against zero over participants (two-sided t-test). We further computed the correlation between the neural signal envelopes and the corresponding band-limited attentional fluctuations.

Event-triggered neural signals. To assess the relationship between event-triggered attention- and neural modulations we averaged over signal amplitudes for switch and stay events separately in each cortical source for a time window of ± 0.3 sec relative to event onset. We z-scored the neuronal activity within each window prior to averaging over events and participants. For visualization purposes we grouped a subset of cortical sources into four regions-of-interest[17,19,22] approximating: early visual areas including V1, medial parietal cortex, medial prefrontal cortex (mPFC) and the dorsolateral prefrontal cortex (dlPFC) including the frontal eye field (FEF). The average ROI activity was statistically compared between event types using a two-sided paired t-test.

References

1. Gold, J. I. & Shadlen, M. N. The neural basis of decision making. *Annu. Rev. Neurosci.* **30**, 535–574 (2007).

2. Donner, T. H., Siegel, M., Fries, P. & Engel, A. K. Buildup of Choice-Predictive Activity in Human Motor Cortex during Perceptual Decision Making. *Current Biology* **19**, 1581–1585 (2009).

3. Siegel, M., Buschman, T. J. & Miller, E. K. Cortical information flow during flexible sensorimotor decisions. *Science* **348**, 1352–1355 (2015).

4. Pinto, L. *et al.* Task-dependent changes in the large-scale dynamics and necessity of cortical regions. *Neuron* **104**, 810–824 (2019).

5. Khilkevich, A. *et al.* Brain-wide dynamics linking sensation to action during decision-making. *Nature* 1–11 (2024).

6. Wilming, N., Murphy, P. R., Meyniel, F. & Donner, T. H. Large-scale dynamics of perceptual decision information across human cortex. *Nature communications* **11**, 5109 (2020).

7. Kang, Y. H. *et al.* Multiple decisions about one object involve parallel sensory acquisition but time-multiplexed evidence incorporation. *Elife* **10**, e63721 (2021).

8. Busemeyer, J. R., Gluth, S., Rieskamp, J. & Turner, B. M. Cognitive and neural bases of multi-attribute, multi-alternative, value-based decisions. *Trends in cognitive sciences* **23**, 251–263 (2019).

9. Russo, J. E. & Dosher, B. A. Strategies for multiattribute binary choice. *Journal of Experimental Psychology: Learning, Memory, and Cognition* **9**, 676 (1983).

10. Noguchi, T. & Stewart, N. In the attraction, compromise, and similarity effects, alternatives are repeatedly compared in pairs on single dimensions. *Cognition* **132**, 44–56 (2014).

11. Noguchi, T. & Stewart, N. Multialternative decision by sampling: A model of decision making constrained by process data. *Psychological review* **125**, 512 (2018).

12. Krajbich, I. Accounting for attention in sequential sampling models of decision making. *Current Opinion in Psychology* **29**, 6–11 (2019).

13. Posner, M. I. Orienting of attention: Then and now. *Q J Exp Psychol (Hove)* **69**, 1864–1875 (2016).

14. Buschman, T. J. & Kastner, S. From behavior to neural dynamics: an integrated theory of attention. *Neuron* **88**, 127–144 (2015).

15. Fiebelkorn, I. C. & Kastner, S. Functional specialization in the attention network. *Annual review of psychology* **71**, 221–249 (2020).

16. Orquin, J. L. & Mueller Loose, S. Attention and choice: A review on eye movements in decision making. *Acta Psychologica* **144**, 190–206 (2013).

17. Fiebelkorn, I. C. & Kastner, S. A Rhythmic Theory of Attention. *Trends in Cognitive Sciences* **23**, 87–101 (2019).

18. Fiebelkorn, I. C., Saalmann, Y. B. & Kastner, S. Rhythmic Sampling within and between Objects despite Sustained Attention at a Cued Location. *Current Biology* **23**, 2553–2558 (2013).

19. VanRullen, R. Perceptual Cycles. *Trends in Cognitive Sciences* **20**, 723–735 (2016).

20. Landau, A. N. & Fries, P. Attention samples stimuli rhythmically. *Curr. Biol.* **22**, 1000–1004 (2012).

21. VanRullen, R., Carlson, T. & Cavanagh, P. The blinking spotlight of attention. *Proceedings of the National Academy of Sciences* **104**, 19204–19209 (2007).

22. Fries, P. Rhythmic attentional scanning. *Neuron* **111**, 954–970 (2023).

23. Rollenhagen, J. E. & Olson, C. R. Low-frequency oscillations arising from competitive interactions between visual stimuli in macaque inferotemporal cortex. *Journal of neurophysiology* **94**, 3368–3387 (2005).

24. Senoussi, M., Moreland, J. C., Busch, N. A. & Dugué, L. Attention explores space periodically at the theta frequency. *Journal of Vision* **19**, 22–22 (2019).

25. Helfrich, R. F. *et al.* Neural Mechanisms of Sustained Attention Are Rhythmic. *Neuron* **99**, 854-865.e5 (2018).

26. Fiebelkorn, I. C., Pinsk, M. A. & Kastner, S. A Dynamic Interplay within the Frontoparietal Network Underlies Rhythmic Spatial Attention. *Neuron* **99**, 842-853.e8 (2018).

27. Mo, C. *et al.* Competing rhythmic neural representations of orientations during concurrent attention to multiple orientation features. *Nature Communications* **10**, 5264 (2019).

28. Kienitz, R. *et al.* Theta Rhythmic Neuronal Activity and Reaction Times Arising from Cortical Receptive Field Interactions during Distributed Attention. *Current Biology* **28**, 2377-2387.e5 (2018).

29. Busch, N. A. & VanRullen, R. Spontaneous EEG oscillations reveal periodic sampling of visual attention. *Proceedings of the National Academy of Sciences* **107**, 16048–16053 (2010).

30. Fiebelkorn, I. C., Pinsk, M. A. & Kastner, S. The mediodorsal pulvinar coordinates the macaque fronto-parietal network during rhythmic spatial attention. *Nature communications* **10**, 1–15 (2019).

31. Gaillard, C. *et al.* Prefrontal attentional saccades explore space rhythmically. *Nat Commun* **11**, 925 (2020).

32. Dugué, L., Roberts, M. & Carrasco, M. Attention reorients periodically. *Current Biology* **26**, 1595–1601 (2016).

33. Bellet, J., Chen, C.-Y. & Hafed, Z. M. Sequential hemifield gating of α-and β-behavioral performance oscillations after microsaccades. *Journal of neurophysiology* **118**, 2789–2805 (2017).

34. Hogendoorn, H. Voluntary saccadic eye movements ride the attentional rhythm. *Journal of cognitive neuroscience* **28**, 1625–1635 (2016).

35. Jepma, M. & Nieuwenhuis, S. Pupil Diameter Predicts Changes in the Exploration–Exploitation Trade-off: Evidence for the Adaptive Gain Theory. *Journal of Cognitive Neuroscience* **23**, 1587–1596 (2011).

36. Dahl, M. J., Mather, M. & Werkle-Bergner, M. Noradrenergic modulation of rhythmic neural activity shapes selective attention. *Trends in cognitive sciences* **26**, 38–52 (2022).

37. Kienitz, R., Schmid, M. C. & Dugué, L. Rhythmic sampling revisited: experimental paradigms and neural mechanisms. *European Journal of Neuroscience* **55**, 3010–3024 (2022).

38. Liu, B., Nobre, A. C. & van Ede, F. Functional but not obligatory link between microsaccades and neural modulation by covert spatial attention. *Nature Communications* **13**, 3503 (2022).

39. Liu, B., Alexopoulou, Z.-S. & van Ede, F. Attentional shifts bias microsaccade direction but do not cause new microsaccades. *Communications Psychology* **2**, 97 (2024).

40. Krauzlis, R. J., Lovejoy, L. P. & Zénon, A. Superior Colliculus and Visual Spatial Attention. *Annu. Rev. Neurosci.* **36**, 165–182 (2013).

41. Krajbich, I., Armel, C. & Rangel, A. Visual fixations and the computation and comparison of value in simple choice. *Nature neuroscience* **13**, 1292–1298 (2010).

42. Mormann, M. & Russo, J. E. Does Attention Increase the Value of Choice Alternatives? *Trends in Cognitive Sciences* **25**, 305–315 (2021).

43. Brouwer, G. J. & Heeger, D. J. Decoding and reconstructing color from responses in human visual cortex. *Journal of Neuroscience* **29**, 13992–14003 (2009).

44. Pfeffer, T. *et al.* Coupling of pupil-and neuronal population dynamics reveals diverse influences of arousal on cortical processing. *elife* **11**, e71890 (2022).

45. Siegel, M., Engel, A. K. & Donner, T. H. Cortical Network Dynamics of Perceptual Decision-Making in the Human Brain. *Front Hum Neurosci* **5**, (2011).

46. Raposo, I. *et al.* Periodic attention deficits after frontoparietal lesions provide causal evidence for rhythmic attentional sampling. *Current Biology* (2023).

47. Buschman, T. J. & Miller, E. K. Serial, covert shifts of attention during visual search are reflected by the frontal eye fields and correlated with population oscillations. *Neuron* **63**, 386–396 (2009).

48. Hafed, Z. M. & Clark, J. J. Microsaccades as an overt measure of covert attention shifts. *Vision research* **42**, 2533–2545 (2002).

49. Moore, T. & Armstrong, K. M. Selective gating of visual signals by microstimulation of frontal cortex. *Nature* **421**, 370–373 (2003).

50. Laubrock, J., Kliegl, R., Rolfs, M. & Engbert, R. When do microsaccades follow spatial attention? *Attention, Perception, & Psychophysics* **72**, 683–694 (2010).

51. Caplette, L., Jerbi, K. & Gosselin, F. Rhythmic information sampling in the brain during visual recognition. *Journal of Neuroscience* **43**, 4487–4497 (2023).

52. Michel, R., Dugué, L. & Busch, N. A. Distinct contributions of alpha and theta rhythms to perceptual and attentional sampling. *European Journal of Neuroscience* **55**, 3025–3039 (2022).

53. Merholz, G., Grabot, L., VanRullen, R. & Dugué, L. Periodic attention operates faster during more complex visual search. *Scientific Reports* **12**, 6688 (2022).

54. Platt, M. L. & Plassmann, H. Multistage valuation signals and common neural currencies. *Neuroeconomics* 237–258 (2014).

55. Saalmann, Y. B., Pinsk, M. A., Wang, L., Li, X. & Kastner, S. The Pulvinar Regulates Information Transmission Between Cortical Areas Based on Attention Demands. *Science* **337**, 753–756 (2012).

56. Griffiths, B. J. *et al.* Rhythmic interactions between the mediodorsal thalamus and prefrontal cortex precede human visual perception. *Nature Communications* **13**, 3736 (2022).

57. Wyart, V., De Gardelle, V., Scholl, J. & Summerfield, C. Rhythmic fluctuations in evidence accumulation during decision making in the human brain. *Neuron* **76**, 847–858 (2012).

58. Tsetsos, K. *et al.* Economic irrationality is optimal during noisy decision making. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 3102–3107 (2016).

59. Peylo, C., Hilla, Y. & Sauseng, P. Cause or consequence? Alpha oscillations in visuospatial attention. *Trends in Neurosciences* **44**, 705–713 (2021).

60. Groh, J. M., Schmehl, M. N., Caruso, V. C. & Tokdar, S. T. Signal switching may enhance processing power of the brain. *Trends in Cognitive Sciences* (2024).

61. Stine, G. M., Trautmann, E. M., Jeurissen, D. & Shadlen, M. N. A neural mechanism for terminating decisions. *Neuron* (2023).

62. Boroujeni, K. B. & Womelsdorf, T. Routing states transition during oscillatory bursts and attentional selection. *Neuron* **111**, 2929–2944 (2023).

63. Hunt, L. T. *et al.* Triple dissociation of attention and decision computations across prefrontal cortex. *Nature neuroscience* **21**, 1471–1481 (2018).

64. Abdalaziz, M., Redding, Z. V. & Fiebelkorn, I. C. Rhythmic temporal coordination of neural activity prevents representational conflict during working memory. *Current Biology* **33**, 1855–1863 (2023).

65. Hipp, J. F. & Siegel, M. Dissociating neuronal gamma-band activity from cranial and ocular muscle activity in EEG. *Front Hum Neurosci* **7**, 338 (2013).

66. Siems, M., Pape, A.-A., Hipp, J. F. & Siegel, M. Measuring the cortical correlation structure of spontaneous oscillatory activity with EEG and MEG. *Neuroimage* **129**, 345–355 (2016).

67. Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational intelligence and neuroscience* **2011**, 156869 (2011).

68. Hipp, J. F., Hawellek, D. J., Corbetta, M., Siegel, M. & Engel, A. K. Large-scale cortical correlation structure of spontaneous oscillatory activity. *Nat. Neurosci.* **15**, 884–890 (2012).

69. Gross, J. *et al.* Dynamic imaging of coherent sources: Studying neural interactions in the human brain. *Proc. Natl. Acad. Sci. U.S.A.* **98**, 694–699 (2001).

70. Engbert, R. & Kliegl, R. Microsaccades uncover the orientation of covert attention. *Vision research* **43**, 1035–1045 (2003).