1
2
3 **Hemispheric divergence of interoceptive processing across psychiatric disorders**
4
5
6

7 Emily M. Adamic[1,2], Adam R. Teed[1], Jason A. Avery[3], Feliberto de la Cruz[4], Sahib S. Khalsa[1,5]
8
9
[1]Laureate Institute for Brain Research, Tulsa, OK, USA, 74136
[2]Department of Biological Sciences, University of Tulsa, Tulsa, OK, USA, 74104
[3]Laboratory of Brain and Cognition, National Institute of Mental Health, Bethesda, MD, USA,
20814
[4]Laboratory for Autonomic Neuroscience, Imaging, and Cognition (LANIC), Department of
Psychosomatic Medicine and Psychotherapy, Jena University Hospital, Jena, Thuringia,
Germany, 07743
[5]Oxley College of Health Sciences, University of Tulsa, Tulsa, OK, USA, 74119

30 **Correspondence:**
31 Sahib Khalsa MD, PhD
32 skhalsa@laureateinstitute.org

38 **Abbreviated title:** Hemispheric divergence of interoception
39
40
41
42 **Version date:** May 6, 2024

Abstract

Interactions between top-down attention and bottom-up visceral inputs are assumed to produce conscious perceptions of interoceptive states, and while each process has been independently associated with aberrant interoceptive symptomatology in psychiatric disorders, the neural substrates of this interface are unknown. We conducted a preregistered functional neuroimaging study of 46 individuals with anxiety, depression, and/or eating disorders (ADE) and 46 propensity-matched healthy comparisons (HC), comparing their neural activity across two interoceptive tasks differentially recruiting top-down or bottom-up processing within the same scan session. During an interoceptive attention task, top-down attention was voluntarily directed towards cardiorespiratory or visual signals, whereas during an interoceptive perturbation task, intravenous infusions of isoproterenol (a peripherally-acting beta-adrenergic receptor agonist) were administered in a double-blinded and placebo-controlled fashion to drive bottom-up cardiorespiratory sensations. Across both tasks, neural activation converged upon the insular cortex, localizing within the granular and ventral dysgranular subregions bilaterally. However, contrasting hemispheric differences emerged, with the ADE group exhibiting (relative to HCs) an asymmetric pattern of overlap in the left insula, with increased or decreased proportions of co-activated voxels within the left or right dysgranular insula, respectively. The ADE group also showed less agranular anterior insula activation during periods of bodily uncertainty (i.e., when anticipating possible isoproterenol-induced changes that never arrived). Finally, post-task changes in insula functional connectivity were associated with anxiety and depression severity. These findings confirm the dysgranular mid-insula as a key cortical interface where attention and prediction meet real-time bodily inputs, especially during heightened awareness of interoceptive states. Further, the dysgranular mid-insula may indeed be a "locus of disruption" for psychiatric disorders.

Introduction

Maintaining the body in a physiologically optimal state is a primary function of the
mammalian nervous system. The neurovisceral architecture accomplishing such feats depends on
the hierarchical central processing of peripheral internal body signals and operates hierarchically
and across heterogenous time scales (Smith et al. 2017), automatically returning the internal state
to a setpoint (i.e., homeostasis) (Cannon 1939) or shifting the internal state away from a setpoint in
anticipation of future deviations (i.e. allostasis) (Sterling 2012). Importantly, these regulatory
operations may depend on the unconscious as well as conscious monitoring of interoceptive states.
Interoceptive sensations may even dominate awareness during significant physiological
disturbances, motivating behaviors that restore bodily states to homeostatic equilibrium (Khalsa
and Lapidus 2016) and forming the basis for emotional experience and numerous cognitive
processes (Adolfi et al. 2017; Carvalho and Damasio 2021; Critchley and Garfinkel 2017; Uddin et
al. 2014). In turn, dysfunctional interoception has been implicated in the pathophysiology of
psychiatric (Bonaz et al. 2021; Khalsa et al. 2018) and neurological disorders (Birba et al. 2022;
Drane et al. 2020), underscoring the importance of identifying the form and function of
interoceptive neurocircuitry in humans, particularly in relation to conscious experience of the inner
body.
Conscious interoceptive perceptions must, to a considerable extent, reflect the organism's
current physiological state. This is hierarchically conveyed to the brain by cascades of peripheral
interoceptive inputs, from vagal and spinal afferent nerves relaying signals transduced from
interoceptors located throughout the major internal organ systems (e.g., cardiovascular,
respiratory, gastrointestinal, urogenital), as well as via direct chemohumoral signaling (Berntson
and Khalsa 2021). Ascending interoceptive pathways converge via brainstem and thalamic nuclei
onto the insular cortex, a structurally and functionally diverse region suggested to be the
integrative hub for interoceptive awareness, emotional feelings, and overall body awareness (Craig
2002, 2009) (for discussions of additional pathways and cortical targets see (Berntson and Khalsa
2021; Khalsa et al. 2009; Mayeli et al. 2023). A principal role for the insular cortex in interoceptive
awareness is supported by previous functional neuroimaging studies from our group, which have
highlighted the insula as the primary region dynamically tracking pharmacological perturbations of
the heart and lungs in both healthy individuals (Hassanpour et al. 2016, 2018), and those with
generalized anxiety disorder (Teed et al. 2022) (for fMRI meta-analyses of cardiac and
gastrointestinal interoception findings see Schulz (2016) and Halani et al. (2020), respectively). Our
studies have identified multiple subregions of the insula in this "bottom-up" processing of afferently
relayed cardiorespiratory signals, including the dysgranular mid and granular posterior sectors, in
general agreement with existing theoretical (Barrett and Simmons 2015; Fermin, Friston, and
Yamawaki 2022) and cytoarchitectonic (Evrard 2019) arguments regarding their function.
While cortical sensitivity to ascending signals is presumptively required for conscious
detection of internal physiological shifts in humans, accumulating evidence also suggests that
subjective perceptions (in any sensory modality) are not simply veridical reflections of these
ascending inputs, faithfully transmitted and reproduced in a dormant sensory cortex (Clark 2013;
Corbetta, Patel, and Shulman 2008; Mechelli et al. 2004; Pace-Schott et al. 2019). Instead, intrinsic
"top-down" processes also appear to modulate the activity of sensory cortices (including the
insula), potentially altering the resulting percept in the process. For example, internal models of the

body that anticipate the sensory consequences of descending regulation can adjust the precision-weighting of prediction error signals in a context-specific manner, increasing or decreasing the sensory processing of interoceptive inputs (Clark 2016; Petzschner et al. 2021; Pezzulo, Barca, and Friston 2015; Seth and Friston 2016). Extending beyond these model-based influences are goal-directed activities (also described previously as the 'attentional spotlight' effect (Brefczynski and DeYoe 1999)), whereby focusing voluntary attention towards certain environmental signals not only alters their conscious experience but selectively enhances neural activity in the responsive area of cortex. With respect to interoception, such goal-directed shifts in attention towards endogenous visceral signals have consistently identified activation of the mid insular cortex in healthy individuals (Avery et al. 2017; Simmons et al. 2013), and revealed abnormal mid insula activity associated with symptoms in those with eating disorders (Kerr et al. 2016), depression (Avery et al. 2014; Burrows et al. 2022), and substance use disorders (Stewart et al. 2020).

Conscious perceptions of interoceptive signals are thus underpinned by reciprocal interactions between bottom-up, ascending interoceptive afferents and the coupled, top-down descending efferent systems that enact adaptive regulation and attentional control (Berntson and Khalsa 2021). Based on the available evidence, the insula is uniquely positioned to serve as a cortical interface between these two information streams for the processing of interoceptive signals. The highly interconnected nature of the mammalian insula with other cortical (prefrontal (Mufson and Mesulam 1982), somatosensory (Augustine 1996), cingulate (Mufson and Mesulam 1982) cortices), subcortical (amygdala, thalamus (Gehrlach et al. 2020)), and brainstem (Gogolla 2017) areas facilitates the ability for salient bottom-up activity to recruit conscious attention mechanisms and/or give rise to perceptions and feelings that can adaptively drive behavior and promote homeostatic regulation (Evrard 2019). Furthermore, disrupted processing in this region may be characteristic of individuals with certain psychiatric disorders. In a previous human functional magnetic resonance imaging (fMRI) meta-analysis, abnormal activation of the left dorsal mid-insula differentiated healthy individuals from groups of individuals with anorexia nervosa, anxiety disorders, depression, bipolar disorder, or schizophrenia, across tasks involving the bottom-up modulation of pain, dyspnea, or hunger or the top-down manipulation of interoceptive attention (Nord, Lawson, and Dalgleish 2021). This insular "locus of disruption" was inferred as a possible transdiagnostic and domain-general neuromarker of psychopathology, and the finding advanced the field by pointing toward the mid-insula as an important area for interoceptive dysfunction and a putative target for clinical intervention (Paulus and Khalsa 2021). Despite the theoretical and empirical convergence related to bottom-up and top-down processing in the insula , no study to date has directly investigated both interoceptive processing streams together in the same sample.

Here, we conducted a pre-registered conjunction analysis of two established neuroimaging tasks preferentially manipulating the assessment of top-down and bottom-up interoception (Adamic et al. 2021). Our hypotheses, based on a voxelwise whole-brain approach, were threefold: 1) broad regions of the dysgranular mid insula would be co-activated when a) engaging top-down interoceptive attention under physiological resting conditions and b) during bottom-up perturbation of peripheral interoceptive signals via the adrenergic drug isoproterenol, 2) the agranular anterior insula would be preferentially selective to manipulations of interoceptive uncertainty, such as those occurring during saline infusions, and 3) relative to healthy individuals,

interoceptive processing in the insula would show lateralized differences across a transdiagnostic
group of individuals with anxiety, depression, and/or eating disorders. To extend the functional
interpretation of convergence, the convergent dysgranular insula regions identified in the initial
analysis were used as seeds in an exploratory whole-brain resting state functional connectivity (FC)
analysis. Finally, divergent voxels selective to the engagement of top-down expectancies about the
body were isolated, focusing on temporal windows of the ISO task when the anticipation of
potential interoceptive change was maximal. We predicted that the agranular insula would be
preferentially engaged during these windows, and not engaged during the peak of perturbation or
during interoceptive attention alone.
165
Methods

Participants

Seventy individuals with primary diagnoses of anxiety, depression, and/or eating disorders
(ADE) and 57 healthy comparisons (HC) were included in the study. Diagnoses were based on DSM-
5 criteria using the Mini-International Neuropsychiatric Interview (Sheehan et al. 1998). Comorbid
anxiety and depression were allowed, but those with panic disorder were excluded to reduce the
likelihood of isoproterenol-induced panic anxiety (Pohl et al. 1988) in the scanner. Participants
with a primary diagnosis of anxiety were further required to have scores greater than seven for the
GAD-7 (Spitzer et al. 2006) and greater than 10 for the Overall Anxiety Severity and Impairment
Scale (OASIS) (Campbell-Sills et al. 2009), indicative of clinically significant anxiety levels.
Participants with eating disorders were required to have met lifetime DSM-5 criteria for anorexia
nervosa but were required to be weight restored as defined by a minimum body mass index of 18.5
or greater for at least 1 month to control for the well-described state effects of acute starvation on
neural activation (Wierenga et al. 2015). We used several self-report scales to measure symptom
severity across the sample due to our interest in transdiagnostic markers of pathology.
Psychotropic medications were allowed as long as there was no change in dosage during the four
weeks preceding the MRI. Finally, a medical evaluation including a 12-lead electrocardiogram read
by a cardiologist ensured all participants were generally healthy, free of cardiorespiratory,
endocrine, metabolic, or other illnesses, as well as free of major neurological disorders or other
psychiatric disorders besides ADE. Additional exclusion criteria involved anything that would
render the participant unfit for the MRI environment, such as inability to remove body piercings or
implanted electrodes. The study was part of a larger project investigating interoception in anorexia
nervosa and generalized anxiety disorder (ClinicalTrials.gov Identifier: NCT02615119). Additional
recruitment, screening, and exclusion information can be found in (Teed et al. 2022), which used a
subset of the current sample for a separate analysis.
All study procedures were approved by the Western Institutional Review Board, and all
participants provided written informed consent before participation. All data were collected at the
Laureate Institute for Brain Research in Tulsa, OK.
194
195 ***Experimental design and statistical analysis***

During a single scanning session, participants completed two fMRI tasks designed to preferentially assess intersecting aspects of cardiorespiratory interoception while measuring blood oxygen level-dependent (BOLD) signals (Figure 1). The Isoproterenol Infusion (ISO) task was always performed before the Visceral Interoceptive Attention (VIA) task, which we considered to preferentially emphasize the bottom-up or top-down processing of interoceptive signals, respectively. The fixed order of these tasks reflected the project's primary emphasis on acquiring the infusion data. Complete details of these paradigms are described in the following sections. Both tasks were immediately preceded by and followed by an eight-minute eyes-open resting state scan. For the resting state scans, participants were instructed to remain as still as possible, to keep their eyes open and fixated on a white cross centered upon a dark screen, and to "clear your mind and do not think about anything in particular." To minimize head motion during scanning, a layer of tape was gently placed over the participant's forehead to provide them with tactile feedback regarding head position. Cardiac and respiratory activity was measured simultaneously at 40Hz during all scans using a pulse oximeter and thoracic respiratory belt, respectively. Finally, participants completed a high-resolution anatomical MRI scan before any resting state or task-based imaging data was collected.

Isoproterenol Infusion Task

Intravenous bolus infusions of the peripherally-acting beta-adrenergic agonist isoproterenol (ISO) were used to manipulate afferent interoceptive signals in a bottom-up fashion. Infusions of 2.0 micrograms (mcg), 0.5 mcg, or Saline were administered 60 seconds into each four-minute infusion scan as per Hassanpour et al. (Hassanpour et al. 2018) and Teed et al. (Teed et al. 2022). Infusion delivery was randomized, double-blinded, and repeated twice for each dose for a total of six infusions (*i.e.,* six infusion scans). Throughout each infusion scan, participants were instructed to focus their attention on the sensations from their heart and lungs while providing real-time, continuous intensity ratings using an MRI-compatible dial (Current Designs Inc.) with their dominant hand, the rating of which was visible to them on the screen while scanning. The intensity rating scale ranged from 0 (none or normal) to 10 (most ever). Thus, this task engaged top-down attention towards cardiorespiratory sensations while parametrically perturbing the bottom-up stream in the ISO trials, with an absence of such perturbation in Saline trials. Following our lab's previous studies (Hassanpour et al. 2018; Teed et al. 2022) we split the time course of this task into pre-defined epochs of interest, and focused on the Anticipation period (seconds 60 to 80, directly following the infusion but before the ISO-induced rise in heart rate) and the Peak period (seconds 80 to 120, encompassing the onset and peak magnitude of ISO-induced heart rate change), and contrasted these with the Baseline period (the initial 45 seconds of the task). Following each scan, participants also retrospectively reported the intensity of perceived cardiac or respiratory sensations, separately, that they felt during the scan, as well as their anxiety or excitement. To familiarize participants with the task and isoproterenol-induced sensations, single-blinded infusions of 1mcg ISO and Saline (one trial each) were given before the MRI session.

Visceral Interoception Attention Task

The Visceral Interoceptive Attention (VIA) task was used to assess the top-down attentional processing of interoceptive signals at rest (*i.e.,* no infusions were delivered). This minimalistic task, adapted from Simmons et al. (Simmons et al. 2013), involved repeated trials of interoceptive or exteroceptive attention delivered across three six-minute runs, with the order of these runs randomized across participants. During the interoceptive attention trials, the words "HEART", "LUNGS", or "STOMACH" were presented on the screen for eight seconds, during which participants were instructed to focus their attention on the naturally-occurring sensations experienced from that organ. During the exteroceptive attention control trials, the word "TARGET" was shown on the screen at changing shades of black to grey, and participants were instructed to pay attention to the degree of color change. After half of all trials in each condition participants were asked to indicate via an MRI-compatible button box the intensity of sensations they had perceived from that organ (interoceptive condition) or the intensity of the color change (exteroceptive condition) on a rating scale of 1 (no sensation/no change in color) to 7 (high intensity).

MRI data acquisition

All scans were conducted in a 3 Tesla General Electric scanner with an eight-channel head coil (GE MR750; GE Healthcare). Anatomical images were acquired via T1-weighted, magnetization-prepared, rapid gradient-echo sequences images (MPRAGE). MPRAGE parameters were: 190 axial slices, slice thickness = 0.9mm, TR/TE = 5/2.012 ms, FOV = 240 x 192mm^2, matrix size 256 x 256, flip angle 8 degrees, inversion time 725 ms, SENSE acceleration R =2, with a sampling bandwidth of 31.2. For the task-based and resting state functional scans, T2*-weighted images were acquired via an echoplanar sequence of the following parameters: single-shot gradient echo planar imaging (EPI) sequence covering the whole brain, obtaining 39 axial slices, 2.9mm thick with no gap and a voxel size of 1.875 x 1.875 x 2.9mm^3, with TR/TE = 2000/27ms, FOV = 240 x 240mm^2, flip angle 78 degrees, SENSE acceleration R = 2 with a 96 x 96 matrix.

fMRI task-based analysis

Each task was analyzed separately in AFNI (Cox 1996), following the previously-published preprocessing pipelines specific to each task (Simmons et al. 2013; Teed et al. 2022). Preprocessing steps common to both tasks included removal of the first four volumes (ISO) or five volumes (VIA) for steady state tissue magnetization, despiking, slice time correction, spatial smoothing (6mm full width at half maximum Gaussian kernel), co-registration to the first volume, scaling of each voxel, and normalization to Talairach space. In the ISO task, temporal fluctuations of cardiac and respiratory frequencies and their first-order harmonics were additionally removed from the imaging data using RETROICOR, implemented via custom code in MATLAB (MathWorks).
After preprocessing, whole-brain individual-level activation maps were created. For the ISO task, this involved regressing out baseline drift and contrasting average BOLD percent signal change (PSC) within each epoch of interest (Anticipation and Peak, separately) against the average PSC during the Baseline period. For the VIA task, this involved a multiple linear regression model that included terms for baseline drift, six motion parameters, and each interoceptive and exteroceptive attention condition, and contrasting PSC associated with cardiorespiratory attention (*i.e.,* collapsed across "Heart" and "Lung" trials) against that of exteroceptive attention ("Target"

279   control condition). This cardiorespiratory contrast was specifically chosen as it best matched the
280   cardiorespiratory perturbation induced by ISO.
281   After the creation of individual-level maps, quality control checks were performed.
Participants were excluded from further group analysis if they had poor quality imaging data
according to the previously established approaches in each task. For the ISO task, individual runs
were excluded if more than 20% of all time points had over 0.3mm motion, based on the Euclidean
norm of the six motion parameters; following this, participants were excluded who were missing
both runs of the Saline or 2.0mcg doses. For the VIA task, all time points with an outlier fraction
greater than 5% or an estimated motion greater than 0.3mm were censored, and any run in which
this censoring removed over 20% of volumes was excluded. If a participant was missing two or
more VIA runs (out of three), they were removed from the analysis. To avoid an unbalanced
conjunction analysis in the following step, participants excluded from one task were also removed
from the other task. The groups were then propensity matched based on age, sex, and BMI to
equalize the sample sizes (*MatchIt* function in R), resulting in a final sample of participants who had
high quality imaging data in *both* tasks.
294   Using this propensity-matched sample, group-level statistical activation maps were created
295   for each task and group separately, using whole-brain voxel-wise t-tests (*3dttest++* in AFNI) with
296   Equitable Threshold And Clustering corrections for an overall False Positive Rate of 5% (Cox 2019).
297   With these group activation maps, we were thus able to identify voxels significantly active at the
298   group level in each experimental condition.
299
300   *Convergence maps*

301   To examine voxels co-activated across the ISO and VIA tasks, we restricted the functional
neuroimaging analysis to the specific portions of each task that best exemplified the interoceptive
processes of interest: namely, Heart + Lung interoceptive attention relative to exteroceptive
attention in the VIA task, and the Peak period of the 2.0mcg dose relative to the Baseline period in
the ISO task. We narrowed the focus of analysis to this specific time period and dose of the ISO task
because it represents, as shown in previous studies, the peak level of cardiorespiratory
perturbation that can be confidently ascertained by subjective report in all individuals (Hassanpour
et al. 2018; Teed et al. 2022). Using the statistically-thresholded group activation maps from each
task, binarized into active and non-active voxels, we created convergence maps by joining the
binary maps across the specified contrasts of each task. These convergence maps thus identified
which voxels were commonly activated, *i.e.*, *convergent*, at the group level during both tasks. Due to
our pre-specified focus on cytoarchitectural processing of the insular cortex, we subdivided these
convergence maps (after converting to MNI space) into three bilateral insular subregions of
interest, defined using the Brainnetome probabilistic cytoarchitectonic atlas (Fan et al. 2016), as
follows: agranular anterior (regions 165 and 167 for the left hemisphere, 166 and 168 for the
right), dysgranular mid (regions 169 and 173 for the left hemisphere, 170 and 174 for the right),
and granular posterior (region 163 and 171 for the left hemisphere, 164 and 172 for the right).
318   Each convergence map was evaluated in four ways: 1) the spatial extent of convergence (*i.e.,*
number of convergent voxels) within each insular subregion, tested for group differences via chi-
square tests, 2) the spatial similarity of patterns of convergence between HC versus ADE groups,

measured via Dice coefficient separately for each hemisphere (using AFNI's *3ddot*), 3) the degree of
task overlap within each group and hemisphere separately, quantified via the overlap coefficient
(*i.e.,* the proportion of the ISO cluster overlapping with the VIA cluster, as per (Bowring, Maumet,
and Nichols 2019), and finally 4) the average PSC across convergent voxels within each
cytoarchitectonic subregion, compared across groups and hemispheres using linear mixed effects
regression (*lmerTest* library in R, with the following equation: percent signal change ~ group *
hemisphere + (1|subject)). In this linear model, the fixed effects of group and hemisphere were
tested with respect to the magnitude of activation within subregion-specific convergent voxels, and
significant main effects were interpreted using post-hoc contrasts between the estimated marginal
means of each condition (*emmeans* library in R). Additionally, linear regression was used to
examine possible associations between PSC in hemisphere- and subregion-specific convergent
voxels and the following variables: trait measures of anxiety and depression, the objective degree of
peripheral perturbation (*i.e.,* mean change in heart-rate during the Peak period of the 2.0mcg dose),
real-time subjective rating of cardiorespiratory intensity (*i.e.,* mean change in dial rating during the
Peak period of the 2.0mcg dose), and retrospective ratings of cardiac and respiratory intensity,
anxiety, and excitement following the 2.0mcg dose. For these associations, we used a dimensional
approach that collapsed across groups, considering all participants on a continuous scale.
338
fMRI resting state analysis

In an exploratory whole-brain resting state functional connectivity (FC) analysis, we aimed
to identify which connections, if any, between convergent regions and the rest of the brain were
altered following the interoceptive manipulations, and if these patterns of connectivity change
differed between groups. We predicted that residual group-specific changes in functional
connectivity of the insular regions activated during the ISO and VIA tasks would be present in the
resting state scan at the end of the scanning session, reflecting additional neural signatures of
interoceptive processing. We did not include a temporal control condition for this exploratory
analysis, and thus we were unable to examine the specificity of insular connectivity changes across
time in relation to test-retest reliability measures of the group's resting state data.
Resting state scans were analyzed in AFNI with a preprocessing pipeline optimized for
connectivity analysis. After discarding the first five images to ensure steady-state tissue
magnetization, we removed spikes in the signal intensity time course, and removed physiological
noise using respiration volume per time (RVT) (Birn et al. 2008) and RETROICOR (Glover, Li, and
Ress 2000) regressors that model slow blood oxygenation level fluctuations and time-locked
cardiac and respiratory artifacts, respectively. The RVT regressors consisted of the RVT function
and four delayed terms at 5, 10, 15, and 20 seconds, and the RETROICOR regressors consisted of
four respiratory and four cardiac generated on a slice-wise basis by AFNI's "RetroTS.m" script (Jo et
al. 2010). Further preprocessing steps included slice-timing correction, rigid body realignment,
warping to MNI space, and smoothing with a 6-mm full-width half-maximum Gaussian kernel. Next,
we applied a high-pass filter to remove all non-relevant ultra-slow fluctuations below 0.01 Hz, and
then further reduced potential contributions of non-neural sources by regressing out the following
nuisance variables: (1) 12 head motion regressors (6 realignment parameters and their
derivatives), (2) voxel-wise local white matter regressors, and (3) three principal components of
ventricle signals (ANATICOR). Following preprocessing, any subject with greater than 10% of

volumes above our motion cut-off of 0.3mm for the Euclidean norm of the motion derivatives was
excluded from further analysis.
From the pre- and post-interoception resting state scans, time series were extracted from
the convergent ROIs within the left and right hemispheres for HC and ADE groups separately, and
correlated these respective time series against those of all other voxels in the brain to create an FC
map for each resting state period (i.e., pre- and post-interoception). The resulting correlation
coefficients were converted to Fisher z-statistics to produce a more normally distributed variable.
To examine group differences in the effect of time (*i.e.,* pre- versus post-interoception tasks) on
resting state FC, all resulting clusters of this group*time interaction analysis were corrected for
multiple comparisons using the *3dClustSim* function in AFNI, using 10,000 Monte Carlo simulations
of whole-brain fMRI data to determine the cluster size at which the false positive probability was
below $\alpha < 0.05$ with an uncorrected voxel-wise threshold of $p < 0.001$, yielding a threshold of 150
contiguous voxels. This approach allowed the identification of regions that changed connectivity
with the convergent insula differently for HC vs. ADE groups following top-down and bottom-up
interoception. Finally, we conducted a correlational analysis to examine whether individual
changes in FC were related to trait measures of anxiety and depression in the transdiagnostic
sample, using the Bonferroni method for multiple comparison correction. These correlations were
performed to help interpret the functional significance of residual changes, based on previous
studies showing trait-related differences following interoception (De la Cruz et al. 2023).
383
384
Divergence maps

To isolate voxels selective to the engagement of top-down expectancies about the body, we
focused on specific time windows of the ISO task were where expectations of potential
interoceptive change were preferentially engaged: namely, the Anticipation period across all doses,
and the Peak period of the Saline dose. The Anticipation period covered the 20 second epoch right
after the intravenous infusion was delivered but before the onset of any cardiorespiratory
perturbation (in the case of an isoproterenol dose administration). Thus, the Anticipation period
represented a time when no peripheral change was occurring, but also one where participants were
unsure about whether or not such a change was coming. This state of uncertainty would continue
into the Peak period of the Saline dose, where again, no peripheral physiological change was
induced. In the setting of isoproterenol dose administration, this state of uncertainty would resolve
when participants correctly identified and rated the onset of changes in cardiorespiratory
sensation. Based on previous studies from our lab using this task, we expected that by the end of
the Saline Peak window some participants would update their beliefs about what had occurred
after realizing that they had not reported experiencing a cardiorespiratory perturbation (i.e., their
dial ratings remained at zero). Together, the Anticipation period and the ensuing Peak period of the
Saline dose share some similarities with the VIA task in their involvement of top-down
interoceptive attention during physiological rest, but critically, they differ in their added
expectancies about possible changes in the state of the body. Thus, by isolating voxels preferentially
active during these ISO anticipatory windows that were not active during VIA top-down attention
alone, it we presumed it would be possible to separate the effect of bodily expectancies from the
general effect of bodily attention.

For this aim, group-level activation maps from the Anticipation period (averaged across all doses) and the Peak period of the Saline dose, both relative to the Baseline period, were examined. Like the convergence map approach described above, these maps were binarized within each period and group separately, transformed to MNI space, reduced to the cytoarchitecturally-subdivided insular cortex, and joined with the binarized activation map for the Heart and Lung relative to Target contrast in the VIA task. Here, our focus was instead on the *divergent* voxels, *i.e.,* on those that were active only during specific periods of the ISO task and that were not co-active during the VIA task. As per our hypothesis regarding anterior insula involvement in interoceptive expectancy processing, these "ISO task only" voxels were restricted to those within the agranular anterior insula, and were evaluated in two ways: 1) spatial extent of activation (*i.e.,* number of voxels) across groups, time periods, and hemispheres, examining differences via chi-square tests with Bonferroni correction for multiple comparisons, and 2) magnitude of activation, examining differences in average PSC across the fixed effects of group, hemisphere, and time period using linear mixed effects regression and interpreting significant main effects using post-hoc contrasts of estimated marginal means (as previously described). Finally, we examined possible associations between average PSC in these expectancy-specific agranular regions and trait measures of anxiety using linear regression, transdiagnostically across groups.

Physiological and behavioral analysis

To test for group differences in the time courses of heart rate and real-time dial ratings during the ISO task (2.0mcg isoproterenol and Saline infusions), we used cluster-based permutation testing (*permutes* package in R) separately for each dose after averaging across both trials. A significant group difference was defined as any contiguous set of time points (two or more continuous seconds) where permutation testing revealed a statistical difference at $p < 0.05$. We used linear regression transdiagnostically to test if the magnitude of heart rate or dial rating change, when averaged across each epoch of interest, was associated with dimensional measures of trait anxiety and depression across groups. Finally, we used two-tailed t-tests separately for each task and condition to test group differences in retrospective ratings of interoceptive intensity, anxiety, and excitement in the ISO task and interoceptive and exteroceptive intensity in the VIA task. This approach reflected our primary interest in investigating group differences in perceptual ratings, rather than performing additional within-group comparisons across modalities or doses.

Code accessibility

Custom AFNI and R scripts are posted on the Open Science Framework preregistration for this project (Adamic et al. 2021).

Results

Of the 127 total participants included in this study, 57 ADE individuals and 46 HC individuals fulfilled criteria for high-quality brain imaging data with both tasks. In line with the predominant female prevalence of these disorders, most of the participants were female. To obtain

equal-sized groups for group-level comparison, propensity matching on age, self-reported sex, and body mass index resulted in 46 individuals per group (Table 1). After propensity matching, the two groups were not significantly different regarding age, body mass index, and sex, but they differed significantly on trait measures of anxiety (State-Trait Anxiety Inventory; STAI), anxiety sensitivity (Anxiety Sensitivity Index; ASI, Total and subscales), depression (Patient Health Questionnaire-9; PHQ-9) (Kroenke, Spitzer, and Williams 2001), and eating disorders (Sick, Control, One, Fat, Food questionnaire; SCOFF) (Kutz et al. 2020). All participants in the ADE group exhibited some form of psychiatric comorbidity. Across the group, 67% had a diagnosis of GAD (current or lifetime), 72% of major depression (current or remitted), and 35% of anorexia nervosa (current or lifetime). Most ADE participants reported taking psychiatric medication (see Table 1 for details).

Convergence analysis

Bilateral patches of the ventral mid-insular cortex were commonly activated during the ISO and VIA tasks (Figure 2) (Table 2). Quantification of the spatial extent of this convergence in the pre-defined probabilistic agranular, dysgranular and granular subregions revealed that these convergent clusters were localized primarily to the dysgranular insula bilaterally (Figure 3A), but with a hemispheric asymmetry across groups: the ADE group showed a greater proportion of overlapping voxels within the left dysgranular insula (χ^2_1 = 26.7, p<0.001 for group comparison) whereas the HC group showed a greater number of overlapping voxels within the right dysgranular insula (χ^2_1 = 26.7, p<0.001 for group comparison) (Figure 3B). Correspondingly, the spatial similarity of these convergently activated voxels between the groups was lower for the left dysgranular insula (Dice coefficient of 0.58) than for the right (Dice coefficient of 0.78) (Figure 2). In addition to these group hemispheric differences in spatial extent, there was a lower amount of overlap between the two tasks in the left dysgranular insula for both groups (left hemisphere overlap coefficients: 0.42 and 0.63 for HC and ADE, respectively; right hemisphere overlap coefficients: 0.83 and 0.78, for HC and ADE, respectively).

Despite these group differences in the spatial extent and spatial similarity of convergence, the magnitude of activation within convergent dysgranular subregions during the Peak of perturbation (ISO 2.0mcg dose) or Heart and Lung attention (VIA task) was not different between groups ($F_{1,90}$ = 0.64 and 0.05, p = 0.43 and 0.82 for main effect of group in ISO and VIA tasks, respectively). However, during both of these task conditions, activation was greater within the right hemisphere than the left ($F_{1,90}$ = 6.13 and 5.10, p = 0.015 and 0.027 for main effect of hemisphere during ISO and VIA tasks, respectively; t_{90} = -2.48 and -2.25, p = 0.02 and 0.03 for post-hoc contrasts between the left versus right hemisphere when collapsed across groups) (Figure 4A). This hemispheric effect on activation magnitude was specific to interoceptive processing, as no similar hemispheric difference was seen during the exteroceptive condition of the VIA task ($F_{1,90}$ =0.68, p=0.41). Furthermore, the magnitude of activation within the right (but not the left) dysgranular convergent region during the peak of cardiorespiratory perturbation was associated with the intensity of real-time cardiorespiratory sensations (*i.e.,* mean dial rating) during this period when examined across the entire sample (Pearson's *r* = 0.12 and 0.21, *p* = 0.25 and 0.046 for the left and right hemispheres, respectively; Figure 4B). Conversely, PSC in either convergent hemisphere during perturbation was not associated with objective measures of physiological arousal (*i.e.,* mean

change in heart rate, Pearson's r = 0.11 and 0.09, p = 0.31 and 0.40 for left and right hemispheres,
respectively), post-scan ratings of cardiac intensity (Pearson's r = 0.16 and 0.20, p = 0.15 and 0.06),
post-scan ratings of respiratory intensity (Pearson's r = 0.12 and 0.06, p = 0.26 and 0.58 for left and
right hemispheres, respectively), post-scan ratings of state anxiety (Pearson's r = 0.14 and 0.08, p =
0.20 and 0.49 for left and right hemispheres, respectively), or trait measures of anxiety sensitivity
(*i.e.,* ASI-Total, Pearson's r = 0.12 and 0.16, p = 0.25 and 0.13 for left and right hemispheres,
respectively).
Finally, pre- to post-task resting state functional connectivity of the convergent regions
showed a pattern of differential change in the ADE compared to HC group: functional connectivity
between the right dysgranular convergent cluster and left middle frontal gyrus increased in the
ADE group only, while the HC group saw no differences (p < 0.05, corrected; Figure 5A and 5B).
This was the only significant cluster showing connectivity differences between groups. When
looking at this insular-IFG connection across the entire sample, the magnitude of functional
connectivity change between these two areas was associated with trait anxiety (Pearson's r = 0.43
and 0.36 for OASIS and GAD7, respectively, p < 0.001) and trait depression (Pearson's r = 0.38, p <
0.001 for PHQ-9) (Figure 5C) but not the magnitude of ISO-induced heart rate (Pearson's r = -0.05,
p = 0.66) or dial rating (Pearson's r = 0.01, p = 0.90) at the 2.0μg dose.
506
Divergence analysis

The Anticipation period of the ISO task engaged large swaths of the bilateral agranular
insular cortex that were not activated at any point during the VIA task (Figure 6A) (Table 3). During
this period, the right hemisphere was activated to a greater spatial extent than the left in both
groups ($\chi^2{}_1$ = 22.7 and 7.3, p < 0.001 and p=0.007 for left vs. right in the HC and ADE groups,
respectively), and to a greater spatial extent in HC versus ADE individuals in the right hemisphere
only ($\chi^2{}_1$ = 7.3, p=0.007 for group comparison). There was no group difference in the spatial extent
in the left hemisphere ($\chi^2{}_1$ = 2.0, p = 0.15) (Figure 6B). Notably, these spatial patterns of activation
were similar between groups (Dice coefficient =0.86 in the left and 0.87 in the right).
During the Peak period of the Saline dose, the spatial extent of activation in the agranular
insula increased to include significantly more voxels, for both groups and hemispheres ($\chi^2{}_1$ = 75.3
and 71.4, p < 0.001 for Anticipation vs. Peak in the left and right hemispheres in HC; $\chi^2{}_1$ = 36.3 and
12.1, p < 0.001 for Anticipation vs. Peak in the left and right hemispheres in ADE) (Figure 6B). This
activation expansion occurred to a greater degree in the HC versus ADE group ($\chi^2{}_1$ = 17.2 and 70.3,
corrected p<0.001 for HC versus ADE in the left and right hemispheres, respectively). Finally,
agranular insula recruitment within the HC group during the Saline Peak period was shifted
towards the right hemisphere (*i.e.*, more active voxels in the right than the left, $\chi^2{}_1$ = 24.3, p <
0.001), while the ADE group showed no hemispheric difference during this period ($\chi^2{}_1$ = 0.10, p =
0.75). Despite this difference in number of activated voxels, the spatial patterns of activation were
still largely overlapping between groups (Dice coefficient = 0.88 in the left, 0.82 in the right).
Along with expansions in the spatial extent of activation, the signal intensity in the
agranular insula increased from the Anticipation period to the Saline Peak period ($F_{1,270}$=28,
p<0.0001 for main effect of epoch), with no differences between groups ($F_{1,270}$ = 0.25,p = 0.62 for
the main effects of group) (Figure 6C). This increase occurred in both hemispheres (t_{270} = 2.7 and
4.8, p = 0.008 and < 0.001 for Saline Peak – Anticipation contrast in the left and right hemispheres,

respectively, when collapsed across groups). Notably, the right hemisphere was activated more
strongly than the left during the Saline Peak period (t_{270} = -3.0, p = 0.003), whereas there was no
difference across hemispheres in the Anticipation period (t_{270} = -0.8, p = 0.41). The magnitude of
agranular insula activation was not related to trait measures of anxiety sensitivity in either epoch
or hemisphere when examined transdiagnostically across groups (ASI * hemispheric interaction
effect: $F_{1,90}$ = 1.9, p = 0.17 for Anticipation, and $F_{1,90}$ = 1.7, p = 0.19 for Saline Peak).
538
Physiological response and perceptual ratings

During the 2.0mcg dose there were large perturbations in heart rate in both groups, and
while the ADE appeared to have a slightly greater response on average, this reached statistical
significance via permutation testing only during two continuous time points in the Peak period
(seconds 98 through 99, t = 1.7 and 1.5, p = 0.044 and 0.045, Figure 7A). However, the ADE group
reported significantly higher real-time dial ratings for the latter portion of the Peak period
extending into much of the ensuing recovery period (seconds 110 through 142, t = 1.8–2.3, p =
0.008–0.042, Figure 7B). During the Saline dose of the ISO task, both heart rate and perceptual dial
ratings remained near baseline levels. While heart rate did not differ between groups (t = -0.53 –
1.64, p = 0.05 – 1.00), the HC group showed small but statistically significant increases in dial
ratings compared to those of the ADE group for portions of the Peak period and the ensuing
recovery period (t = -1.67 – -2.60, p = 0.005 – 0.048) (Figure 7A and 7B). Although there were ISO-
induced differences in continuous measures of cardiorespiratory sensation, there were no group
differences in retrospective cardiac or respiratory intensity ratings for the 2.0 mcg infusion (t_{80} =
1.5 and 1.2 for heart and lungs respectively, p = 0.13 and 0.25) or the saline infusion (t_{80} = 0.3 and
1.2, p = 0.75 and 0.22) (Figure 7C). The ADE group reported greater retrospective anxiety than the
HC group following infusions (t_{85} = 4.8 and 3.8 for 2.0mcg and Saline, respectively, p < 0.001), with
no differences in excitement (t_{85} = 0.2 and 0.9, p = 0.80 and 0.37 for 2.0mcg and Saline,
respectively). During the VIA task, the ADE group reported higher cardiac intensity ratings than the
HC group during heart-focused attention (t_{89} = 2.0, p = 0.04) but lower exteroceptive intensity
ratings during the target condition (t_{90} = 2.7, p < 0.01). There were no group differences in the
intensity of reported lung or stomach sensations (t_{89} = 1.6 and 1.4, p = 0.10 and 0.16) (Figure 7D).
561

Discussion

The current study revealed overlapping patterns of neural activation within the dysgranular
mid-insula during top-down and bottom-up interoceptive processing, corroborating previous
studies examining each task independently (Hassanpour et al. 2018, Simmons et al. 2013), and
supporting the notion of the insula as a key functional interface between goal-directed and sensory-
driven interoceptive streams. Importantly, hemispheric differences in the patterns of dysgranular
activation distinguished individuals with anxiety, depression, and eating disorders from healthy
comparisons, as hypothesized. These hemispheric asymmetries, and the disparate spatial patterns
within the left dysgranular insula, support the conceptualization of this subregion as a "locus of
disruption" for interoceptive symptomatology in these disorders (Nord et al. 2021). The
hemispheric imbalance was concurrent with a heightened perceptual response in the clinical
sample, as well as with changes in right dysgranular insula-to-left middle frontal gyrus functional

connectivity following the interoceptive tasks. Finally, activity within the agranular insula was
selective to periods of increased interoceptive expectancies, aligning with a prediction-generating
role for this region and one that was distinct from the convergent effects seen in the dysgranular
mid-insular subregion.
Top-down processing commonly refers to cognitive influences on perception, such as
attention, memory, and expectation, that putatively help the brain to make sense of the stimuli it
encounters. In contrast, bottom-up processing commonly refers to the stimulus-driven signals that
are processed hierarchically as they reach the brain. With respect to interoception, top-down
processing can refer to the brain's ability to predict or interpret how the internal body should feel
(*i.e.,* the expected interoceptive signals) based on the previous or current environmental context. In
turn, bottom-up processing can refer to the actual state of the body in the current moment. For
instance, top-down influences may include the anticipation of an upcoming meal leading to feelings
of hunger, while bottom-up influences may refer to actual hunger signals from the body at lunch
time, such as hormonal concentrations of ghrelin/leptin, and stomach contractions. The continuous
balance between these processes determines the qualia of conscious perceptions and has been
suggested to be key to health and the optimization of internal and subjective states (Sterling 2012).
However, as processes, they are rarely studied concomitantly in the manner conducted in this
study. The convergence identified between top-down and bottom-up interoceptive processing
within the dysgranular mid-insula suggests that this cortical subregion is an area where further
detail regarding the reciprocal interactions underpinning interoceptive perceptions should be
investigated, particularly in relation to psychiatric disorders involving disturbances of mood,
anxiety, or appetite.
Although the current findings emphasize the role of interoceptive processing within
dysgranular subregions of the insula, a number of experimental and analytic next steps are worth
considering. Due to the broad systems-level of inference allowed by fMRI methodology and the
relatively rudimentary block averaging analytic approach employed, additional measurements are
warranted to establish the cellular and molecular mechanisms underlying the observed
dysgranular co-activation. Such approaches would be important for verifying a true intersection of
interoceptive neurocircuitry in which connecting top-down attention and prediction circuits
directly modulate, or are modulated by, incoming sensory afferences. An alternative outcome could
be identifying an overpassing flow of signals across these circuits within the same region.
Implementing high resolution approaches to imaging these processes via high-field fMRI on a trial-
by-trial basis could inform the laminar architecture underpinning attention-driven modulation of
incoming viscerosensory signals as well as allowing for computational modelling of these processes
at the level of cortical laminar subfields (Kasper et al. 2022). In an interoceptive application of this
analyses employing a respiratory bottom-up perturbation approach, activation of the anterior
insula was associated with the prediction certainty but not prediction error for different inspiratory
breathing loads, with parameters distinguishing between high versus low trait anxiety individuals
(Harrison et al. 2021). This computational approach could be adapted for cardiorespiratory
processing within the dysgranular insula. While typically restricted to high field strengths, such as 7
Tesla, there are emerging reports that laminar imaging can be performed at lower field strengths
(Knudsen et al. 2023). Thus, there are a number of currently feasible steps for further extending the
present findings.
While we did not set out to test predictive coding or other computational models in the
current study, certain computationally relevant theories provide useful frameworks for
interpreting the observed dysgranular co-activation during each imaging task. These include the

Embodied Predictive Interoceptive Coding (EPIC) model (Barrett and Simmons 2015), describing
laminar-specific hypotheses for the interactions between anterior-mediated interoceptive
predictions and posterior-mediated viscerosensation across the antero-posterior axis of the insula,
and the Insula Hierarchical Modular Adaptive Interoception Control model (Fermin et al. 2022)
using the hierarchical tripartite subdivision of the insula and describing parallel networks within
prefrontal cortical and striatal regions that underpin higher-order interoceptive representation. For
additional theoretical discussions of the computational functional neuroanatomy of interoception,
see Allen et al. (2022), Paulus, Feinstein, and Khalsa (2019), and Petzschner et al. (2021).
Collectively, these theories can be interpreted to suggest a model of bidirectional signal flow along
the antero-posterior axis of the insula: while the granular posterior subregions may be the first to
receive ascending visceral input, the agranular anterior subregions continuously issue predictions
regarding the anticipated states of such input (Barrett and Simmons 2015). From this perspective,
the dysgranular mid-insula is well-positioned between the agranular and granular poles to
integrate the signals from each source. Computations within dysgranular layers can link incoming
sensations with top-down expectations (as suggested in the EPIC model), effectively translating
visceral signals into endogenously meaningful interoceptive prediction error signals, and providing
the rest of the brain with an internal context for interpreting the current bodily state (Seth and
Friston 2016).
Applying computational frameworks for interoception to the present findings may also be
useful for interpreting the broad agranular insula activation found only during periods of increased
bodily expectancies (*i.e.*, "divergence"). This subregion's engagement during the Anticipation
window and expansion throughout the Peak period of the Saline dose likely reflects an increase in
predictive processing during these conditions, as both groups expected a potential perturbation
and were subsequently required to update their predictions upon experiencing a physiologically
quiescent state instead. The reduced spatial extent of this activation in the ADE versus HC group
could plausibly stem from an overreliance on top-down predictions and perhaps a pruning of
spatial activation in the associated regions, similar to the functional alterations of neural activity
that have been observed following the acquisition of a motor skill (Wiestler and Diedrichsen 2013).
Such 'efficient' neural exchanges might even result from a chronic inability to adjust confidence in
bodily input in relation to its ambiguity (*i.e.,* increasing confidence in visceral signals during bodily
perturbation), and could occur alongside maladaptive allostatic interoceptive adjustments
described in psychiatric disorders (Barrett, Quigley, and Hamilton 2016; Smith et al. 2020).
Computational theories of interoception can also be applied to the observed co-activation within
the dysgranular mid-insula (i.e., "convergence"). Dysgranular convergence aligns with our proposed
intersectional (as opposed to overpassing) role for signal processing this region, and could
represent the generation of prediction errors as inflows of isoproterenol-induced viscerosensation
are compared to task-specific interoceptive predictions focused by the lens of directed attention.
While internally-directed attention alone might selectively enhance cortical representations of the
viscera across wide swaths of insula, as suggested previously (Avery et al. 2014; Pollatos et al.
2007), these past studies focus narrowly on perceptions of the viscera during physiological
quiescent states when visceral signals are relatively ambiguous. By manipulating cardiorespiratory
signals during the ISO 2.0mcg dose and eliciting unambiguous perceptual states, the current study
effectively spotlights the dysgranular subregion's role in heightened interoceptive awareness and

provides a neuroanatomical focus for further examinations of bi-directional interoceptive
processing in this important region.
665   The right-hemispheric increase in mid-insula activity, as well as the hemispheric
asymmetries and altered functional connectivity found in the ADE group has several implications
for an interoceptive processing role for the dysgranular insula. Previous authors have suggested a
right-hemispheric dominance in interoceptive processing in healthy populations (Craig 2009;
Schulz 2016), and have pointed towards the right anterior insula as the key hub for body awareness
(Craig 2009). While a rightward emphasis in the magnitude of insular activation was found during
ISO-induced cardiorespiratory sensations across both groups, this activity was located more
posteriorly, towards the mid insula (similar to previous isoproterenol fMRI studies (Hassanpour et
al. 2016, 2018)), where it was correlated with cardiorespiratory sensation. While this finding
supports a role for the mid-insula as an integrative hub for interoceptive awareness, it was in
contrast to a lack of clear associations with physiological signals, emotional feelings such as anxiety
or excitement or trait measures of ADE severity, despite the evocation of greater state anxiety
following perturbation in the ADE individuals. Probing the symptomatic relevance these potential
brain-behaviour relationships may call for more refined approaches to the study of interoceptive
perturbation. One option might be to evaluate the intersecting influence of several forms of
perturbation, by studying both exogenous approaches such as isoproterenol infusion, and
endogenous approaches such as mental stressors (i.e., Stroop task (Gianaros et al. 2005)), physical
stressors (i.e., Valsalva (Song et al. 2018) or lower body negative pressure (Kimmerly et al. 2005)),
or social stressors (i.e., Trier Social Stress task (Allen et al. 2017)) within the same individuals.
684   Despite observing several right hemispheric associations across both groups, the ADE group
demonstrated a clear leftward activation asymmetry and distinct activation patterns in the left
dysgranular mid-insula across both the ISO and VIA tasks. These left insula abnormalities are
compatible with the "locus of disruption" identified by Nord et al. (2021), when conducting an fMRI
meta-analysis of existing top-down or bottom-up interoception studies of various psychiatric
populations. However, beyond identifying a plausible region of cortex, the precise mechanisms
underlying this disruption remain unclear. Assuming an intersectional role involving intermingling
of top-down and bottom-up streams, it seems plausible that numerous mechanisms could
contribute to the aberrant processing identified in the current study. Such mechanisms include a
reduced attentional enhancement of ascending visceral input (Avery et al. 2014; Kerr et al. 2016),
and/or an inflexibility in adjusting to the salience of interoceptive signals occurring during
perturbation, as revealed by computational modeling (Smith et al. 2020). Reduced flexibility of
neural processing may also explain the singular finding of altered connectivity between the right
mid-insula and left middle frontal gyrus following the interoceptive tasks in the ADE group, as this
frontal region is likely involved in in attentional control including attention bias to threat or
internal representations (Henseler et al. 2011; Sylvester et al. 2003; White et al. 2016). An inability
to return this coupling to baseline following interoceptive engagement (or a shift in the opposing
direction than HCs) could suggest a dysregulated attentional switching mechanism, particularly
since the degree of connectivity change was associated with trait measures of anxiety and
depression severity.
704   From an affective standpoint, the observed hemispheric divergence in interoceptive
processing is worth considering in the context of the functional lateralization of emotion theories

proposed within the insula. Such theories (Craig, 2005; Davidson et al., 1999, 2004; Picard et al.,
2016), supported by meta-analytic (Murphy, Nimmo-Smith, and Lawrence 2003) and human brain
stimulation Oppenheimer et al. 1992) findings, have attributed the left anterior insula to the
energy-enriching behaviors of the parasympathetic nervous system, and the right anterior insula
with the energy-expending ones of the sympathetic nervous system (Oppenheimer et al. 1992). The
current finding of right-hemispheric dominance for conscious awareness of isoproterenol (an
arousal-like, and thus sympathetic-emulating manipulation), as well as the increased signal
intensity in the right anterior insula during the expectancy of such manipulation, may be
considered aligned with this perspective. Correspondingly, activity within the right insula has been
associated with conscious awareness of threat (Critchley, Mathias, and Dolan 2002), and mouse
models have shown causal evidence for the right anterior insula in orchestrating stress responses
to aversive visceral stimuli (Wu et al. 2020). However, assumptions of autonomic opponency are
likely overly simplistic, and it is more appropriate to consider these systems as functionally coupled
across spinal and vagal systems instead of anatomically distinct (Jänig 2006; Jänig et al. 2017). In
this manner, an admittedly speculative emotional lateralization account of functional coupling
might posit that the right anterior insula generates interoceptive predictions of spinal efference and
the left anterior insula for those of vagal efference; in turn the respective dysgranular areas would
then be specialized towards receiving and computing the prediction errors of these different inputs.
However, since there does not appear to be an neuroanatomical lateralization for afferent tracts
(*i.e.,* spinal input does not have a preference for the right hemisphere, and vagal not for the left)
(Dum, Levinthal, and Strick 2009; Shipley and Sanders 1982), it seems more plausible that the
contribution of interoceptive processing to emotion requires a carefully integrated balance of both
autonomic systems across both hemispheres. Based on the present data, such a balance appears to
be asymmetrically shifted in the setting of psychiatric disorders affecting the anxiety, depression,
and disordered eating spectrums, particularly during modulations of interoceptive state.
There are several limitations and strengths of the present approach to consider. First, while
the quantification of insula asymmetries was strengthened by the use of previously established
task-specific pipelines and justified by cytoarchitectural hypotheses leveraging prominent theories
about interoceptive neurocircuitry, several analysis steps differed between the tasks that could
impact the observed results, including the use of a block versus event-related design and
differences in the regression of physiological noise and motion correction. This pre-specified
approach was intentional. It allowed confidence that the analysis metrics were sound (i.e., that a
voxel counted as co-activated would have good task-validity in relation to the methods of
previously published ISO and VIA studies). However, it also meant that there was variability within-
individuals in how the data was handled for each task. Future approaches involving harmonized
pre-processing steps across both tasks may be considered. Second, the present approach to
determining 'convergence' was admittedly simplistic in relying on co-activation as a determinant.
Applying more sophisticated analysis techniques, such as those amenable to individual-level
pattern analyses (as opposed to group activation maps) would provide greater insight into the
individual responses of each participant. Such approaches are worth considering as they are a core
component of personalized evaluations applying functional neuroimaging to diagnostic validation
or treatment selection (Goldstein-Piekarski et al. 2022). Third, while the use of tripartite insular
subregions was appropriate to the level of resolution in our data, future investigations could

employ high resolution fMRI to directly test the laminar-specific hypotheses of the predictive
coding theories we have discussed. Fourth, while the conceptual approach to the top-
down/bottom-up construct may mask the complexity of the brain (Rauss and Pourtois 2013) and
the process of attention (Hommel et al. 2019), the integration of this concept with several
behaviorally relevant tasks in the same individuals offers a useful and informative heuristic in its
application. Finally, while our results focus on the insula's role in cardiorespiratory awareness, it's
important to note that the insula isn't the only brain area involved in sensing internal body states
(Berntson and Khalsa 2021; Mayeli et al. 2023). Future research should explore structural and
functional relationships between the insula and the rest of the brain; for example, relationships
between white matter tracts (Ghaziri et al. 2017, 2018), grey matter volume (Hatton et al. 2012) ,
and functional coupling (Cauda et al. 2011). To build this more complete picture of interoceptive
processing, other sophisticated network-based and whole-brain analysis approaches will be
required, including those oriented towards the development of clinically-predictive models for
interoceptive symptoms in psychiatric disorders.
763

Conclusion

This study confirms the pivotal role of the dysgranular insula in the processing of top-down
and bottom-up interoceptive inputs, and validates the dysgranular mid-insula as a possible locus of
interoceptive disruption in psychiatric disorders.

Funding:

National Institute of Mental Health NIMH K23MH112949, R01MH127225 (to SSK) and National
Institute of General Medical Sciences NIGMS 1P20GM121312 (to SSK).
772
773

774 **Disclosures/competing interest:**

775 The authors have no financial or non-financial competing interests to disclose.
776
777

778 **Acknowledgements**:

779 The authors would like to thank Maria Puhl and Gabriel Morrison for their contributions to data
780 preprocessing, as well as Valerie Upshaw and Rachel Lapidus for their assistance with data
781 collection.
782
783

784 **Figures and Tables**
785
Table 1. Demographic information for the transdiagnostic anxiety, depression, and/or eating
disorders (ADE) group and the propensity-matched healthy comparison (HC) group. GAD =
generalized anxiety disorder, MDD = major depressive disorder, AN = anorexia nervosa, PTSD =
post-traumatic stress disorder, OCD = obsessive compulsive disorder, SUD = substance use
disorder, BMI = body mass index, STAI = State-Trait Anxiety Inventory, ASI = Anxiety Sensitivity
Index, with a Total Score, and Physical, Cognitive, and Social subscales, PHQ-9 = depression module
of the Patient Health Questionnaire, SCOFF = "Sick, Control, One, Fat, Food" eating disorder
questionnaire.

	HC (n=46)	ADE (n=46)	t	df	p-value
Sex	41 Female 5 Male	45 Female 1 Male	-	-	-
Age	24 (5)	25 (7)	0.57	80.49	0.57
BMI	23.48 (3.26)	23.34 (3.33)	-0.20	89.97	0.84
STAI	30.0 (1.1)	56.0 (1.4)	-14.43	83.01	**<0.0001**
ASI-Total	7.39 (5.26)	13.98 (8.89)	8.89	57.47	**<0.0001**
ASI-Physical	1.15 (1.59)	5.19 (5.81)	5.81	53.38	**<0.0001**
ASI-Cognitive	0.98 (1.47)	7.41 (6.53)	6.52	49.54	**<0.0001**
ASI-Social	5.26 (4.04)	13.83 (6.08)	7.96	78.28	**<0.0001**
PHQ-9	0.78 (1.11)	10.39 (5.98	10.72	48.12	**<0.0001**
SCOFF	0 (0)	1.09 (1.23)	6.01	45	**<0.0001**
Diagnoses	No psychiatric disorders	67% GAD 72% MDD (current or remitted) 35% AN 13% Social anxiety disorder 11% PTSD 4% OCD 4% SUD	-	-	-
Medication use	None	*Taking one or more psychiatric medication:* Serotonergic[1]: 35% Histaminergic[2]: 11% Dopaminergic[3]: 7% SNRI[4]: 7% Benzodiazepene[5]: 4% Miscellaneous[6]: 2% Beta-blocker[7]: 2%	-	-	-

[1]*Escitalopram, Fluoxetine, Trazodone, Sertraline, Paroxetine, Duloxetine*
[2]*Lamotrigine, Hydroxyzine, Doxepin*

[3]*Olanzapine, Aripiprazole, Quetiapine*

[4]*Atomoxetine, Buproprion, Desvenlafaxine*

[5]*Clonazepam, Alprazolam*

[6]*Vortioxetine*

[7]*Propranolol*

807 **Table 2.** Insular regions demonstrating convergent processing between top-down and bottom-up
808 interception**.**

	MNI coordinates (center of mass)			Volume (mm³)
	X	*Y*	*Z*	
Left Insula: ADE	-39	-0.1	-8.2	1736 (217 voxels)
Left Insula: HC	-40	-1.3	-4.8	1240 (155 voxels)
Right Insula: ADE	39	2.3	-12.9	1088 (136 voxels)
Right Insula: HC	39.9	2.2	-10.3	1960 (245 voxels)

812 **Table 3.** Agranular anterior insular regions demonstrating divergent processing during periods of
813 increased bodily expectancies.

		MNI coordinates (center of mass)			Volume (mm³)
		X	*Y*	*Z*	
Anticipation	**Left: ADE**	-31.6 / -31.9	17.9 / 9.2	4.8 / -15	1168 (146 voxels) / 344 (43 voxels)
	Left: HC	-31.5	15.2	-1.2	1688 (211 voxels)
	Right: ADE	35.4	19.1	1.4	1648 (206 voxels)
	Right: HC	37 / 35	13 / 7	-9 / -19	1744 (218 voxels) / 368 (46 voxels)
Saline (Peak)	**Left: ADE**	-33.1	16.7	-1.6	2264 (283 voxels)
	Left: HC	-33.2	15.7	-2.7	2760 (345 voxels)
	Right: ADE	35.4	18.6	-0.9	2120 (265 voxels)
	Right: HC	34.3	16.7	-4.7	3024 (378 voxels)

Figure 1. Experimental design. During a single functional neuroimaging scanning session two
neuroimaging tasks were used to localize the convergence between bottom-up and top-down
interoceptive processing. The Isoproterenol Infusion (ISO) task manipulated interoceptive input in
a bottom-up manner, while the Visceral Interoceptive Attention (VIA) task manipulated
interoceptive attention in a top-down manner. During the ISO task, participants were asked to
attend to their cardiorespiratory sensations while receiving double-blinded infusions of either
isoproterenol 2.0 micrograms (mcg, solid line) or 0.5mcg (not pictured), a rapidly acting peripheral
beta-adrenergic analogue eliciting transiently increased cardiovascular and respiratory signals in a
manner akin to adrenaline, or infusions of saline (dashed line), resulting in no physiological change.
During the VIA task, on-screen cues directed participants to shift their attention towards naturally
occurring body sensations from a particular internal organ (heart, lungs, or stomach, the
interoceptive attention conditions) or the word "TARGET" (the exteroceptive attention condition)
that would flash at different intensities. No infusions were given during this task, so the body
remained at physiological rest. Both tasks were preceded and followed by a resting state scan.
832

833
Figure 2. Convergence between bottom-up and top-down interoceptive processing in healthy
comparisons (HC, top row) and individuals with anxiety, depression, and/or eating disorders (ADE,
bottom row). Each convergence map reflects the only cluster of voxels that were co-activated
across the whole brain during the ISO task, involving perturbation of cardiorespiratory signals, and
the VIA task, involving goal-directed interoceptive attention towards cardiac and respiratory
signals at rest. Numbers on the left and right reflect the Dice similarity coefficient, which are used to
quantify the degree of spatial overlap between groups. Relative to HCs, the ADE group showed
lower spatial similarity for the left insular cortex (Dice coefficient of 0.58), than the right insular
cortex (Dice coefficient of 0.78).
843
844

845
Figure 3. Hemispheric divergence of bottom-up and top-down interoceptive processing across
psychiatric disorders. (A) Hemisphere-specific convergence maps (white outlines) overlaid on a
tripartite probabilistic cytoarchitectonic division of the insular cortex. (B) Quantification of
convergence within each cytoarchitectonic subregion (*i.e.,* number of co-activated voxels in relation
to the total number of voxels within that subregion), showing asymmetric proportional voxel co-
activation in the left and right dysgranular insula between the groups. *indicates significant group
difference for the proportion of co-activated voxels in each subregion via chi-square test. HC:
healthy comparison. ADE: anxiety, depression, and/or eating disorder.
854

855
Figure 4. Hemispheric differences in the activation magnitude of the convergent dysgranular insula
subregion. (A) Co-activated voxels within the right dysgranular insula subregion exhibited a greater
degree of percent signal change during the Peak period of the 2.0mcg dose in the ISO task (left sub
panel) and during the Heart and Lung attention condition in the VIA task (right sub panel). As there
were no main effects of group, mean percent signal change is collapsed across groups. *indicates
p<0.05 for the left versus right post-hoc contrast, following a main effect of hemisphere in the linear
mixed effects regression. (B) Across both groups the mean percent signal change within the right
but not left convergent dysgranular insula subregion was correlated with real-time
cardiorespiratory intensity ratings during the Peak period of the 2.0mcg dose in the ISO task.
*indicates p<0.05 for the Pearson's R correlation coefficient, separately for each hemisphere.
866
867

868
Figure 5. Changes in whole-brain functional connectivity of convergent dysgranular insula
subregions following performance of the interoceptive tasks. (A) Functional connectivity between
the right convergent dysgranular insula subregion and the left middle frontal gyrus showed
opposing effects in the ADE versus HC groups. (B) While functional connectivity between these
regions increased from baseline following interoception in the ADE group, it decreased in the HC
group, resulting in a significant interaction effect. (C) Across the entire sample changes in functional
connectivity were associated with trait measures of anxiety and depression. When examined in
each group individually, associations between connectivity change and trait anxiety occurred in
opposite directions (left and middle panel).

Figure 6. Divergence of top-down processing during interoceptive uncertainty in individuals with anxiety, depression, and/or eating disorders (ADE) and healthy comparisons (HC). (A) Selective activation of the bilateral agranular anterior insula during the Anticipation period (top) and Saline Peak period (bottom) of the ISO task. Both epochs demarcate periods of maximal expectancy about future changes in body state while the body remains at physiological rest (i.e., no ensuring evidence of heart rate or breathing rate increases). These brain areas showed activity only during these

periods of the ISO task and were not active during the Heart and Lung attention part of the VIA task,
indicated they have specialized roles. (B) Quantifying the spatial extent of this agranular activation
(*i.e.,* number of active voxels in relation to total number of voxels in that subregion) revealed that
greater right hemisphere activation during the anticipation window occurred for both groups, and
more so in the HC than the ADE group. During the peak period of Saline this activation covered
more of the bilateral agranular insula in both groups. However, the right hemisphere was activated
more than the left in both groups, and the HC group exhibited more active voxels in the right
hemisphere than the left, and more than the ADE group in both hemispheres, while the ADE group
showed no hemispheric difference during this window. (C) Increased ISO-specific activation of
agranular insular during the anticipation period versus the saline peak period. This pattern,
occurring for both hemispheres and across both groups, was greater in the right than the left
hemisphere.
897

Figure 7. Time courses of heart rate (A) and continuous dial ratings (B) during the 2.0mcg (solid
lines) and Saline (dotted lines) doses of the ISO task. Heart rate changes observed during both
infusions were generally similar between both groups across the majority of the infusion period.

The cluster-based permutation analysis showed a two-second instance of significantly higher heart rate in the ADE group starting at the 98th second of the 2.0mcg dose only during the Peak period, followed by a prolonged increased in dial rating for large portions of the Peak and Recovery periods (seconds 110 through 142, shown via the shaded area). No heart rate differences were observed during the Saline infusion. *indicates p<0.05 for significant group difference using permutation analysis. (C) Retrospective ratings on the ISO task showed no significant group differences in cardiac or respiratory intensity or excitement following saline or the 2.0mcg infusion, while the ADE group reported increased anxiety following both infusions. (D) Sensation ratings during the VIA task. The ADE group reported higher cardiac intensity ratings than the HC group but lower exteroceptive ratings, with no group difference in respiratory or gastric sensation ratings. *indicates p<0.05 for two-sample t-test between groups, separately for each condition. Where appropriate, both the p-value and Cohen's d are shown.

918 **References**

919
920
Adamic, Emily, Jason Avery, Sahib S. Khalsa, and Adam Robert Teed. 2021. "Convergence between Goal-
Directed and Stimulus-Driven Streams of Cardiorespiratory Interoception." doi:
10.17605/OSF.IO/6NXA3.

Adolfi, Federico, Blas Couto, Fabian Richter, Jean Decety, Jessica Lopez, Mariano Sigman, Facundo
Manes, and Agustín Ibáñez. 2017. "Convergence of Interoception, Emotion, and Social
Cognition: A Twofold FMRI Meta-Analysis and Lesion Approach." *Cortex* 88:124–42. doi:
10.1016/j.cortex.2016.12.019.

Allen, Micah, Andrew Levy, Thomas Parr, and Karl J. Friston. 2022. "In the Body's Eye: The
Computational Anatomy of Interoceptive Inference." *PLoS Computational Biology*
18(9):e1010490. doi: 10.1371/journal.pcbi.1010490.

Augustine, J. R. 1996. "Circuitry and Functional Aspects of the Insular Lobe in Primates Including
Humans." *Brain Research. Brain Research Reviews* 22(3):229–44. doi: 10.1016/s0165-
0173(96)00011-2.

Avery, Jason A., Kaiping Burrows, Kara L. Kerr, Jerzy Bodurka, Sahib S. Khalsa, Martin P. Paulus, and W.
Kyle Simmons. 2017. "How the Brain Wants What the Body Needs: The Neural Basis of Positive
Alliesthesia." *Neuropsychopharmacology* 42(4):822–30. doi: 10.1038/npp.2016.128.

Avery, Jason A., Wayne C. Drevets, Scott E. Moseman, Jerzy Bodurka, Joel C. Barcalow, and W. Kyle
Simmons. 2014. "Major Depressive Disorder Is Associated with Abnormal Interoceptive Activity
and Functional Connectivity in the Insula." *Biological Psychiatry* 76(3):258–66. doi:
10.1016/j.biopsych.2013.11.027.

Barrett, Lisa Feldman, Karen S. Quigley, and Paul Hamilton. 2016. "An Active Inference Theory of
Allostasis and Interoception in Depression." *Philosophical Transactions of the Royal Society B:*
Biological Sciences 371(1708):20160011. doi: 10.1098/rstb.2016.0011.

Barrett, Lisa Feldman, and W. Kyle Simmons. 2015. "Interoceptive Predictions in the Brain." *Nature*
Reviews Neuroscience 16(7):419–29. doi: 10.1038/nrn3950.

Berntson, Gary G., and Sahib S. Khalsa. 2021. "Neural Circuits of Interoception." *Trends in Neurosciences*
44(1):17–28. doi: 10.1016/j.tins.2020.09.011.

Birba, Agustina, Hernando Santamaría-García, Pavel Prado, Josefina Cruzat, Agustín Sainz Ballesteros,
Agustina Legaz, Sol Fittipaldi, Claudia Duran-Aniotz, Andrea Slachevsky, Rodrigo Santibañez,
Mariano Sigman, Adolfo M. García, Robert Whelan, Sebastián Moguilner, and Agustín Ibáñez.
2022. "Allostatic-Interoceptive Overload in Frontotemporal Dementia." *Biological Psychiatry*
92(1):54–67. doi: 10.1016/j.biopsych.2022.02.955.

Birn, Rasmus M., Monica A. Smith, Tyler B. Jones, and Peter A. Bandettini. 2008. "The Respiration
Response Function: The Temporal Dynamics of FMRI Signal Fluctuations Related to Changes in
Respiration." *NeuroImage* 40(2):644–54. doi: 10.1016/j.neuroimage.2007.11.059.

Bonaz, Bruno, Richard D. Lane, Michael L. Oshinsky, Paul J. Kenny, Rajita Sinha, Emeran A. Mayer, and
Hugo D. Critchley. 2021. "Diseases, Disorders, and Comorbidities of Interoception." *Trends in*
Neurosciences 44(1):39–51. doi: 10.1016/j.tins.2020.09.009.

Bowring, Alexander, Camille Maumet, and Thomas E. Nichols. 2019. "Exploring the Impact of Analysis
Software on Task FMRI Results." *Human Brain Mapping* 40(11):3362–84. doi:
10.1002/hbm.24603.

Brefczynski, Julie A., and Edgar A. DeYoe. 1999. "A Physiological Correlate of the 'spotlight' of Visual
Attention." *Nature Neuroscience* 2(4):370–74. doi: 10.1038/7280.

Burrows, Kaiping, Danielle C. DeVille, Kelly T. Cosgrove, Rayus T. Kuplicki, Tulsa 1000 Investigators,
Martin P. Paulus, Robin Aupperle, Sahib S. Khalsa, and Jennifer L. Stewart. 2022. "Impact of
Serotonergic Medication on Interoception in Major Depressive Disorder." *Biological Psychology*
169:108286. doi: 10.1016/j.biopsycho.2022.108286.

Campbell-Sills, Laura, Sonya B. Norman, Michelle G. Craske, Greer Sullivan, Ariel J. Lang, Denise A.
Chavira, Alexander Bystritsky, Cathy Sherbourne, Peter Roy-Byrne, and Murray B. Stein. 2009.
"Validation of a Brief Measure of Anxiety-Related Severity and Impairment: The Overall Anxiety
Severity and Impairment Scale (OASIS)." *Journal of Affective Disorders* 112(1–3):92–101. doi:
10.1016/j.jad.2008.03.014.

Cannon, W. B. 1939. *The Wisdom of the Body, 2nd Ed*. Oxford, England: Norton & Co.

Carvalho, Gil B., and Antonio Damasio. 2021. "Interoception and the Origin of Feelings: A New
Synthesis." *BioEssays: News and Reviews in Molecular, Cellular and Developmental Biology*
43(6):e2000261. doi: 10.1002/bies.202000261.

Cauda, Franco, Federico D'Agata, Katiuscia Sacco, Sergio Duca, Giuliano Geminiani, and Alessandro
Vercelli. 2011. "Functional Connectivity of the Insula in the Resting Brain." *NeuroImage* 55(1):8–
23. doi: 10.1016/j.neuroimage.2010.11.049.

Clark, Andy. 2013. "Whatever next? Predictive Brains, Situated Agents, and the Future of Cognitive
Science." *The Behavioral and Brain Sciences* 36(3):181–204. doi: 10.1017/S0140525X12000477.

Clark, Andy. 2016. "Attention Alters Predictive Processing." *The Behavioral and Brain Sciences* 39:e234.
doi: 10.1017/S0140525X15002472.

Corbetta, Maurizio, Gaurav Patel, and Gordon L. Shulman. 2008. "The Reorienting System of the Human
Brain: From Environment to Theory of Mind." *Neuron* 58(3):306–24. doi:
10.1016/j.neuron.2008.04.017.

Cox, R. W. 1996. "AFNI: Software for Analysis and Visualization of Functional Magnetic Resonance
Neuroimages." *Computers and Biomedical Research, an International Journal* 29(3):162–73. doi:
10.1006/cbmr.1996.0014.

Cox, Robert W. 2019. "Equitable Thresholding and Clustering (ETAC): A Novel Method for FMRI
Clustering in AFNI." *BioRxiv* 295931. doi: 10.1101/295931.

Craig, A. D. 2002. "How Do You Feel? Interoception: The Sense of the Physiological Condition of the
Body." *Nature Reviews. Neuroscience* 3(8):655–66. doi: 10.1038/nrn894.

Craig, A. D. (Bud). 2005. "Forebrain Emotional Asymmetry: A Neuroanatomical Basis?" *Trends in
Cognitive Sciences* 9(12):566–71. doi: 10.1016/j.tics.2005.10.005.

Craig, A. D. Bud. 2009. "How Do You Feel--Now? The Anterior Insula and Human Awareness." *Nature
Reviews. Neuroscience* 10(1):59–70. doi: 10.1038/nrn2555.

Critchley, Hugo D., and Sarah N. Garfinkel. 2017. "Interoception and Emotion." *Current Opinion in
Psychology* 17:7–14. doi: 10.1016/j.copsyc.2017.04.020.

Critchley, Hugo D., Christopher J. Mathias, and Raymond J. Dolan. 2002. "Fear Conditioning in Humans:
The Influence of Awareness and Autonomic Arousal on Functional Neuroanatomy." *Neuron*
33(4):653–63. doi: 10.1016/s0896-6273(02)00588-3.

Davidson, Richard J., William Irwin, Richard J. Davidson, William Irwin, Richard J. Davidson, and William
Irwin. 1999. "The Functional Neuroanatomy of Emotion and Affective Style." *Trends in Cognitive
Sciences* 3(1):11–21. doi: 10.1016/S1364-6613(98)01265-0.

Davidson, Richard J., Alexander J. Shackman, and Jeffrey S. Maxwell. 2004. "Asymmetries in Face and
Brain Related to Emotion." *Trends in Cognitive Sciences* 8(9):389–91. doi:
10.1016/j.tics.2004.07.006.

De la Cruz, Feliberto, Adam R. Teed, Rachel C. Lapidus, Valerie Upshaw, Andy Schumann, Martin P.
Paulus, Karl-Jürgen Bär, and Sahib S. Khalsa. 2023. "Central Autonomic Network Alterations in
Anorexia Nervosa Following Peripheral Adrenergic Stimulation." *Biological Psychiatry. Cognitive
Neuroscience and Neuroimaging* 8(7):720–30. doi: 10.1016/j.bpsc.2022.12.009.

Drane, Daniel L., Negar Fani, Mark Hallett, Sahib S. Khalsa, David L. Perez, and Nicole A. Roberts. 2020.
"A Framework for Understanding the Pathophysiology of Functional Neurological Disorder." *CNS
Spectrums* 1–7. doi: 10.1017/S1092852920001789.

Dum, Richard P., David J. Levinthal, and Peter L. Strick. 2009. "The Spinothalamic System Targets Motor
and Sensory Areas in the Cerebral Cortex of Monkeys." *Journal of Neuroscience* 29(45):14223–
35. doi: 10.1523/JNEUROSCI.3398-09.2009.

Evrard, Henry C. 2019. "The Organization of the Primate Insular Cortex." *Frontiers in Neuroanatomy*
13:43. doi: 10.3389/fnana.2019.00043.

Fan, Lingzhong, Hai Li, Junjie Zhuo, Yu Zhang, Jiaojian Wang, Liangfu Chen, Zhengyi Yang, Congying Chu,
Sangma Xie, Angela R. Laird, Peter T. Fox, Simon B. Eickhoff, Chunshui Yu, and Tianzi Jiang. 2016.
"The Human Brainnetome Atlas: A New Brain Atlas Based on Connectional Architecture."
Cerebral Cortex 26(8):3508–26. doi: 10.1093/cercor/bhw157.

Fermin, Alan S. R., Karl Friston, and Shigeto Yamawaki. n.d. "An Insula Hierarchical Network Architecture
for Active Interoceptive Inference." *Royal Society Open Science* 9(6):220226. doi:
10.1098/rsos.220226.

Gehrlach, Daniel A., Caroline Weiand, Thomas N. Gaitanos, Eunjae Cho, Alexandra S. Klein, Alexandru A.
Hennrich, Karl-Klaus Conzelmann, and Nadine Gogolla. 2020. "A Whole-Brain Connectivity Map
of Mouse Insular Cortex." *ELife* 9:e55585. doi: 10.7554/eLife.55585.

Ghaziri, Jimmy, Alan Tucholka, Gabriel Girard, Olivier Boucher, Jean-Christophe Houde, Maxime
Descoteaux, Sami Obaid, Guillaume Gilbert, Isabelle Rouleau, and Dang Khoa Nguyen. 2018.
"Subcortical Structural Connectivity of Insular Subregions." *Scientific Reports* 8(1):8596. doi:
10.1038/s41598-018-26995-0.

Ghaziri, Jimmy, Alan Tucholka, Gabriel Girard, Jean-Christophe Houde, Olivier Boucher, Guillaume
Gilbert, Maxime Descoteaux, Sarah Lippé, Pierre Rainville, and Dang Khoa Nguyen. 2017. "The
Corticocortical Structural Connectivity of the Human Insula." *Cerebral Cortex* 27(2):1216–28. doi:
10.1093/cercor/bhv308.

Glover, Gary H., Tie-Qiang Li, and David Ress. 2000. "Image-Based Method for Retrospective Correction
of Physiological Motion Effects in FMRI: RETROICOR." *Magnetic Resonance in Medicine*
44(1):162–67. doi: 10.1002/1522-2594(200007)44:1<162::AID-MRM23>3.0.CO;2-E.

Gogolla, Nadine. 2017. "The Insular Cortex." *Current Biology: CB* 27(12):R580–86. doi:
10.1016/j.cub.2017.05.010.

Goldstein-Piekarski, Andrea N., Tali M. Ball, Zoe Samara, Brooke R. Staveland, Arielle S. Keller, Scott L.
Fleming, Katherine A. Grisanzio, Bailey Holt-Gosselin, Patrick Stetz, Jun Ma, and Leanne M.
Williams. 2022. "Mapping Neural Circuit Biotypes to Symptoms and Behavioral Dimensions of
Depression and Anxiety." *Biological Psychiatry* 91(6):561–71. doi:
10.1016/j.biopsych.2021.06.024.

Halani, Priyanka Kadam, Uduak U. Andy, Hengyi Rao, and Lily A. Arya. 2020. "Regions of the Brain
Activated in Bladder Filling vs Rectal Distention in Healthy Adults: A Meta-Analysis of
Neuroimaging Studies." *Neurourology and Urodynamics* 39(1):58–65. doi: 10.1002/nau.24221.

Harrison, Olivia K., Laura Köchli, Stephanie Marino, Roger Luechinger, Franciszek Hennel, Katja Brand,
Alexander J. Hess, Stefan Frässle, Sandra Iglesias, Fabien Vinckier, Frederike H. Petzschner,
Samuel J. Harrison, and Klaas E. Stephan. 2021. "Interoception of Breathing and Its Relationship
with Anxiety." *Neuron* 109(24):4080-4093.e8. doi: 10.1016/j.neuron.2021.09.045.

Hassanpour, Mahlega S., W. Kyle Simmons, Justin S. Feinstein, Qingfei Luo, Rachel C. Lapidus, Jerzy
Bodurka, Martin P. Paulus, and Sahib S. Khalsa. 2018. "The Insular Cortex Dynamically Maps
Changes in Cardiorespiratory Interoception." *Neuropsychopharmacology* 43(2):426–34. doi:
10.1038/npp.2017.154.

Hassanpour, Mahlega S., Lirong Yan, Danny J. J. Wang, Rachel C. Lapidus, Armen C. Arevian, W. Kyle
Simmons, Jamie D. Feusner, and Sahib S. Khalsa. 2016. "How the Heart Speaks to the Brain:
Neural Activity during Cardiorespiratory Interoceptive Stimulation." *Philosophical Transactions
of the Royal Society B: Biological Sciences* 371(1708):20160017. doi: 10.1098/rstb.2016.0017.

Hatton, Sean N., Jim Lagopoulos, Daniel F. Hermens, Sharon L. Naismith, Maxwell R. Bennett, and Ian B.
Hickie. 2012. "Correlating Anterior Insula Gray Matter Volume Changes in Young People with

Clinical and Neurocognitive Outcomes: An MRI Study." *BMC Psychiatry* 12(1):45. doi:
10.1186/1471-244X-12-45.

Henseler, Ilona, Sebastian Krüger, Peter Dechent, and Oliver Gruber. 2011. "A Gateway System in Rostral
PFC? Evidence from Biasing Attention to Perceptual Information and Internal Representations."
NeuroImage 56(3):1666–76. doi: 10.1016/j.neuroimage.2011.02.056.

Hommel, Bernhard, Craig S. Chapman, Paul Cisek, Heather F. Neyedli, Joo-Hyun Song, and Timothy N.
Welsh. 2019. "No One Knows What Attention Is." *Attention, Perception, & Psychophysics*
81(7):2288–2303. doi: 10.3758/s13414-019-01846-w.

Jänig, Wilfrid. 2006. *Integrative Action of the Autonomic Nervous System: Neurobiology of Homeostasis*.
Cambridge: Cambridge University Press.

Jänig, Wilfrid, Janet R. Keast, Elspeth M. McLachlan, Winfried L. Neuhuber, and Michelle Southard-
Smith. 2017. "Renaming All Spinal Autonomic Outflows as Sympathetic Is a Mistake." *Autonomic*
Neuroscience: Basic & Clinical 206:60–62. doi: 10.1016/j.autneu.2017.04.003.

Jo, Hang Joon, Ziad S. Saad, W. Kyle Simmons, Lydia A. Milbury, and Robert W. Cox. 2010. "Mapping
Sources of Correlation in Resting State FMRI, with Artifact Detection and Removal." *NeuroImage*
52(2):571–82. doi: 10.1016/j.neuroimage.2010.04.246.

Kasper, Lars, Maria Engel, Jakob Heinzle, Matthias Mueller-Schrader, Nadine N. Graedel, Jonas Reber,
Thomas Schmid, Christoph Barmet, Bertram J. Wilm, Klaas Enno Stephan, and Klaas P.
Pruessmann. 2022. "Advances in Spiral FMRI: A High-Resolution Study with Single-Shot
Acquisition." *NeuroImage* 246:118738. doi: 10.1016/j.neuroimage.2021.118738.

Kerr, Kara L., Scott E. Moseman, Jason A. Avery, Jerzy Bodurka, Nancy L. Zucker, and W. Kyle Simmons.
2016. "Altered Insula Activity during Visceral Interoception in Weight-Restored Patients with
Anorexia Nervosa." *Neuropsychopharmacology* 41(2):521–28. doi: 10.1038/npp.2015.174.

Khalsa, Sahib S., Ralph Adolphs, Oliver G. Cameron, Hugo D. Critchley, Paul W. Davenport, Justin S.
Feinstein, Jamie D. Feusner, Sarah N. Garfinkel, Richard D. Lane, Wolf E. Mehling, Alicia E.
Meuret, Charles B. Nemeroff, Stephen Oppenheimer, Frederike H. Petzschner, Olga Pollatos,
Jamie L. Rhudy, Lawrence P. Schramm, W. Kyle Simmons, Murray B. Stein, Klaas E. Stephan,
Omer Van den Bergh, Ilse Van Diest, Andreas von Leupoldt, Martin P. Paulus, Vivien Ainley,
Obada Al Zoubi, Robin Aupperle, Jason Avery, Leslie Baxter, Christoph Benke, Laura Berner, Jerzy
Bodurka, Eric Breese, Tiffany Brown, Kaiping Burrows, Yoon-Hee Cha, Ashley Clausen, Kelly
Cosgrove, Danielle Deville, Laramie Duncan, Patrice Duquette, Hamed Ekhtiari, Thomas Fine,
Bart Ford, Indira Garcia Cordero, Diamond Gleghorn, Yvette Guereca, Neil A. Harrison, Mahlega
Hassanpour, Tanja Hechler, Aaron Heller, Natalie Hellman, Beate Herbert, Behnaz Jarrahi, Kara
Kerr, Namik Kirlic, Megan Klabunde, Thomas Kraynak, Michael Kriegsman, Juliet Kroll, Rayus
Kuplicki, Rachel Lapidus, Trang Le, Kyle Logie Hagen, Ahmad Mayeli, Amanda Morris, Nasir
Naqvi, Kristina Oldroyd, Christiane Pané-Farré, Raquel Phillips, Tasha Poppa, Willliam Potter,
Maria Puhl, Adam Safron, Margaret Sala, Jonathan Savitz, Heather Saxon, Will Schoenhals, Colin
Stanwell-Smith, Adam Teed, Yuri Terasawa, Katie Thompson, Marisa Toups, Satoshi Umeda,
Valerie Upshaw, Teresa Victor, Christina Wierenga, Colleen Wohlrab, Hung-wen Yeh, Adrian
Yoris, Fadel Zeidan, Vadim Zotev, and Nancy Zucker. 2018. "Interoception and Mental Health: A

Roadmap." *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging* 3(6):501–13. doi:
10.1016/j.bpsc.2017.12.004.

Khalsa, Sahib S., and Rachel C. Lapidus. 2016. "Can Interoception Improve the Pragmatic Search for
Biomarkers in Psychiatry?" *Frontiers in Psychiatry* 7. doi: 10.3389/fpsyt.2016.00121.

Khalsa, Sahib S., David Rudrauf, Justin S. Feinstein, and Daniel Tranel. 2009. "The Pathways of
Interoceptive Awareness." *Nature Neuroscience* 12(12):1494–96. doi: 10.1038/nn.2411.

Kimmerly, Derek S., Deborah D. O'Leary, Ravi S. Menon, Joseph S. Gati, Kevin Shoemaker. 2005. "Cortical
regions associated with autonomic cardiovascular regulation during lower body negative
pressure in humans." *Journal of Physiology* 569(1):331-345.
https://doi.org/10.1113/jphysiol.2005.091637

Knudsen, Lasse, Christopher J. Bailey, Jakob U. Blicher, Yan Yang, Peng Zhang, and Torben E. Lund. 2023.
"Improved Sensitivity and Microvascular Weighting of 3T Laminar FMRI with GE-BOLD Using
NORDIC and Phase Regression." *NeuroImage* 271:120011. doi:
10.1016/j.neuroimage.2023.120011.

Kroenke, K., R. L. Spitzer, and J. B. Williams. 2001. "The PHQ-9: Validity of a Brief Depression Severity
Measure." *Journal of General Internal Medicine* 16(9):606–13. doi: 10.1046/j.1525-
1497.2001.016009606.x.

Kutz, Amanda M., Alison G. Marsh, Craig G. Gunderson, Shira Maguen, and Robin M. Masheb. 2020.
"Eating Disorder Screening: A Systematic Review and Meta-Analysis of Diagnostic Test
Characteristics of the SCOFF." *Journal of General Internal Medicine* 35(3):885–93. doi:
10.1007/s11606-019-05478-6.

Mayeli, Ahmad, Obada Al Zoubi, Evan J. White, Sheridan Chappelle, Rayus Kuplicki, Alexa Morton,
Jaimee Bruce, Ryan Smith, Justin S. Feinstein, Jerzy Bodurka, Martin P. Paulus, and Sahib S.
Khalsa. 2023. "Parieto-Occipital ERP Indicators of Gut Mechanosensation in Humans." *Nature
Communications* 14(1):3398. doi: 10.1038/s41467-023-39058-4.

Mechelli, Andrea, Cathy J. Price, Karl J. Friston, and Alumit Ishai. 2004. "Where Bottom-up Meets Top-
down: Neuronal Interactions during Perception and Imagery." *Cerebral Cortex (New York, N.Y.:
1991)* 14(11):1256–65. doi: 10.1093/cercor/bhh087.

Mufson, Elliott J., and M.-Marsel Mesulam. 1982. "Insula of the Old World Monkey. II: Afferent Cortical
Input and Comments on the Claustrum." *Journal of Comparative Neurology* 212(1):23–37. doi:
https://doi.org/10.1002/cne.902120103.

Murphy, Fionnuala C., Ian Nimmo-Smith, and Andrew D. Lawrence. 2003. "Functional Neuroanatomy of
Emotions: A Meta-Analysis." *Cognitive, Affective, & Behavioral Neuroscience* 3(3):207–33. doi:
10.3758/CABN.3.3.207.

Nord, Camilla L., Rebecca P. Lawson, and Tim Dalgleish. 2021. "Disrupted Dorsal Mid-Insula Activation
During Interoception Across Psychiatric Disorders." *The American Journal of Psychiatry*
178(8):761–70. doi: 10.1176/appi.ajp.2020.20091340.

Oppenheimer, S. M., A. Gelb, J. P. Girvin, and V. C. Hachinski. 1992. "Cardiovascular Effects of Human
Insular Cortex Stimulation." *Neurology* 42(9):1727–32. doi: 10.1212/wnl.42.9.1727.

Oppenheimer, Stephen M., Adrian Gelb, John P. Girvin, and Vladimir C. Hachinski. 1992. "Cardiovascular
Effects of Human Insular Cortex Stimulation." *Neurology* 42(9):1727–1727. doi:
10.1212/WNL.42.9.1727.

Pace-Schott, Edward F., Marlissa C. Amole, Tatjana Aue, Michela Balconi, Lauren M. Bylsma, Hugo
Critchley, Heath A. Demaree, Bruce H. Friedman, Anne Elizabeth Kotynski Gooding, Olivia
Gosseries, Tanja Jovanovic, Lauren A. J. Kirby, Kasia Kozlowska, Steven Laureys, Leroy Lowe,
Kelsey Magee, Marie-France Marin, Amanda R. Merner, Jennifer L. Robinson, Robert C. Smith,
Derek P. Spangler, Mark Van Overveld, and Michael B. VanElzakker. 2019. "Physiological
Feelings." *Neuroscience and Biobehavioral Reviews* 103:267–304. doi:
10.1016/j.neubiorev.2019.05.002.

Paulus, Martin P., Justin S. Feinstein, and Sahib S. Khalsa. 2019. "An Active Inference Approach to
Interoceptive Psychopathology." *Annual Review of Clinical Psychology* 15:97–122. doi:
10.1146/annurev-clinpsy-050718-095617.

Paulus, Martin P., and Sahib S. Khalsa. 2021. "When You Don't Feel Right Inside: Homeostatic
Dysregulation and the Mid-Insular Cortex in Psychiatric Disorders." *The American Journal of
Psychiatry* 178(8):683–85. doi: 10.1176/appi.ajp.2021.21060622.

Petzschner, Frederike H., Sarah N. Garfinkel, Martin P. Paulus, Christof Koch, and Sahib S. Khalsa. 2021.
"Computational Models of Interoception and Body Regulation." *Trends in Neurosciences*
44(1):63–76. doi: 10.1016/j.tins.2020.09.012.

Pezzulo, Giovanni, Laura Barca, and Karl J. Friston. 2015. "Active Inference and Cognitive-Emotional
Interactions in the Brain." *Behavioral and Brain Sciences* 38:e85. doi:
10.1017/S0140525X14001009.

Picard, Rosalind W., Szymon Fedor, and Yadid Ayzenberg. 2016. "Multiple Arousal Theory and Daily-Life
Electrodermal Activity Asymmetry." *Emotion Review* 8(1):62–75. doi:
10.1177/1754073914565517.

Pohl, R., V. K. Yeragani, R. Balon, J. M. Rainey, H. Lycaki, A. Ortiz, R. Berchou, and P. Weinberg. 1988.
"Isoproterenol-Induced Panic Attacks." *Biological Psychiatry* 24(8):891–902. doi: 10.1016/0006-
3223(88)90224-7.

Pollatos, Olga, Rainer Schandry, Dorothee P. Auer, and Christian Kaufmann. 2007. "Brain Structures
Mediating Cardiovascular Arousal and Interoceptive Awareness." *Brain Research* 1141:178–87.
doi: 10.1016/j.brainres.2007.01.026.

Rauss, Karsten, and Gilles Pourtois. 2013. "What Is Bottom-Up and What Is Top-Down in Predictive
Coding?" *Frontiers in Psychology* 4:276. doi: 10.3389/fpsyg.2013.00276.

Schulz, Stefan M. 2016. "Neural Correlates of Heart-Focused Interoception: A Functional Magnetic
Resonance Imaging Meta-Analysis." *Philosophical Transactions of the Royal Society B: Biological
Sciences* 371(1708):20160018. doi: 10.1098/rstb.2016.0018.

Seth, Anil K., and Karl J. Friston. 2016. "Active Interoceptive Inference and the Emotional Brain."
Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences
371(1708):20160007. doi: 10.1098/rstb.2016.0007.

Sheehan, D. V., Y. Lecrubier, K. H. Sheehan, P. Amorim, J. Janavs, E. Weiller, T. Hergueta, R. Baker, and G.
C. Dunbar. 1998. "The Mini-International Neuropsychiatric Interview (M.I.N.I.): The
Development and Validation of a Structured Diagnostic Psychiatric Interview for DSM-IV and
ICD-10." *The Journal of Clinical Psychiatry* 59 Suppl 20:22-33;quiz 34-57.

Shipley, Michael T., and Marc S. Sanders. 1982. "Special Senses Are Really Special: Evidence for a
Reciprocal, Bilateral Pathway between Insular Cortex and Nucleus Parabrachialis." *Brain*
Research Bulletin 8(5):493–501. doi: 10.1016/0361-9230(82)90007-7.

Simmons, W. Kyle, Jason A. Avery, Joel C. Barcalow, Jerzy Bodurka, Wayne C. Drevets, and Patrick
Bellgowan. 2013. "Keeping the Body in Mind: Insula Functional Organization and Functional
Connectivity Integrate Interoceptive, Exteroceptive, and Emotional Awareness." *Human Brain*
Mapping 34(11):2944–58. doi: 10.1002/hbm.22113.

Smith, Ryan, Rayus Kuplicki, Justin Feinstein, Katherine L. Forthman, Jennifer L. Stewart, Martin P.
Paulus, Tulsa 1000 Investigators, and Sahib S. Khalsa. 2020. "A Bayesian Computational Model
Reveals a Failure to Adapt Interoceptive Precision Estimates across Depression, Anxiety, Eating,
and Substance Use Disorders." *PLOS Computational Biology* 16(12):e1008484. doi:
10.1371/journal.pcbi.1008484.

Smith, Ryan, Julian F. Thayer, Sahib S. Khalsa, and Richard D. Lane. 2017. "The Hierarchical Basis of
Neurovisceral Integration." *Neuroscience & Biobehavioral Reviews* 75:274–96. doi:
10.1016/j.neubiorev.2017.02.003.

Song, Xiaopeng, Bhaswati Roy, Gregg C. Fonarow, Mary A. Woo, Rajesh Kumar. 2018. "Brain structural
changes associated with aberrant functional responses to the Valsalva maneuver in heart
failure. *Journal of Neuroscience Research* 96(9). https://doi.org/10.1002/jnr.24264

Spitzer, Robert L., Kurt Kroenke, Janet B. W. Williams, and Bernd Löwe. 2006. "A Brief Measure for
Assessing Generalized Anxiety Disorder: The GAD-7." *Archives of Internal Medicine*
166(10):1092–97. doi: 10.1001/archinte.166.10.1092.

Sterling, Peter. 2012. "Allostasis: A Model of Predictive Regulation." *Physiology & Behavior* 106(1):5–15.
doi: 10.1016/j.physbeh.2011.06.004.

Stewart, Jennifer L., Sahib S. Khalsa, Rayus Kuplicki, Maria Puhl, T1000 Investigators, and Martin P.
Paulus. 2020. "Interoceptive Attention in Opioid and Stimulant Use Disorder." *Addiction Biology*
25(6):e12831. doi: https://doi.org/10.1111/adb.12831.

Sylvester, Ching-Yune C., Tor D. Wager, Steven C. Lacey, Luis Hernandez, Thomas E. Nichols, Edward E.
Smith, and John Jonides. 2003. "Switching Attention and Resolving Interference: FMRI Measures
of Executive Functions." *Neuropsychologia* 41(3):357–70. doi: 10.1016/s0028-3932(02)00167-7.

Teed, Adam R., Justin S. Feinstein, Maria Puhl, Rachel C. Lapidus, Valerie Upshaw, Rayus T. Kuplicki, Jerzy
Bodurka, Olujimi A. Ajijola, Walter H. Kaye, Wesley K. Thompson, Martin P. Paulus, and Sahib S.

Khalsa. 2022. "Association of Generalized Anxiety Disorder With Autonomic Hypersensitivity and
Blunted Ventromedial Prefrontal Cortex Activity During Peripheral Adrenergic Stimulation: A
Randomized Clinical Trial." *JAMA Psychiatry* 79(4):323–32. doi:
10.1001/jamapsychiatry.2021.4225.

Uddin, Lucina Q., Joshua Kinnison, Luiz Pessoa, and Michael L. Anderson. 2014. "Beyond the Tripartite
Cognition–Emotion–Interoception Model of the Human Insular Cortex." *Journal of Cognitive*
Neuroscience 26(1):16–27. doi: 10.1162/jocn_a_00462.

White, Lauren K., Jennifer C. Britton, Stefanie Sequeira, Emily G. Ronkin, Gang Chen, Yair Bar-Haim,
Tomer Shechner, Monique Ernst, Nathan A. Fox, Ellen Leibenluft, and Daniel S. Pine. 2016.
"Behavioral and Neural Stability of Attention Bias to Threat in Healthy Adolescents."
NeuroImage 136:84–93. doi: 10.1016/j.neuroimage.2016.04.058.

Wiestler, Tobias, and Jörn Diedrichsen. 2013. "Skill Learning Strengthens Cortical Representations of
Motor Sequences" edited by J. C. Culham. *ELife* 2:e00801. doi: 10.7554/eLife.00801.

Wu, Yu, Changwan Chen, Ming Chen, Kai Qian, Xinyou Lv, Haiting Wang, Lifei Jiang, Lina Yu, Min Zhuo,
and Shuang Qiu. 2020. "The Anterior Insular Cortex Unilaterally Controls Feeding in Response to
Aversive Visceral Stimuli in Mice." *Nature Communications* 11(1):640. doi: 10.1038/s41467-020-
14281-5.

1236