Diversity-disease relationships in natural microscopic nematode communities

Robbert van Himbeeck[1], Jessica N. Sowa[2], Hala Tamim El Jarkass[3], Wenjia Wu[4, 5], Job Oude Vrielink[1], Joost A. G. Riksen[1], Aaron Reinke[3], Lisa van Sluijs[1, *]

[1]Laboratory of Nematology, Wageningen University, Wageningen, the Netherlands

[2]Department of Biology, West Chester University of Pennsylvania,

West Chester, United States of America

[3]Department of Molecular Genetics, University of Toronto, Toronto, Canada

[4]Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems & CAS

Engineering Laboratory for Vegetation Ecosystem Restoration on Islands and Coastal Zones, South

China Botanical Garden, Chinese Academy of Sciences, Guangzhou, China

[5]South China National Botanical Garden, Guangzhou, China

*Corresponding author

Robbert van Himbeeck; robbert.vanhimbeeck@wur.nl; 0000-0002-1253-219X

Jessica N. Sowa; jsowa@wcupa.edu; 0000-0002-0896-7932

Hala Tamim El Jarkass; hala.tamimeljarkass@mail.utoronto.ca; 0000-0003-0118-5978

Wenjia Wu; wuwenjia@scbg.ac.cn; 0000-0001-7519-7803

Job Oude Vrielink; joboudevrielink@hotmail.com; 0009-0007-8331-816X

Joost A. G. Riksen; joost.riksen@wur.nl

Aaron Reinke; aaron.reinke@utoronto.ca; 0000-0001-7612-5342

Lisa van Sluijs; lisa.vansluijs@wur.nl; 0000-0002-4692-0844; +31317482998

Keywords: biodiversity, dilution effect, *C. elegans*, parasites, community composition, Nanopore sequencing, metabarcoding, microsporidia, virus

Abstract

Biodiversity can affect parasite prevalence, a phenomenon widely studied in macroscopic organisms. However, data from microscopic communities is lacking, despite their essential role in ecosystem functioning and the unique experimental opportunities microscopic organisms offer. Here, we study diversity-disease effects in wild nematode communities by profiting from the molecular tools available in the well-studied model nematode *Caenorhabditis elegans*. Nanopore sequencing was used to characterize nematode community diversity and composition, whereas parasites were identified using nine distinct experimental assays based on fluorescent staining or fluorescent reporter strains. Our results indicate biotic stress is abundant in wild nematode communities. Moreover, in two assays, diversity-disease relations were observed: microsporidia and immune system activation were more often detected in relatively species-poor communities. Other assays, targeting different parasites, were without diversity-disease relations. Together, this study provides the first demonstration of diversity-disease effects in microbial communities and establishes the use of nematode communities as model systems for disease dilution.

Introduction

Ecosystem functioning depends on biodiversity and biodiversity is shaped by parasites [1–3]. Increased (host) biodiversity can theoretically both cause increased (i.e. amplification effect) or decreased (i.e. dilution effect) parasite prevalence [2,4,5]. Generally, amplification effects occur when the parasite becomes more prevalent due to an increased availability of suitable hosts. Dilution effects results from reduced parasite reproduction because parasites experience difficulties finding a high-quality host amongst all species present. In contrast, parasites that replicate in a low-biodiversity population spread easily the because of high chances that the next organism they encounter is also a highly suitable host. Host-parasite relationships have been primarily studied in macroscopic species (e.g. [6–8]), but despite the necessity of well-functioning microscopic communities for ecosystem stability [3], natural host-parasite interactions in microscopic communities are little understood.

Interpretation of diversity-disease-correlations has not always been straightforward, as they depend on complex interactions within the natural community. Diversity-disease effects can be obscured or enhanced by parasite encounters, susceptibility and transmission of the species in the community, the order, timeframe and magnitude in which species enter or disappear from the community, life history traits of the host and host range of the parasite [4,7–21]. Interestingly, microscropic species that are relatively easy to grow and manipulate would present excellent models for studying diversity-disease effects, in particular when information from laboratory experiments can be combined with field observations. A model system where complex natural observations can be paired to controlled experiments will provide new experimental opportunities and biological insight into diversity-disease relations of (micro)organisms.

Nematodes present one of the most diverse and abundant groups of microscopic organisms worldwide [22] and some species possess characteristics that make them excellent model microscopic organisms. Of all nematodes, the free-living bacterivorous nematode *Caenorhabditis elegans* is a key model organism in many biological fields, ranging from developmental biology to neuroscience [23–26]. Today's science not only focuses on the molecular and genetic understanding of this 1mm-sized organism in the laboratory, but considers its natural ecology too which led to the discovery of many naturally-infecting parasites [27]. Parasites infecting *C. elegans* and related bacterivorous species comprise opportunistic bacteria and obligatory parasites including oomycetes, microsporidia and viruses many of which cause deadly infections [27–32]. Recently, complete sequencing of the 18S gene using long-read technology of Nanopore was established as a technique to determine nematode community diversity and composition in large-scale experiments [33]. This enables high-resolution nematode community characterization, and thereby provides a new opportunity to study diversity-disease interaction in natural communities of the model nematode *C. elegans* [33].

In this study, we initially assess the (a)biotic environment where *C. elegans* was detected and thereafter, parasite presence was linked to the species diversity in nematode communities to investigate if diversity-disease correlates occur in microscopic organisms. During the autumn season natural nematode communities were collected from decaying organic matter. Sampling locations were chosen based on locations where *C. elegans* was previously found [34,35], so that molecular tools in this species could be used for efficient discovery of stress induced by bacterial, microsporidian and viral parasites. Morphological and Nanopore 18S gene amplicon sequencing reconstructed the nematode diversity and community networks which were connected to parasite observations.

83   Altogether, our results provide the first indication of disease-diversity relationships within microscopic

84   communities and prospect a new experimental model system.

Methods

Sample collection and nematode extraction

Sampling was performed for five times during Autumn at the following three locations in the Netherlands: private garden (Heelsum), private vegetable garden (Wageningen), and patch of green surrounding a ditch (Renkum) (Supplementary Table S1). The active nematode community was extracted from the plant samples as previously described [36]. Additionally, details and a schematic overview covering the sample collection and nematode extraction are in Supplementary Text S1 and Supplementary Figure S1.

Microsporidian identification in extracted nematode communities

Filtered nematodes were propagated for ~5 days on a 6cm NGM plate seeded with 10x *E. coli* OP50. Nematode communities that perished before this procedure or contained mites were not considered in subsequent analyses. Once the population was composed of a large quantity of non-starved adults, a portion of the plate was chunked onto a new seeded 6cm NGM plate to continually propagate potential infections. The remainder of the plate was then washed with 700 µl of M9 media and placed in a microcentrifuge tube. Samples were allowed to gravity settle for ~1-5 minutes, to remove contaminating bacteria and fungi through supernatant removal. 700 µl of acetone was added to the nematode pellet prior to spinning down in a centrifuge for 30 seconds at 8,000 rcf. The supernatant was discarded, and 500 µl of the chitin binding Direct Yellow 96 (DY96) solution (1x PBST, 0.1% SDS, 20 ug/ml DY96) was added. The samples were incubated for 30 minutes in the dark at 20°C prior to spinning down in a centrifuge for 30 seconds at 8,000 rcf. The supernatant was removed, and 20 µl of EverBrite Mounting Medium (Biotium) was added to the samples prior to mounting on glass slides for imaging using an Axio Imager.M2 (Zeiss). Z-stacks were captured at 63x using an apotome unit with maximum projection. Samples were considered infected with microsporidia if spore clusters were visible in a population. Spore sizes were assessed by measuring the length and width of at least 45 spores using Zen software.

To identify the species of microsporidia present within a nematode sample, molecular characterization was performed. Briefly, 20 large adult nematodes were placed in 10 µl of lysis buffer (50mM KCl, 100mM Tris-HCl, 2.5mM MgCl2, 0.45% NP40) and placed in a thermocycler at 65°C for 60 minutes, followed by 95°C for 15 minutes. 2 µl of lysate was then used as template in a PCR reaction with the forward primer V1F [5' CACCAGGTTGATTCTGCCTGAC-3'] and the reverse primer 1492r [5'-GGTTACCTTGTTACGACTT-3'] or 18sR1492 [5'- GGAAACCTTGTTACGACTT-3'] to amplify microsporidian 18S. Sanger sequencing was performed on amplicons using the same primers listed above.

Indication of present parasites or parasites in the homogenized substrate using fluorescent reporter strains

C. elegans reporter strains AGD926 (zcIs4[*hsp-4::GFP*]), SJ4100 (zcIs13[*hsp-6p::GFP*]) and AU133 (agIs17[*myo-2p::mCherry + irg-1p::GFP*]) were obtained from the Caenorhabditis Genetic Center

(CGC). These strains indicate endoplasmic reticulum (ER) stress, mitochondrial stress or bacterial
virulence respectively (Supplementary Table S2) [37] and were used to explore presence of harmful
components in the microbiota of the homogenized substrate.

Starved populations from the reporter strains (AGD926, SJ4100, AU133) were transferred to
fresh 6cm NGM plates containing *E. coli* OP50 just before nematode extraction from natural
substrates. From each homogenized substrate 200µL blender solution was equally spread over the
plates containing the reporter strains that were then incubated at 20ºC for 48h. After incubation,
reporter nematodes exposed to blender solution were checked to investigate if GFP fluorescence was
higher than GFP expressed by nematodes grown on control plates containing only *E. coli* OP50 [37].
Fluorescence was checked using an Olympus SZX10 microscope with a NIGHTSEA Microscope
Fluorescence Adapter.

136   Observation of intracellular parasites using Intracellular Pathogen Response and viral stress reporter
137   strains

Extracted nematode communities were co-cultured with each of the following four reporter strains to
observe activation of antiparasitic and antiviral pathways: SOW1 and SOW6 that detect upregulation
of IPR genes *eol-1* and *pals-5* and SX2790 and SX2999 that detect upregulation of the antiviral gene
lys-3 (Supplementary Table S2) [38]. Fluorescence was checked once per day for 7 days using an
Olympus SZX10 stereomicroscope equipped with an SZX-RFA stereo fluorescence illuminator unit
and filter sets for GFP and RFP fluorescence viewing. Co-cultures were counted as positive if >10%
of the reporter *C. elegans* were fluorescing.

146   Extraction and screening for potential viruses

Co-cultures that resulted in reporter *C. elegans* fluorescence were washed off of test plates using M9
buffer and concentrated into 1 ml total volume via centrifugation. They were then homogenized via
vortexing for 4min with 30-50 1 mm diameter silicon carbide beads (BioSpec Products). Homogenate
was centrifuged to pellet debris, and supernatant was passed through a 0.22 µm pore size syringe filter
(GenClone). Filtrates were seeded onto plates with SOW6, SOW1, SX2790, or SX2999 reporter
strains and observed for fluorescence each day for 7 days.

154   DNA and RNA extraction

DNA and RNA were isolated from samples that contained at least 100 nematodes at the moment of
flash-freezing. From the 112 samples that met this requirement RNA and DNA was extracted as
described previously by Harkes *et al.* [39]. In short, nematodes were bead homogenized using 3mm
beads in a retch. Then RNA was isolated with a pH 4.5 phenol washing step, followed by a DNA
isolation buffer washing step at pH 8.0. DNA and RNA concentrations were determined via Qubit
fluorometer measurements (Thermo Scientific).

Nematode DNA amplification

After DNA/RNA extractions, 81 samples contained sufficient DNA quantity for 18S rRNA amplification. Nematode communities were characterized using the workflow by van Himbeeck *et al*. (2024), with minor adjustments [33]. EXP-NBD196 (Oxford Nanopore Technologies ltd., UK) barcoded primers 988F (5'- ctcaaagattaagccatgc-3') and 2646R (5'-gctaccttgttacgactttt-3') were used to amplify the near-complete 18S rRNA fragment (~1750bp) [33,40]. PCR was performed in quadruplicate per sample and each reaction consisted of 12.5 µl LongAMP Taq 2x MM, 400 nM of each primer, 3 µl DNA template and 5.5 µl autoclaved Mili-Q water. After amplification all 4 PCR replicates were pooled and DNA amplification was verified on agarose gels. Finally, DNA concentrations were determined via Qubit 4 Fluorometer measurements (Thermo Scientific).

Nematode 18S sequencing, read processing and species identification

The 18S rRNA was successfully amplified for 78 samples that were used for sequencing in four batches where each library contained equimolar ratios of all samples. Unwanted small fragments (<600bp) were removed prior to sequencing using NucleoMag NGS beads (0.5:1 bead:sample ratio) (Macherey Nagel). Library preparation was performed using the SQK-LSK112 kit, following the instructions of the manufacturer. Sequencing was performed on a MinION Mk1C using R9.4.1 flow cells.

Base-calling was performed using Guppy (v 6.2.1) in super accuracy mode (https://community.nanoporetech.com). The basecalled reads were demultiplexed using Guppy barcoder (v 6.2.1.) and adapters and barcodes were removed. Three samples (LvS26, LvS81, and LvS148) were excluded from the analysis at this stage because insufficient reads were obtained. Basecalling quality was controlled using NanoPlot (v1.40.0) (mean Phred score > 15) and Decona (v0.1.3) was used to filter 1400-2600bp reads with a >Q15 quality score [41,42]. Decona then clustered reads at 95% and Medaka consensus sequences were created from clusters with at least 100 reads. NCBI BLAST was used with an in-house database for taxonomic identification of nematode species [33,40]. Identifications with a similarity below 97% were excluded from the dataset. This data processing resulted in a OTU table containing the number of sequencing reads of each nematode taxon per sample. From this table, the number of species per sample was determined by summing the number of taxa with a read count larger than 0.

Data analysis

All data was processed and visualised using the package *tidyverse* and *ggpubr* in custom written scripts in R (v 4.2.1) [43,44]. Generalized linear models (glm), Chi-square and Wilcoxon Rank Sum tests were computed with *R base*. The effect of abiotic factors on community compositions was tested after creating a 'physeq' object in the *phyloseq* package and performing a PERMANOVA based on Bray-Curtis dissimilarity index (n = 10,000) using 'adonis2' from the *vegan* package [45,46]. The *vegan* package was also used for calculating diversity indices (richness and Shannon index (H')) with 'specnumber' and 'diversity' functions [45]. The nematode diversity was normalized (for substrate and community size) to facilitate cross-sample comparisons as the samples collected in this study

varied because of ephemeral nature of the substrates and the consequent opportunistic sampling method that was applied. Therefore, normalized nematode richness was defined as the total number of nematode species (based on sequencing data) found in a sample, divided by the number of individuals (based on microscopy-based count data) per gram wet substrate found in that sample. Nematode co-occurrence networks were based on significant Spearman correlations (p adjust < 0.05) between species (that were collected at least three times) as calculated with the 'rcorr' function in the *Hmisc* package and adjusted by the Holm-Bonferroni method. Networks were visualized in Cytoscape (v3.9.1) when at least 3 species were present within a network.

Data availability

C. elegans strains isolated in this study are available from CaeNDR (caendr.org) [34,35]. Custom written scripts and raw data files can be found on https://git.wur.nl/published_papers/van_himbeeck_2023_nematode_diversity_and_parasites/-/tree/main. The nematode sequencing data is accessible on SRA under BioProject PRJNA1021795. Processed data is included in the Supplementary Tables of this manuscript. Microsporidia 18S sequences are available in NCBI under the following accession numbers: OR636101, OR636102 and OR636103.

Results

Nematode community networks of the nematode C. elegans and co-occurring species

To link parasite presence to interspecies diversity in microorganisms, nematode communities were collected and analyzed. Nematode communities were obtained from decaying organic matter at three locations where *Caenorhabditis elegans* has previously been identified in the Netherlands [34,35] (Supplementary Figure S2, Supplementary Table S1). Nematode communities were subsequently extracted from the substrates, counted and the number of nematode species present was microscopically and molecularly established. Moreover, molecular analysis of the samples allowed for precise determination of the species present. The presence of *C. elegans* was of specific interest in this study, as this species was used as a reporter to monitor parasite presence.

In total 160 nematode communities were collected from ephemeral substrates and 76 of these were further characterised by sequencing (Supplementary Table S3). In these samples 58 different nematode species were molecularly characterised (Supplementary Table S4, S5). For 29 nematodes species only the genus name could be noted, as the used reference database does not always provide sequences annotated to species-level. The majority of the detected taxa were either bacterial or fungal feeders but also insect-parasitic, plant-parasitic and predatory nematodes were present (Supplementary Table S6). The estimated variance in nematode community composition originated for 20% from seasonal change, sampling location, moisture and pH differences (Supplementary table S7). Remaining variance in community compositions is likely in part attributable to biotic interactions, including nematode-bacteria, nematode-parasite, and nematode-nematode interactions [12,47]. Nematode-nematode interactions were examined by searching for specific cooccurrence or exclusion of certain species between samples. Most of the identified nematodes here were either opportunists or part of the basal fauna, although a few species exemplary for progressive succession (~5% of species) were also identified [48]. The species that formed cooccurrence networks were typically part of basal nematode fauna that is maintained over longer periods of time (Supplementary Table S4 and S8; Figure 1A). We did not observe any significant negative correlation between species. *Panagrolaimus subelongatus* (17), *Rhabditophanes sp.* (16), *Panagrolaimusrigidus* (16), *Acrostichus sp.* (15), *Rhabditoides sp.* (13), *Panagrolaimus sp.* (13), *Rhabditella axei* (12), *Pristionchus entomophagus* (10) and *Eucephalobus striatus* (10) co-occurred at least 10 times with *C. elegans* agreeing with previously found coincidences of these species (Félix and Duveau, 2012). Nevertheless, *C. elegans* nematodes did not coincide with these particular species more often than expected based on chance which underlines the opportunistic behaviour of *C. elegans* (Supplementary Table S4). Moreover, *Caenorhabditis* species are known to coexist together in some places [49–51]. Here, only one sample with a second *Caenorhabditis* species was obtained. This decaying kohlrabi contained *C. briggsae* and 7 other nematode species, but not *C. elegans*, despite molecular identification of *C. elegans* on other kohlrabies from the same location. *C. elegans* was detected in more than 71 % of the samples and we found that communities where *C. elegans* is present and absent have similar nematode richness (p=0.56, Wilcoxon Rank-Sum test). These findings support our use of it as a parasite reporter species for communities that differ in nematode diversity.

Figure 1 Nematode-nematode co-occurrences in decaying organic plants – A) Spearman-based (adjusted p < 0.05) co-occurrence networks of species that were present in at least 3 samples and contained at least 3 species. Colours indicate the c-p value of the species depicted (colonizer-persister value; higher values indicate species typically present in more mature nematode communities): yellow indicates c-p value 1, pink indicates c-p value 2, red indicates c-p value 3, blue indicates c-p value 4, green indicates c-p value 5, grey indicates c-p values are not known or variable for this genus [48]. Circle size reflects the number of samples in which a species was observed in the sequencing data. B) Nematode richness (in normalized species richness) in samples where *C. elegans* was absent or present.

Nematode diversity determines parasite presence in natural field populations of nematodes

Collected nematode communities were screened for phenotypes that indicate: i) presence of microsporidia ii) mitochondrial stress, ER stress and stress caused by bacterial virulence [37], iii) IPR activation and iv) presumptive viruses [52] (Figure 2A). Some nematode communities died before they could be screened or could not be maintained successfully in the laboratory, and were excluded from further analysis (Figure 2B). All experiments harboured indications of parasite presence and 80% of the collected samples responded in at least one of the experimental assays (Figure 2C). In most cases, samples were found positive for multiple stressors (Figure 2C). Notably, these numbers likely underrepresent the total number of parasites occurring in the wild as not all may infect the reporter strains used, parasites might be lost during the extraction procedure and nematodes potentially evade parasites in some of the assays [53–55]. Taken together, these data suggest that biotic stress is a common burden for wild nematodes.

Subsequently, we compared parasite presence to nematode diversity for each of the experimental assays (Supplementary Table S9). For two assays, diversity-disease correlations were observed: 1) microsporidia were more often present in communities with lowered nematode richness (Figure 2D) and 2) activation of the *eol-1*-based IPR reporter strain was more often observed in co-cultures when communities with a lowered nematode richness (Figure 2E). In agreement with previous studies, we found that presumptive *C. elegans*-infecting viruses were rare and observed in only 2 (1.5%) of the screened communities [31]. Both communities contained lower than average nematode richness (Figure 2F), but the low incidence of viral infections obstructs drawing solid conclusions

from this observation. For the other experimental assays we did not observe diversity-disease
relationships (Supplementary Figure S3, S4, Supplementary Table S3).

Diversity-disease relationships can be driven by host density, frequency or a combination of
both [4,7,13,14]. To explore which of these may determine diversity-disease relationships in microbial
nematode communities, the data gained in the microsporidian assay was explored further. We
identified microsporidia in 15 out of 76 screened samples (16%) (Supplementary figure S5, S6).
Almost half of these microsporidia isolates were found in *C. elegans* (Supplementary Table S10). The
microsporidian isolates included the previously described species *Nematocida parisii* [28,56], but not
all microsporidia could be molecularly determined and may represent novel species. Moreover,
microsporidia were identified in *Pristionchus uniformus*, *Panagrolaimus rigidus* and in other
nematodes of which the species could not be determined. Here, host density was defined as the
number of host species per wet gram of substrate and host frequency was defined as the proportion of
the host amongst the different nematode species in the sample. Analysis by a generalized linear model
suggested a trend that host density (p = 0.058) as well as the interaction between host density and
frequency (p = 0.072) might affect microsporidian presence in *C. elegans* (Supplementary Table S11),
but additional data would be essential to observe if the same trend remains. As indicated above, key
species may affect spread of parasites, but we could not pinpoint presence of key species in our
relatively small dataset containing only 3 identified microsporidian hosts species (chi-square p > 0.05
for expected versus observed microsporidia infections per species). Microsporidia spores are known to
be resistant to environmental conditions, including pH and desiccation [57,58]. Furthermore, pH has
been shown to induce germination is thought influence the infectivity of microsporidia [59]. Here,
microsporidia were more often observed on alkaline substrates, whereas substrate moisture varied
(Supplementary Figure S7). In conclusion, because pH and nematode diversity were not correlated
(glm, p = 0.46) (Supplementary Table S12), microsporidian presence in our samples was more likely
to be observed in nematode communities with lower richness (possibly dependent on relatively higher
host density- and frequency within the community) and an alkaline environment.

Figure 2 Biotic stress in nematode communities using *C. elegans* as a reporter species – A) The four different experimental assays to screen for biotic stress: i) staining of extracted nematodes to visualise microsporidia, ii) screening of *C. elegans* reporter strains on homogenized sample substrate containing a mixture of microbes, iii) co-culturing *C. elegans* reporter strains with the extracted nematode community and iv) co-culturing *C. elegans* reporter strains on 0.2 µm filtrate of the extracted nematode community B) The fraction of samples that was found to respond to each of the biotic stress indication assays. Because only a selection of samples was screened for virus presence after a 0.2 µm filter step (assay iv) these samples are not depicted here. C) Overlap of biotic stress indications within each of the samples. The plot shows how many samples (count y-axis) responded to specific combinations of biotic stress assays (x-axis). Colours illustrate the number of stress assays a sample was found positive for ranging from five (red) to none (grey). D) Normalized nematode richness for samples in which microsporidia were present or absent. E) Normalized nematode richness

328   for samples which did or did not indicated mitochondrial stress on upregulation of the *C. elegans* IPR
329   gene *eol-1* upon coculture. F) Normalized nematode richness for samples which did or did not contain
330   *C. elegans* viruses.

Discussion

Although microscopic communities are vital for ecosystem functioning and stability, host-parasite relationships and the diversity-disease relationships have been poorly studied in microscopic communities. We argue that understanding diversity-disease effects in microbial communities is essential for predicting ecosystem functioning in a rapidly changing environment. Here we used wild nematode communities to study the effect of host diversity on the occurrence of parasites in microbial communities. Nanopore sequencing was applied for high-resolution characterization of nematodes present within each community. With this method we molecularly defined nematode communities naturally occurring with the important model organism *C. elegans*. Fluorescent reporter tools available in this model organism were used to identify bacterial-, microsporidian- and viral-induced stress inflicted on *C. elegans*. These screenings also discovered potentially new parasite species. In line with the dilution theory, some specialistic parasites were more often found in communities with fewer nematode species, indicating that diversity-disease effects can be found among microscopic organisms.

Specific nematode-parasite interactions may underlie diversity-disease effects

Diversity-disease relationships within the nematode communities were tested via four different biotic stress assays examining nine classes of biotic stress. The vast majority of nematode communities in the collected samples responded in at least one of these tests, suggesting that biotic stress is frequent in wild nematode populations. Within these communities, we found correlation between nematode diversity and intracellular parasite prevalence as 1) samples containing microsporidian parasites had a lower nematode diversity and 2) the *eol-1*-based Intracellular Pathogen Response reporter strain more often responded to communities with a lower diversity. The experimental assays to indicate biotic stress covered diverse possible origins of stress-inducing parasites with varying host specificities. Parasites most prone to experience dilution effects are those that have a narrow host range [13]. Our data corresponds with this theory, as we observed potential dilution effects in microsporidian infected communities. Microsporidia are parasites of most types of animals with the vast majority of species only observed to display host and tissue specificity in one or two closely related hosts [59–61]. Microsporidia that infect nematodes also often display specialized tissue specificity, though there are examples of more generalist species [28,62]. Microsporidian occurrence in communities with lower species richness thus enhances their chance of encountering a high-quality host. Nevertheless, the lethal nature of microsporidia may also explain lower nematode diversity in communities suffering from microsporidia with a broad host range. We also observed potential dilution effects using the *eol-1* reporter strain in *rde-1* immunocompromised *C. elegans*. *Eol-1* is an RNA decapping enzyme that is upregulated during viral and microsporidian infections and is involved in activating enhanced RNAi in response to mitochondrial stress [63–66].

For other biotic stress reporter assays, we did not observed diversity-disease effects. The *hsp-6* reporter strain reports mitochondrial stress just like *eol-1*, but this reporter strain lacked the sensitivity-increasing *rde-1* mutation which is a possible explanation for not different observations between these two reporter strains. Also reporter strains (based on *lys-3*) that respond to viruses did not show a correlation with nematode diversity, despite the narrow host range of viruses [67]. As we have only

identified two *C. elegans* viruses in our screening a larger study would be necessary to draw
conclusions [31]. Another of the reporter strains responding to intracellular parasites (based on *pals-5*)
did not show any diversity-disease relationship. It was noted that there were relatively few sequenced
communities for which expression of this reporter strain was not observed. Finally, bacteria are
usually opportunistic parasites not bound to certain hosts, therefore dilution effects were not expected,
but amplification effects may occur. Nevertheless, these were not observed in the reporters strain that
indicate bacterial virulence and also not in the ER stress reporter strain that detects stress caused by
both generalists and specialists.

Potential mechanistic bases of diversity-disease effects in ephemeral nematode communities

The short-lived ephemeral nematode communities sampled in this study consisted mostly of bacterial-
feeding *r*-strategists that could experience strong competition for food in their temporary environment.
The life history traits of the species present in these nematode communities may determine disease-
diversity effects. Competition between species can drive dilution effects, especially in absence of
predation [4]. Although several predatory species were observed, these were often present at low
density and frequency. An exception was formed for facultative predators like *Pristionchus* [68], but
these may also feed on bacteria in this habitat. Together, low predation and high competition
circumstances suggest possibilities for observation of dilution effects, and we indeed observed a lower
species diversity for nematode communities that were infected with microsporidia. Yet of note,
potential predation by other species, such as mites or fungi was not studied here. Moreover, migration
of ephemeral-specialised nematodes provides a spatial factor in the host-parasite dynamics that might
be compared to that of a fragmented landscape. Fragmentation of ephemeral nematode communities
may further contribute to the observation of diversity-disease effects [15,20,70,71].

Detailed insight into ephemeral nematode community compositions that include C. elegans

To date, a comprehensive overview of nematode species co-occurring with *C. elegans* in its natural
habitats was lacking. Historically, most studies investigated nematode community compositions in soil
and water instead (e.g. for agricultural purposes or environmental monitoring). Here long-read
sequencing data was used to identify community members that occur in the same habitats as *C.*
elegans. We performed (nearly) full 18S sequencing to characterize the total nematode community on
a low taxonomic level (mostly species or genus) [33], which yielded a total of 96 taxa. Subsequent
studies may further broaden knowledge about *C. elegans* community compositions by simultaneously
including other microfauna such as protists and bacteria. In agreement with previous studies, we
identified nematodes that are known to co-occur with *C. elegans* from the genera *Caenorhabditis (C.*
briggsae), Oscheius (O. tipulae and *O. dolichura), Panagrolaimus (P. detritophagus, P. rigidus, P.*
subelongatus), *Rhabditophanes* and *Mesorhabditis (M. spiculigera)* and *Aphelenchoides* in our
communities [49,50]. *E. striatus* is part of the basal nematode fauna, but all of the other species are
species that quickly respond to enriched food resources [48]. The observation that these other
nematodes do not specifically co-occur suggest that *C. elegans* can be considered an opportunist
whose presence may be observed across suitable ecological niches of ephemeral substrates in the
Netherlands.

Conclusions

Biodiversity is important as it not only contributes to ecosystem functioning, but also to ecosystem stability, as exemplified by the diversity-disease relationships. Below-ground microbial communities - such as nematode communities - are vulnerable to shifts and biodiversity loss due to global change [72]. As microbial communities are vital for the majority of ecosystems, it is important to understand their response to a decrease in biodiversity in terms of parasite dynamics. Our study provides the first demonstration of diversity-disease relationships in microscopic nematode communities. Diversity-disease relationships depend on dynamic multifactorial processes in communities, and as such may be strikingly difficult to understand or capture. Studying diversity-disease effects in nematode populations poses a promising outlook for the future. In this study, we have extracted communities from the field through the use of a small plastic bag and have shown these could be observed and subsequently disassembled in the lab. Many collected species were successfully cultured afterwards, indicating follow-up experiments can be used to determine exactly which factors could cause diverse-disease effects to occur in nematode communities. Under laboratory settings, one can deduct or control factors such as species presence, density and select community members (including competitors and predators) based on specific life history traits. Furthermore, nematode parasites can be examined to trace infection and transmission (even in real time) [73,74]. Together, our results invite further characterization of host-parasite interactions in microscopic communities, and open opportunities for using nematodes communities as a diversity-disease model system in the lab and field.

Acknowledgements

The authors thank Semih K. Aslan (primer design), Hans Helder (primer design, feedback on initial results), Stefan Geisen, Mark Sterken and Jan Kammenga (feedback on experimental set-up), Boas Kanis (sampling pilot), David Jordan and Eric Miska (reporter strains), Erik Andersen and Robyn Tanny (CaeNDR database) for their contributions. Some strains were provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). Funding for this research was provided by the Ecology Fund Grant of the Royal Netherlands Academy of Arts and Sciences. This work was supported by a Canadian Institutes of Health Research grant no. 400784 (to A.R.) and H.T.E.J was supported by a University of Toronto Open Fellowship and an Ontario Graduate Scholarship.

Author contribution statement

LvS designed the research. RvH, JNS, HTEJ, WW, JOV, JAG, AR and LvS collected and processed nematode community, parasite and sequencing data. RvH and LvS performed computational analyses. RvH and LvS wrote the first draft of the manuscript, and all authors contributed to the interpretation of results and article writing.

References

1. Buck JC. Indirect Effects Explain the Role of Parasites in Ecosystems. Trends in Parasitology. Elsevier Ltd; 2019. pp. 835–847. doi:10.1016/j.pt.2019.07.007

2. Frainer A, McKie BG, Amundsen PA, Knudsen R, Lafferty KD. Parasitism and the Biodiversity-Functioning Relationship. Trends in Ecology and Evolution. Elsevier Ltd; 2018. pp. 260–268. doi:10.1016/j.tree.2018.01.011

3. Wagg C, Bender SF, Widmer F, Van Der Heijden MGA. Soil biodiversity and soil community composition determine ecosystem multifunctionality. Proc Natl Acad Sci U S A. 2014;111: 5266–5270. doi:10.1073/pnas.1320054111

4. Su M, Jiang Z, Hui C. How Multiple Interaction Types Affect Disease Spread and Dilution in Ecological Networks. Front Ecol Evol. 2022;10.

5. Gómez-Hernández EA, Moreno-Gómez FN, Bravo-Gaete M, Córdova-Lepe F. Concurrent dilution and amplification effects in an intraguild predation eco-epidemiological model. Sci Rep. 2023;13. doi:10.1038/s41598-023-33345-2

6. Levi T, Keesing F, Holt RD, Barfield M, Ostfeld RS. Quantifying dilution and amplification in a community of hosts for tick-borne pathogens. Ecological Applications. 2016;26: 484–498. doi:10.1890/15-0122

7. Keesing F, Ostfeld RS. Dilution effects in disease ecology. Ecol Lett. 2021;24: 2490–2505. doi:10.1111/ele.13875

8. Ferraguti M, Martínez-De la Puente J, Jiménez–Clavero MÁ, Llorente F, Roiz D, Ruiz S, et al. A field test of the dilution effect hypothesis in four avian multi-host pathogens. PLoS Pathog. 2021;17: 1–21. doi:10.1371/journal.ppat.1009637

9. Johnson PTJ, Preston DL, Hoverman JT, Richgels KLD. Biodiversity decreases disease through predictable changes in host community competence. Nature. 2013;494: 230–233. doi:10.1038/nature11883

10. Johnson PTJ, Preston DL, Hoverman JT, LaFonte BE. Host and parasite diversity jointly control disease risk in complex communities. Proc Natl Acad Sci U S A. 2013;110: 16916–16921. doi:10.1073/pnas.1310557110

11. Johnson PTJ, Hartson RB, Larson DJ, Sutherland DR. Diversity and disease: Community structure drives parasite transmission and host fitness. Ecol Lett. 2008;11: 1017–1026. doi:10.1111/j.1461-0248.2008.01212.x

12. Okamoto KW, Amarasekare P, Post DM, Vasseur DA, Turner PE. The interplay between host community structure and pathogen life□history constraints in driving the evolution of host□range shifts. Hawley D, editor. Funct Ecol. 2019;33: 2338–2353. doi:10.1111/1365-2435.13467

13.   Collins CD, Bever JD, Hersh MH. Community context for mechanisms of disease dilution:
insights from linking epidemiology and plant–soil feedback theory. Annals of the New York
Academy of Sciences. Blackwell Publishing Inc.; 2020. pp. 65–85. doi:10.1111/nyas.14325

14.   Faust CL, Dobson AP, Gottdenker N, Bloomfield LSP, McCallum HI, Gillespie TR, et al. Null
expectations for disease dynamics in shrinking habitat: Dilution or amplification? Philosophical
Transactions of the Royal Society B: Biological Sciences. 2017;372.
doi:10.1098/rstb.2016.0173

15.   Liu X, Xiao Y, Lin Z, Wang X, Hu K, Liu M, et al. Spatial scale-dependent dilution effects of
biodiversity on plant diseases in grasslands. Ecology. 2023;104. doi:10.1002/ecy.3944

16.   Chen S, Liu X, He Q, Zhou S. Higher-order interactions on disease transmission can reverse
the dilution effect or weaken the amplification effect to unimodal pattern. Ecol Modell. 2022.
doi:10.1016/j.ecolmodel.2022.110156

17.   Gandy S, Kilbride E, Biek R, Millins C, Gilbert L. No net effect of host density on tick-borne
disease hazard due to opposing roles of vector amplification and pathogen dilution. Ecol Evol.
2022;12. doi:10.1002/ece3.9253

18.   Linske MA, Williams SC, Stafford KC, Ortega IM. Ixodes scapularis (Acari: Ixodidae)
Reservoir Host Diversity and Abundance Impacts on Dilution of Borrelia burgdorferi
(Spirochaetales: Spirochaetaceae) in Residential and Woodland Habitats in Connecticut, United
States. J Med Entomol. 2018;55: 680–690. doi:10.1093/JME/TJX237

19.   Civitello DJ, Cohen J, Fatima H, Halstead NT, Liriano J, McMahon TA, et al. Biodiversity
inhibits parasites: Broad evidence for the dilution effect. Proc Natl Acad Sci U S A. 2015;112:
8667–8671. doi:10.1073/pnas.1506279112

20.   Halliday FW, Rohr JR, Laine AL. Biodiversity loss underlies the dilution effect of biodiversity.
Ecol Lett. 2020;23: 1611–1622. doi:10.1111/ele.13590

21.   Liu X, Chen L, Liu M, García-Guzmán G, Gilbert GS, Zhou S. Dilution effect of plant
diversity on infectious diseases: latitudinal trend and biological context dependence. Oikos.
2020;129: 457–465. doi:10.1111/oik.07027

22.   van den Hoogen J, Geisen S, Routh D, Ferris H, Traunspurger W, Wardle DA, et al. Soil
nematode abundance and functional group composition at a global scale. Nature. 2019;572:
194–198. doi:10.1038/s41586-019-1418-6

23.   Gaertner BE, Phillips PC. Caenorhabditis elegans as a platform for molecular quantitative
genetics and the systems biology of natural variation. Genet Res (Camb). 2010;92: 331–348.
doi:10.1017/S0016672310000601

24.   Marsh EK, May RC. Caenorhabditis elegans, A model organism for investigating immunity.
Applied and Environmental Microbiology. 2012. pp. 2075–2081. doi:10.1128/AEM.07486-11

25.   Hahnel SR, Dilks CM, Heisler I, Andersen EC, Kulke D. Caenorhabditis elegans in
anthelmintic research – Old model, new perspectives. Int J Parasitol Drugs Drug Resist. 2020;
105398. doi:10.1016/j.ijpddr.2020.09.005

26.   Link CD. C. elegans models of age-associated neurodegenerative diseases: Lessons from
523   transgenic worm models of Alzheimer's disease. Exp Gerontol. 2006;41: 1007–1013.
524   doi:10.1016/j.exger.2006.06.059

525   27.   Schulenburg H, Félix M. The Natural Biotic Environment of Caenorhabditis elegans. Genetics.
526   2017;206: 55–86. doi:10.1534/genetics.116.195511

527   28.   Zhang G, Sachse M, Prevost M-C, Luallen RJ, Troemel ER, Felix M-A. A Large Collection of
528   Novel Nematode- Infecting Microsporidia and Their Diverse Interactions with *Caenorhabditis*
529   *elegans* and Other Related Nematodes. PLoS Pathog. 2016;12.
530   doi:10.1371/journal.ppat.1006093

531   29.   Osman GA, Fasseas MK, Koneru SL, Essmann CL, Kyrou K, Srinivasan MA, et al. Natural
532   Infection of C. elegans by an Oomycete Reveals a New Pathogen-Specific Immune Response.
533   Current Biology. 2018;28: 640-648.e5. doi:10.1016/j.cub.2018.01.029

534   30.   Richaud A, Frézal L, Tahan S, Jiang H, Blatter JA, Zhao G, et al. Vertical transmission in
535   Caenorhabditis nematodes of RNA molecules encoding a viral RNA-dependent RNA
536   polymerase. Proc Natl Acad Sci U S A. 2019;116: 24738–24747.
537   doi:10.1073/pnas.1903903116

538   31.   Frézal L, Jung H, Tahan S, Wang D, Félix M-A. Noda-Like RNA Viruses Infecting
539   Caenorhabditis Nematodes: Sympatry, Diversity, and Reassortment. J Virol. 2019;93.
540   doi:10.1128/jvi.01170-19

541   32.   Grover M, Fasseas MK, Essmann C, Liu K, Braendle C, Félix MA, et al. Infection of C.
542   elegans by Haptoglossa Species Reveals Shared Features in the Host Response to Oomycete
543   Detection. Front Cell Infect Microbiol. 2021;11. doi:10.3389/fcimb.2021.733094

544   33.   van Himbeeck R, Cazzaniga SG, van den Elsen S, Oude Vrielink J, Aslan SK, Visser J, et al. A
545   full-length SSU rRNA-based workflow for high-resolution monitoring of nematode
546   communities reveals direct and indirect responses to plant-based manipulations. Soil Biol
547   Biochem. 2024;189. doi:10.1016/j.soilbio.2023.109263

548   34.   Cook DE, Zdraljevic S, Roberts JP, Andersen EC. CeNDR, the Caenorhabditis elegans natural
549   diversity resource. Nucleic Acids Res. 2017;45: D650–D657. doi:10.1093/nar/gkw893

550   35.   Crombie TA, McKeown R, Moya ND, Evans KS, Widmayer SJ, LaGrassa V, et al. CaeNDR,
551   the Caenorhabditis Natural Diversity Resource. Nucleic Acids Res. 2024;52: D850–D858.
552   doi:10.1093/nar/gkad887

553   36.   Van Bezooijen J. Methods and Techniques for Nematology. 2006. Available:
554   https://www.wur.nl/en/show/manual-methods-and-techniques-for-nematology-1.htm

555   37.   Samuel BS, Rowedder H, Braendle C, Félix M-AA, Ruvkun G. Caenorhabditis elegans
556   responses to bacteria from its natural habitats. Proc Natl Acad Sci U S A. 2016/06/19.
557   2016;113: E3941–E3949. doi:10.1073/pnas.1607183113

38. Le Pen J, Jiang H, Di Domenico T, Kneuss E, Kosałka J, Leung C, et al. Terminal
uridylyltransferases target RNA viruses as part of the innate immune system. Nat Struct Mol
Biol. 2018;25: 778–786. doi:10.1038/s41594-018-0106-9

39. Harkes P, Suleiman AKA, van den Elsen SJJ, de Haan JJ, Holterman M, Kuramae EE, et al.
Conventional and organic soil management as divergent drivers of resident and active fractions
of major soil food web constituents. Sci Rep. 2019;9: 1–15. doi:10.1038/s41598-019-49854-y

40. Van Megen H, Van Den Elsen S, Holterman M, Karssen G, Mooyman P, Bongers T, et al. A
phylogenetic tree of nematodes based on about 1200 full-length small subunit ribosomal DNA
sequences. Nematology. 2009;11: 927–950. doi:10.1163/156854109X456862

41. De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. NanoPack: Visualizing
and processing long-read sequencing data. Bioinformatics. 2018;34: 2666–2669.
doi:10.1093/bioinformatics/bty149

42. Oosterbroek S, Doorenspleet K, Nijland R, Jansen L. Decona: From demultiplexing to
consensus for Nanopore amplicon data. ARPHA Conference Abstracts. 2021;4.
doi:10.3897/aca.4.e65029

43. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to the
Tidyverse. J Open Source Softw. 2019;4: 1686. doi:10.21105/joss.01686

44. Kassambara Alboukadel. ggpubr: "ggplot2" Based Publication Ready Plots. In:
https://rpkgs.datanovia.com/ggpubr/. 2023.

45. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan:
Community Ecology Package. 2020. Available: https://cran.r-project.org/package=vegan

46. McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. PLoS One. 2013;8. doi:10.1371/journal.pone.0061217

47. Vandvik V, Skarpaas O, Klanderud K, Telford RJ, Halbritter AH, Goldberg DE. Biotic
rescaling reveals importance of species interactions for variation in biodiversity responses to
climate change. Proceedings of the National Academy of Sciences. 2020;117: 22858–22865.
doi:10.1073/pnas.2003377117

48. Ferris H, Bongers T, De Goede RGM. A framework for soil food web diagnostics: extension of
the nematode faunal analysis concept. Applied Soil Ecology. 2001.

49. Félix M-A, Duveau F. Population dynamics and habitat sharing of natural populations of
Caenorhabditis elegans and C. briggsae. BMC Biol. 2012/06/27. 2012;10: 59.
doi:10.1186/1741-7007-10-59

50. Schulenburg H, Petersen C, Strathmann E, Prahl S, Dirksen P. The prevalence of
Caenorhabditis elegans across 1.5 years in selected North German locations: the importance of
substrate type, abiotic parameters, and Caenorhabditis competitors. BMC Ecol. 2014;14: 4.
doi:10.1186/1472-6785-14-4

51.   Sloat SA, Noble LM, Paaby AB, Bernstein M, Chang A, Kaur T, et al. Caenorhabditis
nematodes colonize ephemeral resource patches in neotropical forests. Ecol Evol. 2022;12.
doi:10.1002/ece3.9124

52.   Jarkass HT El, Mok C, Schertzberg MR, Fraser AG, Troemel ER, Reinke AW. An intestinally
secreted host factor promotes microsporidia invasion of C. Elegans. Elife. 2022;11.
doi:10.7554/eLife.72458

53.   Kaletsky R, Moore RS, Vrla GD, Parsons LR, Gitai Z, Murphy CT. C. elegans interprets
bacterial non-coding RNAs to learn pathogenic avoidance. Nature. 2020;586: 445–451.
doi:10.1038/s41586-020-2699-5

54.   Meisel JD, Kim DH. Behavioral avoidance of pathogenic bacteria by Caenorhabditis elegans.
Trends in Immunology. Elsevier Ltd; 2014. pp. 465–470. doi:10.1016/j.it.2014.08.008

55.   Schulenburg H, Ewbank JJ. The genetics of pathogen avoidance in Caenorhabditis elegans.
Molecular Microbiology. 2007. pp. 563–570. doi:10.1111/j.1365-2958.2007.05946.x

56.   Troemel ER, Félix MA, Whiteman NK, Barrière A, Ausubel FM. Microsporidia are natural
intracellular parasites of the nematode Caenorhabditis elegans. PLoS Biol. 2008;6.
doi:10.1371/journal.pbio.0060309

57.   Fenoy S, Rueda C, Higes M, Martín-Hernández R, Del Aguila C. High-level resistance of
Nosema ceranae, a parasite of the honeybee, to temperature and desiccation. Appl Environ
Microbiol. 2009;75: 6886–6889. doi:10.1128/AEM.01025-09

58.   Santillana-Hayat M, Sarfati C, Fournier S, Chau F, Porcher R, Molina JM, et al. Effects of
chemical and physical agents on viability and infectivity of Encephalitozoon intestinalis
determined by cell culture and flow cytometry. Antimicrob Agents Chemother. 2002;46: 2049–
2051. doi:10.1128/AAC.46.6.2049-2051.2002

59.   Willis AR, Reinke AW. Factors That Determine Microsporidia Infection and Host Specificity.
2022. pp. 91–114. doi:10.1007/978-3-030-93306-7_4

60.   Murareanu BM, Sukhdeo R, Qu R, Jiang J, Reinke AW. Generation of a Microsporidia Species
Attribute Database and Analysis of the Extensive Ecological and Phenotypic Diversity of
Microsporidia. 2021. doi:10.1128/mBio

61.   Bojko J, Reinke AW, Stentiford GD, Williams B, Rogers MSJ, Bass D. Microsporidia: a new
taxonomic, evolutionary, and ecological synthesis. Trends in Parasitology. Elsevier Ltd; 2022.
pp. 642–659. doi:10.1016/j.pt.2022.05.007

62.   Wadi L, Reinke AW. Evolution of microsporidia: An extremely successful group of eukaryotic
intracellular parasites. PLoS Pathog. 2020;16. doi:10.1371/journal.ppat.1008276

63.   Han B, Ma Y, Tu V, Tomita T, Mayoral J, Williams T, et al. Microsporidia interact with host
cell mitochondria via voltage-dependent anion channels using sporoplasm surface protein 1.
mBio. 2019;10. doi:10.1128/mBio.01944-19

64. Mao K, Breen P, Ruvkun G. Mitochondrial dysfunction induces RNA interference in C.
elegans through a pathway homologous to the mammalian RIG-I antiviral response. PLoS Biol.
2020;18. doi:10.1371/journal.pbio.3000996

65. Sowa JN, Jiang H, Somasundaram L, Tecle E, Xu G, Wang D, et al. The Caenorhabditis
elegans RIG-I Homolog DRH-1 Mediates the Intracellular Pathogen Response upon Viral
Infection. Pfeiffer JK, editor. J Virol. 2019;94: e01173-19. doi:10.1128/JVI.01173-19

66. Chen K, Franz CJ, Jiang H, Jiang Y, Wang D. An evolutionarily conserved transcriptional
response to viral infection in Caenorhabditis nematodes. BMC Genomics. 2017;18: 303.
doi:10.1186/s12864-017-3689-3

67. Félix MA, Ashe A, Piffaretti J, Wu G, Nuez I, Bélicard T, et al. Natural and experimental
infection of *Caenorhabditis* nematodes by novel viruses related to nodaviruses. PLoS Biol.
2011/02/02. 2011;9: e1000586. doi:10.1371/journal.pbio.1000586

68. Serobyan V, Ragsdale EJ, Sommer RJ. Adaptive value of a predatory mouth-form in a
dimorphic nematode. Proceedings of the Royal Society B: Biological Sciences. 2014;281.
doi:10.1098/rspb.2014.1334

69. Johnson PTJ, Rohr JR, Hoverman JT, Kellermanns E, Bowerman J, Lunde KB. Living fast and
dying of infection: Host life history drives interspecific variation in infection and disease risk.
Ecol Lett. 2012;15: 235–242. doi:10.1111/j.1461-0248.2011.01730.x

70. Liu X, Chen F, Lyu S, Sun D, Zhou S. Random species loss underestimates dilution effects of
host diversity on foliar fungal diseases under fertilization. Ecol Evol. 2018;8: 1705–1713.
doi:10.1002/ece3.3749

71. Johnson EE, Escobar LE, Zambrana-Torrelio C. An Ecological Framework for Modeling the
Geography of Disease Transmission. Trends in Ecology and Evolution. Elsevier Ltd; 2019. pp.
655–668. doi:10.1016/j.tree.2019.03.004

72. Berlinches de Gea A, Hautier Y, Geisen S. Interactive effects of global change drivers as
determinants of the link between soil biodiversity and ecosystem functioning. Glob Chang
Biol. 2022. doi:10.1111/gcb.16471

73. Pirenne L, Jakobs MAH, Jordan D, Franze K, Miska EA. The Orsay Virus as a model for
population-wide viral infection dynamics. bioRxiv. 2021; 2021.03.02.433572. Available:
https://www.biorxiv.org/content/10.1101/2021.03.02.433572v1%0Ahttps://www.biorxiv.org/c
ontent/10.1101/2021.03.02.433572v1.abstract

74. Shaw CL, Kennedy DA. Developing an empirical model for spillover and emergence: Orsay
virus host range in Caenorhabditis. Proceedings of the Royal Society B: Biological Sciences.
2022;289: 20221165. doi:10.1098/rspb.2022.1165

Supplementary information

Supplementary Text S1 Sample collection and nematode extraction – Detailed description of
sample collection and nematode extractions.

Supplementary Table S1 Summary of sampling location information – The table indicates
location details and number of collected samples at each location.

Supplementary Table S2 *C. elegans* **reporter strain genetic information**

Supplementary Table S3 Collected sample data – Contains information per sample collected
including sample location, data, type, substrate weight, moisture, pH, nematode density and
normalized species richness (based on morphological species identification) and whether or not *C.*
elegans was isolated from the sample.

Supplementary Table S4 Nanopore sequencing data output – The data shows reads per species per
sample.

Supplementary Table S5 Sequencing abbreviations of nematode species – The data shows reads
abbreviations used for sequenced nematode species.

Supplementary Table S6 Overview of molecularly identified nematodes in the sequenced
nematode communities The table indicates the nematode taxon, Maturity Index (according to its c-p
value), feeding preference and percentage of the samples in which this taxon was discovered.

Supplementary Table S7 Abiotic effects on community composition based on a permutation
model

Supplementary Table S8 Co-occurring species in nematode communities – Based on Spearman
correlations and Hochberg-corrected p-values.

Supplementary Table S9 Diversity indices of sequenced samples – Shannon's H, absolute richness
(number of species) and normalised species richness (species corrected for sampling effort) for each
sample based on sequencing.

Supplementary Table S10 Data microsporidian species and hosts – PCR-based host and
microsporidian species identification and morphological characteristics of the nematode species
identified. Some PCRs failed or nematodes were lost during the screening, here the missing data is
indicated with NA.

Supplementary Table S11 Effects of frequency and density on microsporidia of *C. elegans* –
GLM testing dilution and frequency effects on microsporidian occurrence in C. elegans.

Supplementary Table S12 Correlation of normalised richness and pH – GLM testing effect of pH
on sequence-based normalised species richness.