Effects of urbanisation, habitat characteristics, and management on garden pond

biodiversity: findings from a large-scale citizen science survey

Authors

Zsuzsanna Márton[1,2*], Barbara Barta[1,2,3], Csaba F. Vad[1,2], Beáta Szabó[1,2], Andrew J. Hamer[1,2],

Vivien Kardos[1], Csilla Laskai[1,2], Ádám Fierpasz[1,3], Zsófia Horváth[1,2,4]

[1]Institute of Aquatic Ecology, HUN-REN Centre for Ecological Research, Karolina út 29,

1113 Budapest, Hungary

[2]National Multidisciplinary Laboratory for Climate Change, HUN-REN Centre for Ecological

Research, Karolina út 29, 1113 Budapest, Hungary

[3]Doctoral School of Biology, ELTE Eötvös Loránd University, Pázmány Péter sétány 1/c,

1113 Budapest, Hungary

[4]ELTE Eötvös Loránd University, Pázmány Péter sétány 1/c, 1113 Budapest, Hungary

Corresponding author

Zsuzsanna Márton, HUN-REN Institute of Aquatic Ecology, Centre for Ecological Research,

Budapest, Karolina út 29, 1113 Budapest, Hungary

Email: marton.zsuzsanna@ecolres.hu

Abstract

The rapid expansion of urban areas often leads to degradation, fragmentation, and loss of

natural habitats, threatening biodiversity. While urban ponds might contribute substantially to

the biodiversity of urban blue-green infrastructure, the role of garden ponds is still largely

unkown. We lack a comprehensive understanding of how local habitat features, different

forms of management, and urbanisation might impact the biodiversity of these habitats. This

study aimed to reveal the importance of garden ponds via a country-wide online citizen

science survey in Hungary, Central Europe. Data from over 800 pond owners revealed the

occurrence and local frequency of various native animal taxa (amphibians, odonates, and

birds), and introduced animals (e.g., fish). We collected data about pond features and

management practices. We tested the effect of pond features, pond management, and

landscape-level drivers (urbanisation, surrounding wetland coverage) on the presence of

conspicuous animal taxa (adult amphibians and tadpoles, birds, odonates) to identify the

potential drivers of the biodiversity of garden ponds. Key pond features including pond age,

area, aquatic and shoreline vegetation were the most important factors, while algaecide

addition was the most influential management practice negatively affecting amphibian

presence. Urbanisation negatively affected the presence of adult amphibians and their

tadpoles, but it was not associated negatively with the presence of odonates and birds. Our

results indicate the high potential to utilise garden ponds as urban habitats surveyed with the

help of the public. Developing effective urban biodiversity monitoring and conservation

strategies are necessary for a better functioning blue-green infrastructure. The high level of

engagement of pond owners, as in our survey, can create valuable data for achieving these

aims.

1. Introduction

Land use changes including the intensification of agricultural activities, industrial

development, and urbanisation are major drivers of species loss (Storch et al., 2022; Yang et

al., 2024). Among them, urbanisation is a complex driver of global biodiversity loss

(Concepción, 2021; Danneck et al., 2023; McDonald et al., 2019), acting through habitat loss,

fragmentation, and degradation (Grooten & Almond, 2018; Stamenković et al., 2019; Tickner

et al., 2020). It can create strongly selective environments e.g. via pollution (chemical light,

noise), physical barriers obstructing organisms' movement (roads and buildings), often

leading to a limited species pool in urban habitats (Capotorti et al., 2023; Hill et al., 2021;

Schueller et al., 2023). At the same time, urban habitats can also act as refuges for

biodiversity depending on habitat planning and management (Soanes et al., 2019, 2023).

Therefore, it is increasingly important to understand the role of local habitat features and

management measures of the blue-green infrastructure in our efforts to efficiently conserve

and increase urban biodiversity.

Ponds are increasingly recognised as key aquatic habitats, often supporting high levels

of biodiversity within a region or landscape and hosting threatened or endangered species

(Céréghino et al., 2007; Williams et al., 2004). Lentic aquatic habitats less than 5 m deep and

ranging 1 m^2 and 2-5 ha in size are generally considered ponds (Biggs et al., 2005; Céréghino

et al., 2007; Richardson et al., 2022). It has been estimated that freshwater ponds (< 1 km^2)

cover more than 3% of the Earth's surface, and constitute the most widespread type of inland

standing waters (Downing et al., 2006). They often represent the most abundant aquatic

habitats locally in natural landscapes and even in areas with high agricultural activity (Oertli

et al., 2009; Ruggiero et al., 2008).

Garden ponds are popular features that can provide a range of ecosystem services

including microclimate regulation, supporting biodiversity, and cultural services such as

wildlife observation or connecting with nature (Hill et al., 2021; Oertli et al., 2023; Peeters et

al., 2023). Yet, very limited information exists about the exact number and ecological roles of

garden ponds in cities (Hill et al., 2021). Garden ponds can host threatened taxa like

amphibians (Hassall, 2014), which are among the most vulnerable freshwater organisms

worldwide (Hamer & McDonnell, 2008). At the same time, garden ponds may also act as a

source of invasive species due to commercial trade, limited knowledge, and aesthetic

preference thereby threatening native biodiversity (Patoka et al., 2017). To promote efficient

management practices that support native species while minimizing the facilitation of

invasion events, we need better understanding how current management practices affect the

wildlife of these urban habitats. Currently, this represents a general knowledge gap likely

linked to the novelty of this ecosystem type in urban blue-green infrastructure and also their

limited accessibility to researchers.

Citizen (or community) science, i.e., the involvement of members of the public in

scientific research and data production, is becoming a valued and necessary research tool for

scientists to evaluate or gain knowledge about a wide range of scientific topics (Fraisl et al.,

2022). Data collected by citizens and later harmonized and validated by experts can greatly

improve our understanding of species diversity at large spatial and temporal scales as well as

ecological dynamics and the conservation status of species (Steven et al., 2019). Garden

ponds are difficult to access or sample as they are situated on private properties. With the

involvement of their owners, citizen science data collection can not only be achievable, but is

essentially the only feasible way to collect large-scale datasets from these ecosystems. Hence,

for a better understanding of the biodiversity, ecology, and management of garden ponds,

citizen science is an essential tool, which can at the same time lead to a high level of public

engagement (Hamer et al., 2024; Peeters et al., 2023).

To improve our understanding on the ecological roles of garden ponds, we conducted

a countrywide online survey in Central Europe (Hungary). The specific aims of our study

were to 1) explore how frequently wild animals visit or inhabit garden ponds focusing on

taxonomic groups that are conspicuous enough to be easily and reliably identified by the

public (amphibians and their tadpoles, birds, odonates); 2) investigate the frequency of animal

introductions, including potentially invasive ones, to garden ponds (i.e. animals such as fish,

mussels, snails or crayfish introduced by the pond owner); 3) identify the potential drivers of

native biota by testing the reported presence of animals against urbanisation. local pond

features and management practices applied by the owners (for the latter, data was available

from Hamer et al., 2024), with the expectation that urbanisation will have an overall negative

effect on the presence of most taxa.

2. Materials and Methods

2.1. Citizen science survey

An online survey within the citizen science project "MyPond" was launched in 2021

to investigate the biodiversity, local features, and management practices of garden ponds in

Hungary (Central Europe). The link to the questionnaire on the "MyPond" website

(https://mypond.hu/en/) was distributed via traditional media outlets and social media. The

survey consisted of a total of 30 questions: 6 multiple choices, 10 short answers, and 14

closed (yes-no) questions. We collected data on the local features of ponds, including age,

area, substrate of the pond (concrete, rubber pond liner, plastic, and natural, the latter

including soil, gravel, clay etc.), and the presence/absence of aquatic vegetation, shoreline

vegetation, and fish. Local management data included circulation, cleaning, draining, leaf

removal, and use of algaecide. The results from these management data and local pond

features were summarised and published by Hamer et al., 2024.

The biodiversity dataset consisted of four animal taxa, amphibians (here, we refer to

adult amphibians or post-metamorphic juvenile sightings) and their larvae referred to as

tadpoles (reported separately from adult amphibians), birds, and odonates (we used the

Hungarian term „szitakötő" which does not differentiate between damselfiles and

dragonflies). The survey contained a link to the amphibian identification key (illustrated with

drawings) of the BirdLife Hungary to help the citizens correctly identify amphibian species

(MME (2022). https://www.mme.hu/en/node/4440), which is available as a free smartphone

application. The identification key has been developed especially for citizen science

monitoring and it guides a non-scientist volunteer through the most important features

(including shape, colour, and habitat) of amphibians, resulting in species-level identification.

Also, there was an option to name the species of birds they had seen in their gardens.

2.2. Data validation and processing

The questionnaire is still available on the website and collects data for future purposes.

Between 2021 and 2022 (6 July 2021 - 9 September 2022), 834 participants filled out the

questionnaire, of which a validated subset of 753 responses was used for statistical analysis

(Figure 1). Validation was carried out to exclude duplicates and incomplete data. Duplicates

were identified as having the same geolocation or email address provided by the owners.

However, respondents may have multiple ponds in their garden, therefore the length, width

and depth were also inspected before exclusion. If the pond sizes were roughly the same, it

was considered a duplicate. We also excluded ponds with an area over 200 m^2, which we set

as a threshold based on the frequency distribution of pond areas (the largest was 24000 m^2,

but 98% ponds were smaller than 200 m^2). We approximated the pond surface area based on

the lengths and widths (that pond owners provided), by the multiplication of these

measurements. Pond age was used as 0 if the pond was created in the year when the survey

was filled out. In some cases, pond age was missing from the dataset, since the previous

homeowner constructed the pond, and the current owner did not hold specific information on

the exact year of construction. If they provided the information of when they moved in, we

calculated the pond age as the number of years since then.

In the case of the amphibian species dataset, we excluded amphibian species if pond

owners made statements such as "we have not seen that species in a few years". However, we

included species if the owner reported them as "rarely" seen. Furthermore, entries were

excluded from the amphibian species dataset if it only included the name "frog", "toad",

"newt", or "tadpoles", as species identity could not be determined. "Green frogs" were

grouped into the *Pelophylax* spp. complex as the three species that comprise this group are

morphologically difficult to separate, resulting in an amphibian dataset of 332 entries. Rare

species were also cross-checked with the most up-to-date species distribution maps

(https://herpterkep.mme.hu/index.php?lang=en) and retained only if their occurrence had

already been proven within the region of the sighting. Regarding birds, due to the high

number of potential bird species, we decided to group the reported birds from the open-ended

question (Table S1) into larger categories: songbirds, wading birds, swallows, woodpeckers,

ducks, and pigeons. Here, songbirds included thrushes, sparrows, finches, starlings, tits, larks,

and nuthatches while wading birds included herons, egrets and storks. These larger categories

can be still reliably identified by the public and were actually used by many of the participants

in their original responses to this question where they could not fully identify the species they

had seen. For the analysis on the potential drivers of occurrence, we further grouped these

data into three main categories according to their pond use type (see below). Data were

excluded if respondents did not specify the species or categories, resulting in a dataset of 661

entries. The full questionnaire can be found in the supplementary information (Table S1).

The resulting datasets included both occurrence (binary; 0, 1) and local frequency

(ordinal scale; 0, 1, 2, 3) data of each recorded animal taxa (amphibians and their tadpoles,

odonates, and birds). Pond features (pond age, area, substrate, aquatic vegetation, shoreline

vegetation, and fish) and management (cleaning, draining, leaf removal, and the use of

algaecides) were used as binomial and ratio scale data in the statistical analysis (Table 1).

2.3. Landscape-level drivers: urbanisation and wetland coverage data

The area of surrounding urban land and wetland coverage, predicted to be landscape-

level drivers of species occurrence, were calculated using the Ecosystem Map of Hungary

(project KEHOP-430-VEKOP-15-2016-00001, Ministry of Agriculture, 2019). The wetland

coverage category contained pixels of wetlands, marshlands, temporary wetlands, standing

waters, and flowing waters, while the urban land category (hereafter urbanisation) contained

pixels of tall buildings, short buildings, sealed roads, dirt roads, railways, other artificial

surfaces, and urban green spaces with and without trees. The number of raster pixels (20x20

m) was counted within a 1 km radius around each focal pond using the Zonal Histogram tool

(QGIS Development Team, 2022). A 1 km buffer was chosen as a widely used measure for

studying birds (Humphrey et al., 2023) and it also contains the mean and maximum dispersal

distances of most amphibian species expected to occur in Hungary (Kovar et al., 2009). While

we acknowledge that this might have created some overlap in the individual buffer zones of

some of the ponds (mostly in Budapest where ponds were situated closer to each other), given

the average pond distance of 3514 m in the dataset, it still provided a wide and largely

independent gradient of landscape cover for most of the studied ponds. While a few ponds

were less than 1 km apart, resulting in partially overlapping buffer zones, this was only the

case in 0.09% of all the pond pairs, hence the effect of this should have been minimal. The

classification of land coverage was used as defined by the Ecosystem Map of Hungary

(Ministry of Agriculture, 2019). Urbanisation and wetland coverage were then calculated as

the proportion of total pixels within the 1 km radius covered by the respective land cover type

(Oertli & Parris, 2019). All spatial analysis was carried out in QGIS 3.28.1 (QGIS

Development Team, 2022).

2.4. Statistical analysis

To visualise the occurrence and the local frequency of animal taxa, we created

barplots and pie charts using the "ggplot2" R package (Wickham, 2016).

To examine the relationship between amphibian community composition and pond

features, pond management, and landscape-level drivers (urbanisation and wetland coverage),

we performed canonical correspondence analysis (CCA) using the "vegan" R package

(Oksanen, 2017). Prior to the CCA analysis, we excluded ponds with no amphibian species

recorded, and the analysis was based on the amphibian species occurrence data (binomial). To

normalize the distribution of variables belonging to pond features and landscape-level drivers,

we used log transformation for pond area, and square root transformation was applied for

wetland coverage prior to the CCA. Pond management variables were included as binomial

data. Significant explanatory variables among pond features, management, and landscape-

level drivers were identified based on an automatic stepwise (backward and forward) model

selection procedure using permutation tests ("ordistep" function, direction = "both", n = 999

permutations).

To analyse which pond features, management types, or landscape-level drivers had the

strongest effect on the occurrence of each four investigated animal taxa (amphibians,

tadpoles, birds and odonates; Table 1), generalised linear models (GLM) were fitted with a

binomial distribution and logit link function (hereinafter logistic regression model). Model

selection was carried out based on Akaike's Information Criteria (AIC) with the "MASS" R

package (Venables & Ripley, 2002). In the case of birds, we also ran separate models where

bird categories were additionally grouped into three main groups according to the way they

use the ponds: wading birds (herons, egrets, and storks), ducks, and other birds (mostly

including songbirds that mainly use the ponds for drinking). To test for potential spatial

autocorrelation in the resulting final GLMs, we used the Moran's I test from the „DHARMA"

R package (Hartig, 2022). As no significant autocorrelation was found (Table S2), we kept the

original GLM models for the analyses on the effects of pond features, management types, and

landscape-level drivers. Visualisation was carried out with the "ggplot2" (Wickham, 2016)

and "ggbreak" (Xu et al., 2021) R packages.

All statistical analyses and illustrations were made with the R program v 4.2.2 (R Core

Team, 2022).

3. Results

Based on the occurrence data reported by 753 respondents, odonates (in 96.5% of the

ponds) and birds (96.5%) were observed at almost all garden ponds. While the occurrence of

amphibian sightings (81.4%) was higher than that of tadpoles, the latter were still observed in

close to two-thirds of the ponds (62.1%). Among the amphibian species, *Pelophylax* spp. was

the most frequently reported species (29.9%). The most frequently reported birds were

songbirds (88.3%, including tits, blackbirds, sparrows, etc.). Within the introduced animal

taxa, fish were the most frequent (84.4%), while snails, turtles, mussels, and crayfish were all

introduced to less than 6% of the ponds (Figure 2).

Local frequency data of amphibians, birds, and odonates revealed that most (60%,

78.5%, and 86.5%, respectively) of the respondents regularly see these animal taxa in their

ponds. Among them, tadpoles were absent from the highest share of ponds (37.8%), but

amphibian breeding was regularly observed in over 40% of the ponds detected via tadpole

presence (Figure 3).

Participants recorded the presence of 13 amphibian species in garden ponds: three

newt species (*Ichthyosaura alpestris*, *Lissotriton vulgaris*, *Triturus dobrogicus*), nine anurans

(*Bombina bombina*, *Bufo bufo*, *Bufotes viridis*, *Hyla arborea*, *Pelobates fuscus*, *Pelophylax*

spp., *Rana arvalis*, *R. dalmatina*, *R. temporaria*) and one salamander (*Salamandra*

salamandra). Amphibian species composition (based on occurrence data, Table 1) was

significantly influenced by pond age (p = 0.050), pond area (p = 0.043), rubber pond liner (p

= 0.034), shoreline vegetation (p = 0.072), algaecide usage (p = 0.077), the introduction of

fish (p = 0.001) (Figure 4). There was a negative association between *Rana arvalis* and pond

age, i.e., it occurred mostly in newly installed ponds with rubber pond liner and shoreline

vegetation. *Lissotriton vulgaris*, *Rana temporaria,* and *R. dalmatina* responded negatively to

fish presence and algaecide treatment. *Bufotes viridis* was positively associated with

urbanisation. *Salamandra salamandra*, *Ichthyosaura alpestris, Triturus dobrogicus, Bombina*

bombina, Hyla arborea, and *Pelobates fuscus* were positively associated with large, well-

vegetated ponds with a natural substrate, which are only occasionally drained (Figure 4).

Based on the results of the logistic regression models, urbanisation was negatively

associated with the presence of amphibians and their tadpoles, while it generally had a

positive effect on the presence of odonates and birds (Figure 5, Table S3). Wetland coverage

did not have a significant effect in any of the models, although a positive trend was found for

amphibians (Figure 5, odds ratio (OR) = 34.300, p = 0.109). The most frequently significant

pond feature was pond age (logistic regression model of amphibian: OR = 0.256, p < 0.001;

tadpoles: OR = 0.351, p < 0.001; odonates: OR = 0.154, p < 0.001; birds: OR = 0.155, p <

0.001), followed by rubber pond liner (amphibians: OR = 2.150, p = 0.004; tadpoles: OR =

1.997, p = 0.003; odonates: OR = 2.510, p = 0.048; birds: OR = 2.510, p = 0.048; wading

birds: OR = 2.655, p = 0.021) and pond area (amphibians: OR = 1.036, p = 0.001; tadpoles:

OR = 1.014, p = 0.001). Amphibians and their tadpoles, odonates, and birds were less likely

to be present in or at newly installed ponds (0-1 year). Pond area had a positive, while

algaecides had a negative, effect on the presence of both amphibians and their tadpoles. There

was a positive effect of aquatic vegetation on tadpoles, odonates and birds, however, shoreline

vegetation was only significant for the presence of tadpoles (Figure 5, Table S3).

4. Discussion

Our citizen science survey provided comprehensive large-scale data on the presence of

multiple animal taxa of garden ponds, covering an entire European country. Odonates and

birds were observed at almost all 753 garden ponds, but amphibians and their tadpoles were

also common. Local frequency data revealed that most respondents regularly see these animal

taxa. Urbanisation had an overall negative effect on the presence of amphibians and tadpoles;

conversely, odonates and birds showed an unexpected positive association with it. The age of

the ponds was a general predictor, with animals less likely to be present in newly installed

ponds. Among local pond features, pond area and rubber pond liner were the most important

across the animal taxa, mostly with positive effects. Among management types, algaecide

addition had a negative effect on amphibians and their tadpoles. Our data also highlighted that

owners introduce various animal taxa into their ponds, most frequently fish, with potential

consequences for the native biodiversity surrounding garden ponds. Our results overall

illustrated the potentially important role of garden ponds in urban biodiversity and

conservation by providing secondary habitats for multiple animal taxa.

<u>4.1. Effect of urbanisation</u>

Urbanisation had disparate effects across the animal taxa we investigated. It had a

negative effect on the occurrence (presence/absence) of amphibians and their tadpoles, in line

with our original hypothesis and also corroborating earlier findings (Hamer & Parris, 2011;

Rubbo & Kiesecker, 2005). Amphibians are particularly vulnerable to urbanisation and

fragmented landscapes, comprised of roads, houses and buildings that can impede dispersal

between ponds and natural upland habitats that are important for amphibians to complete their

complex life cycle (Hamer & McDonnell, 2008; Scheffers & Paszkowski, 2012). Although

urbanisation generally has a negative effect on amphibian diversity, some species are urban

adapters (McKinney, 2002), favored by larger body sizes and the ability to persist even in

fragmented landscapes (Hahs et al., 2023). For example, *Bufotes viridis* was positively

associated with urbanisation in our study. This species has a relatively large body size and

persists in the centre of European cities and towns (Landler et al., 2023; Mazgajska &

Mazgajski, 2020; Sistani et al., 2021). The *Pelophylax* spp. complex had no discernible

associations with any of the variables, probably because these species are habitat generalists

and can maintain metapopulations in both natural and urban landscapes (Herczeg et al., 2012;

Ficetola and De Bernardi, 2004). Conversely, species such as *Ichthyosaura alpestris* which is

associated with natural ponds and is particularly vulnerable to landscape fragmentation due to

urbanisation (Emaresi et al., 2011; Van Buskirk, 2012), were only recorded in the Bakony

Mountains, a protected forest in western Hungary. *Salamandra salamandra* can persist in

low-density urban areas provided there is sufficient forest nearby. Accordingly, it was

reported from a garden pond adjacent to a large forested area (Bükk Mountains in northern

Hungary) and interestingly, a population exists even at the fringe of the capital city Budapest

(Kiss et al., 2022a).

Conversely, occurrences of odonates and birds were positively associated with

urbanisation in general. These results might seem counterintuitive at first, although there is

evidence that in certain cases urban areas are able to maintain higher odonate species diversity

than agricultural lands, due to a network of public, stormwater, or garden ponds that can

foster the migration of species (Goertzen & Suhling, 2019; Willigalla & Fartmann, 2012).

However, other studies showed that species richness of odonates increased from the city

center to more rural areas, probably due to increasing habitat availability (more water bodies

in the rural areas than in cities) (Holtmann et al., 2018; Willigalla & Fartmann, 2012). Here,

we only recorded the presence of odonates as a group. Therefore, the observed positive

association could also be explained by a few highly tolerant species occurring in urbanised

landscapes, but more detailed follow-up studies will be necessary to link this pattern to

species identities and biodiversity. The same might be true for birds, resulting from the

presence of more generalist bird species such as blackbirds or sparrows, which are common

even in heavily urbanised landscapes (Callaghan et al., 2019; Mennechez & Clergeau, 2006).

Furthermore, birds are more likely to visit garden ponds, puddles, or fountains, in urban areas,

where other natural water sources are scarce (Tryjanowski et al., 2022), which could have

contributed to the pattern we found.

<u>4.2. The most influential pond features</u>

Pond age was overall the most important pond feature shaping the occurrence of

animal taxa. Newly installed ponds (0-1 years) were less occupied by amphibians and their

tadpoles, odonates and birds, which can be due to the lack of vegetation and sediment that

could offer hiding and breeding places to aquatic and semi-aquatic animal taxa (Biggs et al.,

2005; Huikkonen et al., 2020; Soukup et al., 2022). Additionally, colonisation time might be

another factor as animals need time to find and occupy newly installed ponds in the urban

landscape (Cayuela et al., 2020; Coccia et al., 2016; Unglaub et al., 2021). For example, many

odonate species have longer life cycles, so adult life stages may only reach higher densities at

older ponds (Hassall & Thompson, 2008; Stoks & Córdoba-Aguilar, 2012), hence pond age

may increase encounter probabilities with them. Most amphibians and their tadpoles were

significantly associated with older ponds, corresponding to a longer period for species to

locate and colonise these ponds (Moor et al., 2022). Among them, *Rana arvalis* was an

exception, being strongly associated with newly installed ponds that this species can

potentially colonise while moving relatively long distances through the landscape (Brzeziński

& Mętrak, 2014). Many amphibian species prefer well-vegetated or large ponds (Arntzen et

al., 1997; Péntek et al., 2017; Rannap et al., 2013), which was also supported by our results as

Bombina bombina, Pelobates fuscus, Triturus dobrogicus were positively associated with

aquatic vegetation. Aquatic vegetation was also positively associated with tadpoles, odonates,

and bird presence which indicates the habitat structuring rola of aquatic vegetation that

benefits biodiversity (Ansari et al., 2017; Law et al., 2019). Unexpectedly, rubber pond liner

had a significantly positive effect on the presence of amphibians and their tadpoles, and also

on odonates, birds, and wading birds. This may have resulted from the generally larger size of

the rubber-lined ponds as shown by an earlier study (Hamer et al., 2024).

4.3. Management of garden ponds

The use of chemicals, such as pesticides, fertilizers, and algaecides is common around

ponds in agricultural and recreational settings like golf courses, but they can also be applied in

private gardens to improve soil and water quality or control biological processes (Boyd &

Massaut, 1999; Oertli & Parris, 2019). Even though pesticides are intended to target specific

pests, they can affect other non-target organisms via overland runoff, potentially impacting

nearby habitats (Rumschlag et al., 2020; Yamamuro et al., 2019). However, private gardens,

including those with ponds, are characterized by relatively lower levels of pesticide use, due

to which they can even serve as refuges for pollinators (Hall et al., 2017; Siviter et al., 2023).

Furthermore, unlike public ponds that often receive polluted surface runoff (Ivanovsky et al.,

2018; Osman et al., 2019), garden ponds are usually isolated from such events. As a result,

fertilizers and insecticides applied elsewhere in the garden have limited potential to

contaminate the ponds. Additionally, insecticides are also typically applied on garden crops,

and therefore they represent minimal risk to garden pond fauna, as these are usually not

applied on aquatic plants such as water lilies and sedges.

On the other hand, the direct application of algaecides was a common practice among

Hungarian garden pond owners according to our survey (reported from 33% of the ponds).

Our findings imply a negative effect of algaecides on amphibians and their tadpoles. At the

same time, our multivariate analysis showed a strong association between algaecide use and

the presence of fish, and both were negatively aligned with almost all amphibian species.

Hence the negative effect of algaecides on the presence and reproduction of amphibians might

be also linked to the presence of fish in the same ponds. High nutrient inputs in the form of

fish food and a low abundance of aquatic macrophytes, which are competitors of

phytoplankton (Zeng et al., 2022), contributes to high algal production that could

subsequently lead to algaecide application in these ponds. While we did not collect specific

information on the exact types of algaecide used, several respondents, on their own accord,

indicated a specific brand of algae managing substance which uses a natural approach (i.e.

microbial decomposition) rather than artificial chemicals, which points more at the indirect

link through fish.

4.4. Invasion risk of the introduced animals and the role of fish

Based on direct communication with the pond owners, exotic animals are frequently

introduced to garden ponds for ornamental purposes (e.g. goldfish, koi) (Hassall, 2014).

According to our results, fish in general were the most frequently introduced animals, being

consequently present in 84% of ponds. In comparison to our results, studies conducted in the

UK and the Netherlands reported that only ~35-32% of the garden pond owners stocked their

ponds with fish (Loram et al., 2011; Peeters et al., 2023). Besides fish, turtles and several

invertebrates (snails, mussels, and crayfish) were also reported. Previous studies found that

aquaria and garden ponds may act as sources of invasions (C. Chucholl, 2013; Patoka et al.,

2014; Soes et al., 2011; Van Den Neucker et al., 2017), for example in the case of the red-

eared terrapin (*Trachemys scripta elegans*) (Tietz et al., 2023), or signal crayfish

(*Pacifastacus leniusculus*) (F. Chucholl et al., 2021). In this sense, without the indispensable

knowledge and education about invasive species, garden pond owners and garden centres can

pose a threat to native biodiversity. This highlights the importance of outreach activities

targeting pond owners, encouraging more conscious management choices.

We also found that the introduction of fish was negatively associated with all recorded

amphibian species. Fish are important predators of all amphibian life stages and a negative

relationship between fish and amphibian populations has been demonstrated for many

European amphibians (Hartel et al., 2007; Van Buskirk, 2012). But despite the introduction of

fish into 84% of ponds, amphibian reproduction was observed in 62% of the ponds, indicating

that some species are able to coexist with fish (Hamer et al., 2021; Kloskowski et al., 2020).

Aquatic vegetation can provide refugia for amphibians from fish predation (Hartel et al.,

2007), which could be also observed in our dataset, where well-vegetated ponds supported a

higher co-existence of fish and amphibians (73.5%) than poorly vegetated ponds (54%). As

the introduction of fish was a significant predictor of garden pond design, within-pond

features, and management practices in these ponds (Hamer et al., 2024), this result implies a

potential conflict between creating ponds with the main aim of keeping ornamental fish and

their potential to support native biodiversity. However, our results imply that planting dense

aquatic vegetation can mitigate the negative effect of fish on amphibians in garden ponds.

Also, wading birds were frequently (24.3%) reported from the ponds, the majority of

which were stocked with fish (18.1% of the ponds in our survey). Sightings in this group

included grey herons, which regularly utilise garden ponds for foraging on fish (Jakubas &

Manikowska, 2011). This group represents a source of human-nature conflict as, according to

anecdotal evidence and comments given as part of the responses in our survey, they can

damage ponds by puncturing pond liners and prey upon ornamental fish.

4.5. Caveats and relevance of using citizen science data in garden pond surveys

We identified a number of caveats related to the use of citizen science data as part of

our study. Due to reliability issues, we lost 9.7% of the survey responses in the first step of the

validation (excluding duplicates and incomplete data). In the case of amphibian species and

bird types, even with our best efforts to validate the dataset, there is a chance of

misidentification. To overcome this, we collected data on taxa that can easily be identified,

and are considered typical groups representing pond biodiversity (Oertli et al., 2002). We also

provided   an   identification   key   for   amphibian   species   (MME   2022.

https://www.mme.hu/en/node/4440) along with our survey. Another unexpected caveat was

the unit of pond size for example, where some respondents did the conversion of centimeter to

meter or meter to centimeter incorrectly. In these cases, we manually checked the pond size

with Google Maps based on GPS coordinates. Although we can not compare our

questionnaire data to earlier or other regional datasets as a reference, the high number of

ponds in our study provides robust data for the general patterns observed in these poorly

studied secondary habitats. These could later serve as important baselines for more

environmentally friendly management activities, being more beneficial for the local pond

biota.

Increasing pressure for urban development poses a significant threat to blue-green

infrastructure, reducing its extent and complicating its management (Ancillotto et al., 2019).

Citizen science can provide important means for efficient large-scale biodiversity monitoring,

particularly in cases where access to private properties is limited, as was the case in our study.

The great potential of the approach is well-illustrated by the high level of engagement in the

present survey, with over 800 completed forms. Also, citizen science emerges as a powerful

tool for urban planning, as it can contribute to gathering valuable data on urban biodiversity

and utilise it for more efficient conservation strategies. For instance, it could help urban

planning directly by identifying hotspots of aquatic biodiversity or critical areas for the

conservation of key groups like amphibians in urban environments. Amphibians are one of

the central targets for pond creation projects in Europe (De Necker et al., 2024) and there is

increasing evidence that urban ponds, including garden ponds, can contribute to their

conservation in the urban landscape by offering valuable secondary habitats and potentially

acting as stepping stones between larger habitats (Butterworth et al., 2025; Hamer et al., 2021;

Hamer & Parris, 2011; Kiss et al., 2022b). Citizen science data could also provide a feasible

way to assess the green infrastructure of cities and could be used for the long-term monitoring

of the potential effects of management activities on urban biodiversity (Callaghan et al.,

2020). In our study, we collected a large-scale biodiversity dataset of garden ponds, which

could potentially help to identify ponds or larger areas hosting protected species. Such

programs may help authorities to make informed decisions on planning and managing the

blue-green infrastructure to support the biodiversity of protected species, for example by

creating new ponds in parks near to garden ponds with the known occurrence of target

species. Information on the location of these new connecting elements of aquatic habitat

networks can aid the formation of corridors increasing the connectivity between larger aquatic

habitats in cities (Hyseni et al., 2021). Without the active engagement of citizens, such

information would not be available for city planners. Finally, active citizen engagement not

only contributes to large-scale data collection but it is also a tool for environmental education.

Participants may become more conscious of environmental issues and their role in it which

might lead to more active engagement in supporting blue-green infrastructure development.

5. Conclusions

450   To date, there has been limited information available on the biodiversity of garden ponds

451   worldwide. In this study, we present the first large-scale dataset evaluating the effects of

452   urbanisation, local features, and management practices on garden pond biodiversity using

453   responses from 753 pond owners in Hungary. Our results highlighted the importance of

454   garden ponds for urban biodiversity and conservation, as they serve as secondary habitats for

455 multiple animal taxa in urban environments. This implication can be directly communicated

456 to the public while promoting pond installations in urban areas, where the importance of

457 aquatic vegetation can be emphasised. These activities, however, should also highlight

458 relevant facts about invasive species and the most suitable local features and management

459 types to maximise local biodiversity. For effective biodiversity conservation in urban areas,

460 where scarce natural habitats are under constant pressure from degradation and

461 mismanagement, public engagement is becoming increasingly valuable. Citizen science not

462 only provides essential tools for gathering data but could be utilised to further engage citizens

463 in their crucial role as pond managers.

Declaration of competing interest

464 The authors declare that they have no known competing financial interests or personal

465 relationships that could have appeared to influence the work reported in this paper.

References

467 Ancillotto, L., Bosso, L., Salinas-Ramos, V. B., & Russo, D. (2019). The importance of ponds

468   for the conservation of bats in urban landscapes. *Landscape and Urban Planning*, *190*,

469   103607. https://doi.org/10.1016/j.landurbplan.2019.103607

470 Ansari, A. A., Shalini Saggu, S. S., Al-Ghanim, S. M., Abbas, Z. K., Gill, S. S., Khan, F. A.,

471   Dar, M. I., Naikoo, M. I., & Khan, A. A. (2017). Aquatic plant biodiversity: A

472   biological indicator for the monitoring and assessment of water quality. In A. A.

473   Ansari, S. S. Gill, Z. K. Abbas, & M. Naeem (Eds.), *Plant biodiversity: Monitoring,*

474   *assessment and conservation* (1st ed., pp. 218–227). CABI.

475   https://doi.org/10.1079/9781780646947.0218

Arntzen, J. W., Bugter, R. J. F., Cogalniceanu, D., & Wallis, G. P. (1997). The distribution

and conservation status of the Danube crested newt, Triturus dobrogicus. *Amphibia-*

Reptilia, *18*(2), 133–142. https://doi.org/10.1163/156853897X00026

Biggs, J., Williams, P., Whitfield, M., Nicolet, P., & Weatherby, A. (2005). 15 years of pond

assessment in Britain: Results and lessons learned from the work of Pond

Conservation. *Aquatic Conservation: Marine and Freshwater Ecosystems*, *15*(6), 693–

714. https://doi.org/10.1002/aqc.745

Boyd, C. E., & Massaut, L. (1999). Risks associated with the use of chemicals in pond

aquaculture. *Aquacultural Engineering*, *20*(2), 113–132.

https://doi.org/10.1016/S0144-8609(99)00010-2

Brzeziński, M., & Mętrak, M. (2014). Spring migration rates and community structure of

amphibians breeding in an old and newly established midfield ponds. *Folia Zoologica*,

63(3), 161–170. https://doi.org/10.25225/fozo.v63.i3.a3.2014

Butterworth, H. L., Van Helden, B. E., & Close, P. G. (2025). From ponds to patios: Evidence

of frog reproduction and recruitment in residential gardens. *Urban Ecosystems*, *28*(1),

1–13. https://doi.org/10.1007/s11252-024-01653-8

Callaghan, C. T., Major, R. E., Wilshire, J. H., Martin, J. M., Kingsford, R. T., & Cornwell,

W. K. (2019). Generalists are the most urban☐tolerant of birds: A phylogenetically

controlled analysis of ecological and life history traits using a novel continuous

measure of bird responses to urbanization. *Oikos*, *128*(6), 845–858.

https://doi.org/10.1111/oik.06158

Callaghan, C. T., Ozeroff, I., Hitchcock, C., & Chandler, M. (2020). Capitalizing on

opportunistic citizen science data to monitor urban biodiversity: A multi-taxa

framework. *Biological Conservation*, *251*, 108753.

https://doi.org/10.1016/j.biocon.2020.108753

Capotorti, G., Valeri, S., Giannini, A., Minorenti, V., Piarulli, M., & Audisio, P. (2023). On

the Role of Natural and Induced Landscape Heterogeneity for the Support of

Pollinators: A Green Infrastructure Perspective Applied in a Peri-Urban System. *Land*,

12(2), 387. https://doi.org/10.3390/land12020387

Cayuela, H., Valenzuela-Sánchez, A., Teulier, L., Martínez-Solano, Í., Léna, J.-P., Merilä, J.,

Muths, E., Shine, R., Quay, L., Denoël, M., Clobert, J., & Schmidt, B. R. (2020).

Determinants and Consequences of Dispersal in Vertebrates with Complex Life

Cycles: A Review of Pond-Breeding Amphibians. https://doi.org/10.5167/UZH-185243

Céréghino, R., Biggs, J., Oertli, B., & Declerck, S. (2007). The ecology of European ponds:

Defining the characteristics of a neglected freshwater habitat. In B. Oertli, R.

Céréghino, J. Biggs, S. Declerck, A. Hull, & M. R. Miracle (Eds.), *Pond Conservation*

in Europe (pp. 1–6). Springer Netherlands. https://doi.org/10.1007/978-90-481-9088-

1_1

Chucholl, C. (2013). Invaders for sale: Trade and determinants of introduction of ornamental

freshwater crayfish. *Biological Invasions*, *15*(1), 125–141.

https://doi.org/10.1007/s10530-012-0273-2

Chucholl, F., Fiolka, F., Segelbacher, G., & Epp, L. S. (2021). eDNA Detection of Native and

Invasive Crayfish Species Allows for Year-Round Monitoring and Large-Scale

Screening of Lotic Systems. *Frontiers in Environmental Science*, *9*, 639380.

https://doi.org/10.3389/fenvs.2021.639380

Coccia, C., Vanschoenwinkel, B., Brendonck, L., Boyero, L., & Green, A. J. (2016). Newly

created ponds complement natural waterbodies for restoration of macroinvertebrate

assemblages. *Freshwater Biology*, *61*(10), 1640–1654.

https://doi.org/10.1111/fwb.12804

Concepción, E. D. (2021). Urban sprawl into Natura 2000 network over Europe. *Conservation*

Biology, *35*(4), 1063–1072. https://doi.org/10.1111/cobi.13687

Danneck, J., Chin Fung Tiang, D., Ngo Duc, K., Azhar, B., Sanusi, R., Campos☐Arceiz, A.,

& Lechner, A. M. (2023). Conserving urban tropical biodiversity by connecting

networks of green patches. *Integrative Conservation*, inc3.21.

https://doi.org/10.1002/inc3.21

De Necker, L., Brendonck, L., Vanschoenwinkel, B., Florencio, M., Rhazi, L., & Gołdyn, B.

(2024). Lessons from pond creation and restoration projects in Europe. *Restoration*

Ecology, e14342. https://doi.org/10.1111/rec.14342

Downing, J. A., Prairie, Y. T., Cole, J. J., Duarte, C. M., Tranvik, L. J., Striegl, R. G.,

McDowell, W. H., Kortelainen, P., Caraco, N. F., Melack, J. M., & Middelburg, J. J.

(2006). The global abundance and size distribution of lakes, ponds, and

impoundments. *Limnology and Oceanography*, *51*(5), 2388–2397.

https://doi.org/10.4319/lo.2006.51.5.2388

Emaresi, G., Pellet, J., Dubey, S., Hirzel, A. H., & Fumagalli, L. (2011). Landscape genetics

of the Alpine newt (Mesotriton alpestris) inferred from a strip-based approach.

Conservation Genetics, *12*(1), 41–50. https://doi.org/10.1007/s10592-009-9985-y

Ficetola, F., & De Bernardi, F. (2004). Amphibians in a human-dominated landscape: The

community structure is related to habitat features and isolation. *Biological*

Conservation, *119*(2), 219–230. https://doi.org/10.1016/j.biocon.2003.11.004

Fraisl, D., Hager, G., Bedessem, B., Gold, M., Hsing, P.-Y., Danielsen, F., Hitchcock, C. B.,

Hulbert, J. M., Piera, J., Spiers, H., Thiel, M., & Haklay, M. (2022). Citizen science in

environmental and ecological sciences. *Nature Reviews Methods Primers*, *2*(1), 64.

https://doi.org/10.1038/s43586-022-00144-4

Goertzen, D., & Suhling, F. (2019). Urbanization versus other land use: Diverging effects on

dragonfly communities in Germany. *Diversity and Distributions*, *25*(1), 38–47.

https://doi.org/10.1111/ddi.12820

Grooten, M., & Almond, R. E. A. (Eds.). (2018). *Living planet report 2018: Aiming higher*.

WWF--World Wide Fund for Nature.

Hahs, A. K., Fournier, B., Aronson, M. F. J., Nilon, C. H., Herrera-Montes, A., Salisbury, A.

B., Threlfall, C. G., Rega-Brodsky, C. C., Lepczyk, C. A., La Sorte, F. A.,

MacGregor-Fors, I., Scott MacIvor, J., Jung, K., Piana, M. R., Williams, N. S. G.,

Knapp, S., Vergnes, A., Acevedo, A. A., Gainsbury, A. M., … Moretti, M. (2023).

Urbanisation generates multiple trait syndromes for terrestrial animal taxa worldwide.

Nature Communications, *14*(1), 4751. https://doi.org/10.1038/s41467-023-39746-1

Hall, D. M., Camilo, G. R., Tonietto, R. K., Ollerton, J., Ahrné, K., Arduser, M., Ascher, J. S.,

Baldock, K. C. R., Fowler, R., Frankie, G., Goulson, D., Gunnarsson, B., Hanley, M.

E., Jackson, J. I., Langellotto, G., Lowenstein, D., Minor, E. S., Philpott, S. M., Potts,

S. G., … Threlfall, C. G. (2017). The city as a refuge for insect pollinators.

Conservation Biology, *31*(1), 24–29. https://doi.org/10.1111/cobi.12840

Hamer, A. J., Barta, B., Bohus, A., Gál, B., & Schmera, D. (2021). Roads reduce amphibian

abundance in ponds across a fragmented landscape. *Global Ecology and*

Conservation, *28*, e01663. https://doi.org/10.1016/j.gecco.2021.e01663

Hamer, A. J., Barta, B., Márton, Z., Vad, C. F., Szabó, B., Tornero, I., & Horváth, Z. (2024).

Patterns and correlates in the distribution, design and management of garden ponds

along an urban–rural gradient. *Urban Ecosystems*, *27*(5), 1915–1930.

https://doi.org/10.1007/s11252-024-01559-5

572 Hamer, A. J., & McDonnell, M. J. (2008). Amphibian ecology and conservation in the

573  urbanising world: A review. *Biological Conservation*, *141*(10), 2432–2449.

574  https://doi.org/10.1016/j.biocon.2008.07.020

575 Hamer, A. J., & Parris, K. M. (2011). Local and landscape determinants of amphibian

576  communities in urban ponds. *Ecological Applications*, *21*(2), 378–390.

577  https://doi.org/10.1890/10-0390.1

578 Hartel, T., Nemes, S., Cogălniceanu, D., Öllerer, K., Schweiger, O., Moga, C.-I., & Demeter,

579  L. (2007). The effect of fish and aquatic habitat complexity on amphibians.

580  *Hydrobiologia*, *583*(1), 173–182. https://doi.org/10.1007/s10750-006-0490-8

581 Hartig, F. (2022). DHARMa: Residual diagnostics for hierarchical (multi-level/mixed)

582  regression models [Manual].

583 Hassall, C. (2014). The ecology and biodiversity of urban ponds. *WIREs Water*, *1*(2), 187–

584  206. https://doi.org/10.1002/wat2.1014

585 Hassall, C., & Thompson, D. J. (2008). The effects of environmental warming on Odonata: A

586  review. *International Journal of Odonatology*, *11*(2), 131–153.

587  https://doi.org/10.1080/13887890.2008.9748319

588 Herczeg, D., Gallé, R., & Molnár, N. 2012. Effects of urbanization on the occurrence of

589  Anura assemblages in the city of Szeged (Hungary). Tiscia, 39, 3-8.

590 Hill, M. J., Greaves, H. M., Sayer, C. D., Hassall, C., Milin, M., Milner, V. S., Marazzi, L.,

591  Hall, R., Harper, L. R., Thornhill, I., Walton, R., Biggs, J., Ewald, N., Law, A.,

592  Willby, N., White, J. C., Briers, R. A., Mathers, K. L., Jeffries, M. J., & Wood, P. J.

593  (2021). Pond ecology and conservation: Research priorities and knowledge gaps.

594  *Ecosphere*, *12*(12). https://doi.org/10.1002/ecs2.3853

595 Holtmann, L., Juchem, M., Brüggeshemke, J., Möhlmeyer, A., & Fartmann, T. (2018).

596  Stormwater ponds promote dragonfly (Odonata) species richness and density in urban

areas. *Ecological Engineering*, *118*, 1–11.

https://doi.org/10.1016/j.ecoleng.2017.12.028

Huikkonen, I., Helle, I., & Elo, M. (2020). Heterogenic aquatic vegetation promotes

abundance and species richness of Odonata (Insecta) in constructed agricultural

wetlands. *Insect Conservation and Diversity*, *13*(4), 374–383.

https://doi.org/10.1111/icad.12396

Hyseni, C., Heino, J., Bini, L. M., Bjelke, U., & Johansson, F. (2021). The importance of blue

and green landscape connectivity for biodiversity in urban ponds. *Basic and Applied*

Ecology, *57*, 129–145. https://doi.org/10.1016/j.baae.2021.10.004

Ivanovsky, A., Belles, A., Criquet, J., Dumoulin, D., Noble, P., Alary, C., & Billon, G.

(2018). Assessment of the treatment efficiency of an urban stormwater pond and its

impact on the natural downstream watercourse. *Journal of Environmental*

Management, *226*, 120–130. https://doi.org/10.1016/j.jenvman.2018.08.015

Jakubas, D., & Manikowska, B. (2011). The response of Grey Herons *Ardea cinerea* to

changes in prey abundance. *Bird Study*, *58*(4), 487–494.

https://doi.org/10.1080/00063657.2011.608423

Kiss, I., Vörös, J., & Hamer, A. J. (2022a). Larval development and habitat usage of stream-

breeding Fire salamanders in an urban environment. *Acta Zoologica Academiae*

Scientiarum Hungaricae, *68*(4), 321–340.

https://doi.org/10.17109/AZH.68.4.321.2022

Kiss, I., Vörös, J., & Hamer, A. J. (2022b). Movement patterns within an urban population of

fire salamanders highlight the importance of conserving small habitat patches. *Journal*

of Zoology, *316*(4), 240–249. https://doi.org/10.1111/jzo.12949

Kloskowski, J., Nieoczym, M., & Stryjecki, R. (2020). Between-habitat distributions of pond

tadpoles and their insect predators in response to fish presence. *Hydrobiologia*, *847*(5),

1343–1356. https://doi.org/10.1007/s10750-020-04190-5

Kovar, R., Brabec, M., Bocek, R., & Vita, R. (2009). Spring migration distances of some

Central European amphibian species. Amphibia-Reptilia, 30(3), 367–378.

https://doi.org/10.1163/156853809788795236

Landler, L., Burgstaller, S., & Schweiger, S. (2023). Land-use preferences of the European

green toad (Bufotes viridis) in the city of Vienna (Austria): The importance of open

land in urban environments. *Frontiers in Zoology*, *20*(1), 3.

https://doi.org/10.1186/s12983-022-00480-x

Law, A., Baker, A., Sayer, C., Foster, G., Gunn, I. D. M., Taylor, P., Pattison, Z., Blaikie, J.,

& Willby, N. J. (2019). The effectiveness of aquatic plants as surrogates for wider

biodiversity in standing fresh waters. *Freshwater Biology*, *64*(9), 1664–1675.

https://doi.org/10.1111/fwb.13369

Loram, A., Warren, P., Thompson, K., & Gaston, K. (2011). Urban Domestic Gardens: The

Effects of Human Interventions on Garden Composition. *Environmental Management*,

48(4), 808–824. https://doi.org/10.1007/s00267-011-9723-3

Mazgajska, J., & Mazgajski, T. D. (2020). Two amphibian species in the urban environment:

Changes in the occurrence, spawning phenology and adult condition of common and

green toads. *The European Zoological Journal*, *87*(1), 170–179.

https://doi.org/10.1080/24750263.2020.1744743

McDonald, R. I., Mansur, A. V., Ascensão, F., Colbert, M., Crossman, K., Elmqvist, T.,

Gonzalez, A., Güneralp, B., Haase, D., Hamann, M., Hillel, O., Huang, K., Kahnt, B.,

Maddox, D., Pacheco, A., Pereira, H. M., Seto, K. C., Simkin, R., Walsh, B., … Ziter,

C. (2019). Research gaps in knowledge of the impact of urban growth on biodiversity.

Nature Sustainability, *3*(1), 16–24. https://doi.org/10.1038/s41893-019-0436-6

McKinney, M. L. (2002). Urbanization, Biodiversity, and Conservation. *BioScience*, *52*(10),

883. https://doi.org/10.1641/0006-3568(2002)052[0883:UBAC]2.0.CO;2

Mennechez, G., & Clergeau, P. (2006). Effect of urbanisation on habitat generalists: Starlings

not so flexible? *Acta Oecologica*, *30*(2), 182–191.

https://doi.org/10.1016/j.actao.2006.03.002

Ministry of Agriculture, 2019. Ecosystem base map of Hungary. DOI:

10.34811/osz.alapterkep

MME (2022). Amphibian and Reptile identification key. https://www.mme.hu/en/node/4440

Moor, H., Bergamini, A., Vorburger, C., Holderegger, R., Bühler, C., Egger, S., & Schmidt,

B. R. (2022). Bending the curve: Simple but massive conservation action leads to

landscape-scale recovery of amphibians. *Proceedings of the National Academy of*

Sciences, *119*(42), e2123070119. https://doi.org/10.1073/pnas.2123070119

Oertli, B., Céréghino, R., Hull, A., & Miracle, R. (2009). Pond conservation: From science to

practice. *Hydrobiologia*, *634*(1), 1–9. https://doi.org/10.1007/s10750-009-9891-9

Oertli, B., Decrey, M., Demierre, E., Fahy, J. C., Gallinelli, P., Vasco, F., & Ilg, C. (2023).

Ornamental ponds as Nature-based Solutions to implement in cities. *Science of The*

Total Environment, *888*, 164300. https://doi.org/10.1016/j.scitotenv.2023.164300

Oertli, B., Joye, D. A., Castella, E., Juge, R., Cambin, D., & Lachavanne, J.-B. (2002). Does

size matter? The relationship between pond area and biodiversity. *Biological*

Conservation, *104*(1), 59–70. https://doi.org/10.1016/S0006-3207(01)00154-9

Oertli, B., & Parris, K. M. (2019). Review: Toward management of urban ponds for

freshwater biodiversity. *Ecosphere*, *10*(7), e02810. https://doi.org/10.1002/ecs2.2810

Oksanen, J. (2017). Vegan: Ecological diversity. In *R Package Version 2.4-4*.

Osman, M., Wan Yusof, K., Takaijudin, H., Goh, H. W., Abdul Malek, M., Azizan, N. A.,

Ab. Ghani, A., & Sa'id Abdurrasheed, A. (2019). A Review of Nitrogen Removal for

Urban Stormwater Runoff in Bioretention System. *Sustainability*, *11*(19), 5415.

https://doi.org/10.3390/su11195415

Patoka, J., Bláha, M., Kalous, L., & Kouba, A. (2017). Irresponsible vendors: Non-native,

invasive and threatened animals offered for garden pond stocking: Non-native and

threatened animals in garden ponds. *Aquatic Conservation: Marine and Freshwater*

Ecosystems, *27*(3), 692–697. https://doi.org/10.1002/aqc.2719

Patoka, J., Petrtýl, M., & Kalous, L. (2014). Garden ponds as potential introduction pathway

of ornamental crayfish. *Knowledge and Management of Aquatic Ecosystems*, *414*, 13.

https://doi.org/10.1051/kmae/2014019

Peeters, E. T. H. M., Wilhelm, M., Gerritsen, A. A. M., & Seelen, L. M. S. (2023). Ecological

characterisation of urban ponds in the Netherlands: A study based on data collected by

volunteers. *Frontiers of Biogeography*, *15*(1). https://doi.org/10.21425/F5FBG57816

Péntek, A. L., Vad, C. F., Zsuga, K., & Horváth, Z. (2017). Metacommunity dynamics of

amphibians in years with differing rainfall. *Aquatic Ecology*, *51*(1), 45–57.

https://doi.org/10.1007/s10452-016-9597-9

QGIS Development Team, 2022. QGIS Geographic Information System. Open Source

Geospatial Foundation Project. http://qgis.osgeo.org

Rannap, R., Markus, M., & Kaart, T. (2013). Habitat use of the common spadefoot toad

(Pelobates fuscus) in Estonia. *Amphibia-Reptilia*, *34*(1), 51–62.

https://doi.org/10.1163/15685381-00002868

Richardson, D. C., Holgerson, M. A., Farragher, M. J., Hoffman, K. K., King, K. B. S.,

Alfonso, M. B., Andersen, M. R., Cheruveil, K. S., Coleman, K. A., Farruggia, M. J.,

Fernandez, R. L., Hondula, K. L., López Moreira Mazacotte, G. A., Paul, K., Peierls,

B. L., Rabaey, J. S., Sadro, S., Sánchez, M. L., Smyth, R. L., & Sweetman, J. N.

(2022). A functional definition to distinguish ponds from lakes and wetlands.

Scientific Reports, *12*(1), 10472. https://doi.org/10.1038/s41598-022-14569-0

Rubbo, M. J., & Kiesecker, J. M. (2005). Amphibian Breeding Distribution in an Urbanized

Landscape. *Conservation Biology*, *19*(2), 504–511. https://doi.org/10.1111/j.1523-

1739.2005.000101.x

Ruggiero, A., Céréghino, R., Figuerola, J., Marty, P., & Angélibert, S. (2008). Farm ponds

make a contribution to the biodiversity of aquatic insects in a French agricultural

landscape. *Comptes Rendus Biologies*, *331*(4), 298–308.

https://doi.org/10.1016/j.crvi.2008.01.009

Rumschlag, S. L., Mahon, M. B., Hoverman, J. T., Raffel, T. R., Carrick, H. J., Hudson, P. J.,

& Rohr, J. R. (2020). *Consistent Effects of Pesticides on Community Structure and*

Ecosystem Function in Freshwater Systems.

https://doi.org/10.1101/2020.02.03.932442

Scheffers, B. R., & Paszkowski, C. A. (2012). The effects of urbanization on North American

amphibian species: Identifying new directions for urban conservation. *Urban*

Ecosystems, *15*(1), 133–147. https://doi.org/10.1007/s11252-011-0199-y

Schueller, S. K., Li, Z., Bliss, Z., Roake, R., & Weiler, B. (2023). How Informed Design Can

Make a Difference: Supporting Insect Pollinators in Cities. *Land*, *12*(7), 1289.

https://doi.org/10.3390/land12071289

Sistani, A., Burgstaller, S., Gollmann, G., & Landler, L. (2021). The European green toad,

Bufotes viridis, in Donaufeld (Vienna, Austria): Status and size of the population.

Herpetozoa, *34*, 259–264. https://doi.org/10.3897/herpetozoa.34.e75578

Siviter, H., Pardee, G. L., Baert, N., McArt, S., Jha, S., & Muth, F. (2023). Wild bees are

exposed to low levels of pesticides in urban grasslands and community gardens.

Science of The Total Environment, *858*, 159839.

https://doi.org/10.1016/j.scitotenv.2022.159839

Soanes, K., Sievers, M., Chee, Y. E., Williams, N. S. G., Bhardwaj, M., Marshall, A. J., &

Parris, K. M. (2019). Correcting common misconceptions to inspire conservation

action in urban environments. *Conservation Biology*, *33*(2), 300–306.

https://doi.org/10.1111/cobi.13193

Soanes, K., Taylor, L., Ramalho, C. E., Maller, C., Parris, K., Bush, J., Mata, L., Williams, N.

S. G., & Threlfall, C. G. (2023). Conserving urban biodiversity: Current practice,

barriers, and enablers. *Conservation Letters*, *16*(3), e12946.

https://doi.org/10.1111/conl.12946

Soes, M., Majoor, G., & Keulen, S. (2011). Bellamya chinensis (Gray, 1834) (Gastropoda:

Viviparidae), a new alien snail species for the European fauna. *Aquatic Invasions*,

6(1), 97–102. https://doi.org/10.3391/ai.2011.6.1.12

Soukup, P. R., Näslund, J., Höjesjö, J., & Boukal, D. S. (2022). From individuals to

communities: Habitat complexity affects all levels of organization in aquatic

environments. *WIREs Water*, *9*(1), e1575. https://doi.org/10.1002/wat2.1575

Stamenković, O., Stojković Piperac, M., Milošević, D., Buzhdygan, O. Y., Petrović, A.,

Jenačković, D., Đurđević, A., Čerba, D., Vlaičević, B., Nikolić, D., & Simić, V.

(2019). Anthropogenic pressure explains variations in the biodiversity of pond

communities along environmental gradients: A case study in south-eastern Serbia.

Hydrobiologia, *838*(1), 65–83. https://doi.org/10.1007/s10750-019-03978-4

Steven, R., Barnes, M., Garnett, S. T., Garrard, G., O'Connor, J., Oliver, J. L., Robinson, C.,

Tulloch, A., & Fuller, R. A. (2019). Aligning citizen science with best practice:

Threatened species conservation in Australia. *Conservation Science and Practice*,

1(10). https://doi.org/10.1111/csp2.100

Stoks, R., & Córdoba-Aguilar, A. (2012). Evolutionary Ecology of Odonata: A Complex Life

Cycle Perspective. *Annual Review of Entomology*, *57*(1), 249–265.

https://doi.org/10.1146/annurev-ento-120710-100557

Storch, D., Šímová, I., Smyčka, J., Bohdalková, E., Toszogyova, A., & Okie, J. G. (2022).

Biodiversity dynamics in the Anthropocene: How human activities change equilibria

of species richness. *Ecography*, *2022*(4), ecog.05778.

https://doi.org/10.1111/ecog.05778

Tickner, D., Opperman, J. J., Abell, R., Acreman, M., Arthington, A. H., Bunn, S. E., Cooke,

S. J., Dalton, J., Darwall, W., Edwards, G., Harrison, I., Hughes, K., Jones, T.,

Leclère, D., Lynch, A. J., Leonard, P., McClain, M. E., Muruven, D., Olden, J. D., …

Young, L. (2020). Bending the Curve of Global Freshwater Biodiversity Loss: An

Emergency Recovery Plan. *BioScience*, *70*(4), 330–342.

https://doi.org/10.1093/biosci/biaa002

Tietz, B., Penner, J., & Vamberger, M. (2023). Chelonian challenge: Three alien species from

North America are moving their reproductive boundaries in Central Europe. *NeoBiota*,

82, 1–21. https://doi.org/10.3897/neobiota.82.87264

Tryjanowski, P., Jankowiak, Ł., Czechowski, P., Dulisz, B., Golawski, A., Grzywaczewski,

G., Indykiewicz, P., Kwieciński, Z., Mitrus, C., Nowakowski, J. J., Polakowski, M.,

Siekiera, J., Sparks, T. H., & Wuczyński, A. (2022). Summer water sources for

temperate birds: Use, importance, and threats. *The European Zoological Journal*,

89(1), 913–926. https://doi.org/10.1080/24750263.2022.2101699

Unglaub, B., Cayuela, H., Schmidt, B. R., Preißler, K., Glos, J., & Steinfartz, S. (2021).

Context‐dependent dispersal determines relatedness and genetic structure in a patchy

amphibian population. *Molecular Ecology*, *30*(20), 5009–5028.

https://doi.org/10.1111/mec.16114

Van Buskirk, J. (2012). Permeability of the landscape matrix between amphibian breeding

sites. *Ecology and Evolution*, *2*(12), 3160–3167. https://doi.org/10.1002/ece3.424

Van Den Neucker, T., Schildermans, T., & Scheers, K. (2017). The invasive Chinese mystery

snail *Bellamya chinensis* (Gastropoda: Viviparidae) expands its European range to

Belgium. *Knowledge & Management of Aquatic Ecosystems*, *418*, 8.

https://doi.org/10.1051/kmae/2016040

Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S*. Springer New

York. https://doi.org/10.1007/978-0-387-21706-2

Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis. In *Journal of the Royal*

Statistical Society: Series A (Statistics in Society).

Williams, P., Whitfield, M., Biggs, J., Bray, S., Fox, G., Nicolet, P., & Sear, D. (2004).

Comparative biodiversity of rivers, streams, ditches and ponds in an agricultural

landscape in Southern England. *Biological Conservation*, *115*(2), 329–341.

https://doi.org/10.1016/S0006-3207(03)00153-8

Willigalla, C., & Fartmann, T. (2012). Patterns in the diversity of dragonflies (Odonata) in

cities across Central Europe. *European Journal of Entomology*, *109*(2), 235–245.

https://doi.org/10.14411/eje.2012.031

Xu, S., Chen, M., Feng, T., Zhan, L., Zhou, L., & Yu, G. (2021). Use ggbreak to Effectively

Utilize Plotting Space to Deal With Large Datasets and Outliers. *Frontiers in*

Genetics, *12*, 774846. https://doi.org/10.3389/fgene.2021.774846

Yamamuro, M., Komuro, T., Kamiya, H., Kato, T., Hasegawa, H., & Kameda, Y. (2019).

Neonicotinoids disrupt aquatic food webs and decrease fishery yields. *Science*,

366(6465), 620–623. https://doi.org/10.1126/science.aax3442

Yang, L., Xu, H., Pan, S., Chen, W., & Zeng, J. (2024). Identifying the impact of global

human activities expansion on natural habitats. *Journal of Cleaner Production*, *434*,

140247. https://doi.org/10.1016/j.jclepro.2023.140247

Zeng, Q., Wei, Z., Yi, C., He, Y., & Luo, M. (2022). The effect of different coverage of

aquatic plants on the phytoplankton and zooplankton community structures: A study

based on a shallow macrophytic lake. *Aquatic Ecology*, *56*(4), 1347–1358.

https://doi.org/10.1007/s10452-022-09970-w

Web references

https://mypond.hu/en/ - Last accessed: 2023.11.23.

https://herpterkep.mme.hu/index.php?lang=en - Last accessed: 2023.11.23.

List of tables

Table 1 Data types in the validated dataset

Main variable groups	Recorded variables	Data type
Biodiversity - occurrence	**amphibians and their tadpoles, birds, odonates**	Binomial (0, 1)
Biodiversity - local frequency	**amphibians and their tadpoles, birds, odonates**	Ordinal (0, 1, 2, 3, corresponding to no, singular, occasional, and frequent sightings per pond)
Pond management	**circulation, cleaning, draining, leaf removal, algaecide**	Binomial (0, 1)
Pond features	**age, area, substrate (concrete, rubber, plastic, natural), aquatic vegetation, shoreline vegetation, fish presence**	Ratio scale and binomial (0,1) data

List of figures

Figure 1 Location of the 753 garden ponds in Hungary used in the analyses. (Map was created with R v. 4.2.2)

Figure 2 Percentage of all ponds with the occurrence of the animal taxa reported by the citizens. Samples sizes were n=667 in case of types of birds (top right), and n=753 in the other three datasets. For the abbreviation of amphibian species (bottom left figure), see the legend of Figure 4.

Figure 3 Share (%) of each response on the local frequency of animal taxa (amphibians and their tadpoles, odonates and birds) in garden ponds (n=753)

Figure 4 Canonical correspondence analysis (CCA) biplot based on the reported prevalence of amphibian species, environmental predictors, and level of urbanisation (red points: species scores, dark blue arrows: significant predictors, light green arrows: all other predictors). The analysis is based on data from 332 ponds with validated species data. (Abbreviated species names: *I. alpestris = Ichthyosaura alpestris*, *L. vulgaris = Lissotriton vulgaris*, *T. dobrogicus = Triturus dobrogicus*, *B. bombina = Bombina bombina*, *B. bufo = Bufo bufo*, *B. viridis = Bufotes viridis*, *H. arborea = Hyla arborea*, *P. fuscus = Pelobates fuscus*, *R. arvalis = Rana arvalis*, *R. dalmatina = Rana dalmatina*, *R. temporaria = Rana temporaria*, *S. salamandra = Salamandra salamandra*)

Figure 5 Summary of the best-fit logistic regression models (based on AIC) odds ratio (filled circles) and their 95% confidence intervals (whiskers) testing the relationship between the occurrence of animal taxa and pond features and management. Samples sizes were n=753 for

amphibians, odonates, birds, and tadpoles, and n=667 for other birds and wading birds.

(Birds: all birds; Wading birds: herons, egrets and storks; Other birds: songbirds, pigeons,

woodpeckers

Figure 1 Location of the 753 garden ponds in Hungary used in the analyses. (Map was created with R

v. 4.2.2)

Figure 2 Percentage of all ponds with the occurrence of the animal taxa reported by the citizens.

Samples sizes were n=667 in case of types of birds (top right), and n=753 in the other three datasets.

For the abbreviation of amphibian species (bottom left figure), see the legend of Figure 4.

Figure 3 Share (%) of each response on the local frequency of animal taxa (amphibians and their

tadpoles, odonates and birds) in garden ponds (n=753)

Figure 4 Canonical correspondence analysis (CCA) biplot based on the reported prevalence of amphibian species, environmental predictors, and level of urbanisation (red points: species scores, dark blue arrows: significant predictors, light green arrows: all other predictors). The analysis is based on data from 332 ponds with validated species data. (Abbreviated species names: *I. alpestris = Ichthyosaura alpestris*, *L. vulgaris = Lissotriton vulgaris*, *T. dobrogicus = Triturus dobrogicus*, *B. bombina = Bombina bombina*, *B. bufo = Bufo bufo*, *B. viridis = Bufotes viridis*, *H. arborea = Hyla arborea*, *P. fuscus = Pelobates fuscus*, *R. arvalis = Rana arvalis*, *R. dalmatina = Rana dalmatina*, *R. temporaria = Rana temporaria*, *S. salamandra = Salamandra salamandra*)

Figure 5 Summary of the best-fit logistic regression models (based on AIC) odds ratio (filled circles) and their 95% confidence intervals (whiskers) testing the relationship between the occurrence of animal taxa and pond features and management. Samples sizes were n=753 for amphibians, odonates, birds, and tadpoles, and n=667 for other birds and wading birds. (Birds: all birds; Wading birds: herons, egrets and storks; Other birds: songbirds, pigeons, woodpeckers

List of appendices

Table S1 Questions of the "MyPond" online survey

Table S2 Summary statistics of the Moran's I spatial autocorrelation tests used for each GLM model listed in Table S3

Table S3 Summary statistics of the best-fit logistic regression models (based on AIC) testing the relationship between the prevalence of animal groups and pond features and management

Table S1 Questions of the "MyPond" online survey

Questions	Answer options
What kind of animals are there in the pond? (Check all that apply!)	Fish Turtle None Other
Do you see odonates* around the pond?	Yes, often Yes, but only rarely Yes, I've seen them once No
Have you seen amphibians in the pond (e.g. frog, newt, etc.)?	Yes, often Yes, but only rarely Yes, I've seen them once No
If so, what kind?	
Have you seen tadpoles in the pond?	Yes, often Yes, but only rarely Yes, I've seen them once No

Do birds visit the pond?	Yes, often Yes, but only rarely Yes, I've seen them once No
If you have seen them, can you name the type of bird?	
How old is your garden pond?	Made this year 1 year old 2 years old 3 years old 4 years old 5 years old 6 years old 7 years old 8 years old 9 years old 10 years old Older than 10 years I do not remember Made by the previous owner, I bought it with the house

If your answer to the previous question was that you had bought the house with an existing pond, please write approx. how many years ago you bought the house!	
What is the pond's length?	
What is the pond's width?	
Please check the appropriate answer regarding the substrate of the pond! (Check all that apply!)	Rubber pond liner Concrete Plastic basin Other
Did you plant any plants in the garden pond?	Yes, on the shoreline (sedge, iris, etc.) Yes, in the water (water lily, pondweed, etc.) In both places No
Do you drain the water from the pond?	Yes No
Do you clean the pond bed, e.g. to remove sediment?	Yes No

Do you use any other treatments? (Check all that apply!)	Yes, algaecide
	Yes, heating
	Yes, I remove the leaves
	Yes, other
	No
Do you use any equipment to circulate the water (e.g. filter, fountain)?	Yes
	No

*In the case of „odonates", we used the Hungarian term „szitakötő" which does not differentiate between damselfiles and dragonflies.

Table S2 Summary statistics of the Moran's I spatial autocorrelation tests used for each GLM model listed in Table S3

Group	Observed	Expected	sd	p value
Amphibians	0.007	-0.001	0.009	0.355
Tadpoles	0.002	-0.001	0.009	0.461
Odonates	0.011	-0.001	0.009	0.196
Birds	0.011	-0.001	0.009	0.196
Ducks	0.001	-0.001	0.009	0.765
Other birds	0.005	-0.001	0.009	0.464
Wading birds	0.012	-0.001	0.009	0.167

Table S3 Summary statistics of the best-fit logistic regression models (based on AIC) testing the relationship between the prevalence of animal groups and pond features and management

Group	Final model elements	Estimate	Std.Error	z value	p value	F value	Odds ratio	Confidence Interval (95%)
							Odds ratio results	
Amphibians	Age 0-1	-1.261	0.258	-5.262	<0.001	28.830	0.256	0.154-0.426
	Area	0.036	0.011	3.310	0.001	19.426	1.036	1.016-1.061
	Rubber pond liner	0.765	0.266	2.871	0.004	8.644	2.150	1.269-3.618
	Natural substrate	14.442	528.959	0.027	0.978	4.195	1.872E+06	0.000-2.157E+75
	Algaecide	-0.421	0.208	-2.024	0.043	4.368	0.656	0.436-0.988
	Urbanisation	-1.956	0.436	4.487	<0.001	23.693	0.141	0.058-0.326
	Wetland coverage	3.535	2.207	1.601	0.109	3.160	34.300	0.621-3810.007
Tadpoles	Age 0-1	-1.045	0.233	4.483	<0.001	19.557	0.351	0.221-0.553
	Area	0.014	0.004	3.312	0.001	14.316	1.014	1.006-1.024
	Concrete substrate	0.471	0.321	1.468	0.142	2.101	1.601	0.862-3.048
	Rubber pond liner	0.692	0.234	2.954	0.003	8.344	1.997	1.264-3.171
	Algaecide	-0.631	0.167	3.756	<0.001	13.412	0.532	0.382-0.739
	Shoreline vegetation	1.120	0.481	2.326	0.020	5.723	3.066	1.243-8.407

	Aquatic vegetation	0.457	0.222	2.056	0.040	3.987	1.580	1.020-2.446
	Urbanisation	-1.236	0.315	-3.916	<0.001	15.010	0.290	0.155-0.536
Odonates	Age 0-1	-1.866	0.438	-4.257	<0.001	17.374	0.154	0.065-0.368
	Rubber pond liner	0.920	0.465	1.979	0.048	3.785	2.510	0.969-6.118
	Shoreline vegetation	16.047	1102.637	0.015	0.988	4.847	9.321E+06	3.921E-19-9.744E+94
	Aquatic vegetation	1.281	0.461	2.779	0.005	7.443	3.599	1.423-8.833
	Urbanisation	1.236	0.748	1.651	0.099	2.891	3.443	0.799-15.355
Birds	Age 0-1	-1.866	0.438	-4.257	<0.001	17.374	0.155	0.065-0.368
	Rubber pond liner	0.920	0.465	1.979	0.048	3.785	2.510	0.9697-6.118
	Shoreline vegetation	16.047	1102.637	0.015	0.988	4.847	9.321E+06	3.921E-19-9.744E+94
	Aquatic vegetation	1.281	0.461	2.779	0.005	7.443	3.599	1.423-8.833
	Urbanisation	1.236	0.748	1.651	0.099	2.891	3.443	0.799-15.355
Ducks	Urbanisation	-0.667	0.447	-1.492	0.136	2.209	0.512	0.212-1.235
Other birds	Cleaning	0.708	0.328	2.158	0.031	4.754	2.031	1.073-3.925
	Urbanisation	1.239	0.612	2.023	0.043	4.131	3.454	1.044-11.638
Wading birds	Area	-0.006	0.004	-1.513	0.131	2.673	0.993	0.984-1.001
	Rubber pond liner	1.011	0.424	2.382	0.017	6.579	2.751	1.254-6.716

	Plastic substrate	1.104	0.445	2.479	0.013	6.539	3.01	1.288-7.513
	Draining	-0.284	0.198	-1.433	0.152	2.087	0.752	0.506-1.105
	Circulation	-0.341	0.209	-1.632	0.102	2.605	0.711	0.473-1.076