Effects of urbanisation, habitat characteristics, and management on garden pond

biodiversity: a citizen science approach

**Authors**

Zsuzsanna Márton[1,2*], Barbara Barta[1,2,3], Csaba F. Vad[1,2], Beáta Szabó[1,2], Andrew J. Hamer[1,2],

Vivien Kardos[1], Csilla Laskai[1,2], Ádám Fierpasz[1,3], Zsófia Horváth[1,2,4]

[1]Institute of Aquatic Ecology, HUN-REN Centre for Ecological Research, Karolina út 29,

1113 Budapest Hungary

[2]National Multidisciplinary Laboratory for Climate Change, HUN-REN Centre for

Ecological Research, Karolina út 29, 1113 Budapest Hungary

[3]Doctoral School of Biology, ELTE Eötvös Loránd University, Pázmány Péter sétány 1/c,

1113 Budapest, Hungary

[4]ELTE Eötvös Loránd University, Pázmány Péter sétány 1/c, 1113 Budapest, Hungary

**Corresponding author**

Zsuzsanna Márton, HUN-REN Institute of Aquatic Ecology, Centre for Ecological Research,

Budapest, Karolina út 29, 1113 Budapest, Hungary

Email: marton.zsuzsanna@ecolres.hu

**Highlights**

20    ● Over 800 garden pond owners participated in a citizen science project in Hungary.

21    ● Garden ponds can host flagship animal groups, e.g., amphibians, odonates, and birds.

22    ● Pond age, area, and vegetation influenced the occurrence of these animal groups.

23    ● Urbanisation had a negative effect on amphibian communities.

24    ● Algaecide application played a key role in the presence of multiple animal groups.

**Abstract**

The rapid expansion of urban areas leads to loss, fragmentation, and degradation of natural habitats, posing a serious threat to biodiversity. While urban ponds might contribute substantially to the biodiversity of urban blue-green infrastructure, the role of garden ponds, one of their most numerous representatives, is still little known. We lack a comprehensive understanding of how local habitat features, different forms of management, and urbanisation might impact the biodiversity of these habitats. This study aimed to reveal the importance of garden ponds via a country-wide online citizen science survey in Hungary, Central Europe. Based on data from over 800 pond owners, we gained knowledge about the prevalence and local frequency of various native animal groups, including amphibians, odonates, and birds, as well as information regarding introduced animals, such as fish. Additionally, we collected data about pond features and management practices. We tested the effect of pond features, pond management, and landscape-level drivers (urbanisation, surrounding wetland coverage) on the presence of conspicuous animal groups (adult amphibians, birds, odonates, and tadpoles) along with species of adult amphibians to identify the potential drivers of the biodiversity of garden ponds. Key pond features like pond age, area, aquatic and shoreline vegetation were the most important factors, while algaecide addition was the most influential management practice negatively affecting the presence of amphibians. Urbanisation had a

negative effect on the presence of adult amphibians and their tadpoles, but surprisingly, it was

not negatively associated with the presence of odonates and birds. Our results demonstrate the

vital role of garden ponds in urban biodiversity conservation, most notably by providing

secondary habitats and contributing to amphibian conservation, as evidenced by the

occurrence of 13 species. Emphasizing public involvement is key in developing effective

urban biodiversity conservation strategies and recognizing citizens' potential contributions to

a better functioning blue-green infrastructure.

**Graphical abstract**

**Graphical abstract:** Summary of the key results from the MyPond survey. Arrows indicate

significant relationships (blue: positive, red: negative ) between the presence of animal groups and

pond features and management practices)

**Keywords**

Citizen science, Community science, Garden ponds, Urban ponds, Biodiversity, Pond management

## 1. Introduction

Urbanisation is accelerating worldwide, being one of the major drivers of the reshuffling of global biodiversity acting through habitat loss, fragmentation, and degradation (Grooten and Almond, 2018; Stamenković et al., 2019; Tickner et al., 2020). Previous studies have shown that urbanisation often has a negative impact on biodiversity (Concepción, 2021; Danneck et al., 2023; McDonald et al., 2019). It can also create strongly selective environments, leading to a limited species pool in urban compared to natural habitats (Capotorti et al., 2023; Hill et al., 2021; Schueller et al., 2023). Species that can withstand the effects of urbanisation might adapt via genetic (Brans et al., 2018; Brans and De Meester, 2018; Johnson and Munshi-South, 2017) and behavioral changes (Ma et al., 2023; Rivera-Gutierrez et al., 2023; Uchida et al., 2020; Yin et al., 2023), which can foster population persistence and even successful colonization of the habitats in the blue-green infrastructure of urban areas. Consequently, urban habitats can also act as refuges and host considerable biodiversity depending on habitat planning and management (Soanes et al., 2023, 2019). Therefore, it is increasingly important to understand the role of local habitat features and management measures of blue-green infrastructure in our efforts to efficiently conserve and increase urban biodiversity.

Ponds are increasingly recognised as key aquatic habitats, by contributing over-proportionally to regional biodiversity and hosting threatened or endangered species (Céréghino et al., 2007; Williams et al., 2004). While there is no officially recognized international categorization, lentic aquatic habitats that are less than 5 m deep and are between

1 m$^2$ and up to 2-5 ha in size are generally considered ponds (Biggs et al., 2005; Céréghino et

al., 2007; Richardson et al., 2022). There is still much uncertainty about the global number of

ponds, but it has been estimated that small lakes (< 1 km$^2$) and ponds cover more than 3% of

the Earth's surface, and constitute the most widespread type of inland standing waters

(Downing et al., 2006). They often represent the locally most abundant aquatic habitats in

natural landscapes and even in areas under high agricultural activity (Oertli et al., 2009;

Ruggiero et al., 2008), being also among the most numerous components of the blue-green

infrastructure in cities (Oertli et al., 2023).

Garden ponds are popular garden features, which can be numerous in urban areas.

They are created for various reasons, including aesthetic values, microclimate regulation,

keeping ornamental animals, or supporting biodiversity (Oertli et al., 2023). Garden ponds

have the potential to offer many ecosystem services, and small ponds are a way to connect to

nature, observe wildlife, and also to increase the blue-green surface in urban areas (Peeters et

al., 2023). Still, there is only scarce information about the exact number of garden ponds in

cities (Hill et al., 2021), and no standard protocol has been established regarding their

management to support high biodiversity. Although garden ponds are small, their overall

contribution to biodiversity can be quite important (Biggs et al., 2017). For example, they can

host threatened taxa like amphibians (Hassall, 2014), which are among the most vulnerable

freshwater organisms worldwide (Hamer and McDonnell, 2008). However, due to

commercial trade and a lack of knowledge of many exotic species, garden ponds may also act

as a source of invasive species thereby threatening native biodiversity (Patoka et al., 2017).

For efficient management both suitable for the native species and not facilitating such

invasion events, a better understanding of the current management practices and their

connection with the wildlife of these secondary habitats would be essential.

Citizen (or community) science, i.e., the involvement of citizens in scientific research

and data production, is becoming a valued and necessary research tool for scientists to

evaluate or gain knowledge about a wide range of scientific topics (Fraisl et al., 2022),

ranging from archeology (Smith, 2014) through ecology (Johansson et al., 2020; Oertli et al.,

2023; Peeters et al., 2023) to astrophysics (Memarsadeghi et al., 2022). Citizen science has

many shortcomings, such as the lack of standardized methods in the study design or unreliable

data collection without expert knowledge. Still, data collected by citizens and later

harmonized and validated by experts can greatly improve our understanding of species

diversity at large spatial and temporal scales as well as ecological dynamics and the

conservation status of species (Steven et al., 2019). For a better understanding of the

biodiversity, ecology, and management of garden ponds, citizen science is an essential tool.

This is because it allows us to gain data from ponds situated mostly in private gardens and to

monitor details in local management based on data reported directly by the owners.

We conducted an online citizen science survey to explore the importance of garden

ponds by assessing potential links between the biodiversity and management of garden ponds

in Hungary based on data from pond owners. First, based on the resulting large-scale dataset

covering a wide gradient in urbanisation across the country, we aimed to reveal the prevalence

and local frequency (frequency of sightings) of multiple native aquatic and terrestrial animal

groups that visit or use these garden ponds. We focused on taxonomic groups that are

conspicuous enough so that they are easily and reliably identifiable by the public (amphibians,

birds, odonates) and also collected species data on adult amphibians for which we

accompanied our survey with an identification key. In addition, to gain information on

potentially invasive exotic species, we collected information on animals deliberately

introduced by the owners themselves. Second, in an attempt to identify the potential drivers of

native biota, we tested the presence of these organism groups against local pond features and

management practices applied in the garden ponds. We hypothesized that pond management

and features have a direct effect on the prevalence and local frequency of multiple animal

groups and that urbanisation has an overall negative effect on the presence of these animal

groups.

**2.   Materials and Methods**

2.1. Citizen science survey

An online survey within the citizen science project "MyPond" was launched in 2021

to investigate the biodiversity, local features, and management practices of garden ponds in

Hungary (Central Europe). The link to the questionnaire on the "MyPond" website

(https://mypond.hu/en/) was distributed via traditional media outlets and social media. The

survey consisted of a total of 30 questions: 6 multiple choices, 10 short answers, and 14

closed (yes-no) questions. We collected data on the local features of ponds, including age,

area, substrate of the pond (concrete, rubber, plastic, natural), and the presence/absence of

aquatic vegetation, shoreline vegetation, and fish. Besides, local management data included

circulation, cleaning, draining, leaf removal, and use of algaecide. The biodiversity dataset

comprised four animal groups, amphibians (here, we refer to adult amphibians or post-

metamorphic juvenile sightings) and their tadpoles (tadpoles were reported separately from

adult amphibians), birds, and odonates. The survey contained a link to the amphibian

identification guide (drawings) of the Hungarian Ornithological and Nature Conservation

Society   to   help   the   citizens   correctly   identify   amphibian   species

(https://www.mme.hu/en/node/4440), which is available as a free smartphone application.

Also, there was an option to name the type of birds they had seen in their gardens. The

questionnaire is still available on the website and collects data for future purposes. Between

2021 and 2022 (2021 July 6- 2022 September 9), 834 participants filled out the questionnaire,

of which a validated subset of 753 responses was used for statistical analysis (Figure 1).

Validation was carried out to exclude duplicates, incomplete data, and any pond with an area

over 200 $m^2$, which was based on the frequency distribution of pond areas (the largest was

24000 $m^2$, while most (98%) ponds were smaller than 200 $m^2$).

In the case of the amphibian species dataset, we excluded amphibian species if pond

owners made vague statements (e.g. "we have not seen that species in a few years").

However, we included species if the owner reported them as "rarely" seen. Furthermore,

entries were excluded from the amphibian species dataset if it only included the name "frog",

"toad", "newt", or "tadpoles", so species identity could not be determined. "Green frogs"

were grouped into the *Pelophylax* spp. complex as the three species that comprise this group

are morphologically difficult to separate (n = 332). Rare species were also cross-checked with

the most up-to-date species distribution maps (https://herpterkep.mme.hu/index.php?lang=en)

and retained only if their occurrence had already been proven within the region of the

sighting. Species-level data for birds was available from 667 ponds, as those ponds where

citizens reported birds only in general but did not specify the species were excluded from this

dataset. The full questionnaire can be found in the supplementary information (Table S1).

The resulting datasets included both prevalence (binary; 0, 1) and local frequency

(ordinal scale; 0, 1, 2, 3) data of each recorded animal group (amphibians and their tadpoles,

odonates, birds). Pond features and management were used as binomial and ratio scale data in

the statistical analysis (Table 1).

The area of urban land and wetland coverage as landscape-level drivers were

calculated using the Ecosystem Map of Hungary (project KEHOP-430-VEKOP-15-2016-

00001, Ministry of Agriculture, 2019). The number of raster pixels (2x2 m) was counted

within a 1 km radius circle around each focal pond using the Zonal Histogram tool. The

classification of land cover was used as defined by the Ecosystem Map of Hungary (Ministry

of Agriculture, 2019). The wetland coverage category contained pixels of wetlands,

marshlands, temporary wetlands, standing waters, and flowing waters, while the urban land

category (hereinafter urbanisation) contained pixels of tall buildings, short buildings, sealed

roads, dirt roads, railways, other artificial surfaces, and urban green spaces with and without

trees. Urbanisation and wetland coverage were then calculated as the proportion of total pixels

within the 1 km radius covered by the respective land cover type. All spatial analysis was

carried out in QGIS 3.28.1 (QGIS Development Team, 2022).

2.2. Statistical analysis

To visualize the prevalence and the local frequency of animal groups, we created

barplots and pie charts using the "ggplot2" R package (Wickham, 2016).

To examine the relationship between amphibian community composition and pond

features, pond management, and landscape-level drivers (urbanisation and wetland coverage),

we performed canonical correspondence analysis (CCA) using the "vegan" R package

(Oksanen, 2017). The analysis was based on prevalence data (binomial) of the amphibian

species, our only species-level dataset. To normalize the distribution of variables belonging to

pond features and landscape-level drivers, we used log transformation for pond area, and

square root transformation was applied for wetland coverage prior to the CCA. Pond

management variables were included as binomial data. Significant explanatory variables

among pond features, management, and landscape-level drivers were identified based on an

automatic stepwise (backward and forward) model selection procedure using permutation

tests ("ordistep" function, direction = "both", n = 999 permutations).

To analyze which pond features, management types, or landscape-level drivers had the

strongest effect on the prevalence of each animal group (Table 1), generalized linear models

(GLM) were fitted with a binomial distribution and logit link function (hereinafter logistic

regression model). Model selection was carried out based on Akaike's Information Criteria

(AIC) with the "MASS" R package (Venables and Ripley, 2002). To further analyze these

effects on birds, reported bird types were categorized into three main groups (wading birds,

ducks, and other birds, the latter category mostly including songbirds). Visualization was

carried out with the "ggplot2" (Wickham, 2016) and "ggbreak" (Xu et al., 2021) R packages.

All statistical analyses and illustrations were made with the R program v 4.2.2 (R Core

Team, 2022).

## 210    3. Results

Based on the prevalence data reported by 753 respondents, odonates (in 96.5% of the

ponds) and birds (96.5%) were observed at almost all garden ponds. While the prevalence of

amphibian sightings (81.4%) was higher than that of tadpoles, the latter was still observed in

close to two-thirds of the ponds (62.1%). Among the amphibian species, *Pelophylax* spp. was

the most frequently reported species (29.9%). The most frequently reported birds were

songbirds (88.3%, including tits, blackbirds, sparrows, etc.). Within the introduced animal

groups, fish were the most frequent (84.4%), while snails, turtles, mussels, and crayfish were

all introduced to less than 6% of the ponds, respectively (Figure 2).

Local frequency data of amphibians, birds, and odonates revealed that most (60%,

78.5%, and 86.5%, respectively) of the respondents regularly see these animal groups in their

ponds. Tadpoles were regularly present in the smallest share of ponds (40.6%) (Figure 3).

Tadpoles were absent from the highest share of ponds (37.8%), but amphibian breeding was

regularly observed in over 40% of the ponds detected via tadpole presence (Figure 3).

Participants recorded the presence of 13 amphibian species in garden ponds: three

newt species (*Ichthyosaura alpestris*, *Lissotriton vulgaris*, *Triturus dobrogicus*), nine anurans

(*Bombina bombina*, *Bufo bufo*, *Bufo viridis*, *Hyla arborea*, *Pelobates fuscus*, *Pelophylax* spp.,

*Rana arvalis*, *Rana dalmatina*, *Rana temporaria*) and one salamander (*Salamandra*

*salamandra*). Amphibian species composition (based on prevalence data, Table 1) was

significantly influenced by pond age, substrate, shoreline vegetation, algaecide usage, the

introduction of fish, and urbanisation (Figure 4). There was a positive association between

*Rana arvalis* and pond age, i.e., it occurred mostly in newly-installed ponds with rubber

substrate and shoreline vegetation. *Lissotriton vulgaris*, *Rana temporaria,* and *Rana*

*dalmatina* responded negatively to fish presence and algaecide treatment. *Bufo viridis* was

positively associated with urbanisation. *Salamandra salamandra*, *Ichthyosaura alpestris,*

*Triturus dobrogicus, Bombina bombina, Hyla arborea,* and *Pelobates fuscus* were positively

associated with large, well-vegetated ponds with a natural substrate, which are only

occasionally drained (Figure 4).

Urbanisation was negatively associated with the presence of amphibians and their

tadpoles, while it generally did not have a negative effect on the presence of odonates and

birds (Figure 5, Table S2). Pond age, area, and rubber substrate were the most frequently

significant pond features. Amphibians and their tadpoles, odonates, and birds were less likely

to be present in or at newly installed ponds (0-1 year). Pond area had a positive effect on

amphibians and their tadpoles. Algaecide had a negative effect on the presence of both

amphibians and their tadpoles. There was a positive effect of aquatic vegetation on tadpoles,

odonates and birds, however, shoreline vegetation was only significant for the presence of

tadpoles (Figure 5, Table S2).

**4. Discussion**

Our citizen science survey provided comprehensive large-scale data on the presence of

flagship animal groups of garden ponds, covering an entire European country. Odonates and

birds were observed at almost all 753 garden ponds, but amphibians and their tadpoles were

also common. Local frequency data revealed that most respondents regularly see these animal

groups in their ponds. The age of the ponds turned out to be a general predictor, with animals

more likely to be present in older ponds. In addition, area and rubber substrate were the most

important pond features across the animal groups. Among management types, algaecide

addition had a negative effect on amphibians and their tadpoles. Urbanisation had an overall

negative effect on the presence of amphibians and tadpoles; conversely, odonates and birds

showed an unexpected non-negative association. Our data also highlighted that owners

introduce various animal groups into their ponds, most frequently fish, with potential

consequences for the native biodiversity surrounding garden ponds. Our results demonstrated

the crucial role of garden ponds in urban biodiversity and conservation, as they provide

important secondary habitats for multiple animal groups in urban areas.

Urbanisation, i.e., the extent of urban landscape within a 1 km radius around the

ponds, had disparate effects across the animal groups we investigated. It had a negative effect

on the prevalence (presence/absence) of amphibians and their tadpoles in line with our

original hypothesis and also corroborated earlier findings on amphibian prevalence and

reproduction (Hamer and Parris, 2011; Rubbo and Kiesecker, 2005). Urbanisation is

characterized by landscapes fragmented by roads, houses, and buildings that can impede

dispersal between ponds and natural upland habitats that are important for amphibians to

complete their complex life cycle (Hamer and McDonnell, 2008; Scheffers and Paszkowski,

2012). Urbanisation can also lead to habitat degradation such as the loss of aquatic and

shoreline vegetation, which is key for amphibian oviposition and it also provides shelter for

amphibians and their larvae (Hamer and McDonnell, 2008). Although urbanisation generally

has a negative effect on amphibian diversity, some species are urban adapters (McKinney,

2002), favored by larger body sizes and lower dispersal requirements (they can persist in

small habitat patches) (Hahs et al., 2023). For example, *Bufo viridis* was positively associated

with urbanisation. This species has a relatively large body size and persists in the center of

European cities and towns (Landler et al., 2023; Mazgajska and Mazgajski, 2020; Sistani et

al., 2021). The *Pelophylax* spp. complex had no discernible associations with any of the

variables, probably because these species are habitat generalists and can maintain

metapopulations in both natural and urban landscapes (Herczeg et al., 2012; Francesco

Ficetola and De Bernardi, 2004). Conversely, species associated with natural ponds such as

*Ichthyosaura alpestris* were only recorded in the Bakony Mountains (western Hungary) which

is a protected forest, and this species is particularly vulnerable to landscape fragmentation due

to urbanisation (Emaresi et al., 2011; Van Buskirk, 2012). The sighting of *Salamandra*

*salamandra* also occurred adjacent to a large forested area (Bükk Mountains in northern

Hungary) and populations of this species can persist in low-density urban areas provided there

is sufficient forest nearby (Kiss et al., 2022).

On the other hand, urbanisation was not negatively associated with the prevalence of

odonates and birds in general. These results might seem counter-intuitive at first, although

there is evidence that in certain cases urban areas are able to maintain higher odonate species

diversity than agricultural lands, due to a network of public, stormwater, or garden ponds that

can foster the migration of species (Goertzen and Suhling, 2019; Willigalla and Fartmann,

2012). However, other studies also showed that species richness of odonates increased from

the city center to more rural areas, probably due to increasing habitat availability (more water

bodies in the rural areas than in cities) (Holtmann et al., 2018; Willigalla and Fartmann,

2012). Here, we only detected the presence of odonates as a group. Therefore, in our case, this

pattern could also be explained by a few highly tolerant species occurring in urbanized

landscapes, but more detailed follow-up studies will be necessary to link this pattern to

species identities and biodiversity. The same might be true for birds, resulting from the

presence of more generalist bird species such as blackbirds or sparrows, which are common

even in heavily urbanised landscapes (Callaghan et al., 2019; Mennechez and Clergeau,

2006). Furthermore, it has been found that birds were more likely to visit water sources

resembling natural waters, such as garden ponds, puddles, or fountains, in urban areas, where

other natural water sources are scarce (Tryjanowski et al., 2022), which could have

contributed to the pattern we found. Also, wading birds were a group quite frequently (24.3

%) reported from the ponds, which regularly included sightings of grey herons – a species

regularly utilising garden ponds for foraging (Jakubas and Manikowska, 2011).

Pond age was overall the most important pond feature shaping the prevalence of

animal groups. Newly installed ponds were less visited by wild animals, which can be due to

the lack of vegetation that could offer hiding and breeding places to aquatic and semi-aquatic

animal groups (Huikkonen et al., 2020). Additionally, colonization time might be another

factor as animals need time to find and occupy newly installed ponds in the urban landscape

(Cayuela et al., 2020; Coccia et al., 2016; Unglaub et al., 2021). Among the amphibian

species, *Rana arvalis* was strongly associated with newly installed ponds that this species can

potentially colonize while moving relatively long distances through the landscape (Brzeziński

and Mętrak, 2014). Altogether, amphibians and their tadpoles were significantly associated

with older ponds, correlating with a greater period for species to locate and colonize these

ponds (Moor et al., 2022). Many amphibian species prefer well-vegetated or large ponds

(Arntzen et al., 1997; Péntek et al., 2017; Rannap et al., 2013), which was supported by our

results that several species such as *Bombina bombina, Pelobates fuscus, Triturus dobrogicus*

were positively associated with aquatic vegetation in our ponds. Due to the larger size of the

rubber-lined ponds, rubber substrate had a significantly positive effect on the presence of

amphibians and their tadpoles, also odonates, birds, and wading birds (Hamer et al., 2023).

Algaecide addition was the most important form of pond management, influencing the

presence of animal groups, particularly amphibians and their tadpoles. The application of

algaecides in ponds was negatively associated with the prevalence of amphibians and their

tadpoles. This might be related to widely used substances of chemicals produced to reduce

algal growth such as copper, which can inhibit tadpole growth and reduce survival

(Christenson et al., 2014). It has been shown experimentally that tadpoles decrease algal

abundance due to grazing, which suggests that creating ponds that are suitable for amphibians

could provide natural bioremediation in garden ponds and eliminate the need to use algaecides

(Dvorsky et al., 2023). However, other animal groups like odonates or birds did not have a

significant negative association with algaecide addition. Birds use these habitats quite

differently (e.g. for drinking or bathing) (Gibbons et al., 2023) than amphibians, who spend

most of their lifecycle in the water. So birds may be less affected by the quality of water.

Regarding odonates, there can be a number of highly tolerant urban species that might even

withstand elevated levels of copper in the water (Tollett et al., 2009), which could have

contributed to their presence under algaecide addition.

Exotic animals are frequently introduced by the owners of garden ponds for

ornamental purposes (Hassall, 2014). According to our results, fish were the most frequently

introduced animals (e.g. goldfish), being consequently present in 84% of ponds. In

comparison to our finding, a study in the UK reported that only ~35% of the garden pond

owners stocked their ponds with fish (Loram et al., 2011), which was only a third of what we

found in Hungary. Besides fish, turtles and several invertebrates (snails, mussels, and

crayfish) were also reported. Previous studies found that aquaria and garden ponds may act as

sources of invasive species  (Chucholl, 2013; Patoka et al., 2014; Soes et al., 2011; Van Den

Neucker et al., 2017), e.g. the red-eared slider (*Trachemys scripta elegans*) (Tietz et al.,

2023), water hyacinth (*Pontederia crassipes*) (Olden et al., 2021), signal crayfish

(*Pacifastacus leniusculus*) (Chucholl et al., 2021). In this sense, without the indispensable

knowledge and education about invasive species, garden pond owners and garden centers can

pose a threat to native biodiversity. This highlights the importance of outreach activities

targeting pond owners, encouraging more conscious management choices.

We also found that the introduction of fish was negatively associated with all recorded

amphibian species. Fish are important predators of all amphibian life stages and a negative

relationship between fish and amphibian populations has been demonstrated for many

European amphibians (Hartel et al., 2007; Van Buskirk, 2005). Despite the introduction of

fish into 85% of ponds, amphibian reproduction was observed in 60% of the ponds, indicating

that some species are able to co-exist with fish (Hamer et al., 2021; Kloskowski et al., 2020).

In another study focusing on local pond features in the same set of ponds, it was also found

that the introduction of fish was a significant predictor of garden pond design and within-pond

features including management practices (Hamer et al., 2023). This result implies a potential

conflict between creating ponds with the main aim of keeping ornamental fish and the

potential of garden ponds to support native biodiversity.

To this date, there was no information available about the biodiversity of garden ponds

in Hungary, similar to the rest of Eastern Central Europe. This also means a limitation in the

implications of our study, given that we can not compare our questionnaire data to earlier or

other regional datasets as a reference. Another potential caveat is that even with our best

efforts for a validated dataset, there is the potential for species misidentification (especially

with amphibians). To overcome this, we collected data on taxa that can easily be identified,

and are considered flagship groups of pond diversity (Oertli et al., 2002). Concurrently, the

high number of ponds in our analyses provides robust data for the general patterns that can be

observed in these secondary habitats that could inform pond owners on management activities

being more beneficial for the local pond biota.

With the globally accelerating rate of urbanisation and the related loss and alteration

of aquatic habitats, a sound knowledge of the ecological roles of garden ponds is crucial. For

this, citizen science projects can be important tools by providing access to private properties

and data with high spatial coverage. The potential of the approach is well-illustrated by the

high level of engagement in the present survey, with over 800 completed forms. Based on

communication with the respondents, many had a high interest in the habitat quality and

biodiversity of their garden ponds. This suggests there is a high potential to change peoples'

perspectives on the "ideal" garden pond via evidence-based outreach activities. We suggest

the promotion of pond installations in urban areas by highlighting relevant facts about

invasive species and the most suitable local features and management types to maximise local

biodiversity. For effective biodiversity conservation in urban areas, where natural habitats are

under constant pressure from degradation and mismanagement, public input and engagement

are essential, for which citizen science can provide essential tools.

**Declaration of competing interest**

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

**CRediT authorship contribution statement**

*Zsuzsanna Márton*: Data curation, Formal analysis, Visualization, Writing - original draft,

Writing - review & editing

*Barbara Barta*: Conceptualization, Methodology, Data curation, Writing - review & editing

*Csaba F. Vad*: Conceptualization, Methodology, Writing - review & editing

*Beáta Szabó*: Formal analysis, Writing - review & editing

*Andrew J. Hamer*: Data curation, Formal analysis, Writing - review & editing

*Vivien Kardos*: Methodology, Visualization, Writing - review & editing

*Csilla Laskai*: Methodology, Writing - review & editing

*Ádám Fierpasz*: Methodology, Writing - review & editing

*Zsófia Horváth*: Conceptualization, Methodology, Writing - review & editing, Supervision

## Acknowledgments

402 This work would not have been possible without the involvement of 834 citizen scientists
403 filling out our survey. We acknowledge the help of Virág Csiszár in the popularization and
404 design of the homepage and survey.

## Funding

This work was supported by Eötvös Loránd Research Network (currently Hungarian Research
Network), the RRF-2.3.1-21-2022-369-00014 project, and the Sustainable Development and
Technologies National Programme of the Hungarian Academy of Sciences (FFT NP FTA).
Andrew J. Hamer, Barbara Barta, and Ádám Fierpasz acknowledge further support from
OTKA K142296. Zsófia Horváth acknowledges further support from the Bolyai+ Grant [grant
number ÚNKP-22-5-ELTE-84]. Barbara Barta was supported by the ÚNKP-22-3 New
National Excellence Program of the Ministry for Culture and Innovation from the source of
the National Research, Development and Innovation Fund [grant number ÚNKP-22-3-I-
ELTE-568].

## Data availability

416 Data will be deposited in the Figshare database upon acceptance of the article.

417 **Table 1** Data types in the validated dataset

| Main variable groups | Recorded variables | Data type |
|---|---|---|
| **Biodiversity - prevalence** | **amphibians and their tadpoles, birds, odonates** | Binomial (0, 1) |
| **Biodiversity - local frequency** | **amphibians and their tadpoles, birds, odonates** | Ordinal (0, 1, 2, 3, corresponding to no, singular, occasional, and frequent sightings per pond) |
| **Pond management** | **circulation, cleaning, draining, leaf removal, algaecide** | Binomial (0, 1) |
| **Pond features** | **age, area, substrate (concrete, rubber, plastic, natural), aquatic vegetation, shoreline vegetation, fish presence** | Ratio scale data |

420     **Figure 1** Location of the 753 garden ponds in Hungary used in the analyses. (Map was created with R

421                                                              v. 4.2.2)

423     **Figure 2** Percentage of all ponds with the occurrence of the animal groups reported by the citizens.

424     Samples sizes were n=667 in case of types of birds (top right), and n=753 in the other three datasets.

428    **Figure 3** Share (%) of each response on the local frequency of animal groups (amphibians and their

429                tadpoles, odonates and birds) in garden ponds (n=753)

**Figure 4** Canonical correspondence analysis (CCA) biplot based on the reported prevalence of

amphibian species, environmental predictors, and level of urbanisation (red points: species scores,

dark blue arrows: significant predictors, light green arrows: all other predictors). The analysis is based

on data from 332 ponds where participants recorded amphibian species data correctly.

436        **Figure 5** Summary of the best-fit logistic regression models (based on AIC) odds ratio and their 95% confidence interval testing the relationship

437     between the prevalence of animal groups and pond features and management. Samples sizes were n=753 for amphibians, odonates, birds, and tadpoles, and

438                            n=667 for other birds and wading birds.

**References**

Arntzen, J.W., Bugter, R.J.F., Cogalniceanu, D., Wallis, G.P., 1997. The distribution and
conservation status of the Danube crested newt, Triturus dobrogicus. Amphib.-Reptil.
18, 133–142. https://doi.org/10.1163/156853897X00026

Biggs, J., Von Fumetti, S., Kelly-Quinn, M., 2017. The importance of small waterbodies for
biodiversity and ecosystem services: implications for policy makers. Hydrobiologia
793, 3–39. https://doi.org/10.1007/s10750-016-3007-0

Biggs, J., Williams, P., Whitfield, M., Nicolet, P., Weatherby, A., 2005. 15 years of pond
assessment in Britain: results and lessons learned from the work of Pond
Conservation. Aquat. Conserv. Mar. Freshw. Ecosyst. 15, 693–714.
https://doi.org/10.1002/aqc.745

Brans, K.I., De Meester, L., 2018. City life on fast lanes: Urbanization induces an
evolutionary shift towards a faster lifestyle in the water flea *Daphnia*. Funct. Ecol. 32,
2225–2240. https://doi.org/10.1111/1365-2435.13184

Brans, K.I., Stoks, R., De Meester, L., 2018. Urbanization drives genetic differentiation in
physiology and structures the evolution of pace-of-life syndromes in the water flea
*Daphnia magna*. Proc. R. Soc. B Biol. Sci. 285, 20180169.
https://doi.org/10.1098/rspb.2018.0169

Brzeziński, M., Mętrak, M., 2014. Spring migration rates and community structure of
amphibians breeding in an old and newly established midfield ponds. Folia Zool. 63,
161–170. https://doi.org/10.25225/fozo.v63.i3.a3.2014

Callaghan, C.T., Major, R.E., Wilshire, J.H., Martin, J.M., Kingsford, R.T., Cornwell, W.K.,
2019. Generalists are the most urban☐tolerant of birds: a phylogenetically controlled
analysis of ecological and life history traits using a novel continuous measure of bird
responses to urbanization. Oikos 128, 845–858. https://doi.org/10.1111/oik.06158

Capotorti, G., Valeri, S., Giannini, A., Minorenti, V., Piarulli, M., Audisio, P., 2023. On the

Role of Natural and Induced Landscape Heterogeneity for the Support of Pollinators:

A Green Infrastructure Perspective Applied in a Peri-Urban System. Land 12, 387.

https://doi.org/10.3390/land12020387

Cayuela, H., Valenzuela-Sánchez, A., Teulier, L., Martínez-Solano, Í., Léna, J.-P., Merilä, J.,

Muths, E., Shine, R., Quay, L., Denoël, M., Clobert, J., Schmidt, B.R., 2020.

Determinants and Consequences of Dispersal in Vertebrates with Complex Life

Cycles: A Review of Pond-Breeding Amphibians. https://doi.org/10.5167/UZH-

185243

Céréghino, R., Biggs, J., Oertli, B., Declerck, S., 2007. The ecology of European ponds:

defining the characteristics of a neglected freshwater habitat, in: Oertli, Beat,

Céréghino, Régis, Biggs, Jeremy, Declerck, Steven, Hull, A., Miracle, M.R. (Eds.),

Pond Conservation in Europe. Springer Netherlands, Dordrecht, pp. 1–6.

https://doi.org/10.1007/978-90-481-9088-1_1

Christenson, T.A., Horton, M.E., Jackson, B.C., Smith, G.R., Rettig, J.E., 2014. Effects of

Cutrine-Plus® algaecide and predators on wood frog (Lithobates sylvaticus) tadpole

survival and growth. Environ. Sci. Pollut. Res. 21, 12472–12478.

https://doi.org/10.1007/s11356-014-3186-z

Chucholl, C., 2013. Invaders for sale: trade and determinants of introduction of ornamental

freshwater crayfish. Biol. Invasions 15, 125–141. https://doi.org/10.1007/s10530-012-

0273-2

Chucholl, F., Fiolka, F., Segelbacher, G., Epp, L.S., 2021. eDNA Detection of Native and

Invasive Crayfish Species Allows for Year-Round Monitoring and Large-Scale

Screening of Lotic Systems. Front. Environ. Sci. 9, 639380.

https://doi.org/10.3389/fenvs.2021.639380

Coccia, C., Vanschoenwinkel, B., Brendonck, L., Boyero, L., Green, A.J., 2016. Newly

490            created ponds complement natural waterbodies for restoration of macroinvertebrate

491            assemblages. Freshw. Biol. 61, 1640–1654. https://doi.org/10.1111/fwb.12804

492     Concepción, E.D., 2021. Urban sprawl into Natura 2000 network over Europe. Conserv. Biol.

493            35, 1063–1072. https://doi.org/10.1111/cobi.13687

494     Danneck, J., Chin Fung Tiang, D., Ngo Duc, K., Azhar, B., Sanusi, R., Campos Arceiz, A.,

495            Lechner, A.M., 2023. Conserving urban tropical biodiversity by connecting networks

496            of green patches. Integr. Conserv. inc3.21. https://doi.org/10.1002/inc3.21

497     Downing, J.A., Prairie, Y.T., Cole, J.J., Duarte, C.M., Tranvik, L.J., Striegl, R.G., McDowell,

498            W.H., Kortelainen, P., Caraco, N.F., Melack, J.M., Middelburg, J.J., 2006. The global

499            abundance and size distribution of lakes, ponds, and impoundments. Limnol.

500            Oceanogr. 51, 2388–2397. https://doi.org/10.4319/lo.2006.51.5.2388

501     Dvorsky, C., Riddle, K., Boone, M., 2023. Assessing the Impact of Chemical Algae

502            Management Strategies on Anurans and Aquatic Communities. Environ. Toxicol.

503            Chem. 42, 213–224. https://doi.org/10.1002/etc.5514

504     Emaresi, G., Pellet, J., Dubey, S., Hirzel, A.H., Fumagalli, L., 2011. Landscape genetics of

505            the Alpine newt (Mesotriton alpestris) inferred from a strip-based approach. Conserv.

506            Genet. 12, 41–50. https://doi.org/10.1007/s10592-009-9985-y

507     Fraisl, D., Hager, G., Bedessem, B., Gold, M., Hsing, P.-Y., Danielsen, F., Hitchcock, C.B.,

508            Hulbert, J.M., Piera, J., Spiers, H., Thiel, M., Haklay, M., 2022. Citizen science in

509            environmental and ecological sciences. Nat. Rev. Methods Primer 2, 64.

510            https://doi.org/10.1038/s43586-022-00144-4

511     Francesco Ficetola, G., De Bernardi, F., 2004. Amphibians in a human-dominated landscape:

512            the community structure is related to habitat features and isolation. Biol. Conserv. 119,

513            219–230. https://doi.org/10.1016/j.biocon.2003.11.004

Gibbons, E.K., Close, P.G., Van Helden, B.E., Rooney, N.J., 2023. Water in the city:

visitation of animal wildlife to garden water sources and urban lakes. Urban Ecosyst.

26, 1413–1425. https://doi.org/10.1007/s11252-023-01391-3

Goertzen, D., Suhling, F., 2019. Urbanization versus other land use: Diverging effects on

dragonfly    communities    in    Germany.    Divers.    Distrib.    25,    38–47.

https://doi.org/10.1111/ddi.12820

Grooten, M., Almond, R.E.A. (Eds.), 2018. Living planet report 2018: aiming higher. WWF--

World Wide Fund for Nature, Gland, Switzerland.

Hahs, A.K., Fournier, B., Aronson, M.F.J., Nilon, C.H., Herrera-Montes, A., Salisbury, A.B.,

Threlfall, C.G., Rega-Brodsky, C.C., Lepczyk, C.A., La Sorte, F.A., MacGregor-Fors,

I., Scott MacIvor, J., Jung, K., Piana, M.R., Williams, N.S.G., Knapp, S., Vergnes, A.,

Acevedo, A.A., Gainsbury, A.M., Rainho, A., Hamer, A.J., Shwartz, A., Voigt, C.C.,

Lewanzik, D., Lowenstein, D.M., O'Brien, D., Tommasi, D., Pineda, E., Carpenter,

E.S., Belskaya, E., Lövei, G.L., Makinson, J.C., Coleman, J.L., Sadler, J.P., Shroyer,

J., Shapiro, J.T., Baldock, K.C.R., Ksiazek-Mikenas, K., Matteson, K.C., Barrett, K.,

Siles, L., Aguirre, L.F., Armesto, L.O., Zalewski, M., Herrera-Montes, M.I., Obrist,

M.K., Tonietto, R.K., Gagné, S.A., Hinners, S.J., Latty, T., Surasinghe, T.D., Sattler,

T., Magura, T., Ulrich, W., Elek, Z., Castañeda-Oviedo, J., Torrado, R., Kotze, D.J.,

Moretti, M., 2023. Urbanisation generates multiple trait syndromes for terrestrial

animal taxa worldwide. Nat. Commun. 14, 4751. https://doi.org/10.1038/s41467-023-

39746-1

Hamer, A.J., Barta, B., Bohus, A., Gál, B., Schmera, D., 2021. Roads reduce amphibian

abundance in ponds across a fragmented landscape. Glob. Ecol. Conserv. 28, e01663.

https://doi.org/10.1016/j.gecco.2021.e01663

Hamer, A.J., Barta, B., Márton, Z., Vad, C.F., Szabó, B., Tornero, I., Horváth, Z., 2023.

Patterns and correlates in the distribution, design and management of garden ponds

along an urban-rural gradient (preprint). Ecology.

https://doi.org/10.1101/2023.11.30.569344

Hamer, A.J., McDonnell, M.J., 2008. Amphibian ecology and conservation in the urbanising

world: A review. Biol. Conserv. 141, 2432–2449.

https://doi.org/10.1016/j.biocon.2008.07.020

Hamer, A.J., Parris, K.M., 2011. Local and landscape determinants of amphibian

communities in urban ponds. Ecol. Appl. 21, 378–390. https://doi.org/10.1890/10-

0390.1

Hartel, T., Nemes, S., Cogălniceanu, D., Öllerer, K., Schweiger, O., Moga, C.-I., Demeter, L.,

2007. The effect of fish and aquatic habitat complexity on amphibians. Hydrobiologia

583, 173–182. https://doi.org/10.1007/s10750-006-0490-8

Hassall, C., 2014. The ecology and biodiversity of urban ponds. WIREs Water 1, 187–206.

https://doi.org/10.1002/wat2.1014

Herczeg, D., Gallé, R., & Molnár, N. 2012. Effects of urbanization on the occurence

of Anura assemblages in the city of Szeged (Hungary). Tiscia, 39, 3-8.

Hill, M.J., Wood, P.J., Fairchild, W., Williams, P., Nicolet, P., Biggs, J., 2021. Garden pond

diversity: Opportunities for urban freshwater conservation. Basic Appl. Ecol. 57, 28–

40. https://doi.org/10.1016/j.baae.2021.09.005

Holtmann, L., Juchem, M., Brüggeshemke, J., Möhlmeyer, A., Fartmann, T., 2018.

Stormwater ponds promote dragonfly (Odonata) species richness and density in urban

areas. Ecol. Eng. 118, 1–11. https://doi.org/10.1016/j.ecoleng.2017.12.028

Huikkonen, I., Helle, I., Elo, M., 2020. Heterogenic aquatic vegetation promotes abundance

and species richness of Odonata (Insecta) in constructed agricultural wetlands. Insect

Conserv. Divers. 13, 374–383. https://doi.org/10.1111/icad.12396

Jakubas, D., Manikowska, B., 2011. The response of Grey Herons Ardea cinerea to changes
in prey abundance. Bird Study 58, 487–494.
https://doi.org/10.1080/00063657.2011.608423

Johansson, F., Heino, J., Coiffard, P., Svanbäck, R., Wester, J., Bini, L.M., 2020. Can
information from citizen science data be used to predict biodiversity in stormwater
ponds? Sci. Rep. 10, 9380. https://doi.org/10.1038/s41598-020-66306-0

Johnson, M.T.J., Munshi-South, J., 2017. Evolution of life in urban environments. Science
358, eaam8327. https://doi.org/10.1126/science.aam8327

Kiss, I., Vörös, J., Hamer, A.J., 2022. Movement patterns within an urban population of fire
salamanders highlight the importance of conserving small habitat patches. J. Zool.
316, 240–249. https://doi.org/10.1111/jzo.12949

Kloskowski, J., Nieoczym, M., Stryjecki, R., 2020. Between-habitat distributions of pond
tadpoles and their insect predators in response to fish presence. Hydrobiologia 847,
1343–1356. https://doi.org/10.1007/s10750-020-04190-5

Landler, L., Burgstaller, S., Schweiger, S., 2023. Land-use preferences of the European green
toad (Bufotes viridis) in the city of Vienna (Austria): the importance of open land in
urban environments. Front. Zool. 20, 3. https://doi.org/10.1186/s12983-022-00480-x

Loram, A., Warren, P., Thompson, K., Gaston, K., 2011. Urban Domestic Gardens: The
Effects of Human Interventions on Garden Composition. Environ. Manage. 48, 808–
824. https://doi.org/10.1007/s00267-011-9723-3

Ma, L., Liu, Y., Lu, W., Zhang, Zheng, Li, W., Zhang, Ziwei, Zhang, X., Zhu, C., Bai, J., Xu,
Z., Han, Y., Ruan, L., 2023. A highly effective incubation strategy enhanced the urban
bird hatch success. Avian Res. 14, 100074. https://doi.org/10.1016/j.avrs.2022.100074

Mazgajska, J., Mazgajski, T.D., 2020. Two amphibian species in the urban environment:
changes in the occurrence, spawning phenology and adult condition of common and

green       toads.       Eur.       Zool.       J.       87,       170–179.

https://doi.org/10.1080/24750263.2020.1744743

McDonald, R.I., Mansur, A.V., Ascensão, F., Colbert, M., Crossman, K., Elmqvist, T.,

Gonzalez, A., Güneralp, B., Haase, D., Hamann, M., Hillel, O., Huang, K., Kahnt, B.,

Maddox, D., Pacheco, A., Pereira, H.M., Seto, K.C., Simkin, R., Walsh, B., Werner,

A.S., Ziter, C., 2019. Research gaps in knowledge of the impact of urban growth on

biodiversity. Nat. Sustain. 3, 16–24. https://doi.org/10.1038/s41893-019-0436-6

McKinney, M.L., 2002. Urbanization, Biodiversity, and Conservation. BioScience 52, 883.

https://doi.org/10.1641/0006-3568(2002)052[0883:UBAC]2.0.CO;2

Memarsadeghi, N., Joseph, R.D., Kaufmann, J.C., Lee, B.S., 2022. Golomb Patterns,

Astrophysics, and Citizen Science Games. IEEE Access 10, 76125–76135.

https://doi.org/10.1109/ACCESS.2022.3188632

Mennechez, G., Clergeau, P., 2006. Effect of urbanisation on habitat generalists: starlings not

so flexible? Acta Oecologica 30, 182–191. https://doi.org/10.1016/j.actao.2006.03.002

Ministry of Agriculture, 2019. Ecosystem base map of Hungary. DOI:

10.34811/osz.alapterkep

Moor, H., Bergamini, A., Vorburger, C., Holderegger, R., Bühler, C., Egger, S., Schmidt,

B.R., 2022. Bending the curve: Simple but massive conservation action leads to

landscape-scale recovery of amphibians. Proc. Natl. Acad. Sci. 119, e2123070119.

https://doi.org/10.1073/pnas.2123070119

Oertli, B., Céréghino, R., Hull, A., Miracle, R., 2009. Pond conservation: from science to

practice. Hydrobiologia 634, 1–9. https://doi.org/10.1007/s10750-009-9891-9

Oertli, B., Decrey, M., Demierre, E., Fahy, J.C., Gallinelli, P., Vasco, F., Ilg, C., 2023.

Ornamental ponds as Nature-based Solutions to implement in cities. Sci. Total

Environ. 888, 164300. https://doi.org/10.1016/j.scitotenv.2023.164300

Oertli, B., Joye, D.A., Castella, E., Juge, R., Cambin, D., Lachavanne, J.-B., 2002. Does size

matter? The relationship between pond area and biodiversity. Biol. Conserv. 104, 59–

70. https://doi.org/10.1016/S0006-3207(01)00154-9

Oksanen, J., 2017. Vegan: ecological diversity. R Package Version 24-4.

Olden, J.D., Whattam, E., Wood, S.A., 2021. Online auction marketplaces as a global

pathway for aquatic invasive species. Hydrobiologia 848, 1967–1979.

https://doi.org/10.1007/s10750-020-04407-7

Patoka, J., Bláha, M., Kalous, L., Kouba, A., 2017. Irresponsible vendors: Non-native,

invasive and threatened animals offered for garden pond stocking: Non-native and

threatened animals in garden ponds. Aquat. Conserv. Mar. Freshw. Ecosyst. 27, 692–

697. https://doi.org/10.1002/aqc.2719

Patoka, J., Petrtýl, M., Kalous, L., 2014. Garden ponds as potential introduction pathway of

ornamental crayfish. Knowl. Manag. Aquat. Ecosyst. 13.

https://doi.org/10.1051/kmae/2014019

Peeters, E.T.H.M., Wilhelm, M., Gerritsen, A.A.M., Seelen, L.M.S., 2023. Ecological

characterisation of urban ponds in the Netherlands: a study based on data collected by

volunteers. Front. Biogeogr. 15. https://doi.org/10.21425/F5FBG57816

Péntek, A.L., Vad, C.F., Zsuga, K., Horváth, Z., 2017. Metacommunity dynamics of

amphibians in years with differing rainfall. Aquat. Ecol. 51, 45–57.

https://doi.org/10.1007/s10452-016-9597-9

QGIS Development Team, 2022. QGIS Geographic Information System. Open Source

Geospatial Foundation Project. http://qgis.osgeo.org

Rannap, R., Markus, M., Kaart, T., 2013. Habitat use of the common spadefoot toad

(Pelobates fuscus) in Estonia. Amphib.-Reptil. 34, 51–62.

https://doi.org/10.1163/15685381-00002868

R Core Team, 2022. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Richardson, D.C., Holgerson, M.A., Farragher, M.J., Hoffman, K.K., King, K.B.S., Alfonso,

M.B., Andersen, M.R., Cheruveil, K.S., Coleman, K.A., Farruggia, M.J., Fernandez,

R.L., Hondula, K.L., López Moreira Mazacotte, G.A., Paul, K., Peierls, B.L., Rabaey,

J.S., Sadro, S., Sánchez, M.L., Smyth, R.L., Sweetman, J.N., 2022. A functional

definition to distinguish ponds from lakes and wetlands. Sci. Rep. 12, 10472.

https://doi.org/10.1038/s41598-022-14569-0

Rivera-Gutierrez, H.F., Jaramillo-Calle, V., Lopera-Salazar, A., Martinez-Alvarado, D., 2023.

Does learning matter? Birdsong-learning program determines coping strategies for

living in urban noisy environments. Behav. Ecol. Sociobiol. 77, 22.

https://doi.org/10.1007/s00265-023-03292-z

Rubbo, M.J., Kiesecker, J.M., 2005. Amphibian Breeding Distribution in an Urbanized

Landscape. Conserv. Biol. 19, 504–511. https://doi.org/10.1111/j.1523-

1739.2005.000101.x

Ruggiero, A., Céréghino, R., Figuerola, J., Marty, P., Angélibert, S., 2008. Farm ponds make

a contribution to the biodiversity of aquatic insects in a French agricultural landscape.

C. R. Biol. 331, 298–308. https://doi.org/10.1016/j.crvi.2008.01.009

Scheffers, B.R., Paszkowski, C.A., 2012. The effects of urbanization on North American

amphibian species: Identifying new directions for urban conservation. Urban Ecosyst.

15, 133–147. https://doi.org/10.1007/s11252-011-0199-y

Schueller, S.K., Li, Z., Bliss, Z., Roake, R., Weiler, B., 2023. How Informed Design Can

Make a Difference: Supporting Insect Pollinators in Cities. Land 12, 1289.

https://doi.org/10.3390/land12071289

Sistani, A., Burgstaller, S., Gollmann, G., Landler, L., 2021. The European green toad,

Bufotes viridis, in Donaufeld (Vienna, Austria): status and size of the population. Herpetozoa 34, 259–264. https://doi.org/10.3897/herpetozoa.34.e75578

Smith, M.L., 2014. Citizen Science in Archaeology. Am. Antiq. 79, 749–762. https://doi.org/10.7183/0002-7316.79.4.749749

Soanes, K., Sievers, M., Chee, Y.E., Williams, N.S.G., Bhardwaj, M., Marshall, A.J., Parris, K.M., 2019. Correcting common misconceptions to inspire conservation action in urban environments. Conserv. Biol. 33, 300–306. https://doi.org/10.1111/cobi.13193

Soanes, K., Taylor, L., Ramalho, C.E., Maller, C., Parris, K., Bush, J., Mata, L., Williams, N.S.G., Threlfall, C.G., 2023. Conserving urban biodiversity: Current practice, barriers, and enablers. Conserv. Lett. 16, e12946. https://doi.org/10.1111/conl.12946

Soes, M., Majoor, G., Keulen, S., 2011. Bellamya chinensis (Gray, 1834) (Gastropoda: Viviparidae), a new alien snail species for the European fauna. Aquat. Invasions 6, 97–102. https://doi.org/10.3391/ai.2011.6.1.12

Stamenković, O., Stojković Piperac, M., Milošević, D., Buzhdygan, O.Y., Petrović, A., Jenačković, D., Đurđević, A., Čerba, D., Vlaičević, B., Nikolić, D., Simić, V., 2019. Anthropogenic pressure explains variations in the biodiversity of pond communities along environmental gradients: a case study in south-eastern Serbia. Hydrobiologia 838, 65–83. https://doi.org/10.1007/s10750-019-03978-4

Steven, R., Barnes, M., Garnett, S.T., Garrard, G., O'Connor, J., Oliver, J.L., Robinson, C., Tulloch, A., Fuller, R.A., 2019. Aligning citizen science with best practice: Threatened species conservation in Australia. Conserv. Sci. Pract. 1. https://doi.org/10.1111/csp2.100

Tickner, D., Opperman, J.J., Abell, R., Acreman, M., Arthington, A.H., Bunn, S.E., Cooke, S.J., Dalton, J., Darwall, W., Edwards, G., Harrison, I., Hughes, K., Jones, T., Leclère, D., Lynch, A.J., Leonard, P., McClain, M.E., Muruven, D., Olden, J.D., Ormerod, S.J.,

Robinson, J., Tharme, R.E., Thieme, M., Tockner, K., Wright, M., Young, L., 2020.

Bending the Curve of Global Freshwater Biodiversity Loss: An Emergency Recovery

Plan. BioScience 70, 330–342. https://doi.org/10.1093/biosci/biaa002

Tietz, B., Penner, J., Vamberger, M., 2023. Chelonian challenge: three alien species from

North America are moving their reproductive boundaries in Central Europe. NeoBiota

82, 1–21. https://doi.org/10.3897/neobiota.82.87264

Tollett, V.D., Benvenutti, E.L., Deer, L.A., Rice, T.M., 2009. Differential Toxicity to Cd, Pb,

and Cu in Dragonfly Larvae (Insecta: Odonata). Arch. Environ. Contam. Toxicol. 56,

77–84. https://doi.org/10.1007/s00244-008-9170-1

Tryjanowski, P., Jankowiak, Ł., Czechowski, P., Dulisz, B., Golawski, A., Grzywaczewski,

G., Indykiewicz, P., Kwieciński, Z., Mitrus, C., Nowakowski, J.J., Polakowski, M.,

Siekiera, J., Sparks, T.H., Wuczyński, A., 2022. Summer water sources for temperate

birds:   use,   importance,   and   threats.   Eur.   Zool.   J.   89,   913–926.

https://doi.org/10.1080/24750263.2022.2101699

Uchida, K., Shimamoto, T., Yanagawa, H., Koizumi, I., 2020. Comparison of multiple

behavioral traits between urban and rural squirrels. Urban Ecosyst. 23, 745–754.

https://doi.org/10.1007/s11252-020-00950-2

Unglaub, B., Cayuela, H., Schmidt, B.R., Preißler, K., Glos, J., Steinfartz, S., 2021.

Context□dependent dispersal determines relatedness and genetic structure in a patchy

amphibian population. Mol. Ecol. 30, 5009–5028. https://doi.org/10.1111/mec.16114

Van Buskirk, J., 2012. Permeability of the landscape matrix between amphibian breeding

sites. Ecol. Evol. 2, 3160–3167. https://doi.org/10.1002/ece3.424

Van Buskirk, J., 2005. LOCAL AND LANDSCAPE INFLUENCE ON AMPHIBIAN

OCCURRENCE   AND   ABUNDANCE.   Ecology   86,   1936–1947.

https://doi.org/10.1890/04-1237

Van Den Neucker, T., Schildermans, T., Scheers, K., 2017. The invasive Chinese mystery

snail *Bellamya chinensis* (Gastropoda: Viviparidae) expands its European range to

Belgium. Knowl. Manag. Aquat. Ecosyst. 8. https://doi.org/10.1051/kmae/2016040

Venables, W.N., Ripley, B.D., 2002. Modern Applied Statistics with S, Statistics and

Computing. Springer New York, New York, NY. https://doi.org/10.1007/978-0-387-

21706-2

Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis, Journal of the Royal

Statistical Society: Series A (Statistics in Society).

Williams, P., Whitfield, M., Biggs, J., Bray, S., Fox, G., Nicolet, P., Sear, D., 2004.

Comparative biodiversity of rivers, streams, ditches and ponds in an agricultural

landscape in Southern England. Biol. Conserv. 115, 329–341.

https://doi.org/10.1016/S0006-3207(03)00153-8

Willigalla, C., Fartmann, T., 2012. Patterns in the diversity of dragonflies (Odonata) in cities

across Central Europe. Eur. J. Entomol. 109, 235–245.

https://doi.org/10.14411/eje.2012.031

Xu, S., Chen, M., Feng, T., Zhan, L., Zhou, L., Yu, G., 2021. Use ggbreak to Effectively

Utilize Plotting Space to Deal With Large Datasets and Outliers. Front. Genet. 12,

774846. https://doi.org/10.3389/fgene.2021.774846

Yin, L., Wang, C., Han, W., Zhang, C., 2023. Birds' Flight Initiation Distance in Residential

Areas of Beijing Are Lower than in Pristine Environments: Implications for the

Conservation of Urban Bird Diversity. Sustainability 15, 4994.

https://doi.org/10.3390/su15064994

**Web references**

https://mypond.hu/en/ - Last accessed: 2023.11.23.

https://www.mme.hu/en/node/4440 - Last accessed: 2023.10.15.

https://herpterkep.mme.hu/index.php?lang=en - Last accessed: 2023.11.23.