Bone quality following peripubertal growth in a mouse model of transmasculine gender-affirming hormone therapy

Brandon Henry[1,2], Cynthia Dela Cruz[3], Robert W. Goulet[1], Bonnie T. Nolan[1], Conor Locke[1], Vasantha Padmanabhan[4,5], Molly Moravek[5], Ariella Shikanov[3,5], and Megan L. Killian[1]*

1. Department of Orthopaedic Surgery, Michigan Medicine, Ann Arbor, Michigan

2. College of Osteopathic Medicine, Michigan State University, East Lansing, Michigan

3. Department of Biomedical Engineering, University of Michigan, Ann Arbor, Michigan

4. Department of Pediatrics, Michigan Medicine, Ann Arbor, Michigan

5. Department of Obstetrics and Gynecology, Michigan Medicine, Ann Arbor, Michigan

Corresponding author:

*Megan L. Killian, PhD; mlkillia@med.umich.edu

Keywords: gender-affirming hormone therapy, bone mechanics, bone morphometry, mouse model

Abstract

During peri-puberty, bone growth and the attainment peak bone mass is driven predominantly by sex steroids. This is important when treating transgender youth, who have become increasingly present at pediatric clinics. Analogues of gonadotropin-releasing hormone (GnRH) are commonly prescribed to transgender youth prior to starting gender-affirming hormone therapy (GAHT). However, the impact of GnRH agonists on long bones with the addition of GAHT is relatively unknown. To explore this, we developed a trans-masculine model by introducing either GnRHa or vehicle treatment to female-born mice at a pre-pubertal age. This treatment was followed by male GAHT (testosterone, T) or control treatment three weeks later. Six weeks after T therapy, bone quality was compared between four treatment groups: Control (vehicle only), GnRHa-only, GnRHa + T, and T-only. Bone length/size, bone shape, mechanical properties, and trabecular morphology were modulated by GAHT. Independent of GnRHa administration, mice treated with T had shorter femurs, larger trabecular volume and increased trabecular number, higher trabecular bone mineral density, and wider superstructures on the surface of bone (e.g., third trochanters) when compared to control or GnRHa-only mice. In conclusion, prolonged treatment of GnRHa with subsequent GAHT treatment directly affect the composition, parameters, and morphology of the developing long bone. These findings provide insight to help guide clinical approaches to care for transgender youth.

Introduction

Sex steroids play a crucial role in the attainment and maintenance of peak bone density (1). While sex steroids affect individuals during all ages of life, their effects are at their highest and most active during peri-pubertal growth affecting body composition and peak bone mass.(2–4). In peri-pubescent boys, testosterone (T) stimulates periosteal apposition, leading to increased bone width compared with peri-pubescent girls, despite comparable cortical thickness between sexes (2). Estrogen (E) plays a major regulatory role in bone metabolism in both men and women, especially post-puberty, as it maintains bone remodeling and homeostasis (2). During development, estrogen promotes closure of the epiphysial plate during puberty; testosterone can induce similar effects via conversion to estrogen by aromatase (5). The release of sex steroids is regulated by signaling within the hypothalamo-pituitary gonadal (HPG)-axis. Gonadotropin hormone-releasing hormone (GnRH) is secreted by the hypothalamus and stimulates the release of pituitary gonadotoprins that activate gonadal production of sex steroids like T and E. Inhibition of the HPG-axis leads to a hypoestrogenic state, limiting the role of estrogen in bone homeostasis (6,7).

Puberty suppression (specifically for transgender adolescents) has been used historically to lessen the psychological burden of gender dysmorphia by making physical sex characteristics more congruent with the experienced gender (8–10). One way this is accomplished is through administration of a GnRH agonist (GnRHa). Within the HPG- axis GnRHa will suppress the production and section of luteinizing hormone (LH) and follicle stimulating hormone (FSH), which results in arrested ovarian folliculogenesis and steroidogenesis. Adolescent patients with persistent feelings of gender dysphoria that worsens at the onset of puberty are eligible for GnRHa treatment according to International Guidelines (8,11,12). GnRHa can be prescribed as

early as 10-11 yrs for trans boys and 11-12yrs for trans girls (corresponding to Tanner stage 2-3). Around 15-16 yrs of age, T or E can be used as gender affirming hormone therapy (GAHT) for full hormonal transitioning in consenting sufficiently capacitated patients (8,11,12). While much of what we know about GAHT in humans has focused on reproductive health, the effect of GAHT and delayed puberty/prolonged pubertal suppression on bone health are underappreciated. Previously, a decrease in areal bone mineral density (aBMD) Z- scores were reported after GnRHa administration, which were partially restored after GAHT (13).

While the effect of sex hormones on bone morphology and properties have been heavily studied, the impact of induction of the hypoestrogenic state in combination with T treatment in peripubertal mice such as that used in GAHT has not been explored extensively. In a model like ours, Dubois et al reports that GnRHa administration reduced bone mass acquisition, and bone strength in female to male (transmasculine) mice (8). This effect was also reversed by administration of T, achieving parameters equal to control female mice and in some cases male control (8). In this study, we examined the effect of GnRHA treatment with and without testosterone treatment on long bones in mice during peripubertal skeletal growth. Specifically, we measured femoral bone quality in adult female-born mice subjected to pubertal suppression followed by testosterone treatment to assess changes in the morphology of trochanteric, cortical, trabecular bone and its mechanical properties.

Methods and Materials

Experimental Design

C57BL/6N female mice (N=28) and age-matched male mice (N= 3) were used in this study. Female mice were assigned to one of four experimental groups to mimic gender-affirming hormone therapy (GAHT) in humans, and experiments were initiated at postnatal age 26 days

(P26), which is approximately the time female mice enter puberty (Figure 1) (14). Experimental

groups included controls (sham surgery with empty silastic tubing implant; n=7), GnRHa-only

treatment (for pubertal suppression; n=7), T-only treatment (for male transition; n=7), and

GnRHa + T treatment (combined; n=7). For the GnRHa and GnRHa + T groups, mice were

subcutaneously implanted with GnRHa (3.6mg) depot at P26. Control and T-only groups

underwent a sham surgery with no implantation. At 3 weeks post-implantation or sham surgery,

mice were subcutaneously implanted with silastic tubing containing testosterone (10 mg) or

empty silastic tubing (control and GnRHa-only group). Six weeks after T-implantation, all

animals were euthanized (Figure 1). Body weights were recorded at time of euthanasia and one

femur from each mouse was dissected and fixed in 4% cold paraformaldehyde for 24-48hr then

stored in 70% ethanol.

Figure 1. Summary of experimental design. At P26, C57BL/6N female mice were treated with

GnRHa or sham. After 3 weeks, mice were supplemented with Testosterone (n=14) or placebo

(n=14). Animals were euthanized at P89, then compared to control group (Sham + Placebo) and

untreated P89 male mice.

Nano-computed tomography

Fixed femurs were imaged using nano-computed tomography (nanoCT; nanotom-m, phoenix|x-ray, Wunstorf, Germany) to obtain high resolution three-dimensional images of the femurs (parameters: 7.9 mm voxel size; 80kV; 400mA; 0.762mm aluminum filter; average parameter of 3; and integration time of 500ms). Calibration phantoms containing air, water, and a hydroxyapatite mimic (1.69mg/cc; Gammex, Middleton, WI, USA) were included in each scan. Full image volumes were reconstructed using datos|x software (GE, Inspection Technologies, LP, Skaneateles, NY, USA), and gray values were converted to Hounsfield units using the calibration phantoms. Dragonfly software (version 2021.1; Object Research Systems Inc, Montreal, Canada) was used to create a volume of interest (VOI) and a region of interest (ROI) of each imaged femur (Figure 2). A validated fully convolutional neural network (FCNN) was used to segment bone from background (DICE coefficient = 0.983 ± 0.016). A second validated FCNN was used to segment cortical from trabecular bone (Dice coefficient = 0.977 ± 0.023) to generate cortical, trabecular, third trochanter sub-volumes for structural quantification (15). The segmented image files were visually inspected by an expert (B.H.) for thresholding quality and cortical-trabecular segmentation (Figure 2). Erroneously identified voxels were manually corrected (15).

Figure 2. High resolution nanoCT was used to generate volumetric images of femurs (7.9mm voxel size). Analyses were performed on regions of interests including third trochanter, cortical, and trabecular regions.

All images were rotated to align the femoral long axis with the image Z-axis and the third trochanter with the X-axis for analysis. The boundary between cortical (Ct.) and trabecular (Tb.) bone was defined by prior studies (16,17). A boundary formula of the third trochanter was modeled after the cortical and trabecular bone boundary. We defined the start of the third trochanter ROI as the proximal boundary of the cortical bone where the periosteal surface starts to extend outward. The height of the third trochanter (in the long-axis Z dimension) was created by taking the 15% percent of total bone length. Third trochanter diameter was defined as the lateral margin of the third trochanter to the medial border to the femoral shaft (Figure 3A') (18). Standard cortical and trabecular morphometric properties were calculated in each VOI (BV = bone volume; TV = total volume; I_{max} = maximum moment of inertia; Th = thickness, Ar = area, Ma = marrow, Pm = perimeter, BMD = bone mineral density; BMC = bone mineral content; Sp. = spacing, N = number). Data were compared using a two-way ANOVA correcting for multiple

comparisons (Prism v9.0; Graphpad, LaJolla, CA). Data were presented as mean ± standard deviation.

Mechanical Testing

Contralateral femurs (n = 6 females per treatment; n=3 males) were loaded to failure in 4-point bending with a span length of 6.5mm and 2.2mm for the top and bottom fixtures, respectively (MTS Systems, 858 Mini Bionix II) (19,20). Fixtures were positioned at mid-femur with the posterior side of the femur under tension and samples were kept hydrated using PBS. Femur length was measured using calipers from the greater trochanter to the lateral femoral condyle. Femurs were loaded at 0.05mm/sec and loads were measured using a 50lb load cell (Honeywell). Maximum load (N), ultimate load (N), yield load (N), stiffness (N/mm), energy (Nmm), yield displacement (mm), ultimate displacement (mm), and failure displacement (mm) were calculated from the load–displacement curves. Data from female-born mice were compared using two-way ANOVAs and corrected for multiple comparisons using Tukey's multiple comparisons tests (Prism v9.0; Graphpad, LaJolla, CA).

Results

Bone size was influenced by GAHT, testosterone treatment with or without GnRHa administration lead to a significant decrease in femur length (Table 1: source of variation = T, p = 0.0011). Additionally, the shape of superstructures on long bones was influenced by GAHT (Figure 3A). Specifically, the width of the third trochanters was larger in mice treated with testosterone compared to control (without GnRHa or T administration) and GnRHa only treatment (Figure 3B, p = 0.0023). Mice treated with testosterone, with or without GnRHa, also had reduced cortical thickness adjacent to the third trochanter (Figure 3C).

Treatment with testosterone, independent of GnRHa treatment, also led to significant gain in trabecular bone volume and number (Figure 4A and B; Table 2). This change was also associated with decreased trabecular spacing (Figure 4B). Testosterone treatment with or without GnRHa led to increased trabecular bone mineral density (BMD) with testosterone-only reaching comparable levels as trabecular BMD for age-matched male mice (Figure 4C). For most of the trabecular outcomes measured, female-born mice had significantly different trabecular bone properties compared to age-matched male mice, especially without T treatment (Table 2). However, with the inclusion of T treatment in female-born mice, trabecular properties were more similar to age-matched male mice independent of GnRHa treatment (Table 2). Unlike trabecular changes, we did not observe changes to cortical bone properties with GnRHa or T treatment in female mice (Table 3). However, most cortical bone properties, with the exception of cortical BMD, were significantly different between female-born mice with and without GnRHa or T treatment compared to age matched male mice (Table 3).

The effect of testosterone accounted for ~40% of the variation in ultimate displacement (p=0.0013; Fig 5). T treatment accounted for 42.65% of the variation in femur length measured at time of biomechanical testing whereas GnRHa treatment accounted for 14.89% of the variation.

Figure 3. Third trochanter bone parameters in mice prepubertally treated with gonadotropin-releasing hormone analogues (GnRHa) and supplemented with gender-affirming hormones (GAH). (A') Transverse 3D reconstruction of femoral third trochanteric bone from P89 female mice displaying peak third trochanteric length measured in mm. (A'') Full 3D reconstruction of femoral third trochanteric bone region of interest from adult female mice. (B) Third trochanter width was larger in groups treated with T compared to control. (C) Cortical bone was thinner in female mice treated with T compared to control or GnRHa only. Data are presented as biological

replicates (individual dots) and mean +/- standard deviation. Gray horizontal bars represent range for age-matched male control mice, n=3.

Figure 4. Trabecular bone parameters in mice prepubertally treated with gonadotropin-releasing

hormone analogues (GnRHa) and supplemented with gender-affirming hormones (GAH). (A) 3D

reconstruction of femoral trabecular bone from P89 female mice. A representative image from each group is depicted. (B) T-treated femurs showed a significant increase in trabecular bone volume and trabecular bone volume fraction (BV/TV). (C) Trabecular density and number were increased in femurs treated with T; (D) inversely T-treated groups displayed a lower trabecular separation compared to non-T treated mice. Trabecular Data are presented as biological replicates (individual dots) and mean +/- standard deviation. Gray horizontal bars represent range for age-matched male control mice, n=3. ** $p < 0.01$; *** $p < 0.001$, **** $p < 0.0001$.

Figure 5. Ultimate displacement during fracture was significantly higher in female mice treated with T (with or without GnRHa) compared to GnRHa only mice. Data are presented as biological replicates (individual dots) and mean +/- standard deviation. Gray horizontal bars represent range for age-matched male control mice, n=3. * $p = 0.0117$; ** $p < 0.01$.

Table 1. Biomechanical outcomes for femurs following 4-pt bending. * indicates that the effect of GnRHa treatment was significant, p = 0.0371; ** indicates that the effect of T treatment was significant; p = 0.0013; *** indicates group is significantly different than female mice with and without GnRHa or T, p <0.01; ^ indicates group is significantly different than no T or GnRHa females, p<0.01; ↓ indicates group is significantly shorter compared to same GnRHa treatment with no T, p<0.01.

		No GnRHa			GnRHa-treated			Male		
		Mean	95% CI	N	Mean	95% CI	N	Mean	95% CI	N
Yield Load (N)	no T	19.12	19.59	5	19.12	19.59	5	22.95	13.01	3
	with T	18.78	18.57	6	16.98	12.51	5			
Ultimate load (N)	no T	24.21	14.48	5	25.07	4.01	6	31.08	1.92	3
	with T	25.99	9.18	6	25.62	4.83	5			
Stiffness (N/mm)	no T	139.78	13.82	5	153.65	9.82	6	7.01	41.87	3
	with T	151.03	10.40	6	148.35	7.33	5			
Energy (Nmm)	no T	8.49	40.95	5	9.51	44.30	6	7.01	41.87	3
	with T	9.27	26.15	6	9.54	22.94	5			
Yield displacement (mm)*	no T	0.15	17.73	6	0.12	11.97	6	0.13	20.98	3
	with T	0.14	24.03	6	0.13	7.83	5			
Ultimate displacement (mm)	no T	0.27	5.04	5	0.25	6.89	6	0.25	5.93	3
	with T**	0.29	7.25	6	0.29	7.60	5			
Femur length (mm)	no T	15.08**	2.36	6	15.34	1.55	6	15.76	2.08	3
	with T	14.53****↓	1.93	6	14.76***↓	1.74	5			

Table 2. Trabecular bone morphometry with and without GnRHA and T in female mice. If GnRHa was a source of variation (SoV) for morphometric outcomes, the associated p-values were noted in the morphometric column. If T was a SoV for morphometric outcomes, the signficance was noted in T-treatment columns as: * = p <0.05; *** = p <0.001; **** = p <0.0001. For outcomes in which male mice were significantly different than respective treatment groups, p-values are indicated in the treatment column as: * = p <0.05; *** = p <0.001; **** = p <0.0001.

		No GnRHa			GnRHa-treated			Male		
		Mean	95% CI	N	Mean	95% CI	N	Mean	95% CI	N
Tb anisotropy MIL	no T	0.377	57.657	7	0.432	40.617	7	0.338	11.722	3
	with T	0.362	32.750	7	0.375	25.403	7			
Tb BV (mm^3) GnRHa SoV, p<0.001	no T	0.094**	12.782	7	0.087***	25.118	7	0.289	2.277	3
	with T****	0.253	34.641	7	0.211	44.360	7			
Tb TV (mm^3) GnRHa SoV, p<0.001	no T	2.535*	9.778	7	2.632*	3.943	7	3.398	5.992	3
	with T	2.861	20.840	7	2.703	7.904	7			
BV/TV GnRHa SoV p=0.023	no T	0.037	11.319	7	0.033*	22.453	7	0.085	6.736	3
	with T****	0.091	37.625	7	0.079	44.000	7			
TbTh (mm) GnRHa SoV p=0.0003	no T	0.041**	9.088	7	0.043*	7.902	7	0.049	2.462	3
	with T	0.044*	2.444	7	0.045	6.494	7			
TbSp (mm) GnRHa SoV p=0.298	no T	0.308	19.371	7	0.322*	9.428	7	0.228	11.471	3
	with T****	0.201	17.221	7	0.225	17.940	7			
TbN (1/mm)	no T	2.920	13.131	7	2.754*	8.502	7	3.629	9.711	3
	with T****	4.139	12.645	7	3.766	14.186	7			
Tb BMD (mg HA/cm^3) GnRHa SoV p=0.0007	no T	111.975**	13.904	7	103.929**	22.457	7	228.359	15.178	3
	with T****	224.018	27.406	7	185.092	30.431	7			
Tb BMC (mg HA)	no T	0.285***	18.946	7	0.275****	25.015	7	0.772	11.490	3
	with T****	0.601	31.250	7	0.500	32.426	7			

		Mean	95% CI	N	Mean	95% CI	N	Mean	95% CI	N
GnRHa SoV p<0.0001										
Tb TMD (mg HA/cm^3)	no T	1146.584**	12.827	7	1259.507	7.184	7	1522.848	3.775	3
	with T*	1370.588**	8.297	7	1281.477	8.818	7			
Tb TMC (mg HA)	no T	0.108***	21.266	7	0.110***	30.551	7	0.440	5.262	3
	with T****	0.355	41.321	7	0.277	51.834	7			

Table 3. Cortical bone morphometry for female mice with and without GnRHa and T treatment. For outcomes in which male mice were significantly different than respective treatment groups, p-values are indicated in the treatment column as: * = p <0.05; *** = p <0.001; **** = p <0.0001.

		No GnRHa			GnRHa-treated			Male		
		Mean	95% CI	N	Mean	95% CI	N	Mean	95% CI	N
Cortical BMD (mg/cm^3)	no T	2284.874	2.574	7	2271.668	1.997	7	2310.110	2.770	3
	with T	2240.032	3.655	7	2221.611	2.838	7			
Cortical TV (mm^3)	no T	1.072****	3.963	7	1.113****	2.533	7	1.416	5.644	3
	with T	1.041****	7.007	7	1.051****	8.390	7			
Ct. thickness (mm^3)	no T	0.190****	2.555	7	0.193****	2.198	7	0.218	3.454	3
	with T	0.190****	2.432	7	0.188****	5.857	7			
Ct. perimeter (mm)	no T	7.881****	2.777	7	8.026***	1.624	7	8.970	4.782	3
	with T	8.0701***	4.445	7	8.112**	3.964	7			
Ct. area (mm^2)	no T	0.698****	2.988	7	0.722****	2.344	7	0.894	3.626	3
	with T	0.714****	5.569	7	0.711****	7.664	7			
Marrow area (mm^2)	no T	0.854**	6.036	7	0.872**	3.896	7	1.055	12.704	3
	with T	0.885**	9.116	7	0.897*	7.384	7			
I$_{max}$ (mm^4)	no T	0.1712****	8.649	7	0.187****	5.845	7	0.301	11.650	3
	with T	0.190****	13.354	7	0.192****	16.371	7			

Discussion

The effect of long pubertal suppression in transgender youth followed by sex steroid administration on bone growth and quality is a contemporary topic and important to investigate for the long-term health of transgender youth. Peak bone mass and body composition are strongly affected by sex steroids during puberty. High peak bone mass at skeletal maturity can predict a lower rate of fracture risk later in life (1–5). This is of special consideration for transgender youth who may elect to suppress puberty using GnRHa to delay or inhibit secondary sex characteristics. Estrogen plays a key role in bone regulation and growth, and conversion of testosterone to estrogen via aromatase or estrogen treatment in states of estrogen deficiencies can lead to increased bone remodeling. (21–25). Estrogen treatment can accelerate skeletal maturity, as well as induce senescence of proliferating growth plate cells, leading to early fusion of the epiphysis (21). We believe that testosterone conversion via aromatase could be an explanation for our finding of a decrease of femur lengths in our T treated groups (Table 1). Peak longitudinal growth velocity is also correlated with levels of growth hormone (GH) and insulin-like growth factor 1 (IGF-1) (5). Estrogen has been shown to increase GH secretion in boys and girls during puberty, and this increase is observed earlier in girls (5). Coincidentally, testosterone has a similar effect on GH and IGF-1 but much of the effect of testosterone is mediated through its aromatization to estrogen by the enzyme aromatase (5,26–28). Future studies should explore how changes in GH and IGF-1 are associated with gender affirming care in the context of bone health.

GnRHa can also induce a hypoestrogenic state resulting in bone density loss and reduced bone health (6). Prior to the widespread use of estrogen "add-back" therapy , some studies showed immediate bone density loss ranging from 2%-6% after 6 months of GnRHa therapy, but

this bone loss was recoverable (6). While the literature focused on the impact of gender-affirming therapy is growing, few have explored the effects of a hypoestrogenic period followed by subsequent GAH administration. Of note, in an adolescent mouse model of GAHT like ours, Dubois et al. showed that treatment of GnRHa led to fat mass accumulation, reduced lean mass and muscle strength, reduced of bone mass acquisition and bone strength, and increased bone marrow adiposity (8). Additionally, they found that T administration in prepubertal female mice can reverse the hypoestrogenic effects of GnRHa treatment (8). In the present study, we evaluated long bone (femur) quality in female-born mice subjected to peripubertal suppression followed by testosterone treatment. We found T-treatment with or without GnRHa during peripubertal growth had a strong effect on trabecular bone but only a marginal effect on cortical bone. Previous studies have demonstrated that bones with a well-developed trabecular network have increased bone strength (7–9). In our study, androgen administration led to a significant increase in trabecular bone density and number. We found that GnRHa treatment also resulted in longer femurs and, conversely, T administration led to shorter femurs, which could be explained by androgen-mediated acceleration of growth plate closure (29).

We also observed a visual difference in the shape of long bones, specifically focusing on the size of the third trochanter. With T treatment, we found the trochanter extended further from the centroid of the bone, and this may be caused by increased muscle contractility of the large gluteus maximus muscle. In young adults, gender affirming care for transgender men has been shown to improve knee extension and flexion strength (30). Further studies will begin to explore if there is an actual difference in trochanter size, begin analysis of a transfeminine model.

In animals and in humans, bone mass measurements in adulthood can be used to assess both peak bone mass acquisition as well as monitor the subsequent decline caused by aging or

gonadal insufficiency (22). Men with aromatase deficiency and women in menopause have lower bone mass, highlighting the impact of estrogen on bone maintenance (23). Conversely, women with polycystic ovarian syndrome (PCOS) and hyperandrogenism have increased trabecular BMD, and women with androgen insensitivity have lower BMD, suggesting that testosterone regulates BMD in women as well (23). In animal models, removal of gonadal sex hormone production in neonatal mice (e.g., ovariectomy or orchiectomy) can lead to restricted femoral bone growth during peripubertal development (23–25).

In summary this work demonstrates that delayed puberty in female-born mice does not significantly impair long bone quality or strength into young adulthood, and supplementation with gender affirming care (e.g., T) can lead to increased trabecular bone density. These gender-affirming therapies also influence the shape and length of long bones like the femur, which may correlate with changes in skeletal muscle function. These findings can help guide clinicians in their treatment of transgender adolescents by increasing the understanding of the effects of hormone treatment on long bones during a critical period of development.

Acknowledgements: This work was supported by funding from the National Institutes of Health NIAMS (R01AR079367 to MLK and P30AR069620) and NICHD (R01 HD098233 to MM, AS and VP); the National Science Foundation (CAREER 1944448 to MLK). Schematics designed using Biorender.

1. Rothman MS, Iwamoto SJ. Bone Health in the Transgender Population. Clin Rev Bone Miner Metab. 2019 Jun;17(2):77–85.

2. Fighera TM, Ziegelmann PK, Rasia da Silva T, Spritzer PM. Bone Mass Effects of Cross-Sex Hormone Therapy in Transgender People: Updated Systematic Review and Meta-Analysis. J Endocr Soc. 2019 May 1;3(5):943–64.

3. Almeida M, Laurent MR, Dubois V, Claessens F, O'Brien CA, Bouillon R, et al. Estrogens and Androgens in Skeletal Physiology and Pathophysiology. Physiol Rev. 2017 Jan;97(1):135–87.

4. Loomba-Albrecht LA, Styne DM. Effect of puberty on body composition: Curr Opin Endocrinol Diabetes Obes. 2009 Feb;16(1):10–5.

5. Singh D, Sanyal, Chattopadhyay N. The role of estrogen in bone growth and formation: changes at puberty. Cell Health Cytoskelet. 2010 Dec;1.

6. Sauerbrun-Cutler MT, Alvero R. Short- and long-term impact of gonadotropin-releasing hormone analogue treatment on bone loss and fracture. Fertil Steril. 2019 Nov;112(5):799–803.

7. Ortmann O, Weiss J, Diedrich K. Gonadotrophin-releasing hormone (GnRH) and GnRH agonists: mechanisms of action. Reprod Biomed Online. 2002 Jan;5:1–7.

8. Dubois V, Ciancia S, Doms S, El Kharraz S, Sommers V, Kim NR, et al. Testosterone Restores Body Composition, Bone Mass, and Bone Strength Following Early Puberty Suppression in a Mouse Model Mimicking the Clinical Strategy in Trans Boys. J Bone Miner Res. 2023 Jun 8;jbmr.4832.

9. Chew D, Anderson J, Williams K, May T, Pang K. Hormonal Treatment in Young People With Gender Dysphoria: A Systematic Review. 2018;141(4).

10. Martinerie L, Condat A, Bargiacchi A, Bremont-Weill C, De Vries MC, Hannema SE. MANAGEMENT OF ENDOCRINE DISEASE: Approach to the management of children and adolescents with Gender Dysphoria. Eur J Endocrinol. 2018 Nov;179(5):R219–37.

11. Coleman E, Radix AE, Bouman WP, Brown GR, De Vries ALC, Deutsch MB, et al. Standards of Care for the Health of Transgender and Gender Diverse People, Version 8. Int J Transgender Health. 2022 Aug 19;23(sup1):S1–259.

12. Hembree WC, Cohen-Kettenis PT, Gooren L, Hannema SE, Meyer WJ, Murad MH, et al. Endocrine Treatment of Gender-Dysphoric/Gender-Incongruent Persons: An Endocrine Society* Clinical Practice Guideline. J Clin Endocrinol Metab. 2017 Nov 1;102(11):3869–903.

13. Ciancia S, Dubois V, Cools M. Impact of gender-affirming treatment on bone health in transgender and gender diverse youth. Endocr Connect. 2022 Nov 1;11(11):e220280.

14. Piekarski DJ, Boivin JR, Wilbrecht L. Ovarian Hormones Organize the Maturation of Inhibitory Neurotransmission in the Frontal Cortex at Puberty Onset in Female Mice. Curr Biol. 2017 Jun;27(12):1735-1745.e3.

15. Bouxsein ML, Boyd SK, Christiansen BA, Guldberg RE, Jepsen KJ, Müller R. Guidelines for assessment of bone microstructure in rodents using micro-computed tomography. J Bone Miner Res. 2010 Jun 7;25(7):1468–86.

16. Bell KL, Garrahan N, Kneissel M, Loveridge N, Grau E, Stanton M, et al. Cortical and cancellous bone in the human femoral neck: Evaluation of an interactive image analysis system. Bone. 1996 Nov;19(5):541–8.

17. Bell KL, Loveridge N, Power J, Garrahan N, Meggitt BF, Reeve J. Regional differences in cortical porosity in the fractured femoral neck. Bone. 1999 Jan;24(1):57–64.

18. Hamrick MW, McPherron AC, Lovejoy CO, Hudson J. Femoral morphology and cross-sectional geometry of adult myostatin-deficient mice. Bone. 2000 Sep;27(3):343–9.

19. Patton DM, Bigelow EMR, Schlecht SH, Kohn DH, Bredbenner TL, Jepsen KJ. The relationship between whole bone stiffness and strength is age and sex dependent. J Biomech. 2019 Jan;83:125–33.

20. Jepsen KJ, Centi A, Duarte GF, Galloway K, Goldman H, Hampson N, et al. Biological constraints that limit compensation of a common skeletal trait variant lead to inequivalence of tibial function among healthy young adults. J Bone Miner Res. 2011 Dec;26(12):2872–85.

21. Weise M, De-Levi S, Barnes KM, Gafni RI, Abad V, Baron J. Effects of estrogen on growth plate senescence and epiphyseal fusion. Proc Natl Acad Sci. 2001 Jun 5;98(12):6871–6.

22. Glatt V, Canalis E, Stadmeyer L, Bouxsein ML. Age-Related Changes in Trabecular Architecture Differ in Female and Male C57BL/6J Mice. J Bone Miner Res. 2007 May 7;22(8):1197–207.

23. Ornoy A. Gender dependent effects of testosterone and 17/3- estradiol on bone growth and modelling in young mice. :16.

24. Marquez Hernandez RA, Ohtani J, Fujita T, Sunagawa H, Kawata T, Kaku M, et al. Sex hormones receptors play a crucial role in the control of femoral and mandibular growth in newborn mice. Eur J Orthod. 2011 Oct 1;33(5):564–9.

25. Seidlová-Wuttke D, Jarry H, Pitzel L, Wuttke W. Effects of estradiol-17β, testosterone and a black cohosh preparation on bone and prostate in orchidectomized rats. Maturitas. 2005 Jun;51(2):177–86.

26. Mauras N. Growth hormone and sex steroids. Endocrinol Metab Clin North Am. 2001 Sep;30(3):529–44.

27. Veldhuis JD, Roemmich JN, Rogol AD. Gender and Sexual Maturation-Dependent Contrasts in the Neuroregulation of Growth Hormone Secretion in Prepubertal and Late Adolescent Males and Females—A General Clinical Research Center-Based Study. 2000;85(7).

28. Juul A. The effects of oestrogens on linear bone growth. Hum Reprod Update. 2001 May 1;7(3):303–13.

29. Nilsson LO, Boman A, Sävendahl L, Grigelioniene G, Ohlsson C, Ritzén EM, et al. Demonstration of Estrogen Receptor-β Immunoreactivity in Human Growth Plate Cartilage. J Clin Endocrinol Metab. 1999 Jan;84(1):370–3.

30. Wiik A, Lundberg TR, Rullman E, Andersson DP, Holmberg M, Mandić M, et al. Muscle Strength, Size, and Composition Following 12 Months of Gender-affirming Treatment in Transgender Individuals. J Clin Endocrinol Metab. 2020 Mar 1;105(3):e805–13.