1   Novel Evolution of the Mineralocorticoid Receptor in Humans compared to Chimpanzees,

2   Gorillas and Orangutans

4   Yoshinao Katsu[1,2,] *, Jiawen Zhang[2], Michael E. Baker[3,4,] *

5   [1] Faculty of Science

6   Hokkaido University

7   Sapporo, Japan

8   [2] Graduate School of Life Science

9   Hokkaido University

10   Sapporo, Japan

11   [3] Division of Nephrology-Hypertension

12   Department of Medicine, 0693

13   University of California, San Diego

14   9500 Gilman Drive

15   La Jolla, CA

16   Center for Academic Research and Training in Anthropogeny (CARTA) [4]

17   University of California, San Diego

18   La Jolla, CA

*Correspondence to
Y. Katsu; E-mail: ykatsu@sci.hokudai.ac.jp (YK)
M. E. Baker; E-mail: mbaker@health.ucsd.edu (MEB)

Abstract

We identified five distinct full-length human mineralocorticoid receptor (MR) genes

containing either 984 amino acids (MR-984) or 988 amino acids (MR-988), which can be

distinguished by the presence or absence of Lys, Cys, Ser, Trp (KCSW) in their DNA-binding

domain (DBD) and mutations at codons 180 and 241 in their amino terminal domain (NTD).

Two human MR-KCSW genes contain either (Val-180, Val-241) or (Ile-180, Val-241) in their

NTD and three human MR-984 genes contain either (Ile-180, Ala-241) or (Val-180, Val-241) or

(Ile-180, Val-241). Human MR-KCSW with (Ile-180, Ala-241) has not been cloned. In

contrast, chimpanzees contain four MRs: two MR-988s with KCSW in their DBD or two MR-

984s without KCSW in their DBD. Chimpanzee MRs only contain (Ile180, Val-241) in their

NTD. A chimpanzee MR with either (Val-180, Val-241) or (Ile-180, Ala-241) in the NTD has

not been cloned. Gorillas and orangutans each contain one MR-988 with KCSW in the DBD and

one MR-984 without KCSW, and these MRs only contain (Ile-180, Val-241) in their NTD. A

gorilla MR or orangutan MR with either (Val-180, Val-241) or (Ile-180, Ala-241) in the NTD

has not been cloned. Together, these data suggest that human MRs with (Val-180, Val-241) or

(Ile-180, Ala-241) in the NTD evolved after humans and chimpanzees diverged from their

common ancestor. Considering the multiple functions in human development of the MR in

kidney, brain, heart, skin, and lungs, as well as MR activity in interaction with the glucocorticoid

receptor, we suggest that the evolution of human MRs that are absent in chimpanzees may have

been important in the evolution of humans from chimpanzees. Studies are underway to

characterize transcriptional activation of the five human MRs by aldosterone, cortisol, and other

corticosteroids for comparison with each other to elucidate the roles of these MRs in human

physiology.

Running title: Evolution of the DNA-binding domain of the Mineralocorticoid Receptor

Introduction

The mineralocorticoid receptor (MR) is a ligand-activated transcription factor, belonging

to the nuclear receptor family, a diverse group of transcription factors that arose in multicellular

animals [1–5]. The traditional physiological function of the MR is to maintain electrolyte

balance by regulating sodium and potassium transport in epithelial cells in the kidney and colon

[6–10]. In addition, the MR has important physiological functions in many other tissues,

including brain, heart, skin and lungs [10–16].

The MR and its paralog, the glucocorticoid receptor (GR), descended from an ancestral

corticoid receptor (CR) in a cyclostome (jawless fish) that evolved about 550 million years ago

at the base of the vertebrate line [17–24]. A descendent of this ancestral steroid receptor, the CR

in lamprey (*Petromyzon marinus*), is activated by aldosterone [25,26] and other corticosteroids

[25,26]. Lampreys contain two CR isoforms, which differ only in the presence of a four amino

acid insert Thr, Arg, Gln, Gly (TRQG) in their DNA-binding domain (DBD) [27] (Figure 1).

We found that several corticosteroids had a similar half-maximal response (EC50) for lamprey

CR1 and CR2 [26]. However, these corticosteroids had a lower fold-activation of transcription

for CR1, which contains the four amino acid insert, than for CR2 suggesting that the deletion of

the four amino acid sequence in CR2 selected for increased transcriptional activation by

corticosteroids of CR2 [26,27].

A distinct MR and GR first appear in sharks and other cartilaginous fishes

(Chondrichthyes) [19,21,28–31]. The DBD in elephant shark MR and GR lacks the four amino

acid sequence found in lamprey CR1 [26] (Figure 1). We inserted this four-residue sequence

from lamprey CR1 into the DBD in elephant shark MR and GR and found that in HEK293 cells

co-transfected with the TAT3 promoter, the mutant elephant shark MR and GR had lower

transcriptional activation by corticosteroids than did their wild-type elephant shark MR and GR

counterparts, indicating that the insertion of the four amino acid sequence into the DBD of wild-

type elephant shark MR and GR had a similar effect on transcriptional activation as the KCSW

insert had in the DBD of lamprey CR1 [27].

We then analyzed the DBD sequence of human MR, which had been cloned, sequenced

and characterized by Arriza et al. [32], and found that like elephant shark MR, this human MR

(MR1) lacks a four-residue segment in its DBD. This human MR has been widely studied

[8,10,12,13,33–35]. Unexpectedly, our BLAST [36] search with the DBD from this human MR

found a second, previously described, human MR splice variant with a KCSW insert (MR-

KCSW) in its DBD [27,37–39] (Figure 1). A homology model of human MR with KCSW in the

DBD constructed by Wickert et al. [39] found no distortion by the four amino acids on adjacent

secondary structures of the DBD in human MR.

As described later, further BLAST searches found two full-length human MRs with this

insert (MR-KCSW) and three full-length human MRs without this insert. The three human MRs

without the KCSW insert contain either (Ile-180, Ala-241) or (Val-180, Val-241) or (Ile-180,

Val-241) in their amino terminal domain (NTD) (Figure 2). The two human MR-KCSW splice

variants contain either (Val-180, Val-241) or (Ile-180, Val-241) in their NTD (Figure 2). A

human MR-KCSW with (Ile-180, Ala-241) has not been cloned.

Transcriptional activation of any human MR-KCSW by corticosteroids has not been

reported. To remedy this deficiency, we have studies underway to characterize transcriptional

activation of the five full-length human MRs by a panel of corticosteroids. Here, in this brief

report, we describe our evolutionary analysis of human MR from a comparison of five full-

length human MRs with four full-length MRs in chimpanzees and two full-length MRs in

gorillas, and orangutans. We find that chimpanzees, gorillas, and orangutans lack an MR with

either (Ile-180, Ala-241) or (Val-180, Val-241) in their NTD. We propose that MRs with these

amino acids in the NTD evolved in humans after divergence of humans from chimpanzees. Due

to the multiple functions in human development of the MR alone [11,15,16,40], as well as due to

the interaction of the MR with the GR [41–45], we suggest that the evolution of three distinct

MRs in humans that are absent in chimpanzees may have been important in the evolution of

humans from chimpanzees.

```
Human MR1:          CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Human MR-KCSW:      CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG KCSWQHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Rat MR:            CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Rat MR-KCSW:       CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG KCSWQHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Platypus MR:       CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Platypus MR-KCSW:  CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG KCSWQHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Turtle MR:         CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Turtle MR-KCSW:    CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG KCSWQHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Xenopus MR:        CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QH SYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Xenopus MR-KCSR:   CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG KCSRQHSYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Lungfish MR:       CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACR VRKCLQAGM
Zebrafish MR:      CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACR VRKCLQAGM
Elephant shark MR:CLVCSDEASGCHYGVLTCGSCKVFFKRAVEG----QQNYLCAGRNDCIIDKIRRKNCPACRLRKCLKAGM
Lamprey CR1:       CLICSDEASGCHYGVLTCGSCKVFFKRAVEGTRQGQHNYLCAGRNDCIIDKIRRKNCPACRLRKCIQAGM
Lamprey CR2:       CLICSDEASGCHYGVLTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRLRKCIQAGM
```

Figure 1. Comparison of the DNA-binding domain on human MR1, human MR-KCSW,

rat MR, rat MR-KCSW, Platypus MR, Platypus MR-KCSW, Turtle MR, Turtle MR-

KCSW, Xenopus MR, Xenopus MR-KCSR, lungfish MR, zebrafish MR, elephant shark

MR, and lamprey CR1 and CR2.

Figure legend: The DBD of the human MR-KCSW splice variant has an insertion of four amino

acids that is absent in human MR1. Otherwise, the rest of the sequences of human MR and

human MR-KCSW are identical. Moreover, except for the four amino acid insert in the DBD of

human MR, the DBDs of rat MR, platypus MR and turtle MR are identical. The four amino acid

splice variant in the DBD of *Xenopus* MR differs from these MRs by one amino acid.

Differences between the DBD sequence in human MR and selected vertebrate MRs are shown in

red. Protein accession numbers are AAA59571 for human MR, XP_011530277 for human MR-

KCSW, NP_037263 for rat MR, XP_038953451 for rat MR-KCSW, XP_007669969 for

platypus MR, XP_016083764 for platypus MR-KCSW, XP_043401725 for turtle MR,

XP_037753298 for turtle MR-KCSW, NP_001084074 for *Xenopus* MR, XP_018098693 for

Xenopus MR-KCSR, BCV19931 for lungfish MR, NP_001093873 for zebrafish MR,

XP_007902220 for elephant shark MR, XP_032811370 for lamprey CR1, and XP_032811371

for lamprey CR2.

Results and Discussion

Humans have five full-length MRs.

A BLAST [36] search of GenBank retrieved five distinct full-length human MR genes

(Figure 2). We retrieved three full-length human MRs without a KCSW insert in the DBD.

These three human MRs contain either (Ile-180, Ala-241), (Val-180, Val-241) or (Ile-180, Val-

241) in the NTD (Figure 2). We also retrieved two human MR-KCSWs containing either (Val-

180, Val-241) or (Ile-180, Val-241) in their NTD. A human MR-KCSW sequence with (Ile-180,

Ala-241) has not been deposited in GenBank.

Figure 2. Comparison of five full-length human MRs.

Figure legend: There are three human MRs with 984 amino acids. They contain either valine-180 and valine-241, isoleucine-180 and valine-241 or isoleucine-180 and alanine-241 in the amino-terminal domain (NTD). There are two human MRs with 988 amino acids and KCSW in the DBD and either valine-180 and valine-241 or isoleucine-180 and valine-241 in the NTD. A human MR with KCSW and isoleucine-180 and alanine-241 in the NTD was not found in GenBank.

Chimpanzees have four full-length MRs.

Chimpanzees contain two full-length MRs that have DBDs that are identical to the DBD in human MRs without the KCSW insert and two MRs that have DBDs that are identical to the DBD in human MR-KCSW (Figure 1, Figure 3). All four chimpanzee MRs contain (Ile-180, Val-241) in their NTD, which is similar to the NTD in one human MR (Figure 2). Chimpanzees lack an MR with either (Ile-180, Ala-241) or (Val-180, Val-241) in their NTD. The different

149 chimpanzee MR sequences contain a Ser-591 to Asn-591 mutation in their NTD (Figure 3).

150 Human MRs contain a serine-591 corresponding to serine-591 in chimpanzee MR.

```
MR1:       METKGYHSLPEGLDMERRWGQVSQAVEHSSLGPTERTDENNYMEIVNVSCVSGAIPNNSTQGSSKEKHELLPCLQQDNNRPGILTSDIKT  90
MR2-KCSW:   METKGYHSLPEGLDMERRWGQVSQAVEHSSLGPTERTDENNYMEIVNVSCVSGAIPNNSTQGSSKEKHELLPCLQQDNNRPGILTSDIKT  90
MR3:       METKGYHSLPEGLDMERRWGQVSQAVEHSSLGPTERTDENNYMEIVNVSCVSGAIPNNSTQGSSKEKHELLPCLQQDNNRPGILTSDIKT  90
MR4-KCSW:   METKGYHSLPEGLDMERRWGQVSQAVEHSSLGPTERTDENNYMEIVNVSCVSGAIPNNSTQGSSKEKHELLPCLQQDNNRPGILTSDIKT  90
           ******************************************************************************************

MR1:       ELESKELSATVAESMGLYMDSVRDADYSYEQQNQQGSMSPAKIYQNVEQLVKFYKGNGHRPSTLSCVNRPLRSFMSDSGSSVNGGVMRAI  180
MR2-KCSW:   ELESKELSATVAESMGLYMDSVRDADYSYEQQNQQGSMSPAKIYQNVEQLVKFYKGNGHRPSTLSCVNRPLRSFMSDSGSSVNGGVMRAI  180
MR3:       ELESKELSATVAESMGLYMDSVRDADYSYEQQNQQGSMSPAKIYQNVEQLVKFYKGNGHRPSTLSCVNRPLRSFMSDSGSSVNGGVMRAI  180
MR4-KCSW:   ELESKELSATVAESMGLYMDSVRDADYSYEQQNQQGSMSPAKIYQNVEQLVKFYKGNGHRPSTLSCVNRPLRSFMSDSGSSVNGGVMRAI  180
           ******************************************************************************************

MR1:       VKSPIMCHEKSPSVCSPLNMTSSVCSPAGINSVSSTTASFGNFPVHSPITQGTPLTCSPNVENRGSRSHSPAHASNVGSPLSSPLSSMKS  270
MR2-KCSW:   VKSPIMCHEKSPSVCSPLNMTSSVCSPAGINSVSSTTASFGNFPVHSPITQGTPLTCSPNVENRGSRSHSPAHASNVGSPLSSPLSSMKS  270
MR3:       VKSPIMCHEKSPSVCSPLNMTSSVCSPAGINSVSSTTASFGNFPVHSPITQGTPLTCSPNVENRGSRSHSPAHASNVGSPLSSPLSSMKS  270
MR4-KCSW:   VKSPIMCHEKSPSVCSPLNMTSSVCSPAGINSVSSTTASFGNFPVHSPITQGTPLTCSPNVENRGSRSHSPAHASNVGSPLSSPLSSMKS  270
           ******************************************************************************************

MR1:       SISSPPSHCSVKSPVSSPNNVTLRSSVSSPANINNSRCSVSSPSNTNNRSTLSSPAASTVGSICSPVNNAFSYTASGTSAGSSTLRDVVP  360
MR2-KCSW:   SISSPPSHCSVKSPVSSPNNVTLRSSVSSPANINNSRCSVSSPSNTNNRSTLSSPAASTVGSICSPVNNAFSYTASGTSAGSSTLRDVVP  360
MR3:       SISSPPSHCSVKSPVSSPNNVTLRSSVSSPANINNSRCSVSSPSNTNNRSTLSSPAASTVGSICSPVNNAFSYTASGTSAGSSTLRDVVP  360
MR4-KCSW:   SISSPPSHCSVKSPVSSPNNVTLRSSVSSPANINNSRCSVSSPSNTNNRSTLSSPAASTVGSICSPVNNAFSYTASGTSAGSSTLRDVVP  360
           ******************************************************************************************

MR1:       SPDTQEKGAQEVPFPKTEEVESAISNGVTGQLNIVQYIKPEPDGAFSSSCLGGNSKINSDSSFSVPIKQESTKHSCSGTSFKGNPTVNPF  450
MR2-KCSW:   SPDTQEKGAQEVPFPKTEEVESAISNGVTGQLNIVQYIKPEPDGAFSSSCLGGNSKINSDSSFSVPIKQESTKHSCSGTSFKGNPTVNPF  450
MR3:       SPDTQEKGAQEVPFPKTEEVESAISNGVTGQLNIVQYIKPEPDGAFSSSCLGGNSKINSDSSFSVPIKQESTKHSCSGTSFKGNPTVNPF  450
MR4-KCSW:   SPDTQEKGAQEVPFPKTEEVESAISNGVTGQLNIVQYIKPEPDGAFSSSCLGGNSKINSDSSFSVPIKQESTKHSCSGTSFKGNPTVNPF  450
           ******************************************************************************************

MR1:       PFMDGSYFSFMDDKDYYSLSGILGPPVPGFDGNCEGSGFPVGIKQEPDDGSYYPEASIPSSAIVGVNSGGQSFHYRIGAQGTISLSRSAR  540
MR2-KCSW:   PFMDGSYFSFMDDKDYYSLSGILGPPVPGFDGNCEGSGFPVGIKQEPDDGSYYPEASIPSSAIVGVNSGGQSFHYRIGAQGTISLSRSAR  540
MR3:       PFMDGSYFSFMDDKDYYSLSGILGPPVPGFDGNCEGSGFPVGIKQEPDDGSYYPEASIPSSAIVGVNSGGQSFHYRIGAQGTISLSRSAR  540
MR4-KCSW:   PFMDGSYFSFMDDKDYYSLSGILGPPVPGFDGNCEGSGFPVGIKQEPDDGSYYPEASIPSSAIVGVNSGGQSFHYRIGAQGTISLSRSAR  540
           ******************************************************************************************

MR1:       DQSFQHLSSFPPVNTLVESWKSHGDLSSRRSDGYPVLEYIPENVSSSTLRSVSTGSSRPSKICLVCGDEASGCHYGVVTCGSCKVFFKRA  630
MR2-KCSW:   DQSFQHLSSFPPVNTLVESWKSHGDLSSRRSDGYPVLEYIPENVSSSTLRSVSTGSSRPSKICLVCGDEASGCHYGVVTCGSCKVFFKRA  630
MR3:       DQSFQHLSSFPPVNTLVESWKSHGDLSSRRSDGYPVLEYIPENVSSSTLRNVSTGSSRPSKICLVCGDEASGCHYGVVTCGSCKVFFKRA  630
MR4-KCSW:   DQSFQHLSSFPPVNTLVESWKSHGDLSSRRSDGYPVLEYIPENVSSSTLRNVSTGSSRPSKICLVCGDEASGCHYGVVTCGSCKVFFKRA  630
           ************************************************** ***********************************

MR1:       VEG----QHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGMNLGARKSKKLGKLKGIHEEQPQQQQPPPPPPPPQSPEEGTTYIAPAKE  716
MR2-KCSW:   VEGKCSWQHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGMNLGARKSKKLGKLKGIHEEQPQQQQPPPPPPPPQSPEEGTTYIAPAKE  720
MR3:       VEG----QHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGMNLGARKSKKLGKLKGIHEEQPQQQQPPPPPPPPQSPEEGTTYIAPAKE  716
MR4-KCSW:   VEGKCSWQHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGMNLGARKSKKLGKLKGIHEEQPQQQQPPPPPPPPQSPEEGTTYIAPAKE  720
           ***     *********************************************************************************

MR1:       PSVNTALVPQLSTISRALTPSPVMVLENIEPEIVYAGYDSSKPDTAENLLSTLNRLAGKQMIQVVKWAKVLPGFKNLPLEDQITLIQYSW  806
MR2-KCSW:   PSVNTALVPQLSTISRALTPSPVMVLENIEPEIVYAGYDSSKPDTAENLLSTLNRLAGKQMIQVVKWAKVLPGFKNLPLEDQITLIQYSW  810
MR3:       PSVNTALVPQLSTISRALTPSPVMVLENIEPEIVYAGYDSSKPDTAENLLSTLNRLAGKQMIQVVKWAKVLPGFKNLPLEDQITLIQYSW  806
MR4-KCSW:   PSVNTALVPQLSTISRALTPSPVMVLENIEPEIVYAGYDSSKPDTAENLLSTLNRLAGKQMIQVVKWAKVLPGFKNLPLEDQITLIQYSW  810
           ******************************************************************************************

MR1:       MCLSSFALSWRSYKHTNSQFLYFAPDLVFNEEKMHQSAMYELCQGMHQISLQFVRLQLTFEEYTIMKVLLLLSTIPKDGLKSQAAFEEMR  896
MR2-KCSW:   MCLSSFALSWRSYKHTNSQFLYFAPDLVFNEEKMHQSAMYELCQGMHQISLQFVRLQLTFEEYTIMKVLLLLSTIPKDGLKSQAAFEEMR  900
MR3:       MCLSSFALSWRSYKHTNSQFLYFAPDLVFNEEKMHQSAMYELCQGMHQISLQFVRLQLTFEEYTIMKVLLLLSTIPKDGLKSQAAFEEMR  896
MR4-KCSW:   MCLSSFALSWRSYKHTNSQFLYFAPDLVFNEEKMHQSAMYELCQGMHQISLQFVRLQLTFEEYTIMKVLLLLSTIPKDGLKSQAAFEEMR  900
           ******************************************************************************************

MR1:       TNYIKELRKMVTKCPNNSGQSWQRFYQLTKLLDSMHDLVSDLLEFCFYTFRESHALKVEFPAMLVEIISDQLPKVESGNAKPLYFHRK  984
MR2-KCSW:   TNYIKELRKMVTKCPNNSGQSWQRFYQLTKLLDSMHDLVSDLLEFCFYTFRESHALKVEFPAMLVEIISDQLPKVESGNAKPLYFHRK  988
MR3:       TNYIKELRKMVTKCPNNSGQSWQRFYQLTKLLDSMHDLVSDLLEFCFYTFRESHALKVEFPAMLVEIISDQLPKVESGNAKPLYFHRK  984
MR4-KCSW:   TNYIKELRKMVTKCPNNSGQSWQRFYQLTKLLDSMHDLVSDLLEFCFYTFRESHALKVEFPAMLVEIISDQLPKVESGNAKPLYFHRK  988
           ************************************************************************************
```

Figure 3. Multiple alignment of four chimpanzee MRs.

Figure legend: The four chimpanzee MRs were aligned with Clustal Omega [46]. MR1 and MR2-KCSW have serine-591 and MR3 and MR4-KCSW have asparagine-591 in the NTD. In contrast to human MRs, chimpanzee MRs only contain an isoleucine at codon 180 and a valine

156 at codon 241. Chimpanzees also contain an MR with either a serine or an asparagine at codon

157 591, while human MRs only contain a serine at codon 591.

159 **Gorillas and orangutans have two full-length MRs.**

160    Gorillas and orangutans each contain two full-length MRs: one MR with 984 amino acids

and one MR with 988 amino acids. The MRs with 988 amino acids have the KCSW sequence in

the DBD. All full-length gorilla MRs and orangutan MRs have isoleucine-180 and valine-241 in

the amino-terminal domain (NTD) (Figure 4). A gorilla MR or orangutan MR with either valine-

180 and valine-241 or isoleucine-180 and alanine-241 in the NTD was not found in GenBank.

Figure 4. Comparison of gorilla and orangutan MR sequences.

Figure legend: Gorilla MRs and Orangutan MRs conserve an isoleucine-180, valine-241 pair

corresponding to isoleucine-180 and valine-241 in the NTD of human MR. A gorilla MR or an

orangutan MR with either valine-180 and valine-241 or isoleucine-180 and alanine-241 in the

NTD was not found in GenBank. Gorillas and orangutans also only contain an MR with a serine

at codon 591. Human MRs only contain a serine at codon 591.

Evolutionary divergence of human and chimpanzee MRs.

Our analysis of human and chimpanzee MRs indicates that a human MR with either (Ile-180, Ala-241) or (Val-180, Val-241) in the NTD evolved after the divergence of humans and chimpanzees from a common ancestor. The physiological consequences of (Ile-180, Ala-241) or (Val-180, Val-241) in the NTD of human MR remain to be elucidated. Moreover, the absence of an asparagine at codon 591 in human MR is intriguing considering the presence of an asparagine at codon 591 in the MR in chimpanzees. As a first step, to determine if there are differences in transcriptional activation of these MRs by corticosteroids, we have cloned these human MR sequences and are screening them for transcriptional activation by aldosterone, cortisol, and other corticosteroids.

Evolution of the MR DBD in basal terrestrial vertebrates

The evolution of an MR with a DBD containing a four amino acid insert in human MR and other terrestrial MRs was surprising to us (Figure 1). We did not expect to find the KCSW insert in human MR at the position homologous to the position of TRQG in the DBD of lamprey CR because an insert at this position was not present in the DBD of elephant sharks and lungfish MRs (Figure 1). Moreover, the rest of the DBD sequence is highly conserved in terrestrial vertebrates (Figure 1) suggesting that the evolution of a four amino acid sequence in the DBD has an important function in terrestrial vertebrates. The evolution of the KCSR sequence into the DBD *Xenopus* MR (Figure 1) places the re-emergence of this motif close to the origin of terrestrial vertebrates. Evolution of aldosterone synthesis in lungfish is thought to be important in the evolution of terrestrial vertebrates [4,6,21,47–49]. The evolution of KCSR in Xenopus

MR in the homologous position to the DBD in human MR suggests that insertion of KCSR in

Xenopus MR also may have been important early in the evolution of terrestrial vertebrates.

Moreover, turtles contain an MR with KCSW in the homologous position in DBD (Figure 1) of

human MR DBD, indicating that KCSW also has an ancient origin in MRs in terrestrial

vertebrates. Indeed, the conservation of KCSW in the MR DBD in turtles and humans suggests

an important function for KCSW in terrestrial vertebrates.

Funding: This work was supported by Grants-in-Aid for Scientific Research from the Ministry

of Education, Culture, Sports, Science and Technology of Japan (23K05839) to Y.K., and the

Takeda Science Foundation to Y.K.

Competing interests: The authors have declared that no competing interests exist.

Contributions.

Yoshinao Katsu: Investigation, Conceptualization, Supervision, Formal Analysis, Writing –

original draft, Writing – review & editing.

Jiawen Zhang: Data curation, Investigation, Methodology.

Michael E. Baker: Conceptualization; Formal analysis, Supervision, Writing – original draft,

Writing – review & editing.

References

1. Evans RM. The steroid and thyroid hormone receptor superfamily. Science.
1988;240(4854):889-895. doi:10.1126/science.3283939.

2. Bridgham JT, Eick GN, Larroux C, et al. Protein evolution by molecular tinkering:
diversification of the nuclear receptor superfamily from a ligand-dependent ancestor. PLoS
Biol. 2010;8(10):e1000497. Published 2010 Oct 5. doi:10.1371/journal.pbio.1000497.

3. Baker ME, Nelson DR, Studer RA. Origin of the response to adrenal and sex steroids: Roles
of promiscuity and co-evolution of enzymes and steroid receptors. J Steroid Biochem Mol
Biol. 2015;151:12-24. doi:10.1016/j.jsbmb.2014.10.020.

4. Baker ME. Steroid receptors and vertebrate evolution. Mol Cell Endocrinol.
2019;496:110526. doi:10.1016/j.mce.2019.110526.

5. Evans RM, Mangelsdorf DJ. Nuclear Receptors, RXR, and the Big Bang. Cell. 2014 Mar
27;157(1):255-66. doi: 10.1016/j.cell.2014.03.012. PMID: 24679540; PMCID:
PMC4029515.

6. Rossier BC, Baker ME, Studer RA. Epithelial sodium transport and its control by
aldosterone: the story of our internal environment revisited. Physiol Rev. 2015;95(1):297-
340. doi:10.1152/physrev.00011.2014.

7. Hanukoglu I, Hanukoglu A. Epithelial sodium channel (ENaC) family: Phylogeny, structure-
function, tissue distribution, and associated inherited diseases. Gene. 2016;579(2):95-132.
doi:10.1016/j.gene.2015.12.061.

8. Lifton RP, Gharavi AG, Geller DS. Molecular mechanisms of human hypertension. Cell.
2001;104(4):545-556. doi:10.1016/s0092-8674(01)00241-0.

9. Shibata S. 30 YEARS OF THE MINERALOCORTICOID RECEPTOR: Mineralocorticoid
receptor and NaCl transport mechanisms in the renal distal nephron. J Endocrinol.
2017;234(1):T35-T47. doi:10.1530/JOE-16-0669.

10. Grossmann C, Almeida-Prieto B, Nolze A, Alvarez de la Rosa D. Structural and molecular
determinants of mineralocorticoid receptor signalling. Br J Pharmacol. 2021 Nov 22. doi:
10.1111/bph.15746. Epub ahead of print. PMID: 34811739.

11. Hawkins UA, Gomez-Sanchez EP, Gomez-Sanchez CM, Gomez-Sanchez CE. The
ubiquitous mineralocorticoid receptor: clinical implications. Curr Hypertens Rep.
2012;14(6):573-580. doi:10.1007/s11906-012-0297-0.

12. Jaisser F, Farman N. Emerging Roles of the Mineralocorticoid Receptor in Pathology:
Toward New Paradigms in Clinical Pharmacology. Pharmacol Rev. 2016;68(1):49-75.
doi:10.1124/pr.115.011106.

13. Funder J. 30 YEARS OF THE MINERALOCORTICOID RECEPTOR: Mineralocorticoid
receptor activation and specificity-conferring mechanisms: a brief history. J Endocrinol.
2017 Jul;234(1):T17-T21. doi: 10.1530/JOE-17-0119. Epub 2017 May 22. PMID:
28533421.

14. Gomez-Sanchez EP. Brain mineralocorticoid receptors in cognition and cardiovascular
homeostasis. Steroids. 2014 Dec;91:20-31. doi: 10.1016/j.steroids.2014.08.014. PMID:
25173821; PMCID: PMC4302001.

15. Joëls M, de Kloet ER. 30 YEARS OF THE MINERALOCORTICOID RECEPTOR: The
brain mineralocorticoid receptor: a saga in three episodes. J Endocrinol. 2017
Jul;234(1):T49-T66. doi: 10.1530/JOE-16-0660. PMID: 28634266.

16. Paul SN, Wingenfeld K, Otte C, Meijer OC. Brain mineralocorticoid receptor in health and
disease: From molecular signalling to cognitive and emotional function. Br J Pharmacol.
2022 Jul;179(13):3205-3219. doi: 10.1111/bph.15835. Epub 2022 Apr 7. PMID: 35297038;
PMCID: PMC9323486.

17. Shimeld SM, Donoghue PC. Evolutionary crossroads in developmental biology: cyclostomes
(lamprey and hagfish). Development. 2012 Jun;139(12):2091-9. doi: 10.1242/dev.074716.
PMID: 22619386.

18. Close DA, Yun SS, McCormick SD, Wildbill AJ, Li W. 11-deoxycortisol is a corticosteroid
hormone in the lamprey. Proc Natl Acad Sci U S A. 2010 Aug 3;107(31):13942-7. doi:
10.1073/pnas.0914026107. Epub 2010 Jul 19. PMID: 20643930; PMCID: PMC2922276.

19. Thornton JW. Evolution of vertebrate steroid receptors from an ancestral estrogen receptor
by ligand exploitation and serial genome expansions. Proc Natl Acad Sci U S A. 2001 May
8;98(10):5671-6. doi: 10.1073/pnas.091553298. Epub 2001 May 1. PMID: 11331759;
PMCID: PMC33271.

20. Smith JJ, Kuraku S, Holt C, Sauka-Spengler T, Jiang N, Campbell MS, Yandell MD,
Manousaki T, Meyer A, Bloom OE, Morgan JR, Buxbaum JD, Sachidanandam R, Sims C,
Garruss AS, Cook M, Krumlauf R, Wiedemann LM, Sower SA, Decatur WA, Hall JA,
Amemiya CT, Saha NR, Buckley KM, Rast JP, Das S, Hirano M, McCurley N, Guo P,
Rohner N, Tabin CJ, Piccinelli P, Elgar G, Ruffier M, Aken BL, Searle SM, Muffato M,
Pignatelli M, Herrero J, Jones M, Brown CT, Chung-Davidson YW, Nanlohy KG, Libants
SV, Yeh CY, McCauley DW, Langeland JA, Pancer Z, Fritzsch B, de Jong PJ, Zhu B,
Fulton LL, Theising B, Flicek P, Bronner ME, Warren WC, Clifton SW, Wilson RK, Li W.
Sequencing of the sea lamprey (Petromyzon marinus) genome provides insights into
vertebrate evolution. Nat Genet. 2013 Apr;45(4):415-21, 421e1-2. doi: 10.1038/ng.2568.
Epub 2013 Feb 24. PMID: 23435085; PMCID: PMC3709584.

21. Baker ME, Katsu Y. 30 YEARS OF THE MINERALOCORTICOID RECEPTOR:
Evolution of the mineralocorticoid receptor: sequence, structure and function. J Endocrinol.
2017;234(1):T1-T16. doi:10.1530/JOE-16-0661.

22. York JR, McCauley DW. The origin and evolution of vertebrate neural crest cells. Open
Biol. 2020 Jan;10(1):190285. doi: 10.1098/rsob.190285. Epub 2020 Jan 29. PMID:
31992146; PMCID: PMC7014683.

23. Kuratani S. Evo-devo studies of cyclostomes and the origin and evolution of jawed
vertebrates. Curr Top Dev Biol. 2021;141:207-239. doi: 10.1016/bs.ctdb.2020.11.011.
Epub 2020 Dec 13. PMID: 33602489.

24. Janvier P. microRNAs revive old views about jawless vertebrate divergence and evolution.
Proc Natl Acad Sci U S A. 2010 Nov 9;107(45):19137-8. doi: 10.1073/pnas.1014583107.
Epub 2010 Nov 1. PMID: 21041649; PMCID: PMC2984170.

25. Bridgham JT, Carroll SM, Thornton JW. Evolution of hormone-receptor complexity by
molecular exploitation. Science. 2006;312(5770):97-101. doi:10.1126/science.1123348.

26. Katsu Y, Lin X, Ji R, Chen Z, Kamisaka Y, Bamba K, Baker ME. N-terminal domain
influences steroid activation of the Atlantic sea lamprey corticoid receptor. J Steroid
Biochem Mol Biol. 2023 Apr;228:106249. doi: 10.1016/j.jsbmb.2023.106249. Epub 2023
Jan 13. PMID: 36646152.

27. Katsu Y, Zhang J, Baker ME. Reduced steroid activation of elephant shark GR and MR after
inserting four amino acids from the DNA-binding domain of lamprey corticoid receptor-1.
PLoS One. 2023 Aug 23;18(8):e0290159. doi: 10.1371/journal.pone.0290159. PMID:
37611044; PMCID: PMC10446182.

28. Venkatesh B, Lee AP, Ravi V, Maurya AK, Lian MM, Swann JB, Ohta Y, Flajnik MF,
Sutoh Y, Kasahara M, Hoon S, Gangu V, Roy SW, Irimia M, Korzh V, Kondrychyn I, Lim
ZW, Tay BH, Tohari S, Kong KW, Ho S, Lorente-Galdos B, Quilez J, Marques-Bonet T,
Raney BJ, Ingham PW, Tay A, Hillier LW, Minx P, Boehm T, Wilson RK, Brenner S,
Warren WC. Elephant shark genome provides unique insights into gnathostome evolution.
Nature. 2014 Jan 9;505(7482):174-9. doi: 10.1038/nature12826. Erratum in: Nature. 2014
Sep 25;513(7519):574. Erratum in: Nature. 2020 Dec;588(7837):E15. PMID: 24402279;
PMCID: PMC3964593.

29. Katsu Y, Kohno S, Oka K, et al. Transcriptional activation of elephant shark
mineralocorticoid receptor by corticosteroids, progesterone, and spironolactone. Sci Signal.
2019;12(584):eaar2668. Published 2019 Jun 4. doi:10.1126/scisignal.aar2668, (n.d.).

30. Fonseca E, Machado AM, Vilas-Arrondo N, Gomes-Dos-Santos A, Veríssimo A, Esteves P,
Almeida T, Themudo G, Ruivo R, Pérez M, da Fonseca R, Santos MM, Froufe E, Román-
Marcote E, Venkatesh B, Castro LFC. Cartilaginous fishes offer unique insights into the
evolution of the nuclear receptor gene repertoire in gnathostomes. Gen Comp Endocrinol.
2020 Sep 1;295:113527. doi: 10.1016/j.ygcen.2020.113527. Epub 2020 Jun 8. PMID:
32526329.

31. Carroll SM, Bridgham JT, Thornton JW. Evolution of hormone signaling in elasmobranchs
by exploitation of promiscuous receptors. Mol Biol Evol. 2008;25(12):2643-2652.
doi:10.1093/molbev/msn204.

32. Arriza JL, Weinberger C, Cerelli G, et al. Cloning of human mineralocorticoid receptor
complementary DNA: structural and functional kinship with the glucocorticoid receptor.
Science. 1987;237(4812):268-275. doi:10.1126/science.3037703.

33. Fuller PJ. Novel interactions of the mineralocorticoid receptor. Mol Cell Endocrinol. 2015
Jun 15;408:33-7. doi: 10.1016/j.mce.2015.01.027. Epub 2015 Feb 7. PMID: 25662276.

34. Hellal-Levy C, Couette B, Fagart J, Souque A, Gomez-Sanchez C, Rafestin-Oblin M.
Specific hydroxylations determine selective corticosteroid recognition by human
glucocorticoid and mineralocorticoid receptors. FEBS Lett. 1999;464(1-2):9-13.
doi:10.1016/s0014-5793(99)01667-1.

35. Geller DS, Farhi A, Pinkerton N, et al. Activating mineralocorticoid receptor mutation in
hypertension exacerbated by pregnancy. Science. 2000;289(5476):119-123.
doi:10.1126/science.289.5476.119.

36. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J
Mol Biol. 1990 Oct 5;215(3):403-10. doi: 10.1016/S0022-2836(05)80360-2. PMID:
2231712.

37. Bloem LJ, Guo C, Pratt JH. Identification of a splice variant of the rat and human
mineralocorticoid receptor genes. J Steroid Biochem Mol Biol. 1995 Nov;55(2):159-62.
doi: 10.1016/0960-0760(95)00162-s. PMID: 7495694.

38. Wickert L, Watzka M, Bolkenius U, Bidlingmaier F, Ludwig M. Mineralocorticoid receptor
splice variants in different human tissues. Eur J Endocrinol. 1998 Jun;138(6):702-4. doi:
10.1530/eje.0.1380702. PMID: 9678540.

39. Wickert L, Selbig J, Watzka M, Stoffel-Wagner B, Schramm J, Bidlingmaier F, Ludwig M.
Differential mRNA expression of the two mineralocorticoid receptor splice variants within
the human brain: structure analysis of their different DNA binding domains. J
Neuroendocrinol. 2000 Sep;12(9):867-73. doi: 10.1046/j.1365-2826.2000.00535.x. PMID:
10971811.

40. de Kloet ER, Joëls M. Brain mineralocorticoid receptor function in control of salt balance
and stress-adaptation. Physiol Behav. 2017 Sep 1;178:13-20. doi:
10.1016/j.physbeh.2016.12.045. Epub 2017 Jan 13. PMID: 28089704.

41. Liu W, Wang J, Sauter NK, Pearce D. Steroid receptor heterodimerization demonstrated in
vitro and in vivo. Proc Natl Acad Sci U S A. 1995 Dec 19;92(26):12480-4. doi:
10.1073/pnas.92.26.12480. PMID: 8618925; PMCID: PMC40381.

42. Trapp T, Holsboer F. Heterodimerization between mineralocorticoid and glucocorticoid
receptors increases the functional diversity of corticosteroid action. Trends Pharmacol Sci.
1996 Apr;17(4):145-9. doi: 10.1016/0165-6147(96)81590-2. PMID: 8984741.

43. Mifsud KR, Reul JM. Acute stress enhances heterodimerization and binding of corticosteroid
receptors at glucocorticoid target genes in the hippocampus. Proc Natl Acad Sci U S A.
2016 Oct 4;113(40):11336-11341. doi: 10.1073/pnas.1605246113. Epub 2016 Sep 21.
PMID: 27655894; PMCID: PMC5056104.

44. Pooley JR, Rivers CA, Kilcooley MT, Paul SN, Cavga AD, Kershaw YM, Muratcioglu S,
Gursoy A, Keskin O, Lightman SL. Beyond the heterodimer model for mineralocorticoid
and glucocorticoid receptor interactions in nuclei and at DNA. PLoS One. 2020 Jan
10;15(1):e0227520. doi: 10.1371/journal.pone.0227520. PMID: 31923266; PMCID:
PMC6953809.

45. Kiilerich P, Triqueneaux G, Christensen NM, et al. Interaction between the trout
mineralocorticoid and glucocorticoid receptors in vitro. J Mol Endocrinol. 2015;55(1):55-
68. doi:10.1530/JME-15-0002.

46. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
Remmert M, Söding J, Thompson JD, Higgins DG. Fast, scalable generation of high-
quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol. 2011
Oct 11;7:539. doi: 10.1038/msb.2011.75. PMID: 21988835; PMCID: PMC3261699.

47. Joss JM. Lungfish evolution and development. Gen Comp Endocrinol. 2006 Sep
15;148(3):285-9. doi: 10.1016/j.ygcen.2005.10.010. Epub 2005 Dec 7. PMID: 16337631.

48. Rossier BC. Osmoregulation during Long-Term Fasting in Lungfish and Elephant Seal: Old
and New Lessons for the Nephrologist. Nephron. 2016;134(1):5-9. doi:
10.1159/000444307. Epub 2016 Feb 23. PMID: 26901864.

49. McCormick SD, Bradshaw D. Hormonal control of salt and water balance in vertebrates.
Gen Comp Endocrinol. 2006 May 15;147(1):3-8. doi: 10.1016/j.ygcen.2005.12.009. Epub
2006 Feb 2. PMID: 16457828.