1   Novel Evolution of the Mineralocorticoid Receptor in Humans compared to Chimpanzees,

2   Gorillas and Orangutans

4   Yoshinao Katsu[1,2,] *, Jiawen Zhang[2], Michael E. Baker[3,4,] *

5   [1] Faculty of Science

6   Hokkaido University

7   Sapporo, Japan

8   [2] Graduate School of Life Science

9   Hokkaido University

10   Sapporo, Japan

11   [3] Division of Nephrology-Hypertension

12   Department of Medicine, 0693

13   University of California, San Diego

14   9500 Gilman Drive

15   La Jolla, CA

16   Center for Academic Research and Training in Anthropogeny (CARTA) [4]

17   University of California, San Diego

18   La Jolla, CA

*Correspondence to
Y. Katsu; E-mail: ykatsu@sci.hokudai.ac.jp (YK)
M. E. Baker; E-mail: mbaker@health.ucsd.edu (MEB)

Abstract

Five distinct full-length mineralocorticoid receptor (MR) genes have been identified in humans.

These human MRs can be distinguished by the presence or absence of an in-frame insertion of 12

base pairs coding for Lys, Cys, Ser, Trp (KCSW) in their DNA-binding domain (DBD) and the

presence of two amino acid mutations in their amino terminal domain (NTD). Two human MRs

with the KCSW insertion (MR-KCSW) and three human MRs without KCSW in the DBD have

been identified. The three human MRs without KCSW contain either (Ile-180, Ala-241) or (Val-

180, Val-241) or (Ile-180, Val-241) in their NTD. The two human MRs with KCSW contain

either (Val-180, Val-241) or (Ile-180, Val-241) in their NTD. Human MR-KCSW with (Ile-180,

Ala-241) has not been cloned. In contrast, chimpanzees contain two MRs with KCSW and two

MRs without KCSW in their DBD and both contain only Ile180, Val-241 in their NTDs. Each

pair of chimpanzee MRs differ at another amino acid in the NTD. A chimpanzee MR with either

Val-180, Val-241 or Ile-180, Ala-241 in the NTD has not been cloned. Gorillas and orangutans

each contain one MR with KCSW in the DBD and one MR without KCSW. Both gorilla and

orangutan MRs contain I-180, Val-241 in their NTD. Neither Val-180, Val-241 nor Ile-180,

Ala-241 are found in the NTD in either a gorilla MR or an orangutan MR. These data suggest

that human MRs with Val-180, Val-241 or Ile-180, Ala-241 in the NTD evolved after humans

and chimpanzees diverged from their common ancestor. These unique human MRs may have

had a role in the divergent evolution of humans from chimpanzees. Studies are underway to

characterize transcriptional activation of the five human MRs by aldosterone, cortisol, and other

corticosteroids for comparison with each other to elucidate the roles of these MRs in human

physiology.

Running title: Evolution of the DNA-binding domain of the Mineralocorticoid Receptor

Introduction

52   The mineralocorticoid receptor (MR) is a ligand-activated transcription factor, belonging

53   to the nuclear receptor family, a diverse group of transcription factors that arose in multicellular

54   animals [1–5]. The traditional physiological function of the MR is to maintain electrolyte

55   balance by regulating sodium and potassium transport in epithelial cells in the kidney and colon

56   [6–10]. In addition, the MR has important physiological functions in many other tissues,

57   including brain, heart, skin and lungs [10–16].

58   The MR and its paralog, the glucocorticoid receptor (GR), descended from an ancestral

corticoid receptor (CR) in a cyclostome (jawless fish) that evolved about 550 million years ago

at the base of the vertebrate line [17–24]. A descendent of this ancestral steroid receptor, the CR

in lamprey (*Petromyzon marinus*), is activated by aldosterone [25,26] and other corticosteroids

[25,26]. Lampreys contain two CR isoforms, which differ only in the presence of a four amino

acid insert Thr, Arg, Gln, Gly (TRQG) in their DNA-binding domain (DBD) [27] (Figure 1).

We found that several corticosteroids had a similar half-maximal response (EC50) for lamprey

CR1 and CR2 [26]. However, these corticosteroids had a lower fold-activation of transcription

for CR1, which contains the four amino acid insert, than for CR2 suggesting that the deletion of

the four amino acid sequence in CR2 selected for increased transcriptional activation by

corticosteroids of CR2 [26,27].

A distinct MR and GR first appear in sharks and other cartilaginous fishes

(Chondrichthyes) [19,21,28–31]. The DBD in elephant shark MR and GR lacks the four amino

acid sequence found in lamprey CR1 [26] (Figure 1). We inserted this four-residue sequence

from lamprey CR1 into the DBD in elephant shark MR and GR and found that in HEK293 cells

co-transfected with the TAT3 promoter, the mutant elephant shark MR and GR had lower

transcriptional activation by corticosteroids than did their wild-type elephant shark MR and GR

counterparts, indicating that the insertion of the four amino acid sequence into the DBD of wild-

type elephant shark MR and GR had a similar effect on transcriptional activation as the KCSW

insert had in the DBD of lamprey CR1 [27].

We then analyzed the DBD sequence of human MR, which had been cloned, sequenced

and characterized by Arriza et al. [32], and found that like elephant shark MR, this human MR

(MR1) lacks a four-residue segment in its DBD. This human MR has been widely studied

[8,10,12,13,33–35]. Unexpectedly, our BLAST [36] search with the DBD from this human MR

found a second, previously described, human MR splice variant with a KCSW insert (MR-

KCSW) in its DBD [27,37–39] (Figure 1). As described later, further BLAST searches found

two full-length human MRs with this insert (MR-KCSW) and three full-length human MRs

without this insert. The three human MRs without the KCSW insert contain either (Ile-180, Ala-

241) or (Val-180, Val-241) or (Ile-180, Val-241) in their amino terminal domain (NTD). The

two human MR-KCSW splice variants contain either (Val-180, Val-241) or (Ile-180, Val-241) in

their NTD. A human MR-KCSW with (Ile-180, Ala-241) has not been cloned.

Transcriptional activation of human MR-KCSW by corticosteroids has not been reported.

To remedy this deficiency, we have studies underway to characterize transcriptional activation of

the five full-length human MRs by a panel of corticosteroids. Here, in this brief report, we

describe our evolutionary analysis of human MR from a comparison of five full-length human

MRs with four full-length MRs in chimpanzees and two full-length MRs in gorillas, and

orangutans. We find that chimpanzees, gorillas, and orangutans lack an MR with either (Ile-180,

Ala-241) or (Val-180, Val-241) in their NTD. We propose that MRs with these amino acids in

the NTD evolved in humans after divergence of humans from chimpanzees. Due to the multiple

functions in human development of the MR alone [11,15,16,40], as well as due to the interaction

of the MR with the GR [41–45], we suggest that the evolution of three distinct MRs in humans

that are absent in chimpanzees may have been important in the evolution of humans from

chimpanzees.

```
Human MR1:          CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Human MR-KCSW:       CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG KCSWQHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Rat MR:             CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Rat MR-KCSW:        CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG KCSWQHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Platypus MR:        CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Platypus MR-KCSW:   CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG KCSWQHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Turtle MR:          CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Turtle MR-KCSW:     CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG KCSWQHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Xenopus MR:         CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QH SYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Xenopus MR-KCSR:    CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG KCSRQHSYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGM
Lungfish MR:        CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACR VRKCLQAGM
Zebrafish MR:       CLVCGDEASGCHYGVVTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACR VRKCLQAGM
Elephant shark MR:CLVCSDEASGCHYGVLTCGSCKVFFKRAVEG----QQNYLCAGRNDCIIDKIRRKNCPACRLRKCLKAGM
Lamprey CR1:        CLICSDEASGCHYGVLTCGSCKVFFKRAVEGTRQGQHNYLCAGRNDCIIDKIRRKNCPACRLRKCIQAGM
Lamprey CR2:        CLICSDEASGCHYGVLTCGSCKVFFKRAVEG----QHNYLCAGRNDCIIDKIRRKNCPACRL RKCIQAGM
```

Figure 1. Comparison of the DNA-binding domain on human MR1, human MR-KCSW,

rat MR, rat MR-KCSW, Platypus MR, Platypus MR-KCSW, Turtle MR, Turtle MR-

KCSW, Xenopus MR, Xenopus MR-KCSR, lungfish MR, zebrafish MR, elephant shark

MR, and lamprey CR1 and CR2.

Figure legend: The DBD of the human MR-KCSW splice variant has an insertion of four amino

acids that is absent in human MR1. Otherwise, the rest of the sequences of human MR and

human MR-KCSW are identical. Moreover, except for the four amino acid insert in the DBD of

human MR, the DBDs of rat MR, platypus MR and turtle MR are identical. The four amino acid

splice variant in the DBD of *Xenopus* MR differs from these MRs by one amino acid.

Differences between the DBD sequence in human MR and selected vertebrate MRs are shown in

red. Protein accession numbers are AAA59571 for human MR, XP_011530277 for human MR-

KCSW, NP_037263 for rat MR, XP_038953451 for rat MR-KCSW, XP_007669969 for

platypus MR, XP_016083764 for platypus MR-KCSW, XP_043401725 for turtle MR,

XP_037753298 for turtle MR-KCSW, NP_001084074 for *Xenopus* MR, XP_018098693 for

Xenopus MR-KCSR, BCV19931 for lungfish MR, NP_001093873 for zebrafish MR,

XP_007902220 for elephant shark MR, XP_032811370 for lamprey CR1, and XP_032811371

for lamprey CR2.

Results and Discussion

Humans have five full-length MRs

A BLAST [36] search of GenBank retrieved five distinct full-length human MR genes

(Figure 2). We retrieved three full-length human MRs without a KCSW insert in the DBD.

These three human MRs contain either (Ile-180, Ala-241), (Val-180, Val-241) or (Ile-180, Val-

241) in the NTD (Figure 2). We also retrieved two human MR-KCSWs containing either (Val-

180, Val-241) or (Ile-180, Val-241) in their NTD. A human MR-KCSW sequence with (Ile-180,

Ala-241) has not been deposited in GenBank.

Figure 2. Comparison of five full-length human MRs.

Figure legend: There are three human MRs with 984 amino acids. They contain either valine-

180 and valine-241, isoleucine-180 and valine-241 or isoleucine-180 and alanine-241 in the

amino-terminal domain (NTD). There are two human MRs with 988 amino acids and KCSW in

the DBD and either valine-180 and valine-241 or isoleucine-180 and valine-241 in the NTD. A

human MR with isoleucine-180 and alanine-241 and a KCSW insert in the NTD was not found

in GenBank.

Chimpanzees have four full-length MRs

Chimpanzees contain two full-length MRs that have DBDs that are identical to the DBD

in human MRs without the KCSW insert and two MRs that have DBDs that are identical to the

DBD in human MR-KCSW (Figure 1, Figure 3). All four chimpanzee MRs contain (Ile-180,

Val-241) in their NTD, which is similar to the NTD in one human MR (Figure 2). Chimpanzees

lack an MR with either (Ile-180, Ala-241) or (Val-180, Val-241) in their NTD. The different

chimpanzee MR sequences contain a Ser-591 to Asn-591 mutation in their NTD (Figure 3).

Human MRs contain a serine-591 corresponding to serine-591 in chimpanzee MR.

```
MR1:       METKGYHSLPEGLDMERRWGQVSQAVEHSSLGPTERTDENNYMEIVNVSCVSGAIPNNSTQGSSKEKHELLPCLQQDNNRPGILTSDIKT  90
MR2-KCSW:   METKGYHSLPEGLDMERRWGQVSQAVEHSSLGPTERTDENNYMEIVNVSCVSGAIPNNSTQGSSKEKHELLPCLQQDNNRPGILTSDIKT  90
MR3:       METKGYHSLPEGLDMERRWGQVSQAVEHSSLGPTERTDENNYMEIVNVSCVSGAIPNNSTQGSSKEKHELLPCLQQDNNRPGILTSDIKT  90
MR4-KCSW:   METKGYHSLPEGLDMERRWGQVSQAVEHSSLGPTERTDENNYMEIVNVSCVSGAIPNNSTQGSSKEKHELLPCLQQDNNRPGILTSDIKT  90
           ******************************************************************************************

MR1:       ELESKELSATVAESMGLYMDSVRDADYSYEQQNQQGSMSPAKIYQNVEQLVKFYKGNGHRPSTLSCVNRPLRSFMSDSGSSVNGGVMRAI 180
MR2-KCSW:   ELESKELSATVAESMGLYMDSVRDADYSYEQQNQQGSMSPAKIYQNVEQLVKFYKGNGHRPSTLSCVNRPLRSFMSDSGSSVNGGVMRAI 180
MR3:       ELESKELSATVAESMGLYMDSVRDADYSYEQQNQQGSMSPAKIYQNVEQLVKFYKGNGHRPSTLSCVNRPLRSFMSDSGSSVNGGVMRAI 180
MR4-KCSW:   ELESKELSATVAESMGLYMDSVRDADYSYEQQNQQGSMSPAKIYQNVEQLVKFYKGNGHRPSTLSCVNRPLRSFMSDSGSSVNGGVMRAI 180
           ******************************************************************************************

MR1:       VKSPIMCHEKSPSVCSPLNMTSSVCSPAGINSVSSTTASFGNFPVHSPITQGTPLTCSPNVENRGSRSHSPAHASNVGSPLSSPLSSMKS 270
MR2-KCSW:   VKSPIMCHEKSPSVCSPLNMTSSVCSPAGINSVSSTTASFGNFPVHSPITQGTPLTCSPNVENRGSRSHSPAHASNVGSPLSSPLSSMKS 270
MR3:       VKSPIMCHEKSPSVCSPLNMTSSVCSPAGINSVSSTTASFGNFPVHSPITQGTPLTCSPNVENRGSRSHSPAHASNVGSPLSSPLSSMKS 270
MR4-KCSW:   VKSPIMCHEKSPSVCSPLNMTSSVCSPAGINSVSSTTASFGNFPVHSPITQGTPLTCSPNVENRGSRSHSPAHASNVGSPLSSPLSSMKS 270
           ******************************************************************************************

MR1:       SISSPPSHCSVKSPVSSPNNVTLRSSVSSPANINNSRCSVSSPSNTNNRSTLSSPAASTVGSICSPVNNAFSYTASGTSAGSSTLRDVVP 360
MR2-KCSW:   SISSPPSHCSVKSPVSSPNNVTLRSSVSSPANINNSRCSVSSPSNTNNRSTLSSPAASTVGSICSPVNNAFSYTASGTSAGSSTLRDVVP 360
MR3:       SISSPPSHCSVKSPVSSPNNVTLRSSVSSPANINNSRCSVSSPSNTNNRSTLSSPAASTVGSICSPVNNAFSYTASGTSAGSSTLRDVVP 360
MR4-KCSW:   SISSPPSHCSVKSPVSSPNNVTLRSSVSSPANINNSRCSVSSPSNTNNRSTLSSPAASTVGSICSPVNNAFSYTASGTSAGSSTLRDVVP 360
           ******************************************************************************************

MR1:       SPDTQEKGAQEVPFPKTEEVESAISNGVTGQLNIVQYIKPEPDGAFSSSCLGGNSKINSDSSFSVPIKQESTKHSCSGTSFKGNPTVNPF 450
MR2-KCSW:   SPDTQEKGAQEVPFPKTEEVESAISNGVTGQLNIVQYIKPEPDGAFSSSCLGGNSKINSDSSFSVPIKQESTKHSCSGTSFKGNPTVNPF 450
MR3:       SPDTQEKGAQEVPFPKTEEVESAISNGVTGQLNIVQYIKPEPDGAFSSSCLGGNSKINSDSSFSVPIKQESTKHSCSGTSFKGNPTVNPF 450
MR4-KCSW:   SPDTQEKGAQEVPFPKTEEVESAISNGVTGQLNIVQYIKPEPDGAFSSSCLGGNSKINSDSSFSVPIKQESTKHSCSGTSFKGNPTVNPF 450
           ******************************************************************************************

MR1:       PFMDGSYFSFMDDKDYYSLSGILGPPVPGFDGNCEGSGFPVGIKQEPDDGSYYPEASIPSSAIVGVNSGGQSFHYRIGAQGTISLSRSAR 540
MR2-KCSW:   PFMDGSYFSFMDDKDYYSLSGILGPPVPGFDGNCEGSGFPVGIKQEPDDGSYYPEASIPSSAIVGVNSGGQSFHYRIGAQGTISLSRSAR 540
MR3:       PFMDGSYFSFMDDKDYYSLSGILGPPVPGFDGNCEGSGFPVGIKQEPDDGSYYPEASIPSSAIVGVNSGGQSFHYRIGAQGTISLSRSAR 540
MR4-KCSW:   PFMDGSYFSFMDDKDYYSLSGILGPPVPGFDGNCEGSGFPVGIKQEPDDGSYYPEASIPSSAIVGVNSGGQSFHYRIGAQGTISLSRSAR 540
           ******************************************************************************************

MR1:       DQSFQHLSSFPPVNTLVESWKSHGDLSSRRSDGYPVLEYIPENVSSSTLRSVSTGSSRPSKICLVCGDEASGCHYGVVTCGSCKVFFKRA 630
MR2-KCSW:   DQSFQHLSSFPPVNTLVESWKSHGDLSSRRSDGYPVLEYIPENVSSSTLRSVSTGSSRPSKICLVCGDEASGCHYGVVTCGSCKVFFKRA 630
MR3:       DQSFQHLSSFPPVNTLVESWKSHGDLSSRRSDGYPVLEYIPENVSSSTLRNVSTGSSRPSKICLVCGDEASGCHYGVVTCGSCKVFFKRA 630
MR4-KCSW:   DQSFQHLSSFPPVNTLVESWKSHGDLSSRRSDGYPVLEYIPENVSSSTLRNVSTGSSRPSKICLVCGDEASGCHYGVVTCGSCKVFFKRA 630
           *************************************************** *************************************

MR1:       VEG----QHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGMNLGARKSKKLGKLKGIHEEQPQQQQPPPPPPPPQSPEEGTTYIAPAKE 716
MR2-KCSW:   VEGKCSWQHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGMNLGARKSKKLGKLKGIHEEQPQQQQPPPPPPPPQSPEEGTTYIAPAKE 720
MR3:       VEG----QHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGMNLGARKSKKLGKLKGIHEEQPQQQQPPPPPPPPQSPEEGTTYIAPAKE 716
MR4-KCSW:   VEGKCSWQHNYLCAGRNDCIIDKIRRKNCPACRLQKCLQAGMNLGARKSKKLGKLKGIHEEQPQQQQPPPPPPPPQSPEEGTTYIAPAKE 720
           ***     **********************************************************************************

MR1:       PSVNTALVPQLSTISRALTPSPVMVLENIEPEIVYAGYDSSKPDTAENLLSTLNRLAGKQMIQVVKWAKVLPGFKNLPLEDQITLIQYSW 806
MR2-KCSW:   PSVNTALVPQLSTISRALTPSPVMVLENIEPEIVYAGYDSSKPDTAENLLSTLNRLAGKQMIQVVKWAKVLPGFKNLPLEDQITLIQYSW 810
MR3:       PSVNTALVPQLSTISRALTPSPVMVLENIEPEIVYAGYDSSKPDTAENLLSTLNRLAGKQMIQVVKWAKVLPGFKNLPLEDQITLIQYSW 806
MR4-KCSW:   PSVNTALVPQLSTISRALTPSPVMVLENIEPEIVYAGYDSSKPDTAENLLSTLNRLAGKQMIQVVKWAKVLPGFKNLPLEDQITLIQYSW 810
           ******************************************************************************************

MR1:       MCLSSFALSWRSYKHTNSQFLYFAPDLVFNEEKMHQSAMYELCQGMHQISLQFVRLQLTFEEYTIMKVLLLLSTIPKDGLKSQAAFEEMR 896
MR2-KCSW:   MCLSSFALSWRSYKHTNSQFLYFAPDLVFNEEKMHQSAMYELCQGMHQISLQFVRLQLTFEEYTIMKVLLLLSTIPKDGLKSQAAFEEMR 900
MR3:       MCLSSFALSWRSYKHTNSQFLYFAPDLVFNEEKMHQSAMYELCQGMHQISLQFVRLQLTFEEYTIMKVLLLLSTIPKDGLKSQAAFEEMR 896
MR4-KCSW:   MCLSSFALSWRSYKHTNSQFLYFAPDLVFNEEKMHQSAMYELCQGMHQISLQFVRLQLTFEEYTIMKVLLLLSTIPKDGLKSQAAFEEMR 900
           ******************************************************************************************

MR1:       TNYIKELRKMVTKCPNNSGQSWQRFYQLTKLLDSMHDLVSDLLEFCFYTFRESHALKVEFPAMLVEIISDQLPKVESGNAKPLYFHRK 984
MR2-KCSW:   TNYIKELRKMVTKCPNNSGQSWQRFYQLTKLLDSMHDLVSDLLEFCFYTFRESHALKVEFPAMLVEIISDQLPKVESGNAKPLYFHRK 988
MR3:       TNYIKELRKMVTKCPNNSGQSWQRFYQLTKLLDSMHDLVSDLLEFCFYTFRESHALKVEFPAMLVEIISDQLPKVESGNAKPLYFHRK 984
MR4-KCSW:   TNYIKELRKMVTKCPNNSGQSWQRFYQLTKLLDSMHDLVSDLLEFCFYTFRESHALKVEFPAMLVEIISDQLPKVESGNAKPLYFHRK 988
           ********************************************************************************************
```

Figure 3. Multiple alignment of four chimpanzee MRs.

Figure legend: The four chimpanzee MRs were aligned with Clustal Omega [46]. MR1 and

MR2-KCSW have serine-591 and MR3 and MR4-KCSW have asparagine-591 in the NTD.

Gorillas and orangutans have two full-length MRs

Gorillas and orangutans each contain two full-length MRs: one MR with 984 amino acids

and one MR with 988 amino acids. The MRs with 988 amino acids have the KCSW sequence in

the DBD. All full-length gorilla MRs and orangutan MRs have isoleucine-180 and valine-241 in

the amino-terminal domain (NTD) (Figure 4). A gorilla MR or orangutan MR with either valine-

180 and valine-241 or isoleucine-180 and alanine-241 in the NTD was not found in GenBank.

Figure 4. Comparison of gorilla and orangutan MR sequences.

Figure legend: Gorilla MRs and Orangutan MRs conserve an isoleucine-180, valine-241 pair

corresponding to isoleucine-180 and valine-241 in the NTD of human MR. A gorilla MR or an

orangutan MR with either valine-180 and valine-241 or isoleucine-180 and alanine-241 in the

NTD was not found in GenBank.

Evolutionary divergence of human and chimpanzee MRs

Our analysis of human and chimpanzee MRs indicates that human MR with either (Ile-

180, Ala-241) or (Val-180, Val-241) in the NTD evolved after the divergence of humans and

chimpanzees from a common ancestor. The physiological consequences of (Ile-180, Ala-241) or

(Val-180, Val-241) in the NTD of human MR remain to be elucidated. As a first step, to

determine if there are differences in transcriptional activation of these MRs by corticosteroids,

we have cloned these human MR sequences and are screening them for transcriptional activation

by aldosterone, cortisol, and other corticosteroids.

Evolution of the MR DBD in basal terrestrial vertebrates

The evolution of an MR with a DBD containing a four amino acid insert in human MR

and other terrestrial MRs was surprising to us (Figure 1). We did not expect to find the KCSW

insert in human MR at the position homologous to the position of TRQG in the DBD of lamprey

CR because an insert at this position was not present in the DBD of elephant sharks and lungfish

MRs (Figure 1). Moreover, the rest of the DBD sequence is highly conserved in terrestrial

vertebrates (Figure 1) suggesting that the evolution of this four amino acid sequence in the DBD

has an important function in terrestrial vertebrates. The evolution of the KCSR sequence into the

DBD *Xenopus* MR (Figure 1) places the re-emergence of this motif close to the origin of

terrestrial vertebrates. Moreover, turtles contain an MR with KCSW in the homologous position

in DBD (Figure 1) of human MR DBD, indicating that KCSW also has an ancient origin in

terrestrial vertebrate MRs. The conservation of KCSW in the MR DBD in turtles and humans

suggests an important function for KCSW in terrestrial vertebrates.

Funding: This work was supported by Grants-in-Aid for Scientific Research from the Ministry

of Education, Culture, Sports, Science and Technology of Japan (23K05839) to Y.K., and the

Takeda Science Foundation to Y.K.

Competing interests: The authors have declared that no competing interests exist.

Contributions.

Yoshinao Katsu: Investigation, Conceptualization, Supervision, Formal Analysis, Writing –

original draft, Writing – review & editing.

Jiawen Zhan: Data curation, Investigation, Methodology.

Michael E. Baker: Conceptualization; Formal analysis, Supervision, Writing – original draft,

Writing – review & editing.

References

1. Evans RM. The steroid and thyroid hormone receptor superfamily. Science.
1988;240(4854):889-895. doi:10.1126/science.3283939.

2. Bridgham JT, Eick GN, Larroux C, et al. Protein evolution by molecular tinkering:
diversification of the nuclear receptor superfamily from a ligand-dependent ancestor. PLoS
Biol. 2010;8(10):e1000497. Published 2010 Oct 5. doi:10.1371/journal.pbio.1000497.

3. Baker ME, Nelson DR, Studer RA. Origin of the response to adrenal and sex steroids: Roles
of promiscuity and co-evolution of enzymes and steroid receptors. J Steroid Biochem Mol
Biol. 2015;151:12-24. doi:10.1016/j.jsbmb.2014.10.020.

4. Baker ME. Steroid receptors and vertebrate evolution. Mol Cell Endocrinol.
2019;496:110526. doi:10.1016/j.mce.2019.110526.

5. Evans RM, Mangelsdorf DJ. Nuclear Receptors, RXR, and the Big Bang. Cell. 2014 Mar
27;157(1):255-66. doi: 10.1016/j.cell.2014.03.012. PMID: 24679540; PMCID:
PMC4029515.

6. Rossier BC, Baker ME, Studer RA. Epithelial sodium transport and its control by
aldosterone: the story of our internal environment revisited. Physiol Rev. 2015;95(1):297-
340. doi:10.1152/physrev.00011.2014.

7. Hanukoglu I, Hanukoglu A. Epithelial sodium channel (ENaC) family: Phylogeny, structure-
function, tissue distribution, and associated inherited diseases. Gene. 2016;579(2):95-132.
doi:10.1016/j.gene.2015.12.061.

8. Lifton RP, Gharavi AG, Geller DS. Molecular mechanisms of human hypertension. Cell.
2001;104(4):545-556. doi:10.1016/s0092-8674(01)00241-0.

9. Shibata S. 30 YEARS OF THE MINERALOCORTICOID RECEPTOR: Mineralocorticoid
receptor and NaCl transport mechanisms in the renal distal nephron. J Endocrinol.
2017;234(1):T35-T47. doi:10.1530/JOE-16-0669.

10. Grossmann C, Almeida-Prieto B, Nolze A, Alvarez de la Rosa D. Structural and molecular
determinants of mineralocorticoid receptor signalling. Br J Pharmacol. 2021 Nov 22. doi:
10.1111/bph.15746. Epub ahead of print. PMID: 34811739.

11. Hawkins UA, Gomez-Sanchez EP, Gomez-Sanchez CM, Gomez-Sanchez CE. The
ubiquitous mineralocorticoid receptor: clinical implications. Curr Hypertens Rep.
2012;14(6):573-580. doi:10.1007/s11906-012-0297-0.

12. Jaisser F, Farman N. Emerging Roles of the Mineralocorticoid Receptor in Pathology:
Toward New Paradigms in Clinical Pharmacology. Pharmacol Rev. 2016;68(1):49-75.
doi:10.1124/pr.115.011106.

13. Funder J. 30 YEARS OF THE MINERALOCORTICOID RECEPTOR: Mineralocorticoid
receptor activation and specificity-conferring mechanisms: a brief history. J Endocrinol.
2017 Jul;234(1):T17-T21. doi: 10.1530/JOE-17-0119. Epub 2017 May 22. PMID:
28533421.

14. Gomez-Sanchez EP. Brain mineralocorticoid receptors in cognition and cardiovascular
homeostasis. Steroids. 2014 Dec;91:20-31. doi: 10.1016/j.steroids.2014.08.014. PMID:
25173821; PMCID: PMC4302001.

15. Joëls M, de Kloet ER. 30 YEARS OF THE MINERALOCORTICOID RECEPTOR: The
brain mineralocorticoid receptor: a saga in three episodes. J Endocrinol. 2017
Jul;234(1):T49-T66. doi: 10.1530/JOE-16-0660. PMID: 28634266.

16. Paul SN, Wingenfeld K, Otte C, Meijer OC. Brain mineralocorticoid receptor in health and
disease: From molecular signalling to cognitive and emotional function. Br J Pharmacol.
2022 Jul;179(13):3205-3219. doi: 10.1111/bph.15835. Epub 2022 Apr 7. PMID: 35297038;
PMCID: PMC9323486.

17. Shimeld SM, Donoghue PC. Evolutionary crossroads in developmental biology: cyclostomes
(lamprey and hagfish). Development. 2012 Jun;139(12):2091-9. doi: 10.1242/dev.074716.
PMID: 22619386.

18. Close DA, Yun SS, McCormick SD, Wildbill AJ, Li W. 11-deoxycortisol is a corticosteroid
hormone in the lamprey. Proc Natl Acad Sci U S A. 2010 Aug 3;107(31):13942-7. doi:
10.1073/pnas.0914026107. Epub 2010 Jul 19. PMID: 20643930; PMCID: PMC2922276.

19. Thornton JW. Evolution of vertebrate steroid receptors from an ancestral estrogen receptor
by ligand exploitation and serial genome expansions. Proc Natl Acad Sci U S A. 2001 May
8;98(10):5671-6. doi: 10.1073/pnas.091553298. Epub 2001 May 1. PMID: 11331759;
PMCID: PMC33271.

20. Smith JJ, Kuraku S, Holt C, Sauka-Spengler T, Jiang N, Campbell MS, Yandell MD,
Manousaki T, Meyer A, Bloom OE, Morgan JR, Buxbaum JD, Sachidanandam R, Sims C,
Garruss AS, Cook M, Krumlauf R, Wiedemann LM, Sower SA, Decatur WA, Hall JA,
Amemiya CT, Saha NR, Buckley KM, Rast JP, Das S, Hirano M, McCurley N, Guo P,
Rohner N, Tabin CJ, Piccinelli P, Elgar G, Ruffier M, Aken BL, Searle SM, Muffato M,
Pignatelli M, Herrero J, Jones M, Brown CT, Chung-Davidson YW, Nanlohy KG, Libants
SV, Yeh CY, McCauley DW, Langeland JA, Pancer Z, Fritzsch B, de Jong PJ, Zhu B,
Fulton LL, Theising B, Flicek P, Bronner ME, Warren WC, Clifton SW, Wilson RK, Li W.
Sequencing of the sea lamprey (Petromyzon marinus) genome provides insights into
vertebrate evolution. Nat Genet. 2013 Apr;45(4):415-21, 421e1-2. doi: 10.1038/ng.2568.
Epub 2013 Feb 24. PMID: 23435085; PMCID: PMC3709584.

21. Baker ME, Katsu Y. 30 YEARS OF THE MINERALOCORTICOID RECEPTOR:
Evolution of the mineralocorticoid receptor: sequence, structure and function. J Endocrinol.
2017;234(1):T1-T16. doi:10.1530/JOE-16-0661.

22. York JR, McCauley DW. The origin and evolution of vertebrate neural crest cells. Open
Biol. 2020 Jan;10(1):190285. doi: 10.1098/rsob.190285. Epub 2020 Jan 29. PMID:
31992146; PMCID: PMC7014683.

23. Kuratani S. Evo-devo studies of cyclostomes and the origin and evolution of jawed
vertebrates. Curr Top Dev Biol. 2021;141:207-239. doi: 10.1016/bs.ctdb.2020.11.011.
Epub 2020 Dec 13. PMID: 33602489.

24. Janvier P. microRNAs revive old views about jawless vertebrate divergence and evolution.
Proc Natl Acad Sci U S A. 2010 Nov 9;107(45):19137-8. doi: 10.1073/pnas.1014583107.
Epub 2010 Nov 1. PMID: 21041649; PMCID: PMC2984170.

25. Bridgham JT, Carroll SM, Thornton JW. Evolution of hormone-receptor complexity by
molecular exploitation. Science. 2006;312(5770):97-101. doi:10.1126/science.1123348.

26. Katsu Y, Lin X, Ji R, Chen Z, Kamisaka Y, Bamba K, Baker ME. N-terminal domain
influences steroid activation of the Atlantic sea lamprey corticoid receptor. J Steroid
Biochem Mol Biol. 2023 Apr;228:106249. doi: 10.1016/j.jsbmb.2023.106249. Epub 2023
Jan 13. PMID: 36646152.

27. Katsu Y, Zhang J, Baker ME. Reduced steroid activation of elephant shark GR and MR after
inserting four amino acids from the DNA-binding domain of lamprey corticoid receptor-1.

PLoS One. 2023 Aug 23;18(8):e0290159. doi: 10.1371/journal.pone.0290159. PMID:
37611044; PMCID: PMC10446182.

28. Venkatesh B, Lee AP, Ravi V, Maurya AK, Lian MM, Swann JB, Ohta Y, Flajnik MF,
Sutoh Y, Kasahara M, Hoon S, Gangu V, Roy SW, Irimia M, Korzh V, Kondrychyn I, Lim
ZW, Tay BH, Tohari S, Kong KW, Ho S, Lorente-Galdos B, Quilez J, Marques-Bonet T,
Raney BJ, Ingham PW, Tay A, Hillier LW, Minx P, Boehm T, Wilson RK, Brenner S,
Warren WC. Elephant shark genome provides unique insights into gnathostome evolution.
Nature. 2014 Jan 9;505(7482):174-9. doi: 10.1038/nature12826. Erratum in: Nature. 2014
Sep 25;513(7519):574. Erratum in: Nature. 2020 Dec;588(7837):E15. PMID: 24402279;
PMCID: PMC3964593.

29. Katsu Y, Kohno S, Oka K, et al. Transcriptional activation of elephant shark
mineralocorticoid receptor by corticosteroids, progesterone, and spironolactone. Sci Signal.
2019;12(584):eaar2668. Published 2019 Jun 4. doi:10.1126/scisignal.aar2668, (n.d.).

30. Fonseca E, Machado AM, Vilas-Arrondo N, Gomes-Dos-Santos A, Veríssimo A, Esteves P,
Almeida T, Themudo G, Ruivo R, Pérez M, da Fonseca R, Santos MM, Froufe E, Román-
Marcote E, Venkatesh B, Castro LFC. Cartilaginous fishes offer unique insights into the
evolution of the nuclear receptor gene repertoire in gnathostomes. Gen Comp Endocrinol.
2020 Sep 1;295:113527. doi: 10.1016/j.ygcen.2020.113527. Epub 2020 Jun 8. PMID:
32526329.

31. Carroll SM, Bridgham JT, Thornton JW. Evolution of hormone signaling in elasmobranchs
by exploitation of promiscuous receptors. Mol Biol Evol. 2008;25(12):2643-2652.
doi:10.1093/molbev/msn204.

32. Arriza JL, Weinberger C, Cerelli G, et al. Cloning of human mineralocorticoid receptor
complementary DNA: structural and functional kinship with the glucocorticoid receptor.
Science. 1987;237(4812):268-275. doi:10.1126/science.3037703.

33. Fuller PJ. Novel interactions of the mineralocorticoid receptor. Mol Cell Endocrinol. 2015
Jun 15;408:33-7. doi: 10.1016/j.mce.2015.01.027. Epub 2015 Feb 7. PMID: 25662276.

34. Hellal-Levy C, Couette B, Fagart J, Souque A, Gomez-Sanchez C, Rafestin-Oblin M.
Specific hydroxylations determine selective corticosteroid recognition by human
glucocorticoid and mineralocorticoid receptors. FEBS Lett. 1999;464(1-2):9-13.
doi:10.1016/s0014-5793(99)01667-1.

35. Geller DS, Farhi A, Pinkerton N, et al. Activating mineralocorticoid receptor mutation in
hypertension exacerbated by pregnancy. Science. 2000;289(5476):119-123.
doi:10.1126/science.289.5476.119.

36. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J
Mol Biol. 1990 Oct 5;215(3):403-10. doi: 10.1016/S0022-2836(05)80360-2. PMID:
2231712.

37. Bloem LJ, Guo C, Pratt JH. Identification of a splice variant of the rat and human
mineralocorticoid receptor genes. J Steroid Biochem Mol Biol. 1995 Nov;55(2):159-62.
doi: 10.1016/0960-0760(95)00162-s. PMID: 7495694.

38. Wickert L, Watzka M, Bolkenius U, Bidlingmaier F, Ludwig M. Mineralocorticoid receptor
splice variants in different human tissues. Eur J Endocrinol. 1998 Jun;138(6):702-4. doi:
10.1530/eje.0.1380702. PMID: 9678540.

39. Wickert L, Selbig J, Watzka M, Stoffel-Wagner B, Schramm J, Bidlingmaier F, Ludwig M.
Differential mRNA expression of the two mineralocorticoid receptor splice variants within
the human brain: structure analysis of their different DNA binding domains. J
Neuroendocrinol. 2000 Sep;12(9):867-73. doi: 10.1046/j.1365-2826.2000.00535.x. PMID:
10971811.

40. de Kloet ER, Joëls M. Brain mineralocorticoid receptor function in control of salt balance
and stress-adaptation. Physiol Behav. 2017 Sep 1;178:13-20. doi:
10.1016/j.physbeh.2016.12.045. Epub 2017 Jan 13. PMID: 28089704.

41. Liu W, Wang J, Sauter NK, Pearce D. Steroid receptor heterodimerization demonstrated in
vitro and in vivo. Proc Natl Acad Sci U S A. 1995 Dec 19;92(26):12480-4. doi:
10.1073/pnas.92.26.12480. PMID: 8618925; PMCID: PMC40381.

42. Trapp T, Holsboer F. Heterodimerization between mineralocorticoid and glucocorticoid
receptors increases the functional diversity of corticosteroid action. Trends Pharmacol Sci.
1996 Apr;17(4):145-9. doi: 10.1016/0165-6147(96)81590-2. PMID: 8984741.

43. Mifsud KR, Reul JM. Acute stress enhances heterodimerization and binding of corticosteroid
receptors at glucocorticoid target genes in the hippocampus. Proc Natl Acad Sci U S A.
2016 Oct 4;113(40):11336-11341. doi: 10.1073/pnas.1605246113. Epub 2016 Sep 21.
PMID: 27655894; PMCID: PMC5056104.

44. Pooley JR, Rivers CA, Kilcooley MT, Paul SN, Cavga AD, Kershaw YM, Muratcioglu S,
Gursoy A, Keskin O, Lightman SL. Beyond the heterodimer model for mineralocorticoid
and glucocorticoid receptor interactions in nuclei and at DNA. PLoS One. 2020 Jan
10;15(1):e0227520. doi: 10.1371/journal.pone.0227520. PMID: 31923266; PMCID:
PMC6953809.

45. Kiilerich P, Triqueneaux G, Christensen NM, et al. Interaction between the trout
mineralocorticoid and glucocorticoid receptors in vitro. J Mol Endocrinol. 2015;55(1):55-
68. doi:10.1530/JME-15-0002.

46. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
Remmert M, Söding J, Thompson JD, Higgins DG. Fast, scalable generation of high-
quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol. 2011
Oct 11;7:539. doi: 10.1038/msb.2011.75. PMID: 21988835; PMCID: PMC3261699.