Model cyanobacterial consortia reveal a consistent core microbiome independent of inoculation source or cyanobacterial host species

Andreja Kust[1,2], Jackie Zorz[1,2], Catalina Cruañas Paniker[2,#], Keith Bouma-Gregson[3], Netravathi Krishnappa[2], Wendy Liu[1], Jillian F. Banfield[1,2,4,5], Spencer Diamond[1,2,*]

[1]Department of Earth and Planetary Science, University of California, Berkeley, Berkeley, CA, USA.
[2]Innovative Genomics Institute, University of California, Berkeley, CA, USA.
[3]U.S. Geological Survey, California Water Science Center, Sacramento, CA, USA.
[4]Department of Environmental Science, Policy and Management, University of California, Berkeley, CA, USA.
[5]School of Earth Sciences, University of Melbourne, Melbourne, Victoria, Australia.

[#]Current Address: Department of Life Sciences, Imperial College London, London, United Kingdom.

*Corresponding author:
Spencer Diamond
Innovative Genomics Institute
University of California - Berkeley
Innovative Genomics Institute Building
2151 Berkeley Way, Berkeley, CA 94720
Phone: (818) 268-5776
Email: sdiamond@berkeley.edu

Key Words: Cyanobacteria, Microbiology, Metagenomics, Microbial Ecology, Synthetic Communities.

Abstract

Cyanobacteria are integral to biogeochemical cycles, influence climate processes, and hold promise for commercial applications. In natural habitats, they form complex consortia with other microorganisms, where interspecies interactions shape their ecological roles. Although *in vitro* studies of these consortia have significantly advanced our understanding, they often lack the biological replication needed for robust statistical analysis of shared microbiome features and functions. Moreover, the microbiomes of many model cyanobacterial strains, which are central to our understanding of cyanobacterial biology, remain poorly characterized. Here, we expanded on existing *in vitro* approaches by co-culturing five well-characterized model cyanobacterial strains with microorganisms filtered from three distinct freshwater sources, generating 108 stable consortia. Metagenomic analyses revealed that, despite host and inoculum diversity, these consortia converged on a similar set of non-cyanobacterial taxa, forming a 25-species core microbiome. The large number of stable consortia in this study enabled statistical validation of both previously observed and newly identified core microbiome functionalities in micronutrient biosynthesis, metabolite transport, and anoxygenic photosynthesis. Furthermore, core species showed significant enrichment of plasmids, and functions encoded on plasmids suggested plasmid-mediated roles in symbiotic interactions. Overall, our findings uncover the potential microbiomes recruited by key model cyanobacteria, demonstrate that laboratory-enriched consortia retain many taxonomic and functional traits observed more broadly in phototroph-heterotroph assemblages, and show that model cyanobacteria can serve as robust hosts for uncovering functional roles underlying cyanobacterial community dynamics.

Introduction

In nature cyanobacteria form complex microbial communities [1] where inter-microbial interactions are critical for the resilience, stability, and environmental impact of cyanobacterial assemblages [2–7]. Cyanobacteria-associated microorganisms can act as fixed carbon sinks, modulate cyanobacterial productivity [8], and contribute to the biosynthesis of energetically costly compounds that benefit the entire microbial community [4]. These associated microbes may also play key roles in nutrient remineralization, enhancing the bioavailability of essential nutrients [6], and help mitigate oxidative stress [9]. Conversely, cyanobacteria-associated microorganisms may engage in competitive interactions, including competition with cyanobacteria for inorganic nutrients [9–12]. Cyanobacterial consortia share taxonomic and functional similarities with other phototroph-associated consortia [13] including those of the plant rhizosphere [14, 15], diatoms [16–18] and algae [19], suggesting that common ecological principles drive the assembly of these systems. However, a comprehensive understanding of the dynamics that drive cyanobacterial community assembly, and the roles of core microbial members within these communities, is still evolving [20–26].

Previous studies have provided important insights into cyanobacterial communities using cultivation-independent methods [6, 27–31], synthetic community assemblages [8, 32–35] and isolation and co-cultivation of cyanobacteria with their native microbial partners [21–23, 25, 26, 36–38]. However, *in vitro* consortia, which have made important contributions to our understanding of cyanobacterial communities [8, 21–25, 32–36], often contain few bacterial species and when assembled from isolates can lack taxonomic diversity that accurately reflects natural consortia. Alternatively, naturally sourced communities are not typically cultivated under standardized inoculation conditions or monitored longitudinally. Furthermore, the potential composition and function of microbiomes associated with cyanobacterial model organisms,

model species critical to our understanding of cyanobacterial biology, are largely
uncharacterized. To address these limitations, we built on previous methodologies to establish a
large number of *in vitro* communities grown under standardized laboratory conditions where
model cyanobacterial species were used as hosts to enrich co-associated microbiomes from
diverse natural inoculation sources. We chose well-characterized model cyanobacterial strains
as community hosts including *Synechocystis* spp. [39, 40], *Synechococcus elongatus* [41, 42],
and *Nostoc* spp. [43], where information about co-associated communities in natural
environments is limited [32, 33, 44]. This system builds upon previous approaches by balancing
the simplicity and genetic tractability of model cyanobacteria with the complexity of inoculating
these hosts with species derived from environmental samples.
109
Ultimately this work resulted in the establishment of 108 stable *in vitro* consortia. Using 16S
rRNA gene amplicon and genome-resolved metagenomic profiling, we observed that despite
the taxonomic composition of the freshwater source microbiomes and diversity of cyanobacterial
hosts, the resulting communities rapidly stabilized and became enriched with taxa commonly
found in natural freshwater cyanobacterial and other phototrophic symbiotic assemblages [6, 25,
45, 46]. We monitored the stability of these communities over time, including post-
cryopreservation revival, confirming their robustness, reproducibility, and ability to be transferred
or re-utilized for additional study. The large degree of replication in this study enabled the
identification of a 25 species core microbiome, which has not been previously evaluated using *in*
vitro efforts. Comparison of core microbiome metagenome assembled genomes (MAGs) to
other MAGs recovered in this study confirmed statistically significant enrichment of metabolic
traits previously noted observationally in other *in vitro* studies [12, 16, 25, 28, 47] and identified
additional functions in porphyrin metabolism and cyanobacterial alkane degradation associated
with the core microbiome. Core microbiome members were also significantly enriched in
putative plasmids, which encoded functionality potentially supportive of phototroph-heterotroph
interactions [48]. Overall, our results uncover the diversity of microbiomes recruited by model
cyanobacterial species, demonstrate that laboratory-enriched consortia using model strains
retain many of the taxonomic and functional features seen in broader phototroph–heterotroph
systems, and establish model cyanobacteria as effective platforms for uncovering new
functional roles driving cyanobacterial community dynamics.
130

Materials and Methods

132
Cyanobacterial selection, inoculation, growth, and passaging
We selected five model cyanobacterial strains—*Synechococcus elongatus* PCC 7942 (7942),
Synechococcus elongatus UTEX 3055 (3055), *Nostoc* spp., PCC 7120 (A7120), *Synechocystis*
spp., PCC 6803 (6803), and *Leptolyngbya* spp**.,** BL0902 (L0902)—based on their genetic
tractability, metabolic diversity, and relevance to cyanobacterial biology and biotechnology.
Axenic model cyanobacterial strains were inoculated with bacterial biomass collected from three
freshwater bodies (**Table S1**), and were grown in BG-11 [49] medium at 24°C under a 12:12
light cycle and passaged at both weekly and bi-weekly intervals for 12 weeks. Samples were
collected at each passage and rigorous standardized assessments of culture contamination,
and host dominance were applied to determine if a co-culture should be retained in the study
(**Table S2**). For comprehensive details, see **Supplementary Methods** and **Fig. 1A**.
144
DNA extraction and sequencing
DNA for shotgun metagenomic sequencing was extracted from inoculation source material and
biomass of co-culture samples at day 84 of passaging using a modified Qiagen DNeasy
PowerSoil Pro Kit protocol. DNA from co-culture samples was sequenced with paired-end reads

of 150 bp, whereas DNA from the source material was sequenced with paired-end reads of 250
bp on a NovaSeq System (Illumina). We used longer sequencing reads (250 bp) and a larger
library insert size for complex environmental source samples to improve assembly and contig
recovery [50].
153
For 16S rRNA gene amplicon sequencing, genomic DNA was extracted [51] from co-culture
samples of all passages. The extracted DNA was amplified for the V4–V5 regions of the 16S
rRNA gene using Illumina-compatible primers and sequenced on a MiSeq System (Illumina)
with a read depth of 50,000 reads per sample (mean = 46,098 reads per sample). This depth
enabled frequent monitoring of community changes and complemented shotgun metagenomic
sequencing, which was employed for species-level resolution and functional analysis of gene
content. For comprehensive details, see **Supplementary Methods.**
161
16S rRNA gene data processing and statistical analysis
Amplicon sequence variants (ASVs) were generated from fastq files using the USEARCH-
UNOISE3 pipeline [52], with modifications, to identify a total of 2,126 ASVs. Taxonomy was
assigned using a naive Bayes classifier trained on the SILVA v138 SSU [53] reference
database. We ultimately analyzed 869 rRNA gene amplicon samples (**Table S3**) that
represented samples collected from all longitudinal passages of co-culture samples determined
to be un-contaminanted, dominated by the expected host cyanobacterium, and also analyzed
with shotgun genome-resolved metagenomics at day 84. Alpha-diversity (Richness, Shannon)
was assessed with rarefied ASV counts (**Table S4**) using linear mixed-effects models. Beta-
diversity was analyzed with additive log ratio (ALR) transformed ASVs and Aitchison distances
(**Table S5**). Differential abundance was tested with Maaslin2 (**Tables S6**). For comprehensive
details, see **Supplementary Methods** and **Code Availability.**
174
Assembly independent marker gene analysis of shotgun metagenomic data
As recovering MAGs from complex source samples was challenging, we also employed an
analysis where marker genes were directly identified from shotgun metagenomic reads to more
accurately compare diversity and taxonomic composition between source samples and co-
cultures. This was conducted using ribosomal proteins as in prior work [54]. Using SingleM
v0.13.2 [55], unique Ribosomal protein L6 (rpL6) were identified and clustered at 95%
nucleotide identity to produce rpL6 OTUs that could be compared across samples (**Table S7**).
Over/under-enrichment of bacterial orders in cyanobacterial communities relative to source
samples was evaluated with a permutation-based method (**Table S8**). For comprehensive
details, see **Supplementary Methods** and **Code Availability**.
185
Metagenome assembly, binning, bin de-replication
Metagenome assembly and binning were conducted using methods similar to [50]. However,
scaffolds > 5000 bp were reassembled with COBRA v50 [56], and validated circular contigs
were re-incorporated into the assembly. Genome binning was performed on a sample by
sample basis and only used scaffolds > 2,500 bp using MetaBat2, maxbin2, CONCOCT, and
vamb [57–60]. Best MAGs across the four methods for each sample were selected using
DasTool [61] and evaluated with checkM [61, 62]. Only MAGs with ≥60% completeness and
contamination of ≤10% were retained. MAGs were de-replicated at the species level and
representative species MAGs were selected using dRep [63], considering MAGs as the same
species if ANI was ≥ 95% across ≥ 10% of the MAG length. This resulted in a non-redundant set
of 537 MAGs (**Table S9**). Taxonomic classification was conducted using GTDB-tk [64] classify
workflow against the GTDB-R214 taxonomy. For comprehensive details, see **Supplementary**
Methods.
199

Phylogenetic analysis

GToTree v1.6.34 [65] was employed to detect, align, trim, and concatenate 120 universal single
copy genes [66] from non-redundant MAGs as well as an archaeal outgroup species *Acidianus*
hospitalis W1 (GCF_000213215.1_2). MAGs with < 10% of markers were excluded. Maximum-
likelihood phylogenetic reconstruction was conducted with IQ-TREE version 1.6.12 [67]. The
best-fit evolutionary models for each marker protein partition were determined using
ModelFinder Plus. Tree topology significance was assessed using ultrafast bootstrap analysis
with 1,000 replicates and the Shimodaira-Hasegawa approximate likelihood ratio test (SH-
aLRT) with 1,000 replicates. The resulting phylogenetic tree was displayed using iTol v6.8.1
[68]. For comprehensive details, see **Supplementary Methods.**

MAG abundance mapping and diversity analysis

Shotgun metagenomic reads from each sample were mapped to species representative MAGs
with Bowtie2 [69] and coverage quantified with CoverM [70]. MAGs were considered present in
a sample if they met stringent coverage and breadth criteria (**Table S10**). Metagenomic samples
were removed from analysis when the expected host cyanobacterium comprised ≤ 90 % of the
cyanobacterial fraction resulting in 121 shotgun metagenomic samples (108 cyanobacterial
cultures and 13 source environment samples) for downstream analysis.
Beta-diversity was assessed using Aitchison distance of MAG mapped read counts, and
permutational analysis of variance (PERMANOVA) was used to estimate the influence of
sample type (culture or source) and location (**Table S11**). For comprehensive details, see
Supplementary Methods and **Code Availability.**

Core microbiome criteria

The concept of a core microbiome lacks a universally accepted definition [71, 72], so we
developed an operational framework to identify core microbiome species (MAGs) across co-
cultures based on two key criteria: (i) a core species must reproducibly co-occur (was
considered present in a co-culture by MAG mapping criteria) with at least four cyanobacterial
host species across three independent samples for each host (a minimum of 12 samples); and
(ii) it must be consistently recruited (was considered present in a co-culture by MAG mapping
criteria) from at least two distinct environmental sources across at least three independent
samples per environmental inoculum source (a minimum of 6 samples). This operational
definition ensures core species are consistently represented across both host diversity and
environmental variability, reducing the likelihood of sampling artifacts or host-specific
associations. To asses the relatedness of core microbiome MAGs assigned to the same
species, we performed digital DNA-DNA hybridization (dDDH) analysis using the Genome-to-
Genome Distance Calculator [73, 74](**Tables S12**).

Population level single nucleotide variant (SNV) analysis

Population-level SNV analysis was performed on 108 cyanobacterial co-culture samples using
inStrain v1.6.4 [75], generating 3,088 pairwise comparisons for 133 MAGs, including 555
comparisons involving core microbiome species. No identical strain sharing was detected
across 176 comparisons between geographically distinct inocula (popANI ≥ 99.999%; **Table**
S13). Beta regression modeling showed significant effects of inoculum source and host strain
on MAG microdiversity (adjusted P ≤ 0.05; **Table S14**). For comprehensive details, see
Supplementary Methods, Supplementary Results, and **Code Availability.**

Assessment of co-culture community taxonomic overlap with other studies

We compiled a set of 1,843 MAGs that passed completeness and contamination criteria used in
our study (920 distinct species) from 10 genome-resolved studies involving diverse
cyanobacterial or plant rhizosphere samples (**Tables S15 and S16**). These included studies of

non-axenic cyanobacteria cultures [23, 76–78], environmental cyanobacterial consortia [27, 28,
30, 79], and rhizosphere derived bacteria [80, 81]. Identical species (≥ 95% genome ANI)
between the 920 species-representative MAGs from other studies and the 319 MAGs observed
in co-cultures of this study were identified using dRep [63]. Additionally we evaluated higher
level taxonomic overlap between species-representative MAGs present in our co-cultures and
those of other studies at the class and genus level using taxonomic classifications produced
using GTDB-tk [64] classify workflow against the GTDB-R214 taxonomy.
258
Functional annotation and core microbiome functional enrichment analysis
Predicted proteins from all 537 MAGs were annotated using KOfam HMMs via kofamscan [82],
and METABOLIC [83]. Alkane degradation genes were identified using hmmsearch [84] with
HMMs from CANT-HYD [85]. To assess core microbiome functional enrichment (n = 25 MAGs)
compared to non-core MAGs (n = 512 MAGs), Fisher tests were conducted on KEGG orthologs
(KOs) with p-values adjusted for multiple testing (FDR ≤ 0.05). Two tests were performed: one
using all MAGs (n = 537 MAGs) and another constrained to Pseudomonadota, Bacteroidota,
and Spirochaetota MAGs (n = 391 MAGs) to account for taxonomic bias in the core microbiome
MAG set. Broader functional categories were assigned, and differential representation was
analyzed using Fisher tests with multiple testing correction. For comprehensive details, see
Tables S17-S20, **Supplementary Methods,** and **Code Availability**.
270
Identification, curation, and analysis of putative mobile genetic elements
Contigs predicted to be mobile genetic elements (MGEs) were identified across 537 MAGs
using geNomad [86] filtering for contigs ≥12 kb and FDR ≤ 0.05 (**Table S21**). We evaluated all
contigs ≥12 kb that were confidently identified as putative plasmids, given the high frequency of
circular element fragmentation in short-read metagenomic assemblies. Plasmid assignment to
MAGs was based on a guilt-by-association approach, where plasmids co-binned with a MAG
were considered associated. Plasmid content was compared between core, auxiliary, and
source genomes using Dunn's test. All plasmid contigs were aligned to the IMG/PR database
[87] using BLASTn for identification (**Table S22**). Genes related to conjugation, mobilization,
and replication were identified using HMMs from Conjscan [88] and Pfam [89] via hmmsearch
[84]. Plasmids were classified as mobilizable or conjugative based on the presence of key
genes (e.g., MOBX, T4CP, virB4). General functional annotation was performed identically as
for MAGs detailed above.
284
For MGE-associated functional enrichment, plasmid-contigs were removed from the 537 MAGs,
and KEGG orthology (KO) enrichment analysis was repeated. Differential KO enrichment was
compared between the full and MGE-removed analyses using Fisher's test, with FDR correction
(FDR ≤ 0.05) (**Tables S18 and S19**). Fold changes in KO enrichment were calculated between
the original and MGE-removed analyses. For comprehensive details, see **Supplementary**
Methods, and **Code Availability**.
291

Results

293
Inoculation of model cyanobacterial strains with bacteria from diverse freshwater
sources established stable *in vitro* communities
A total of 108 consortia were retained at the end of 12 weeks of passaging being free of
contamination and dominated by the original cyanobacterial host **(Fig. S1 and Supplementary**
Methods). The retention of consortia in the experiment was influenced by the cyanobacterial
host, inoculum source, and the interaction between host strain and passage rate. Among the

tested strains, 6803, L0902, and A7120 exhibited the highest retention rates (**Supplementary**
Results).
302
Profiling of 16S rRNA gene amplicons showed that despite differences in inoculum source and
host strain, communities rapidly stabilized within the first 28 days (**Figs. 1B-D and S2**).
Cryopreservation and revival confirmed the long-term stability of the communities, with diversity
metrics and taxonomic composition returning to pre-cryopreservation values **(Fig. 1B-D)**. The
final consortia set evaluated for this study (84 days post inoculation) closely resembled natural
cyanobacterial communities, displaying a consistent taxonomic structure (**Fig. 1, S3, and**
Supplementary Results). These results confirm that model cyanobacterial species can form
stable, reproducible consortia *in vitro*, simulating natural microbiomes [6, 9, 19, 90].
311
Genomic resolution enabled comprehensive characterization of *in vitro* communities
Shotgun sequencing of stable consortia (84 days) and source samples yielded 537 non-
redundant species-level MAGs, with 59.8% classified as high-quality (completeness of ≥90%,
contamination of ≤5%) and 40.2% as medium-quality (completeness of ≥60%, contamination of
≤10% (**Table S9**). These MAGs, spanning 17 phyla, primarily Bacteroidota and
Pseudomonadota (**Fig. 2A**), included 84.2% of MAGs without species-level representatives in
existing genomic databases, and59 MAGs unclassified beyond the family level (**Table S9**). The
recovered MAGs effectively represented the *in vitro* communities, with 86.2% of co-culture
reads mapping back to MAGs compared to only 9.3% of source sample reads mapping back to
MAGs on average **(Fig. 2A, Table S2, Supplementary Results**).
322
Specific taxonomic groups become enriched in stable *in vitro* communities
We directly compared diversity between environmental source samples and co-cultures. This
analysis was conducted both by comparing MAG diversity (**Fig. S5**) and performing an
assembly-independent comparison of ribosomal protein L6 marker OTU diversity (L6OTU; **Fig.**
2B and S4), to overcome low MAG recovery from source samples. Both analyses revealed
significantly lower alpha-diversity for *in vitro* consortia relative to source samples (**Fig. S4 and**
S5). Using L6OTU analysis we found that 22 orders were overrepresented, whereas 59 were
underrepresented within *in vitro* consortia relative to source environments. Bacterial orders from
the Pseudomonadota and Bacteroidota were the most frequently enriched in co-cultures, and
bacterial orders within the Candidate Phyla Radiation (CPR) and Omnitrophota were
underrepresented (**Fig. 2B, Table S8**). Despite differences in inoculum sources, the *in vitro*
environment consistently favored the enrichment of specific taxa (**Supplementary Results**).
335
Inoculum source and cyanobacterial host strain drive inter-community variation
Stable co-cultures were predominantly composed of their original cyanobacterial host strain,
accompanied by prevalent Pseudomonadota and Bacteroidota species, along with less common
phyla such as Spirochaetota, Actinomycetota, and Armatimonadota (**Fig. 3A-B and Table S10**).
Between co-culture samples, we found that the source of inoculum (R^2 = 0.23, P < 0.0001;
PERMANOVA), the cyanobacterial host strain (R^2 = 0.11, P < 0.0001; PERMANOVA), and the
interaction between the inoculum source and host strain (R^2 = 0.08, P < 0.0001; PERMANOVA),
all contributed to statistically significant variations in community composition, with inoculum
source having the strongest influence (**Fig. S5G-I**). Despite these variations, communities
converged toward a broadly similar taxonomic structure with enrichment of major taxonomic
groups relative to source samples (**Fig. 2B and 3A-B**). Comparison of source samples and co-
cultures revealed distinct clustering, with *in vitro* community samples tightly clustered relative to
samples of source environments (**Fig. S4G, S5G-I, and Table S11**). These observations
suggest that even though differences between *in vitro* communities exist, they are substantially
smaller than the variations introduced during community formation from source inocula. Thus,

even though inoculum source and host strain shape community diversity, the resulting consortia
maintain high-level taxonomic coherence across different host strains and inocula.
353
Core species identified across model cyanobacterial communities
Although there is no universally accepted definition of a core microbiome, the consistent
taxonomic composition and developmental patterns in our *in vitro* cyanobacterial communities
(**Fig. 1B, 3A-B, and Fig. S3**) allowed us to define a 25 species core microbiome using genome-
resolved data (**See Methods**). These species were reproducibly detected across multiple
communities with different cyanobacterial hosts (>= 4 host species) and environmental inocula
(>= 2 source environments; **Fig. S6**). Hereinafter we refer to non-core species detected in
cultures as auxiliary species and species only detected in environmental samples as source
species (**Fig. 2A and Table S9**). Core microbiome species averaged 1.0 ± 1.3% (Range: 0 -
7.3%) of the total community abundance, increasing to 14.7 ± 19.4% (Range: 0 - 82.4%) when
considering only non-cyanobacterial taxa **(Fig. 3C)**. Individual core species were present in
18.5% - 68.5% of all samples (**Table S9 and Fig. S6**). The majority of core species belonged to
taxonomic orders significantly enriched in the L6OTU analysis (**Fig. 2B**) with the most numerous
being Rhizobiales (n = 7 species), Rhodobacterales (n = 5 species) and Sphingomonadales (n
= 5 species) (**Fig. 3D**). Phylogenetically, core species were closely clustered, with significantly
lower phylogenetic distances between their MAGs on average relative to auxiliary or source
inoculum species (**Fig. 2A and S7**). Single nucleotide variant (SNV) analysis further confirmed
that identical species were independently selected into *in vitro* communities from distinct
geographic water sources, with no evidence of strain-sharing between environments (**Fig. S8,**
Table S13-S14, Supplementary Methods and Results). Eighteen core species lacked a
representative in public databases (**Table S9**), marking an expansion of genomic knowledge for
organisms that may occupy a generalized niche in cyanobacterial communities.
376
Taxonomic similarity between *vitro* and natural phototrophic consortia
We identified 29 (9.1%) shared species, **(Fig. 3E and S9**) between our *in vitro* communities and
MAGs compiled from other studies on cyanobacterial consortia (n = 1,843 MAGs). Among
these, three MAGs are core microbiome members identified in our study. In total, 22 (76%)
shared species also belonged to taxonomic orders enriched during *in vitro* community
establishment in this study (**Fig. 2B, 3E**). On average, this study shared more species with other
studies (3.2 ± 6.3 species) than the studies did with each other (0.7 ± 2.9 species). Most
pairwise comparisons between studies identified no shared species (42/55 comparisons). The
species shared with this study were primarily from freshwater-associated cyanobacterial
communities, with no species overlap in saltwater, soda lake, or plant rhizosphere communities.
387
At higher taxonomic levels we observed broad overlap between studies. Out of the 48 observed
orders in this study, 37 (77%) were shared with other studies (**Fig. S9B**), with the orders
enriched during *in vitro* community establishment in this study being the most commonly shared
(P = 0.018; Wilcoxon test, **Fig. S9C**). These findings support that the *in vitro* communities
developed here reasonably represent natural cyanobacterial consortia and share considerable
taxonomic overlap with other phototroph-heterotroph systems.
394
Enriched metabolic functions in the core cyanobacterial microbiome
We identified metabolic functions enriched in core microbiome MAGs (n = 25 MAGs) relative to
all non-core MAGs (n = 512 MAGs), and a constrained set of non-core MAGs from three phyla
in the core microbiome Bacteroidota, Pseudomonadota, and Spirocheaotota (n = 366 MAGs). In
both analyses, we found significant enrichment of 1073 and 1065 KEGG orthology groups
(KOs), respectively (FDR ≤ 0.05; Fisher test; **Tables S17-S19**). Enriched functional categories
included cofactor metabolism (Fisher test; OR_{All} = 1.6, FDR_{All} = 0.007; $OR_{Constrained}$ = 1.9,

FDR$_{Constrained}$ = 0.0002) and membrane transport (Fisher test; OR$_{All}$ = 1.7, FDR$_{All}$ = 1.9 e-6;
OR$_{Constrained}$ = 1.7, FDR$_{Constrained}$ = 1.1e^{-5}). (**Fig. 4A, Table S19**). The constrained analysis
mirrored the full analysis and additionally identified signal transduction as (Fisher test;
OR$_{Constrained}$ = 1.7, FDR$_{Constrained}$ = 0.006) a significantly overrepresented functional category in
the core microbiome (**Fig. 4A**).

Within cofactor metabolism, 39 KOs in the porphyrin metabolic pathway were significantly
enriched in core microbiome MAGs including those involved in the production of vitamin B12,
chlorophylls, and hemes. The gene encoding 5-aminolevulinate synthase (K00643) which
facilitates one-step synthesis of 5-Aminolevulinate (ALA) from glycine was highly enriched in
core MAGs (OR$_{All}$ = 9.9, FDR$_{All}$ = 6.5 e-6; Fisher test), whereas non-core MAGs favored the
more complex three-step pathway from glutamate (K02492; Fisher test; OR$_{All}$ = 24.6, FDR$_{All}$ =
3.3 e-8). The gene *cobC* (K02225), involved in vitamin B12 synthesis, showed the strongest
core microbiome association (Fisher test; OR$_{All}$ = 24.6, FDR$_{All}$ = 3.3 e-8), alongside other
enzymes in the vitamin B12 and bacteriochlorophyll synthesis pathways (**Fig. 4B and Table**
S18).

We also found enrichment of KOs related to anoxygenic photosystems, with 80% (20/25) of
core microbiome members encoding these components (K08929/K08928/K13991; **Fig. 5, Table**
S18 and S20). This may indicate that light driven energy production is an important metabolic
strategy of cyanobacteria-associated bacteria. Electron donors for these anoxygenic
phototrophs are likely organic substrates [91], which agrees with functional predictions. Only
one member of the core microbiome, a *Georhizobium* species, encoded RuBisCO for carbon
fixation.

Genes for phosphonate degradation (K06162-6) and transport (K02041-4) were significantly
enriched in core MAGs, offering an alternate source of phosphorus to the community. KOs
involved in methylamine (K15228-9), nitrile (K01721/K20807), and urea (K01428-30) utilization
were also enriched in core MAGs, putatively expanding available carbon and nitrogen sources
(**Fig. 5 and Table S18 and S20**). Additionally, many core microbiome MAGs that encoded
methylamine to formaldehyde degradation also had genes to oxidize formaldehyde to formate
and CO_2. Finally, core microbiome members were enriched in several genes related to
degradation of cyanobacteria-derived long-chain alkanes. All cyanobacterial strains in our study
encoded KOs (K14330 and K14331) responsible for long-chain alkane biosynthesis [92, 93],
wheres core microbiome members were enriched in genes (*LadA*, *AhyA*, *AlkB*, and *Cyp153*) for
the degradation of these compounds [85] (**Fig. 5 and Table S18 and S20**). This suggests that
core microbiome species may be utilizing these cyanobacteria-produced alkanes as an energy
source.

Core microbiome MAGs are enriched in putative plasmid elements
Extrachromosomal replicons and plasmids have been shown to encode functions that mediate
phototroph-heterotroph symbioses [48]. Here we assessed if there were differences in the
frequency and distribution of putative plasmid elements associated with the MAGs resolved in
our study. We identified 850 putative plasmid contigs across 537 species-representative MAGs
(**Table S21**). Even though 98% of putative plasmids were not circularized, we found 15%
encoded plasmid-specific replication domains and 26% encoded proteins for plasmid
mobilization (**Fig. S10A**). In our comparison with the comprehensive IMG/PR database, 81
plasmids from our dataset matched entries in the database. Of these 81 matching plasmids,
80% were recovered from metagenomic datasets, primarily from aquatic environments, with
only 7 being assessed as putatively complete (**Table S22**).

452
We found that core microbiome MAGs had a significantly higher proportion of putative plasmid
contigs than auxiliary (Dunn Test = 3.7, *P* = 0.0003) or source MAGs (Dunn Test = 6.5, *P* = 0)
(**Fig 6A**). As bacteria from the order Rhizobiales are known to encode large numbers of
extrachromosomal plasmids [94, 95] we also performed the comparison only between MAGs of
this order. We found that core Rhizobiales MAGs contained a significantly larger proportion of
putative plasmids than auxiliary (Dunn Test = 3.0, *P* = 0.0036) or source (Dunn Test = 2.4, *P* =
0.0253) MAGs **(Fig. S10D)**.
460
Putative plasmids of the core microbiome encode functions facilitating symbioses
To assess functional contribution of putative plasmids in core microbiome MAGs, we repeated
our functional enrichment analysis, excluding plasmid-predicted contigs from all MAGs. We
identified 197 KEGG Orthology (KO) groups that were no longer significantly enriched in the
core microbiome (**Tables S18 and S19**), indicating that these KOs were enriched on putative
plasmids. Functional category analysis of this KO set revealed significant enrichment of signal
transduction (Fisher test; OR = 0.7, FDR = 0.038), membrane transport (Fisher test; OR = 0.6,
FDR = 0.038), and carbohydrate metabolism (Fisher test; OR = 0.5, FDR = 0.049) on putative
plasmids of the core microbiome (**Fig. 6B**).
470
Enriched plasmid-specific functions on core microbiome plasmids (**Fig. 6C and Table S18**)
included those for plasmid replication (*repC*) and conjugal transfer (*TrbBCDFGI*). Carbohydrate
metabolic functions included those mediating the interconversion and metabolism of sugars
commonly found in cyanobacterial exopolysaccharides [96, 97], such as xylose (*XylABD*),
rhamnose (*RhaAM*), and fucose (K18335, K07046, and *FucA*). Transporters for importing
xylose (*XylG*) and simple sugars (*AbcSS* and *AbcMS*) were also enriched. Additionally, putative
plasmids in the core microbiome may enable the utilization of peptides as alternative carbon
and nitrogen sources as evidenced by the enrichment of dipeptide (*DppBCF*) and oligopeptide
(*OppBC*) transport systems, and genes (*Mut*, *PccA*, and *BkdAB*) involved in converting 2-
oxobutanoate to succinyl-CoA, linking amino acid degradation to the TCA cycle.
481
We identified plasmid-enriched genes that may mediate symbiotic interactions and biofilm
formation. These genes include multiple flagellar structural proteins (FlgBDK and FlhAR), a
flagellar regulatory protein (RpoD), and components of an Imp-like type VI secretion system
(ImpBCFM) (**Fig. 6C and Table S18**). Furthermore, we identified a set of plasmid-enriched KOs
involved in exopolysaccharide biosynthesis (*exoM* and *GumL*). We discovered comprehensive
gene clusters for the biosynthesis of succinoglycan and alginate on a large (407 kb) putative
plasmid (**Fig. 6D and Table S23**) associated with a Rhizobiales MAG from the core microbiome
(StrawCreek_S_L0902_W_A_idba_concoct_4). Although this element was not fully circularized,
it encoded plasmid-specific replication factors (*repABC*) and a complete conjugal transfer
system, including a relaxase protein (TraI) and type IV secretion apparatus (*trb*).
492
493 # Discussion
494
In this work, we built on existing approaches [21–23, 25, 26, 36–38] to generate and analyze
108 *in vitro* cyanobacterial consortia using well-characterized, model cyanobacterial hosts. This
approach provides three core advances for the *in vitro* study of cyanobacterial communities.
First, leveraging cyanobacterial host species that serve as key model systems in cyanobacterial
research [43] significantly enhances our previously poor understanding of the microbiomes they
recruit. Second, extensive replication enabled the identification of a core microbiome across
multiple taxonomically diverse cyanobacterial hosts. Third, leveraging this core microbiome, we

confirmed, statistically rather than solely by observation, enrichment of functional traits
associated with core species, and, furthermore, uncovered metabolic functions and the potential
for plasmids to mediate phototroph-heterotroph symbioses in these systems.
505
The cyanobacterial hosts used in this study were selected for their diversity and importance as
model organisms for advancing our understanding of cyanobacterial biology [41, 42, 98–103].
Despite their value as models, little is known about their behavior in ecological contexts.
Whereas previous studies have utilized some of these species in two-member co-culture
systems [8, 32, 33, 35], this study provides the insights into more complex microbial
communities potentially recruited by these species from natural environments. We provide
strong evidence that not only can *in vitro* communities rapidly stabilize around these model
hosts (**Fig. 1B-D and S3**), but that these communities exhibit strong taxonomic overlaps with
freshwater cyanobacteria in natural habitats [28, 30], cultivated in lower-throughput laboratory
systems [23, 78], and share higher level taxonomic overlap with plant rhizosphere associated
communities (**Fig. 3E, S9**). Furthermore, evidence supporting host-specific selective pressures
was observed, as species co-occurring with the same host exhibited significantly higher strain-
level ANI across co-cultures compared to the same species associated with different hosts **(Fig.
S9)**.
520
By inducing co-culture diversity through variation of both cyanobacterial host species and
environmental inoculum sources, consistently shared features could be robustly identified. This
diversity was leveraged across consortia to define a 25-species core microbiome (**Fig. 3D**)
identified across five model cyanobacterial host strains. Despite the limitation of using
freshwater cyanobacterial species exclusively, the core microbiome identified here provides a
foundation for understanding organisms that commonly co-occur across an array of
taxonomically and functionally diverse cyanobacteria. Although some core species identified
here were also present in cyanobacterial consortia from other studies **(Fig. 3E)**, our hypothesis
is that the specific identities of these core microbiome members are less critical than the
functional traits and "microbial archetypes" they represent. Therefore, comparing the genomic
content of core and non-core species provides a framework for statistically identifying functional
traits that play a broader role in supporting cyanobacterial ecosystems.
533
Functional genomic analysis of the core microbiome revealed a significant enrichment of
metabolic pathways associated with the biosynthesis and potential cross-feeding of resource-
intensive compounds (e.g., vitamin B12), the enhancement of nitrogen and phosphorus
availability, and anoxygenic phototrophy (**Figs. 4 and 5**). Even though some of these pathways,
such as anoxygenic photosynthesis [28, 30], have been previously reported [12, 16] this study
provides robust statistical evidence for their specific enrichment within a core microbiome.
Collectively, these findings support the hypothesis that core microbiome functions facilitate
metabolic cooperation centered around phototrophy. This is further reinforced by the
predominance of porphyrin cofactor metabolism among enriched functions, including numerous
enzymes involved in vitamin B12 and bacteriochlorophyll biosynthesis pathways **(Fig. 4)**. The
core microbiome exhibits enrichment in ALA synthase, which enables a more energetically
efficient route for ALA production from glycine, potentially advantageous in communities where
porphyrin compounds are critical resources. The importance of such exchanges is underscored
by studies demonstrating enhanced cyanobacterial growth when supplemented with the
porphyrin precursor ALA [104]. Although vitamin B12 cross-feeding in microbial communities
has been studied intensively [105–107], the shared production and exchange of other
energetically demanding and metabolically linked molecules, such as chlorophyll precursors,
remain poorly characterized. The enrichment of anoxygenic phototrophy within the core
microbiome, consistent with observations from other studies [91], may suggest that non-

competitive photosynthetic [108] partnerships are important in cyanobacterial microbiomes for
exchanging essential and costly metabolic precursors common to all organisms. Finally,
although multiple studies have posited the function of cyanobacterial long-chain alkanes [109],
this study shows the enrichment of long-chain alkane degradation pathways in core microbiome
organisms (**Fig. 5**). This capability was specifically prevalent in those core microbiome
organisms that also encoded phototropic capabilities. This observation would support that
cyanobacteria produce these compounds to specifically select and modulate their surrounding
communities by creating a metabolic niche.
561
Even though plasmids have been studied in microbial symbioses [110, 111], their roles within
cyanobacterial consortia remain largely unexplored. Here, core microbiome genomes exhibited
a higher prevalence of plasmids compared to non-core members (**Fig. 6**). Functionally core
microbiome plasmids encoded gene clusters linked to the biosynthesis of exopolysaccharides
such as succinoglycan, alginate, and xanthan gum-like compounds, polymers known to play
essential roles in plant-bacterial interactions [110, 112]. Succinoglycan is critical for nitrogen-
fixing Rhizobiales [113, 114], and alginate forms hydrogels that enhance biofilm formation [115–
117]. In particular, alginate encapsulation of *Synechococcus elongatus* has been shown to
improve both sucrose export and co-culture stability in laboratory settings [32]. Additionally, core
microbiome plasmids contained a two-component nitrate regulatory system (*NasST*), gene
clusters for mobility machinery, and Type VI secretion systems, features commonly found in
plant-associated bacteria [118], suggesting potential roles in colonization, competition, and
biofilm formation. Beyond colonization and biofilm forming functionality, plasmids in core
microbiome species encoded metabolic pathways for pentose and hexose sugar utilization,
including rhamnose and xylose metabolism, key components of cyanobacterial
exopolysaccharides [96, 97] (**Fig. 6, S10, Tables S17-18, S22**). Collectively, these findings
support the hypothesis that plasmids contribute to core microbiome establishment and
maintenance by enhancing biofilm formation and niche adaptation.The data further highlight
adaptive advantages plasmids may confer within cyanobacterial consortia, providing new
insights into how plasmids may shape these microbial interactions.
582
Even though compelling, several limitations of our approach warrant discussion. The absence of
cyanobacteria-free controls, limited our ability to disentangle effects solely attributable to media
or laboratory conditions. However, we observed substantial taxonomic overlap (**Fig. 3E and S9**)
between our co-cultures and other freshwater cyanobacterial and rhizosphere consortia [6, 28,
45, 46, 103]. Furthermore, the use of naturally sourced inocula carried the potential for
introducing diverse eukaryotic organisms, including flagellates, ciliates, fungi, and phototrophic
diatoms and algae. Although we rigorously screened cultures for readily detectable eukaryotes
(**See Methods**), we did not assess the diversity or function of eukaryotic species present within
final co-cultures, which on average constituted 0.64±1.89 % of assembled data on an
assembled contig size basis (**Table S2**). Our objective was to focus on the prokaryotic
interactions of model cyanobacteria, given the limited existing data on these hosts' recruited
microbiomes. Despite these constraints, our findings and related work [19] underscore the value
of *in vitro* cyanobacterial model systems for revealing insights into phototroph–heterotroph
interactions. Although methods for cultivating phytoplankton consortia are well-established [19,
23, 28, 38, 47], our framework and biobanked stable co-cultures offer a controlled yet
ecologically relevant platform to advance understanding of freshwater microbial community
dynamics. Future investigations using complementary methods such as long-read sequencing,
mesocosm experiments, cyanobacteria-free controls, and inclusion of eukaryotic organisms, will
help refine our understanding of aquatic microbiome functionality.
602

Data Availability

Raw sequencing read data generated as part of this project are available under the NCBI BioProject accession number PRJNA1271998. All metagenome bins and putative plasmid sequences recovered in this study have been deposited in FigShare under the following DOIs: 10.6084/m9.figshare.26148823 and 10.6084/m9.figshare.26148811

Code Availability

Code used in the analysis for this paper are available at the following GitHub repository: https://github.com/SDmetagenomics/Kust_Cyano_2023

Reporting Summary

Further information on research design is available in the Nature Research Reporting Summary linked to this article.

Acknowledgements

We would like to thank Prof. Susan Golden, Prof. James Golden, and Dr. Arnaud Taton for providing the cyanobacterial strains used in this study. Eel River sampling occurred at the UC Angelo Coast Range Reserve. Funding for this project was provided in part by the Gordon and Betty Moore Foundation (GBMF9321). Funding for this project was also provided in part by the Shurl and Kay Curci Foundation.

Author Contributions

AK, JFB, and SD conceived overall experimental concepts and design. AK and SD designed the sampling and cultivation strategy. AK, KBG, and SD collected water samples. AK, CCP, WL, and SD performed cultivation and sampling. AK, CCP, NK, and SD performed sequencing and sequence processing. AK and SD performed sequence assembly and genomic binning. AK, JZ, and SD performed annotation and statistical analysis on amplicon and metagenomic datasets. AK, JZ, and SD wrote the manuscript. AK, JZ, JFB, and SD edited the manuscript

Competing Interests

The authors declare no competing interests.

Corresponding Author

All correspondence should be addressed to Spencer Diamond (sdiamond@berkeley.edu)

References

1. Paerl HW, Pinckney JL. A mini-review of microbial consortia: Their roles in aquatic production and biogeochemical cycling. *Microb Ecol* 1996; **31**: 225–247.

2. Conover AE, Morando M, Zhao Y, Semones J, Hutchins DA, Webb EA. *Alphaproteobacteria* facilitate *Trichodesmium* community trimethylamine utilization. *Environ Microbiol* 2021; **23**: 6798–6810.

3. Zhang Z, Fan X, Peijnenburg WJGM, Zhang M, Sun L, Zhai Y, et al. Alteration of dominant cyanobacteria in different bloom periods caused by abiotic factors and species interactions. *J Environ Sci* 2021; **99**: 1–9.

4. Smith DJ, Kharbush JJ, Kersten RD, Dick GJ. Uptake of phytoplankton-derived carbon and cobalamins by novel Acidobacteria genera in *Microcystis* blooms inferred from metagenomic and metatranscriptomic evidence. *Appl Environ Microbiol* 2022; **88**: e01803-21.

5. Cai H, McLimans CJ, Jiang H, Chen F, Krumholz LR, Hambright KD. Aerobic anoxygenic phototrophs play important roles in nutrient cycling within cyanobacterial *Microcystis* bloom microbiomes. *Microbiome* 2024; **12**: 1–19.

6. Cook KV, Li C, Cai H, Krumholz LR, Hambright KD, Paerl HW, et al. The global interactome. *Limnol Oceanogr* 2020; **65**: S194–S207.

7. Li W, Baliu-Rodriguez D, Premathilaka SH, Thenuwara SI, Kimbrel JA, Samo TJ, et al. Microbiome processing of organic nitrogen input supports growth and cyanotoxin production of *Microcystis aeruginosa* cultures. *ISME J* 2024; **18**: wrae082.

8. Hays SG, Yan LLW, Silver PA, Ducat DC. Synthetic photosynthetic consortia define interactions leading to robustness and photoproduction. *J Biol Eng* 2017; **11**: 4.

9. Kim M, Shin B, Lee J, Park HY, Park W. Culture-independent and culture-dependent analyses of the bacterial community in the phycosphere of cyanobloom-forming *Microcystis aeruginosa*. *Sci Rep* 2019; **9**: 20416.

10. Dick GJ, Duhaime MB, Evans JT, Errera RM, Godwin CM, Kharbush JJ, et al. The genetic and ecophysiological diversity of *Microcystis*. *Environ Microbiol* 12 2021; **23**: 7278–7313.

11. Jackrel SL, Schmidt KC, Cardinale BJ, Denef VJ. Microbiomes reduce their host's sensitivity to interspecific interactions. *mBio* 2020; 11.

12. Baker D, Godwin CM, Khanam M, Burtner AM, Dick GJ, Denef VJ. Variation in resource competition traits among strains is affected by their microbiomes. *mLife* 2023; **2**: 401–415.

13. Durán P, Flores-Uribe J, Wippel K, Zhang P, Guan R, Melkonian B, et al. Shared features and reciprocal complementation of the *Chlamydomonas* and *Arabidopsis* microbiota. *Nat Commun* 2022; **13**: 1–14.

14. Ling N, Wang T, Kuzyakov Y. Rhizosphere bacteriome structure and functions. *Nat Commun* 2022; **13**: 1–13.

15. Morella NM, Weng FC-H, Joubert PM, Metcalf CJE, Lindow S, Koskella B. Successive passaging of a plant-associated microbiome reveals robust habitat and host genotype-dependent selection. *Proc Natl Acad Sci USA* 2020; **117**: 1148–1159.

16. Amin SA, Hmelo LR, van Tol HM, Durham BP, Carlson LT, Heal KR, et al. Interaction and signalling between a cosmopolitan phytoplankton and associated bacteria. *Nature* 2015; **522**: 98–101.

17. Kojadinovic-Sirinelli M, Villain A, Puppo C, Sing SF, Prioretti L, Hubert P, et al. Exploring the microbiome of the 'star' freshwater diatom *Asterionella formosa* in a laboratory context. *Environ Microbiol* 2018; **20**: 3601–3615.

18. Sison-Mangus MP, Jiang S, Tran KN, Kudela RM. Host-specific adaptation governs the interaction of the marine diatom, *Pseudo-nitzschia* and their microbiota. *ISME J* 2014; **8**: 63–76.

19. Jackrel SL, Yang JW, Schmidt KC, Denef VJ. Host specificity of microbiome assembly and its fitness effects in phytoplankton. *ISME J* 2020; **15**: 774–788.

20. Nelson C, Giraldo-Silva A, Garcia-Pichel F. A symbiotic nutrient exchange within the cyanosphere microbiome of the biocrust cyanobacterium, *Microcoleus vaginatus. ISME J* 2021; **15**: 282–292.

21. Zhu L, Zancarini A, Louati I, De Cesare S, Duval C, Tambosco K, et al. Bacterial communities associated with four cyanobacterial genera display structural and functional differences: evidence from an experimental approach. *Front Microbiol* 2016; **7**: 1662.

22. Cole JK, Hutchison JR, Renslow RS, Kim Y-M, Chrisler WB, Engelmann HE, et al. Phototrophic biofilm assembly in microbial-mat-derived unicyanobacterial consortia: model systems for the study of autotroph-heterotroph interactions. *Front Microbiol* 2014; **5**: 109.

23. Zhao L, Lin L-Z, Zeng Y, Teng W-K, Chen M-Y, Brand JJ, et al. The facilitating role of phycospheric heterotrophic bacteria in cyanobacterial phosphonate availability and *Microcystis* bloom maintenance. *Microbiome* 2023; **11**: 142.

24. Halary S, Duperron S, Demay J, Duval C, Hamlaoui S, Piquet B, et al. Metagenome-based exploration of bacterial communities associated with cyanobacteria strains isolated from thermal muds. *Microorganisms* 2022; 10.

25. Duxbury SJN, Raguideau S, Rosko J, Cremin K, Coates M, Quince C, et al. Reproducible spatial structure formation and stable community composition in the cyanosphere predicts metabolic interactions. *bioRxiv* 2022.

26. Haines M, Vadlamani A, Daniel Loty Richardson W, Strous M. Pilot-scale outdoor trial of a cyanobacterial consortium at pH 11 in a photobioreactor at high latitude. *Bioresour Technol* 2022; **354**: 127173.

27. Bouma-Gregson K, Olm MR, Probst AJ, Anantharaman K, Power ME, Banfield JF. Impacts of microbial assemblage and environmental conditions on the distribution of anatoxin-a producing cyanobacteria within a river network. *ISME J* 2019; **13**: 1618–1634.

28. Pérez-Carrascal OM, Tromas N, Terrat Y, Moreno E, Giani A, Marques LCB, et al. Single-colony sequencing reveals microbe-by-microbiome phylosymbiosis between the cyanobacterium *Microcystis* and its associated bacteria. *Microbiome* 2021,**12**: 9.

29. Zhu C-M, Zhang J-Y, Guan R, Hale L, Chen N, Li M, et al. Alternate succession of aggregate-forming cyanobacterial genera correlated with their attached bacteria by co-pathways. *Sci Total Environ* 2019; **688**: 867–879.

30. Li H, Bhattarai B, Barber M, Goel R. Stringent response of cyanobacteria and other bacterioplankton during different stages of a harmful cyanobacterial bloom. *Environ Sci Technol* 2023; **57**: 16016–16032.

31. Smith DJ, Berry MA, Cory RM, Johengen TH, Kling GW, Davis TW, et al. Heterotrophic bacteria dominate catalase expression during *Microcystis* blooms. *Appl Environ Microbiol* 2022; **88**: e0254421.

32. Taylor L. Weiss, Eric J. Young, Daniel C. Ducat. A synthetic, light-driven consortium of cyanobacteria and heterotrophic bacteria enables stable polyhydroxybutyrate production. *Metabolic Engineering* 2017; **44**: 236–245.

33. Zuñiga C, Li T, Guarnieri MT, Jenkins JP, Li C-T, Bingol K, et al. Synthetic microbial communities of heterotrophs and phototrophs facilitate sustainable growth. *Nat Commun* 2020; **11**: 1–13.

34. Weissberg O, Aharonovich D, Sher D. Phototroph-heterotroph interactions during growth and long-term starvation across *Prochlorococcus* and *Alteromonas* diversity. *ISME J* 2022; 17: **227**–237.

35. Christie-Oleza JA, Sousoni D, Lloyd M, Armengaud J, Scanlan DJ. Nutrient recycling facilitates long-term stability of marine microbial phototroph–heterotroph interactions. *Nat Microbiol* 2017; **2**: 1–10.

36. Wu Q, Zhang Y, Li Y, Li J, Zhang X, Li P. Comparison of community composition between *Microcystis* colony-attached and free-living bacteria, and among bacteria attached with *Microcystis* colonies of various sizes in culture. *Aquat Ecol* 2019; **53**: 465–481.

37. Vardaka E, Kormas KA, Katsiapi M, Genitsaris S, Moustaka-Gouni M. Molecular diversity of bacteria in commercially available '*Spirulina*' food supplements. *PeerJ* 2016; 4.

38. Yancey CE, Kiledal EA, Chaganti SR, Denef VJ, Errera RM, Evans JT, et al. The Western Lake Erie culture collection: A promising resource for evaluating the physiological and genetic diversity of *Microcystis* and its associated microbiome. *Harmful Algae* 2023; **126**: 102440.

39. Ikeuchi M, Tabata S. *Synechocystis* sp. PCC 6803 - a useful tool in the study of the genetics of cyanobacteria. *Photosynth Res* 2001; **70**: 73–83.

40. Mehdizadeh Allaf M, Peerhossaini H. Cyanobacteria: model microorganisms and beyond. *Microorganisms* 2022; 10.

41. Rubin BE, Wetmore KM, Price MN, Diamond S, Shultzaberger RK, Lowe LC, et al. The essential gene set of a photosynthetic organism. *Proc Natl Acad Sci USA* 2015; **112**: E6634–43.

800 42. Diamond S, Jun D, Rubin BE, Golden SS. The circadian oscillator in *Synechococcus*
801   *elongatus* controls metabolite partitioning during diurnal growth. *Proc Natl Acad Sci USA*
802   2015; **112**: E1916–25.
803
804 43. Zeng X, Zhang C-C. The making of a heterocyst in cyanobacteria. *Annu Rev Microbiol* 2022;
805   **76**: 597–618.
806
807 44. Cahill B, Straka L, Maldonado Ortiz J, Krajmalnik-Brown R, Rittmann BE. Effects of light
808   intensity on soluble microbial products produced by *Synechocystis* sp. PCC 6803 and
809   associated heterotrophic communities. *Algal Res* 2019; **38**: 101409.
810
811 45. Seymour JR, Amin SA, Raina J-B, Stocker R. Zooming in on the phycosphere: the
812   ecological interface for phytoplankton–bacteria relationships. *Nat Microbiol* 2017; **2**: 1–12.
813
814 46. Steinrücken P, Jackson S, Müller O, Puntervoll P, Kleinegris DMM. A closer look into the
815   microbiome of microalgal cultures. *Front Microbiol* 2023; **14**: 1108018.
816
817 47. Jackrel SL, White JD, Evans JT, Buffin K, Hayden K, Sarnelle O, et al. Genome evolution
818   and host-microbiome shifts correspond with intraspecific niche divergence within harmful
819   algal bloom-forming *Microcystis aeruginosa*. *Mol Ecol* 2019; **28**: 3994–4011.
820
821 48. Wardell GE, Hynes MF, Young PJ, Harrison E. Why are rhizobial symbiosis genes mobile?
822   *Philos Trans R Soc Lond B Biol Sci* 2022; **377**: 20200471.
823
824 49. Guillard RRL, Lorenzen CJ. Yellow-green algae with chlorophyllide c. *J Phycol* 1972; **8**: 10–
825   14.
826
827 50. Diamond S, Andeer PF, Li Z, Crits-Christoph A, Burstein D, Anantharaman K, et al.
828   Mediterranean grassland soil C–N compound turnover is dependent on rainfall and depth,
829   and is mediated by genomically divergent microorganisms. *Nat Microbiol* 2019; **4**: 1356–
830   1367.
831
832 51. Clerico EM, Ditty JL, Golden SS. Specialized techniques for site-directed mutagenesis in
833   cyanobacteria. *Methods Mol Biol* 2007; **362**: 155–171.
834
835 52. Edgar RC. UNOISE2: improved error-correction for Illumina 16S and ITS amplicon
836   sequencing. *bioRxiv* 2016; 081257.
837
838 53. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal
839   RNA gene database project: improved data processing and web-based tools. *Nucleic Acids*
840   *Res* 2013; **41**: D590–6.
841
842 54. Olm MR, Crits-Christoph A, Diamond S, Lavy A, Matheus Carnevali PB, Banfield JF.
843   Consistent metagenome-derived metrics verify and delineate bacterial species boundaries.
844   *mSystems* 2020; 5.
845
846 55. Woodcroft BJ, Aroney STN, Zhao R, Cunningham M, Mitchell JAM, Blackall L, et al. SingleM
847   and Sandpiper: Robust microbial taxonomic profiles from metagenomic data. *bioRxiv* 2024;
848   2024.01.30.578060
849

850 56. Chen L, Banfield JF. COBRA improves the completeness and contiguity of viral genomes
851   assembled from metagenomes. *Nat Microbiol* 2024; **9**: 737–750.
852
853 57. Wu YW, Simmons BA, Singer SW. MaxBin 2.0: an automated binning algorithm to recover
854   genomes from multiple metagenomic datasets. *Bioinformatics* 2016; **32**: 605–607.
855
856 58. Alneberg J, Bjarnason BS, Bruijn ID, Schirmer M, Quick J, Ijaz UZ, et al. Binning
857   metagenomic contigs by coverage and composition. *Nat Methods* 2014; **11**: 1144–1146.
858
859 59. Kang DD, Froula J, Egan R, Wang Z. MetaBAT, an efficient tool for accurately reconstructing
860   single genomes from complex microbial communities. *PeerJ* 2015; **3**: e1165.
861
862 60. Nissen JN, Johansen J, Allesøe RL, Sønderby CK, Armenteros JJA, Grønbech CH, et al.
863   Improved metagenome binning and assembly using deep variational autoencoders. *Nat*
864   *Biotechnol* 2021; **39**: 555–560.
865
866 61. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, et al. Recovery of
867   genomes from metagenomes via a dereplication, aggregation and scoring strategy. *Nat*
868   *Microbiol* 2018; 3: 836–843.
869
870 62. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the
871   quality of microbial genomes recovered from isolates, single cells, and metagenomes.
872   *Genome Res* 2015; **25**: 1043–1055.
873
874 63. Olm MR, Brown CT, Brooks B, Banfield JF. dRep: a tool for fast and accurate genomic
875   comparisons that enables improved genome recovery from metagenomes through de-
876   replication. *ISME J 7* 2017; **11**: 2864–2868.
877
878 64. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify genomes
879   with the Genome Taxonomy Database. *Bioinformatics* 2020; **36**: 1925–1927.
880
881 65. Lee MD. GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics* 2019; **35**:
882   4162–4164.
883
884 66. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A, Hugenholtz P. GTDB: an
885   ongoing census of bacterial and archaeal diversity through a phylogenetically consistent,
886   rank normalized and complete genome-based taxonomy. *Nucleic Acids Res* 2021; **50**:
887   D785–D794.
888
889 67. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective
890   stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 2015; **32**:
891   268–274.
892
893 68. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
894   display and annotation. *Nucleic Acids Res* 2021; **49**: W293–W296.
895
896 69. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 2012;
897   **9**: 357–359.
898

899 70. Aroney STN, Newell RJP, Nissen JN, Camargo AP, Tyson GW, Woodcroft BJ. CoverM: read
900   alignment statistics for metagenomics. *Bioinformatics* 2025; **41**:4.
901

902 71. Neu AT, Allen EE, Roy K. Defining and quantifying the core microbiome: Challenges and
903   prospects. *Proc Natl Acad Sci USA* 2021; 118.
904

905 72. Dong C, Shao Q, Zhang Q, Yao T, Huang J, Liang Z, et al. Preferences for core microbiome
906   composition and function by different definition methods: Evidence for the core microbiome
907   of *Eucommia ulmoides* bark. *Sci Total Environ* 2021; **790**: 148091.
908

909 73. Auch AF, von Jan M, Klenk H-P, Göker M. Digital DNA-DNA hybridization for microbial
910   species delineation by means of genome-to-genome sequence comparison. *Stand Genomic*
911   *Sci* 2010; **2**: 117–134.
912

913 74. Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M. Genome sequence-based species
914   delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics*
915   2013; **14**: 60.
916

917 75. Olm MR, Crits-Christoph A, Bouma-Gregson K, Firek BA, Morowitz MJ, Banfield JF. inStrain
918   profiles population microdiversity from metagenomic data and sensitively detects shared
919   microbial strains. *Nat Biotechnol* 2021; 39: 727–736.
920

921 76. Berthold M, Albrecht M, Campbell DA, Omar NM. Draft genomes of 3 cyanobacteria strains
922   and 17 co-habiting proteobacteria assembled from metagenomes. *Microbiol Resour*
923   *Announc* 2023; **12**: e0046023.
924

925 77. Cornet L, Bertrand AR, Hanikenne M, Javaux EJ, Wilmotte A, Baurain D. Metagenomic
926   assembly of new (sub)polar Cyanobacteria and their associated microbiome from non-
927   axenic cultures. *Microb Genom* 2018; 4.
928

929 78. Li Q, Lin F, Yang C, Wang J, Lin Y, Shen M, et al. A large-scale comparative metagenomic
930   study reveals the functional interactions in six bloom-forming -epibiont communities. *Front*
931   *Microbiol* 2018; **9**: 746.
932

933 79. Zorz JK, Sharp C, Kleiner M, Gordon PMK, Pon RT, Dong X, et al. A shared core
934   microbiome in soda lakes separated by large distances. *Nat Commun* 2019; **10**: 4230.
935

936 80. Levy A, Salas Gonzalez I, Mittelviefhaus M, Clingenpeel S, Herrera Paredes S, Miao J, et al.
937   Genomic features of bacterial adaptation to plants. *Nat Genet* 2017; **50**: 138–150.
938

939 81. Bai L, Cui J, Jie W, Cai B. Analysis of the community compositions of rhizosphere fungi in
940   soybeans continuous cropping fields. *Microbiol Res* 2015; **180**: 49–56.
941

942 82. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, et al. KofamKOALA:
943   KEGG Ortholog assignment based on profile HMM and adaptive score threshold.
944   *Bioinformatics* 4 2020; **36**: 2251–2252.
945

946 83. Zhou Z, Tran PQ, Breister AM, Liu Y, Kieft K, Cowley ES, et al. METABOLIC: high-
947   throughput profiling of microbial genomes for functional traits, metabolism, biogeochemistry,
948   and community-scale functional networks. *Microbiome* 2022; **10**: 33.
949

84. Eddy SR. Accelerated profile HMM searches. *PLoS Comput Biol* 2011; **7**: e1002195.

85. Khot V, Zorz J, Gittins DA, Chakraborty A, Bell E, Bautista MA, et al. CANT-HYD: A curated database of phylogeny-derived Hidden Markov Models for annotation of marker genes involved in hydrocarbon degradation. *Front Microbiol* 2021; **12**: 764058.

86. Camargo AP, Roux S, Schulz F, Babinski M, Xu Y, Hu B, et al. Identification of mobile genetic elements with geNomad. *Nat Biotechnol* 2024; **42**: 1303–1312.

87. Camargo AP, Call L, Roux S, Nayfach S, Huntemann M, Palaniappan K, et al. IMG/PR: a database of plasmids from genomes and metagenomes with rich annotations and metadata. *Nucleic Acids Res* 2024; **52**: D164–D173.

88. Cury J, Abby SS, Doppelt-Azeroual O, Néron B, Rocha EPC. Identifying conjugative plasmids and integrative conjugative elements with CONJscan. *Methods Mol Biol* 2020; **2075**: 265–283.

89. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, et al. Pfam: The protein families database in 2021. *Nucleic Acids Res* 2021; **49**: D412–D419.

90. Smith DJ, Tan JY, Powers MA, Lin XN, Davis TW, Dick GJ. Individual *Microcystis* colonies harbour distinct bacterial communities that differ by *Microcystis* oligotype and with time. *Environ Microbiol* 2021; **23**: 3020–3036.

91. Yurkov VV, Beatty JT. Aerobic anoxygenic phototrophic bacteria. *Microbiol Mol Biol Rev* 1998; **62**: 695–724.

92. Schirmer A, Rude MA, Li X, Popova E, del Cardayre SB. Microbial biosynthesis of alkanes. *Science* 2010; **329**: 559–562.

93. Lea-Smith DJ, Biller SJ, Davey MP, Cotton CAR, Perez Sepulveda BM, Turchyn AV, et al. Contribution of cyanobacterial alkane production to the ocean hydrocarbon cycle. *Proc Natl Acad Sci USA* 2015; **112**: 13591–13596.

94. Leinberger J, Holste J, Bunk B, Freese HM, Spröer C, Dlugosch L, et al. High potential for secondary metabolite production of CP157, isolated from the crustacean. *Front Microbiol* 2021; **12**: 688754.

95. Czarnecki J, Chapkauskaitse E, Bos J, Sentkowska D, Wawrzyniak P, Wyszyńska A, et al. Differential localization and functional specialization of centromere-like sites in replicons of Alphaproteobacteria. *Appl Environ Microbiol* 2022; **88:** e0020722.

96. De Philippis R, Vincenzini M. Exocellular polysaccharides from cyanobacteria and their possible applications. *FEMS Microbiol Rev* 1998; **22**: 151–175.

97. Mota R, Flores C, Tamagnini P. Cyanobacterial Extracellular Polymeric Substances (EPS). In: Oliveira JM, Radhouani H, Reis RL (eds). *Polysaccharides of Microbial Origin*. 2021. Springer, Cham, pp 1–28.

999   98. Yang Y, Lam V, Adomako M, Simkovsky R, Jakob A, Rockwell NC, et al. Phototaxis in a wild
1000   isolate of the cyanobacterium. *Proc Natl Acad Sci USA* 2018; **115**: E12378–E12387.
1001

1002   99. Schatz D, Nagar E, Sendersky E, Parnasa R, Zilberman S, Carmeli S, et al. Self-
1003   suppression of biofilm formation in the cyanobacterium *Synechococcus elongatus*. *Environ*
1004   *Microbiol* 2013; **15**: 1786–1794.
1005

1006   100. Cohen SE, Golden SS. Circadian rhythms in cyanobacteria. *Microbiol Mol Biol Rev* 2015;
1007   **79**: 373–385.
1008

1009   101. Kaneko T, Sato S, Kotani H, Tanaka A, Asamizu E, Nakamura Y, et al. Sequence analysis of
1010   the genome of the unicellular cyanobacterium *Synechocystis* sp. strain PCC6803. II.
1011   Sequence determination of the entire genome and assignment of potential protein-coding
1012   regions (supplement*). DNA Res* 1996; **3**: 185–209.
1013

1014   102. Lázaro S, Fernández-Piñas F, Fernández-Valiente E, Blanco-Rivero A, Leganés F. pbpB, a
1015   gene coding for a putative penicillin-binding protein, is required for aerobic nitrogen fixation
1016   in the cyanobacterium *Anabaena* sp. strain PCC7120. *J Bacteriol* 2001; **183**: 628–636.
1017

1018   103. Taton A, Lis E, Adin DM, Dong G, Cookson S, Kay SA, et al. Gene transfer in *Leptolyngbya*
1019   sp. strain BL0902, a cyanobacterium suitable for production of biomass and bioproducts*.*
1020   *PLoS One* 2012; **7**: e30901.
1021

1022   104. Promotive effect of 5-aminolevulinic acid on the growth and photosynthesis of *Spirulina*
1023   *platensis*. Journal of Fermentation and Bioengineering 1995; **79**: 453–457.
1024

1025   105. Grant MAA, Kazamia E, Cicuta P, Smith AG. Direct exchange of vitamin B12 is
1026   demonstrated by modelling the growth dynamics of algal-bacterial cocultures. *ISME J* 2014;
1027   **8**: 1418–1427.
1028

1029   106. Sultana S, Bruns S, Wilkes H, Simon M, Wienhausen G. Vitamin B is not shared by all
1030   marine prototrophic bacteria with their environment. *ISME J* 2023; **17**: 836–845.
1031

1032   107. Hessler T, Huddy RJ, Sachdeva R, Lei S, Harrison STL, Diamond S, et al. Vitamin
1033   interdependencies predicted by metagenomics-informed network analyses and validated in
1034   microbial community microcosms. *Nat Commun* 2023; **14**: 4768.
1035

1036   108. Koblížek M. Ecology of aerobic anoxygenic phototrophs in aquatic environments. *FEMS*
1037   *Microbiol Rev* 2015; **39**: 854–870.
1038

1039   109. Parveen H, Yazdani SS. Insights into cyanobacterial alkane biosynthesis. *J Ind Microbiol*
1040   *Biotechnol* 2022; 49.
1041

1042   110. Finks SS, Martiny JBH. Plasmid-encoded traits vary across environments*. mBio* 2023; **14**:
1043   e0319122.
1044

1045   111. Finks SS, Moudgalya P, Weihe C, Martiny JBH. The contribution of plasmids to trait diversity
1046   in a soil bacterium. *ISME Commun* 2024; **4**: ycae025.
1047

1048   112. Acosta-Jurado S, Fuentes-Romero F, Ruiz-Sainz J-E, Janczarek M, Vinardell J-M. Rhizobial
1049   exopolysaccharides: genetic regulation of their synthesis and relevance in symbiosis with

1050   legumes. *Int J Mol Sci* 2021; 22.

1051

1052   113.Skorupska A, Janczarek M, Marczak M, Mazur A, Król J. Rhizobial exopolysaccharides:
1053   genetic control and symbiotic functions. *Microb Cell Fact* 2006; **5**: 1–19.

1054

1055   114.Reuber TL, Walker GC. Biosynthesis of succinoglycan, a symbiotically important
1056   exopolysaccharide of *Rhizobium meliloti*. *Cell* 1993; **74**: 269–280.

1057

1058   115.Allison DG, Sutherland IW. The role of exopolysaccharides in adhesion of freshwater
1059   bacteria. *Microbiology* 1987; **133**: 1319–1327.

1060

1061   116.Urtuvia V, Maturana N, Acevedo F, Peña C, Díaz-Barrera A. Bacterial alginate production: an
1062   overview of its biosynthesis and potential industrial production. *World J Microbiol Biotechnol*
1063   2017; **33**: 198.

1064

1065   117.Boyd A, Chakrabarty AM. *Pseudomonas aeruginosa* biofilms: role of the alginate
1066   exopolysaccharide. *J Ind Microbiol* 1995; **15**: 162–168.

1067

1068   118.Bernal P, Llamas MA, Filloux A. Type VI secretion systems in plant-associated bacteria.
1069   *Environ Microbiol* 2018; **20**: 1–15.

1070

1071 **Figure Legends**

1072

Figure 1 | Assessment of *in vitro* community stability over the 119 d passaging
experiment. (A) Experimental design for cyanobacterial *in vitro* community cultivation. Water
samples were collected from three freshwater locations (two sites per location) resulting in six
source inocula. Bacterial fractions were prefiltered, then concentrated using 0.2 μm and 0.1 μm
filters, combined, and used to inoculate five cyanobacterial host strains in triplicate. Each host
was paired with six source inocula twice for passaging at both 7- and 14-day intervals over 12
weeks, resulting in 180 initial cultures. At each passage, samples were collected for microscopic
analysis, OD measurements, and 16S rRNA gene amplicon sequencing. After the 84 d
passage, samples were collected for metagenomic sequencing and cryopreservation. The
frozen consortia were then regrown, passaged, and harvested as before. **(B)** Order-level
community composition for *in vitro* communities at 14 d intervals over 119 d, based on 16S
rRNA gene amplicon sequencing averaged across all samples. Orders found in the core
microbiome are indicated with an asterisk (*). In all plots the vertical red line separates pre-
cryopreservation passaging (left) and post-cryopreservation passaging (right). **(C)** Change in
Aitchinson distance between previous passage time points for *in vitro* communities at 14 d
intervals over 119 d. Gray dots give change for each community compared to itself in the
preceding time point, purple squares indicate estimated marginal means, and error bars
represent 95% CI of marginal mean. Statistical significance between comparisons estimated
using linear mixed effects models with comparisons not sharing the same letter being
significantly different (FDR ≤ 0.05). **(D)** Observed richness of *in vitro* communities at 14 d
intervals over 119 d (total ASV counts). Gray dots give values for each community at a given
time point, blue squares indicate estimated marginal means, and error bars represent 95% CI of
marginal mean. Statistical significance between time points estimated using linear mixed effects
models with time points not sharing the same letter being significantly different (FDR ≤ 0.05).

1097
Figure 2 | Maximum likelihood phylogenetic tree and enrichment of microbial orders. (A)
A phylogenetic tree of 537 MAGs in our study constructed using a concatenated alignment of
120 bacterial-specific marker genes. An archaeal reference genome (GCF_000213215.1)
served as an outgroup for rooting the tree. Nodes marked with black dots represent a high
bootstrap support (≥ 95%, calculated using ufboot with n = 1000 iterations). The outer ring
highlights core microbiome MAGs (in blue), and the inner ring differentiates MAGs observed in
in vitro cultures (orange) and those exclusively observed in source samples (purple). Branch
tips with cyanobacterial host species are denoted by yellow stars and have no outer ring colors.
Scale bar indicates the number of nucleotide substitutions per site. **(B)** Microbial orders showing
significant over or under enrichment when comparing ribosomal protein L6 marker gene OTU
(rpL6 OTU) diversity between true *in vitro* community compositions and randomly sampled *in*
vitro communities (n = 10,000 permutations) where all observed rpL6 OTUs were sampled. The
Z-score indicates the level of enrichment of an order in observed communities relative to
communities generated from random sampling. Only orders with statistically significant
enrichment values are shown (FDR ≤ 0.05; permutation test). Orders positioned on the left of
the red line are underrepresented in true consortia, and those on the right of the line are over-
represented in true consortia. Blue squares identify taxa found in the core microbiome. Bars are
colored by phylum as indicated in the key of Fig. 2A with additional colors added and noted in
the key of Fig. 2B for minor phyla (Other) and Omnitrophota. Also see Tables S8 and S9.
1117
Figure 3 | Microbial community composition, core microbiome, and cross community
overlap. (A) Phylum level relative abundance of *in vitro* communities based on mapping
samples to a database of 537 species-level MAGs. Sample compositions are averaged within
cyanobacterial strain for each environmental inoculum. **(B)** Phylum level taxonomic composition
of *in vitro* communities with cyanobacteria removed. **(C)** Composition of *in vitro* communities
with cyanobacteria removed depicting the relative fraction assigned to core and auxiliary
microbiome MAGs. **(D)** Cyanobacterial host association and order-level taxonomy of the 25
genomes identified as core microbiome members. The numbers above the colored stacked bars
indicate the total number of core species shared. The colored stacked bars represent the order-
level taxonomy of these species. Connected dots below each stacked bar indicate the set of
cyanobacterial hosts that share the species above. The numbers in parentheses on the right of
the plot indicate the total number of species detected in at least three cultures for each host
strain. The legend on the right of the figure provides the color code for each order in the stacked
bar plot. An asterisk (*) indicates orders that were significantly enriched in co-cultured
communities relative to source environment samples (Also See Fig. 2B). **(E)** The number and
taxonomy of species shared between the communities of our study and those of other studies
where genome-resolved analysis of cyanobacterial consortia was conducted. The numbers
above the colored stacked bars indicate the total number species shared. The colored stacked
bars indicate the order-level taxonomy of these species. Blue asterisks above the stacked bar
plots indicate that a core microbiome species identified in our study was shared. Connected
dots below each stacked bar indicate the set of studies that share the species above. Numbers
in parentheses on the right of the plot indicate the total number of species recovered in each
study. The green and blue color shading under the connected dot plots indicates the type and

origin of cyanobacterial community for the compared studies. The coloring of orders in the
stacked bar plot of this figure are the same as in Figure 3D.
1143
Figure 4 | Metabolic functions enriched in the core microbiome. (A) Functional categories
of statistically over or under enriched KEGG functions in the core microbiome. The enrichment
odds-ratio for each functional category is shown for the set of KEGG functions with differential
enrichment when the core microbiome MAGs were compared to all 512 non-core MAGs (red)
and the 366 non-core MAGs in the phyla *Bacteroidota*, *Proteobacteria*, and *Spircheaotota*
(blue). Bars are solid colors for categories with statistically significant over or under
representation (FDR ≤ 0.05; Fisher test) and transparent if not significant. **(B)** Metabolic reaction
map for porphyrin compound biosynthetic pathways. Numbers indicate the KEGG KO
associated with a reaction. This map shows enrichment data from the comparison of core MAGs
to all 512 non-core MAGs. The size of colored circles indicates the percentage of MAGs in core
(green; n = 25) or non-core (blue; n = 512) sets that encode the respective reaction. Circles are
solid colors if a KEGG KO was detected as significantly over or under enriched in the core
microbiome (FDR ≤ 0.05; Fisher test), and transparent if not significant. Also see Tables S16-
S18.
1158
Figure 5 | Comparison of metabolic functions in cyanobacterial hosts and core
microbiome species. Encoded metabolic potential for specific biogeochemical turnover
reactions identified in cyanobacterial host species MAGs and the 25 core microbiome MAGs.
Specific metabolic functions (left) are separated by higher level metabolic categories (right).
Functions marked with and asterisk (*) had at least one associated KO with significant
enrichment in the core microbiome relative to all 512 non-core MAGs. Each column represents
an individual MAG in the core microbiome, MAGs are grouped at different phylum levels, and
colored by order level taxonomy. Identification of the potential for a specific function in each
MAG is indicated by a colored box, and boxes share order and color. Criteria for identifying
specific functions can be found in Table S19.
1169
Figure 6 | Predicted mobile genetic element distribution and functional potential. (A)
Proportion of contigs predicted to be mobile genetic elements in each MAG of the core,
auxiliary, and source microbiome. Points are colored by phylum level taxonomy of the MAG.
Box plots behind points indicate median, first, and third quartiles of data. Statistical differences
between groups are indicated by letters, and groups not sharing the same letter are significantly
different (Statistic$_{Core-v-Aux}$ = 3.7, FDR < 0.001; Statistic$_{Core-v-Source}$ = 6.5, FDR < 0.0001;
Statistic$_{Aux-v-Source}$ = 6.6, FDR < 0.0001; Dunn Test). **(B)** Functional category analysis for the set
of KEGG KOs that becomes non-significant when all MGE contigs are removed from MAGs.
Functional categories that are significantly over or underrepresented in this set of KOs are
colored accordingly (FDR ≤ 0.05; Fisher test). X-axis indicates the natural logarithm of the
Fisher test enrichment score. Also see Tables S15, S16 and S20. **(C)** Individual KEGG KOs
from membrane transport, carbohydrate metabolism, and signal transduction categories that
become non-significant when the enrichment analysis for core microbiome functions is repeated
with predicted MGE contigs removed from all MAGs. The x-axis indicates the -log10(FDR) for
each KO in the original enrichment analysis. The y-axis indicates the fold change in the odds-

ratio for a KO between the original analysis and the analysis conducted with all MGE contigs
removed. **(D)** A representative plasmid contig (407,520 bp) co-binned with an *Allorhizobium*
MAG from the core microbiome (StrawCreek_S_L0902_W_A_idba_concoct_4). Genes are
colored by functional category. Dashed lines indicate a break in the contig for visualization
purposes. Also see Table S23.

A Sampling locations

Serial filtration

Biomass for DNA + cultivation

5 Axenic cyanobacterial strains

n=1, 2, 3,...180

6 bacterial inocula

7 and 14 d ppassaging

Eel River
Discovery Bay
Strawberry Creek

0 10 20 mi

Amplicon sequencing

Cryopreservation

Re-growth for 4 weeks

Cryopreservation

Shotgun + amplicon sequencing

Amplicon sequencing

Passaging 84 d

B

Order
- Burkholderiales
- Chitinophagales*
- Cyanobacteria
- Cytophagales*
- Fimbriimonadales
- Flavobacteriales
- Rhizobiales*
- Rhodobacterales*
- Sphingobacteriales
- Sphingomonadales*
- Xanthomonadales
- Other

C

D

A

Core Genomes
Source_Cultures

Legend:
- Core genomes (blue)
- Cultures (orange)
- Source only (purple)
- Host cyanobacteria (yellow star)

Phyla:
- CPR
- CSP1-3
- Chloroflexota
- Actinomycetota
- Bacillota
- Armatimonadota
- Cyanobacteria
- Planctomycetota
- Chlamydiota
- Verrucomicrobiota
- Gemmatimonadota
- Bacteroidota
- Acidobacteriota
- Spirochaetota
- Myxococcota
- Bdellovibrionota
- Gammaproteobacteria
- Alphaproteobacteria

B

Z Score

Legend:
- Core microbiome (blue)
- Omnitrophota (pink)
- Other (grey)

Figure 3

Order

- Acetobacterales*
- Chitinophagales*
- Cytophagales*
- GCA-2729495*
- Leptospirales*
- NS11-12g*
- Pseudomonadales*
- Rhizobiales*
- Rhodobacterales*
- Sphingomonadales*
- Actinomycetales*
- Burkholderiales
- CACIAM−22H2
- Caulobacterales*
- Ferrovibrionales
- Fimbriimonadales*
- Flavobacterales*
- Gemmatimonadales
- M30B66
- UBA7662

Phylum

- Cyanobacteria
- Armatimonadota
- Spirochaetota
- Bacteroidota
- Pseudomonadota
- Chlamydiota
- Acidobacteriota
- Actinobacteriota
- Firmicutes
- Verrucomicrobiota
- Gemmatimonadota
- Planctomycetota

Microbial Genome Group

- Auxilary
- Core

Figure 5