1   **An isogenic *E. coli* population gives rise to multiple persister phenotypes**

2   **Authors:** K M Taufiqur Rahman[1], Ruqayyah Amaratunga[1], Abhyudai Singh[3], Tahmina Hossain[1*] and
3   Nicholas C. Butzin[1,2*]
4
Affiliations:
[1]Department of Biology and Microbiology; South Dakota State University; Brookings, South Dakota,
USA
[2]Department of Chemistry and Biochemistry, South Dakota State University, Brookings, South Dakota,
USA
[3]Electrical & Computer Engineering, University of Delaware, Newark, Delaware, USA
11
12   *Corresponding author. Email: *nicholas.butzin@gmail.com; tahminatan.40@gmail.com*
13

Abstract

Bacterial persisters are a multidrug-tolerant subpopulation capable of surviving and resuscitating after
lethal antibiotic treatment, leading to relapsing infections and the emergence of antibiotic resistance. We
hypothesize that multiple persister subpopulations are within an isogenic *Escherichia coli* population,
allowing them to survive lethal antibiotic stress. We conducted transcriptomic analyses at multiple time
points with lethal ampicillin (Amp) antibiotic, and as expected, several genes were differentially
expressed over time. We identified a subset of genes consistently upregulated by comparing transcription
levels at different time points of Amp-treated to untreated. Some genes had previously been associated
with persisters, while others were new. Subsequently, network analysis showed the gene response
between networks, but could not map hypothetical genes. So, we overexpressed seven hypothetical genes,
which resulted in slow growth or no growth, indicating that high production harmed the cell. We then
made single gene knockouts, which dramatically reduced persister level by ~4-6 fold at 3 h and ~10-15
fold at 6 h of Amp treatment. However, no significant difference in survival rates was observed at 24 h,
indicating the presence of multiple persister subpopulations. Our mathematical model demonstrated a 20-
fold decrease in the slow-decaying fraction in the mutant, suggesting the importance of decay kinetics in
bacterial survival. These results support the existence of multiple subpopulations of persisters, each
characterized by distinct decay rates. These results challenge the idea of complete dormancy, suggest the
presence of intricate, multifaceted survival mechanisms, and indicate that the persister population itself is
heterogeneous.
33

34   **Keywords**
35   Antibiotic tolerance, antibiotic persistence, antibiotic resistance, phenotypic heterogeneity, cell-state
36   transitions, phenotypic switching
37

Significance statement
Bacterial persisters, a subpopulation known for their multidrug tolerance and ability to survive lethal
antibiotic treatments, have long posed challenges in understanding their formation and long-term survival.
They are a driving force of antibiotic resistance, so it is paramount that we learn more about them as the
antibiotic resistance problem continues to grow. Our study challenges the long-held consensus that
persisters are completely dormant and are of one single population. Our results clearly show that
persisters are not as dormant as once thought, and multiple populations of persisters form during lethal
antibiotic treatment despite the cells being genetically identical. We use wet lab and dry lab
(mathematical modeling) to demonstrate these new findings.
47

Introduction
Bacteria inhabit diverse and harsh environments, requiring them to adapt to stress by altering their
metabolic states, a phenomenon widely known as persistence[1, 2]. Persistence is a bet-hedging strategy
employed by all free-living bacteria to endure stress[2]. Persisters are a small fraction of the bacterial

population that arises from inherent phenotypic heterogeneity (switching or transition) rather than genetic
mutations. They show reduced cellular activity and do not divide under stress[3]. When the threat subsides,
persister cells can resume their growth, leading to recurrent infections that often require prolonged and
repeated antibiotic therapy. Recent studies have demonstrated that persisters also have a high mutation
rate[4], which has been linked to the evolution of antibiotic resistance, a pressing public health distress. It is
the leading cause of death worldwide, outpacing other pathogens, including diseases such as HIV or
malaria[5]. Global death projections are now estimated at 10M/year by 2050[6, 7] unless substantial measures
are taken to curb this disturbing trend. Here, we specifically focus on probing the persister survival
mechanism under antibiotic stress.
61
Despite being discovered over eight decades ago, the molecular mechanisms responsible for persistence
are still unclear and debated. An excellent approach to determine the essential genes and networks for
persistence is to study the transcriptome profile —comprising the entirety of mRNA synthesized by cells
at a specific time—of persister cells through RNA sequencing (RNA-Seq). Nevertheless, due to their low
abundance and intrinsic transient nature, it is enormously challenging to study their transcriptomic profile.
Thus, many previous transcriptomic studies have relied on sub-lethal concentrations or short-term
exposure to lethal antibiotic concentrations[8, 9]. The populations under scrutiny often encompass short-term
tolerant cells (dividing, slow-growing cells) but not necessarily persisters. It's worth noting that short-term
tolerance can mask persistence, and empirical evidence has demonstrated their distinct nature[10]. Our
recent work reveals that high persistence levels do not correlate with high short-term tolerance levels.
Hence, it becomes imperative to use lethal antibiotic concentrations when investigating the transcriptional
regulatory mechanisms of persisters[11].
74
The genes related to antibiotic survival are heavily disputed because of conflicting results and different
interpretations of data. Some argue protein synthesis is slowed in persisters due to slower transcription
and translation rates[12], while others provide evidence for ample protein synthesis[13]. Other metabolic
processes such as oxidative respiration[14] and antibiotic efflux[15] are also debated[12, 14-17]. The role of
adenosine triphosphate (ATP) and the tricarboxylic acid (TCA) cycle is particularly controversial[14, 16-19].
Much of the debate comes from looking at mixed tolerant and persister populations, and only individual
genes (not networks or clusters of genes).
82
In this study, we isolated RNA from persisters treated with high doses of an antibiotic for sufficient time
to kill short-term tolerant cells. We used time-course assays followed by mutation studies to test our
hypothesis that the persisters transcriptome is not static but changes over time. Our study also highlights
that differentially expressed commonly upregulated genes could be a potential regulator for persisters.
This study is a follow-up to a previous study where we showed that *E. coli* employs an array of networks
to survive antibiotic exposure. These networks collaborate in regulating the cellular response, with no
single gene or network solely controlling this intricate response[20]. Here, we conducted a network biology
approach to illustrate how gene responses within a network change or adapt at different treatment time
points: 3 h, 6 h, and 24 h. Through network centrality analysis, we identified potential key hubs. Our
results showed that the gene network responses vastly differ across these three time points. We employed
a mutation study with selected hypothetical genes to gain deeper insights and further validate our
potential target. We showed that the knockout of a single gene (or the alteration of a single network) can
reduce the ability of a population to survive antibiotics at specific time points but not for long-term
antibiotic stress. We also used mathematical modeling to elucidate the survival mechanism of persisters in
mutant strains when exposed to antibiotic stress, demonstrating the existence of multiple subpopulations
of persisters.
99
Results and Discussion
The overview of the experimental plan is shown in Fig. 1. We harvested RNA from a persister
subpopulation treated with high doses of a bactericidal antibiotic (ampicillin, a β-lactam antibiotic that

inhibits transpeptidase enzyme involved in cell-wall synthesis)[21] for sufficient time to kill short-term
tolerant cells and then subjected for sequencing to analyze the transcriptome profile of the persisters. We
previously showed that the *E. coli* Dh5αZ1 strain displayed a biphasic cell death curve, where the short-
term cells died quickly in the first phase, and the remaining cells (persisters) died slowly and steadily,
beginning at ~2 h. We used rapid filtration[22, 23] for larger samples, followed by flash freezing. Rapid
filtration allows us to isolate and wash cells quickly. Large-scale pelleting can be slow, the transcriptome
may alter over time, and RNA can be degraded during harvesting. We have empirically shown that
removing dead cells requires several washes when pelleting (but not with rapid filtration)[24]. Evidence
supports that most RNA from dead cells is unstable after death[25-27], and RNA is sensitive to intra- and
extra-cellular RNases (or ribonucleases) degradation. This is why RT-qPCR is often preferred for
detecting food pathogens or bacteria on surfaces over DNA[25, 26, 28-39]. The relative stability of DNA
compared to RNA facilitates its detection from dead cells over extended periods. We complemented our
approach with mutational studies (gene knockouts and gene overexpression) to validate our
transcriptomic findings further.

118
119
Fig. 1. Schematic diagram of the experimental design. Stationary phase cells were diluted 1:100 times in
pre-warmed MMB+ media and treated with Amp (0.1 mg/mL) for 24 h, and samples were isolated for
RNA sequencing before antibiotic treatment (0 h) and after antibiotic treatment at 3 h, 6 h, and 24 h. Cells
were isolated using rapid filtration to minimize dead cells. Subsequent steps involved RNA extraction,
RNA sequencing, and gene network analysis using bioinformatics tools. Overexpression and mutational
studies were conducted to test the network analysis and validate the transcriptome-derived findings.

126
Time-series persister transcriptome analysis showed differential expression of several genes. We
compared the transcriptome profiles at different time points to investigate the dynamic changes and/or
existence of multiple persister subpopulations in response to lethal antibiotic treatment. We identified
substantial differences in gene expression by comparing the expression patterns of a non-treated
population at 0 h with persister subpopulations at 3 h, 6 h, and 24 h. Our findings revealed a notable
number of differentially expressed genes at each time point: 378 at 3 h, 1478 at 6 h, and 310 at 24 h (Fig.
2 a). Interestingly, these genes showed diverse patterns of upregulation and downregulation. At 3 h, 160
genes were upregulated, and 218 were downregulated. This trend continued at 6 h, where 620 genes were
upregulated and 858 were downregulated, and at 24 h, 143 genes were upregulated, and 167 genes
downregulated (Fig. 2 a). This observation defies the commonly held hypothesis of a global metabolic
slowdown in persister cells. The fact that upregulated genes did not substantially outnumber
downregulated genes indicates a complex regulatory mechanism governing gene expression under
antibiotic stress. Here, the interactive heatmap showed a clear expression difference of all gene sets at all
time points (Fig. 2 b). This observation led us to focus on a particular set of genes, showing a similar
expression profile, and we found 27 genes were commonly upregulated at all time points (Fig. 2 c).
Among these 27 genes, 8 were identified as hypothetical (Table S1), while 7 were annotated as ribosomal

proteins. There is a variation in ribosome-associated proteins between persister cells and normal cells.
This causes the suppression of transcriptional activity in persister cells[40]. However, our particular interest
was hypothetical genes. These hypothetical genes have not been previously associated with antibiotic
survival. We assume these genes may play a pivotal role in the adaptive response of persister cells,
signifying their potential role in antibiotic survival. We wondered if these upregulated genes were solely
due to the antibiotic we used, so we tested another type of antibiotic. Of the 27 genes identified with the 6
h Amp treatment, 22 are also commonly upregulated with ciprofloxacin (Cip) treatment at 6 h (Fig S1).
This observation demonstrated that the commonly upregulated genes follow consistency with different
types of antibiotics; Amp targets cell wall formation[21], while Cip targets DNA synthesis[41].Our goal was to
determine how these genes were related to persister survival. We build off what we knew from previous
persister and transcriptomic analysis studies[24, 42-48]. Previous studies identified several persister-related
genes including *oxyR, dnaK, sucB, relA, rpoS, clpB, mqsR,* and *recA*, which were found to be related to
prominent persister formation with multiple antibiotics treatment[42]. We found that these genes showed
fluctuating responses at different time points with Amp and Cip treatments. However, our study
particularly focused on commonly upregulated genes, including the hypothetical genes and their response
to long-term persister survival.

159
160
Fig. 2. Comparative transcriptome profile analysis of the population before and after antibiotic treatment
shows that the gene expression profile is not the same over time. **(a)** Volcano plots for 0 h (non-persister
population) vs 3 h (persister population); 0 h vs 6 h, and 0 h vs 24 h. **(b)** Heatmap visualization for gene
expression shows that many genes are differently expressed at different points in Amp treatment. Each
row corresponds to a gene, and each column represents treatment times. The intensity of the color at each
intersection point indicates the level of gene expression in each time point, with red denoting high
expression and blue denoting low expression. The intensity of the colors at each intersection point
signifies the gene expression levels in the respective sample. **(c)** Venn diagram for transcriptome
comparison of commonly upregulated, downregulated, and stable genes on 0 h vs 3 h, 0 h vs 6 h, and 0 h
vs 24 h. We focused on the 27 commonly upregulated genes, hypothesizing that they likely are important
for persister long-term survival. Differentially expressed genes are defined to be those with p-adjusted

values <0.1 and fold change higher or lower than ±2 (these same parameter values were employed for all
subsequent figures).

174
Topological variation of differentially expressed genes during antibiotic treatment. Using a cluster-
centric top-down approach, we analyzed the changes in gene network response and identified common
key network hubs at different time points. Although there are certain limitations of studying protein or
gene networks and their pattern of interconnections (refers to network topology) due to minimal
resources for predicting network interactions and dynamic changes depending on cell types[49, 50], this type
of analysis can be a powerful tool to explore interactions that are challenging to identify by other means.
The emergence of systems biology and mathematical modeling, encompassing graph theory, and neural
networks, has facilitated the integration of fundamental cellular components (such as genes, RNAs,
proteins, etc.) into a unified intricate network[51]. Using this systematic approach, we provided valuable
insights into the network responses of differentially expressed genes with lethal doses of antibiotics
treatment at different time points (Fig. 3 a). We constructed the gene network and identified major
clusters of genes using the molecular complex detection (MCODE) method[52], which detects highly
densely connected nodes in a network. We found a significant fluctuation of gene network response at
different time points: significant variation of edges (connections) (Fig. 3 b i) and the associated nodes
(genes) that form gene clusters (Fig. 3 b ii and Fig. S2 a-c) at different time points. The number of
upregulated genes was higher at 6 h compared to 3 h and 24 h treatments. In comparison, the number of
downregulated genes was significantly lower at 24 h, indicating a fluctuation of gene network response
(Fig. 3 b ii). The stable genes in the clusters may be responsible for maintaining the dormant state, while
the downregulated genes may play a role in preserving cellular resources during dormancy. Notably, it is
predicted that a network or cluster will undergo minimal to no significant alterations under normal
conditions. However, when subjected to external perturbations, there is a high likelihood of experiencing
substantial changes in gene expression and network reconfiguration. Based on these observations, we
have two competing hypotheses:
Hypothesis 1: Long-time antibiotic exposure may significantly alter network response.
Hypothesis 2: Little or no alterations in the network response. In other words, any variations observed in
the network's behavior are due to random chance or noise rather than deliberate or systematic changes in
gene expression or network response.

202
Network analysis shows a dramatic change in gene network response (Fig. 3 b and Fig. S2 a-c). This
means that the overall network connectivity of a gene has a greater influence on its expression, and it
seems likely that multiple genes or pathways influence how bacteria survive against stress. These results
support Hypothesis 1, that multiple persister subpopulation exists, and due to stress, network
reconfiguration occurs. This happens due to a 'one-to-many' gene interaction rather than a 'one-to-one'
interaction, facilitated by cell-state transitions and network switching. Similar to cancer cell metastasis[53,
54], the dynamic network switching of genes may allow reversible phenotypic plasticity (cell-state
transition), causing resistance to drug and survival benefits to the cells. Next, we focused on identifying
the subset of genes with the strongest control in the entire network and the potential roles of the key genes
(Table S1) in the network at different time points.

Network centrality analysis highlights potential key hubs. Our goal was to identify key gene hubs that
are important to surviving antibiotics. We analyzed each gene's network centrality (such as degree,
betweenness, and closeness) to demonstrate the gene attributes in a network. These classic network
centrality[55] initially developed for characterizing social networks that have been transposed to biological
networks to determine the hierarchy of nodes or edges in an intricate network. Thus, it provides
information on a gene's importance in a central network based on connectivity, or information flow
throughout the network. A high centrality score indicates that a node has a larger-than-average number of
connections in a network. We compared the centrality score of 27 genes we identified that are commonly

upregulated throughout a time course of antibiotic treatment to the entire gene sets of the network at
different time points. One of the key aspects of our analysis was the comparison of the centrality scores of
differentially expressed genes with the average centrality score of all genes within the network. At 3 h, a
substantial 23% of expressed genes showed high betweenness centrality scores (Fig. 3 c), while at 6 h and
24 h, the proportion decreased slightly to 21% (Fig. S3 i-ii).

Similarly, when considering closeness centrality, ~33% of genes displayed high scores at 3 h (Fig. 3 c).
This percentage slightly diminished to ~32% at 6 h and ~31% at 24 h (Fig. S3 i-ii). These results indicate
that a substantial portion of genes maintains a central position regarding their ability to interact with other
genes within the network rapidly. A robust node (genes) plays a significant role as this node is minimally
affected by disruption of other nodes in a network[56], and this might be the central gene for bacterial
persisters. What's particularly intriguing is the overlap between genes showing high betweenness and
closeness centrality at all time points. Most of the commonly upregulated genes (excluding the
hypothetical genes) showed high betweenness and closeness centrality, suggesting their importance to
persister survival. This observation suggests that these genes might play critical regulatory roles early in
the process, thereby contributing to the long-term survival of persister cells.

It was unsurprising that 8 genes of the 27 commonly upregulated genes showed lower centrality scores
than all genes' average scores because they are hypothetical genes. Hypothetical genes often represent
novel or less-studied elements within the network; thus, we often do not know what controls their
expression or how their protein productions interact with the expression of other genes. The involvement
of these genes should not be disregarded, as they might serve as vital regulatory players in processes yet
to be fully understood. But it does highlight the importance of studying how hypothetical genes are
related to persisters. In addition, we recognize that the multi-pronged network approach and the resulting
data have limitations and can potentially identify false positives that can compromise the prediction of
essential nodes in a network. So, experimental validation of this data is indispensable. We considered this
aspect by investigating 7 hypothetical genes, recognizing their contribution to the persister survival
through wet-lab mutational studies and overexpression experiments (Fig. 4 a-b).

Identifying gene clusters with potential biological functions. We investigated the roles of various genes
at different time points by integrating biological interpretation and functional gene grouping by ClueGO,
a Cytoscape plug-in[57, 58] (Table S2 and Fig. S4). We focused on known oxidative, cellular, and antibiotic
response genes. We found gene (both up and downregulated) clusters with significant fluctuation at
different time points (Table S2), suggesting a potential shift in the cellular dynamics or adaptation
mechanisms. At 6 h, the highest number of upregulated genes known to be related to oxidative, cellular,
and antibiotic response genes were identified. Surprisingly, no known oxidative, cellular, and antibiotic
response genes were upregulated at 24 h, This indicates that other genes are more important at 24 h, or the
oxidative, cellular, and antibiotic response proteins produced at 6 h were sufficient (intact and functional)
to remain by 24 h, thus no increased expression is required. Our observations strongly suggest that the
response of genes to antibiotic treatment is dynamic, and the changes in the network response are possibly
linked to the long-term survival of persister cells.

263
Fig. 3. a. This figure illustrates the construction of a gene interaction network model using upregulated,
downregulated, and stable genes. Each gene is represented as a node, and the interactions between genes
are depicted as edges. The network was analyzed using MCODE to identify gene clusters at different time
points. The clusters were then compared to showcase variations in the gene network responses. Key nodes
within the network were identified through centrality analyses involving node degree, betweenness, and
closeness. **b.** Network analysis shows dynamic topological variation in gene networks at different
treatment time points. **i.** Significant fluctuation in edges: comparing the number of edges within the

network highlights fluctuations relative to the total number of edges. At 6 h treatment time-point, the number of edges is higher than 3 h and 24 h, indicating dynamic network changes. **ii.** Top-ranking clusters are shown at three distinct time points: 3 h (total clusters: 62), 6 h (total clusters: 59), and 24 h (total clusters: 53). Here, we highlight the top rank cluster for each time point, and the remaining clusters for each time-point shown in Fig. S2 a-c. At 3 h, the top rank cluster constituted 321 nodes and 1,115 edges, and the expression of genes vastly fluctuated (i.e., stable: 293; up: 7; down: 21). At 6 h, 278 nodes and 1,882 edges were detected with variable gene expression (i.e., stable: 246; up:19; down:13); and 24 h the top rank cluster was associated with 242 nodes and 914 edges with highly fluctuated gene expression (i.e., 229, up: 6, down: 7). The results indicate that the total number of nodes and edges of the entire networks varied at 3 h, 6 h, and 24 h. **c.** Network analysis predicts key hubs. Identification of potential network hubs at 3 h: Degree (k) vs. Betweenness Centrality (BC) and Degree (k) vs. Closeness Centrality (CC) analyses identify commonly upregulated genes as potential network hubs. The remaining 6 h and 24 h centrality analysis is described in Fig. S3 i-ii. Degree centrality refers to the distribution of nodes' degrees (number of connections) in a network. It provides insights into the network structure and can reveal important connectivity, centrality, and overall network behavior characteristics. Betweenness centrality (also known as a bottleneck) quantifies how often a node appears on the shortest paths between pairs of other nodes in the network[59], whereas closeness centrality quantifies how quickly a node can interact with other nodes in the network[60].

Overexpression of hypothetical genes decreases growth rate. We cloned all the selected hypothetical genes under IPTG/Ara promoter (Fig. 4 b i). When we overexpressed the genes, most of them had significant growth defects. This shows that overexpression disrupts normal cellular processes. Notably, the overexpression of two genes, *yjbE* and *yajI*, led to the cells entering a nondividing state. This state lasted several hours before cells began growing, likely because of a mutation. To explore this, growing cells were subcultured without an inducer and then tested to see if they went into a non-dividing state again. Induced and uninduced cultures had no growth defects, suggesting a mutation (data not shown). Given the well-established connections between the growth rate and both short-term and long-term survival under antibiotic stress[61-64], we chose not to employ overexpression studies to investigate these genes' involvement in persistence. Instead, we employed gene editing to create knockout mutants, enabling us further to explore the role of these hypothetical genes in persistence.

Knockout of hypothetical genes reduces persister survival under antibiotic treatment initially (3 h and 6 h), but not in the long run (24 h or longer). We focused on hypothetical genes not previously shown to play a role in persistence. We first mutated hypothetical gene *yjbE* because of its dynamic expression profile, and it showed the highest expression levels at 6 h (>10-fold) and 24 h (>7-fold) of all the hypotheticals. The knockout mutant was constructed CRISPR/cas9-based genome editing tool[65] (we modified the plasmid and the protocol for this knockout to get the mutants in our strain of *E. coli*, see Methods) to test the effect(s) of the given genes at different time points (Fig. 4 b ii-iii). The *yjbE* knockout mutants did not show any growth defects under normal conditions, which contrasts with the growth defects observed upon gene overexpression. We wanted to know if the other hypothetical genes would act similarly, so we knocked out all of them but *ycfJ* and *ygfB* because getting those knockouts were challenging. All seven knockout mutants did not show any growth defects under normal conditions, which contrasts with the growth defects observed upon gene overexpression. This outcome was unsurprising, given that we grew the cells in defined media with ample nutrition and a well-controlled environment. If these genes are associated with persistence or environmental stress survival, it would be unlikely to observe growth defects under optimal environmental conditions. However, under lethal antibiotic stress, based on our transcriptional analysis, we would expect a dramatic decrease in persister levels in the mutants.

As expected, there was a dramatic decrease in persister levels in the mutants following Amp treatment compared to the wild type: 4-5-fold reduction at 3 h, and 10-14-fold reduction at 6 h (Fig. 4 b ii-iii).

Remarkably, the persister levels in the mutants did not significantly differ from the wild type at 24 h. This
result supports the existence of multiple persister subpopulations in an isogenic *E. coli* population, and
different persister subpopulations have unique survival mechanisms, each associated with varying decay
rates. This led us to propose two plausible hypotheses (Fig. 5a):
(1) Within a given population, multiple phenotypes of persisters result in different subpopulations, and
each phenotype dynamically stems from non-persisters. These distinct phenotypes emerge during the
logarithmic growth phase or the rapid death phase (of the biphasic death curve), given that susceptible
cells perish by the end of the rapid death phase.
(2) One persister phenotype forms dynamically from non-persister populations and subsequently adapts
sequentially or non-sequentially to multiple other phenotypes. This could happen during the fast or slow
death phase.

Of these two hypotheses, we favor the second one, which involves an adaptive response, as it aligns with
our transcriptomic findings. This intriguing possibility suggests that the persister state is highly dynamic
and adaptable, and cells have multiple strategies to survive antibiotic stress. Intuitively, both persister and
non-persister cells undergo cell-state transitions, giving rise to various persister phenotypes that facilitate
bet-hedging and resistance to antibiotic onslaught. However, with the current data, it remains unclear
whether the occurrence of various persister phenotypes is a result of random or reversible phenotypic
switching, and if these transitions occur before or after antibiotic treatment.

Mutants and wild-type strains showed similar growth rates (Fig. 4b ii), suggesting that any difference in
their death rates would be unrelated to their ability to grow and divide. We observed significant variations
in persister levels at different time points between wild-type and mutant strains (Fig. 4b iii), illustrating
the presence of multiple subpopulations of persisters. When a gene is removed, consequently disrupting
certain survival mechanisms, it eliminates persister phenotypes reliant on those mechanisms, leading to
decreased early time point persistence levels (at 3 h and 6 h). This phenomenon accounts for the observed
reduction in early persistence among mutant strains. Furthermore, it provides compelling evidence that
the selected hypothetical genes regulate the persister cell's long-term survival and antibiotic stress
response. However, this decrease does not seem to have a significant long-term effect on the cells. In the
long run (24 h to 72 h), the persister phenotypes with the slowest decay rates survive regardless of gene
disruptions.

Generally, the identification and characterization of persister cells have been based on biphasic death
curves[66]. However, the binary nature of this curve may not accurately represent the diverse array of decay
rates among persister cells. A recent study demonstrated the advantages of employing multi-phase killing
curves. Depending on the intricate interplay of decay rates within persister cell subpopulations, these
curves can manifest in various phasic (triphasic, tetraphasic, etc.) patterns. This approach provides more
accurate information as it accounts for bacterial survival times that cannot be properly predicted through a
singular focus on a biphasic death curve[64]. Our results suggest that the long-term behavior of persisters
resembles those of multi-phase killing curves, featuring multiple decay rates (Fig. 4b iii and Fig. 5b).
These findings indicate that persister cells are not a homogeneous entity, but consist of multiple distinct
subpopulations, each with its unique survival characteristics and decay rates. These challenge the "one-
size-fits-all" approach as it has become increasingly evident that persister cells are heterogeneous. They
do not conform to a uniform response to antibiotics, which a simple biphasic curve might imply. Instead,
the presence of multiple decay rates within persister populations highlights the need for a more subtle and
comprehensive approach to understand the intricate mechanism of these cells.

We developed a simple mathematical model (described in the Methods) to explore further the variations
in decay fraction rate between *E. coli* wild type and mutant strains in response to lethal antibiotic
treatment. We plotted the bacterial survival fractions over time, showing that different persister
subpopulations had distinct decay rates. Notably, the slow-decay fraction of the mutant strain exhibited a

significant 20-fold decrease, emphasizing the importance of decay kinetics in bacterial survival. This
result again supports our hypothesis that multiple subpopulations of persisters exist, each with distinct
decay rates (Fig. 5b).
376
377   It is well established that persisters form due to phenotypic heterogeneity in the general *E. coli*
378   population. We identified here that the persister population itself has its own heterogeneity and can switch
379   between different cell-states. This heterogeneity (diversity) in the persister population allows
380   subpopulations of persisters to survive antibiotic stress, but they die at different times, which is evident by
381   the different decay rates.

a. Experimental workflow

i. Overexpression → Growth rate comparison

ii. Gene knockout (CRISPR-Cas9) → Growth rate comparison → iii. Persister level comparison

Gene editing → Plasmid curing

b.

i. Overexpression slows the growth rate

ii. WT and mutant grow at a similar rate

iii. Mutants have fewer persisters at 3 h and 6 h than WT

Fig. 4. a. Experimental workflow. **i.** Hypothetical genes were under an inducible promoter ($P_{lac/ara-1}$) and in *E. coli* DH5αZ1 and grown with or without arabinose; uninduced: 0% induced: 0.2% arabinose. **ii.** Hypothetical genes were knocked out using the CRISPR/CAS-9 system, and the growth rate of each mutant was compared with the wild-type strain. **iii.** The persister level of the mutant was compared to the wild type at 3 time points: 3 h, 6 h, and 24 h (normalized by time 0 for each strain). **b.** Results: **i.** Overexpression of the hypothetical genes slows growth or causes cells to enter a non-growth state. **ii.** Wild-type and knockout mutant growth rates are nearly identical in the same media. **iii.** Persister level is significantly reduced in mutants at 3 h and 6 h compared to wild type. However, no significant decrease was observed at 24 h. Persister level comparison was shown up to 72 h for *yjbE* gene, and 24 h for all other mutants. These results show that the 7 hypothetical genes help cells survive in the first 6 h but are not helpful by 24 h. *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.*

a. Competing hypotheses

Rejected hypothesis: One phenotypical
type of persisters is in a given population

N: Non-persister population
P: Persister population
K: Decay rate
K_0: Decay rate is faster than K_1, K_2, K_x
⊘: Cell death

Hypothesis: Multiple persister phenotypes within a given population

1. Multiple persister phenotypes form from non-persisters

2. Multiple persister phenotypes form sequentially or non-sequentially

b. The mathematical model shows multiple decay rates are required
to reconstruct *in vivo* results

407
Fig. 5. a. Model figures demonstrating competing hypotheses for multiple persister subpopulations
408
and decay rates a. Null hypothesis: Only one phenotypical type of persisters exists within a population.
409
The persister population (P) is formed from the non-persister population (N). Non-persisters and
410
persisters alternate in the process of persister formation and revival. Alternative hypothesis: (1) Multiple
411
persister phenotypes (P_1, P_2, P_X) exist, and all phenotypes form from non-persisters (N). (2) A single
412
phenotype of persisters forms from non-persister cells, and eventually, that phenotype sequentially or
413
non-sequentially adapts to other phenotypes. Persister formation and revival involve a dynamic
414
interchange between non-persisters and persister subpopulations and simultaneous alternation within the
415
persister subpopulations. In any scenario, non-persisters (susceptible cells) have the highest decay rate
416

(K_0) compared to the decay rate of any persister subpopulation. **b.** The plot shows the dynamics of the
bacterial surviving fraction in response to lethal antibiotic exposure as predicted by the model (see
Methods). The model was constructed considering three different subpopulations: a fast-decaying (non-
persister population (N) having susceptible and tolerant cells), a slow-decaying (persister subpopulation 1
(P_1)), and a very slow or non-decaying population (persister subpopulation 2 (P_2)). The decay rate of the
wild type and the *yjbE* mutant was calculated individually and concurrently, showing a 20-fold decrease
in the slow-decaying fraction (P_1) in the mutant, while the non-decaying fraction (P_2) remains the same in
the mutant. It indicates that multiple and distinct persister subpopulation maintain their unique decay
rates. The plot axes represent percent survival in log scale (Y-axis), time (h) (X-axis).
426
Outlook
Persisters are often described as metabolically dormant and controlled by one or a few sets of genes[15, 16,
67]. However, although persisters are dormant with respect to growth, they render an active response
against antibiotics[16, 20]. Here, we argued that persisters have an altered cellular activity rather than being
in a dormant state. We mutated 7 hypothetical genes and observed their effect on persister survival, which
validated the biological relevance of our transcriptomic analysis. Our result also shows that not a single
gene is responsible for persister formation; it's a total network response, and persister can adjust their
transcriptome to survive lethal antibiotic stress. Our 7 mutational experiments and mathematical modeling
results clearly demonstrate that more than one subpopulation of persisters can develop from an isogenic
E. coli population. Another line of evidence supporting this claim is obvious; persister cells continue to
die at a slow decay rate (Fig. 1, see death curve). If persisters were of one identical population, the
population should die relatively at the same rate. But we always see a slow decay rate because different
subpopulations die quicker than others.
440
What is the selective advantage of having multiple subpopulations of persisters? We propose that,
logically, this is a good strategy for survival. Antibiotic-resistant bacteria have molecular mechanisms
available to detect a specific threat, respond, and allow for continued growth. However, non-resistant
bacteria do not contain the genes to detect a particular threat, so individual cells react to the danger in
their own manner. Most cells die, but those that can persist will survive longer. Some persister cells (a
subpopulation) have responded in a manner that will allow them to survive longer than other persister
subpopulations. This is what our results suggest. It is also possible that one subpopulation of persisters is
converting to another subpopulation of persisters, but currently, we have no evidence for this. A simple
analogy to this scenario is to imagine boats at sea encountering a devastating, fast-moving storm. An
aircraft courier may be large enough and stable enough to weather the storm; they are resistant to it. Small
boats would not be resistant to the storm but can be persistent by different manners. For example, boats
may speed in different directions, and those who move away from the storm will survive longer than
those who stay put or head into the storm. Those who move completely out of the storm have a higher
chance of surviving than those who only run in front of it because it may eventually catch them. Others
may radio for help, and so on. Bacterial persisters sample different methods to survive, and this
heterogeneity in response is what makes them so difficult to kill.
457
We have previously hypothesized that the presence of a single persister gene is unlikely, but disruption of
essential cellular networks leads to persister formation. The results shown here support this. We also
hypothesize a "ground state" where cells have entered a non-growing, reduced activity state, but are not
quite persisters. Once in the ground state (a transition stage), cells can shift into persistence (a more
dormant state) or return to the dividing phase. Cells can reach the ground state by several different means,
but the ground state occurs when cells encounter internal or external stress. In this manner, the condition
that the cell entered the ground state determines the type of persister that will form. We make this
hypothesis based on the fact that persister-related genes are constantly identified, and if one mechanism is
reasoned for persister formation by one research group, another's group disclaimed that mechanism[42-46,
68](suggesting not one gene is responsible for persister formation). The minimal genome contains mainly

essential genes (less than 500 genes) and still form persisters (other microbes contain genes that have
been to be persister-related, but the minimal genome lacks them)[69], slowed growth either externally[64] or
internally[2, 70] appears to increase persister levels (showing that both internal and external stress alters
persister levels). Our results clearly show that more than one subpopulation of persisters develops because
of different stress response genes. These genes affect the type of persister and how long they will survive
antibiotics.
474
In addition, persisters are present at substantially higher levels in stationary phase compared to log
phase[71]. Although different phases can be identified in batch cultures (in this order): lag, log, stationary,
death, and long-term stationary (LTS phase), it has been proposed that many bacteria exist in the LTS
phase longer than any other phase in natural environments[72, 73]. *E. coli* cultures have been retained in LTS
stage (without additional nutrients) for more than 5 years while maintaining viability[72]. Although
persisters were identified first, persisters and LTS cells share similar phenotypes. They are unlikely to be
in the same state because LTS have been shown to mutate to survive, while persisters can survive without
mutation. It is possible that LTS could be cells in the ground state. Here, we specifically use the term
"ground state" because the term describes a low-activity state. Though we do not have enough evidence to
declare the "ground state" hypothesis correct, we have ample evidence to put it forward to the scientific
community.
486
Methods
Microbial strains and media. *Escherichia coli* DH5αZ1 containing the p24KmNB82 plasmid was used
in this study. DH5αZ1 is a derivative of *E .coli* K12 strain, and it has been used in our previous
persistence studies[24, 74]. The cultures were grown in MMB+ as described in our previous work[11, 24]. We
used 0.5% glycerol as the carbon source and each of the 20 canonical amino acids (0.04 mg/mL) for fast-
growing cultures. Cultures were grown in MMB+/kanamycin (Km, 0.025 mg/mL) or on Miller's lysogeny
broth (LB) agar plates/Km (0.025 mg/mL). All cultures were incubated at 37 °C and shaken at 300 rpm.
494
495   **Persister assay.** The assay was done as previously described[11, 24]. Briefly, overnight cultures were diluted
1/100 into fresh media and grown until they reached between OD_{600} ~0.1- 0.2. A reduced volume of
culture (10 mL) was aliquoted into three 100 mL flasks, and grown for ~2-3 h until the culture reached to
mid-log phase having ~1E+8 CFU/mL (~OD 0.5). After that, the culture was treated with a bactericidal
concentration of Amp (0.1 mg/mL) for 3 h at 250 rpm, 37°C. Persister percentage was calculated by
comparing CFUs per milliliter (CFU/mL) before antibiotic treatment to CFU/mL after antibiotic
treatment. Plates were incubated at 37ºC for 48 h, then scanned using a flatbed scanner. Custom scripts
were used to identify and count bacterial colonies[75, 76]. For the time-kill curve, antibiotic treatment was
extended to 24 h, and the number of persisters was recorded at 3 h, 6 h, and 24 h.
503
504   **RNA sample preparation and sequencing.** We used our previously developed methodology to harvest
RNA from persisters[20]. Cultures are first treated with lethal Amp (0.1 mg/ml) concentrations, and their
transcriptomes are isolated and compared to an untreated population. In brief, **s**amples were treated with
Tween (0.2%) and chloramphenicol (0.005 mg/ml) then immediately filtered out of solution (using rapid
filtration) and resuspended in lysis buffer containing 25 mM Tris pH 8.0, 25 mM NH4Cl, 10 mM
MgOAc, 0.8% Triton X-100, 0.1 U/μL RNase-free DNase I, 1 U/μL Superase-In, ~ 60 U/μl ReadyLyse
Lysozyme, 1.55 mM Chloramphenicol, and 17 μM 5′-guanylyl imidodiphosphate (GMPPNP); Rapid
filtration allows for the harvesting of any whole-cell; this minimizes RNA degradation because samples
are immediately frozen in liquid nitrogen have suspension in lysis buffer. The solution was then frozen in
liquid nitrogen and stored at − 80 °C. Frozen pellets were thawed on ice and centrifuged for 20 min at
16,000×g at 4 °C with 0.3% Sodium Deoxychorate to pellet particulate matter. The supernatant was then
removed, and then RNA was column purified (Qiagen miRNeasy cat. No. 217004) before removing

rRNA (Invitrogen RiboMinus cat no. K155004) and prepping the cDNA libraries (NEBNext Ultra II cat

no. E7103S). Libraries were multiplexed, then pooled and sequenced by Illumina HiSeq paired-end

sequencing by GENEWIZ. The data is analyzed using a combination of Geneious Prime NGS software[77],

custom Python scripts modified from our previous transcriptome analyses[78], and the DESeq2 R package[79].

We consider significant changes in gene expression to be greater than two-fold with an adjusted p-value

<0.1. InteractiVenn[80] was used to generate Venn-diagram.

Network construction and data analyses. We constructed the network and retrieved nodes degree,

betweenness and closeness centrality scores using Cytoscape (v3.10.0) StringApp (v2.0.1)[57, 81, 82].

Molecular Complex Detection (MCODE)_v2.0.3 method[52] was used to identify the molecular clusters of

the entire network. MCODE used the graph theoretic clustering algorithm to identify densely connected

regions in a large protein-protein interaction network (PPIn) representing molecular complexes[52]. We also

investigated the functional roles of oxidative, cellular, and antibiotic stress response genes through

ClueGO (v2.5.10), a Cytoscape plug-in[57, 58]. We used default parameter values during network

construction via String App (i.e., Network type: full String network, Confidence cutoff: 0.4) and selective

parameters values during MCODE analysis like, Find Clusters: in Whole Network; Network Scoring - a)

Include Loops: Turn off, b) Degree Cutoff: 10; Cluster Finding- a) Haircut: Turn on, b) Fluff: Turn off, c)

Node Score Cutoff: 0.2, d) K-Core: 2, e) Max. Depth: 100. During ClueGO analysis, default parameter

values were used such as Analysis Mode: Functional Analysis, Visual Style: Groups, Ontologies:

Biological and cellular process.

Overexpression of hypothetical genes. Since our control strain contains the plasmid p24KmNB82 (CFP-

LAA), we cloned the seven selected hypothetical genes under the same promoter ($P_{lac/ara-1}$) and ribosome

binding sites as in REF[83]. Our control strain Dh5αZ1 produces two proteins (genes are on the genome),

AraC and LacI, which are used to control the expression of genes through the $P_{lac/ara-1}$ promoter. AraC is

an activator in the presence of arabinose, while LacI is a repressor of this promoter. Isopropyl β-D-1-

thiogalactopyranoside (IPTG) acts as a repressor of LacI, thus allowing for activating the $P_{lac/ara-1}$

promoter. We used both arabinose (0.2%) and IPTG (0.100 mM) to express the hypothetical genes.

Knockout of hypothetical genes. To knockout the hypothetical genes, we used the clustered regularly

interspaced short palindromic repeats (CRISPR)-associated nuclease 9 (Cas9)-based genome editing tool

pEcCas/pEcgRNA system explained in ref[65], but had to modify the protocol and plasmid to get the system

to work efficiently with our *E. coli* strain. The pEcgRNA plasmid was originally designed for

spectinomycin selection; however, the *E. coli* strain we used in our work contains a streptomycin resistant

gene. To facilitate effective selection, the pEcgRNA plasmid was modified by introducing an ampicillin-

resistant gene at the BglII/XbaI restriction sites. In brief, 0.040 mL of pre-induced (with 1% arabinose)

competent cells containing pEcCas plasmid was added with pEcgRNA plasmid (200 ng - 800 ng), and

donor dsDNA (200 ng - 800 ng). Then, the mixture was electroporated in a precooled 2 mm Gene Pulser

cuvette (Bio-Rad, Hurcules, USA) at 1.8kV, 25uF, 200 ohms. After electroporation, freshly prepared 1mL

of SOC media (containing 1.5% arabinose) was added, and the culture was shaken for 1 h at 250 rpm,

37°C. Then, the culture was plated on LB/Amp (0.1 mg/mL)/Km (0.05 mg/mL) plates and incubated at 37

°C for 24 h. Each knockout was confirmed by colony PCR and sequencing. After the knockout, the edited

clone was inoculated into 10 ml of LB with Km (0.05 mg/mL), and 100 mM rhamnose and grew for 12

hours at 37 °C to cure pEcgRNA plasmid. Then, the culture was again transferred to 10 ml of LB with no

antibiotics and grew for overnight at 37 °C to cure pEcCas plasmid. Then, the culture was plated on LB

agar with 5% sucrose plate and incubated overnight at 37 °C. Finally, ~48 colonies from sucrose plates

were streaked on LB, LB/Amp (0.1 mg/mL), and LB/Kan (0.05 mg/mL) plates and incubated at 37 °C

overnight. Colonies that only grew on LB plates were selected as our knockout strain. Finally, we

transformed the p24KmNB82 plasmid[24] in each knockout strain to keep it similar to our control strain.

Determination of growth curve. Overnight cultures were diluted into OD 0.1 (measured in

SpectronicTM 200E), and 0.03 mL of diluted cultures were inoculated into individual wells containing

0.27 mL of SP16 media in a 96-Well Optical-Bottom Plate with Polymer Base (Thermo Fisher) to

measure OD at 600 nm using FLUOstar Omega microplate reader.

Mathematical model. To model the dynamics of the bacterial surviving fraction in response to lethal

antibiotic exposure, we consider three different subpopulations: a fast-decaying (non-persister population

having susceptible and tolerant cells), a slow-decaying (persister subpopulation 1), and a very slow or non-

decaying population (persister subpopulation 2). In persister subpopulation 2, we considered the time scale

of decay to be zero in the experimental timespan since the death rate is so low after 24 h based on our

experimental result. But it is notable to mention that in real scenarios, the decay rate cannot be zero because

no cells can live forever, and at some point, cells will die. However, let the fraction of cells in these

populations be f_1, f_2 and f_3, respectively, with corresponding decay rates of γ_1, γ_2, and 0. The fraction of

surviving cells $f(t)$ at time t after antibiotic exposure is then given by:

$$f(t) = f_3 + f_2 e^{-\gamma_2 t} + + f_1 e^{-\gamma_1 t}$$

with $f_1 = 1 - f_2 - f_3$. The parameters $f_3, f_2, \gamma_2, \gamma_1$ are estimated by performing a least square fit of $f(t)$

with data.

We initially fitted this equation to data from WT and the yjbE mutant individually and realized the biggest

difference was in f_2, the fraction of cells in the slow-decaying state. Next, we fitted the data from both WT

and mutant simultaneously, keeping f_3, γ_2, γ_1 fixed between the two genotypes while allowing f_2 to vary

between them. Our fitting yields $f_3 \approx 2.5 \times 10^{-6}, \gamma_1 \approx 2.6\ hr^{-1}, \gamma_2 \approx 0.24\ hr^{-1}$ with $f_2 \approx 2 \times 10^{-3}$

(for WT) and $f_2 \approx 10^{-4}$ (for mutant). The model fits are shown in Fig. 5b, where the change in a single

parameter, a 20-fold decrease in the slow-decaying fraction in the mutant, is sufficient to capture the decay

kinetics of bacterial survival.

Statistical analysis. All data is presented in the manuscript as mean ± SEM of at least three independent

biological replicates. Statistical significance was assessed using an f-test to determine variance ($p < 0.05$

was considered to have significant variance), followed by a two-tailed t-test with unequal variances (if F

statistic > F critical value) or equal variances (if F statistic < F critical value).

Competing interests

No conflict.

Authors' contributions

K.R., T.H., R.A., performed the experiments, and ran data analysis. A.S. constructed mathematical

modeling. N.C.B. and T.H. formalized the research concept and designed the experiments as well as

planned and directed the project. All authors contributed to discussing and editing the manuscript.

Acknowledgments

We would like to thank Sudeshna Roy, Thomas Ebrahem, and Isaac Kovash, for their contributions to

some experimental works. This work is supported by the National Science Foundation award numbers

1922542, 2240028, and 1849206, and by a USDA National Institute of Food and Agriculture Hatch

project grant number SD00H653-18/project accession no. 1015687.

Data availability

All the data that supports the findings of this study are included in this published article (and its

supplementary information files). Also, any additional data is available from the corresponding author

upon request (email: nicholas.butzin@gmail.com).

References:

1. Engl, C., *Noise in bacterial gene expression.* Biochemical Society Transactions, 2019. **47**(1): p. 209-217.
2. Balaban, N.Q., et al., *Bacterial persistence as a phenotypic switch.* Science, 2004. **305**(5690): p. 1622-1625.
3. Lewis, K., *Persister cells: molecular mechanisms related to antibiotic tolerance.* Handb Exp Pharmacol, 2012(211): p. 121-33.
4. Windels, E.M., et al., *Bacterial persistence promotes the evolution of antibiotic resistance by increasing survival and mutation rates.* The ISME journal, 2019. **13**(5): p. 1239-1251.
5. Murray, C.J., et al., *Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis.* The Lancet, 2022. **399**(10325): p. 629-655.
6. Huemer, M., et al., *Antibiotic resistance and persistence—Implications for human health and treatment perspectives.* EMBO reports, 2020. **21**(12): p. e51034.
7. Control, C.f.D. and Prevention, *Antibiotic resistance threats in the United States, 2019*. 2019: US Department of Health and Human Services, Centres for Disease Control and ….
8. Long, Y., et al., *Identification of novel genes that promote persister formation by repressing transcription and cell division in Pseudomonas aeruginosa.* Journal of Antimicrobial Chemotherapy, 2019. **74**(9): p. 2575-2587.
9. Pu, Y., et al., *Enhanced efflux activity facilitates drug tolerance in dormant bacterial cells.* Molecular cell, 2016. **62**(2): p. 284-294.
10. Ronneau, S., P.W. Hill, and S. Helaine, *Antibiotic persistence and tolerance: not just one and the same.* Curr Opin Microbiol, 2021. **64**: p. 76-81.
11. Hossain, T., A. Singh, and N.C. Butzin, *Escherichia coli cells are primed for survival before lethal antibiotic stress.* Microbiology Spectrum, 2023: p. e01219-23.
12. Wood, T.K., S.J. Knabel, and B.W. Kwan, *Bacterial persister cell formation and dormancy.* Appl Environ Microbiol, 2013. **79**(23): p. 7116-21.
13. Yin, L., et al., *Bacterial Longevity Requires Protein Synthesis and a Stringent Response.* mBio, 2019. **10**(5).
14. Orman, M.A. and M.P. Brynildsen, *Inhibition of stationary phase respiration impairs persister formation in E. coli.* Nat Commun, 2015. **6**: p. 7983.
15. Pu, Y., et al., *Enhanced Efflux Activity Facilitates Drug Tolerance in Dormant Bacterial Cells.* Mol Cell, 2016. **62**(2): p. 284-94.
16. Kim, J.S. and T.K. Wood, *Persistent Persister Misperceptions.* Front Microbiol, 2016. **7**: p. 2134.
17. Prax, M. and R. Bertram, *Metabolic aspects of bacterial persisters.* Front Cell Infect Microbiol, 2014. **4**: p. 148.
18. Wang, Y., et al., *Inactivation of TCA cycle enhances Staphylococcus aureus persister cell formation in stationary phase.* Sci Rep, 2018. **8**(1): p. 10849.
19. Orman, M.A. and M.P. Brynildsen, *Establishment of a method to rapidly assay bacterial persister metabolism.* Antimicrob Agents Chemother, 2013. **57**(9): p. 4398-409.
20. Deter, H.S., T. Hossain, and N.C. Butzin, *Antibiotic tolerance is associated with a broad and complex transcriptional response in E. coli.* Sci Rep, 2021. **11**(1): p. 6112.
21. Fernandes, R., P. Amador, and C. Prudêncio, *β-Lactams: chemical structure, mode of action and mechanisms of resistance.* Reviews and Research in Medical Microbiology, 2013. **24**(1): p. 7-17.
22. Bordag, N., et al., *Fast Filtration of Bacterial or Mammalian Suspension Cell Cultures for Optimal Metabolomics Results.* PLoS One, 2016. **11**(7): p. e0159389.
23. Mohammad, F., R. Green, and A.R. Buskirk, *A systematically-revised ribosome profiling method for bacteria reveals pauses at single-codon resolution.* Elife, 2019. **8**.

24.   Deter, H.S., T. Hossain, and N.C. Butzin, *Antibiotic tolerance is associated with a broad and*
complex transcriptional response in E. coli. Scientific reports, 2021. **11**(1): p. 1-15.

25.   McIngvale, S.C., D. Elhanafi, and M.A. Drake, *Optimization of reverse transcriptase PCR to detect*
viable Shiga-toxin-producing Escherichia coli. Appl Environ Microbiol, 2002. **68**(2): p. 799-806.

26.   Klein, P.G. and V.K. Juneja, *Sensitive detection of viable Listeria monocytogenes by reverse*
transcription-PCR. Appl Environ Microbiol, 1997. **63**(11): p. 4441-8.

27.   McKillip, J.L., L.A. Jaykus, and M. Drake, *Nucleic acid persistence in heat-killed Escherichia coli*
O157:H7 from contaminated skim milk. J Food Prot, 1999. **62**(8): p. 839-44.

28.   Sheridan, G.E., et al., *Detection of mRNA by reverse transcription-PCR as an indicator of viability*
in Escherichia coli cells. Appl Environ Microbiol, 1998. **64**(4): p. 1313-8.

29.   Szabo, E.A. and B.M. Mackey, *Detection of Salmonella enteritidis by reverse transcription-*
polymerase chain reaction (PCR). Int J Food Microbiol, 1999. **51**(2-3): p. 113-22.

30.   Yaron, S. and K.R. Matthews, *A reverse transcriptase-polymerase chain reaction assay for*
detection of viable Escherichia coli O157:H7: investigation of specific target genes. J Appl
Microbiol, 2002. **92**(4): p. 633-40.

31.   de Wet, S.C., et al., *An improved method for RNA extraction from carcass samples for detection*
of viable Escherichia coli O157:H7 by reverse-transcriptase polymerase chain reaction. Lett Appl
Microbiol, 2008. **47**(5): p. 399-404.

32.   D'Souza, D.H., F.J. Critzer, and D.A. Golden, *Real-time reverse-transcriptase polymerase chain*
reaction for the rapid detection of Salmonella using invA primers. Foodborne Pathog Dis, 2009.
6(9): p. 1097-106.

33.   Techathuvanan, C., F.A. Draughon, and D.H. D'Souza, *Real-time reverse transcriptase PCR for the*
rapid and sensitive detection of Salmonella typhimurium from pork. J Food Prot, 2010. **73**(3): p.
507-14.

34.   Miller, N.D., F.A. Draughon, and D.H. D'Souza, *Real-time reverse-transcriptase--polymerase chain*
reaction for Salmonella enterica detection from jalapeno and serrano peppers. Foodborne
Pathog Dis, 2010. **7**(4): p. 367-73.

35.   Techathuvanan, C. and D.H. D'Souza, *Optimization of rapid Salmonella enterica detection in*
liquid whole eggs by SYBR green I-based real-time reverse transcriptase-polymerase chain
reaction. Foodborne Pathog Dis, 2011. **8**(4): p. 527-34.

36.   Zhou, M., et al., *Development of a sigDE-based real-time reverse-transcriptase PCR for the*
detection of viable Salmonella enterica. Foodborne Pathog Dis, 2014. **11**(7): p. 537-44.

37.   Wang, L. and A. Mustapha, *EMA-real-time PCR as a reliable method for detection of viable*
Salmonella in chicken and eggs. J Food Sci, 2010. **75**(3): p. M134-9.

38.   Wang, L., Y. Li, and A. Mustapha, *Detection of viable Escherichia coli O157:H7 by ethidium*
monoazide real-time PCR. J Appl Microbiol, 2009. **107**(5): p. 1719-28.

39.   Malorny, B., C. Bunge, and R. Helmuth, *A real-time PCR for the detection of Salmonella*
Enteritidis in poultry meat and consumption eggs. J Microbiol Methods, 2007. **70**(2): p. 245-51.

40.   Cho, J., et al., *E scherichia coli persister cells suppress translation by selectively disassembling*
and degrading their ribosomes. Molecular Microbiology, 2015. **95**(2): p. 352-364.

41.   Campoli-Richards, D.M., et al., *Ciprofloxacin: a review of its antibacterial activity,*
pharmacokinetic properties and therapeutic use. Drugs, 1988. **35**: p. 373-447.

42.   Wu, N., et al., *Ranking of persister genes in the same Escherichia coli genetic background*
demonstrates varying importance of individual persister genes in tolerance to different
antibiotics. Frontiers in microbiology, 2015. **6**: p. 1003.

43.   Zhang, Y., *Persisters, persistent infections and the Yin–Yang model.* Emerging microbes &
infections, 2014. **3**(1): p. 1-10.

44. Li, T., et al., *Identification of novel genes involved in Escherichia coli persistence to tosufloxacin.*
Frontiers in Cellular and Infection Microbiology, 2020. **10**: p. 581986.
45. Bhatia, R.P., et al., *Transcriptomic profiling of Escherichia coli K-12 in response to a compendium*
of stressors. Scientific Reports, 2022. **12**(1): p. 8788.
46. Mohiuddin, S.G., A. Massahi, and M.A. Orman, *High-throughput screening of a promoter library*
reveals new persister mechanisms in Escherichia coli. Microbiology Spectrum, 2022. **10**(1): p.
e02253-21.
47. Sulaiman, J.E., C. Hao, and H. Lam, *Specific enrichment and proteomics analysis of Escherichia*
coli persisters from rifampin pretreatment. Journal of proteome research, 2018. **17**(11): p. 3984-
3996.
48. Sulaiman, J.E. and H. Lam, *Proteomic investigation of tolerant Escherichia coli populations from*
cyclic antibiotic treatment. Journal of proteome research, 2020. **19**(2): p. 900-913.
49. Han, J.-D.J., et al., *Evidence for dynamically organized modularity in the yeast protein–protein*
interaction network. Nature, 2004. **430**(6995): p. 88-93.
50. Ellis, J.D., et al., *Tissue-specific alternative splicing remodels protein-protein interaction*
networks. Molecular cell, 2012. **46**(6): p. 884-892.
51. Brasch, M.A., J.L. Hartley, and M. Vidal, *ORFeome cloning and systems biology: standardized*
mass production of the parts from the parts-list. Genome research, 2004. **14**(10b): p. 2001-2009.
52. Bader, G.D. and C.W. Hogue, *An automated method for finding molecular complexes in large*
protein interaction networks. BMC bioinformatics, 2003. **4**(1): p. 1-27.
53. Groves, S.M. and V. Quaranta, *Quantifying cancer cell plasticity with gene regulatory networks*
and single-cell dynamics. Frontiers in Network Physiology, 2023. **3**.
54. Saxena, K., et al., *Cancer: More than a geneticist's Pandora's box.* Journal of Biosciences, 2022.
47(2): p. 21.
55. Freeman, L.C., *Centrality in social networks: Conceptual clarification.* Social network: critical
concepts in sociology. Londres: Routledge, 2002. **1**: p. 238-263.
56. Scardoni, G., et al., *Node interference and robustness: performing virtual knock-out experiments*
on biological networks: the case of leukocyte integrin activation network. PLoS One, 2014. **9**(2):
p. e88938.
57. Shannon, P., et al., *Cytoscape: a software environment for integrated models of biomolecular*
interaction networks. Genome research, 2003. **13**(11): p. 2498-2504.
58. Bindea, G., et al., *ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology*
and pathway annotation networks. Bioinformatics, 2009. **25**(8): p. 1091-1093.
59. Zaoli, S., P. Mazzarisi, and F. Lillo, *Betweenness centrality for temporal multiplexes.* Scientific
reports, 2021. **11**(1): p. 4919.
60. Koschützki, D. and F. Schreiber, *Centrality analysis methods for biological networks and their*
application to gene regulatory networks. Gene regulation and systems biology, 2008. **2**: p. GRSB.
S702.
61. Rolinson, G.N., *Effect of-lactam antibiotics on bacterial cell growth rate.* Microbiology, 1980.
120(2): p. 317-323.
62. Lee, A.J., et al., *Robust, linear correlations between growth rates and β-lactam–mediated lysis*
rates. Proceedings of the National Academy of Sciences, 2018. **115**(16): p. 4069-4074.
63. Udekwu, K.I., et al., *Functional relationship between bacterial cell density and the efficacy of*
antibiotics. Journal of antimicrobial chemotherapy, 2009. **63**(4): p. 745-757.
64. Svenningsen, M.S., et al., *Existence of log-phase Escherichia coli persisters and lasting memory of*
a starvation pulse. Life Science Alliance, 2022. **5**(2).
65. Li, Q., et al., *A modified pCas/pTargetF system for CRISPR-Cas9-assisted genome editing in*
Escherichia coli. Acta Biochimica et Biophysica Sinica, 2021. **53**(5): p. 620-627.

66. Balaban, N.Q., et al., *Definitions and guidelines for research on antibiotic persistence.* Nature
Reviews Microbiology, 2019. **17**(7): p. 441-448.
67. Lewis, K., *Persister cells, dormancy and infectious disease.* Nat Rev Microbiol, 2007. **5**(1): p. 48-
56.
68. Pontes, M.H. and E.A. Groisman, *Slow growth determines nonheritable antibiotic resistance in*
Salmonella enterica. Sci Signal, 2019. **12**(592).
69. Hossain, T., et al., *Antibiotic tolerance, persistence, and resistance of the evolved minimal cell,*
Mycoplasma mycoides JCVI-Syn3B. Iscience, 2021. **24**(5): p. 102391.
70. Fridman, O., et al., *Optimization of lag time underlies antibiotic tolerance in evolved bacterial*
populations. Nature, 2014. **513**(7518): p. 418-421.
71. Keren, I., et al., *Persister cells and tolerance to antimicrobials.* FEMS Microbiol Lett, 2004. **230**(1):
p. 13-8.
72. Finkel, S.E., *Long-term survival during stationary phase: evolution and the GASP phenotype.* Nat
Rev Microbiol, 2006. **4**(2): p. 113-20.
73. Finkel, S.E. and R. Kolter, *Evolution of microbial diversity during prolonged starvation.* Proc Natl
Acad Sci U S A, 1999. **96**(7): p. 4023-7.
74. Deter, H.S., et al., *Proteolytic queues at ClpXP increase antibiotic tolerance.* ACS Synthetic
Biology, 2019.
75. Deter, H.S., et al., *A Cell Segmentation/Tracking Tool Based on Machine Learning*, in *Computer*
Optimized Microscopy. 2019, Springer. p. 399-422.
76. Datla, U.S., et al., *The spatiotemporal system dynamics of acquired resistance in an engineered*
microecology. Scientific reports, 2017. **7**(1): p. 1-9.
77. Kearse, M., et al., *Geneious Basic: an integrated and extendable desktop software platform for*
the organization and analysis of sequence data. Bioinformatics, 2012. **28**(12): p. 1647-1649.
78. Deter, H.S., et al., *Mechanisms for Differential Protein Production in Toxin-Antitoxin Systems.*
Toxins (Basel), 2017. **9**(7).
79. Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for*
RNA-seq data with DESeq2. Genome Biol, 2014. **15**(12): p. 550.
80. Heberle, H., et al., *InteractiVenn: a web-based tool for the analysis of sets through Venn*
diagrams. BMC bioinformatics, 2015. **16**: p. 1-7.
81. Szklarczyk, D., et al., *The STRING database in 2023: protein–protein association networks and*
functional enrichment analyses for any sequenced genome of interest. Nucleic acids research,
2023. **51**(D1): p. D638-D646.
82. Doncheva, N.T., et al., *Cytoscape StringApp: network analysis and visualization of proteomics*
data. Journal of proteome research, 2018. **18**(2): p. 623-632.
83. Datla, U.S., et al., *The spatiotemporal system dynamics of acquired resistance in an engineered*
microecology. Scientific Reports, 2017. **7**(1): p. 16071.