FUNCTIONAL CONNECTIVITY, TISSUE MICROSTRUCTURE AND T2 AT 11.1 TESLA

DISTINGUISHES NEUROADAPTIVE DIFFERENCES IN TWO CONTUSIVE BRAIN INJURY

MODELS IN RATS

**Rohan S. Kommireddy[2], **Shray Mehra[2], Marjory Pompilus[2], Rawad Daniel Arja[1], Tian Zhu[1],

Zhihui Yang[1], Yueqiang Fu[1], Jiepei Zhu[1], Firas Kobeissy[1], *Kevin K.W. Wang[1], *Marcelo Febo[2]

(*co-corresponding authors; **equal contributions)

[1]Department of Neurobiology, Center for Neurotrauma, Multiomics & Biomarkers (CNMB),

Neuroscience Institute, Morehouse School of Medicine, Atlanta, GA 30310-1458, USA;

[2] Department of Psychiatry, McKnight Brain Institute, University of Florida, 1149 South Newell

Drive, Gainesville, FL 32611, USA

Short title: Neuroimaging of contusion brain injury rat models

*Co-Corresponding authors: Kevin K.W. Wang, PhD and Marcelo Febo, PhD

Wang Contact information: [1]Department of Neurobiology, Center for Neurotrauma, Multiomics

& Biomarkers (CNMB), Neuroscience Institute, Morehouse School of Medicine, Atlanta, GA

30310-1458; Office: (404) 752 1085; Email: kwang@msm.edu

Febo Contact information: Department of Psychiatry, P.O. Box 100256, Gainesville, Florida

32611; Office: (352) 294 4911; Email: febo@ufl.edu

Acknowledgements: This study was funded by a National Institute on Neurological Disorders

and Stroke grant (UG3 NS106938, PI: Kevin Wang). The contents of this manuscript are solely

the responsibility of the authors and do not necessarily represent the official views of the funding

agencies. This work was performed in the McKnight Brain Institute at the National High

Magnetic Field Laboratory's AMRIS Facility, which is supported by National Science Foundation

Cooperative Agreement No. DMR-1644779 and the State of Florida.

ABSTRACT

The damage caused by contusive traumatic brain injuries (TBIs) is thought to involve breakdown in neuronal communication through focal and diffuse axonal injury along with alterations to the neuronal chemical environment, which adversely affects neuronal networks beyond the injury epicenter(s). In the present study, functional connectivity along with brain tissue microstructure coupled with T2 relaxometry were assessed in two experimental TBI models in rat, controlled cortical impact (CCI) and lateral fluid percussive injury (LFPI). Rats were scanned on an 11.1 Tesla scanner on days 2 and 30 following either CCI or LFPI. Naive controls were scanned once and used as a baseline comparison for both TBI groups. Scanning included functional magnetic resonance imaging (fMRI), diffusion weighted images (DWI), and multi-echo T2 images. fMRI scans were analyzed for functional connectivity across laterally and medially located region of interests (ROIs) across the cortical mantle, hippocampus, and dorsal striatum. DWI scans were processed to generate maps of fractional anisotropy, mean, axial, and radial diffusivities (FA, MD, AD, RD). The analyses focused on cortical and white matter (WM) regions at or near the TBI epicenter. Our results indicate that rats exposed to CCI and LFPI had significantly increased contralateral intra-cortical connectivity at 2 days post-injury. This was observed across similar areas of the cortex in both groups. The increased contralateral connectivity was still observed by day 30 in CCI, but not LFPI rats. Although both CCI and LFPI had changes in WM and cortical FA and diffusivities, WM changes were most predominant in CCI and cortical changes in LFPI. Our results provide support for the use of multimodal MR imaging for different types of contusive and skull-penetrating injury.

INTRODUCTION

Traumatic brain injury (TBI) afflicts millions worldwide, with mortality rates over 30%[1].

Although mild-to-moderate repetitive closed head injuries are the most prevalent among the

different types of TBI, focal contusions and skull penetrating injuries caused by forceful blows,

jolts, bumps to the head, neck, or spinal cord result in devastating outcomes. The long-term,

lifelong sensorimotor, emotional, and cognitive impairments, and permanent disruptions to brain

function and structure, with severe TBI exert a heavy emotional and financial toll on patients and

caregivers. The heterogeneity of damage to the brain with severe TBI makes it difficult to

clinically manage. The pathophysiology of contusive injuries involves mechanistically

heterogeneous and complex cellular and molecular mechanisms extending beyond the primary

site of injury. These so-called secondary injuries can involve aberrant inflammatory

neuroimmune activity, diffuse axonal injury (DAI) and other forms of white matter (WM) damage,

blood-brain barrier (BBB) disruption, microvasculature bleeds, intracranial hematomas, cerebral

and subdural hemorrhage and edema, epileptiform activity, and other adaptive central nervous

system (CNS) pathologies[2]. Clinical TBI treatment and management relies on the earliest

possible interventions to curb the progression of these secondary injuries[3]. Despite the reported

neuroprotective effects of experimental drugs in animal models of TBI, none have proven

effective in clinical trials[4]. This failure can be attributed in part to a lack of data on *in vivo*,

minimally invasive translational CNS markers to bridge preclinical mechanisms to clinical TBI[4].

Functional neuroimaging offers translational data that may help close in on clinically

relevant mechanisms of disease progression and behavioral recovery in TBI, especially when

analyzed together with other quantitative magnetic resonance imaging modalities. For instance,

neuroimaging of a diverse cohort of mild-to-severe TBI identified stronger deactivations of the

default mode network (DMN) regions during cognitive task performance and a negative

relationship between posterior cingulate functional connectivity and mean diffusivity (MD) in

splenium[5]. Similar brain areas were observed to have reduced connectivity following mild TBI[6],

although hyper-connectivity within DMN regions has also been reported[7] and is associated with

metacognitive recovery following moderate to severe TBI[8]. Both increases and decreases in

functional connectivity between limbic, sensorimotor, and association cortical areas have been

observed over extended periods following moderate to severe TBI[9], and in mild concussive

injuries[10]. Individuals sustaining sports-related mild concussions have decreased long-distance

and increased local functional connectivity[11]. Thus, despite heterogeneity in the extent of direct

or secondary damage caused by TBI, consistency can be said to emerge from results showing

temporally evolving functional connectivity reductions and increases across cortical subregions.

Whether or not such changes are specifically restricted to cognitive, motor, sensory, or

association cortices over time remains unclear. Animal studies recapitulating mild concussive

injuries caused by sports injuries or automobile accidents are necessary to model prevalent

forms of TBI in humans[12,13]. Like clinical TBI, however, these produce diffuse and heterogenous

pathology that, while invaluable to determining construct validity of preclinical TBI models, have

so far shown limited ability to reveal fundamental mechanisms involved in cortical reorganization

observed with contusions and penetrating cortical injuries. Preclinical studies using controlled

cortical impact (CCI) and lateral fluid percussive impact (LFPI) models consistently demonstrate

functional reorganization of the cortex near and distant from the TBI epicenter[14]. The large-scale

functional network reorganization continues over 30 to 60 days[15,16] and may involve excitatory-

inhibitory rebalancing following TBI[17]. These cortical changes are likely accompanied by

changes in WM microstructure[18] and chemical changes measurable by tissue T2 relaxometry[19].

The objective of the present study was to assess functional connectivity together with brain

tissue microstructure and T2 relaxometry in CCI and LFPI models to evaluate salient similarities

and differences between these models.

MATERIALS AND METHODS

Subjects

Female and male Sprague-Dawley rats (220-300g) were obtained from Charles River Laboratories (Raleigh, NC, USA.). Rats were housed in sex-matched pairs in a temperature and humidity-controlled vivarium with a 12 h light cycle (lights on at 0700 h) and food and water provided ad libitum. Rats were randomly assigned to one of three experimental conditions: controls (15 male and 8 female rats), CCI (19 male and 12 female rats), and LFPI (7 male and 5 female rats). A subgroup of CCI rats was imaged 2 days post-injury (16 male and 7 female rats) and another group (3 male and 5 female rats) at 30 days. All LFPI rats were imaged 2 days post-injury, and 7 males were scanned once again at 30 days. All procedures received prior approval from the Institutional Animal Care and Use Committee of the University of Florida and followed all applicable NIH guidelines.

Controlled Cortical Impact

The CCI procedure was carried out using a Leica Impact One device (Leica Microsystems Inc). Anesthetic levels were induced with 4% isoflurane gas mixed with 100% oxygen and maintained under 2% isoflurane for the rest of the procedure. The body temperature was regulated to 37°C using a thermal pad while rats were prepared for surgery on a stereotaxic frame. A parasagittal craniectomy (center at anteroposterior [AP], -4.0 mm; L, +2.8 mm from lambda) 5 mm in diameter was performed to expose the brain and allow impactor tip access to the cortical surface. The impactor had a 4-mm flat-face tip. CCI at a depth of 1.5 mm at 4 m/s and a dwell time of 240ms was carried out. All injuries occurred in the right hemisphere. The surgical area was sutured, and recovery was monitored by tail pinch and righting reflexes. Control rats were naïve and received no sham surgical procedures.

Lateral Fluid Percussion Injury

To model diffuse and focal, open-head TBI, and as part of the translational outcomes project in

neurotrauma (TOP-NT), we used the rat LFPI model. LFPI is a well-characterized model of TBI

that captures clinically relevant deficits and pathologies[20-22]. LFPI also applies a localized,

mechanical insult through a parasagittal craniectomy; however, LFPI is administered via a

pressurized fluid pulse (2.0 ±0.2atm). This injury results in a moderated diffuse and focal injury

on the right sensorimotor cortex and expansive and progressing white matter injury[2]. Even in

cases where limited histopathology is observed, LFPI subjects exhibit manifold neurobehavioral

deficits[21,23]. This is consistent with the literature where the majority of TBI cases do not present

overt pathology on imaging assessments, yet patients experience lasting and debilitating

symptoms.

Magnetic Resonance Imaging

Images were collected in an 11.1 Tesla MRI scanner (Magnex Scientific Ltd., Oxford, UK)

with a Resonance Research Inc. gradient set (RRI BFG-240/120-S6, maximum gradient

strength of 1000 mT/m at 325 Amps and a 200 μs risetime; RRI, Billerica, MA) and controlled by

a Bruker Paravision 6.01 console (Bruker BioSpin, Billerica, MA). A custom-made 2.5 cm x 3.5

cm quadrature radiofrequency (RF) surface transmit/receive coil tuned to 470.7MHz (^{1}H

resonance) was used for B1 excitation and signal detection (RF engineering lab, Advanced

Magnetic Resonance Imaging and Spectroscopy Facility, Gainesville, FL). Rats were scanned

under a continuous flow of 1.5 % isoflurane (delivered at 0.1L/min mixed with medical-grade air

containing 70% nitrogen and 30% oxygen). Respiratory rates were monitored continuously, and

body temperature was maintained at 36-37°C using a warm water recirculation system (SA

Instruments, Inc., New York).

For each rat, a high-resolution T2 weighted anatomical scan followed by a functional

magnetic resonance imaging (fMRI) scan, a diffusion weighted imaging (DWI) scan, and a multi-

echo time image series for T2 mapping were acquired. T2-weighted Turbo Rapid Acquisition

with Refocused Echoes (TurboRARE) scans were acquired with the following parameters:

effective echo time (TE) = 37 ms, repetition time (TR) = 5 seconds, RARE factor = 16, number

of averages = 14, field of view (FOV) of 24 mm x 18 mm and 0.9 mm thick slice, and a data

matrix of 256 x 192 and 25 interleaved ascending coronal (axial) slices covering the entire brain

from the rostral-most extent of the anterior frontal cortical surface, caudally towards the upper

brainstem and cerebellum. Functional images were collected using a single-shot spin-echo echo

planar imaging (EPI) sequence with the following parameters: TE = 15 ms, TR = 2 seconds, 300

repetitions, FOV = 24 x 18 mm and 0.9 mm thick slice, and a data matrix of 64 x 48 with 25

interleaved ascending coronal slices in the same space as the T2 anatomical. Respiratory rates,

isoflurane concentration, body temperature, lighting, and room conditions were kept constant

across subjects. Diffusion images used a 4-shot spin echo EPI readout with TE=18ms, TR=4

seconds, 4 averages, FOV = 24 x 18 mm and 0.9 mm slices and a data matrix of 128 x 96 x 25,

30 directions with a b value = 900 s^2/mm and 4 b = 0 scan, pulse duration = 3 ms, and gradient

separation = 8 ms. Multi-slice multi-echo scans were collected with TE=6-60 ms in 6 ms steps

per image volume, TR=2 seconds, 2 averages, FOV = 24 x 18 mm with 0.9 mm slices and a

data matrix of 128 x 128 x 25.

Functional MRI Processing

Image processing was conducted using software tools available in Analysis of Functional

NeuroImages (AFNI)[24], FSL[25], and Advanced Normalization Tools (ANTs)[26] and manual mask

drawing and editing functions in ITKSNAP[27]. Masks outlining the brains of rats on fMRI, DWI,

and T2 anatomical scans were generated using *3dAutomask* in AFNI. These were used for

brain extraction in various steps in the image processing workflows. The processing of fMRI

scans included the following steps: (1) timeseries spike identification and removal using

3dDespike in AFNI, (2) motion correction using *3dvolreg* in AFNI, (3) removal of slow signal

drifts (<0.009Hz) using *3dTproject* in AFNI, (4) identification and suppression of nuisance

signals using *fsl_regfilt* in FSL, (5) low-pass filtering (>0.12Hz) and spatial smoothing (0.6mm

FWHM), and (6) linear affine followed by non-linear registrations of cropped fMRI scans to

cropped anatomical scans using *flirt* in FSL and the *antsRegistrationSyNQuick* script for ANTs.

Anatomical co-registration matrices were applied to fMRI scans to align these to a multi-T2 scan

template upon which ROIs were created. The template was generated by first using rigid-body

and affine co-registration steps and then a final nonlinear symmetric normalization step in ANTs.

A total of 52 bilateral ROIs were manually created in ITKSNAP. These spanned in an axial slice

arrangement the medial and lateral regions of the rostral-to-caudal cortical mantle,

hippocampus and dorsal striatum (**Figure 1A-C**). These ROIs were used for signal extraction for

seed based functional connectivity.

This generated 52 individual ROI text files per subject that contained L2-normalized

resting state signals as a vector of 300 data points. The timeseries files were used in cross-

correlations and in calculations of Pearson r coefficients for every pairwise combinations of

ROIs (using *corrcoef* function in MATLAB). The resulting number of pairwise correlations was

1,274 per subject (after removing 52 self-correlations). Correlation coefficients were Fisher's

transform to ensure normality prior to statistical analyses. The 52 ROIs and template were

visualized in 3D using BrainNet viewer [28]. Weighted undirected matrices derived from the

Pearson correlations were analyzed using Brain Connectivity Toolbox [29] in MATLAB

(Mathworks, Natick, MA). Global graph metrics were calculated for edge density thresholds

ranging from 2-40%. Global network measures for this density range were converted to area

under the curve (AUC) values prior to statistical analyses. Node-specific (ROI) network

measures were calculated at a 12% threshold. The functional network analysis was focused on

measuring node strength (sum of edge weights/node) in subregions of the cortex, hippocampus

and striatum.

Diffusion Weighted Image Processing

ITKSNAP was used to segment contralateral and ipsilateral cortical and white matter ROIs. Diffusion MRI scans were processed using tools available on FSL [30]. DWI scans were first screened for volumes with low signal-to-noise or artefacts. Eddy correction was used for adjusting slight movements during image acquisition and gradient files rotated according to the motion correction vectors. After eddy correction, tensor element reconstruction and estimates of eigenvectors and eigenvalues were performed using weighted least squares regression on *dtifit* in FSL. This last step generated independent images of FA, mean, axial and radial diffusivities (MD, AD, and RD, respectively).

T2 Map processing

T2 parametric maps were created using the MRI processor plugin for ImageJ[31,32]. The simplex minimization method was used for the non-linear fit of multi-TR images to the equation $S_{TE} = S_0 * e^{-TE/T2}$). ITKSNAP was used to segment contralateral and ipsilateral cortical and white matter ROIs, similar to DWI scans. From T2 maps, the mean T2 is calculated per ROI and exported for statistical analysis.

Statistical Analysis

Statistical analyses and data plotting were carried out using GraphPad Prism 9. Functional connectivity data were analyzed using analysis of variance (ANOVA) with repeated measures. A two-stage step-up false discovery rate (FDR) method was used for multiple comparisons correction. DWI scalars and T2 data were analyzed using a two-way ANOVA with Tukey's honest significant difference post hoc test.

RESULTS

Differential Patterns of Recovery of Functional Connectivity in CCI and LFPI

Functional connectivity within and between contra and ipsilateral nodes was evaluated in the medial and lateral areas of the cortex, as well as the hippocampus and striatum, on days 2 and 30 post-injury (ROIs are shown in **Figure 1**). First, contra-to-ipsilateral homotopic connectivity was examined across bilateral ROI representations in the cortex, but no differences were found between controls and TBI rats (data not shown). **Figure 2** highlights the average node strength in 3D connectome maps of control and CCI. The 3D maps illustrate a higher node strength in the contralateral cortex of CCI animals relative to controls. The greater node strength is observed on both days 2 and 30. The 3D maps also show an increase in intra-hemispheric functional connectivity (an increase in the number of node-connecting edges) in the contralateral cortex. A closer evaluation of the CCI group in the heat maps in **Figure 2B** revealed that in the contralateral medial cortex, the rostral and most caudal areas are significantly greater in node strength on days 2 and 30 compared to controls (group x ROI: $F_{102,3468}=2.4$, p<0.0001; **Figure 2B** shows FDR corrected pairwise t-test comparisons). The lateral subregion of the contralateral cortex of CCI rats had significantly greater node strength on both days 2 and 30 post-injury compared to controls but in a limited single slice. In the ipsilateral cortex, the medial subregion had significantly lower node strength on both days 2 and 30 compared to controls (**Figure 2B** shows FDR-corrected pairwise t-test comparisons). This was limited to the more caudal areas of the cortex, within the range of the injury epicenter (**Figure 2B**). The CCI-induced reductions in node strength extended to the lateral areas of the ipsilateral cortex (**Figure 2B**).

Rats that received LFPI had increased node strength in the medial and lateral regions of the contralateral cortex (**Figure 3A**). The increased node strength in the contralateral cortex was observed 2 days, but not 30 days, post-injury (group x ROI: $F_{102,2937}=3.3$, p<0.0001; **Figure 3B** shows FDR corrected pairwise t-test comparisons). The ipsilateral cortex had significant

reductions in node strength, most prominently in the medial cortical areas, although lateral

areas of the posterior region of the cortex also had low node strength in LFPI rats relative to

controls (**Figure 3B** shows FDR corrected pairwise t-test comparisons). The observed

reductions in ipsilateral cortical node strength in LFPI rats were observed on day 2 post-injury

and recovered to control levels in both lateral and medial areas of the cortex by day 30 (**Figure**

3B).

Figure 4 summarizes the functional changes observed in cortical regions of CCI and

LFPI rats. Both CCI and LFPI rats had significant contralateral increases and ipsilateral

decreases in node strength, which we use here as a surrogate marker of functional connectivity.

The observed functional changes were observed in primary sensory areas (e.g.,

somatosensory, auditory, visual) and primary motor cortex on day 2 post-injury in both models

(**Figure 4A** and **4C**). By 30 days after injury, significant decreases in ipsilateral node strength

are still observed near the injury epicenter in CCI rats. No differences were observed in LFPI

rats.

Distinct White Matter and Cortical Diffusion Changes and 30-day Recovery in CCI and

LFPI Models

Figure 5A shows a representative DWI scan collected in a control rat at 11.1 Tesla,

illustrating the slice range for the analyzed scalar maps. We analyzed FA, MD, AD, and RD in

ipsi and contralateral WM and cortex. The ROIs included the cortical TBI epicenter and its

corresponding section of corpus callosum WM tissue. CCI rats had reduced FA in ipsilateral

WM (group x day: $F_{1,94}=29.6$, p<0.0001; post hoc tests: day 2 controls vs day 2 CCI p < 0.0001,

day 2 CCI vs day 30 CCI, p < 0.0001) and contralateral WM (group x day: $F_{1,94}=7.3$, p=0.008;

post hoc tests: day 2 controls vs day 2 CCI p < 0.0001, day 2 CCI vs day 30 CCI, p = 0.005)

and in the ipsilateral cortex on day 2 post-injury (group effect: $F_{1,94}=7.5$, p=0.008; post hoc test:

day 2 controls vs day 2 CCI p = 0.008) (**Figure 5B**). FA returned to control levels in these brain

tissues by day 30 post-CCI. MD in ipsilateral WM was greater in CCI compared to controls on

day 2, but not day 30 (group effect: $F_{1,94}$=11.2, p=0.001; post hoc test: day 2 control vs day 2

CCI, p = 0.01). No differences in MD between control and CCI rats were observed in

contralateral WM, and ipsilateral and contralateral cortex (**Figure 5C**). AD in ipsilateral and

contralateral WM was reduced in CCI rats on day 30 (group x day (ipsi): $F_{1,94}$=13.1, p=0.0005;

post hoc tests: day 2 CCI vs day 30 CCI, p = 0.0001; day 30 control vs day 30 CCI, p = 0.04;

group x day (contra): $F_{1,93}$=7.9, p=0.006; post hoc tests: day 2 CCI vs day 30 CCI, p = 0.004;

day 30 controls vs day 30 CCI, p = 0.009). No differences in AD between controls and CCI rats

were observed in the cortex (**Figure 5D**). RD in ipsilateral WM was reduced in CCI rats on day

30 (group x day: $F_{1,89}$=9.5, p=0.003; post hoc test: day 2 CCI vs day 30 CCI, p = 0.002). No

differences in RD between controls and CCI rats were observed in contralateral WM and cortex

(**Figure 5E**).

Compared to the control groups, LFPI rats had reduced FA in ipsilateral WM on day 2

post-TBI (group x day: F1,75=7.9, p = 0.006; post hoc test: day 2 controls vs day 2 LFPI, p <

0.0001; day 2 LFPI vs day 30 LFPI, p = 0.009) and in the ipsilateral cortex on day 30 (group

effect: $F_{1,75}$=11.7, p=0.001; post hoc test: day 30 controls vs day 30 LFPI, p = 0.01)(**Figure 6A**).

No differences in FA between controls and LFPI rats were observed in contralateral WM and

contralateral cortex (**Figure 6A**). MD increased compared to controls in ipsilateral cortex on day

30 post-TBI (group effect: $F_{1,75}$=14.6, p=0.0003; post hoc test: day 30 controls vs day 30 LFPI, p

= 0.01) and was reduced in contralateral cortex on day 2 post-injury in LFPI rats (group effect:

$F_{1,75}$=12.3, p=0.0008; post hoc test: day 2 control vs day 2 LFPI, p = 0.03) (**Figure 6B**). No

differences in MD between control and LFPI rats were observed in WM ROIs (**Figure 6B**). AD

was lower in contralateral WM (group effect: $F_{1,75}$=9.6, p=0.0003; post hoc tests: day 2 LFPI vs

day 30 LFPI, p = 0.03) and greater in the contralateral cortex on day 30 in LFPI rats compared

to controls (group x day interaction: $F_{1,75}$=34, p<0.0001; post hoc tests: day 2 LFPI vs day 30

LFPI, p < 0.0001; day 30 controls vs day 30 LFPI, p < 0.0001) No differences in AD between

control and LFPI rats were observed in ipsilateral ROIs (**Figure 6C**). RD was lower in

contralateral WM (group effect: $F_{1,72}=16.3$, p = 0.0001; post hoc tests: day 2 controls vs day 2

LFPI, p = 0.04; day 30 controls vs day 30 LFPI, p = 0.02) and contralateral cortex of LFPI

compared to controls on days 2 and 30 (group effect: $F_{1,72}=22.5$, p < 0.0001; post hoc tests: day

2 controls vs day 2 LFPI, p = 0.008; day 30 controls vs day 30 LFPI, p = 0.006). A similar

difference was observed in ipsilateral WM, although post hoc tests were not significant (group

effect: $F_{1,72}=9.4$, p = 0.003) (**Figure 6D**). Compared to the control group, LFPI rats had higher

RD on day 30 post-TBI (group effect: $F_{1,72}=16.1$, p = 0.0001; post hoc tests: day 2 LFPI vs day

30 LFPI, p < 0.0001; day 30 controls vs day 30 LFPI, p < 0.0001).

Distinct White Matter and Cortical T2 Relaxation Changes and 30-day Recovery in CCI

and LFPI Models

Figure 7A shows a representative T2 map collected in a control rat at 11.1 Tesla,

illustrating the slice range for the analysis. We analyzed tissue T2 values (in ms) in ipsilateral

and contralateral WM and cortex. Compared to controls, T2 was increased in ipsilateral WM

(group x day: $F_{1,53}=12.4$, p = 0.0009; post hoc tests: day 2 controls vs day 2 CCI, p < 0.0001;

day 2 CCI vs day 30 CCI, p < 0.0001), contralateral WM (group effect: $F_{1,53}=7.7$, p = 0.008; post

hoc test: day 2 controls vs day 2 CCI, p = 0.007) and in ipsilateral cortex (group x day: $F_{1,53}=6.5$,

p = 0.01; post hoc tests: day 2 controls vs day 2 CCI, p = 0.004; day 2 CCI vs day 30 CCI, p =

0.04) of CCI rats on day 2 post-TBI. T2 values in these brain areas of CCI rats recovered to

control levels on day 30 (**Figure 7B**). No differences in T2 between control and CCI rats were

observed in the contralateral cortex. Compared to controls, T2 values in ipsilateral WM (group

effect: $F_{1,47}=12.4$, p = 0.001; post hoc tests: day 2 controls vs day 2 LFPI, p = 0.03) and

contralateral WM (group effect: $F_{1,47}=5.8$, p = 0.02; post hoc tests: day 2 controls vs day 2 CCI,

p = 0.04) were greater in LFPI rats on day 2. T2 in these WM areas of LFPI rats recovered to

control values by day 30 post-injury (**Figure 7C**). No differences in T2 between control and LFPI

rats were observed in ipsilateral and contralateral cortex (**Figure 7C**).

341   **DISCUSSION**

The results show important similarities and differences in cortical functional connectivity

between CCI and LFPI groups. Both CCI and LFPI rats had significantly increased contralateral

intra-cortical connectivity at 2 days post-injury. This was observed across a similar section of the

cortex in the medial and lateral areas of the cortical mantle. The increased contralateral

connectivity was still observed by day 30 in CCI, but not LFPI rats. While we did not include

additional measures, based on the existing hypothesis suggesting excitatory-inhibitory

rebalancing[14], it is possible that such changes in neuroadaptive processing continue in CCI

animals, and it is established in LFPI. As expected, and reported previously in CCI rats by Yang

et al[15], we observed functional connectivity reductions in the ipsilateral cortex in both groups.

However, once again, we observed that reduced functional connectivity at or near the injury site

is no longer observed by day 30 in LFPI rats, but it endures in both medial and lateral cortical

areas in CCI rats. The cortical areas affected in CCI and LFPI included cross-modal sensory

regions and motor cortex. These results suggest that comparisons between CCI and LFPI might

provide insight into potential cortical neuronal adaptive mechanisms that may be important to

functional recovery. To determine links between cortical functional changes and tissue

microstructure, we assessed DTI-processed DWI scans. Although both CCI and LFPI had

changes in WM and cortical FA and diffusivities, WM changes were most predominant in CCI

and cortical changes in LFPI. This result could be related to the above proposition that LFPI

rats had an appreciable recovery by day 30, possibly in association with a rebalancing of

neuronal activity across the sensorimotor cortex. The hypothesized changes could involve

axonal modifications to compensate for functional losses, as observed in recovery processes in

other injury models[33-35], and this, in turn, can impact axial and radial diffusivities. In the

contralateral cortex of LFPI rats we observed an increased axial diffusivity and reduced radial

diffusivity, which is suggestive of an increase in the directionality of fibers. Finally, T2 values in

ipsilateral- and contralateral WM and cortex were similar in CCI and LFPI rats. The increased

T2 values on day 2 recovered to control levels by day 30. The similarities between ipsilateral

and contralateral regions in both models are interesting. While increased edematous tissue at 2

days could increase T2 at the epicenter, it may be plausible that this also infiltrates the

contralateral region[19]. Histological assessments, particularly 3D whole-brain methods may be

able to provide mechanistic insight.

Changes in cortical excitability have been reported in rats exposed to either CCI or LFPI,

and this has been largely attributed to the development of post-trauma epilepsy (PTE). Both

spontaneous and evoked discharges were reported in layer 5 pyramidal neurons, which

additionally showed increased burst durations during the first 2 weeks post-CCI[36]. Reduced

frequency of spontaneous inhibitory postsynaptic currents (sIPSC), possibly involving reduced

gamma amino butyric acid (GABA) release locally, has also been reported[17]. Increases in

functional connectivity in the contralateral cortex in both LFPI and CCI models may thus

correspond to widespread changes in excitability involved in PTE. However, there is evidence

suggesting the mechanisms may be more nuanced. For example, decreased functional

connectivity in the cortex has been reported in rats also experiencing shorter latencies to seize

and increased interictal epileptic discharges (IEDs)[37]. Besides the role altered neurovascular

coupling is likely to play in both reductions and increases in functional connectivity following TBI,

increased spontaneous firing across multiple neocortical areas should substantially elevate

neuronal oxygen demand and increase blood oxygenation level-dependent (BOLD) signal

fluctuations. Synchronous BOLD activity is expected to increase non-specifically, and in a

widespread manner, in this scenario. Harris and co-workers reported evidence that this

hyperexcitability can be reduced by local inhibitory effects of infused potassium in the

contralateral cortex [38]. They posit that this hyperexcitability may be due to the loss of ipsilateral

cross-callosal excitatory fibers that modulate contralateral inhibitory neurons (contributing to

overexcitability by releasing inhibitory modulation)[38]. The relationship between changes in

neuronal excitability and synchronous BOLD activity is not a linear one. Inhibition of

mitochondrial function reduces both spontaneous neuronal discharges and functional

connectivity, while elevated mitochondrial activity increases neuronal discharges but still

reduces cortical functional connectivity[39]. Our neuroimaging results, and behavioral results

reported by others[40], suggest significant functional recovery over 30 days or more, further

reinforcing the notion of compensatory neuroadaptations that may involve reorganization of

neuronal activity in the cortex of CCI and LFPI rats. PTE-related changes may be accompanied

by changes in functional network topology (perhaps contributing to behavioral recovery)[14,15].

The observed functional connectivity changes can also be attributed to altered

neurovascular coupling[41]. Increased BBB permeability, astrocytosis, synaptic loss, and damage

to microvessels are indicative of a failing neurovascular unit. A result of TBI-induced

neurovascular changes is an alteration in cerebral blood flow (CBF). CBF changes have been

reported early – within hours - post-injury and at 1 year[42,43]. The CBF changes with TBI involve

changes in hypercapnia reactivity, suggesting changes in smooth muscle vasodilatory

responsiveness in contralateral and ipsilateral cortices[44-46]. At 3 h, CBF is reduced globally by

CCI by as much as 85% in ipsilateral and 49% in contralateral cortex[47]. Mild LFPI produced

slight but significant differences in CBF and cerebral blood volume (CBV) between ipsilateral

and contralateral cortex at 12 hours post-injury[42]. On the other hand, at 1-year post-injury, CCI

reduced CBF at the TBI epicenter, including part of the hippocampus. However, no differences

in CBF were observed in contralateral cortex[43]. This latter result suggests a certain degree of

neurovascular recovery takes place in off-epicenter sites of the cortex, particularly in the

contralateral cortex. Thus, functional connectivity changes in these off-TBI regions are likely to

involve neuronal activity changes and are not entirely due to altered CBF mechanisms.

Our DTI analysis focused on corpus callosum segments circumventing the TBI epicenter

and in the contralateral cortex. We also assessed the cortical epicenter area and corresponding

contralateral cortex to evaluate tissue microstructural differences at 2 and 30 days. Based on

the literature, the reported changes in DTI scalars, either using a similar ROI analysis or whole-

brain statistics, appear highly heterogeneous across laboratories. Using the CCI model, Harris

and colleagues reported ipsilateral WM FA reductions and corresponding diffusivity changes at

1 week, which persisted through 4 weeks[18]. The results were supported by histological data

showing demyelinated fibers[18]. This contrasts with our present results using an a priori ROI

approach and the difference is likely due to their more comprehensive WM tract-based statistics

approach. Changes in WM can surpass localized changes near the epicenter, particularly at

later stages post-injury[18]. Conversely, CCI-induced reductions in FA in callosal WM recovered to

control levels by day 28 post-TBI, but MD persisted to this time point[48]. Others have reported

reductions in FA that were observed 6 months after CCI exposure in rats[16]. This was

accompanied by decreases in AD and increased RD[16]. The observed reductions in FA (and

increased MD) are associated with significant increases in activated microglia marker

translocator protein 18kDa (TSPO), *in vivo*[49]. Yet another study using CCI model reports

reductions in FA and increases in MD at 2 days but not at 3 h or 7 days or later[50]. While these

results call for improvements in the standardization of DWI methods to facilitate cross-laboratory

comparisons, they also suggest the possibility that the neuroadaptations in WM carry large

inter-subject (and perhaps, intra-subject) variability. The cellular and molecular changes guiding

WM restructuring and axonal remodeling following TBI are likely to involve a certain degree of

spatial and temporal 'randomness', even with focal contusion models, that could make it difficult

to consistently measure the same results using DWI/DTI across and within research

laboratories. Our results also indicate increases in T2 in ipsilateral and contralateral cortex and

WM at day 2 that is no longer observed on day 30. CCI-induced increased T2 correlates with

tissue edema and this in turn has been linked with increases in aquaporin-4 in cortex[19]. The

temporal evolution of changes in T2 varies across laboratories as well. Recovery of T2 to

control levels in the perilesional or contralateral cortex has been reported at 7 days[50], 14 days[45],

and 30 days (present study), or not recovery by a 42nd day of imaging post-CCI[19]. Manninen et

al report increases in T2 at day 2 post-FPI and reductions in T2 compared to naïve controls by

day 7, with the same effect observed again on day 21[51]. The latter could be a sign of permanent

tissue damage at later time points.

In conclusion, our results provide additional support for the use of multimodal MR

imaging to determine the *in vivo*, progressive changes in function and tissue microstructure

following contusive and skull penetrating injury. Both fMRI and DWI are potentially capable of

distinguishing pathological changes between LFPI and CCI models. However, comparisons with

the literature strongly suggest the need for methodological standardization, data harmonization,

and statistical approaches to facilitate cross-laboratory comparisons, feature extraction, and/or

classifications with enough information to evaluate potentially important preclinical findings

relevant to clinical TBI[52,53].

REFERENCES

1.   Demlie TA, Alemu MT, Messelu MA, et al. Incidence and predictors of mortality among

traumatic brain injury patients admitted to Amhara region Comprehensive Specialized Hospitals,

northwest Ethiopia, 2022. BMC Emerg Med 2023;23(1):55, doi:10.1186/s12873-023-00823-9

2.   Bramlett HM, Dietrich WD. Long-Term Consequences of Traumatic Brain Injury: Current

Status of Potential Mechanisms of Injury and Neurological Outcomes. J Neurotrauma

2015;32(23):1834-48, doi:10.1089/neu.2014.3352

3.   Vella MA, Crandall ML, Patel MB. Acute Management of Traumatic Brain Injury. Surg

Clin North Am 2017;97(5):1015-1030, doi:10.1016/j.suc.2017.06.003

4.   Wang KK, Yang Z, Zhu T, et al. An update on diagnostic and prognostic biomarkers for

traumatic brain injury. Expert Rev Mol Diagn 2018;18(2):165-180,

doi:10.1080/14737159.2018.1428089

5.   Sharp DJ, Beckmann CF, Greenwood R, et al. Default mode network functional and

structural connectivity after traumatic brain injury. Brain 2011;134(Pt 8):2233-47,

doi:10.1093/brain/awr175

6.   Alhourani A, Wozny TA, Krishnaswamy D, et al. Magnetoencephalography-based

identification of functional connectivity network disruption following mild traumatic brain injury. J

Neurophysiol 2016;116(4):1840-1847, doi:10.1152/jn.00513.2016

7.   Bernier RA, Roy A, Venkatesan UM, et al. Dedifferentiation Does Not Account for

Hyperconnectivity after Traumatic Brain Injury. Front Neurol 2017;8(297,

doi:10.3389/fneur.2017.00297

8.   Grossner EC, Bernier RA, Brenner EK, et al. Enhanced default mode connectivity

predicts metacognitive accuracy in traumatic brain injury. Neuropsychology 2019;33(7):922-933,

doi:10.1037/neu0000559

9.   Hillary FG, Slocomb J, Hills EC, et al. Changes in resting connectivity during recovery

from severe traumatic brain injury. Int J Psychophysiol 2011;82(1):115-23,

doi:10.1016/j.ijpsycho.2011.03.011

10.   Virji-Babul N, Hilderman CG, Makan N, et al. Changes in functional brain networks

following sports-related concussion in adolescents. J Neurotrauma 2014;31(23):1914-9,

doi:10.1089/neu.2014.3450

11.   Cao C, Slobounov S. Alteration of cortical functional connectivity as a result of traumatic

brain injury revealed by graph theory, ICA, and sLORETA analyses of EEG signals. IEEE Trans

Neural Syst Rehabil Eng 2010;18(1):11-9, doi:10.1109/TNSRE.2009.2027704

12.   Sakthivel R, Criado-Marrero M, Barroso D, et al. Fixed Time-Point Analysis Reveals

Repetitive Mild Traumatic Brain Injury Effects on Resting State Functional Magnetic Resonance

Imaging Connectivity and Neuro-Spatial Protein Profiles. J Neurotrauma 2023;40(19-20):2037-

2049, doi:10.1089/neu.2022.0464

13. Kulkarni P, Morrison TR, Cai X, et al. Neuroradiological Changes Following Single or

Repetitive Mild TBI. Front Syst Neurosci 2019;13(34, doi:10.3389/fnsys.2019.00034

14. Harris NG, Verley DR, Gutman BA, et al. Disconnection and hyper-connectivity underlie

reorganization after TBI: A rodent functional connectomic analysis. Exp Neurol 2016;277(124-

138, doi:10.1016/j.expneurol.2015.12.020

15. Yang Z, Zhu T, Pompilus M, et al. Compensatory functional connectome changes in a

rat model of traumatic brain injury. Brain Commun 2021;3(4):fcab244,

doi:10.1093/braincomms/fcab244

16. Mohamed AZ, Cumming P, Nasrallah FA. Traumatic brain injury augurs ill for prolonged

deficits in the brain's structural and functional integrity following controlled cortical impact injury.

Sci Rep 2021;11(1):21559, doi:10.1038/s41598-021-00660-5

17. Yang L, Benardo LS, Valsamis H, et al. Acute injury to superficial cortex leads to a

decrease in synaptic inhibition and increase in excitation in neocortical layer V pyramidal cells. J

Neurophysiol 2007;97(1):178-87, doi:10.1152/jn.01374.2005

18. Harris NG, Verley DR, Gutman BA, et al. Bi-directional changes in fractional anisotropy

after experiment TBI: Disorganization and reorganization? Neuroimage 2016;133(129-143,

doi:10.1016/j.neuroimage.2016.03.012

19. Zheng T, Du J, Yuan Y, et al. Effect of Low Intensity Transcranial Ultrasound (LITUS) on

Post-traumatic Brain Edema in Rats: Evaluation by Isotropic 3-Dimensional T2 and Multi-TE T2

Weighted MRI. Front Neurol 2020;11(578638, doi:10.3389/fneur.2020.578638

20. Alder J, Fujioka W, Lifshitz J, et al. Lateral fluid percussion: model of traumatic brain

injury in mice. J Vis Exp 2011;54), doi:10.3791/3063

21.   Lifshitz J, Rowe RK, Griffiths DR, et al. Clinical relevance of midline fluid percussion

brain injury: Acute deficits, chronic morbidities and the utility of biomarkers. Brain Inj

2016;30(11):1293-1301, doi:10.1080/02699052.2016.1193628

22.   Radabaugh H, Bonnell J, Schwartz O, et al. Use of Machine Learning to Re-Assess

Patterns of Multivariate Functional Recovery after Fluid Percussion Injury: Operation Brain

Trauma Therapy. J Neurotrauma 2021;38(12):1670-1678, doi:10.1089/neu.2020.7357

23.   Kochanek PM, Bramlett HM, Dixon CE, et al. Operation Brain Trauma Therapy: 2016

Update. Mil Med 2018;183(suppl_1):303-312, doi:10.1093/milmed/usx184

24.   Cox RW. AFNI: software for analysis and visualization of functional magnetic resonance

neuroimages. Comput Biomed Res 1996;29(3):162-73, doi:10.1006/cbmr.1996.0014

25.   Jenkinson M, Bannister P, Brady M, et al. Improved optimization for the robust and

accurate linear registration and motion correction of brain images. Neuroimage 2002;17(2):825-

41, doi:10.1016/s1053-8119(02)91132-8

26.   Klein A, Andersson J, Ardekani BA, et al. Evaluation of 14 nonlinear deformation

algorithms applied to human brain MRI registration. Neuroimage 2009;46(3):786-802,

doi:10.1016/j.neuroimage.2008.12.037

27.   Yushkevich PA, Piven J, Hazlett HC, et al. User-guided 3D active contour segmentation

of anatomical structures: significantly improved efficiency and reliability. Neuroimage

2006;31(3):1116-28, doi:10.1016/j.neuroimage.2006.01.015

28.   Xia M, Wang J, He Y. BrainNet Viewer: a network visualization tool for human brain

connectomics. PLoS One 2013;8(7):e68910, doi:10.1371/journal.pone.0068910

29.   Rubinov M, Sporns O. Complex network measures of brain connectivity: uses and

interpretations. Neuroimage 2010;52(3):1059-69, doi:10.1016/j.neuroimage.2009.10.003

30.   Smith SM, Jenkinson M, Woolrich MW, et al. Advances in functional and structural MR

image analysis and implementation as FSL. Neuroimage 2004;23 Suppl 1(S208-19,

doi:10.1016/j.neuroimage.2004.07.051

31.   Schmidt KF, Ziu M, Schmidt NO, et al. Volume reconstruction techniques improve the

correlation between histological and in vivo tumor volume measurements in mouse models of

human gliomas. J Neurooncol 2004;68(3):207-15, doi:10.1023/b:neon.0000033364.43142.bf

32.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image

analysis. Nat Methods 2012;9(7):671-5, doi:10.1038/nmeth.2089

33.   Recanzone GH, Merzenich MM, Jenkins WM, et al. Topographic reorganization of the

hand representation in cortical area 3b owl monkeys trained in a frequency-discrimination task.

J Neurophysiol 1992;67(5):1031-56, doi:10.1152/jn.1992.67.5.1031

34.   Xerri C, Coq JO, Merzenich MM, et al. Experience-induced plasticity of cutaneous maps

in the primary somatosensory cortex of adult monkeys and rats. J Physiol Paris 1996;90(3-

4):277-87, doi:10.1016/s0928-4257(97)81438-6

35.   Xerri C, Merzenich MM, Peterson BE, et al. Plasticity of primary somatosensory cortex

paralleling sensorimotor skill recovery from stroke in adult monkeys. J Neurophysiol

1998;79(4):2119-48, doi:10.1152/jn.1998.79.4.2119

36.   Yang L, Afroz S, Michelson HB, et al. Spontaneous epileptiform activity in rat neocortex

after controlled cortical impact injury. J Neurotrauma 2010;27(8):1541-8,

doi:10.1089/neu.2009.1244

37.   Mishra AM, Bai X, Sanganahalli BG, et al. Decreased resting functional connectivity after

traumatic brain injury in the rat. PLoS One 2014;9(4):e95280, doi:10.1371/journal.pone.0095280

38.   Verley DR, Torolira D, Pulido B, et al. Remote Changes in Cortical Excitability after

Experimental Traumatic Brain Injury and Functional Reorganization. J Neurotrauma

2018;35(20):2448-2461, doi:10.1089/neu.2017.5536

39.   Sanganahalli BG, Herman P, Hyder F, et al. Mitochondrial functional state impacts

spontaneous neocortical activity and resting state FMRI. PLoS One 2013;8(5):e63317,

doi:10.1371/journal.pone.0063317

40.   Andrews BT, Barbay S, Townsend J, et al. Unrepaired decompressive craniectomy

worsens motor performance in a rat traumatic brain injury model. Sci Rep 2020;10(1):22242,

doi:10.1038/s41598-020-79155-8

41.   Wu J, He J, Tian X, et al. Activation of the Hedgehog Pathway Promotes Recovery of

Neurological Function After Traumatic Brain Injury by Protecting the Neurovascular Unit. Transl

Stroke Res 2020;11(4):720-733, doi:10.1007/s12975-019-00771-2

42.   Schneider G, Fries P, Wagner-Jochem D, et al. Pathophysiological changes after

traumatic brain injury: comparison of two experimental animal models by means of MRI.

MAGMA 2002;14(3):233-41, doi:10.1007/BF02668217

43.   Kochanek PM, Hendrich KS, Dixon CE, et al. Cerebral blood flow at one year after

controlled cortical impact in rats: assessment by magnetic resonance imaging. J Neurotrauma

2002;19(9):1029-37, doi:10.1089/089771502760341947

44.   Forbes ML, Hendrich KS, Kochanek PM, et al. Assessment of cerebral blood flow and

CO_2 reactivity after controlled cortical impact by perfusion magnetic resonance imaging using

arterial spin-labeling in rats. J Cereb Blood Flow Metab 1997;17(8):865-74,

doi:10.1097/00004647-199708000-00005

45.   Long JA, Watts LT, Li W, et al. The effects of perturbed cerebral blood flow and

cerebrovascular reactivity on structural MRI and behavioral readouts in mild traumatic brain

injury. J Cereb Blood Flow Metab 2015;35(11):1852-61, doi:10.1038/jcbfm.2015.143

46.   Zhang F, Sprague SM, Farrokhi F, et al. Reversal of attenuation of cerebrovascular

reactivity to hypercapnia by a nitric oxide donor after controlled cortical impact in a rat model of

traumatic brain injury. J Neurosurg 2002;97(4):963-9, doi:10.3171/jns.2002.97.4.0963

47.   Hendrich KS, Kochanek PM, Williams DS, et al. Early perfusion after controlled cortical

impact in rats: quantification by arterial spin-labeled MRI and the influence of spin-lattice

relaxation time heterogeneity. Magn Reson Med 1999;42(4):673-81, doi:10.1002/(sici)1522-

2594(199910)42:4<673::aid-mrm8>3.0.co;2-b

48.   Yu M, Wang M, Yang D, et al. Dynamics of blood brain barrier permeability and tissue

microstructure following controlled cortical impact injury in rat: A dynamic contrast-enhanced

magnetic resonance imaging and diffusion kurtosis imaging study. Magn Reson Imaging

2019;62(1-9, doi:10.1016/j.mri.2019.01.017

49.   Missault S, Anckaerts C, Blockx I, et al. Neuroimaging of Subacute Brain Inflammation

and Microstructural Changes Predicts Long-Term Functional Outcome after Experimental

Traumatic Brain Injury. J Neurotrauma 2019;36(5):768-788, doi:10.1089/neu.2018.5704

50.   Li W, Long JA, Watts L, et al. Spatiotemporal changes in diffusion, T2 and susceptibility

of white matter following mild traumatic brain injury. NMR Biomed 2016;29(7):896-903,

doi:10.1002/nbm.3536

51.   Manninen E, Chary K, Lapinlampi N, et al. Acute thalamic damage as a prognostic

biomarker for post-traumatic epileptogenesis. Epilepsia 2021;62(8):1852-1864,

doi:10.1111/epi.16986

52.   Bhagavatula S, Cabeen R, Harris NG, et al. Image data harmonization tools for the

analysis of post-traumatic epilepsy development in preclinical multisite MRI studies. Epilepsy

Res 2023;195(107201, doi:10.1016/j.eplepsyres.2023.107201

53.   Nielson JL, Paquette J, Liu AW, et al. Topological data analysis for discovery in

preclinical spinal cord injury and traumatic brain injury. Nat Commun 2015;6(8581,

doi:10.1038/ncomms9581

FIGURES

Figure 1. Regions of interest were manually created over a multi-subject template to extract fMRI signals for connectivity analysis. A) Coronal view of the distribution of bilateral cortical, hippocampal and striatal ROIs. B) 3D renditions of the ROIs to illustrate the sampling. C) Nomenclature used to identify ROIs individually across the cortical mantle, medially and laterally in each hemisphere, and in subcortical regions. Abbreviations: C-MC, contralateral-medial cortex; Ip-MC, ipsilateral-medial cortex; C-LC, contralateral-lateral cortex; Ip-LC, ipsilateral-lateral cortex; C-Str, contralateral striatum; Ip-Str, ipsilateral striatum; C-Hpc, contralateral hippocampus; Ip-Hpc, ipsilateral hippocampus.

Figure 2. Controlled cortical impact (CCI) increased node strength in contralateral cortex and reduced it in ipsilateral cortex. A) group averaged 3D functional connectome maps illustrate contralateral increases in node strength on days 2 and 30 post-TBI and reductions in ipsilateral cortex. Node size is representative of the calculated ROI node strength and edges (connecting lines) are Fisher's z-transformed Pearson correlation coefficient (maps threshold is 12% edge density; scale bar besides each map indicates z range). B) Heat maps of node strength across the sampled ROIs. Left y axis shows ROI name and x axis group. Asterisks on the right indicate significant difference between CCI day 2 vs controls* and CCI day 30 vs controls (FDR corrected $p<0.05$). The scale bar to the left indicates averaged node strength value.

Figure 3. Lateral fluid percussion injury (LFPI) increased node strength in contralateral cortex and reduced it in ipsilateral cortex. A) group averaged 3D functional connectome maps illustrate contralateral increases in node strength on days 2 post-TBI and reductions in ipsilateral cortex. Node size is representative of the calculated ROI node strength and edges (connecting lines) are Fisher's z-transformed Pearson correlation coefficient (maps threshold is 12% edge density; scale bar besides each map indicates z range). B) Heat maps of node strength across the sampled ROIs. Left y axis shows ROI name and x axis group. Asterisks on the right indicate significant difference between CCI day 2 vs controls* and CCI day 30 vs CCI day 2 (FDR corrected p<0.05). The scale bar to the left indicates averaged node strength value.

Figure 4. Cortical brain areas affected by CCI vs LFPI. The brain regions were identified by overlaying group averaged ROI's showing significant differences between either CCI or LFPI and controls onto a rat brain template and parcellation[13]. A) Regions with increases and decreases in node strength on day 2 following CCI include sensorimotor areas in both left and right hemispheres. B) By day 30 after CCI, parietal and visual areas shown decreases in ipsilateral cortex. C) Regions affected by LFPI include many of the same cortical areas as CCI on day 2. D) Functional connectivity differences are no longer measured on day 30 following LFPI.

Figure 5. White matter (WM) and cortical tissue diffusion changes at 2 and 30 days following CCI. A) Representative diffusion tensor maps of a control animal. Shown are a B0 image, fractional anisotropy (FA) grey scale map and FA with red green blue (RGB) color coded overlay of 1st eigenvector map, and mean, axial and radial diffusivities (MD, AD, and RD, respectively). B-E) FA, MD, AD, and RD in ipsi and contralateral WM and cortex. Data are shown as box and whisker plots. Lines and asterisks indicate significant differences at p<0.05 (Tukey's multiple comparison post hoc test following two-way ANOVA).

Figure 6. White matter (WM) and cortical tissue diffusion changes at 2 and 30 days following LFPI. A-D) FA, MD, AD, and RD in ipsi and contralateral WM and cortex. Data are shown as box and whisker plots. Lines and asterisks indicate significant differences at p<0.05 (Tukey's multiple comparison post hoc test following two-way ANOVA).

Figure 7. White matter (WM) and cortical tissue T2 relaxation at 2 and 30 days following CCI or LFPI. A) Representative T2 maps of a control animal. A-B) T2 (in ms) in ipsi and contralateral WM and cortex of control, CCI and LFPI exposed rats on days 2 and 30 post-TBI. Data are shown as box and whisker plots. Lines and asterisks indicate significant differences at p<0.05 (Tukey's multiple comparison post hoc test following two-way ANOVA).