GOAT: efficient and robust identification of geneset enrichment

Frank Koopmans

Department of Molecular and Cellular Neurobiology, Center for Neurogenomics and Cognitive

Research, Amsterdam Neuroscience, VU University, 1081 HV Amsterdam, The Netherlands.

ABSTRACT

Geneset enrichment analysis is foundational to the interpretation of high throughput biology. We here

present GOAT (https://github.com/ftwkoopmans/goat), a parameter-free permutation-based algorithm

for geneset enrichment analysis of preranked genelists. Estimated geneset p-values are well calibrated

under the null hypothesis and invariant to geneset size. GOAT consistently identifies more Gene

Ontology terms in real-world datasets than current methods and is available as an R package and online

tool.

MAIN TEXT

High throughput studies of biological systems, using experimental assays such as proteomics and RNA-

sequencing, are widely used to identify differentially expressed genes (DEGs) across experimental

conditions. To perform systematic and unbiased interpretation of the genes associated with each

experimental condition, researchers commonly employ geneset enrichment analysis (also referred to as

geneset analysis) to identify which biological processes and (sub)cellular components in the Gene

Ontology (GO) database are overrepresented among these DEGs[1-4].

The vast majority of GO tools employ the Over-Representation Analysis (ORA) approach, subjecting each

geneset (GO term) to a Fisher's exact test using a contingency table constructed from the significant and

non-significant genes in the user's genelist[5-8]. Main disadvantages are; 1) Information on gene

scores/ranks is lost when partitioning the genelist into the foreground set that contains significant genes

and the background set. 2) This requires an arbitrary cutoff for significance and this parameter impacts

the ORA results. 3) ORA critically depends on using the correct background list but mistakes therein are

easily made[9].

The GSEA algorithm[10, 11] is a commonly used permutation based approach for geneset enrichment

analysis that can be applied to a preranked genelist, i.e. a table of gene identifiers and respective

effectsizes or p-values estimated by a statistical model. While GSEA is more sensitive than ORA,

permutation based algorithms take significantly more computation time and require highly optimized

implementations that are not easily ported to other programming languages and webtools, which

hampers their adoption.

We here present GOAT, the Geneset Ordinal Association Test, a novel algorithm for geneset enrichment

testing in preranked genelists that uses bootstrapping to generate competitive null hypotheses. GOAT

uses squared gene rank values (of p-values or effectsizes) as gene scores to boost top-ranked genes in

the input genelist, which are considered most important when interpreting study outcomes in biological

context (Methods). Geneset scores are defined as the mean of their respective gene scores. With this

approach, gene score distributions are right-skewed (long tail for high gene scores) and so are the

empirical geneset null distributions for small genesets. Following the central limit theorem, this

converges to a normal distribution for large genesets (Figure S1). Empirical null distribution for each

geneset size (number of assigned genes) are estimated through extensive bootstrapping procedures and

a skewed normal distribution is fitted to each (Methods). This concludes the first step; for a given

genelist of N genes we obtained the parameters of a skew-normal that represent the null. In the second

step, we compute geneset p-values by testing their score against the respective precomputed null

distribution. We have precomputed null distributions for all genelists of length 100-20 000 and all

genesets sizes, so a typical application of GOAT only requires the second part of the algorithm which

completes in one second for any genelist (Figure S2).

We first validated that geneset p-values estimated by GOAT are accurate under the null hypothesis, i.e.

no surprisingly strong p-values should be found when analyzing random genelists[9, 12]. We generated

synthetic genelists of 500, 2000, 6000 and 10 000 genes and applied the GOAT algorithm to test for

enrichment across 200 000 randomly generated genesets of sizes 10, 20, 50, 100, 200 and 1000.

Expecting geneset enrichment algorithms to yield uniform geneset p-values when testing against many

random genesets, this allowed us to check for potential bias in small or large genesets. Benchmarking

results confirmed that geneset p-values estimated by GOAT accurately match expected values regardless

of genelist length or geneset size (Figure 1).

Next, we applied similar analyses to GSEA (fGSEA implementation) and found inaccuracies under default

settings that could be rescued by greatly increasing the number of permutations that the algorithm

performs (Figure S3).

After confirming that GOAT is accurate, we applied it to various real-world datasets to compare its

sensitivity to GSEA (fGSEA implementation) and ORA and found that GOAT identified significantly more

GO terms (Figure 2). Using a genelist with effectsizes as input allows GOAT and GSEA to evaluate if there

is enrichment in either gene up- or down-regulation. In contrast, using a genelist with p-values will

perform a one-dimensional test of enrichment per geneset (there is no information on up/down-

regulation) and this lead to a strong reduction in significant genesets as also observed in previous

studies[13].

Using gene effectsizes as input, GOAT identified 27~272% more genesets across all datasets compared to

GSEA. With gene p-values as input, GOAT outperformed GSEA by 39~254% except for the Higginbotham

et al. dataset where GSEA and GOAT identified 3 and 2 genesets in total, respectively. Comparing against

ORA, with gene p-values as input for GOAT for fair comparison (ORA uses a one-dimensional input

genelist), GOAT identified 33~1433% more genesets except for the Higginbotham et al. dataset where

ORA and GOAT identified 7 and 2 genesets in total, respectively.

When comparing geneset p-values between GOAT, ORA and GSEA, we found the same trend of

enrichment but with generally stronger p-values for GOAT which strengthens our confidence that the

GOAT approach does not yield unrelated genesets but instead exhibits both correlation to existing

methods and increased power (Figure S4).

To evaluate the performance of GOAT in the analysis of small genelists, we applied MS-DAP[14] to obtain

differential expression statistics for 378 proteins from Shisa6 wildtype *versus* knockout

immunoprecipitation[15]. When testing only for down-regulation, GOAT identified 7 synaptic CC terms,

GSEA 1 and ORA 2. Using the SynGO synaptic knowledgebase[16] instead of GO, GOAT identified 9 CC

terms, compared to 6 for GSEA and 5 for ORA. GOAT identified CC and BP terms related to postsynaptic

membranes and regulation of neurotransmitters, as expected from the known function of Shisa6[15].

While gene sampling-based algorithms for geneset enrichment analysis are not new[4], GOAT takes a next

step by combining; 1) a gene score metric with a non-linear increase in importance for top-ranking

genes, 2) efficient and accurate modeling of null distributions as skew-normal (instead of relying on

nonparametric tests) and 3) a caching strategy that allows us to precompute the null distributions (the

part of the algorithm with highest computational complexity), thereby reducing the application of GOAT

to a simple test that completes in one second even when a large genelist of 10 000 genes is tested

against the GO database.

Taken together, the GOAT algorithm is fast, portable, parameter-free, robust and more sensitive than

existing methods for geneset analysis. Two implementations of GOAT are made available to serve a wide

audience; an R package for use in bioinformatic workflows and a user-friendly online tool.

Figure 1. GOAT p-values are accurate under the null hypothesis in simulations that generate genelists

and genesets of various sizes.

Random genelists of 500, 2000, 6000 and 10 000 genes (N) were generated to check for potential bias in

genelist size (rows). 200 000 Synthetic genesets of 10, 20, 50, 100, 200 or 1000 genes were generated by

random sampling of k genes from the respective genelist to check for potential bias in geneset size (line

colors). The y-axis shows the observed geneset p-value for a given p-value threshold on the x-axis (both

on -log10 scale, p-values are shown as-is without multiple testing correction). Since randomized genesets

were used in these simulations, expected values are on the diagonal (dashed line).

Figure 2. GOAT identifies more significant GO terms compared to GSEA and ORA across 6 omics

studies.

Differential expression analysis results from each study (panels) were subjected to 5 methods (colors) to

identify enriched GO terms. GO MF, CC and BP represent the respective Gene Ontology domains

Molecular Functions, Cellular Components and Biological Processes. The x-axis shows the number of

significant genesets identified by each method after Bonferroni adjustment.

REFERENCES

1.   Maciejewski, H. Gene set analysis methods: statistical models and methodological differences.
Brief Bioinform **15**, 504-518 (2014).
2.   Nam, D. & Kim, S.Y. Gene-set approach for expression pattern analysis. *Brief Bioinform* **9**, 189-
197 (2008).
3.   Hung, J.H., Yang, T.H., Hu, Z., Weng, Z. & DeLisi, C. Gene set enrichment analysis: performance
evaluation and usage guidelines. *Brief Bioinform* **13**, 281-291 (2012).
4.   Maleki, F., Ovens, K., Hogan, D.J. & Kusalik, A.J. Gene Set Analysis: Challenges, Opportunities, and
Future Research. *Front Genet* **11**, 654 (2020).
5.   Huang da, W., Sherman, B.T. & Lempicki, R.A. Systematic and integrative analysis of large gene
lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57 (2009).
6.   Kuleshov, M.V. et al. Enrichr: a comprehensive gene set enrichment analysis web server 2016
update. *Nucleic Acids Res* **44**, W90-97 (2016).
7.   Mi, H., Poudel, S., Muruganujan, A., Casagrande, J.T. & Thomas, P.D. PANTHER version 10:
expanded protein families and functions, and analysis tools. *Nucleic Acids Res* **44**, D336-342
(2016).
8.   Kolberg, L. et al. g:Profiler-interoperable web service for functional enrichment analysis and gene
identifier mapping (2023 update). *Nucleic Acids Res* **51**, W207-W212 (2023).
9.   Wijesooriya, K., Jadaan, S.A., Perera, K.L., Kaur, T. & Ziemann, M. Urgent need for consistent
standards in functional enrichment analysis. *PLoS Comput Biol* **18**, e1009935 (2022).
10.   Subramanian, A. et al. Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550
(2005).
11.   Korotkevich, G. et al. Fast gene set enrichment analysis. *bioRxiv*, 060012 (2021).
12.   Tamayo, P., Steinhardt, G., Liberzon, A. & Mesirov, J.P. The limitations of simple gene set
enrichment analysis assuming gene independence. *Stat Methods Med Res* **25**, 472-487 (2016).
13.   Hong, G., Zhang, W., Li, H., Shen, X. & Guo, Z. Separate enrichment analysis of pathways for up-
and downregulated genes. *J R Soc Interface* **11**, 20130950 (2014).
14.   Koopmans, F., Li, K.W., Klaassen, R.V. & Smit, A.B. MS-DAP Platform for Downstream Data
Analysis of Label-Free Proteomics Uncovers Optimal Workflows in Benchmark Data Sets and
Increased Sensitivity in Analysis of Alzheimer's Biomarker Data. *J Proteome Res* **22**, 374-386
(2023).
15.   Klaassen, R.V. et al. Shisa6 traps AMPA receptors at postsynaptic sites and prevents their
desensitization during synaptic activity. *Nat Commun* **7**, 10682 (2016).
16.   Koopmans, F. et al. SynGO: An Evidence-Based, Expert-Curated Knowledge Base for the Synapse.
Neuron **103**, 217-234 e214 (2019).

METHODS

GOAT algorithm

Test statistics in the input genelist are transformed to gene scores as follows; A) if the user provides p-

values for each gene, the score for each gene is defined as *rank(-pvalue)^2*. B) if effectsizes are provided

instead, gene scores are defined as *rank(effectsize)^2* to test genesets for enrichment of up-regulated

genes (where higher effectsize is better). Analogously, we can generate gene scores to test genesets for

enrichment of down-regulated or absolute effectsizes. Rank transformation increases the robustness of

the user-provided input gene scores and ensures that downstream geneset scoring and computation of

null distributions always operates on a similar type of gene score distribution, regardless of the input

genelist p-value/effectsize distribution. Specifically, we use squared ranks instead of plain rank values to

attribute more weight (importance) to the genes with strongest p-values (or effectsizes) when scoring

genesets.

Geneset scores are defined as the mean score of respective genes; for each user-provided geneset we

iterate all assigned genes, lookup the corresponding gene scores in the input genelist and compute their

mean value.

Competitive null distributions are generated for genesets of each size (N, the number of genes) by

bootstrapping 500 000 random genesets; randomly select N genes from the input genelist, compute the

geneset score (mean of respective gene scores), store this random geneset score, repeat. Because gene

scores were computed as squared rank values, the gene score distribution is skewed with a long tail to

high scores (Figure S1a). Following the central limit theorem[1], the null distributions generated by taking

the mean value of N gene scores 500 000 times over will yield normal distributions when N is large

(Figure S1b). However, when N is small (e.g. 10 genes) the respective null distribution yields a skewed

normal distribution (Figure S1c). Thus, we model the expected gene scores under the null hypothesis by

fitting a skew-normal distribution to the set of 500 000 random gene scores that represent expected

values for a geneset of size N.

Because the rank-transformed gene score distribution is always the same for a genelist of the same size,

so are the respective null distributions at each geneset size N. Hence, we can cache the parameters for

our empirical null distribution and re-use these for efficient application of the GOAT algorithm. With the

cached null parameters (mu, sigma, xi for a skew-normal) in hand, geneset significance testing only

requires the computation of a geneset score (mean of respective gene scores) and testing it against the

given skew-normal.

Finally, we estimate geneset p-values by comparing their scores to size-matched null distributions and

apply multiple testing correction.

GOAT implementation

We have implemented the GOAT algorithm in an open source R package that is available at

https://github.com/ftwkoopmans/goat. Genesets can be automatically downloaded from the GO

database using the Bioconductor[2] infrastructure. Alternatively, GO genesets can be imported directly

from 'gene2go' files that are available through NCBI so users can control which version of the GO

annotations they use and reproduce their analyses at any time using the exact same geneset definitions

as input independent of Bioconductor versions. To work with alternative geneset definitions beyond GO,

users may import genesets from the synaptic gene knowledgebase SynGO[3] or import genesets in the

commonly used GMT format. Crucial parts of the (bootstrapping part of the) GOAT algorithm were

implemented in optimized C++ code to enable efficient generation of permutation-based null

distributions. As a result, finding enriched GO terms with the full GOAT algorithm using 500 000

bootstrap iterations completes in the order of seconds on a typical personal computer (Figure S2). Using

the precomputed null distributions, GOAT completes in a second.

Multiple testing correction in the GOAT R package and GOAT online, is implemented as a two-step

process; First, we apply Bonferroni adjustment independently applied per geneset 'source' (i.e. GO

MF/CC/BP). However, users may also opt for the Benjamini-Hochberg procedure. Second, Bonferroni

adjustment is applied according to the number of geneset sources that were tested. For example, when

working with the GO database a p-value correction (in addition to the first step) is applied to account for

3 separate that are applied tests across GO domains MF, CC and BP. In Figure 1, significant genesets at

Bonferroni adjustment at $\alpha = 0.05$ are shown.

Comparative analyses with ORA (i.e. Fisher's exact or hypergeometric test) and GSEA (using the fGSEA R

package) described in this manuscript were also performed using the GOAT R package, in which we

integrated both algorithms to facilitate benchmarking using the same codebase and workflow.

Specifically, fGSEA version 1.22.0 was used in the analyses presented here.

Datasets and databases

For the Colameo proteomics and gene expression data[4], we used the results from the "Compartment"

test available in supplementary data and defined the foreground genes for ORA as proteins with adjusted

p-value of at most 10^{-4} on account of the large number of significant hits even at this stringent threshold.

The proteomics dataset yielded a genelist with 3935 genes, the gene expression genelist contained

11922 genes. For all other studies we used the same threshold for adjusted p-values as reported in the

original study when performing ORA, which was 5% FDR in all cases. The Sahadevan gene expression was

imported from Table S8 (condition "24 hours") which resulted in a genelist of 13849 genes[5]. The

Higginbotham proteomics data was imported from Table S2 which resulted in a genelist of 2715 genes[6].

The Hondius proteomics dataset was imported from Table S8 (condition "Control vs Tangle") which

resulted in a genelist of 2261 genes[7]. The Wingo proteomics data was imported from Table S4 which

resulted in a genelist of 8170 genes[8]. The Klaassen proteomics data was obtained from the MS-DAP[9]

GitHub repository (https://github.com/ftwkoopmans/msdap), processed with MS-DAP[9] version 1.0.6 to

obtain differentially expressed proteins and finally mapped to a genelist of 378 genes.

The reported proteins/genes in each study were mapped to NCBI Entrez human gene identifiers using

the GOAT R package. For records that mapped to the same gene (e.g. protein isoforms) we retained the

single entry per gene with strongest p-value.

All datasets are bundled with the GOAT R package for the convenience of repeating presented analyses

without having to download the source data and repeat data preparations; after loading the goat R

package, issue the R statement *data(goat_example_datasets)*. The R code that was used to process

these data tables into the genelists used in this manuscript is provided at the GOAT GitHub repository, as

is all code that generated the figures presented in this manuscript.

Gene Ontology data from NCBI gene2go, downloaded at 2023-11-01, was loaded into the GOAT and this

version of the GO database was used for all GO analyses. For SynGO, release 1.2 was obtained from

https://www.syngoportal.org and genesets were loaded into GOAT.

REFERENCES FOR METHODS

1.   Taleb, N.N. arXiv:2001.10488 (2020).
2.   Huber, W. et al. Orchestrating high-throughput genomic analysis with Bioconductor. *Nat*
246   *Methods* **12**, 115-121 (2015).
247   3.   Koopmans, F. et al. SynGO: An Evidence-Based, Expert-Curated Knowledge Base for the Synapse.
248   *Neuron* **103**, 217-234 e214 (2019).
249   4.   Colameo, D. et al. Pervasive compartment-specific regulation of gene expression during
250   homeostatic synaptic scaling. *EMBO Rep* **22**, e52094 (2021).
251   5.   Sahadevan, S. et al. Synaptic FUS accumulation triggers early misregulation of synaptic RNAs in a
252   mouse model of ALS. *Nat Commun* **12**, 3027 (2021).
253   6.   Higginbotham, L. et al. Integrated proteomics reveals brain-based cerebrospinal fluid biomarkers
254   in asymptomatic and symptomatic Alzheimer's disease. *Sci Adv* **6** (2020).
255   7.   Hondius, D.C. et al. The proteome of granulovacuolar degeneration and neurofibrillary tangles in
256   Alzheimer's disease. *Acta Neuropathol* **141**, 341-358 (2021).
257   8.   Wingo, A.P. et al. Shared proteomic effects of cerebral atherosclerosis and Alzheimer's disease
258   on the human brain. *Nat Neurosci* **23**, 696-700 (2020).
259   9.   Koopmans, F., Li, K.W., Klaassen, R.V. & Smit, A.B. MS-DAP Platform for Downstream Data
260   Analysis of Label-Free Proteomics Uncovers Optimal Workflows in Benchmark Data Sets and
261   Increased Sensitivity in Analysis of Alzheimer's Biomarker Data. *J Proteome Res* **22**, 374-386
262   (2023).
263   10.   Klaassen, R.V. et al. Shisa6 traps AMPA receptors at postsynaptic sites and prevents their
264   desensitization during synaptic activity. *Nat Commun* **7**, 10682 (2016).

265