Cognitive Phenotype Shifts in Risk-Taking: Interplay of Non-Suicidal Self-Injury Behaviors and Intensified Depression

Yi-Long Lu[1,#], Yuqi Ge[2,3,#], Mingzhu Li[2,3], Shutian Liang[1], Xiaoxi Zhang[4], Yupeng Sui[4], Lei Yang[2,3], Xueni Li[2,3], Yuyanan Zhang[2,3], Weihua Yue[2,3,5], Hang Zhang[1,5,6,7,8*], Hao Yan[2,3,*]

[1] School of Psychological and Cognitive Sciences and Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing, China.
[2] Peking University Sixth Hospital, Peking University Institute of Mental Health, Beijing, China.
[3] NHC Key Laboratory of Mental Health (Peking University) and National Clinical Research Center for Mental Disorders (Peking University Sixth Hospital), Beijing, China.
[4] Department of Mental Health, Changzhi Medical College, Changzhi, China.
[5] PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing, China.
[6] Peking-Tsinghua Center for Life Sciences, Peking University, Beijing, China.
[7] National Key Laboratory of General Artificial Intelligence, Peking University, Beijing, China.
[8] Chinese Institute for Brain Research, Beijing, China.

[#]: co-first authors.
*: corresponding authors. Email: hao_y@bjmu.edu.cn (H.Y.), hang.zhang@pku.edu.cn (H.Z.)

ABSTRACT

Background

Non-suicidal self-injury (NSSI) behavior is significantly prevalent in both adolescents and psychiatric populations, with the highest risk observed in individuals with major depressive disorder (MDD). NSSI can be considered a result of risky decision-making in response to negative emotions, where individuals choose self-harm over other harmless alternatives, suggesting a potential decision-making deficit in NSSI individuals. This study delves into the complex relationship between NSSI and depression severity in decision-making and its cognitive underpinnings.

Method

We assessed decision behaviors in MDD patients with and without NSSI and healthy controls using the Balloon Analogue Risk Task, which involves risk-taking, learning, and exploration in uncertain scenarios. Using computational modeling, we dissected the nuanced cognitive dimensions influencing decision behaviors. A novel statistical method was developed to elucidate the interaction effects between NSSI and depression severity.

Result

Contrary to common perceptions, we found that individuals with NSSI behaviors were typically more risk-averse, but as depressive symptoms intensified, they began to perceive less risk and meanwhile behave more randomly.

Conclusions

This research offers new insights into the cognitive aspects of NSSI and depression, underscoring the importance of recognizing the influence of comorbid mental disorders in deciphering the cognitive underpinnings of such behaviors, especially in the context of prevalent cross-diagnostic phenomena like NSSI behaviors.

KEYWORDS

Non-suicidal self-injury; Major depressive disorder; Decision under risk; Hierarchical

Bayesian modeling; Risk aversion; Cognitive phenotype

INTRODUCTION

Non-suicidal self-injury (NSSI), prevalent in adolescents (1) and psychiatric populations (2,3), particularly in individuals with depression (4,5), involves deliberate self-harm behaviors (e.g., cutting, burning, scratching, and banging or hitting) without suicidal intent. Though distinct from suicidal behavior, NSSI is crucial to understand due to its strong association with future suicidal tendencies (6,7). NSSI has been described as a morbid form of self-help (8), often used as a coping mechanism for intense emotional distress (9–12). Similar to suicidal actions, choosing to self-harm instead of engaging in more harmless coping solutions (e.g., talking to a friend, practicing mindfulness, or seeking professional counseling) reflects a maladaptive decision-making process (13). Our work, therefore, focuses on the decision characteristics of individuals with NSSI, aiming to identify the potentially multi-faceted cognitive dysfunctions behind NSSI decisions.

Despite the initial hypothesis linking higher impulsivity to NSSI, especially given its prevalence among adolescents, this connection has found little support in research that controls for psychiatric confounds (14,15). A more nuanced perspective considers risky decision-making (16), through which perspective individuals with NSSI would evaluate various emotion-coping options, each with its own set of risks, rewards, and losses. These individuals effectively outweigh the emotional relief from self-harm against its negative consequences (e.g., pain, bleeding, and scarring), and prefer it over more harmless solutions. However, studies using standard risky decision tasks, such as the Iowa Gambling Task, have yielded inconsistent results in differentiating between NSSI and non-NSSI individuals (14,15,17). One reason for these mixed findings could be the inherent stochasticity of risky decision tasks, where different choices lead to variable experiences and thus exert different influences on subsequent decision behaviors, rendering simple behavioral measures less effective in capturing the complexity of individuals' decision characteristics. Furthermore, the

frequent comorbidity of NSSI with conditions like depression (18–20) adds another layer of complexity, potentially confounding the interpretation of the findings. For example, behavioral differences observed in such tasks might not solely reflect NSSI characteristics but also varying levels of depression among individuals. These limitations highlight the necessity of using more advanced methods to understand NSSI-related decision characteristics.

In the present study, we investigated individuals with major depressive disorder (MDD) exhibiting NSSI behavior on at least one day within the past year (D+NSSI), contrasting them with MDD individuals without NSSI (D) and healthy controls (HC). Our approach involved the use of the Balloon Analogue Risk Task (BART, 21) alongside computational modeling to provide a comprehensive understanding of NSSI-related decision characteristics and their modulation by depression. The BART, which simulates real-world risk-taking behavior by requiring participants to pump a virtual balloon for potential rewards, assesses not only attitudes toward risk or loss, as widely known, but also delves into broader cognitive dimensions. These include participants' prior beliefs about risk, their learning from outcomes, and behavioral consistency under risk and uncertainty. Using computational modeling, we were able to disentangle these distinct cognitive dimensions from participants' decision behaviors, by estimating the model parameters for each group based on the hierarchical Bayesian method. Furthermore, we developed a novel statistical method to isolate the effects of NSSI, depression level, and their interaction on the model parameters. This allowed us to unravel the complex interplay of NSSI and depression in shaping individual decision characteristics, offering insights that are otherwise unattainable.

METHODS AND MATERIALS

Participants

From April 2022 to Sept 2023, we recruited 101 outpatients or inpatients diagnosed with depression from Peking University Sixth Hospital, including 58 cases who engaged in NSSI behaviors at least one episode in the past 12 months (D+NSSI) and 42 cases without NSSI (D) behaviors. Additionally, 157 healthy controls (HC) were recruited from two universities. The study had been approved by the medical ethics committee of Peking University Sixth Hospital and all participants provided informed consent.

Fifteen participants from the HC group were excluded due to a history of NSSI behavior. Furthermore, one participant from the HC group and one from the D+NSSI group were excluded for consistently inflating the balloons to the same extent in the decision task. One participant from the D group was excluded for data missing. As a result, a total of 241 participants were included in the final data analysis.

All D+NSSI participants had engaged in NSSI behaviors for at least one episode in the past 12 months, whereas all D participants had no history of self-injury. Additional criteria for inclusion were: a) Individuals aged 16 to 25 years old; b) those diagnosed with major depressive disorder according to the International Classification of Diseases, Tenth Edition (ICD-10). Exclusion criteria included: a) patients engaging in suicidal self-injury; c) current diagnosis of other mental disorders such as pervasive developmental disorders, psychotic disorders, manic episodes, substance dependence, or obsessive-compulsive disorders; and d) presence of severe and unstable physical illness.

Supplementary analyses were performed to compare the subsets of participants who engaged in NSSI behaviors for at least five episodes and those fewer than five episodes in the past 12 months ($N = 30$ for D+NSSI[≥5] and $N = 27$

D+NSSI[<5]). Little differences in decision characteristics were found between these two sub-groups of participants.

Psychiatric Assessment

The Zung self-rating depression scale (SDS) (34) and the Zung self-rating anxiety scale (SAS) (35) were used to assess depression and anxiety levels in this investigation, and raw scores were used. The SDS and SAS are both 20-item scales. Raw scores of 40 and 36 are typically considered indicative of the presence of clinical depression and anxiety, respectively (36,37).

Task and Design

In the Balloon Analogue Risk Task (BART) shown in Figure 1A, each trial presents participants with a virtual balloon on their computer screen. Participants can press a specific key to "pump" the balloon, which adds one point to their prize pool for each pump. However, every pump also carries a risk of causing the balloon to pop. If the balloon pops, the points in the prize pool for that balloon are lost, and the trial ends. Thus, participants must choose between pumping the balloon to potentially increase their reward point or collecting the accumulated points to conclude the trial safely. The total points collected across all trials would be converted into monetary compensation at the end of the experiment.

Each participant completed 30 trials with red balloons and 30 with blue balloons, presented in a random order. The two different colors represent two levels of risk of popping, corresponding to different optimal numbers of pumps to maximize expected reward (see Inset in Figure 1A). One color of balloon is associated with an equal chance of popping on any pump between the 1st and the 32nd ("32-balloon"), while the other between the 1st and 128th ("128-balloon"). In the illustration of Figure 1A, red balloons are 32-balloons and blue balloons are 128-balloons. The real color-risk correspondence was counterbalanced between participants. Participants were

not informed the probability of popping or which balloon would be riskier, and therefore had to learn from their own experience in the experiment.

The experiment consisted of three blocks of 20 trials: the first block was a randomized mix of 10 red and 10 blue balloon trials, and the subsequent two blocks consisted of 20 trials of each balloon color. The order of the last two blocks was counterbalanced across participants. In total, it took approximately 10 minutes.

Statistical Analysis

We conducted the following linear mixed-effects model (LMM) analysis on the number of pumps N_{pump} in the non-popping trials to test the potential influence of major depressive disorder (D) and NSSI behavior (NSSI) on participants' decision:

$$N_{pump} \sim 1 + \mathrm{B} * h * \mathrm{Group} + (1 + h | \mathrm{paticipant}), \tag{1}$$

where B stands for the balloon type (–1 for "32-balloon" and 1 for "128-balloon"), h is the index of the balloon in the experiment, Group denotes categorical variables for the three participant groups, HC, D, and D+NSSI. The $(\cdot | \mathrm{paticipant})$ denotes the random-effects terms that vary with participants. To further examine the potential different effects of NSSI<5 versus NSSI≥5, we conducted a supplementary LMM analysis:

$$N_{pump} \sim 1 + \mathrm{B} * h * D * \mathrm{Group}^4 + (1 + h | \mathrm{paticipant}), \tag{2}$$

where Group^4 indicated 4 participant groups, that is, HC, D, D+NSSI[≥5], and D+NSSI[<5]. All data were normalized on the group level before entering LMMs. All the LMMs were implemented in R 4.0.3.

Computational Modeling

The computational model we used to fit participants' decision behaviors in BART (illustrated in Figure 2A) is an adaptation of the Exponential-Weight Mean–Variance (EWMV) model of Park et al. (25). Below we first introduce the basic modeling idea and specific assumptions of EWMV, and then our adaptation—EWMV with variance standardization (EWMV-VS).

Basic modeling idea. After comparing a comprehensive set of computational models they constructed for BART, Wallsten et al. (38) found that human participants' behaviors in BART are best captured by a model with three key assumptions for the belief updating and decision processes: 1) the agent believes that the probability of balloon popping could vary with balloon colors but is constant across different pumps for the same color of balloon, 2) the agent updates the probability of pop before each trial based on previous experience with popping and successful inflating, and 3) each agent has specific attitudes towards loss or risk. Following this stream of assumptions, Park et al. (25) further proposed the EWMV model, which has closely-related mathematical forms and a comparable predictive power, but whose parameters can be more reliably recovered from human behavior, thus allowing for a more accurate characterization of individual differences.

EWMV. According to the EWMV model, the agent starts with a specific prior probability (parameter ϕ) for balloon popping, and updates its belief based on the numbers of popping and successful inflating previously observed for the balloons that matched the color of the current balloon. In the agent's belief updating process, the relative weights for the prior probability and experience are approximated by exponential weighting functions:

$$p_h = \phi \exp\left(-\xi \sum_{h'=1}^{h-1} m_{h'}\right) + \left(1 - \exp\left(-\xi \sum_{h'=1}^{h-1} m_{h'}\right)\right) P_{h-1}. \qquad (3)$$

Here p_h denotes the pop probability per pump for the h-th balloon in its color, ξ is the parameter for learning rate. With $m_{(\cdot)}$ denoting the number of pumps for a specific balloon indicated by the subscript, $\sum_{h'=1}^{h-1} m_{h'}$ is the total number of pumps made to all previous $h-1$ balloons in the same color. For these $h-1$ balloons, P_{h-1} is the observed pop probability per pump:

$$P_{h-1} = \frac{\sum_{h'=1}^{h-1} d_{h'}}{\sum_{h'=1}^{h-1} m_{h'}}, \qquad (4)$$

where d_h indicates whether balloon h finally popped (1) or not (0).

Suppose the agent has pumped the current balloon for k times, accumulating k points. At this moment, pumping corresponds to a gamble with probability $1 - p_h$ to gain 1 more point but probability p_h to lose the k points; collecting is to win or lose nothing. To decide whether to pump or to collect the accumulated points, the agent calculated the expected utility of pumping relative to collecting:

$$U_{h,k} = [(1 - p_h) - p_h \lambda k] + \rho p_h (1 - p_h)(1 + \lambda k)^2. \qquad (5)$$

The first part of the equation, $[(1 - p_h) - p_h \lambda k]$, is the sum of expected gain and expected subjective loss, with loss scaled by the loss aversion parameter λ, that is, how many units of gain one unit of loss equals to. The second part is the risk discounting term, with the variance of the gamble multiplied by the risk preference parameter ρ.

The agent's probability of choosing pumping is a softmax function of this expected utility:

$$r_{h,k} = \frac{1}{1 + e^{-\beta U_{h,k}}}, \qquad (6)$$

where the parameter β characterizes choice consistency. The final choice of pumping or collecting is a Bernoulli random variable based on $r_{h,k}$.

EWMV with variance standardization. When the probability of balloon popping is relatively small (e.g., 1/32 and 1/128), there could be strong correlation between the mean and variance terms in the computation of expected utility (Eq. 5). As a result, the risk preference parameter ρ cannot be accurately estimated. Similar issues would still occur if the risk preference parameter is incorporated in expected utility not as the scaling factor of variance but through the more classical form of a utility function (39,40). To alleviate this issue, we adopt the standardization method (40) to perform variance standardization on the expected utility:

$$U_{h,k} = \frac{[(1 - p_h) - p_h \lambda k] + \rho p_h (1 - p_h)(1 + \lambda k)^2}{\sqrt{p_h (1 - p_h)}(1 + \lambda k)}. \qquad (7)$$

The standardized $U_{h,k}$ is then entered into Eq. 6 to obtain the probability of choosing pumping. Similar to EWMV, there are five free parameters in our EWMV-VS model: prior belief of popping ϕ, learning rate ξ, choice consistency β, loss aversion λ, and risk preference ρ.

Model Fitting. Hierarchical Bayesian modeling was used for parameter estimation, which allows more accurate estimation of group-level effects. The model parameters were estimated from participants' choice data using the Markov Chain Monte Carlo method implemented by the PyMC package (version 5.9.2) on python 3.11. Four independent chains were run, each with 1000 samples after an initial burn-in of 1000 samples, which resulted in a good convergence of $\hat{R} \leq 1.01$ for all estimated parameters. The 95% highest density intervals (HDI) were calculated for the group-level effects.

Effects-on-Parameters Analysis. To assess the joint effects of NSSI and depression severity on individuals' decision characteristics, we developed an Effects-on-Parameters Analysis (EPA). Specifically, we built a hierarchical Bayesian model based on EWMV-VS, where each of the five parameters of EWMV-VS is generated from a linear combination of NSSI, SDS, and their interaction term. The 15 coefficients were then estimated from participants' choice data via hierarchical Bayesian modeling as described above. For the control analyses, we also added the age and education year as regressors to the group-level linear model. The SDS, age and education year were normalized before entering the model.

RESULTS

Demographic and clinical characteristics

A total of 241 participants were included in data analysis, including 142 healthy controls (HC), and 99 patients diagnosed with major depressive disorder (MDD),

among whom were 57 cases with comorbid NSSI (D+NSSI) who engaged in NSSI behaviors at least one episode in the past 12 months and 42 reporting no history of NSSI (D). Participants' gender, age, education year, self-rating anxiety and depression scales were recorded (Table 1). As expected, the three groups of participants significantly differed in the depression level (SDS, $F(2, 238) = 198.908$, $p < 0.001$), with patients higher than HC (*post-hoc* tests, $p < 0.001$ for D and D+NSSI) and D+NSSI higher than D ($p < 0.001$). Similar differences were found in participants' anxiety level (SAS, $F(2, 238) = 123.718$, $p < 0.001$; D+NSSI > D > HC, all $p < 0.001$). Because anxiety is highly comorbid with depression (22), there was a high correlation between the SDS and SAS scores in our participants with MDD (Pearson's $r = 0.893$, $p < 0.001$), which made their effects practically undistinguishable. Therefore, we will report the effects of SDS only. The three groups had similar gender ratios ($\chi^2(2) = 3.306$, $p = 0.191$). They had small but significant differences in age ($F(2, 238) = 10.803$, $p < 0.001$) and education year ($F(2, 238) = 7.313$, $p < 0.001$). We reported analyses that regressed out the potential confounding effects of age and education year.

Following previous studies (23), here we used the term NSSI in a more general way, referring to individuals who engaged in NSSI behaviors in the past year instead of more specifically to individuals with a separate diagnosis of NSSI disorder in the Diagnostic and Statistical Manual for Mental Disorders, Fifth Edition (DSM-5, 24). Nevertheless, we also report supplementary analyses that compare the subgroups of participants who met and who did not meet the NSSI frequency criterion for DSM-5 diagnosis according to the self-reported NSSI history.

Table 1. Demographic and clinical characteristics

	Total	HC	D	D+NSSI	χ^2/F	*p*-value
	(*N* = 241)	(*N* = 142)	(*N* = 42)	(*N* = 57)		
N by gender					3.306	0.191

Male	78 (32.4%)	49 (34.5%)	16 (38.1%)	13 (22.8%)		
Female	163 (67.6%)	93 (65.5%)	26 (61.9%)	44 (77.2%)		
Age (year)	19.80±2.36	19.30±2.05	21.10±2.57	20.09±2.54	10.803	<0.001
Education year	13.49±2.11	13.11±1.77	14.43±2.49	13.75±2.36	7.313	<0.001
SAS	39.66±14.69	31.33±6.30	45.90±12.46	55.82±15.48	123.718	<0.001
SDS	45.34±16.83	34.93±8.00	51.90±11.56	66.42±13.96	198.908	<0.001

Note. Demographic or scale scores are in mean ± standard deviation. SAS: self-rating anxiety scale; SDS: self-rating depression scale.

Group Differences in Behavioral Measures

On each trial of BART (Figure 1A), participants saw a virtual balloon on the screen and need to decide whether to pump (to potentially increase reward points) or collect points and end the trial. Each pump could increase the balloon's size and thus the reward points by 1 but came with the risk that it may pop and nullify the points. Each trial thus presented a continual gamble between potential reward and loss. Each participant completed 60 trials of 30 red and 30 blue balloons, initially not knowing which color represented a higher risk. The color with higher risk would pop after 1 to 32 pumps ("32-balloon") while the lower-risk color after 1 to 128 pumps ("128-balloon"). The number of pumps that maximizes expected reward is respectively 16 and 64 for the 32- and 128-balloons (Figure 1A, inset). Including these two different levels of risk in the experimental design allowed us to infer participants' prior beliefs and learning as well as their attitudes towards loss and risk from their pumping decisions.

Group differences in N_{pump}. Following the common practice of BART (21), a primary behavioral measure we used is participants' number of pumps in non-popping trials (denoted N_{pump}). Consistent with classical findings, for both levels of

risk, participants pumped much less than the optimal number (Figure 1BC).

Meanwhile, when we plot how the mean N_{pump} of each participant group changed over trials (Figure 1B), we can see differences between different groups as well as trends of learning, at least for the 128-balloons. A linear mixed-effect model (LMM) analysis was performed on N_{pump} to identify the effects of participants group, balloon type, and trial number as well as their interactions. This LMM analysis showed a significant main effect of participant group (F(2, 235.63)= 3.68, p = 0.027). Post-hoc analysis (multi-comparison corrected) further indicated that the D+NSSI group pumped fewer than the HC group (b = –0.213, SE = 0.078, p_{tukey} = 0.018).

Learning effects. We also observed significant learning effects for all groups. The mean N_{pump} for 128-balloons was larger than that of 32-balloons (F(1, 232.18)= 296.01 p < 0.001). As the experiment proceeded, participants' N_{pump} increased overall (F(1, 231.74)= 57.98, p < 0.001), and changed differently under different balloon type (interaction of balloon type by trial number, F(1, 8466.29)= 340.76, p < 0.001), with the N_{pump} increasing over trials for 128-balloons (simple main effect, b = 0.261, SE = 0.018, p < 0.001) but staying constant for 32-balloons. We did not find significant differences between different group of participants in the learning effects (interaction of participants group by trial number, F(2, 231.84)= 0.07, p = 0.94).

Group differences in σ_{pump} and N_{gather}. To provide a more comprehensive behavioral measurement for participants' decision characteristics in BART, we also calculated the standard deviation of N_{pump} across trials (denoted σ_{pump}) and the number of non-popping (i.e., reward gathered) trials (denoted N_{gather}) for each balloon type and participant group (Figure 1DE). Mixed-design ANOVAs for these two measures showed a significant interaction of participant group by balloon type for σ_{pump} (F(2, 238) = 3.10, p = 0.047), whereas no significant interaction was observed for N_{gather} (F(2, 238) = 0.476, p = 0.622). We also observed significant main effects

of participant group for σ_{pump} (F(2, 238) = 3.39, p = 0.035) and N_{gather} (F(2, 238) = 4.88, p = 0.008), other than the obvious differences between different balloon types (F(1, 238) = 187.33, p < 0.001 for σ_{pump} and F(1, 238) = 679.75, p < 0.001 for N_{gather}). According to simple main effects, participants in different groups had significantly different σ_{pump} for 128-balloons (F(2, 238) = 3.44, p = 0.034), but not for 32-balloons (F(2, 238) = 0.73, p = 0.484). The following *post-hoc* comparisons further indicates that for 128-balloons, participants in the D+NSSI group had smaller σ_{pump} than both the HC (t = −3.46, p_{holm} = 0.004) and D groups (t = −2.68, p_{holm} = 0.038). Compared to the HC group, participants in the D+NSSI group also had larger N_{gather} (t = 3.09, p_{holm} = 0.007).

Taken together, these behavioral measures of BART suggest that the HC, D, and D+NSSI groups had distinct decision patterns, though all groups similarly learned from their experiences. Because all behavioral measures are the joint consequence of multiple cognitive components (e.g., learning, risk attitude, etc.), the underlying cognitive differences remained elusive. We will disentangle the cognitive dimensions below, with the help of computational modeling.

Figure 1. Balloon Analogue Risk Task (BART) and behavioral measures for the three groups of participants. A. Schematic representation of the BART, where at each moment participants choose to either pump the balloon to potentially accumulate one more reward point or to collect their accumulated points. If the balloon pops, the reward points accumulated for the balloon would be lost. The risk of balloon popping increases with each pump and varies with the color of the balloon. A new balloon starts after the old balloon popping or being collected. Each participant completed 30 red and 30 blue

balloons (trials) in a random order. **Inset at top right**: Expected gain as a function of the number of pumps. The optimal numbers for the two balloon types (max pump before popping: 32 and 128), as indicated by the dashed vertical lines, are respectively 16 and 64. **B–E** plot the behavioral measures for the three groups of participants ($N = 241$). Color codes: light-green for the 142 healthy controls (HC), blue-green for the 42 participants with major depressive disorder (MDD) but no NSSI behavior (D), and dark-blue for the 57 participants with MDD and NSSI (D+NSSI). **B**. How the mean number of pumps in non-popping trials (N_{pump}) changed over trials. Curves in dashed lines and solid lines are respectively for the balloon types 32 and 128. Shadings denote 1 SE. **C**. Results of the linear mixed-effects model analysis on N_{pump}. The standardized coefficient b is plotted separately for each group (color coded). Error bars represent 1 SE of the coefficients. **D**. Standard deviation of N_{pump} across trials (σ_{pump}) for each balloon type and participant group. **E**. The number of non-popping (i.e., reward gathered) trials (N_{gather}) for each balloon type and participant group. Error bars denote 1 SE. *: $p < 0.05$. **: $p < 0.01$. ***: $p < 0.001$. Asterisks on the left or right side of an ensemble of coefficients in **C** denote significant deviations from 0 for all these coefficients. The asterisks in **E** indicate a main effect between the HC and D+NSSI groups, regardless of the balloon type.

Five Cognitive Dimensions Decomposed by the Computational Model

The computational model we used to fit participants' decision behaviors is an adaptation of the Exponential-Weight Mean–Variance (EWMV) model of Park et al. (25), termed EWMV-VS (see Methods), which outperforms its precedent in predictive power (see Supplemental Table S1) and has a good parameter recovery performance (Supplemental Figure S1). As illustrated in Figure 2A, the model starts with an agent's initial belief about the likelihood of balloon popping (ϕ), which is updated with each successive trial based on the outcomes of previous balloons. This update is modulated by a learning rate (ξ), controlling the speed at which new

information influences the agent's belief of balloon popping. Confronted with the choice to pump and potentially increase their tally by one point or to collect and secure their current pool of points, the agent evaluates this decision by considering the risk of popping and losing all accumulated points for the balloon. The model quantifies the agent's aversion to loss (λ) and their inclination towards risk (ρ) to calculate the expected utility of each pump, while the agent's propensity to stay consistent in their choices is represented by the parameter β. When fitted to participants' decision behaviors, the five parameters of the EWMV-VS model—prior belief of pop ϕ, learning rate ξ, choice consistency β, loss aversion λ, and risk preference ρ—can thus represent participants' characteristics on five distinct cognitive dimensions.

Some of these model parameters have intuitive behavioral implications, as indicated by the correlations between individual participants' model parameters and their behavioral measures (Figure 2B). For example, the average number of pumps in the initial trials of the 128-balloons, labeled "first 5 pumps", serves as a behavioral proxy for the model's parameter for prior belief of pop (ϕ). Similarly, "ΔN_{gather}", the differences in non-pop trials between the 128- and 32-balloons, is a behavioral reflection of the individual's learning rate (ξ), demonstrating how participants adjust their strategies based on experience. Though less intuitive or not unique to any specific parameter, the N_{pump}, σ_{pump}, and the standardized σ_{pump}/N_{pump}, are respectively closely related to loss aversion (λ), risk preference (ρ), and choice consistency (β).

Figure 2. The computational model and its parameters' behavioral implications. A.

Illustration of the computational model used to simulate participant decisions in BART.

The model agent integrates its prior belief of balloon popping (ϕ) with previous

observations up to the last trial (i.e., balloon $h-1$), resulting in the updated belief of

P_{pop}. How fast the prior belief is wiped off by increasing observations is determined by the

learning rate (ξ). For the current balloon h that has accumulated k points, the agent is

now faced with the decision of either pumping to gain 1 more point but with a chance of

P_{pop} losing all k points, or collecting the points. The agent's loss aversion (λ) and risk

preference (ρ) contribute to its expected utility of pumping the balloon. The probability of

choosing pumping is further modulated by the agent's choice consistency (β). **B**. Scatter

plots showing how individuals' model parameters (prior belief of pop ϕ, learning rate ξ,

choice consistency β, loss aversion λ, and risk preference ρ) correlated with their

behavioral measures in the BART. "First 5 pump" refers to the mean number of pump in

the first 5 trials of the 128-balloons, which reflects the individual's prior belief of pop.

"ΔN_{gather}" refers to the difference in the number of reward-gathered trials (N_{gather})

between the 128- and 32-balloons, which reflects how much the individual learns from

experience. The N_{pump} and σ_{pump}, as in Figure 1, respectively refer to the mean and

standard deviation of the number of pumps, which and the combination of which are

closely related to loss aversion, risk preference, and choice consistency. Data points are color-coded to indicate whether the behavioral measure is elicited from the 128-balloons (in gray-blue) or the difference between the 128- and 32-balloons (in gray). Each dot denotes one individual participant. The correlation coefficient (r) and p-value on each panel is based on Pearson's correlation.

Revealing Group-Level Cognitive Differences: Hierarchical Bayesian Modeling

To assess differences in each cognitive dimension across the three groups, we fit the EWMV-VS model to participants' decision behaviors using hierarchical Bayesian modeling (see Methods). Among the five parameters (Figure 3A), we observed significant group differences in the parameters of choice consistency and risk preference. In particular, when compared to participants with MDD but no NSSI behavior (i.e., the D group), the D+NSSI group exhibited a higher choice consistency (difference $M = 0.24$, 95% HDI [0.03, 0.44]). The D+NSSI group also had a lower risk preference than the HC group (difference $M = -0.75$, 95% HDI [-1.5, -0.029]), indicating a more cautious approach to risk-taking.

These findings, rooted in our model analysis, echo the behavioral patterns described earlier, specifically the reduced variability in the number of pumps (σ_{pump}) among the D+NSSI group (Figure 1D). While behavioral data alone might suggest this decreased variation could stem from diminished learning, implying more uniform behavior from early to late trials, our model-based analysis mostly rules out this possibility.

Figure 3. Estimated model parameters for different participant groups. A. The 95% highest density interval (HDI) of the posterior density of the group-level model parameters estimated from hierarchical Bayesian model fitting. Significant differences between groups were found for choice consistency (β) and risk preference (ρ). See text. **B–D** show the joint effects of NSSI and depression severity (as measured by the Self-Rating Depression Scale, SDS) on model parameters, where the participants in analysis were limited to the D and D+NSSI groups with an overlapping range of SDS. Each dot denotes one individual participant. The shading on the regression line denotes the 95% HDI. **B**. Interaction (b_{IA}) between NSSI and SDS on prior belief of pop (ϕ). Higher depression severity was associated with a lower belief in pop likelihood for the NSSI group, while the reverse for the non-NSSI group. **C**. Interaction (b_{IA}) between NSSI and SDS on choice consistency (β). Higher depression severity was associated with less consistent choices in the NSSI group, while the reverse for the non-NSSI group. **D**. Main effect of NSSI (b_N) for risk preference (ρ). The NSSI group was less tolerant of risk than the non-NSSI group. The asterisk (*) denotes that the 95% HDI of the group difference did not contain 0 (similar to 0.05-level significant differences in the frequentist statistics such as t tests).

Unraveling NSSI and Depression Interplay: Effects-on-Parameters Analysis

We noted that the D+NSSI group had a significantly higher depression severity than the D group (as indicated by the SDS score in Table 1), despite both groups being diagnosed with MDD and with SDS scores exceeding the cut-off point for clinical significance (26). To distinguish the impact of NSSI from depression severity, we combined two strategies. First, we selected from the D and D+NSSI groups the participants whose SDS fell into an overlapping range ($N = 81$, with 37 D and 44 D+NSSI). Second, we developed a novel Effects-on-Parameters Analysis (EPA, see Methods), which includes NSSI, SDS, and their interaction as linear predictors of the EWMV-VS model's parameters in hierarchical Bayesian modeling, providing a granular view of their joint effects on individuals' decision characteristics.

By applying EPA to the 81 participants' decision data, we found that NSSI was still associated with a decrease in risk preference ($M = -1.1$, 95% HDI [-2.1, -0.09]), even when depression severity was controlled (Figure 3D). NSSI and SDS jointly influenced the prior belief of pop and choice consistency in a more complicated way, showing as interaction effects. Specifically, the prior belief of pop increased with SDS for participants without NSSI but decreased with SDS for those with NSSI behaviors (Figure 3B; $M = -0.68$, 95% HDI [-1.2, -0.19]). Choice consistency rose with SDS among participants without NSSI but fell among those with NSSI (Figure 3C; $M = -0.28$, 95% HDI [-0.5, -0.03]). These effects were robust, which were similar when the effects of age and education year were controlled (Supplemental Figures S2 and S3).

In other words, participants with NSSI behaviors were more averse towards risk in general. However, as their depressive symptoms became more severe, they appeared to perceive the environment as less risky and were more random in their choices.

Table 2. Participants included in the effects-on-parameters analysis

	Total	D	D+NSSI	t/χ^2	*p*-value
	(N=81)	(N=37)	(N=44)		
N by gender				2.987	0.084
Male	23(28.4%)	14(37.8%)	9(20.5%)		
Female	58(71.6%)	23(62.2%)	35(79.5%)		
Age(year)	20.85±2.52	21.00±2.63	20.73±2.44	−0.484	0.630
Education year	14.27±2.38	14.24±2.49	14.30±2.32	0.098	0.922
SAS	49.22±11.74	47.97±11.61	50.27±11.87	0.877	0.383
SDS	58.09±9.96	54.81±8.81	60.84±10.12	2.832	0.006

Note. Demographic or scale scores are in mean ± standard deviation. SAS: self-rating anxiety scale; SDS: self-rating depression scale. The *t* values for age, education year, SAS, and SDS indicates the differences between the D+NSSI and the D groups.

Similar Decision Behaviors in MDD Patients with NSSI < 5 and NSSI ⩾ 5

Participants who had NSSI behaviors but less than five episodes in the past 12 months would not be diagnosed as the NSSI disorder (DSM-5, 24). The D+NSSI group in our analyses above comprised cases with varying frequencies of NSSI. To test the possibility that they might have different decision characteristics, we performed supplementary analyses where the D+NSSI group was divided into two sub-groups according to the self-reported NSSI frequency in the past 12 months, denoted D+NSSI$^{<5}$ and D+NSSI$^{\geq 5}$ (27 and 30 participants, respectively).

To examine the effects of NSSI<5 versus NSSI≥5, we first performed an LMM analysis on N_{pump} similarly as before, except that the regressor of participant group included four instead of three categories (see Methods). All the significant effects identified in the previous LMM were replicated in this new LMM. That is, there were significant main effect of balloon type (F(1, 230.33)= 231.00, $p < 0.001$), trial number

($F(1, 228.64)= 50.97$, $p < 0.001$) as well as their interaction. The participant group also had a significant effect ($F(3, 234.29)= 2.65$, $p = 0.049$), but *post-hoc* differences between the groups failed to reach significance.

Hierarchical Bayesian modeling analysis for four participant groups also suggests that the NSSI≥5 and NSSI<5 groups had similar model parameters (Figure 4C). Here we separately estimated the differences between D and HC, between the two NSSI groups and D, and between NSSI≥5 and NSSI<5 in each of the five parameters. Similar to what we reported earlier for NSSI as one group, participants with NSSI had an increased choice consistency than the D group (difference $M =$ 0.24, 95% HDI [0.01, 0.51]) and a lower risk preference than the HC group (difference $M = –0.11$, 95% HDI [–2.1, –0.02]). No parameters showed significant differences between the NSSI≥5 and NSSI<5 groups (Supplemental Table S2).

Figure 4. Behavioral measures and estimated model parameters for four participant groups. A. How the mean number of pumps in non-popping trials (N_{pump}) changed over trials for four participant groups. Curves in dashed lines and solid lines are respectively for the balloon types 32 and 128. Shadings denote 1 SE. **B**. LMM model

results. b is the standardized coefficient of the linear mixed-effects model. Error bars represent 1 SE of the coefficients. **C**. The 95% highest density interval (HDI) of the posterior density of the group-level model parameters estimated from hierarchical Bayesian model fitting. The asterisk (*) denotes that the 95% HDI of the group difference did not contain 0 (similar to 0.05-level significant differences in the frequentist statistics such as *t* tests). No significant differences were found for the parameters of D+NSSI$^{<5}$ and D+NSSI$^{\geq5}$ groups. See text. *: $p < 0.05$. **: $p < 0.01$. ***: $p < 0.001$. Asterisks on the left or right side of an ensemble of coefficients in **B** denote significant deviations from 0 for all these coefficients.

DISCUSSION

In the present study, we investigated the decision characteristics of MDD patients with or without NSSI behavior in BART (21), a classic decision-making task that involves learning and exploration as well as risk-taking. In addition to model-free behavioral measures, we used computational modeling to elicit five different cognitive phenotypes from individuals' decision behaviors, providing a comprehensive understanding of NSSI-related decision characteristics and their modulation by depression. We found that the presence of NSSI behaviors is associated with a general aversion to risk. However, as depressive symptoms escalate, individuals with NSSI tend to perceive their environment as less risky and are more variable in their choices, potentially resulting in greater behavioral unpredictability.

Concerning the decision characteristics related to NSSI, little consensus has been reached (15,27), partly because confounds like depression were frequent in previous studies (28). Our finding that individuals with NSSI behaviors are more risk-averse instead of more "impulsive" as commonly perceived is consistent with that of Dillahunt et al. (14). They studied adolescents with and without self-injury history who

were otherwise almost matched in their clinical profile, showing that those with self-injury history made less pumps in BART, seeming to be "risk-averse". In our study, computational modeling allows us to extend beyond such behavioral observation and to precisely identify the cognitive phenotype that leads to the specific behavioral phenomenon, excluding alternative explanations such as an aversion to loss or failure to learn from the environment.

The Effects-on-Parameters analysis we developed further enables us to disentangle the effects of NSSI and depression severity and to uncover interaction effects on the model parameters. The prior belief of risk and choice consistency are the two cognitive dimensions that are influenced by such interactions. The former determines an individual's inclination to explore uncertain options, while the latter is related to the tendency of random exploration (29). Together, our results suggest that when their depressive symptoms deepen, while individuals without NSSI may reduce their exploration of the environment, those with NSSI may even increase their exploration and behavioral randomness, which can be risky behaviors under some occasions.

Such intricate interplay between NSSI and depression may have led to the controversial results documented in previous studies (15). Since NSSI is highly prevalent in mental disorders (5,30), our findings emphasize the need for a comprehensive approach to isolating the influence of comorbid mental disorders, in order to decipher the cognitive underpinnings of decision behaviors in individuals with NSSI behaviors.

To account for the potential heterogeneity among the participants with NSSI behaviors in our study, we classified the participants into two groups (i.e., the NSSI≥5 and NSSI<5 groups) according to the frequency criteria in the DSM-V for the NSSI disorder (24), and repeated all the analyses. The main findings were similar. Furthermore, no significant differences were observed between the two groups.

While the reduction in sample size resulting from sub-grouping might have diminished the statistical power, the findings indicate a potential link between the general risk aversion we reported and the occurrence of Non-Suicidal Self-Injury (NSSI) behaviors, at least in patients with MDD. Nevertheless, we recognize that our findings may not necessarily apply to the persistence of NSSI behaviors that can lead to significant functional impairment, which will require further investigation in a more targeted cohort of individuals with NSSI disorder.

Previous research suggests that behaviors related to risky decision-making vary among younger and older adults (31). There is also evidence that adolescents are more likely to engage in risky behaviors than adults (32,33). The age range in our study was relatively narrow but covered mid-late adolescence to emerging adults. Considering the significant (though small) differences between groups in age, we included age and its highly correlated educational level as confounding variables in supplementary modeling analysis to remove their potential impact, where our major findings still hold. Besides, the D+NSSI group in our study, despite being younger than the D group, exhibits stronger risk aversion, which is in the direction reverse to the age-related effects suggested by previous research. Therefore, the likelihood of our findings being age-driven is relatively low.

In conclusion, NSSI behaviors manifest a general aversion to risk. However, as depressive symptom severity increases, individuals perceive a decrease in environmental risks and demonstrate more unpredictable behaviors. This suggests a complex interaction between depressive symptoms and NSSI behaviors in decision-making. On one hand, overestimating risks in unfamiliar environments may diminish the likelihood of exploring alternative means to cope with emotions. On the other hand, depression amplifies engagement in risky behaviors.

ACKNOWLEDGMENTS

We extend our gratitude to all subjects who participated in this study. This study was supported by the Clinical Medicine Plus X – Young Scholars Project, Peking University the Fundamental Research Funds for the Central Universities (PKU2023LCXQ023) for HY and HZ; HY was partly supported by the National Natural Science Foundation of China (82071505), and HZ was partly supported by the National Natural Science Foundation of China (32171095), National Science and Technology Innovation 2030 Major Program (2022ZD0204803), and funding from Peking-Tsinghua Center for Life Sciences.

CONFICTS OF INTEREST

The authors have no conflicts of interest to declare.

REFERENCES

1. Lim K-S, Wong CH, McIntyre RS, Wang J, Zhang Z, Tran BX, *et al.* (2019): Global Lifetime and 12-Month Prevalence of Suicidal Behavior, Deliberate Self-Harm and Non-Suicidal Self-Injury in Children and Adolescents between 1989 and 2018: A Meta-Analysis. *IJERPH* 16: 4581.

2. Glenn CR, Klonsky ED (2013): Nonsuicidal self-injury disorder: an empirical investigation in adolescent psychiatric patients. *J Clin Child Adolesc Psychol* 42: 496–507.

3. Skegg K, Wu B, Zhang H, Chen J, Chen J, Liu Z, *et al.* (2005): Self-harm. *Gen Psych* 366: 1471–1483.

4. Haw C, Hawton K, Houston K, Townsend E (2001): Psychiatric and personality disorders in deliberate self-harm patients. *Br J Psychiatry* 178: 48–54.

5. Yan H, Zhang Y, Lu Z, Li M, Ge Y, Mei D, *et al.* (2023): High-risk group and functional subtypes of non-suicidal self-injury in young adults with mental disorders. *Front Psychiatry* 14: 1098178.

6. Barrocas AL, Jenness JL, Davis TS, Oppenheimer CW, Technow JR, Gulley LD, *et al.* (2011): Developmental perspectives on vulnerability to nonsuicidal self-injury in youth. *Adv Child Dev Behav* 40: 301–336.

7. Mars B, Heron J, Klonsky ED, Moran P, O'Connor RC, Tilling K, *et al.* (2019): Predictors of future suicide attempt among adolescents with suicidal thoughts or non-suicidal self-harm: a population-based birth cohort study. *The Lancet Psychiatry* 6: 327–337.

8. Favazza AR (1998): The Coming of Age of Self-Mutilation. *The Journal of Nervous & Mental Disease* 186: 259–268.

9. Ge Y, Xiao Y, Li M, Yang L, Song P, Li X, *et al.* (2023): Maladaptive cognitive regulation moderates the mediating role of emotion dysregulation on the

association between psychosocial factors and non-suicidal self-injury in depression. *Front Psychiatry* 14: 1279108.

10. Hasking P, Whitlock J, Voon D, Rose A (2017): A cognitive-emotional model of NSSI: using emotion regulation and cognitive processes to explain why people self-injure. *Cognition and Emotion* 31: 1543–1556.

11. Perez J, Venta A, Garnaat S, Sharp C (2012): The Difficulties in Emotion Regulation Scale: Factor Structure and Association with Nonsuicidal Self-Injury in Adolescent Inpatients. *J Psychopathol Behav Assess* 34: 393–404.

12. Wolff JC, Thompson E, Thomas SA, Nesi J, Bettis AH, Ransford B, *et al.* (2019): Emotion dysregulation and non-suicidal self-injury: A systematic review and meta-analysis. *Eur Psychiatry* 59: 25–36.

13. Nock MK, Mendes WB (2008): Physiological arousal, distress tolerance, and social problem-solving deficits among adolescent self-injurers. *Journal of Consulting and Clinical Psychology* 76: 28–38.

14. Dillahunt AK, Feldman DA, Thomas LR, Farstead BW, Frandsen SB, Lee S, *et al.* (2022): Self-Injury in Adolescence Is Associated with Greater Behavioral Risk Avoidance, Not Risk-Taking. *J Clin Med* 11: 1288.

15. Hamza CA, Willoughby T, Heffer T (2015): Impulsivity and nonsuicidal self-injury: A review and meta-analysis. *Clinical Psychology Review* 38: 13–24.

16. Platt ML, Huettel SA (2008): Risky business: the neuroeconomics of decision making under uncertainty. *Nat Neurosci* 11: 398–403.

17. Schatten HT, Andover MS, Armey MF (2015): The roles of social stress and decision-making in non-suicidal self-injury. *Psychiatry Research* 229: 983–991.

18. Wilkinson P, Kelvin R, Roberts C, Dubicka B, Goodyer I (2011): Clinical and psychosocial predictors of suicide attempts and nonsuicidal self-injury in the Adolescent Depression Antidepressants and Psychotherapy Trial (ADAPT). *Am J Psychiatry* 168: 495–501.

19. Zubrick SR, Hafekost J, Johnson SE, Sawyer MG, Patton G, Lawrence D (2017): The continuity and duration of depression and its relationship to non-suicidal self-harm and suicidal ideation and behavior in adolescents 12-17. *J Affect Disord* 220: 49–56.

20. Chai Y, Luo H, Wong GHY, Tang JYM, Lam T-C, Wong ICK, Yip PSF (2020): Risk of self-harm after the diagnosis of psychiatric disorders in Hong Kong, 2000-10: a nested case-control study. *Lancet Psychiatry* 7: 135–147.

21. Lejuez CW, Read JP, Kahler CW, Richards JB, Ramsey SE, Stuart GL, *et al.* (2002): Evaluation of a behavioral measure of risk taking: The Balloon Analogue Risk Task (BART). *Journal of Experimental Psychology: Applied* 8: 75–84.

22. Dozois DJA, Dobson KS, Westra HA (2004): The comorbidity of anxiety and depression, and the implications of comorbidity for prevention. *The Prevention of Anxiety and Depression: Theory, Research, and Practice.* Washington, DC, US: American Psychological Association, pp 261–280.

23. Turner BJ, Dixon-Gordon KL, Austin SB, Rodriguez MA, Zachary Rosenthal M, Chapman AL (2015): Non-suicidal self-injury with and without borderline personality disorder: Differences in self-injury and diagnostic comorbidity. *Psychiatry Research* 230: 28–35.

24. American Psychiatric Association (2013): *Diagnostic and Statistical Manual of Mental Disorders*, Fifth Edition. American Psychiatric Association. https://doi.org/10.1176/appi.books.9780890425596

25. Park H, Yang J, Vassileva J, Ahn W-Y (2021): Development of a novel computational model for the Balloon Analogue Risk Task: The exponential-weight mean–variance model. *Journal of Mathematical Psychology* 102: 102532.

26. Dunstan DA, Scott N (2019): Clarification of the cut-off score for Zung's self-rating depression scale. *BMC Psychiatry* 19: 177.

27. Allen KJD, Fox KR, Schatten HT, Hooley JM (2019): Frequency of nonsuicidal self-injury is associated with impulsive decision-making during criticism. *Psychiatry Research* 271: 68–75.

28. Sauder CL, Derbidge CM, Beauchaine TP (2016): Neural responses to monetary incentives among self-injuring adolescent girls. *Dev Psychopathol* 28: 277–291.

29. Wilson RC, Geana A, White JM, Ludvig EA, Cohen JD (2014): Humans use directed and random exploration to solve the explore–exploit dilemma. *Journal of Experimental Psychology: General* 143: 2074–2081.

30. Ose SO, Tveit T, Mehlum L (2021): Non-suicidal self-injury (NSSI) in adult psychiatric outpatients – A nationwide study. *Journal of Psychiatric Research* 133: 1–9.

31. Worthy DA, Gorlick MA, Pacheco JL, Schnyer DM, Maddox WT (2011): With Age Comes Wisdom. *Psychol Sci* 22: 1375–1380.

32. Defoe IN, Dubas JS, Figner B, van Aken MAG (2015): A meta-analysis on age differences in risky decision making: Adolescents versus children and adults. *Psychological Bulletin* 141: 48–84.

33. Tymula A, Rosenberg Belmaker LA, Roy AK, Ruderman L, Manson K, Glimcher PW, Levy I (2012): Adolescents' risk-taking behavior is driven by tolerance to ambiguity. *Proc Natl Acad Sci USA* 109: 17135–17140.

34. Zung WWK (1971): A Rating Instrument For Anxiety Disorders. *Psychosomatics* 12: 371–379.

35. ZUNG WWK (1965): A Self-Rating Depression Scale. *Arch Gen Psychiatry* 12: 63.

36. Dunstan DA, Scott N (2018): Assigning Clinical Significance and Symptom Severity Using the Zung Scales: Levels of Misclassification Arising from Confusion between Index and Raw Scores. *Depression Research and Treatment* 2018: 1–13.

37. Zung WWK (1973): From Art to Science. *Arch Gen Psychiatry* 29: 328.

38. Wallsten TS, Pleskac TJ, Lejuez CW (2005): Modeling Behavior in a Clinically Diagnostic Sequential Risk-Taking Task. *Psychological Review* 112: 862–880.

39. Krefeld-Schwalb A, Pachur T, Scheibehenne B (2022): Structural parameter interdependencies in computational models of cognition. *Psychological Review* 129: 313–339.

40. Olschewski S, Sirotkin P, Rieskamp J (2022): Empirical underidentification in estimating random utility models: The role of choice sets and standardizations. *Brit J Math & Statis* 75: 252–292.