Tuning of cortical color mechanism revealed using steady-state visually evoked potentials

Dylan J. Watts[1†], Ana Rozman[1†], Lucy P. Somers[1], Bora Gunel[1], Chris Racey[1], Katie Barnes[1] and Jenny M.
Bosten[1*]

[1]School of Psychology, University of Sussex, Brighton UK
[†]Joint first authors
*Corresponding author: j.bosten@sussex.ac.uk

Abstract

Color information is thought to be received by the primary visual cortex via two dominant retinogeniculate
pathways, one signals color variation between teal and red, and the other signals color variation between
violet and lime. This representation is thought to be transformed in the cortex so that there are a number
of different cell populations representing a greater variety of hues. However, the properties of cortical
color mechanisms are not well understood. In four experiments, we characterized the tuning functions of
cortical color mechanisms by measuring the intermodulation of steady-state visually evoked potentials
(SSVEPs). Stimuli were isoluminant chromatic checkerboards where odd and even checks flickered at
different frequencies. As hue dissimilarity between the odd and even checks increased, the amplitude of
an intermodulation component (I_1) at the sum of the two stimulus frequencies decreased, revealing
cortical color tuning functions. In Experiment 1 we found similar broad tuning functions for 'cardinal' and
intermediate color axes, implying that the cortex has intermediately tuned color mechanisms. In
Experiment 2 we found similar broad tuning functions for 'checkerboards' with no perceptible edges
because the checks were formed from single pixels (~0.096°), implying that the underlying neural
populations do not rely on spatial chromatic edges. In Experiment 3 we manipulated check size and found
that color tuning functions were consistent across check sizes used. In Experiment 4 we measured full 360°
tuning functions for a 'cardinal' cortical color mechanism and found evidence for opponent color
responses. The observed cortical color tuning functions were consistent with those measured using
psychophysics and electrophysiology, implying that tracking intermodulation using SSVEPs provides a
useful method for measuring them.

Color vision, cortical color mechanisms, SSVEPs, intermodulation components, tuning functions, cardinal color
axes, intermediate color axes

1.0 Introduction
Decades of research in the neural mechanisms of human color vision have produced a relatively good
understanding of the representation of color at a receptor level, and in the retinogeniculate color pathways.
However, the cortical mechanisms underlying color perception are much less well understood, and though
color representation in early cortical areas has been studied using electrophysiology in animal models, there
has been relatively little research on the properties of cortical color mechanisms in humans measured directly
using neuroimaging. In particular, there is a debate over how far the separation of color information into
distinct retinogeniculate pathways remains segregated in the cortex (Schluppeck and Engel, 2002;
Gegenfurtner, 2003; Kaneko et al., 2020; Nunez et al., 2021), and over the number and tuning of cortical
color mechanisms (Gegenfurtner, 2003; Emery et al., 2017). We used a novel approach using steady-state
visually evoked potentials (SSVEPs) to measure and characterize the tuning functions of cortical color
mechanisms in humans.

Normal human color vision begins with the relative activations of three types of retinal cone photoreceptors,
sensitive to short (S), medium (M) or long (L) wavelengths of light. Cone signals are then thought to be
combined postreceptorally by distinct populations of bipolar cells, setting up 'cardinal' color pathways which
remain segregated in the retina and LGN and at least as far is the input layers of V1 (Gouras, 1974; Chatterjee
and Callaway, 2003; Li et al., 2022). The S/(L+M) pathway carried by small bistratified ganglion cells and
though koniocellular layers of the LGN signals color differences between lime green and violet. The L/(L+M)

pathway carried by the midget retinal ganglion cells and through parvocellular layers of the LGN signals color differences between teal and red (Solomon and Lennie, 2007; Conway et al., 2018).

To what extent retinogeniculate opponent color representations in early visual cortical areas are retained and to what extent they are transformed is not fully understood. Although early psychophysical evidence was consistent with a 'cardinal' representation of color inherited from the retinogeniculate mechanisms (Boynton and Kambe, 1980; Krauskopf et al., 1982; Krauskopf and Farell, 1990), later evidence favored a 'higher order' color representation based on color mechanisms responsive to hues that are intermediate between the poles of the retinogeniculate color mechanisms (Krauskopf et al., 1986; Webster and Mollon, 1991; Gegenfurtner and Kiper, 1992; Krauskopf and Gegenfurtner, 1992; Hansen and Gegenfurtner, 2013). There has also been fMRI evidence in support of both a 'cardinal' color representation in V1 (Engel et al., 1997) and in support of the existence of intermediately tuned color mechanisms (Goddard et al., 2010; Kuriki et al., 2015).

Studies aiming to measure the tuning functions of color selective cortical neurons have been conducted using indirect psychophysical methods in humans and directly using electrophysiology in primates. Results of psychophysical studies have typically shown broad tuning (Sankeralli and Mullen, 1997; Lindsey and Brown, 2004) especially when 'off-axis looking' (stimulus detection by neighboring color mechanisms than the one being measured) is minimized (D'Zmura and Knoblauch, 1998; Hansen and Gegenfurtner, 2006). Electrophysiological recordings in primates have revealed diversity in the preferred hues of color selective neurons in V1 (Lennie et al., 1990; Hanazawa et al., 2000; Hass and Horwitz, 2013), V2 (Kiper et al., 1997; Solomon and Lennie, 2005), V3 (Gegenfurtner et al., 1997), V4 (Bohon et al., 2016) as well as in downstream color selective areas in the ventral visual pathway (Komatsu et al., 1992; Conway et al., 2007). Color tuning functions of individual color-responsive cells measured electrophysiologically have been found to have a range of half-amplitude bandwidths in hue angle from about 30°-170° in V1 (Wachtler et al., 2003) and 20°-85° in V2 (Kiper et al., 1997). The average bandwidth of color tuning functions tends to decrease through the hierarchy of cortical visual areas (Gegenfurtner, 2003), but large populations of broadly tuned neurons are still found in V4 (Bohon et al., 2016).

The color tuning properties of visual cortical neurons can of course interact with tuning properties to other stimulus features. For instance, many cortical neurons are parallelly tuned to both color and luminance (Johnson et al., 2001; Schluppeck and Engel, 2002), spatial frequency (Lennie et al., 1990) or orientation (Leventhal et al., 1995; Friedman et al., 2003; Garg et al., 2019). One distinction that has been relatively well studied is between so-called single- and double color opponent neurons (Shapley and Hawken, 2011). Double opponent neurons respond to stimuli containing sharp chromatic edges but respond weakly or not at all to full-field or low spatial frequency chromatic stimuli, while single opponent neurons show the opposite pattern (Livingstone and Hubel, 1984; Johnson et al., 2008). Depending on the spatial properties of the chosen stimuli, color tuning results may therefore vary, because different populations of cells may be targeted that are tuned to combinations of color and other stimulus features. In particular, the stimuli with chromatic edges may preferentially activate double opponent neurons, while stimuli without edges may preferentially activate single opponent neurons.

In humans, other than by inference from the results of psychophysical methods, there is one existing study by Chen et al. (2021) that measured cortical color tuning functions using neuroimaging, specifically by measuring steady -state visually-evoked potentials (SSVEPs). SSVEPs occur in response to visual stimuli that are flickering at a precise temporal frequency (Regan, 1966; Norcia et al., 2015), and are typically analyzed in the frequency domain where EEG noise is segregated into many frequency bands, providing high signal to noise for brain responses at the stimulus frequency (Rossion and Boremanse, 2011) and a sensitive index of visual responses in the cortex. Chen et al. (2021) used a method for measuring cortical color tuning functions based on chromatic noise masking, where psychophysical results have shown that the strength of masking increases with the color similarity of the target and mask (Gegenfurtner and Kiper, 1992; D'Zmura and

103 Knoblauch, 1998). They presented central binary chromatic noise targets with surround binary chromatic
104 noise masks, and measured occipital SSVEP amplitudes as a function of the angular difference between the
105 color axes from which target and surround colors were sampled, finding broad color tuning functions
106 centered at both cardinal and intermediate color axes.
107
In the current study we exploited intermodulation of SSVEPs as an alternative scheme for measuring cortical
tuning functions (Regan and Regan, 1987). Intermodulation components in SSVEP occur when there are two
or more stimuli flickering at different frequencies presented simultaneously, and are thought to provide an
index of the degree to which shared neural resources process the different stimuli (Regan and Regan, 1989;
Norcia et al., 2015). We presented flickering checkerboard stimuli where odd checks flickered at one
frequency and remained at a fixed chromaticity across trials, and even checks flickered at a different
frequency and varied in chromaticity. We measured SSVEP amplitudes at the intermodulation frequency,
which we expected to be maximal when odd and even checks have the same chromaticity and are therefore
processed by the same color mechanisms, and to decrease with the color difference between odd and even
checks. In a series of four experiments, we measured cortical color tuning functions at cardinal and
intermediate color axes (Experiment 1), when the stimulus lacked perceptible chromatic edges (Experiment
2), as a function of the size of checkerboard checks (Experiment 3), and around the full hue circle (Experiment
4). Our results reveal that bandwidths of cortical color tuning functions do not differ between cardinal and
intermediate axes, implying the action of intermediately tuned color mechanisms; that surprisingly, tuning
functions do not depend on the presence of perceptible chromatic edges, implying that they probe color
mechanisms with low spatial opponency.
124
125 **2.0 General Methods**
126 Provided in this section are methods common to all four experiments.
127
2.1 Participants
Participants in all experiments had normal color vision assessed using an 'Anomaloskop' anomaloscope
(Oculus, Wetzlar, Germany; Experiments 1-3) or the Ishihara Plates Test (Experiment 4), and were recruited
via email and word-of-mouth. Participants provided written informed consent and were compensated £10
per hour for their time. The study received ethical approval from the Sussex University Science and
Technology Cross Schools Research Ethics Committee (ER/DJW41), and adhered to the tenants of the World
Medical Association's Declaration of Helsinki (2013) with the exception that it was not pre-registered.
135
2.2 Equipment
Electroencephalographic data were acquired using 64-channel Waveguard caps (ANT neuro, Hengelo,
Netherlands) via a high-speed 64-channel ANT Neuro amplifier sampling at 1000 Hz. EEG data were
recorded in ASA lab (ANT Neuro, Hengelo, Netherlands). Additional electrodes were used to measure
horizontal (HEOG) and vertical (VEOG) eye-movements to allow for the removal of blink artefacts. A custom
photodiode input to the EEG amplifier was fitted to check for accurate stimulus timing (see Supplementary
Information 7).
143
Stimuli were displayed on a GDM FW900 CRT monitor (Sony, Tokyo, Japan). The monitor was gamma-
corrected using a luminance meter (Konica-Minolta, Tokyo, Japan) and color calibrated using a PR650
SpectraScan spectroradiometer (PhotoResearch, Dorking, Surrey). Stimuli were presented using a ViSaGe
MKII Stimulus Generator (Cambridge Research Systems, Rochester, UK) using Matlab 2017 (The Mathworks,
Nantick, CA). Triggers to the EEG amplifier from the ViSaGe via BNC and parallel port connections. The
monitor had a resolution of 800 x 600 pixels and ran at a frame rate of 160 Hz.
150
151 *2.3 Heterochromatic flicker photometry*
152 Heterochromatic flicker photometric measurements were made in order to calculate stimuli that are
153 isoluminant for individual observers. Stimuli consisted of a 2D annulus with an outer diameter of 41.9° and

inner diameter of 5.7°, which oscillated at 11.4 Hz between two colors (Figure 1). On each trial, the intensity of one color was fixed, and participants adjusted the intensity of the other color until they perceived minimum flicker (the point of isoluminance). On the basis of these measurements, stimuli in the four experiments were made isoluminant to ensure that they isolated chromatic mechanisms and did not stimulate luminance mechanisms.

Figure 1. Stimulus properties. (a) Stimuli were isoluminant chromatic flickering checkerboards. Even checks flickered at F_A (6.66 Hz) between equal energy white and a 'variable' chromaticity that depended on the color condition. Odd checks flickered at F_B (8 Hz) between equal energy white and a 'fixed' chromaticity. A central 5.7° disc was metameric with equal energy white and did not flicker. (b) Stimulus chromaticities specified in a version of the MacLeod-Boynton (1979) chromaticity diagram based on the Stockman, MacLeod and Johnson (1993) cone fundamentals. For Experiment 1 the fixed chromaticity was either the S/(L+M) axis in a decremental direction (left panel), an intermediate axis (middle panel) or the L/(L+M) axis in an incremental direction (right panel). In each panel, the modulation made by the fixed chromaticity and equal energy white is indicated by the solid line. For each fixed color axis there were 6 color conditions, where the variable chromaticity differed from the fixed chromaticity in steps of 18° from 0° to 90°. Modulations between variable chromaticities and equal energy white are indicated by the dotted lines. For Experiments 2 and 3 the S/(L+M) axis in a decremental direction was tested (left panel). For Experiment 4 the fixed chromaticity was along the S/(L+M) axis in a decremental direction (left panel) and there were 12 color conditions in 30° steps between 0° and 300° from the fixed chromaticity (not shown).

2.4 Stimuli

The stimuli for all four experiments consisted of flickering checkerboards extending 76.8° horizontally and 48° vertically (Figure 1a), with a viewing distance of 30 cm. Each check was square and measured 4.8° x 4.8°. Even checks flickered between equal energy white and a fixed hue at 8 Hz, eliciting SSVEPs at 8 Hz and higher harmonics. Odd checks flickered between equal energy white and a variable hue at 6.66 Hz, eliciting SSVEPs at 6.66 Hz and higher harmonics. Flicker was square wave between equal energy white and the edge of the maximum saturation circle allowed by the monitor's gamut (Figure 1b). There was a central disc metameric with equal energy white of 5.7° with a central fixation point which did not flicker, to avoid the macular pigment (Figure 1a).

2.5 Procedure

Participants first completed heterochromatic flicker photometry task. The consistency of intensity settings was assessed, and if found to range >20%, the task was repeated. Participants were then fitted with facial electrodes and an EEG cap in accordance with the 10/20 system (Jurcak et al., 2007). We ensured that impedances for electrodes of interest were < 10 kΩ and for other electrodes < 20kΩ. In the experiment, participants viewed a series of 10 second trials (see details of methods for each separate experiment). They were instructed not to blink excessively during trials and to blink between trials. Participants controlled the inter-trial interval by pressing a button on a keyboard when they were ready for the next trial. Raw EEG data were monitored in real time for signs of fatigue or other issues.

2.6 EEG Analysis: pre-processing

Pre-processing was conducted in MATLAB, using Letswave 7 (https://nocions.github.io/letswave7/) for Experiments 1 and 4, and Letswave 6 for Experiments 2 and 3 (https://nocions.github.io/letswave6/). First, raw EEG data were imported and mapped to the ANT neuro standard 64-channel electrode coordinate system. Raw data were visually inspected for excessive noise resulting from equipment failures. Problematic channels were removed. No exclusions of channels were made from our analysis cluster, consisting of Oz, O1, O2, Pz, P1, P2, P3, P4, POz, PO3, and PO4 (Figure 2d). Overview of tuning functions at other posterior and occipital channels is presented in Supplementary Information 5 (Supplementary Figure 6), demonstrating signal transitivity across the area. Signals were re-referenced to the grand average of all remaining cortical electrodes. In addition to our analysis cluster, channels FPz, HEOG and VEOG were extracted to detect eye movements. Consistency in temporal profile of our stimuli was ensured by inspecting the decomposed frequency profile of the photodiode which was attached to the screen (see Supplementary Information 7). A Butterworth filter for 50 Hz noise and a high-pass filter (<0.1 Hz) were applied. Data were then segmented into trials using an epoch size of 9000 s, which included an integer number of stimulus repetitions at both flicker frequencies. Segmented trials were baseline corrected within the 1 to 1.2 interval. No ocular artefacts were removed in Experiments 1 and 4 (see Supplementary Information 2 for a comparison of signal amplitudes with and without ocular artefact removal). For Experiments 2 and 3, ocular artefacts were removed by visual inspection if there were fewer than 2 blinks per color condition. Alternatively, Independent Components Analysis (ICA) was applied if there were two or more blinks per condition. Recordings for an individual observer were merged in cases where data was collected across multiple sessions. Trials from the same condition for each observer were then averaged and converted to the frequency domain by fast Fourier transformation (FFT). Amplitudes for each observer and each color condition were extracted at I_1 (14.66 Hz), which is an intermodulation component at the sum of the two stimulus frequencies. Amplitudes were also extracted at F_B (8 Hz), the frequency of the checks of fixed chromaticity – changes in amplitude with the chromaticity of the F_A (6.66 Hz) checks reflects chromatic masking, which has previously been used to infer cortical color tuning functions (Chen and Gegenfurtner, 2021).

3.0 Experiment 1

In Experiment 1 we sought to measure cortical color tuning functions using SSVEP, and to determine whether there are differences in cortical color tuning functions between 'cardinal' color axes aligned with the retinogeniculate color channels L/(L+M) and S/(L+M) and an intermediate color axis.

3.1 Experiment 1: Methods

Eleven participants (4 female) completed two experimental sessions, each lasting 2.5 hours. Three color axes were tested: L/(L+M) increments, S/(L+M) decrements and an intermediate axis (Figure 1b). For each axis, the chromaticity of checks flickering at F_A was fixed along that axis, and the chromaticity of checks flickering at F_B varied on each trial between six hue angles to the fixed axis (0°, 18°, 36°, 54°, 72° and 90°; Figure 2a). There were 180 trials in each session presented in a random order, totaling 360 trials.

Figure 2. Experiment 1 results. (a) Amplitude profile for an example condition (S/(L+M) axis, 0° variable chromaticity), averaged over the 11 participants. (b) SNR profile for the same condition. Fundamental stimulus frequencies and their harmonics are labelled in black and intermodulation components in red. (c) Mean amplitude of the intermodulation component at 14.66 Hz as a function of the hue angle between the fixed and variable chromaticities for the S/(L+M) axis (green solid line), the intermediate axis (orange dotted line) and the L/(L+M) axis (red dashed line). Error bars are 95% confidence intervals. (d) Mean amplitude at the stimulus frequency of the fixed chromaticity at 8 Hz as a function of the hue angle between the fixed and variable chromaticities for the S/(L+M) axis (green solid line), the intermediate axis (orange dotted line) and the L/(L+M) axis (red dashed line). Error bars are within-participant 95% confidence intervals (O'Brien and Cousineau, 2015), jittered along the x-axis for visibility. (e) Schematic representation of analysis cluster electrodes.

3.2 Experiment 1: Results

Strong signals were observed at the two fundamental frequencies, their harmonics ($F_A(1)$ and $F_B(1)$), and at I_1 (e.g. Figures 2a and b). There were additional amplitude peaks at other harmonics of F_A and F_B and at other intermodulation frequencies. We extracted I_1 and F_B for further analyses.

Figure 2 shows the effect of hue dissimilarity between odd and even checks on SSVEP amplitudes at I_1 (panel c) and F_B (panel d) for the three color axes. Tuning functions were similar in shape for all three axes, with a broad half-amplitude bandwidth of approximately 90°. Two-way repeated-measures ANOVAs were conducted to test the effect of color axis and hue angle differences on the amplitudes of I_1 and F_B. Mauchly's test of sphericity was violated for both frequencies ($X^2(54) = 106.50$ for I_1 and $X^2(54) = 115.40$ for F_B, $p <$

.001), therefore Greenhouse-Geisser corrected tests were used (ε =.311 for I_1 and .260 for F_B). The ANOVAs
showed that for both I_1 and F_B there were significant main effects of hue angle difference ($F(1.19,11.92) =$
21.59 for I_1 and $F(1.84,15.49) = 23.18$ for F_B, $p <.001$, $\eta^2_p =.683$ for I_1 and $\eta^2_p = .699$ for F_B). Results from post-
hoc tests (Supplementary Information 1, Supplementary Tables 1 and 2) show that mean SSVEP amplitude
decreased significantly with hue angle difference for I_1 and increased significantly with hue angle difference
for F_B.
263
There was a significant main effect of color axis for F_B, ($F(1.14, 11.36) = 10.83$, $p = .006$, $\eta^2_p =.520$), but not
for I_1 ($F(1.45,14.51) =3.17$, $p =.084$, $\eta^2p =.241$). Post-hoc tests using Bonferroni correction indicated that there
was a significant difference between L/(L+M) and S/(L+M) axes ($M = -.247$, $p < .05$). There was no significant
interaction between axis and hue angle difference either for I_1 ($F(3.11,31.06) = 2.74$, $p = .058$) nor for F_B
($F(2.60, 26.03) = 2.40$, $p = .098$). Thus, the effect of hue angle on SSVEP amplitude was consistent across all
three axes for both frequencies, implying that tuning functions were shaped similarly for all three axes.
270

271
272 **Figure 3. Results of Experiment 2. SSVEP amplitude is plotted as a function of hue angle dissimilarity for single-pixel**
273 **checkerboard stimuli. (a) for I_1 at 14.66 Hz. (a) for F_B at 8 Hz. Error bars are within-participant 95% confidence intervals**
274 **(O'Brien and Cousineau, 2015).**
275
4.0 Experiment 2
The isoluminant checkerboard stimuli in Experiment 1 have strong chromatic edges. The S/(L+M) axis, and
thus neural populations responding to chromatic edges may underlie the results. To test the effect of
chromatic edges on cortical color tuning functions measured using SSVEP we reduced the size of the 'checks'
to single pixels (invisible to participants). If tuning functions change from those measured in Experiment 1
when the checkerboard is constructed from single pixels, then the tuning functions measured in Experiment
1 are attributable to neural populations responsive to chromatic edges.
283
4.1 Experiment 2: Methods
Eleven participants (6 female) completed a single 2.5-hour experimental session. Stimuli consisted of single
pixel checkerboards (.096° x .096°), with the same spatial organization as depicted in Figure 1a. Individual
checks were indistinguishable to the observer. The chromaticity of checks flickering at F_B was fixed, and the
chromaticity of checks flickering at F_A varied between six hue angles to the fixed chromaticity (0°, 18°, 36°,
54°, 72° and 90°). There were 120 experimental trials, where hue conditions were presented in a random
order. Since the S/(L+M) axis produced somewhat larger amplitudes in Experiment 1 than the other two axes,
Experiment 2 probed the S/(L+M) axis only.
292
4.2 Experiment 2: Results

Figure 3 shows the amplitudes of I_1 and F_B against the difference in hue angle between odd and even single
pixel checks. One-way repeated-measures ANOVAs examined the effect of hue angle difference on the
amplitudes of I_1 and F_B. Mauchly's test of sphericity was violated for both frequencies ($X^2(14)$ = 72.16 for I_1
and $X^2(14)$ = 34.53 for F_B, p <.001), therefore Greenhouse-Geisser corrected tests were used (ε = .279 for I_1
and ε = .510 for F_B). There was a significant effect of hue angle difference on signal amplitude for I_1 ($F(1.40,$
13.96) = 9.56, p =.005, η^2_p = .489), but not for F_B ($F(2.55, 25.48)$ = .465, p = .679, η^2_p =.044). Results from post-
hoc tests (see Supplementary Information 1, Supplementary Tables 3 and 4) indicated that mean SSVEP
amplitude decreased with hue angle difference for I_1, but did not change with hue angle difference for F_B.
302
303 **5.0 Experiment 3**
304 Since we observed a difference in the shape of the tuning function for F_B between the 4.8° checks presented
305 in Experiment 1 and the indivisible 0.096° checks presented in Experiment 2, in Experiment 3 we
306 systematically manipulated check size. We measured tuning functions at I_1 and at F_B at 4 check sizes, varying
307 between the check sizes used in Experiments 1 and 2.
308
5.1 Experiment 3: Methods
Eleven participants (5 female) completed two 2.5-hour experimental sessions. There were 4 check size
conditions: 0.096° (as in Experiment 2), 0.4°, 1.3° and 4.8° (as in Experiment 1). To limit the duration of testing
sessions there were four hue angle difference conditions rather than the 6 used in Experiments 1 and 2.
These were 0°, 30°, 60° and 90°. Checks flickering at F_B (8 Hz) at a fixed chromaticity modulated between
equal energy white and a decrement along S/(L+M) axis (left panel of Figure 1b). 2 testing sessions were
conducted separated by at least one day. There were 160 experimental trials per session (10 trials for each
combination of check size and hue angle difference), totaling 320 trials (20 trials in total for each stimulus
type).
318

319
320 **Figure 4. Results of Experiment 3. SSVEP amplitude is plotted as a function of hue angle dissimilarity (x-axis) and check**
321 **size (legend). (a) for I_1 at 14.66 Hz. (b) for F_B at 8 Hz. Error bars are within-participant 95% confidence intervals (O'Brien**
322 **and Cousineau, 2015), jittered along the x-axis for visibility.**
323
5.2 Experiment 3: Results
Figure 4 shows the effect of check size and hue angle difference on amplitudes at I_1 and F_B. The figure shows
that for I_1 check size does not affect the shape of the tuning functions observed. However, F_B, in agreement
with the results of Experiment 2 there is no tuning for the single pixel checkerboard, but similar tuning
functions as observed in the results of Experiment 1 for the larger check sizes.

We conducted two-way repeated-measures ANOVAs to test the effect of hue angle hue angle difference and
check size on the amplitudes at I_1 and F_B. Mauchly's test of sphericity was violated for both frequencies ($X^2(44)$
= 81.52, p = .002 for I_1; $X^2(44)$ = 134.05, $p < .001$ for F_B). Greenhouse-Geisser corrected tests were therefore
used (ε = .367 for I_1; ε = .251 for F_B). There were significant main effects of check size and hue angle on signal
amplitude for I_1 ($F(2.32, 23.21) = 3.61$, p= .038, η^2_p = .265 for check size; $F(1.24, 12.39) = 33.76$, p <.001, η^2_p
=.771 for hue angle difference), but no significant interaction ($F(3.30, 33.01) = 1.55$, p =.218, η^2_p =.134). Post-
hoc tests (Supplementary Information 1, Supplementary Tables 5 and 6) indicated that mean amplitude in
Condition 3 was somewhat stronger than in other conditions, regardless of hue angle, and that mean SSVEP
amplitude decreased with hue angle for I_1, regardless of check size. For F_B there was a significant main effect
of hue angle difference ($F(1.27, 12.67) = 11.30$, p = .004, η^2_p =.530), but there was no significant main effect
of check size ($F(1.51, 15.06) = .428$, p =.604, η^2_p =.041). Results from post-hoc tests (Supplementary
Information 1, Supplementary Tables 5 and 6) indicated that mean SSVEP amplitude increased with hue angle
difference. There was a significant interaction between check size and hue angle for F_B ($F(2.26, 22.58) = 3.39$,
p = .046, η^2_p =.253). Post-hoc analysis (Supplementary Information 1, Supplementary Table 6) indicated that
the shape of the tuning function for F_B differed between check size conditions.

6.0 Experiment 4
The tuning functions observed in the results of Experiments 1-3 are broadband but do not reach an
asymptote at the largest hue angle differences we measured of 90°. In Experiment 4 we aimed to characterize
a tuning function for a cortical color mechanism around the entire hue circle. We kept the fixed chromaticity
checks along the S/(L+M) axis in a decremental direction but varied the variable chromaticity checks around
the full hue circle in 12 steps.

6.1 Experiment 4: Methods
Ten participants (7 female) completed Experiment 4 in one experimental session lasting about 3 hours.
Stimuli were isoluminant flickering checkerboards with a check size of 1.3° (which elicited the largest signals
in Experiment 3). The fixed chromaticity flickering at F_B (8 Hz) was a decrement on the S/(L+M) axis (first
panel of Figure 1b). The chromaticity of the variable checks flickering at F_A (6.66 Hz) spanned the full hue
circle in steps of 30° (hue angle differences ranged from 0° to 330°). Each of the 12 hue angle conditions was
presented 20 times, totaling 240 trials. SSVEP amplitudes were measured at I_1 and at F_B.

360
361

362
363
364 **Figure 5. Results of Experiment 4. (a) tuning functions plotted at I_1 (full line) and F_B (dashed line), where the fixed**
365 **chromaticity was an S/(L+M) decrement (180°), and the variable chromaticity varied around the full hue circle. Error**
366 **envelopes are within-participant 95% confidence intervals (O'Brien and Cousineau, 2015). Tuning functions for**
367 **decrement in S/(L+M) mechanism comparing Experiment 4 results with those from Experiments 1-3 are plotted in**

370
371 *6.2 Experiment 4: Results*
372 Results from Experiment 4 are presented in Figure 5a. There is a peak in I$_1$ centered at 0° hue angle difference,
373 as expected from the results of Experiments 1-3. However, there is a second smaller peak centered at a hue
374 angle difference of 180°. For F$_B$ there is a minimum at a hue angle difference of 0° as expected from the
375 results of Experiments 1 and 3. There is no evidence of a second minimum at a hue angle difference of 180°.

376
One-way repeated measures ANOVAs were used to investigate the amplitude differences between hue angle
difference conditions at I$_1$ and at F$_B$. There was a significant effect of hue angle difference on amplitudes at
I$_1$ ($F(11, 99) = 18.065$, $p < .001$, $\eta^2_p = .667$) and at F$_B$ ($F(11, 99) = 9.851$, $p < .001$, $\eta^2_p = .523$). Post hoc
comparisons (Supplementary Information 1, Supplementary Tables 7 and 8) showed that amplitudes at I$_1$ for
a 0° hue angle difference were significantly higher than amplitudes at all other angles, except for 330°. The
amplitude for a hue angle difference of 180° was not significantly different from those of its most proximal
hue angle difference conditions, 150° and 210°.

385
386 **Figure 6. Signal heatmaps and tuning functions at intermodulation components surpassing 2 SNR in Experiment 4.**
387 **SNR is presented as a function of Frequency in Hz with a range of 0 – 25. The components and their sources (nature**

388   **of interaction between F_A and F_B) are labelled above individual heatmap and tuning function plots. The individual**
389   **heatmaps and tuning function plots are scaled to match in area under the curve.**
390
7.0 Analyses of additional intermodulation components
It is thought that different intermodulation components may arise from separate subpopulations of neurons
(Regan, 1966; Appelbaum et al., 2008; Gordon et al., 2019). For example, the subpopulation of neurons
responding to the difference between F_A and F_B might be separate from the population responding to the
sum of F_A and F_B, even though they both arise from interactions to the same signals (F_A and F_B). These
potentially different populations may also demonstrate distinct color tuning and may be differently spatially
distributed across the cortex. We therefore conducted additional analyses of the data from Experiments 1
and 4 to plot tuning functions for other intermodulation components and visualize their locations in cortical
electrode maps. Firstly, we extracted SNR profiles for a grand mean across observers in the 0° hue difference
condition (where intermodulation components are expected to be greatest). Intermodulation components
(i.e., those where the frequency peak is composed of either an addition or subtraction of multiples of F_A and
F_B) were extracted when the SNR peak surpassed a criterion of 2. This procedure identified 2 qualifying
components in Experiment 1 (Supplementary Information 4, Supplementary Figure 5) and 12 qualifying
components in Experiment 4 (Figure 6). One response at 4 Hz (labelled orange in Figure 6) could either be an
intermodulation component ($3F_B - 3F_A$) or a subharmonic of F_B ($F_B/2$). Since we found no peak at the
equivalent subharmonic of F_A, this peak is more likely to be an intermodulation component. Tuning functions
were calculated for the channels in the analysis cluster and visualized using scaled line plots. Tuning functions
for different intermodulation components had different shapes (e.g., compare tuning functions for F_A+F_B and
for $2F_A+2F_B$). However, since the criterion SNR was applied to the group average, data from some participants
had high noise, impacting the group average tuning functions.
411
Spatial signal distributions were visualized using amplitude scalp heatmaps with the grand mean across
observers, using a relativized scale between frequencies so that the maximum values are always plotted in
yellow and minimum values in dark blue. Comparison of the cortical surface heatmaps for different
intermodulation components revealed that their locations are broadly consistent, both for Experiment 1 and
Experiment 4. However, we found some substantial differences in tuning functions for different
intermodulation components. In results for Experiment 4, Some functions show a *reduction* in amplitude for
the opponent hue (e.g., 26.66 Hz and 28 Hz), in contrast to the characteristic peak observed in I_1. At the 20
Hz intermodulation component, tuning appears to be reversed, with a higher relative peak for the opponent
compared to the proximal hues (Figure 6). Overall, peaks are less pronounced for intermodulation
components other than I_1, but nonetheless may reveal different color tuning properties for different neural
subpopulations.
423
8.0 Discussion
Our results show that human cortical color tuning functions can be effectively measured by capturing
intermodulation of SSVEPs in EEG. We separately frequency tagged two colors spatially interleaved in
checkerboard patterns and extracted signals at the intermodulation frequency (I_1). As predicted, SSVEP
amplitudes at I_1 significantly and systematically reduced with hue angle dissimilarity. All of the tuning
functions we measured by extracting I_1 were broad, with half-amplitude bandwidths of about 90° in hue
angle. This bandwidth is concordant with existing reported bandwidths of color tuning functions measured
psychophysically (Sankeralli and Mullen, 1997; Lindsey and Brown, 2004) and electrophysiologically
(Wachtler et al., 2003). The results of Experiment 1 showed similar tuning functions for cardinal and
intermediate color axes. The results of Experiments 2 and 3 showed that the shape of color tuning functions
did not depend on whether or not the stimulus contained perceptible edges or the sizes of checkerboard
checks in the stimulus. The results of Experiment 4 showed that tuning functions around the whole hue circle
exhibit a bipolar shape with similar 90° half-amplitude bandwidths at both peaks. We interpret each of these
findings in turn.
438

To assess the independence or interaction between color representations along 'cardinal' retinogeniculate
color axes and intermediate color axes, in Experiment 1 we measured tuning functions centered on both
cardinal axes and on an intermediate axis. If colored stimuli specified along an intermediate axis are
represented by a combination of activities in the two cardinal mechanisms, then we would expect observed
'tuning functions' along the intermediate axis to have a larger bandwidth and a smaller peak amplitude than
tuning functions for stimuli with equivalent contrast along the cardinal axes. However, in the results of
Experiment 1 we found very similar peak amplitudes and bandwidths for tuning functions centered along an
intermediate axis as for the cardinal axes, implying the action of an independent intermediately tuned
cortical color mechanism. How are the cardinal retinogeniculate color mechanisms combined in the cortex
to produce intermediate color mechanisms? Gegenfurtner and colleagues have suggested that broad tuning
with a half-amplitude bandwidth greater than 30° typically results from a linear combination of cone-
opponent signals (Hansen and Gegenfurtner, 2006; Chen and Gegenfurtner, 2021). However, nonlinear
combinations of cone-opponent signals could also result in broad tuning functions along intermediate axes
depending on how the inputs are combined. In their 'hue sweep' SSVEP study, Kaneko et al. (2020) found
that SSVEP amplitudes in response to colors along intermediate axes could be greater than SSVEP amplitudes
in response to colors along the cardinal axes, which they interpreted as strong evidence that intermediately
tuned color mechanism result from nonlinear combinations of the cardinal mechanisms. In our Experiment
1 results, the SSVEP amplitudes of I_1 along the intermediate axis were approximately midway between those
of the two cardinal axes. This pattern of results, according to Kaneko et al. is in favor of a nonlinear
combination of cardinal mechanisms. But the results of both studies could also be explained by linear
combination followed by independent gain control for each second stage color mechanism. A linear
combination of cardinal mechanisms to form intermediate mechanisms would be in accordance with recent
optical imaging evidence for linear combination in macaque V1 (Li et al., 2022).
462
To differentiate the contributions of single- and double-opponent neurons to the cortical color tuning
functions that we measured in Experiment 1, in Experiments 2 and 3 we manipulated chromatic edges in the
stimulus. In Experiment 2, our 'single pixel' checkerboard stimulus had no perceptible chromatic edges and
should therefore excite only single-opponent neurons. We found, surprisingly, that tuning functions based
on I_1 had the same shape and bandwidth as the tuning function based on the large checks with chromatic
edges in Experiment 1. In Experiment 3 we manipulated check size systematically over 4 levels, and found no
effect on the bandwidth of I_1 with check size. We found that check sizes of 1.3° generated slightly higher
amplitudes than the other check sizes. However, the lack of significant interaction between check size and
hue angle dissimilarity suggests that the larger amplitudes could result from responses by other spatial-
frequency tuned neural populations that are not hue-angle specific. As well as SSVEP amplitudes at I_1 we also
extracted amplitudes at F_B, the frequency (8 Hz) at which the 'fixed chromaticity' was modulated. Chen and
Gegenfurtner (2021) had previously plotted cortical color tuning functions using this frequency, reasoning
that amplitudes reduce with chromatic masking if the underlying color channel is sensitive to both
modulating colors. We found, in contrast to results at our main target I_1, that tuning functions plotted using
F_B did depend on chromatic edges. For the larger checks used in Experiments 1 and 3 (similar to those used
by Chen and Gegenfurtner), tuning functions derived from amplitudes at the frequency of the fixed
chromaticity F_B were roughly the inverse of tuning functions derived from I_1, implying that both reveal the
action of the same chromatic mechanisms. However, in Experiments 2 and 3, the shapes of the observed
tuning functions derived from F_B differed substantially from those derived from I_1, suggesting that they arise
from distinct chromatic mechanisms. For our single pixel checkerboard stimuli, tuning functions derived from
I_1 matched those measured using larger check sizes, but tuning functions derived from F_B were flat. The
mechanism revealed by F_B is therefore dependent on spatial chromatic edges, but the mechanism revealed
by I_1 is not. The neural populations underlying F_B-derived tuning functions may be double-opponent while
those underlying I_1-derived tuning functions may be single-opponent.
487
In Experiments 1-3 the chromaticity of the "variable" checks was manipulated between hue angle differences
of 0° and 90°. To measure the shape of a color tuning function around the full hue circle, in Experiment 4 we

manipulated the chromaticity of the variable checks between hue angle differences of 0° and 360°.
Surprisingly, we found a second peak in I_1 at a hue angle difference of 180°, where the hue of the variable
chromaticity was opponent to the hue of the fixed chromaticity. The second peak had about half the
amplitude of the primary peak centered at a hue angle difference of 0°. We did not observe this pattern for
tuning functions based on F_B, where we observed a single minimum at 0° and a single maximum at 180°.
What could explain this pattern of findings? We assume, as argued by Chen and Gegenfurtner (2021) that
tuning functions plotted by extracting F_B are based on chromatic masking. Is chromatic masking more
effective for colors 90° distinct from the target than for colors 180° distinct from the target? We have not
identified any psychophysical data that could address this hypothesis, but it seems unlikely that this would
be the case: at 90° masking should be minimal as mask and target chromaticities are carried by orthogonally
tuned color mechanisms. However, target signals will also be influenced by simultaneous contrast, where a
surround chromaticity pushes perception of a target towards the surround's complementary color (Ekroll
and Faul, 2012). Simultaneous color contrast induced by surround S-cone increment checks would enhance
color signals from S-cone increment checks, but simultaneous color contrast induced by L/(L+M) increment
or decrement checks would not affect color signals from S-cone increment checks as it would act orthogonally
to the S-cone axis. The single peak in tuning for F_B could therefore be explained by a combination of masking
and simultaneous contrast.

For tuning functions plotted by extracting I_1, the second peak in amplitude at a hue angle difference of 180°
must arise when the same neural populations are driven by both checks. The checks with fixed color
modulated in chromaticity between an S-cone decrement and equal energy white. The variable color checks
at a 180° hue angle difference modulated between equal energy white and an S-cone increment. One
possibility is that a color mechanism with bipolar tuning (Conway et al., 2002; Chatterjee and Callaway, 2003)
responds to both sets of checks, leading to the second intermodulation peak observed at a hue angle
difference of 180°. A second possibility is that the intermodulation could arise within rectified S-cone on and
off channels, which would be consistent with multiple unipolar channels for color in the cortex (Malkoc et
al., 2005; Emery et al., 2017, 2023). S-cone off channels would respond to the onset of S-cone decrement
checks (at F_A) and to the offset of S-cone increment checks (at F_B). S-cone on channels would show the
opposite pattern of responses. Intermodulations could thus occur separately within on and off channels but
these separate responses would both be at the same intermodulation frequency and would not be separable
in our results. Further work could attempt to measure responses in S-cone increment and decrement
channels separately by manipulating the temporal waveform (Krauskopf and Zaidi, 1986; Stockman et al.,
2017).

SSVEP responses as captured by EEG reflect activity from multiple neural populations. However, it may be
possible to capture the activity of different neural populations by investigating different intermodulation
components (Regan, 1966; Appelbaum et al., 2008; Alp et al., 2016; Gordon et al., 2019). We therefore
plotted color tuning functions for all the intermodulation components that met our criterion of 2 SNR in
Experiments 1 and 4. By eye, we can see two distinct types of tuning function in the results of Experiment 4
(Figure 6). Tuning functions based on some of the intermodulation components show a second peak at a 180°
hue angle difference as for I_1 ($3F_B$-F_A, $2F_A$+F_B and F_A+$2F_B$), while tuning functions based on others do not have
the second peak at a 180° hue angle difference (F_B-F_A, $3F_B$-$3F_A$, $6F_B$-$3F_A$, $2F_A$+$2F_B$ and F_A+$3F_B$). It is possible that
these different groups of intermodulation components reflect the actions of different neural populations
with different color tuning properties. However, the scalp map distributions of the different components are
not obviously different (Figure 6). We also did not find any categorical effect of difference intermodulation
components versus sum intermodulation as has been suggested in the literature (e.g., the tuning functions
for F_A+F_B and $2F_A$+$2F_B$ are both sum intermodulation components but have different shapes).

Our investigation of intermodulation of EEG signals has revealed broadband cortical color tuning functions.
We found that cortical tuning is consistent across both cardinal and intermediate color axes, implying the
action of intermediately tuned cortical color mechanisms. Surprisingly, the tuning functions we measured

with intermodulation of EEG do not rely on chromatic edges, suggesting that they arise from single-opponent neural populations. Our tuning functions are consistent with those derived psychophysically, from single cell electrophysiology and from color masking in EEG, showing that intermodulation SSVEP is a promising method for probing cortical color representation in humans.

Acknowledgements
The work was funded by ERC grant 949242 COLOURCODE awarded to JMB.

References
Alp N, Kogo N, Van Belle G, Wagemans J, Rossion B (2016) Frequency tagging yields an objective neural signature of Gestalt formation. Brain and Cognition 104:15–24.

Appelbaum LG, Wade AR, Pettet MW, Vildavski VY, Norcia AM (2008) Figure-ground interaction in the human visual cortex. Journal of Vision 8:8–8.

Bohon KS, Hermann KL, Hansen T, Conway BR (2016) Representation of Perceptual Color Space in Macaque Posterior Inferior Temporal Cortex (the V4 Complex). eneuro 3:ENEURO.0039-16.2016.

Boynton RM, Kambe N (1980) Chromatic difference steps of moderate size measured along theoretically critical axes. Color Research and Application 5:13–23.

Chatterjee S, Callaway EM (2003) Parallel colour-opponent pathways to primary visual cortex. Nature 426:668–671.

Chen J, Gegenfurtner KR (2021) Electrophysiological evidence for higher-level chromatic mechanisms in humans. Journal of Vision 21:12.

Conway BR, Eskew RT, Martin PR, Stockman A (2018) A tour of contemporary color vision research. Vision Research 151:2–6.

Conway BR, Hubel D, Livingstone M (2002) Color Contrast in Macaque V1. Cerebral Cortex 12:915–925.

Conway BR, Moeller S, Tsao DY (2007) Specialized Color Modules in Macaque Extrastriate Cortex. Neuron 56:560–573.

D'Zmura M, Knoblauch K (1998) Spectral bandwidths for the detection of color. Vision Research 38:3117–3128.

Ekroll V, Faul F (2012) Basic Characteristics of Simultaneous Color Contrast Revisited. Psychol Sci 23:1246–1255.

Emery KJ, Volbrecht VJ, Peterzell DH, Webster MA (2017) Variations in normal color vision. VI. Factors underlying individual differences in hue scaling and their implications for models of color appearance. Vision Research 141:51–65.

Emery KJ, Volbrecht VJ, Peterzell DH, Webster MA (2023) Fundamentally different representations
of color and motion revealed by individual differences in perceptual scaling. Proceedings of
the National Academy of Sciences 120:e2202262120.

Engel S, Zhang X, Wandell B (1997) Colour tuning in human visual cortex measured with functional
magnetic resonance imaging. Nature 388:68–71.

Friedman HS, Zhou H, von der Heydt R (2003) The coding of uniform colour figures in monkey
visual cortex. The Journal of Physiology 548:593–613.

Garg AK, Li P, Rashid MS, Callaway EM (2019) Color and orientation are jointly coded and spatially
organized in primate primary visual cortex. Science 364:1275–1279.

Gegenfurtner KR (2003) Cortical mechanisms of colour vision. Nat Rev Neurosci 4:563–572.

Gegenfurtner KR, Kiper DC (1992) Contrast detection in luminance and chromatic noise. Journal of
the Optical Society of America A 9:1880–1888.

Gegenfurtner KR, Kiper DC, Levitt JB (1997) Functional Properties of Neurons in Macaque Area V3.
Journal of Neurophysiology 77:1906–1923.

Goddard E, Mannion DJ, McDonald JS, Solomon SG, Clifford CWG (2010) Combination of
subcortical color channels in human visual cortex. Journal of Vision 10:25–25.

Gordon N, Hohwy J, Davidson MJ, Van Boxtel JJA, Tsuchiya N (2019) From intermodulation
components to visual perception and cognition-a review. NeuroImage 199:480–494.

Gouras P (1974) Opponent-colour cells in different layers of foveal striate cortex. The Journal of
Physiology 238:583–602.

Hanazawa A, Komatsu H, Murakami I (2000) Neural selectivity for hue and saturation of colour in
the primary visual cortex of the monkey. European Journal of Neuroscience 12:1753–1763.

Hansen T, Gegenfurtner KR (2006) Higher level chromatic mechanisms for image segmentation.
Journal of Vision 6:5–5.

Hansen T, Gegenfurtner KR (2013) Higher order color mechanisms: Evidence from noise-masking
experiments in cone contrast space. Journal of Vision 13:26–26.

Hass CA, Horwitz GD (2013) V1 mechanisms underlying chromatic contrast detection. Journal of
Neurophysiology 109:2483–2494.

Johnson EN, Hawken MJ, Shapley R (2001) The spatial transformation of color in the primary visual
cortex of the macaque monkey. Nature neuroscience 4:409–416.

Johnson EN, Hawken MJ, Shapley R (2008) The Orientation Selectivity of Color-Responsive
Neurons in Macaque V1. J Neurosci 28:8096.

Jurcak V, Tsuzuki D, Dan I (2007) 10/20, 10/10, and 10/5 systems revisited: Their validity as relative
head-surface-based positioning systems. NeuroImage 34:1600–1611.

Kaneko S, Kuriki I, Andersen SK (2020) Steady-State Visual Evoked Potentials Elicited from Early
Visual Cortex Reflect Both Perceptual Color Space and Cone-Opponent Mechanisms.
Cerebral Cortex Communications 1:tgaa059.

Kiper DC, Fenstemaker SB, Gegenfurtner KR (1997) Chromatic properties of neurons in macaque
area V2. Visual Neuroscience 14:1061–1072.

Komatsu H, Ideura Y, Kaji S, Yamane S (1992) Color selectivity of neurons in the inferior temporal
cortex of the awake macaque monkey. J Neurosci 12:408.

Krauskopf J, Farell B (1990) Influence of colour on the perception of coherent motion. Nature
348:328–331.

Krauskopf J, Gegenfurtner K (1992) Color discrimination and adaptation. Vision research 32:2165–
2175.

Krauskopf J, Williams DR, Heeley DW (1982) Cardinal directions of color space. Vision research
22:1123–1131.

Krauskopf J, Williams DR, Mandler MB, Brown AM (1986) Higher order color mechanisms. Vision
research 26:23–32.

Krauskopf J, Zaidi Q (1986) Induced desensitization. Vision research 26:759–762.

Kuriki I, Sun P, Ueno K, Tanaka K, Cheng K (2015) Hue Selectivity in Human Visual Cortex Revealed
by Functional Magnetic Resonance Imaging. Cerebral Cortex 25:4869–4884.

Lennie P, Krauskopf J, Sclar G (1990) Chromatic Mechanisms in Striate Cortex of Macaque. The
Journal of Neuroscience 10:649–669.

Leventhal AG, Thompson KG, D Liu, Y Zhou, SJ Ault (1995) Concomitant sensitivity to orientation,
direction, and color of cells in layers 2, 3, and 4 of monkey striate cortex. J Neurosci
15:1808.

Li P, Garg AK, Zhang LA, Rashid MS, Callaway EM (2022) Cone opponent functional domains in
primary visual cortex combine signals for color appearance mechanisms. Nat Commun
13:6344.

Lindsey DT, Brown AM (2004) Masking of grating detection in the isoluminant plane of DKL color
space. Visual Neuroscience 21:269–273.

Livingstone MS, Hubel DH (1984) Anatomy and physiology of a color system in the primate visual
cortex. The Journal of neuroscience : the official journal of the Society for Neuroscience
4:309–356.

MacLeod DI, Boynton RM (1979) Chromaticity diagram showing cone excitation by stimuli of equal
luminance. Journal of the Optical Society of America 69:1183–1186.

645 Malkoc G, Kay P, Webster MA (2005) Variations in normal color vision. IV. Binary hues and hue
646   scaling. Journal of the Optical Society of America A, Optics, image science, and vision
647   22:2154–2168.

648 Norcia AM, Appelbaum LG, Ales JM, Cottereaur BR, Rossion B (2015) The steady state visual
649   evoked potential in vision research: A review. Journal of Vision 15:1–46.

650 Nunez V, Gordon J, Shapley RM (2021) A multiplicity of color-responsive cortical mechanisms
651   revealed by the dynamics of cVEPs. Vision Research 188:234–245.

652 O'Brien F, Cousineau D (2015) Erratum to "Representing Error bars in within-subject designs in
653   typical software packages." TQMP 11:126–126.

654 Regan D (1966) Some characteristics of average steady-state and transient responses evoked by
655   modulated light. Electroencephalography and Clinical Neurophysiology 20:238–248.

656 Regan D, Regan MP (1987) Nonlinearity in human visual responses to two-dimensional patterns,
657   and a limitation of fourier methods. Vision Research 27:2181–2183.

658 Regan MP, Regan D (1989) Objective Investigation of Visual Function Using a Nondestructive
659   Zoom-FFT Technique for Evoked Potential Analysis. Can j neurol sci 16:168–179.

660 Rossion B, Boremanse A (2011) Robust sensitivity to facial identity in the right human occipito-
661   temporal cortex as revealed by steady-state visual-evoked potentials. Journal of Vision
662   11:1–21.

663 Sankeralli MJ, Mullen KT (1997) Postreceptoral chromatic detection mechanisms revealed by noise
664   masking in three-dimensional cone contrast space. Journal of the Optical Society of
665   America A 14:2633–2646.

666 Schluppeck D, Engel SA (2002) Color Opponent Neurons in V1: A Review and Model Reconciling
667   Results from Imaging and Single-Unit Recording. Journal of Vision 2:5.

668 Shapley R, Hawken MJ (2011) Color in the Cortex: single- and double-opponent cells. Vision
669   Research 51:701–717.

670 Solomon SG, Lennie P (2005) Chromatic Gain Controls in Visual Cortical Neurons. J Neurosci
671   25:4779.

672 Solomon SG, Lennie P (2007) The machinery of colour vision. Nature Reviews Neuroscience 8:276–
673   286.

674 Stockman A, Henning GB, Rider AT (2017) Linear–nonlinear models of the red–green chromatic
675   pathway. Journal of Vision 17:7.

676 Stockman A, MacLeod D, Johnson N (1993) Spectral sensitivities of the middle- and long-
677   wavelength sensitive cones. Journal of the Optical Society of America, A 10:2491–2521.

678 Wachtler T, Sejnowski TJ, Albright TD (2003) Representation of Color Stimuli in Awake Macaque
679   Primary Visual Cortex. Neuron 37:681–691.

680   Webster MA, Mollon JD (1991) Changes in colour appearance following post-receptoral
681   adaptation. Nature 349:235–238.