Tuning of cortical color mechanism revealed using steady-state visually evoked potentials

Dylan J. Watts[1†], Ana Rozman[1†], Lucy P. Somers[1], Bora Gunel[1], Chris Racey[1], Katie Barnes[1] and Jenny M. Bosten[1*]

[1]School of Psychology, University of Sussex, Brighton UK
[†]Joint first authors
[*]Corresponding author: j.bosten@sussex.ac.uk

Abstract

Color information is thought to be received by the primary visual cortex via two dominant retinogeniculate pathways, one signals color variation between teal and red, and the other signals color variation between violet and lime. This representation is thought to be transformed in the cortex so that there are a number of different cell populations representing a greater variety of hues. However, the properties of cortical color mechanisms are not well understood. In four experiments, we characterized the tuning functions of cortical color mechanisms by measuring the intermodulation of steady-state visually evoked potentials (SSVEPs). Stimuli were isoluminant chromatic checkerboards where odd and even checks flickered at different frequencies. As hue dissimilarity between the odd and even checks increased, the amplitude of an intermodulation component (I_1) at the sum of the two stimulus frequencies decreased, revealing cortical color tuning functions. In Experiment 1 we found similar broad tuning functions for 'cardinal' and intermediate color axes, implying that the cortex has intermediately tuned color mechanisms. In Experiment 2 we found similar broad tuning functions for 'checkerboards' with no perceptible edges because the checks were formed from single pixels (~0.096°), implying that the underlying neural populations do not rely on spatial chromatic edges. In Experiment 3 we manipulated check size and found that color tuning functions were consistent across check sizes used. In Experiment 4 we measured full 360° tuning functions for a 'cardinal' cortical color mechanism and found evidence for opponent (bipolar) color responses. The observed cortical color tuning functions were consistent with those measured using psychophysics and electrophysiology, implying that tracking intermodulation using SSVEPs provides a useful method for measuring them.

Color vision, cortical color mechanisms, SSVEPs, intermodulation components, tuning functions, cardinal color axes, intermediate color axes

1.0 Introduction

Decades of research in the neural mechanisms of human color vision have produced a relatively good understanding of the representation of color at a receptor level, and in the retinogeniculate color pathways. However, the cortical mechanisms underlying color perception are much less well understood, and though color representation in early cortical areas has been studied using electrophysiology in animal models, there has been relatively little research on the properties of cortical color mechanisms in humans measured directly using neuroimaging. In particular, there is a debate over how far the separation of color information into distinct retinogeniculate pathways remains segregated in the cortex, and over the number and tuning of cortical color mechanisms. We used a novel approach using steady-state visually evoked potentials (SSVEPs) to measure and characterize the tuning functions of cortical color mechanisms in humans.

Normal human color vision begins with the relative activation of three types of retinal cone photoreceptors, sensitive to short (S), medium (M) or long (L) wavelengths of light. Cone signals are then thought to be combined postreceptorally by distinct populations of bipolar cells, setting up 'cardinal' color pathways which remain segregated at the retina and LGN and at least as far is the input layers of V1. The S/(L+M) pathway carried by small bistratified ganglion cells and though koniocellular layers of the LGN signals color differences between lime green and violet. The L/(L+M) pathway carried by the midget retinal ganglion cells and through parvocellular layers of the LGN signals color differences between teal and red (Solomon and Lennie, 2007; Conway et al., 2018).

To what extent retinogeniculate opponent color representations are retained in early visual cortical areas are retained and to what extent they are transformed is not fully understood. Although early psychophysical evidence was consistent with a 'cardinal' representation of color inherited from the retinogeniculate mechanisms (Boynton and Kambe, 1980; Krauskopf et al., 1982; Krauskopf and Farell, 1990), later evidence favored a 'higher order' color representation based on color mechanisms responsive to hues that are intermediate between the poles of the retinogeniculate color mechanisms (Krauskopf et al., 1986; Webster and Mollon, 1991; Gegenfurtner and Kiper, 1992; Krauskopf and Gegenfurtner, 1992; Hansen and Gegenfurtner, 2013). There has also been fMRI evidence in support of both a 'cardinal' color representation in V1 (Engel et al., 1997) and in support of the existence of intermediately tuned color mechanisms (Goddard et al., 2010; Kuriki et al., 2015). Another unresolved question about cortical color representation is to what extent it is bipolar (where two 'opponent' hues are represented with the achromatic represented somewhere in the middle of the mechanism's response range), or unipolar (where color variation between a single hue and the achromatic point is represented by each mechanism). Chromatic signals in the input layers of V1 must inherit the bipolar color representations of the retinogeniculate color pathways, but psychophysical evidence (Malkoc et al., 2005; Emery et al., 2017, 2023) and electrophysiological evidence from animal models (Gegenfurtner, 2003; Solomon and Lennie, 2007) is consistent with the idea that there are multiple 'monopolar' color mechanisms, each tuned to a particular hue and responding to neighboring hues within a given bandwidth.

Studies aiming to measure the tuning functions of color selective cortical neurons have been conducted mostly using indirect psychophysical methods in humans and directly using electrophysiology in primates. Results of psychophysical studies have typically shown broad tuning (Sankeralli and Mullen, 1997; Lindsey and Brown, 2004) especially when 'off-axis looking' (stimulus detection by neighboring color mechanisms than the one being measured) is minimized (D'Zmura and Knoblauch, 1998; Hansen and Gegenfurtner, 2006). Electrophysiological recordings in primates have revealed diversity in the preferred hues of color selective neurons in V1 (Lennie et al., 1990; Hanazawa et al., 2000; Hass and Horwitz, 2013), V2 (Kiper et al., 1997; Solomon and Lennie, 2005), V3 (Gegenfurtner et al., 1997), V4 (Bohon et al., 2016) as well as in downstream color selective areas in the ventral visual pathway (Komatsu et al., 1992; Conway et al., 2007). Color tuning functions of individual color-responsive cells measured electrophysiologically have been found to have a range of half-amplitude bandwidths in hue angle from about 30°-170° in V1 and 30°-60° in V2. The average bandwidth of color tuning functions tends to decrease through the hierarchy of cortical visual areas (Gegenfurtner, 2003), but large populations of broadly tuned neurons are still found in V4 (Bohon et al., 2016).

The color tuning properties of visual cortical neurons can of course interact with tuning properties to other stimulus features. For instance, many cortical neurons are parallelly tuned to both color and luminance (Johnson et al., 2001), spatial frequency (Lennie et al., 1990) or orientation (Leventhal et al., 1995; Friedman et al., 2003). One distinction that has been relatively well studied is between so-called single- and double color opponent neurons (Shapley and Hawken, 2011). Double opponent neurons respond to stimuli containing sharp chromatic edges but respond weakly or not at all to full-field or low spatial frequency chromatic stimuli, while single opponent neurons show the opposite pattern (Livingstone and Hubel, 1984; Johnson et al., 2008). Depending on the spatial properties of the chosen stimuli, color tuning results may therefore vary, because different populations of cells may be targeted that are tuned to combinations of color and other stimulus features. In particular, the stimuli with chromatic edges may preferentially activate double opponent neurons, while stimuli without edges may preferentially activate single opponent neurons.

In humans, other than by inference from the results of psychophysical methods, there is one existing study by Chen et al. (2021) that measured cortical color tuning functions using neuroimaging, specifically by measuring steady -state visually-evoked potentials (SSVEPs). SSVEPs occur in response to visual stimuli that are flickering at a precise temporal frequency (Regan, 1966; Norcia et al., 2015), and are typically analyzed in

the frequency domain where EEG noise is segregated into many frequency bands, providing high signal to noise for brain responses at the stimulus frequency (Rossion and Boremanse, 2011) and a sensitive index of visual responses in the cortex. Chen et al. (2021) used a method for measuring cortical color tuning functions based on chromatic noise masking, where psychophysical results have shown that the strength of masking increases with the color similarity of the target and mask (Gegenfurtner and Kiper, 1992; D'Zmura and Knoblauch, 1998). They presented central binary chromatic noise targets with surround binary chromatic noise masks, and measured occipital SSVEP amplitudes as a function of the angular difference between the color axes from which target and surround colors were sampled, finding broad color tuning functions centered at both cardinal and intermediate color axes.

In the current study we exploit intermodulation of SSVEPs as an alternative scheme for measuring cortical tuning functions (Regan and Regan, 1987). Intermodulation components in SSVEP occur when there are two or more stimuli flickering at different frequencies presented simultaneously, and are thought to provide an index of the degree to which shared neural resources process the different stimuli (Regan and Regan, 1989; Norcia et al., 2015). We presented flickering checkerboard stimuli where odd checks flickered at one frequency and remained at a fixed chromaticity across trials, and even checks flickered at a different frequency and varied in chromaticity. We measured SSVEP amplitudes at the intermodulation frequency, which we expected to be maximal when odd and even checks have the same chromaticity and are therefore processed by the same color mechanisms, and to decrease with the color difference between odd and even checks. In a series of four experiments, we measured cortical color tuning functions at cardinal and intermediate color axes (Experiment 1), when the stimulus lacked perceptible chromatic edges (Experiment 2), as a function of the size of checkerboard checks (Experiment 3), and around the full hue circle (Experiment 4). Our results reveal that bandwidths of cortical color tuning functions do not differ between cardinal and intermediate axes, implying the action of intermediately tuned color mechanisms; that surprisingly, tuning functions do not depend on the presence of perceivable chromatic edges, implying that they probe color mechanisms with low spatial opponency; and that there is a substantial bipolar component in human cortical color representations.

2.0 General Methods
Provided in this section are methods common to all four experiments.

2.1 Participants
Participants in all experiments had normal color vision assessed using an 'Anomaloskop' anomaloscope (Oculus, Wetzlar, Germany; Experiments 1-3) or the Ishihara Plates Test (Experiment 4), and were recruited via email and word-of-mouth. Participants provided written informed consent and were compensated £10 per hour for their time. The study received ethical approval from the Sussex University Science and Technology Cross Schools Research Ethics Committee (ER/DJW41), and adhered to the tenants of the World Medical Association's Declaration of Helsinki (2013) with the exception that it was not pre-registered.

2.2 Equipment
Electroencephalographic data were acquired using 64-channel Waveguard caps (ANT neuro, Hengelo, Netherlands) via a high-speed 64-channel ANT Neuro amplifier sampling at 1000 Hz. EEG data were recorded in ASA lab (ANT Neuro, Hengelo, Netherlands). Additional electrodes were used to measure horizontal (HEOG) and vertical (VEOG) eye-movements to allow for the removal of blink artefacts. A custom photodiode input to the EEG amplifier was fitted to check for accurate stimulus timing (see Supplementary Information 7).

Stimuli were displayed on a GDM FW900 CRT monitor (Sony, Tokyo, Japan). The monitor was gamma-corrected using a luminance meter (Konica-Minolta, Tokyo, Japan) and color calibrated using a PR650 SpectraScan spectroradiometer (PhotoResearch, Dorking, Surrey). Stimuli were presented using a ViSaGe MKII Stimulus Generator (Cambridge Research Systems, Rochester, UK) using Matlab 2017 (The Mathworks,

Nantick, CA). Triggers to the EEG amplifier from the ViSaGe via BNC and parallel port connections. The monitor had a resolution of 800 x 600 pixels and ran at a frame rate of 160 Hz.

2.3 Heterochromatic flicker photometry

Heterochromatic flicker photometric measurements were made in order to calculate stimuli that are isoluminant for individual observers. Stimuli consisted of a 2D annulus with an outer diameter of 41.9° and inner diameter of 5.7°, which oscillated at 11.4 Hz between two colors (Figure 1). On each trial, the intensity of one color was fixed, and participants adjusted the intensity of the other color until they perceived minimum flicker (the point of isoluminance). On the basis of these measurements, stimuli in the four experiments were made isoluminant to ensure that they isolated chromatic mechanisms and did not stimulate luminance mechanisms.

Figure 1. Stimulus properties. (a) Stimuli were isoluminant chromatic flickering checkerboards. Even checks flickered at F_A (6.66 Hz) between equal energy white and a 'variable' chromaticity that depended on the color condition. Odd checks flickered at F_B (8 Hz) between equal energy white and a 'fixed' chromaticity. A central 5.7° disc was metameric with equal energy white and did not flicker. (b) Stimulus chromaticities specified in a version of the MacLeod-Boynton (1979) chromaticity diagram based on the Stockman, MacLeod and Johnson (1993) cone fundamentals. For Experiment 1 the fixed chromaticity was either the S/(L+M) axis in a decremental direction (left panel), an intermediate axis (middle panel) or the L/(L+M) axis in an incremental direction (right panel). In each panel, the modulation made by the fixed chromaticity and equal energy white is indicated by the solid line. For each fixed color axis there were 6 color conditions, where the variable chromaticity differed from the fixed chromaticity in steps of 18° from 0° to 90°. Modulations between variable chromaticities and equal energy white are indicated by the dotted lines. For Experiments 2 and 3 the S/(L+M) axis in a decremental direction was tested (left panel). For Experiment 4 the fixed chromaticity was along the S/(L+M) axis in a decremental direction (left panel) and there were 12 color conditions in 30° steps between 0° and 300° from the fixed chromaticity (not shown).

2.4 Stimuli

The stimuli for all four experiments consisted of flickering checkerboards extending 76.8° horizontally and 48° vertically (Figure 1a), with a viewing distance of 30 cm. Each check was square and measured 4.8° x 4.8°. Even checks flickered between equal energy white and a fixed hue at 8 Hz, eliciting SSVEPs at 8 Hz and higher harmonics. Odd checks flickered between equal energy white and a variable hue at 6.66 Hz, eliciting SSVEPs

at 6.66 Hz and higher harmonics. Flicker was square wave between equal energy white and the edge of the maximum saturation circle allowed by the monitor's gamut (Figure 1b). There was a central disc metameric with equal energy white of 5.7° with a central fixation point which did not flicker, to avoid the macular pigment (Figure 1a).

2.5 Procedure
Participants first completed heterochromatic flicker photometry task. The consistency of intensity settings was assessed, and if found to range >20%, the task was repeated. Participants were then fitted with facial electrodes and an EEG cap in accordance with the 10/20 system (Jurcak et al., 2007). We ensured that impedances for electrodes of interest were < 10 kΩ and for other electrodes < 20kΩ. In the experiment, participants viewed a series of 10 second trials (see details of methods for each separate experiment). They were instructed not to blink excessively during trials and to blink between trials. Participants controlled the inter-trial interval by pressing a button on a keyboard when they were ready for the next trial. Raw EEG data were monitored in real time for signs of fatigue or other issues.

2.6 EEG Analysis: pre-processing
Pre-processing was conducted in MATLAB, using Letswave 7 (https://nocions.github.io/letswave7/) for Experiments 1 and 4, and Letswave 6 for Experiments 2 and 3 (https://nocions.github.io/letswave6/). First, raw EEG data were imported and mapped to the ANT neuro standard 64-channel electrode coordinate system. Raw data were visually inspected for excessive noise resulting from equipment failures. Problematic channels were removed. No exclusions of channels were made from our analysis cluster, consisting of Oz, O1, O2, Pz, P1, P2, P3, P4, POz, PO3, and PO4 (Figure 2d). Signals were re-referenced to the grand average of all remaining cortical electrodes. In addition to our analysis cluster, channels FPz, HEOG and VEOG were extracted to detect eye movements. Consistency in temporal profile of our stimuli was ensured by inspecting the decomposed frequency profile of the photodiode which was attached to the screen (see Supplementary Information 7). A Butterworth filter for 50 Hz noise and a high-pass filter (<0.1 Hz) were applied. Data were then segmented into trials using an epoch size of 9000 s, which included an integer number of stimulus repetitions at both flicker frequencies. Segmented trials were baseline corrected within the 1 to 1.2 interval. No ocular artefacts were removed in Experiments 1 and 4 (see Supplementary Information 2 for a comparison of signal amplitudes with and without ocular artefact removal). For Experiments 2 and 3, ocular artefacts were removed by visual inspection if there were fewer than 2 blinks per color condition. Alternatively, Independent Components Analysis (ICA) was applied if there were two or more blinks per condition. Recordings for an individual observer were merged in cases where data was collected across multiple sessions. Trials from the same condition for each observer were then averaged and converted to the frequency domain by fast Fourier transformation (FFT). Amplitudes for each observer and each color condition were extracted at I_1 (14.66 Hz), which is an intermodulation component at the sum of the two stimulus frequencies. Amplitudes were also extracted at F_B (8 Hz), the frequency of the checks of fixed chromaticity – changes in amplitude with the chromaticity of the F_A (6.66 Hz) checks reflects chromatic masking, which has previously been used to infer cortical color tuning functions (Chen and Gegenfurtner, 2021).

3.0 Experiment 1
In Experiment 1 we sought to measure cortical color tuning functions using SSVEP, and to determine whether there are differences in cortical color tuning functions between 'cardinal' color axes aligned with the retinogeniculate color channels L/(L+M) and S/(L+M) and an intermediate color axis.

3.1 Experiment 1: Methods
Eleven participants (4 female) completed two experimental sessions, each lasting 2.5 hours. Three color axes were tested: L/(L+M) increments, S/(L+M) decrements and an intermediate axis (Figure 1b). For each axis, the chromaticity of checks flickering at F_A was fixed along that axis, and the chromaticity of checks

flickering at F_B varied on each trial between six hue angles to the fixed axis (0°, 18°, 36°, 54°, 72° and 90°; Figure 2a). There were 180 trials in each session presented in a random order, totaling 360 trials.

Figure 2. Experiment 1 results. (a) Amplitude profile for an example condition (S/(L+M) axis, 0° variable chromaticity), averaged over the 11 participants. (b) SNR profile for the same condition. Fundamental stimulus frequencies and their harmonics are labelled in black and intermodulation components in red. (c) Mean amplitude of the intermodulation component at 14.66 Hz as a function of the hue angle between the fixed and variable chromaticities for the S/(L+M) axis (green solid line), the intermediate axis (orange dotted line) and the L/(L+M) axis (red dashed line). Error bars are 95% confidence intervals. (d) Mean amplitude at the stimulus frequency of the fixed chromaticity at 8 Hz as a function of the hue angle between the fixed and variable chromaticities for the S/(L+M) axis (green solid line), the intermediate axis (orange dotted line) and the L/(L+M) axis (red dashed line). Error bars are within-participant 95% confidence intervals (O'Brien and Cousineau, 2015), jittered along the x-axis for visibility. (e) Schematic representation of analysis cluster electrodes.

3.2 Experiment 1: Results

Strong signals were observed at the two fundamental frequencies, their harmonics ($F_A(1)$ and $F_B(1)$), and at I_1 (e.g. Figures 2a and b). There were additional amplitude peaks at other harmonics of F_A and F_B and at other intermodulation frequencies. We extracted I_1 and F_B for further analyses.

Figure 2 shows the effect of hue dissimilarity between odd and even checks on SSVEP amplitudes at I_1 (panel c) and F_B (panel d) for the three color axes. Tuning functions were similar in shape for all three axes, with a broad half-amplitude bandwidth of approximately 90°. Two-way repeated-measures ANOVAs were

conducted to test the effect of color axis and hue angle differences on the amplitudes of I_1 and F_B. Mauchly's test of sphericity was violated for both frequencies ($X^2(54) = 106.50$ for I_1 and $X^2 (54) = 115.40$ for F_B, $p < .001$), therefore Greenhouse-Geisser corrected tests were used ($\varepsilon =.311$ for I_1 and $.260$ for F_B). The ANOVAs showed that for both I_1 and F_B there were significant main effects of hue angle difference ($F(1.19,11.92) = 21.59$ for I_1 and $F(1.84,15.49) = 23.18$ for F_B, $p <.001$, $\eta^2_p =.683$ for I_1 and $\eta^2_p = .699$ for F_B). Results from post-hoc tests (Supplementary Information 1, Supplementary Tables 1 and 2) show that mean SSVEP amplitude decreased significantly with hue angle difference for I_1 and increased significantly with hue angle difference for F_B.

There was a significant main effect of color axis for F_B, ($F(1.14, 11.36) = 10.83$, $p = .006$, $\eta^2_p =.520$), but not for I_1 ($F(1.45,14.51) =3.17$, $p =.084$, $\eta^2_p =.241$). Post-hoc tests using Bonferroni correction indicated that there was a significant difference between L/(L+M) and S/(L+M) axes ($M = -.247$, $p < .05$). There was no significant interaction between axis and hue angle difference either for I_1 ($F(3.11,31.06) = 2.74$, $p = .058$) nor for F_B ($F(2.60, 26.03) = 2.40$, $p = .098$). Thus, the effect of hue angle on SSVEP amplitude was consistent across all three axes for both frequencies, implying that tuning functions were shaped similarly for all three axes.

Figure 3. Results of Experiment 2. SSVEP amplitude is plotted as a function of hue angle dissimilarity for single-pixel checkerboard stimuli. (a) for I_1 at 14.66 Hz. (a) for F_B at 8 Hz. Error bars are within-participant 95% confidence intervals (O'Brien and Cousineau, 2015).

4.0 Experiment 2
The isoluminant checkerboard stimuli in Experiment 1 have strong chromatic edges. The S/(L+M) axis, and thus neural populations responding to chromatic edges may underlie the results. To test the effect of chromatic edges on cortical color tuning functions measured using SSVEP we reduced the size of the 'checks' to single pixels (invisible to participants). If tuning functions change from those measured in Experiment 1 when the checkerboard is constructed from single pixels, then the tuning functions measured in Experiment 1 are attributable to neural populations responsive to chromatic edges.

4.1 Experiment 2: Methods
Eleven participants (6 female) completed a single 2.5-hour experimental session. Stimuli consisted of single pixel checkerboards (.096° x .096°), with the same spatial organization as depicted in Figure 1a. Individual checks were indistinguishable to the observer. The chromaticity of checks flickering at F_B was fixed, and the chromaticity of checks flickering at F_A varied between six hue angles to the fixed chromaticity (0°, 18°, 36°, 54°, 72° and 90°). There were 120 experimental trials, where hue conditions were presented in a random order. Since the S/(L+M) axis produced somewhat larger amplitudes in Experiment 1 than the other two axes, Experiment 2 probed the S/(L+M) axis only.

4.2 Experiment 2: Results

Figure 3 shows the amplitudes of I_1 and F_B against the difference in hue angle between odd and even single pixel checks. One-way repeated-measures ANOVAs examined the effect of hue angle difference on the amplitudes of I_1 and F_B. Mauchly's test of sphericity was violated for both frequencies ($X^2(14) = 72.16$ for I_1 and $X^2(14) = 34.53$ for F_B, $p < .001$), therefore Greenhouse-Geisser corrected tests were used ($\varepsilon = .279$ for I_1 and $\varepsilon = .510$ for F_B). There was a significant effect of hue angle difference on signal amplitude for I_1 ($F(1.40, 13.96) = 9.56$, $p = .005$, $\eta^2p = .489$), but not for F_B ($F(2.55, 25.48) = .465$, $p = .679$, $\eta^2_p = .044$). Results from post-hoc tests (see Supplementary Information 1, Supplementary Tables 3 and 4) indicated that mean SSVEP amplitude decreased with hue angle difference for I_1, but did not change with hue angle difference for F_B.

5.0 Experiment 3

Since we observed a difference in the shape of the tuning function for F_B between the 4.8° checks presented in Experiment 1 and the indivisible 0.096° checks presented in Experiment 2, in Experiment 3 we systematically manipulated check size. We measured tuning functions at I_1 and at F_B at 4 check sizes, varying between the check sizes used in Experiments 1 and 2.

5.1 Experiment 3: Methods

Eleven participants (5 female) completed two 2.5-hour experimental sessions. There were 4 check size conditions: 0.096° (as in Experiment 2), 0.4°, 1.3° and 4.8° (as in Experiment 1). To limit the duration of testing sessions there were four hue angle difference conditions rather than the 6 used in Experiments 1 and 2. These were 0°, 30°, 60° and 90°. Checks flickering at F_B (8 Hz) at a fixed chromaticity modulated between equal energy white and a decrement along S/(L+M) axis (left panel of Figure 1b). 2 testing sessions were conducted separated by at least one day. There were 160 experimental trials per session (10 trials for each combination of check size and hue angle difference), totaling 320 trials (20 trials in total for each stimulus type).

Figure 4. Results of Experiment 3. SSVEP amplitude is plotted as a function of hue angle dissimilarity (x-axis) and check size (legend). (a) for I_1 at 14.66 Hz. (b) for F_B at 8 Hz. Error bars are within-participant 95% confidence intervals (O'Brien and Cousineau, 2015), jittered along the x-axis for visibility.

5.2 Experiment 3: Results

Figure 4 shows the effect of check size and hue angle difference on amplitudes at I_1 and F_B. The figure shows that for I_1 check size does not affect the shape of the tuning functions observed. However, F_B, in agreement

with the results of Experiment 2 there is no tuning for the single pixel checkerboard, but similar tuning functions as observed in the results of Experiment 1 for the larger check sizes.

We conducted two-way repeated-measures ANOVAs to test the effect of hue angle hue angle difference and check size on the amplitudes at I_1 and F_B. Mauchly's test of sphericity was violated for both frequencies ($X^2(44)$ = 81.52, p = .002 for I_1; $X^2(44)$ = 134.05, $p < .001$ for F_B). Greenhouse-Geisser corrected tests were therefore used (ε = .367 for I_1; ε = .251 for F_B). There were significant main effects of check size and hue angle on signal amplitude for I_1 ($F(2.32, 23.21)$ = 3.61, p= .038, η^2_p = .265 for check size; $F(1.24, 12.39)$ = 33.76, $p <.001$, η^2_p =.771 for hue angle difference), but no significant interaction ($F(3.30, 33.01)$ = 1.55, p =.218, η^2_p =.134). Post-hoc tests (Supplementary Information 1, Supplementary Tables 5 and 6) indicated that mean amplitude in Condition 3 was somewhat stronger than in other conditions, regardless of hue angle, and that mean SSVEP amplitude decreased with hue angle for I_1, regardless of check size. For F_B there was a significant main effect of hue angle difference ($F(1.27, 12.67)$ = 11.30, p = .004, η^2_p =.530), but there was no significant main effect of check size ($F(1.51, 15.06)$ = .428, p =.604, η^2_p =.041). Results from post-hoc tests (Supplementary Information 1, Supplementary Tables 5 and 6) indicated that mean SSVEP amplitude increased with hue angle difference. There was a significant interaction between check size and hue angle for F_B ($F(2.26, 22.58)$ = 3.39, p = .046, η^2_p =.253). Post-hoc analysis (Supplementary Information 1, Supplementary Table 6) indicated that the shape of the tuning function for F_B differed between check size conditions.

6.0 Experiment 4
The tuning functions observed in the results of Experiments 1-3 are broadband but do not reach an asymptote at the largest hue angle differences we measured of 90°. In Experiment 4 we aimed to characterize a tuning function for a cortical color mechanism around the entire hue circle. We kept the fixed chromaticity checks along the S/(L+M) axis in a decremental direction but varied the variable chromaticity checks around the full hue circle in 12 steps.

6.1 Experiment 4: Methods
Ten participants (7 female) completed Experiment 4 in one experimental session lasting about 3 hours. Stimuli were isoluminant flickering checkerboards with a check size of 1.3° (which elicited the largest signals in Experiment 3). The fixed chromaticity flickering at F_B (8 Hz) was a decrement on the S/(L+M) axis (first panel of Figure 1b). The chromaticity of the variable checks flickering at F_A (6.66 Hz) spanned the full hue circle in steps of 30° (hue angle differences ranged from 0° to 330°). Each of the 12 hue angle conditions was presented 20 times, totaling 240 trials. SSVEP amplitudes were measured at I_1 and at F_B.

Figure 5. Results of Experiment 4. (a) tuning functions plotted at I_1 (full line) and F_B (dashed line), where the fixed chromaticity was an S/(L+M) decrement (180°), and the variable chromaticity varied around the full hue circle. Error

envelopes are within-participant 95% confidence intervals (O'Brien and Cousineau, 2015). Tuning functions for decrement in S/(L+M) mechanism comparing Experiment 4 results with those from Experiments 1-3 are plotted in Panel (b) for I_1 (14.66 Hz) and Panel (c) for F_B (8 Hz). For Experiment 3, results are shown only for the 0.096° and 1.3° check size condition.

6.2 Experiment 4: Results
Results from Experiment 4 are presented in Figure 5a. There is a peak in I_1 centered at 0° hue angle difference, as expected from the results of Experiments 1-3. However, there is a second smaller peak centered at a hue angle difference of 180°. For F_B there is a minimum at a hue angle difference of 0° as expected from the results of Experiments 1 and 3. There is no evidence of a second minimum at a hue angle difference of 180°.

One-way repeated measures ANOVAs were used to investigate the amplitude differences between hue angle difference conditions at I_1 and at F_B. There was a significant effect of hue angle difference on amplitudes at I_1 ($F(11, 99) = 18.065$, $p < .001$, $\eta^2_p = .667$) and at F_B ($F(11, 99) = 9.851$, $p < .001$, $\eta^2_p = .523$). Post hoc comparisons (Supplementary Information 1, Supplementary Tables 7 and 8) showed that amplitudes at I_1 for a 0° hue angle difference were significantly higher than amplitudes at all other angles, except for 330°. The amplitude for a hue angle difference of 180° was not significantly different from those of its most proximal hue angle difference conditions, 150° and 210°.

Figure 6. Signal heatmaps and tuning functions at intermodulation components surpassing 2 SNR in Experiment 4. SNR is presented as a function of Frequency in Hz with a range of 0 – 25. The components and their sources (nature of interaction between F_A and F_B) are labelled above individual heatmap and tuning function plots. The individual heatmaps and tuning function plots are scaled to match in area under the curve.

7.0 Analyses of additional intermodulation components
It is thought that different intermodulation components may arise from separate subpopulations of neurons (Regan, 1966; Gordon et al., 2019). For example, the subpopulation of neurons responding to the difference between F_A and F_B might be separate from the population responding to the sum of F_A and F_B, even though they both arise from interactions to the same signals (F_A and F_B). These potentially different populations may

also demonstrate distinct color tuning and may be differently spatially distributed across the cortex. We therefore conducted additional analyses of the data from Experiments 1 and 4 to plot tuning functions for other intermodulation components and visualize their locations in cortical electrode maps. Firstly, we extracted SNR profiles for a grand mean across observers in the 0° hue difference condition (where intermodulation components are expected to be greatest). Intermodulation components (i.e., those where the frequency peak is composed of either an addition or subtraction of multiples of F_A and F_B) were extracted when the SNR peak surpassed a criterion of 2. This procedure identified 2 qualifying components in Experiment 1 (Supplementary Information 4, Supplementary Figure 5) and 13 qualifying components in Experiment 4 (Figure 6). Spatial signal distribution was visualized using amplitude scalp heatmaps with the grand mean across observers, using a relativized scale between frequencies so that the maximum values are always plotted in yellow and minimum values in dark blue. The tuning functions were calculated for the channels in the analysis cluster and visualized using scaled line plots.

Comparison of the cortical surface heatmaps for different intermodulation components revealed that their locations are broadly consistent, both for Experiment 1 and Experiment 4. However, we found some substantial differences in tuning functions for different intermodulation components. In results for Experiment 4, Some functions show a *reduction* in amplitude for the opponent hue (e.g., 26.66 Hz and 28 Hz), in contrast to the characteristic peak observed in I_1. At the 20 Hz intermodulation component, tuning appears to be reversed, with a higher relative peak for the opponent compared to the proximal hues (Figure 6). Overall, peaks are less pronounced for intermodulation components other than I_1, but nonetheless may reveal different color tuning properties for different neural subpopulations.

8.0 Draft Discussion

Our results reveal that cortical color tuning functions can be successfully measured using SSVEP. As predicted, SSVEP amplitudes at the intermodulation frequency (I_1) significantly and systematically reduced with hue angle dissimilarity, over all the color axes and check sizes we tested. This was the case for both 'cardinal' color axes as well as an 'intermediate' color axis. The shape of color tuning functions that we measured by extracting amplitudes of I_1 did not depend on whether or not the stimulus contained perceptible edges. Therefore, these color tuning functions do not rely on neural mechanisms subserving chromatic spatial edges. When measured around the entire hue circle, the tuning function exhibited a bipolar shape with a similar half-amplitude bandwidth at both peaks.

Our finding that cortical color tuning bandwidths do not vary between the cardinal and intermediate color axes confirms the action of intermediately tuned cortical chromatic mechanisms. Chen and Gegenfurtner (2021) suggested that broad tuning with a half-amplitude bandwidth greater than 30° typically results from a linear combination of cone-opponent signals (Hansen and Gegenfurtner, 2006). However, nonlinear combinations of cone-opponent signals could also result in broad tuning functions along intermediate axes. As we did not observe higher SSVEP amplitudes along the intermediate axis than along both cardinal axes, linear summation of signals from cone-opponent mechanisms cannot explain the tuning functions we have measured for the intermediate axis. Our suggestion that tuning functions for the intermediate axis arise from nonlinear combinations of cone-opponent mechanisms aligns with results from single-unit recordings from primate V1 (Horwitz, 2020; Wachtler et al., 2003).

Chen and Gegenfurtner (2021) plotted cortical color tuning functions using a similar stimulus to ours as the amplitude of the frequency at which 'fixed' chromaticity modulated, reasoning that amplitude reduces with chromatic masking if the color channel is sensitive to both modulating colors. In the stimulus that best replicated theirs (Experiment 1), tuning functions derived from amplitudes at the frequency of the fixed chromaticity F_B were roughly the inverse of tuning functions derived from I_1, implying that they both reveal the action of the same chromatic mechanisms. However, in Experiments 2 and 4, the shapes of the observed tuning functions derived from F_B differed substantially from those derived from I_1. In our 'single pixel' checkerboard stimulus without perceptible spatial edges, tuning functions derived from I_1 matched those

measured using larger check sizes, but tuning functions derived from F_B were flat. This implies that the chromatic mechanisms revealed by I_1 and F_B are distinct: the mechanism revealed by F_B is dependent on spatial chromatic edges, but the mechanism revealed by I_1 is not. In the results of Experiment 4, tuning functions derived from I_1 are bipolar, but tuning functions derived from F_B are unipolar.

SSVEP responses as captured by EEG reflect activity from multiple neural populations. Therefore, we cannot be sure how early in the cortical hierarchy the tuning functions we have captured emerge (Kaneko et al., 2020). We found bipolar color tuning in the results of Experiment 4, which may imply that signals from I_1 primarily reflect early stages of cortical processing (e.g., V1), where many neurons remain opponently tuned (Engel et al., 1997).

Our findings offer electrophysiological evidence in accordance with previous psychophysical, EEG and fMRI accounts of higher-order chromatic mechanisms (D'Zmura and Knoblauch, 1998; Kuriki et al., 2015; Chen and Gegenfurtner, 2021). Our results reveal consistent cortical tuning across both cardinal and intermediate color axes for color mechanisms that do not require chromatic edges and exhibit some bipolar tuning. The observed tuning functions are consistent with those derived psychophysical and neurophysiological methods, showing that SSVEP is a promising method for probing cortical color tuning in humans.

Acknowledgements

The work was funded by ERC grant 949242 COLOURCODE awarded to JMB.

References

Bohon KS, Hermann KL, Hansen T, Conway BR (2016) Representation of Perceptual Color Space in Macaque Posterior Inferior Temporal Cortex (the V4 Complex). eneuro 3:ENEURO.0039-16.2016.

Boynton RM, Kambe N (1980) Chromatic difference steps of moderate size measured along theoretically critical axes. Color Research and Application 5:13–23.

Chen J, Gegenfurtner KR (2021) Electrophysiological evidence for higher-level chromatic mechanisms in humans. Journal of Vision 21:12.

Conway BR, Eskew RT, Martin PR, Stockman A (2018) A tour of contemporary color vision research. Vision Research 151:2–6.

Conway BR, Moeller S, Tsao DY (2007) Specialized Color Modules in Macaque Extrastriate Cortex. Neuron 56:560–573.

D'Zmura M, Knoblauch K (1998) Spectral bandwidths for the detection of color. Vision Research 38:3117–3128.

Emery KJ, Volbrecht VJ, Peterzell DH, Webster MA (2017) Variations in normal color vision. VI. Factors underlying individual differences in hue scaling and their implications for models of color appearance. Vision Research 141:51–65.

Emery KJ, Volbrecht VJ, Peterzell DH, Webster MA (2023) Fundamentally different representations of color and motion revealed by individual differences in perceptual scaling. Proceedings of the National Academy of Sciences 120:e2202262120.

Engel S, Zhang X, Wandell B (1997) Colour tuning in human visual cortex measured with functional magnetic resonance imaging. Nature 388:68–71.

Friedman HS, Zhou H, von der Heydt R (2003) The coding of uniform colour figures in monkey visual cortex. The Journal of Physiology 548:593–613.

Gegenfurtner KR (2003) Cortical mechanisms of colour vision. Nat Rev Neurosci 4:563–572.

Gegenfurtner KR, Kiper DC (1992) Contrast detection in luminance and chromatic noise. Journal of the Optical Society of America A 9:1880.

Gegenfurtner KR, Kiper DC, Levitt JB (1997) Functional Properties of Neurons in Macaque Area V3. Journal of Neurophysiology 77:1906–1923.

Goddard E, Mannion DJ, McDonald JS, Solomon SG, Clifford CWG (2010) Combination of subcortical color channels in human visual cortex. Journal of Vision 10:25–25.

Gordon N, Hohwy J, Davidson MJ, Van Boxtel JJA, Tsuchiya N (2019) From intermodulation components to visual perception and cognition-a review. NeuroImage 199:480–494.

Hanazawa A, Komatsu H, Murakami I (2000) Neural selectivity for hue and saturation of colour in the primary visual cortex of the monkey. European Journal of Neuroscience 12:1753–1763.

Hansen T, Gegenfurtner KR (2006) Higher level chromatic mechanisms for image segmentation. Journal of Vision 6:5–5.

Hansen T, Gegenfurtner KR (2013) Higher order color mechanisms: Evidence from noise-masking experiments in cone contrast space. Journal of Vision 13:26–26.

Hass CA, Horwitz GD (2013) V1 mechanisms underlying chromatic contrast detection. Journal of Neurophysiology 109:2483–2494.

Horwitz GD (2020) Signals Related to Color in the Early Visual Cortex. Annual Review of Vision Science 6:287–311.

Johnson EN, Hawken MJ, Shapley R (2001) The spatial transformation of color in the primary visual cortex of the macaque monkey. Nature neuroscience 4:409–416.

Johnson EN, Hawken MJ, Shapley R (2008) The Orientation Selectivity of Color-Responsive Neurons in Macaque V1. J Neurosci 28:8096.

Jurcak V, Tsuzuki D, Dan I (2007) 10/20, 10/10, and 10/5 systems revisited: Their validity as relative head-surface-based positioning systems. NeuroImage 34:1600–1611.

Kaneko S, Kuriki I, Andersen SK (2020) Steady-State Visual Evoked Potentials Elicited from Early Visual Cortex Reflect Both Perceptual Color Space and Cone-Opponent Mechanisms. Cerebral Cortex Communications 1:tgaa059.

Kiper DC, Fenstemaker SB, Gegenfurtner KR (1997) Chromatic properties of neurons in macaque area V2. Visual Neuroscience 14:1061–1072.

Komatsu H, Ideura Y, Kaji S, Yamane S (1992) Color selectivity of neurons in the inferior temporal cortex of the awake macaque monkey. J Neurosci 12:408.

Krauskopf J, Farell B (1990) Influence of colour on the perception of coherent motion. Nature 348:328–331.

Krauskopf J, Gegenfurtner K (1992) Color discrimination and adaptation. Vision research 32:2165–2175.

Krauskopf J, Williams DR, Heeley DW (1982) Cardinal directions of color space. Vision research 22:1123–1131.

Krauskopf J, Williams DR, Mandler MB, Brown AM (1986) Higher order color mechanisms. Vision research 26:23–32.

Kuriki I, Sun P, Ueno K, Tanaka K, Cheng K (2015) Hue Selectivity in Human Visual Cortex Revealed by Functional Magnetic Resonance Imaging. Cerebral Cortex 25:4869–4884.

Lennie P, Krauskopf J, Sclar G (1990) Chromatic Mechanisms in Striate Cortex of Macaque. The Journal of Neuroscience 10:649–669.

Leventhal AG, Thompson KG, D Liu, Y Zhou, SJ Ault (1995) Concomitant sensitivity to orientation, direction, and color of cells in layers 2, 3, and 4 of monkey striate cortex. J Neurosci 15:1808.

Lindsey DT, Brown AM (2004) Masking of grating detection in the isoluminant plane of DKL color space. Visual Neuroscience 21:269–273.

Livingstone MS, Hubel DH (1984) Anatomy and physiology of a color system in the primate visual cortex. The Journal of neuroscience : the official journal of the Society for Neuroscience 4:309–356.

MacLeod DI, Boynton RM (1979) Chromaticity diagram showing cone excitation by stimuli of equal luminance. Journal of the Optical Society of America 69:1183–1186.

Malkoc G, Kay P, Webster MA (2005) Variations in normal color vision. IV. Binary hues and hue scaling. Journal of the Optical Society of America A, Optics, image science, and vision 22:2154–2168.

Norcia AM, Appelbaum LG, Ales JM, Cottereaur BR, Rossion B (2015) The steady state visual evoked potential in vision research: A review. Journal of Vision 15:1–46.

O'Brien F, Cousineau D (2015) Erratum to "Representing Error bars in within-subject designs in typical software packages." TQMP 11:126–126.

Regan D (1966) Some characteristics of average steady-state and transient responses evoked by modulated light. Electroencephalography and Clinical Neurophysiology 20:238–248.

Regan D, Regan MP (1987) Nonlinearity in human visual responses to two-dimensional patterns, and a limitation of fourier methods. Vision Research 27:2181–2183.

Regan MP, Regan D (1989) Objective Investigation of Visual Function Using a Nondestructive Zoom-FFT Technique for Evoked Potential Analysis. Can j neurol sci 16:168–179.

Rossion B, Boremanse A (2011) Robust sensitivity to facial identity in the right human occipito-temporal cortex as revealed by steady-state visual-evoked potentials. Journal of Vision 11:1–21.

Sankeralli MJ, Mullen KT (1997) Postreceptoral chromatic detection mechanisms revealed by noise masking in three-dimensional cone contrast space. Journal of the Optical Society of America A 14:2633–2646.

Shapley R, Hawken MJ (2011) Color in the Cortex: single- and double-opponent cells. Vision Research 51:701–717.

Solomon SG, Lennie P (2005) Chromatic Gain Controls in Visual Cortical Neurons. J Neurosci 25:4779.

Solomon SG, Lennie P (2007) The machinery of colour vision. Nature Reviews Neuroscience 8:276–286.

Stockman A, MacLeod D, Johnson N (1993) Spectral sensitivities of the middle- and long-wavelength sensitive cones. Journal of the Optical Society of America, A 10:2491–2521.

Wachtler T, Sejnowski TJ, Albright TD (2003) Representation of color stimuli in awake macaque primary visual cortex. Neuron, 20:681-91.

Webster MA, Mollon JD (1991) Changes in colour appearance following post-receptoral adaptation. Nature 349:235–238.