Transcriptional complexity in the insect central complex: single nuclei RNA-sequencing of adult brain neurons derived from type 2 neuroblasts

Derek Epiney[#], Gonzalo N Morales Chaya[#], Noah R Dillon[#], Sen-Lin Lai[*], and Chris Q. Doe[*]

Institute of Neuroscience, Howard Hughes Medical Institute, University of Oregon, Eugene, OR 97403

Neuroblast, adult brain neurons, central complex, type 2 lineage, glia, homeodomain transcription factors, single nuclei RNA-sequencing, transcriptional profiling

\#   equal contribution
*   Corresponding authors: *cdoe@uoregon.edu*, *slai@uoregon.edu*

Abstract

In both invertebrates such as *Drosophila* and vertebrates such as mouse or human, the brain contains the most diverse population of cell types of any tissue. It is generally accepted that transcriptional diversity is an early step in generating neuronal and glial diversity, followed by the establishment of a unique gene expression profile that determines morphology, connectivity, and function. In *Drosophila*, there are two types of neural stem cells, called Type 1 (T1) and Type 2 (T2) neuroblasts. In contrast to T1 neuroblasts, T2 neuroblasts generate intermediate neural progenitors (INPs) that expand the number and diversity of cell types. The diversity of T2-derived neurons contributes a large portion of the central complex (CX), a conserved brain region that plays a role in sensorimotor integration. Recent work has revealed much of the connectome of the CX, but how this connectome is assembled remains unclear. Mapping the transcriptional diversity of neurons derived from T2 neuroblasts is a necessary step in linking transcriptional profile to the assembly of the adult brain. Here we perform single nuclei RNA sequencing of T2 neuroblast-derived adult neurons and glia. We identify clusters containing all known classes of glia, clusters that are male/female enriched, and 161 neuron-specific clusters. We map neurotransmitter and neuropeptide expression and identify unique transcription factor combinatorial codes for each cluster (presumptive neuron subtype). This is a necessary step that directs functional studies to determine whether each transcription factor combinatorial code specifies a distinct neuron type within the CX. We map several columnar neuron subtypes to distinct clusters and identify two neuronal classes (NPF+ and AstA+) that both map to two closely related clusters. Our data support the hypothesis that each transcriptional cluster represents one or a few closely related neuron subtypes.

Introduction

In all organisms, the brain has arguably the most complex cellular diversity, from human (Siletti et al., 2023) to *Drosophila* (Davie et al., 2018; Franconville et al., 2018; Hulse et al., 2021). Neuronal diversity is essential for the assembly and function of the adult brain, yet the "parts list" of different neuronal and glial cell types remains incomplete. In *Drosophila*, the laterally-positioned optic lobes have been well characterized for transcriptionally distinct neurons and glia (Konstantinides et al., 2018; Konstantinides et al., 2022; Özel et al., 2022). Less is known about the central brain, with only whole brain transcriptomes (Davie et al., 2018; Li et al., 2022) and a midbrain transcriptome reported (Croset et al., 2018).

The central brain contains many diverse neurons that populate important neuropils, such as the mushroom body (MB) used for learning and memory (Sgammeglia and Sprecher, 2022), or the central complex (CX) used for celestial navigation and sensory-motor integration (Fisher, 2022), among other behaviors. The central brain neurons are all generated from neural stem cells, called neuroblasts (NBs). There are two types of NBs that generate central brain neurons: Type 1 (T1) and Type 2 (T2) NBs. T1 NBs undergo asymmetric divisions to self-renew and generate a ganglion mother cells (GMCs), which each produce two post-mitotic neurons (Pollington et al., 2023; Yu et al., 2010; Yu et al., 2013). There are ~100 T1 NBs per larval central brain lobe that each generate 20-100 neurons and glia (Ito et al., 2013; Yu et al., 2013). T2 NBs have a more complex division pattern than T1 NBs (Bello et al., 2008; Boone and Doe, 2008; Bowman et al., 2008). T2 NBs undergo asymmetric division to self-renew and generate an intermediate neural progenitor (INP); each INP undergoes 4-6 divisions to self-renew and generate a GMC and its two neuron or glial progeny (Bello et al., 2008; Boone and Doe, 2008; Bowman et al., 2008). Thus, each T2 NB division will generate 8-12 progeny cells. Most T2 NBs generate ~500 neurons (Yu et al., 2013), have highly complex neuronal cell types based on morphology and connectivity (Hulse et al., 2020; Wang et al., 2014; Yang et al., 2013; Yu et al., 2013), and produce the columnar neurons of the adult CX (Boyan and Williams, 2011; Kandimalla et al., 2023). Subsequently, we will call neurons born from type I NBs "T1-derived" and neurons born from type 2 NBs "T2-derived."

Although single cell RNA-seq (scRNA-seq) has been done on adult central brain neurons and glia (Croset et al., 2018), the transcriptomic profile of T1 versus T2 neuronal and glial progeny has not yet been characterized. Furthermore, the T2 lineages generate diverse neuronal progeny (Hulse et al., 2020; Wang et al., 2014; Yang et al., 2013; Yu et al., 2013) but focused transcriptional profiling of T2-derived progeny has not been reported. In this report, we generate a single nuclei (snRNA-seq) atlas of T2-derived progeny of the adult central brain. We use several complementary methods to link neuron identity to UMAP clusters. Our data will facilitate linking neuronal transcriptomes with connectomes to gain a molecular understanding of CX development, connectivity, and behavior.

Results

Generation of a transcriptomic atlas of adult central brain neurons and glia

The adult central brain (i.e. brain without optic lobes) is composed of neuronal and glial progeny derived from both T1 and T2 NBs. We first generate a transcriptomic atlas of all central brain neurons derived from both T1 and T2 NBs. The flies were of a genotype that gave permanent lineage tracing of T2 NB-derived progeny in the adult (Figure 1A, A'). This allowed us to first analyze all central brain T1- and T2-derived neurons and glia (Figure 1) and then focus on neurons and glia produced specifically by T2-derived lineages.

We dissociated nuclei from 7-day-old adult fly central brains with optic lobes removed using methods previously described (Li et al., 2022; McLaughlin et al., 2021). The cDNA libraries were then prepared from dissociated nuclei with split-pool-based barcoding for single-nuclei transcription profiling (Rosenberg et al., 2018). We sequenced a total of 30,699 nuclei at 589 median genes per nuclei from the T1+T2 central brain.

To distinguish T2-derived progeny from T1-derived progeny, we expressed *worniu-gal4,asense-gal80* to drive expression of FLP recombinase (FLP) specifically in T2 NBs. This resulted in the flip out of a stop cassette and thus continuous expression of Gal4 under the ubiquitous *actin5C* enhancer. After removal of the stop cassette in T2 NBs, the *actin5c-gal4* can continuously drive expression of the reporter genes (RFP or GFP) and FLP in the T2-derived neuronal and glial progeny in adult brains (Figure 1A). We assigned 3,125 nuclei which have expression of RFP, GFP, or FLP as being derived from T2 lineages. We assigned 27,574 triple-negative nuclei as derived from the T1 lineages. We used Seurat for filtering, integrating, and clustering the atlas to provide transcriptionally unique cell clusters (Stuart et al., 2019). We used uniform manifold approximation and projection (UMAP)-based dimension reduction to visualize the 152 clusters (Figure 1B). The most enriched or least expressed genes (marker genes) for each cluster are shown in Supp. Table 1.

We used known markers (Croset et al., 2018; Davie et al., 2018) to identify distinct cell types in the central brain, including glia, mushroom body neurons, olfactory projection neurons, clock neurons, Poxn+ neurons, serotonergic neurons, dopaminergic neurons, octopaminergic neurons, corazonergic neurons, and hemocytes (Figure 1B). We found a cluster that represents optic lobe neurons, probably due to contamination during dissection when optic lobe tissue was not completely removed. We didn't observe any cluster exclusively containing progeny from T1 or T2 NB lineages (Figure 1C, Supp Figure 1). We conclude that both T1 and T2 generate transcriptionally similar cells, despite their different developmental origins. Alternatively, deeper sequencing may resolve a single cluster into two clusters with each derived from T1 or T2 lineages. We next searched for cluster-defining genes in the T2 enriched clusters (Figure 1C') and found high expression of *AstA* (*Allatostatin A*), *SerT* (*Serotonin transporter*), *Tk* (*Tachykinin*), and *Vmat* (*Vesicular monoamine transporter*) (Figure 1D). These enriched marker genes suggest that *AstA*+ neurons, serotonergic neurons, and *TK*+ neurons may be primarily derived from T2 lineages.

We explored the central brain atlas for glial cell types and their gene expression. We focused on the 3,409 glial nuclei from clusters that expressed the pan glial marker *repo* (Campbell et al., 1994; Xiong et al., 1994) in the T1+T2 atlas (Figure 1E). Each cluster contained a mix of T1 and T2 glial progeny (Figure 1E'). We identified six known glial cell types (astrocytes, cortex, ensheathing, surface glial and the two subtypes: perineurial and subperineurial) based on canonical glial makers (Figure 1F; Supp Table 2). We identified a glial cluster that expressed genes associated with extracellular matrix, *vkg* and *Col4a1* (Figure 1F), which have previously been identified as pan

surface glial markers (DeSalvo et al., 2014; Hindle and Bainton, 2014). Interestingly, the two surface glial subtypes, perineurial and subperineurial clusters, do not express these markers at cluster defining levels, and conversely the surface glia cluster 5 does not express perineurial or subperineurial specific makers (Figure 1F, Supp Table 3). Differential gene expression analysis between T1 and T2 glial progeny did not show differences across any glial cell types (Supp Table 4). We conclude that the adult central brain contains known glial cell types with no differences in gene expression between T1 and T2-derived glia.

Generation of a T1- and T2-specific cell atlas

We explored the diversity of T1 and T2-derived neurons by generating T1- and T2-specific cell atlases. We identified T1-derived neurons by *excluding* cells co-expressing T2-specific markers *FLP+/GFP+/RFP+* plus *repo+* glial clusters. We then generate a T1 neuron atlas containing 22,807 T1-derived neurons that form 114 clusters. Marker genes of each cluster are shown in Supp. Table 5. We identified the neurons that are known to be generated by T1 NBs, including MB neurons and olfactory projection neurons (Croset et al., 2018; Davie et al., 2018; Li et al., 2022). We identified other T1-derived neurons, including clock neurons, Poxn+ neurons, and neurons that release dopamine, serotonin, octopamine/tyramine, and other neuropeptides (Supp. Figure 2) (Croset et al., 2018; Davie et al., 2018). We gathered a list of genes that represented the 10 most significantly enriched genes from each cluster and calculated the scaled averaged expression of each gene from each cluster (Supp. Figure 2). We found that each cluster strongly expressed a unique combination of genes. We conclude that our T1-derived nuclei contain the expected abundance of neuronal diversity seen in previous scRNA-seq atlases (Croset et al., 2018; Davie et al., 2018; Li et al., 2022).

In our whole brain atlas described above, T2-derived cells represent a minor contribution to the atlas due to their low nuclei numbers relative to the T1-derived neurons (Raji and Potter, 2021). To generate a comprehensive T2 atlas, we needed to increase the percentage of T2-derived nuclei for RNAseq. We used fluorescence-activated cell sorting (FACs) to select nuclei that are labeled by T2-specific permanent lineage tracing (Figure 1A,B). We then selected the nuclei *in silico* that showed expression of UAS-transgenes (FLP, GFP, or RFP, see above) to eliminate contamination of T1 nuclei during FACS. In total, we sequenced 61,118 T2 nuclei. We included 3,125 UAS-transgene labeled nuclei from the dissociated central brain without sorting from the T1+T2 atlas (see above). We used integration in Seurat to generate an atlas containing 64,243 nuclei that formed 198 clusters. T2 NBs have been estimated to produced ~5000 neurons/glia in the central brain (Ito et al., 2013; Yang et al., 2013), with ~1,800 neurons in the central complex (Hulse et al., 2020; Schlegel et al., 2024), giving our atlas ~12x coverage. We next filtered out the *repo+* glial cell clusters to generate a T2 neuron cell atlas with 50,148 non-glial nuclei forming 161 clusters (Figure 2A). Marker genes of each cluster are shown in Supp. Table 6.

We calculated the scaled average expression of top 10 significantly enriched genes gathered from each T2 neuron cluster and found each T2 neuron cluster strongly expressed a unique suite of genes (Figure 2B). We conclude that each cluster within our T2 neuron atlas represents a transcriptionally unique cell type. For the remainder of the paper, we focus solely on the diversity of T2-derived neurons and glia.

T2 neuroblasts generate all major classes of fast-acting neurotransmitters

An important aspect of neuronal identity and function is fast-acting neurotransmitter expression. We determined the neuronal populations that expressed seven fast-acting neurotransmitters: glutamatergic, cholinergic, GABAergic, tyraminergic, dopaminergic, serotonergic and octopaminergic in T2-derived neurons (Figure 3A-G). In neurons, we found that cholinergic neurons were most abundant (21%), followed by glutamatergic neurons (12%), GABAergic neurons (9%), tyraminergic neurons (2.6%), dopaminergic neurons (1.2%), serotonergic neurons (1.2%), and octopaminergic neurons (0.7%). Additionally, 12% of neurons were co-expressing 2 or 3 neurotransmitters. The remaining 40% in our atlas were not expressing any neurotransmitter with a log normalized expression <2 (Figure 3H); this is similar to the ratios in the adult midbrain (Croset et al., 2018) and optic lobes (Konstantinides et al., 2022). The nuclei co-expressing two or more fast-acting neurotransmitters may reflect an authentic feature of these neurons, as see in other systems (see Discussion). Alternatively, our double- and triple-positive neurons may be false positives (see Discussion).

T2 neuroblasts generate all major classes of glia

We next explored the complete T2 atlas for glial cell types and their gene expression. We sub-clustered from the T2 atlas for 12,315 glial nuclei from clusters that expressed the pan glial marker *repo* (Figure 4A) (Campbell et al., 1994; Xiong et al., 1994). Similar to our T1+T2 glial atlas, we identified glial cell types: astrocytes, cortex, ensheathing, astrocyte-like, and two surface glial subtypes, perineurial and subperineurial, based on canonical glial makers (Figure 4B; Supp. Table 7). Interestingly, we identified chiasm glia that may represent the migratory chiasm glia generated from the DL1 lineage (Viktorin et al., 2013; see Discussion). We conclude that our T2 atlas contains the expected glial cell types.

T2 neuroblasts generate sex-biased cell types

We noticed that several cell types in the T2 glial atlas showed unequal number of male and female nuclei (Figure 5A-B, E-F). We identified sex-biased clusters by normalizing the number of input nuclei between male and female samples and compared the proportion of nuclei by sex within each cluster to identify sex-biased clusters (Figure 5B,F). We found that the T2 glial atlas contained two female enriched clusters: ensheathing/astrocyte (cluster 7) and chiasm (cluster 5), and one male enriched cluster: the astrocyte-like (cluster 3) (Figure 5B). We determined differential genes expressed between male and female nuclei across all glia (Figure 5C; Supp Table 8). We found female nuclei expressed higher levels of genes including the female-specific genes *yp1, yp2,* and *yp3* (Figure 5C)(Warren et al., 1979). Male nuclei expressed higher levels of genes including the male-specific genes lncRNA:rox1/2 (Figure 5C) (Amrein and Axel, 1997; Meller et al., 1997). We next looked at expression within the sex-biased clusters. We found similar results that male nuclei expressed high levels of the male-specific *lncRNA:roX1/2* and low expression of the female-specific

genes *yp1, yp2,* and *yp3* compared to the female nuclei (Figure 5D; see Discussion). We conclude that male and female adult T2 glia have sex-specific differences in gene expression within the same glial cell type.

We next explored sex differences among neurons (Figure 5E-E"). We identified 14 neuronal clusters with disproportionate amounts of male and female nuclei (Figure 5F). We first tested the differential genes expressed between male and female nuclei across all T2-derived neurons (Figure 5G; Supp Table 9). Similar to the T2 glia sex differences, we found female nuclei expressed higher levels of the female-specific genes *yp1, yp2,* and *yp3* (Figure 5G) (Warren et al., 1979). Male nuclei expressed higher levels of the male-specific genes lncRNA:rox1/2 (Figure 5C) (Amrein and Axel, 1997; Meller et al., 1997). We next looked at expression levels within the sex-biased clusters. We found similar results that male nuclei expressed high levels of the male-specific *lncRNA:roX1/2* and low expression of the female-specific genes *yp1, yp2,* and *yp3* compared to the female nuclei (Figure 5H; see Discussion). We conclude that male and female adult T2 neurons have sex-specific differences in gene expression within the same neuronal subtype.

T2 neuroblasts generate neurons expressing a diverse array of neuropeptides

Neuropeptides are often used as markers to distinguish different neuronal populations. We examined whether individual neuropeptide-encoding genes were exclusively expressed within single clusters, representing single neuronal subtypes. Among the 49 neuropeptides in *Drosophila*, we identified 13 with enriched expression in a limited number of clusters, making them suitable as cluster-defining markers (Figure 6A). For all clusters significantly expressing cluster-defining neuropeptides, we assessed their co-expression with specific fast-acting neurotransmitters (Figure 6A). We found that all neuropeptide-expressing clusters, co-express one or more neurotransmitters (see Figure 3H).

These data raise several questions. What is the relationship between neuropeptide expression and cluster identity? Can two clusters express the same neuropeptide? Can two neuropeptides be co-expressed by the same cluster? Our data show that all of these patterns exist. For example, in some cases, one cluster expresses two or more neuropeptide genes (e.g. clusters 100 and 128 express NPF and AstA); conversely, we also observe one neuropeptide expressed in multiple clusters (e.g. Tk, 2 clusters; Mip, 4 clusters) (Figure 6A).

Determining the relationship between cluster identity, neuropeptide expression and neuronal morphology required a method for specifically labelling of T2 peptidergic neurons, as antibody staining could be contaminated with T1-derived neurons. We developed a genetic approach to selectively visualize neuropeptide expression exclusively in neurons originating from T2 NB lineages. This technique involves driving FLP recombinase (FLP) under the control of the T2-specific driver *wor-GAL4,ase-GAL80*. The T2-specific FLP catalyzes the excision of a stop codon in the *lexAop-FRT-stop-FRT-myr::gfp* transgene, allowing existing neuropeptide-2A-LexA transgenes (Deng et al., 2019) to drive GFP expression specifically in neuropeptide-positive T2-derived neurons (Figure 6B). This method revealed the morphology of specific T2-derived peptidergic neurons.

We first assayed neuronal morphology of Ms+ neurons in clusters 157 and 160. We found that Ms-2A-LexA-expressing neurons project to two regions of the central complex: the entire ellipsoid body and the dorsal fan-shaped body (Figure 6C-C", Supp. Movie 1). Currently we can't determine whether Ms+

neurons in clusters 157 and 160 project to different CX neuropils, or whether neurons from both clusters
share projections into both neuropils.

Next, we assayed neurons in two clusters (100 and 128) that express the same two neuropeptides,
AstA and NPF (Figure 6D-E). We wanted to determine if these clusters contain two cell types with similar
gene expression, or a single cell type that co-expressed both neuropeptides. We used NPF-2A-lexA and
AstA-2A-lexA to map the morphology of each class of neurons. We found that they were generated at
indistinguishable numbers and morphology: both had ~20 neurons projecting to a lateral domain of the
ellipsoid body and the same two layers of the fan-shaped body (Figure 6D-E; Supp. Movies 2-3). We
conclude that similar neurons can co-express two different neuropeptides.

We investigated whether specific transcription factors (TFs), or TF combinatorial codes, might
correlate with the expression of cluster-defining neuropeptides, and thus provide candidates for a
regulatory relationship where each cluster may have a TF code that drives transcription of a specific
neuropeptides. To address this, we performed an unbiased correlation analysis of TF and neuropeptide
expression patterns across all cells in our dataset to identify potential regulatory relationships.
Interestingly, we found high correlations of multiple TFs to each neuropeptide (Figure 6F, Supp. Fig 3, Supp.
Table 10). Our data provide a stepping-stone to the analysis of upstream TF combinatorial codes that drive
neuropeptide expression.

Each neuron cluster consists of a unique combination of transcription factors

It is generally thought that neuronal diversity is generated by TF combinatorial codes that are
uniquely expressed in neuronal subtypes, with an emphasis on homeodomain TFs (Reilly et al.,
2020; Sagner et al., 2021). Thus, we assayed each class of TFs to determine how many were
expressed in cluster-specific (i.e. neuron-specific) patterns. We found that most clusters expressed
a unique TF combination (Figure 7; Supp. Table 11). This is true for all zinc-finger TFs (Figure 7A) and
helix-turn-helix TFs (Figure 7B). Nearly all clusters expressed a unique combination of
homeodomain TFS (Figure 7C), with only a few clusters sharing a homeodomain code (Figure 7C').
In contrast, basic domain, unspecified domain, and high mobility group TFs were more
promiscuously expressed (Figure 7D). We conclude that zinc finger, helix-turn-helix, and
homeodomain TFs are good candidates for forming unique combinatorial codes that should be
tested for a role in generating neuronal diversity.

Linking neurons to UMAP clusters

We sought to link well characterized CX neurons to their transcriptomes within the T2 atlas. We chose
columnar neurons due to known cell type markers and genetic access. We used four complementary
approaches to identify them, described below.

(1) We assigned identities of neuronal clusters based on the expression profiles of neuropeptides and
neurotransmitters (Figure 6) characterized in the literature (Wolff et al., 2024). We further examined these
neurons of their expression of selective TFs and markers, including *bsh*, *cut*, *DIP-β*, *Eip93F*, *Imp*, *runt*, *syp*,
toy, and *zfh2*. The results are summarized in Supp Table 11.

(2) We used a published bulk RNAseq dataset to profile selective CX neurons (Turner-Evans et al., 2020; Wolff et al., 2024) and compared the expression profiles to our snRNA-seq dataset with a published algorithm (Davis et al., 2020) (Figure 8A). We generated a heatmap based on the coefficient of the expression profiles to predict the identity of each cluster. We predicted that E-PG neurons were in cluster 75 and 95, P-EG neurons were in cluster 102, hDeltaK neurons were in cluster 100, FB6A were in cluster 162, and FB7A were in cluster 40 (Figure 8A). The prediction of E-PG neurons arising from cluster 75 and 95 was validated by expression of the transcription factor *dac* (Dillon and Doe, 2024); other neuron-cluster assignments await experimental validation.

(3) We assayed the enhancers in highly specific split-Gal4 lines to determine if the associated genes were expressed in the same neuron subtype as the split-Gal4 line (Figure 8B). For example, the stable split line SS65380 is expressed in the FB2A neurons, and the enhancer DNA used to make the split lines are associated with the *wnt10* and *lmpt* genes. These two genes show high expression in cluster 149 (Figure 8B), allowing us to predict that the FB2A neurons are in cluster 149. A similar approach was used to generate candidate neuron-cluster predictions for the following neurons: FB8G, FB6H, hDeltaA, hDeltaD, hDeltaH, and FB1C (Figure 8B). These predictions can be used to focus experimental validation on a small group of neurons; however, the percentage of validated neurons-clusters awaits experimental analysis.

(4) The fourth approach was based on using known or novel markers of CX neurons together with Gal4/LexA lines that are regulated by known gene enhancers. TFs known to be expressed by PF-R, P-EN and P-FN neurons are Toy and Runt, respectively (Sullivan et al., 2019). We assigned PF-R neurons to cluster 105 based on being Toy+ Runt- and being labeled by Gal4/LexA driven by the enhancer of *rho* gene (Figure 8C, left and middle columns). We assigned P-FN to clusters 38, 46, and 123 based on being Toy- Runt+ and being labeled by split line SS00191 associated with *shakB* and *Pkc53E* genes. Lastly, we assigned P-EN neurons to clusters 66 and 130 by their expression of Toy- Runt+ and labeled by *Gɣ30A-lexA* (Figure 8C, left and middle columns). This approach identified novel TF markers for P-EN, PF-R and P-FN neurons (Figure 8C, right columns), which we validated by antibody staining (Figure 8D-H) (Dillon et al., 2024). The results of cluster identity assignment are summarized in Figure 9 (Supp Table 12).

The differential expression of TFs probably contributes to the morphological or connectivity diversity found within each class of CX neurons (Hulse et al., 2021; Turner-Evans et al., 2020; Wolff and Rubin, 2018; Wolff et al., 2015). We conclude that our T2 atlas can be used to link known neuron subtypes to their transcriptome (Figure 9). This is a necessary step in determining how cluster-specific gene expression regulates neuron-specific functional properties such as morphology, connectivity, and physiology.

Discussion

We used a genetic approach to express RFP in the progeny of T2 NBs in the adult central brain. We excluded optic lobes by manual dissection, and thus our data are complementary to recent scRNA-seq data from the adult optic lobe (Konstantinides et al., 2022). Similarities include neurons

expressing predominantly single fast-acting neurotransmitters (discussed below). Both datasets
show major classes of neurons including MB neurons, olfactory projection neurons, clock neurons,
Poxn+ neurons, serotonergic neurons, dopaminergic neurons, octopaminergic neurons,
corazonergic neurons, and hemocytes. As expected, we did not detect neurons expressing motor
neuron markers. The MB neurons that appear in the T2-derived progeny may be due to off-target
expression of our Gal4 driver at stages we have not assayed, or more likely, due to ambient RNA
released during the dissociation and sorting steps.
342
Interestingly, we observed fewer distinct neuronal clusters in the T1-derived population (114
clusters; Supp. Figure 2) than in the T2-derived neuronal population (161 clusters; Figure 2). This
could be due to the T1-derived progeny containing a large population of transcriptionally similar
Kenyon cells of the MB, whereas T2-derived clusters contain fewer, but transcriptionally more
diverse, cells than T1-derived progeny. It would be interesting to know whether T1 NB lineages are
generally less diverse, or whether the Kenyon cells are exceptional in their lack of transcriptional
diversity, despite the presence of protein gradient in young vs. old neurons (Liu et al., 2015). Our
results are consistent with lineage analysis of T2 NBs, which make distinct neurons across the
temporal axis of T2 NBs, plus unique neurons across the INP temporal axis (Doe, 2017; Ito et al.,
2013; Sullivan et al., 2019; Wang et al., 2014; Yang et al., 2013).
353
354 *Neurotransmitters and neuropeptides*
355
We observed that most fast-acting neurotransmitters are uniquely expressed in neuronal
populations, but we also observe a smaller group of neurons that express two or three
neurotransmitters. This includes excitatory and inhibitory neurons, e.g. 2.9% of all neurons express
genes necessary for cholinergic and GABAergic neurotransmitters (Figure 3). Co-expression of
neurotransmitters has been reported in other systems (Granger et al., 2020; Granger et al., 2023;
Lozovaya et al., 2018; Meye et al., 2016; Saunders et al., 2015; Shabel et al., 2014; Spitzer, 2015;
Takács et al., 2018) and it may be biologically relevant in the T2-derived neurons. Alternatively, co-
expression may be due to background from RNAs released during dissociation.
Neuropeptides are profoundly important for a vast array of behaviors (Schoofs et al., 2017), and
thus our mapping of neuropeptide expression to distinct neuronal subtypes may facilitate a
functional analysis linking neuron identity, neuropeptide expression, and behavior.
367
368 *Glial identities*
369
We identified six classes of glia in the T1+T2 glial atlas that reflects previous scRNAseq datasets
(Figure 1E-F) (Konstantinides et al., 2018; Lago-Baldaia et al., 2023). Interestingly, we detected a
small population of *vkg+* surface glia specific to the whole atlas (T1+T2) and not the T2 glia atlas.
The lack of differential gene expression between T1- and T2-derived glia suggests these cell types
are similar despite different developmental origins. It will be for future work to understand how
different progenitors produce similar glial fates.

Similar to the T1+T2 glial atlas, our T2 atlas captured the known glial subtypes found in other scRNAseq datasets (Figure 4) (Konstantinides et al., 2018; Lago-Baldaia et al., 2023). The T2 glia atlas contains more clusters than the T1+T2 atlas that may be due to the nearly 4x the number of glia captured (T1+T2 atlas 3,409 nuclei; T2 atlas – 12,315 nuclei). Interestingly, we identified the astrocyte-like glia of the central brain previously described (Awasaki et al., 2008). Additionally, we captured chiasm glia that have been shown to be derived from the T2 lineage DL1 and migrate into the optic lobe (Viktorin et al., 2013). These nuclei could represent either chiasm glia that did not migrate out of the central brain or optic lobe tissue that was not completely removed during dissections. It will be interesting to investigate how these T2-derived migratory chiasm glia compare to the optic-lobe derived chiasm glia, because of their different developmental origins.

Sex differences found in the adult neurons and glia

The T2 glia and neuron atlases contained clusters with disproportionate enrichment of female and male nuclei when normalized for sample input. We found the expected differential expression of the yolk proteins (*yp1, yp2, yp3*) enriched in female nuclei and *rox1/2* enriched in male nuclei (Amrein and Axel, 1997; Meller et al., 1997; Warren et al., 1979). Surprisingly, we found additional female and male-specific gene enriched in both neurons and glia including *ATPsyndelta* and undescribed, computationally predicted genes (CGs) (Figure 5). These genes may reflect sex-specific differences in the adult central brain and may provide insight into how behavioral circuits are linked to sex-specific behaviors. Future work should aim to characterize and test these genes.

TF codes

Our atlas showed zinc finger TFs and homeodomain TFs as forming combinatorial codes specific for distinct cell types. This raises the possibility that these TF combinations determine neuronal functional properties, by establishing and maintaining neuronal identities. This is consistent with data from *C. elegans* (Hobert, 2021; Reilly et al., 2020), mammalian spinal cord (Briscoe et al., 2000; Sagner et al., 2021), *Drosophila* optic lobe (Holguera and Desplan, 2018; Konstantinides et al., 2018), leg motor neurons (Baek et al., 2013; Enriquez et al., 2015), and VNC (Soffers et al., 2024). Interestingly, we found more zinc finger TFs expressed in unique combinatorial codes compared to homeodomain TFs; this may be a caution to focusing too narrowly on homeodomain TFs function. We note that our results are mostly correlative and await functional analysis of the TF combinatorial codes found in our dataset. It is also possible that the current sparse TF expression patterns are due, in part, to false negative results due to low read depths. This can be resolved by increasing read depth or by performing functional assays.

Mapping neurons to clusters

In this study, we used several complementary approaches to map identified neurons to their transcriptomic UMAP cluster. First, we used bulk-seq data from specific neurons with our atlas to match neuron to cluster. Second, we used enhancer elements from neuron-specific split Gal4 lines

to link enhancer-associated genes to each hemi-driver with common expression in a cluster. Third,

we used known or novel TFs and other molecular markers that are expressed by a cluster as an

entry point to look for other cluster specific genes. Each of these methods provides candidate

neuron-cluster association, which requires experimental validation. It is not clear which method is

more robust at identifying functional neuron-cluster associations. Nevertheless, we have validated

a number of neuron-cluster associations using the third approach: beginning with a single gene-

cluster correlation. In the future, our data can be used in two pipelines for linking neuron subtype to

transcriptome. First, start with at least one validated marker for a cluster, then search all labeled

clusters for candidate TFs that co-express with the validated marker, followed by validation via

RNA in situ (or antibodies). Second, we can collect neurons of interest with FACS or antibody

selection (Davis et al., 2020), and perform bulk RNA sequencing, followed by looking for clusters

enriched for the TFs found in bulk sequencing data, and then validation via RNA in situ (or

antibodies). Notably, the second pipeline requires no prior markers beyond the Gal4 or LexA line

expressed in single neuronal populations. Both pipelines result in a transcriptome that can be used

to identify (a) TFs for their role in neuronal specification and/or maintenance, (b) cell surface

molecules that may regulate neuronal morphology and/or connectivity, or (c) functionally relevant

genes encoding ion channels, neuropeptides, receptors, and signaling pathways.

Figure 1. Cell atlas of central brain with single nuclei RNAseq.
(**A-A'**) Genetics (A) to label progeny (A') derived from T2 NBs. Dashed lines show the boundary of optic lobes and central brain, and the optic lobes were removed during dissection.
(**B**) Central brain atlas labeled with known cell types. Abbreviations: CLK, clock neurons; CRZ, Corazonergic neurons; DOP, dopaminergic neurons; HEM, hemocytes; MBN, mushroom body neurons; OCTY, octopaminergic-tyraminergic neurons; OL, optic lobe; OPN, olfactory projection neurons; SER: serotoninergic neurons.
(**C**) Central brain atlas labeled by NB (T1 or T2) lineage. Dash line-outlined box shows the region enriched with the cells derived from T2 NBs, and the identity are shown at the bottom-right box.
(**D**) Dot plot of top 5 marker genes of the T2-enriched clusters.
(**E**) Atlas of central brain glia labeled with known cell types. (**E'**) T1 and T2 derived cells colored cyan and red respectfully.
(**F**) Dot plot of known and top marker genes of glial clusters.

Figure 2. Cell atlas of neurons produced by T1 or T2 NBs.
(**A**) Cell atlas from T2 NBs. (**B**) Heatmap of scaled average expression of top 10 markers genes from each T2 neuronal cluster.

Figure 3. Expression of fast-acting neurotransmitters in neurons derived from T2NSC lineages.
(**A-G**) UMAP distribution plots demonstrate the expression of the following neurotransmitters: (**A**) *vesicular glutamate transporter* (VGlut, glutamatergic neurons), (**B**) *vesicular acetylcholine transporter* (VAChT, cholinergic neurons), (**C**) *glutamic acid decarboxylase 1* (Gad1, GABAergic neurons), (**D**) *tyramine β hydroxylase* (Tbh, tyraminergic neurons), (**E**) *tyrosine 3-monooxygenase* (Ple, dopaminergic neurons), (**F**) *serotonin transporter* (SerT, serotonergic neurons) (**G**) *tyrosine decarboxylase* (Tdc2, octopaminergic neurons). All plots have a minimum cutoff value set at 0.
(**H**) The UpSet quantifies the number of cells in the atlas that express each neurotransmitter gene with a scaled expression > 2 (Gu, 2022).

Figure 4. Glial cell types present in the T2-derived cell atlas.
(**A**) Sub-clustered of 12,315 nuclei from Repo+ T2 clusters in UMAP distribution.
(**B**) Dot plot of validated glial subtype markers to identify glial clusters by differential gene expression.

Figure 5. Sex differences present in the T2-derived glia and neurons.
(**A-A''**) T2 glia in UMAP distribution across samples: (**A**) T2 Female (3,910 nuclei), (**A'**) T2 Male (4,385 nuclei), (**A''**) Female and Male mixed (4,020 nuclei).
(**B**) Biased clusters for male to female ratio for number of nuclei in the T2 glia clusters.
(**C**) Differential expression between male and female T2 glia.
(**D**) Heatmap of top differential gene expression between male and female nuclei within glial clusters.
(**E-E''**) T2 neuron in UMAP distribution across samples: (**E**) T2 Female (8,151 nuclei), (**E'**) T2 Male (16,201 nuclei), (**E''**) Female and Male mixed (25,796 nuclei).
(**F**) Biased clusters for male to female ratio for number of nuclei in the T2 neuron clusters.
(**G**) Differential expression between male and female T2 neurons.
(**H**) Heatmap of top differential gene expression between male and female nuclei within neuron clusters.

Figure 6. Neuropeptide expression in T2-derived neurons.
(**A**) Dot plot showing expression of 13 neuropeptides and their co-expression with the 7 fast-acting neurotransmitters (red dashed line) across selected clusters.
(**B**) Genetic scheme to map neuropeptide expression in T2-derived adult neurons.

(C-E) Neuropeptide-expressing neurons labelled with GFP in three-dimensional projections. **(C'-E')** Fan-shaped body projections. **(C"-E")** Ellipsoid body projections. nc82 counterstains (magenta) in the brain for neuropil projections.

(F) Heatmap showing the top 5 transcription factors most strongly correlated with each neuropeptide across all cells (Sigorelli, 2024). Scale bar represents 20 μm.

Figure 7. Combinations of transcription factor expression in T2-derived neurons.
Binarized heatmaps of positive **(A)** zinc finger **(B)** helix-turn-helix **(C)** homeodomain TF markers in T2-derived neurons. Clusters are sorted by similarity based on Jaccard index scores.
(C`) Top five marker genes for clusters which had non-unique combinations of homeodomain TF expression.
(D) Percentage of clusters with unique combinations of TF expression based on TF class (zinc finger TFs = 100% (161 unique clusters), helix-turn-helix TFs = 100% (161 unique clusters), homeodomain TFs = 93.8% (150 unique clusters), basic domain TFs = 69.6% (112 unique clusters), unidentified DNA binding domain TFs = 59% (95 unique clusters), high motility group TFs = 34.2% (55 unique clusters).

Figure 8. Mapping T2-derived neurons to UMAP clusters.
(A) Heatmap of the coefficients (r) of expression profiles of single cell clusters (columns) to the expressions of known central complex neurons (rows) profiled with bulk RNA sequencing. The matching of the cluster to the central complex neurons is determined by the highest coefficient value (gray box) for each cluster. FB6A(1) and FB6A(2) are labeled and profiled with two different split-gal4 drivers (Wolff et al., 2024). Cluster 75 and 95 identified as E-PG were verified in Dillon et al., 2024.
(B) Scaled average expression of the enhancer genes of split-gal4 drivers (columns) in each cluster (rows). The name of central complex neurons (CX) labeled by split-gal4 drivers are listed at the top. The cluster identity was determined by the combination of positive scaled average expression of the enhancer genes (black boxes).
(C) Dotplot of the known marker genes (*toy*, *runt*) expressed in central complex neurons (PF-R, P-FN, and P-EN), lexA-driver enhancer gene (*rho*, *Gγ30A*), split-gal4-driver (SS00191) enhancer genes (*shakB*, *Pkc53E*), and newly identified marker genes.
(D) PF-R (*R37G12-Gal4 UAS-V5*) neuronal cell bodies labeled with V5, and co-stained with marker genes boxed in **(C)**.
(E,F) PF-N (*R16D01-Gal4 UAS-V5*) neuronal cell bodies labeled with V5, and co-stained with marker genes boxed in **(C)**.
(G) PE-N (*R12D09-Gal4 UAS-V5*) neuronal cell bodies labeled with V5, and co-stained with marker genes boxed in **(C)**.
(H) Percentage of marker genes in PE-N and PF-N. Scale bar: 5 μm in all panels.

Figure 9. Summary.
Asterisk, known neuron-cluster associations; others are predicted neuron-cluster associations. Underline, neurotransmitter expressing clusters. Abbreviations: *AstC*, Allatostatin C+ neurons; *CCAP*, *Crustacean cardioactive peptide*+ neurons; DOPA, dopaminergic neurons; *dsx*, *doublesex*+ neurons; MBN, mushroom body neurons; OCTY, octopaminergic-tyraminergic neurons; *Proc*, Proctolin+ neurons; *Tk*, Tachykinin+ neurons; SER, serotonergic neurons. E-PG, FB1C, FB2A, FB2B, FB2l_ab, FB3C, FB4K, FB4L, FB6A, FB6H, FB7B, FB8B, FB8G, FR1, hΔA, hΔD, hΔE, hΔH, hΔI, hΔK, lbSps-P, P-EG, P-EN, P-FGs, P-FN, PF-R, vΔC and vΔE are the names of central complex neurons (Hulse et al., 2021).

Supplemental Figure 1. Nuclei numbers T1 vs T2. Vertical shades indicate clusters with T2/(T1+T2) > 50%.

Materials and Methods

Fly genetics

20xUAS-FLPG5.PEST;worniu-gal4,asense-gal80; Act5c(FRT.CD2)gal4 (Syed et al., 2017); *20xUAS-FLPG5.PEST;worniu-gal4,asense-gal80; lexAop(FRT.stop)-mCD8:GFP* (this study); *Tl{2A-lexA::GAD}AstA[2A-lexA]/TM3,Sb[1]* (RRID:BDSC_84356); *Tl{2A-lexA::GAD}Ms[2A-lexA]/TM3,Sb[1]* (RRID:BDSC_84403); *Tl{2A-lexA::GAD}NPF[2A-lexA]/TM3,Sb[1]* (RRID:BDSC_84422); *UAS-RedStinger* (RRID:BDSC_8545); *UAS-unc84-2xGFP* (Henry et al., 2012); *GMR12D09-lexA/CyO* (RRID:BDSC_54419); *GMR16D01-lexA* (RRID:BDSC_52503); *GMR37G12-lexA* (RRID:BDSC_52765); *13xLexAop2-IVS-myr::smGdP-V5* (RRID:BDSC_62215).

Single nuclei isolation, library preparation, and sequencing

We first used the split-pool method (Parse Bioscience, Seattle, WA, USA) to barcode RNAs from whole brain to generate a library which includes both T1 and T2 progeny. This library, T1+T2, was used for analysis in Figure 1 and 5. To increase the number of T2 nuclei for snRNAseq, we labeled T2 nuclei by crossing *20xUAS-FLPG5.PEST;worniu-gal4,asense-gal80; Act5c(FRT.CD2)gal4* (Syed et al., 2017) to *UAS-RedStinger* (RFP) or *UAS-unc84-2xGFP* (GFP) flies. The adult flies were aged for 1 week at 25 °C before dissection. Equal amounts of male and female central brains were dissected (without optic lobes) at room temperature within 1 hour, flash-frozen in liquid nitrogen, and then stored at -80 °C. Nuclei dissociated from the frozen, dissected brains was performed according to published protocols (McLaughlin et al., 2022). RFP+ or GFP+ nuclei were collected by sorting dissociated nuclei with SONY-SH800 with 100 µm chip. We performed three rounds of sorting and snRNAseq. In the first round, we select GFP+ nuclei and used particle-templated instant partitions to capture single nuclei to generate cDNA library (Fluent BioSciences, Waterton, MA). In the second and third rounds we selected RFP+ nuclei and used split-pool method to generate barcoded cDNA libraries from each individual nucleus. All libraries were sequenced with pair-ends reads 150bp on Illumina Novaseq 6000 (University of Oregon's Genomics and Cell Characterization Core Facility).

snRNA-seq analysis

Our bioinformatic analysis was performed using pipeline from Fluent BioSciences , Parse Bioscience, and the Seurat R package (Hao et al., 2024; Satija et al., 2015). Briefly, pipelines from Fluent BioSciences and Parse Biosciences were used to perform demultiplexing, alignment, filtering, counting of barcodes and UMIs with an output being a cell-by-genes matrix of counts. We aligned our sequences to a custom reference genome by adding *flpD5* (RRID:Addgene_32132), *redstinger* (RRID:Addgene_46165), and *unc84sfGFP* (RRID:Addgene_46023) (+SV40 tail) sequences and annotations to the *Drosophila* genome release BDGP6.32.109. To further ensure that only high-quality cells were retained, we removed any cells with fewer than 200 genes or more than 2500 genes expressed and more than 5% mitochondrial RNA. For the T2 atlas, snRNA-seq data from three rounds of sorting and barcoding were integrated in Seurat with Anchor-based RPCA integration to generate an integrated dataset, and the downstream analysis was performed with the default parameters. The UMAP was generated with resolution 12. The integrated dataset was used for analyses.

Glia analyses

The T1+T2 glia atlas dataset was derived from the T1+T2 *repo*+ glial clusters and re-clustered to represent the glial subtypes. The T2 glia atlas was derived from the T2 *repo*+ glial clusters and re-clustered to represent the glial subtypes. Cell identity was determined by the validated markers for glia shown to differentially expressed between the clusters. Identities were assigned based on

625   expression of these validated markers. Overlapping identities were assigned if multiple subtype
626   markers were expressed within a single cluster.
627
<u>Sex differences</u>
We determined sex-biased clusters within the T2 glia and neuron atlases by identifying clusters that
were disproportionate after normalizing the number of inputs for 'female' and 'male' samples
respectively. The 'female and male' mixed samples were excluded from the analyses as we could
not differentiate the sex origin of these nuclei. To determine differential gene expression between
females and males, we pseudo bulked nuclei by aggregating the snRNAseq for comparison
between sex and clusters. Heatmaps were generated in Seurat using the Scillus package
(https://github.com/xmc811/Scillus) to display the heatmap.
636
<u>Transcription factor combinatorial analysis</u>
We used the Seurat function FindAllMarkers to find positively differentially expressed TFs in all clusters. We
then removed TFs which were not significantly differentially expressed. We gave a value of 1 to any TF that
was found to be a positive marker in each cluster and a value of 0 in clusters which it was not a positive
marker. Next, we found the number of unique combinations of markers for 6 classes (zinc finger, helix-turn-
helix, homeodomain, basic domain, unidentified DNA binding domain, and high motility group) of TFs. The
Jaccard index between each cluster was then calculated and clusters were sorted from most to least similar.
The python packages Matplotlib and Seaborn were then used to generate heatmaps.
645
646   <u>Data and code availability</u>
647   All data and code can be found at https://github.com/dgepiney/2023_Doe_Drosophila_
648   Central_Brain_RNAseq/.
649
<u>Immunohistochemistry and imaging</u>
Standard methods were used for adult brain staining (Sullivan et al., 2019). Primary antibodies were
chicken anti-V5 at 1 µg/mL (Fortis Life Sciences, Waltham, MA), rabbit anti-Lim1 (1:500) (gift from
C. Desplan, New York University), guinea pig anti-Rx (1:500) (gift from C. Desplan, New York
University), and mouse anti-Cut at 2 µg/mL (2B10, DSHB, Iowa City, IA) . Secondary antibodies
were donkey whole IgG antibody conjugated with DyLight405, Alexa Fluor 488, Rhodamine Red-X
(RRX), or Alexa Fluor 647 (Jackson ImmunoResearch Laboratories Inc., West Grove, PA) and used at
5 µg/mL. The antibody stained brains were mounted in DPX (https://www.janelia.org/project-
team/flylight/protocols) on poly-L-Lysine coated coverslip (Corning, Glendale, AZ) and imaged with
Zeiss confocal LSM800 with software Zen.
660
661   <u>Figure production</u>
662   We used Imaris (Bitplane, Abingdon, UK) for confocal image processing, and Illustrator (Adobe, San
663   Jose, CA) to assemble figures.
664
665   **Ethical Approval and Consent to participate**
666   No vertebrate animals were used.
667
668   **Consent for publication**
669   All authors consent for this to be published
670
671   **Availability of data and materials**

672 All data and materials will be placed in the appropriate public repository (e.g. github or Bloomington stock
673 center) upon acceptance for publication
674
Authors' contributions
DE, NRD, GNMC and SL performed dissections.
DE, GNMC, and SL performed FACS, and RNAseq data integration.
NRD generated Figure 1, 4, 5, ST3, ST4, ST5, ST7, ST8, and ST9.
GNMC generated Figures 3, 6, SF3, ST10, SM1, SM2, and SM3.
DE generated Figures 7, 8, and ST12.
SLL generated Figures 1, 2, 8, 9, SF1, SF2, ST1, ST2, ST6, ST11, and ST12.
All authors contributed to writing the manuscript. CQD supervised the project.
683

684 **Acknowledgements**
685 We thank Jason Carriere, and Peter Newstein for comments on the manuscript. We thank the
686 Desplan lab (NYU) for reagents. We thank Jason Carriere (University of Oregon Genomics and Cell
687 Characterization Core Facility) for technical assistances in cDNA library preparation and
688 sequencing. Flies were obtained from the Bloomington Stock Center.
689

690 **Competing Interests**
691 None.
692

693 **Funding**

Funder	Grant reference number	Author
Howard Hughes Medical Institute	None	CQD, SLL, NRD, DE
NIH Training Grant	5-T32-HD07348	NRD
NIH Training Grant	5-T32-GM149387	GNMC
NIH	R01-HD27056	CQD, NRD, DE, GNMC

699 The funders had no role in study design, data collection and interpretation, or the decision to
700 submit the work for publication.
701

Author ORCIDs
Derek Epiney   https://orcid.org/0000-0002-6056-1817
Gonzalo N Morales Chaya   https://orcid.org/0009-0005-4263-1323
Noah R Dillon   https://orcid.org/0000-0002-6708-4208
Sen-Lin Lai   https://orcid.org/0000-0002-7531-283X
Chris Q Doe   https://orcid.org/0000-0001-5980-8029

References

Amrein, H. and Axel, R. (1997). Genes expressed in neurons of adult male Drosophila. *Cell* **88**, 459–469.

Awasaki, T., Lai, S.-L., Ito, K. and Lee, T. (2008). Organization and postembryonic development of glial cells in the adult central brain of Drosophila. *J. Neurosci. Off. J. Soc. Neurosci.* **28**, 13742–13753.

Baek, M., Enriquez, J. and Mann, R. S. (2013). Dual role for Hox genes and Hox co-factors in conferring leg motoneuron survival and identity in Drosophila. *Development* **140**, 2027–38.

Bello, B. C., Izergina, N., Caussinus, E. and Reichert, H. (2008). Amplification of neural stem cell proliferation by intermediate progenitor cells in Drosophila brain development. *Neural Dev* **3**, 5.

Boone, J. Q. and Doe, C. Q. (2008). Identification of Drosophila type II neuroblast lineages containing transit amplifying ganglion mother cells. *Dev Neurobiol* **68**, 1185–95.

Bowman, S. K., Rolland, V., Betschinger, J., Kinsey, K. A., Emery, G. and Knoblich, J. A. (2008). The tumor suppressors Brat and Numb regulate transit-amplifying neuroblast lineages in Drosophila. *Dev Cell* **14**, 535–46.

Boyan, G. and Williams, L. (2011). Embryonic development of the insect central complex: insights from lineages in the grasshopper and Drosophila. *Arthropod Struct. Dev.* **40**, 334–348.

Briscoe, J., Pierani, A., Jessell, T. M. and Ericson, J. (2000). A Homeodomain Protein Code Specifies Progenitor Cell Identity and Neuronal Fate in the Ventral Neural Tube. *Cell* **101**, 435–445.

Campbell, G., Goring, H., Lin, T., Spana, E., Andersson, S., Doe, C. Q. and Tomlinson, A. (1994). RK2, a glial-specific homeodomain protein required for embryonic nerve cord condensation and viability in Drosophila. *Development* **120**, 2957–66.

Croset, V., Treiber, C. D. and Waddell, S. (2018). Cellular diversity in the Drosophila midbrain revealed by single-cell transcriptomics. *eLife* **7**, e34550.

Davie, K., Janssens, J., Koldere, D., De Waegeneer, M., Pech, U., Kreft, Ł., Aibar, S., Makhzami, S., Christiaens, V., Bravo González-Blas, C., et al. (2018). A Single-Cell Transcriptome Atlas of the Aging Drosophila Brain. *Cell* **174**, 982-998.e20.

Davis, F. P., Nern, A., Picard, S., Reiser, M. B., Rubin, G. M., Eddy, S. R. and Henry, G. L. (2020). A genetic, genomic, and computational resource for exploring neural circuit function. *eLife* **9**, e50901.

Deng, B., Li, Q., Liu, X., Cao, Y., Li, B., Qian, Y., Xu, R., Mao, R., Zhou, E., Zhang, W., et al. (2019). Chemoconnectomics: Mapping Chemical Transmission in Drosophila. *Neuron* **101**, 876-893.e4.

DeSalvo, M. K., Hindle, S. J., Rusan, Z. M., Orng, S., Eddison, M., Halliwill, K. and Bainton,
R. J. (2014). The Drosophila surface glia transcriptome: evolutionary conserved blood-brain
barrier processes. *Front. Neurosci.* **8**, 346.

Dillon, N. R. and Doe, C. Q. (2024). Castor is a temporal transcription factor that specifies early
born central complex neuron identity. *Dev. Camb. Engl.* dev.204318.

Dillon, N. R., Manning, L., Hirono, K. and Doe, C. Q. (2024). Seven-up acts in neuroblasts to
specify adult central complex neuron identity and initiate neuroblast decommissioning. *Dev.*
Camb. Engl. **151**, dev202504.

Doe, C. Q. (2017). Temporal Patterning in the Drosophila CNS. *Annu Rev Cell Dev Biol* **33**, 219–
240.

Enriquez, J., Venkatasubramanian, L., Baek, M., Peterson, M., Aghayeva, U. and Mann, R. S.
(2015). Specification of individual adult motor neuron morphologies by combinatorial
transcription factor codes. *Neuron* **86**, 955–70.

Fisher, Y. E. (2022). Flexible navigational computations in the Drosophila central complex. *Curr.*
Opin. Neurobiol. **73**, 102514.

Franconville, R., Beron, C. and Jayaraman, V. (2018). Building a functional connectome of the
Drosophila central complex. *eLife* **7**, e37017.

Granger, A. J., Wang, W., Robertson, K., El-Rifai, M., Zanello, A. F., Bistrong, K., Saunders,
A., Chow, B. W., Nuñez, V., Turrero García, M., et al. (2020). Cortical ChAT+ neurons
co-transmit acetylcholine and GABA in a target- and brain-region-specific manner. *eLife* **9**,
e57749.

Granger, A. J., Mao, K., Saulnier, J. L., Hines, M. E. and Sabatini, B. L. (2023). Developmental
regulation of GABAergic gene expression in forebrain cholinergic neurons. *Front. Neural*
Circuits **17**, 1125071.

Gu, Z. (2022). Complex heatmap visualization. *iMeta* **1**, e43.

Hao, Y., Stuart, T., Kowalski, M. H., Choudhary, S., Hoffman, P., Hartman, A., Srivastava, A.,
Molla, G., Madad, S., Fernandez-Granda, C., et al. (2024). Dictionary learning for
integrative, multimodal and scalable single-cell analysis. *Nat. Biotechnol.* **42**, 293–304.

Henry, G. L., Davis, F. P., Picard, S. and Eddy, S. R. (2012). Cell type-specific genomics of
Drosophila neurons. *Nucleic Acids Res.* **40**, 9691–9704.

Hindle, S. J. and Bainton, R. J. (2014). Barrier mechanisms in the Drosophila blood-brain barrier.
Front. Neurosci. **8**, 414.

Hobert, O. (2021). Homeobox genes and the specification of neuronal identity. *Nat. Rev. Neurosci.*
22, 627–636.

₇₇₇ **Holguera, I. and Desplan, C.** (2018). Neuronal specification in space and time. *Science* **362**, 176–
₇₇₈   180.

₇₇₉ **Hulse, B. K., Haberkern, H., Franconville, R., Turner-Evans, D. B., Takemura, S., Wolff, T.,**
₇₈₀   **Noorman, M., Dreher, M., Dan, C., Parekh, R., et al.** (2020). A connectome of the
₇₈₁   Drosophila central complex reveals network motifs suitable for flexible navigation and
₇₈₂   context-dependent action selection. *bioRxiv* 2020.12.08.413955.

₇₈₃ **Hulse, B. K., Haberkern, H., Franconville, R., Turner-Evans, D., Takemura, S., Wolff, T.,**
₇₈₄   **Noorman, M., Dreher, M., Dan, C., Parekh, R., et al.** (2021). A connectome of the
₇₈₅   Drosophila central complex reveals network motifs suitable for flexible navigation and
₇₈₆   context-dependent action selection. *eLife* **10**, e66039.

₇₈₇ **Ito, M., Masuda, N., Shinomiya, K., Endo, K. and Ito, K.** (2013). Systematic analysis of neural
₇₈₈   projections reveals clonal composition of the Drosophila brain. *Curr Biol* **23**, 644–55.

₇₈₉ **Kandimalla, P., Omoto, J. J., Hong, E. J. and Hartenstein, V.** (2023). Lineages to circuits: the
₇₉₀   developmental and evolutionary architecture of information channels into the central
₇₉₁   complex. *J. Comp. Physiol. A Neuroethol. Sens. Neural. Behav. Physiol.* **209**, 679–720.

₇₉₂ **Konstantinides, N., Kapuralin, K., Fadil, C., Barboza, L., Satija, R. and Desplan, C.** (2018).
₇₉₃   Phenotypic Convergence: Distinct Transcription Factors Regulate Common Terminal
₇₉₄   Features. *Cell* **174**, 622-635.e13.

₇₉₅ **Konstantinides, N., Holguera, I., Rossi, A. M., Escobar, A., Dudragne, L., Chen, Y.-C., Tran,**
₇₉₆   **T. N., Martínez Jaimes, A. M., Özel, M. N., Simon, F., et al.** (2022). A complete temporal
₇₉₇   transcription factor series in the fly visual system. *Nature* **604**, 316–322.

₇₉₈ **Lago-Baldaia, I., Cooper, M., Seroka, A., Trivedi, C., Powell, G. T., Wilson, S. W., Ackerman,**
₇₉₉   **S. D. and Fernandes, V. M.** (2023). A Drosophila glial cell atlas reveals a mismatch
₈₀₀   between transcriptional and morphological diversity. *PLoS Biol.* **21**, e3002328.

₈₀₁ **Li, H., Janssens, J., De Waegeneer, M., Kolluru, S. S., Davie, K., Gardeux, V., Saelens, W.,**
₈₀₂   **David, F. P. A., Brbić, M., Spanier, K., et al.** (2022). Fly Cell Atlas: A single-nucleus
₈₀₃   transcriptomic atlas of the adult fruit fly. *Science* **375**, eabk2432.

₈₀₄ **Liu, Z., Yang, C. P., Sugino, K., Fu, C. C., Liu, L. Y., Yao, X., Lee, L. P. and Lee, T.** (2015).
₈₀₅   Opposing intrinsic temporal gradients guide neural stem cell production of varied neuronal
₈₀₆   fates. *Science* **350**, 317–20.

₈₀₇ **Lozovaya, N., Eftekhari, S., Cloarec, R., Gouty-Colomer, L. A., Dufour, A., Riffault, B., Billon-**
₈₀₈   **Grand, M., Pons-Bennaceur, A., Oumar, N., Burnashev, N., et al.** (2018). GABAergic
₈₀₉   inhibition in dual-transmission cholinergic and GABAergic striatal interneurons is abolished
₈₁₀   in Parkinson disease. *Nat. Commun.* **9**, 1422.

₈₁₁ **McLaughlin, C. N., Brbić, M., Xie, Q., Li, T., Horns, F., Kolluru, S. S., Kebschull, J. M.,**
₈₁₂   **Vacek, D., Xie, A., Li, J., et al.** (2021). Single-cell transcriptomes of developing and adult
₈₁₃   olfactory receptor neurons in Drosophila. *eLife* **10**, e63856.

McLaughlin, C. N., Qi, Y., Quake R., S., Luo, L. and Li, H. (2022). Isolation and RNA
sequencing of single nuclei from Drosophila tissues: STAR Protocols. *STAR Procotols* **3**,
101417.

Meller, V. H., Wu, K. H., Roman, G., Kuroda, M. I. and Davis, R. L. (1997). roX1 RNA paints
the X chromosome of male Drosophila and is regulated by the dosage compensation system.
Cell **88**, 445–457.

Meye, F. J., Soiza-Reilly, M., Smit, T., Diana, M. A., Schwarz, M. K. and Mameli, M. (2016).
Shifted pallidal co-release of GABA and glutamate in habenula drives cocaine withdrawal
and relapse. *Nat. Neurosci.* **19**, 1019–1024.

Özel, M. N., Gibbs, C. S., Holguera, I., Soliman, M., Bonneau, R. and Desplan, C. (2022).
Coordinated control of neuronal differentiation and wiring by a sustained code of
transcription factors. 2022.05.01.490216.

Pollington, H. Q., Seroka, A. Q. and Doe, C. Q. (2023). From temporal patterning to neuronal
connectivity in Drosophila type I neuroblast lineages. *Semin. Cell Dev. Biol.* **142**, 4–12.

Raji, J. I. and Potter, C. J. (2021). The number of neurons in Drosophila and mosquito brains.
PLOS ONE **16**, e0250381.

Reilly, M. B., Cros, C., Varol, E., Yemini, E. and Hobert, O. (2020). Unique homeobox codes
delineate all the neuron classes of C. elegans. *Nature* **584**, 595–601.

Rosenberg, A. B., Roco, C. M., Muscat, R. A., Kuchina, A., Sample, P., Yao, Z., Graybuck, L.
T., Peeler, D. J., Mukherjee, S., Chen, W., et al. (2018). Single-cell profiling of the
developing mouse brain and spinal cord with split-pool barcoding. *Science* **360**, 176–182.

Sagner, A., Zhang, I., Watson, T., Lazaro, J., Melchionda, M. and Briscoe, J. (2021). A shared
transcriptional code orchestrates temporal patterning of the central nervous system. *PLoS*
Biol. **19**, e3001450.

Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. and Regev, A. (2015). Spatial reconstruction
of single-cell gene expression data. *Nat. Biotechnol.* **33**, 495–502.

Saunders, A., Granger, A. J. and Sabatini, B. L. (2015). Corelease of acetylcholine and GABA
from cholinergic forebrain neurons. *eLife* **4**, e06412.

Schlegel, P., Yin, Y., Bates, A. S., Dorkenwald, S., Eichler, K., Brooks, P., Han, D. S., Gkantia,
M., dos Santos, M., Munnelly, E. J., et al. (2024). Whole-brain annotation and multi-
connectome cell typing of Drosophila. *Nature* **634**, 139–152.

Schoofs, L., De Loof, A. and Van Hiel, M. B. (2017). Neuropeptides as Regulators of Behavior in
Insects. *Annu. Rev. Entomol.* **62**, 35–52.

Sgammeglia, N. and Sprecher, S. G. (2022). Interplay between metabolic energy regulation and
memory pathways in Drosophila. *Trends Neurosci.* **45**, 539–549.

849 **Shabel, S. J., Proulx, C. D., Piriz, J. and Malinow, R.** (2014). Mood regulation. GABA/glutamate
850   co-release controls habenula output and is modified by antidepressant treatment. *Science* **345**,
851   1494–1498.

852 **Sigorelli, A.** (2024). Tools for Descriptive Statistics.

853 **Siletti, K., Hodge, R., Mossi Albiach, A., Lee, K. W., Ding, S.-L., Hu, L., Lönnerberg, P.,**
854   **Bakken, T., Casper, T., Clark, M., et al.** (2023). Transcriptomic diversity of cell types
855   across the adult human brain. *Science* **382**, eadd7046.

856 **Soffers, J. H., Beck, E., Sytkowski, D., Maughan, M., Devasi, D., Zhu, Y., Wilson, B., Chen, Y.-**
857   **C. D., Erclik, T., Truman, J. W., et al.** (2024). A library of lineage-specific driver lines
858   connects developing neuronal circuits to behavior in the Drosophila Ventral Nerve Cord.
859   *BioRxiv Prepr. Serv. Biol.* 2024.11.27.625713.

860 **Spitzer, N. C.** (2015). Neurotransmitter Switching? No Surprise. *Neuron* **86**, 1131–1144.

861 **Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., Hao, Y.,**
862   **Stoeckius, M., Smibert, P. and Satija, R.** (2019). Comprehensive Integration of Single-Cell
863   Data. *Cell* **177**, 1888-1902.e21.

864 **Sullivan, L. F., Warren, T. L. and Doe, C. Q.** (2019). Temporal identity establishes columnar
865   neuron morphology, connectivity, and function in a Drosophila navigation circuit. *eLife* **8**,.

866 **Syed, M. H., Mark, B. and Doe, C. Q.** (2017). Steroid hormone induction of temporal gene
867   expression in Drosophila brain neuroblasts generates neuronal and glial diversity. *eLife* **6**, 6.

868 **Takács, V. T., Cserép, C., Schlingloff, D., Pósfai, B., Szőnyi, A., Sos, K. E., Környei, Z., Dénes,**
869   **Á., Gulyás, A. I., Freund, T. F., et al.** (2018). Co-transmission of acetylcholine and GABA
870   regulates hippocampal states. *Nat. Commun.* **9**, 2848.

871 **Turner-Evans, D. B., Jensen, K. T., Ali, S., Paterson, T., Sheridan, A., Ray, R. P., Wolff, T.,**
872   **Lauritzen, J. S., Rubin, G. M., Bock, D. D., et al.** (2020). The Neuroanatomical
873   Ultrastructure and Function of a Biological Ring Attractor. *Neuron* **108**, 145-163.e10.

874 **Viktorin, G., Riebli, N. and Reichert, H.** (2013). A multipotent transit-amplifying neuroblast
875   lineage in the central brain gives rise to optic lobe glial cells in Drosophila. *Dev Biol* **379**,
876   182–94.

877 **Wang, Y. C., Yang, J. S., Johnston, R., Ren, Q., Lee, Y. J., Luan, H., Brody, T., Odenwald, W.**
878   **F. and Lee, T.** (2014). Drosophila intermediate neural progenitors produce lineage-
879   dependent related series of diverse neurons. *Development* **141**, 253–8.

880 **Warren, T. G., Brennan, M. D. and Mahowald, A. P.** (1979). Two processing steps in maturation
881   of vitellogenin polypeptides in Drosophila melanogaster. *Proc. Natl. Acad. Sci. U. S. A.* **76**,
882   2848–2852.

Wolff, T. and Rubin, G. M. (2018). Neuroarchitecture of the Drosophila central complex: A catalog of nodulus and asymmetrical body neurons and a revision of the protocerebral bridge catalog. *J. Comp. Neurol.* **526**, 2585–2611.

Wolff, T., Iyer, N. A. and Rubin, G. M. (2015). Neuroarchitecture and neuroanatomy of the Drosophila central complex: A GAL4-based dissection of protocerebral bridge neurons and circuits. *J. Comp. Neurol.* **523**, 997–1037.

Wolff, T., Eddison, M., Chen, N., Nern, A., Sundaramurthi, P., Sitaraman, D. and Rubin, G. M. (2024). Cell type-specific driver lines targeting the Drosophila central complex and their use to investigate neuropeptide expression and sleep regulation. *BioRxiv Prepr. Serv. Biol.* 2024.10.21.619448.

Xiong, W. C., Okano, H., Patel, N. H., Blendy, J. A. and Montell, C. (1994). repo encodes a glial-specific homeo domain protein required in the Drosophila nervous system. *Genes Dev.* **8**, 981–94.

Yang, J. S., Awasaki, T., Yu, H. H., He, Y., Ding, P., Kao, J. C. and Lee, T. (2013). Diverse neuronal lineages make stereotyped contributions to the Drosophila locomotor control center, the central complex. *J Comp Neurol* **521**, 2645–62.

Yu, H. H., Kao, C. F., He, Y., Ding, P., Kao, J. C. and Lee, T. (2010). A complete developmental sequence of a Drosophila neuronal lineage as revealed by twin-spot MARCM. *PLOS Biol.* **8**,.

Yu, H. H., Awasaki, T., Schroeder, M. D., Long, F., Yang, J. S., He, Y., Ding, P., Kao, J. C., Wu, G. Y., Peng, H., et al. (2013). Clonal development and organization of the adult Drosophila central brain. *Curr Biol* **23**, 633–43.

T2 neuron atlas

A

50,148 neuronal nuclei, 161 clusters

UMAP2
UMAP1

Top cluster-defining genes

B

Genes

Clusters

Scaled average expression

Figure. T2 glia sex differences and T2 neuron sex differences panels (A–H).

A Bulk seq vs. snRNAseq

B split-gal4 lines

C

		Known markers		Enhancer genes			Newly identified markers			
		toy	runt	rho	shakB	Pkc53E	Gy30A	Tdc2	cut	Lim1	Rx

H

	Lim1+	Rx+	Cut+
P-FN (R16D01)	97 ± 2 %	84 ± 7 %	0 ± 0 %
P-EN (R12D09)	98 ± 3 %	87 ± 18 %	100 ± 0 %