Newly Sequenced Genomes Reveal Patterns of Gene Family Expansion in Select Dragonflies (Odonata: Anisoptera)

Ethan R. Tolman[*,1,2,3], Christopher D. Beatty[1,4], Paul B. Frandsen[5,6,7,] Jonas Bush[8], Or R. Bruchim[9], Ella Simone Driever[9], Kathleen M. Harding[10], Dick Jordan[3], Manpreet K. Kohli[1,3,11], Jiwoo Park[9], Seojun Park[9], Kelly Reyes[12,13], Mira Rosario[9], Jisong L. Ryu[9], Vincent Wade[13,14], Anton Suvorov[2], Jessica L. Ware[1]

1. American Museum of Natural History, Department of Invertebrate Zoology, New York, New York
2. Virginia Tech, Department of Biological Sciences, Blacksburg, Virginia
3. Conservation Connection Foundation, Boise, Idaho, USA
4. Program for Conservation Genomics, Department of Biology, Stanford University, Stanford, California, USA
5. Department of Plant and Wildlife Sciences, Brigham Young University, Provo, Utah, USA
6. LOEWE Centre for Translational Biodiversity Genomics (LOEWE-TBG), Frankfurt, Germany
7. Data Science Lab, Office of the Chief Information Officer, Smithsonian Institution, Washington, District of Columbia, USA
8. Huck Institute of Life Sciences, The Pennsylvania State University, University Park, Pennsylvania, USA
9. Timberline High School, Boise, Idaho, USA
10. Redwood City, California, USA
11. Baruch College, Department of Natural Sciences, New York, New York, USA
12. The Young Women's Leadership School, Manhattan, New York, USA
13. American Museum of Natural History, Science Research Mentorship Consortium, New York, New York, USA
14. Manhattan Village Academy, Manhattan, NY, USA

*Corresponding author: etolman@amnh.org; 2085139703; 829 Canyon Ridge Rd #303, Blacksburg, VA 24060

Abstract

Gene family expansions play a key role in shaping patterns of biodiversity across the tree of life. In Insecta, expanded gene families have broadly been tied to diet, pesticide resistance, immune response and survival in extreme environments. Patterns of gene family evolution are of particular interest in Odonata (dragonflies and damselflies), which are the first lineage to fly, and are one of the most exceptional predators. In Odonata, expansions of opsin genes are correlated with the diversification of the herbivorous insects that Odonata prey upon, but general trends in gene family expansion and contraction have not been studied in this order. Here, we show that two families of Anisoptera (dragonflies), Libellulidae and Petaluridae, have expanded gene families related to their unique life history and diversification patterns. Libellulidae share expanded gene families related to immune response, desiccation response, and processing free radicals, which are all related to the success of Libellulidae in low quality water bodies. The Petaluridae have expanded gene families involved in sensory perception, which can be tied to the unique semi-terrestrial lifestyle of the nymphs of this family. Furthermore, Odonata as a whole has a remarkably high (λ=.19) level of gene turnover compared to other insect orders, a phenomenon that will merit further exploration. These results are an important step towards understanding why Libellulidae is, generally, a species rich family of short lived species that are highly tolerant to poor water quality, while Petaluridae is a species poor family of habitat and behavioral specialists.

Introduction

In dragonflies, why did some families diversify into many species while others remained species poor? The most diverse family in Anisoptera, Libellulidae, likely originated in the late Paleogene (~20-40Ma)(Suvorov et al., 2021) and has since radiated into over 1500 known species. Libellulidae inhabit a wide variety of aquatic habitats, and some taxa are very tolerant of poor water quality (Šigutová et al., 2022). Although generation times have not been studied for most of the Libellulidae, they are generally understood to have, on average, a relatively rapid life-cycle compared to other families of Odonata (Tennessen, 2019)

In contrast to Libellulidae, the least diverse family within Anisoptera is Petaluridae, which originated in the mid-to-late Jurassic (~150-175Ma)(Ware et al., 2014a; Kohli et al., 2021; Suvorov et al., 2021; Tolman et al., 2024), and contains only 11 extant species, with little fossil evidence of widespread extinction (Ware et al., 2014a; Tolman et al., 2024). The ages of the individual species lineages within Petaluridae range between 15 and 115 million years (Ware et al., 2014a; Tolman et al., 2024). While Libellulidae inhabit a broad range of habitats, Petaluridae only inhabit fens with high water quality (Turner, 1970; Baird, 2012; Ware et al., 2014a; Baird, 2019). The lifespan of Petaluridae varies from 5 to 20 years depending on the species, quite possibly making them the longest lived dragonflies (Baird, 2012).

To date, no effort has been undertaken to associate genomic traits with the differing evolutionary and ecological histories of Libellulidae and Petaluridae. With the exception of micro-chromosomes, the genomes of these two families differ very little on a structural scale, as the genomes of the petalurid *Tanypteryx hageni* (Selys, 1879) and libellulid *Pantala flavescens* (Fabricius, 1798) share widespread synteny with both each other, and two distantly related species of Zygoptera (Tolman, Beatty, et al., 2023). Previous research has shown that gene family evolution can be linked to ecological and evolutionary diversification (Luo et al., 2018; Breeschoten et al., 2022; Hu et al., 2023; Kuang et al., 2023; Kwak et al., 2023). Identifying these gene family expansions can and should lead to further research into the functional genomics of these species, which may be tied to adaptations to their different habitats. With this information we can also understand how these species may respond to shifting habitats during climate change (Kim et al., 2022).

Adaptive gene family evolution in insecta

Major morphological, physiological and behavioral adaptations are thought to have influenced the

remarkable success of insects: flight, herbivory and complete metamorphosis are all suggested to be

dramatic drivers of diversity. However, the genomic adaptations that have driven diversity in this group are

less well-known. A better understanding of adaptive gene family expansion may help explain why insects

have been so successful (Luo et al., 2018); work on a number of insect genomes has already been revealing

in this context. While not meant to be exhaustive, the following are pertinent examples. The expansion of

gene families associated with detoxification can be related to polyphagy in the Lepidoptera family

Noctuidae (Breeschoten et al., 2022), and in species of Thysanoptera (Hu et al., 2023). Detoxifying genes

have also duplicated in the weevil *Rhynchophorus ferrugineus* (Olivier, 1790), as have genes related to

insecticide resistance (Hazzouri et al., 2020). Expansion of gene families related to nutrient metabolism,

development, stress response, reproduction, and immune processes may play a major role in the

adaptability of the fall armyworm, *Spodoptera frugiperda* (Smith, 1797)(Gao et al., 2023). The expansion

of sensory related gene families can be associated with host differences in Hymenoptera (Kuang et al.,

2023). The Antarctic midge *Parochlus stenenii* (Gercke, 1889) has undergone many gene family

expansions tied to processes such as the innate immune system, unfolded protein response, DNA

packaging, protein folding, and unsaturated fatty acid biosynthesis that are likely adaptive to the frigid

environment it inhabits (Kim et al., 2022). The expansion of transcription factors, proteases, odorant

receptors, and horizontally transferred genes involved in nutritional symbioses probably play an important

role in interactions with plants and bacteria in the hemipteran *Bactericera cockerelli* (Šulc, 1909) (Kwak et

al., 2023). Gene family expansion has also been linked to adaptations necessary for life in the ocean in the

family Hermatobatidae (Wang et al., 2023). Duplications of genes within the cytochrome p450 family, and

families of olfactory binding proteins are associated with pollution tolerance and mate finding in the

caddisfly S*tenopsyche tienmushanensis* (Hwang, 1957) (Luo et al., 2018). Opsins are highly diverse, with

numerous expansions and contractions across the insect tree of life, but the evolutionary processes driving

these patterns largely remain unclear (Guignard et al., 2022) . However, in Odonata, opsin duplication and

loss has been correlated with both larval and adult behavior, and have been correlated with the

diversification of flowering plants, and herbivorous insects which Odonata prey upon (Futahashi et al.,

2015; Suvorov et al., 2017).

To better understand the gene family evolution of the dragonflies in Libellulidae and Petaluridae,

and how gene expansion is related to the evolutionary and ecological histories of the two families, we

sequenced and annotated the genomes of the libellulid *Pachydiplax longipennis* (Burmeister, 1839) and the

petalurid *Uropetala carovei* (White, 1846). We then included the annotations in a gene family expansion

analysis of all publicly available genomes of Odonata (two Libellulidae, one Petaluridae, and three

Zygoptera) that were generated with PacBio HiFi data at the time of analysis. Because it is relevant for

understanding gene family expansion, we also briefly discuss the life history of each of the five Anisoptera

used in the analysis below.

Pachydiplax longipennis (Libellulidae)

Perhaps no dragonfly interacts with humans more than the monotypic species *Pachydiplax longennis*. *P.*

longipennis has far more iNaturalist observations (90,304 as of October 2023), than any other Anisoptera

(*Erythemis simplicicollis* is the second most observed species, with 82,504 observations as of October

2023). *P. longipennis* is found throughout the continental United States, Mexico, and several Caribbean

Islands. In recent years, its range has expanded northward with a warming climate (Lis et al., 2020). *P.*

longipennis can be observed in localities with a highly diverse assemblage of Odonata, as well as bodies of

water with few species, suggesting a high tolerance to poor water quality(Tollett et al., 2009). While it has

not been formally studied, gbif occurrences suggest that *P. longipennis* is well adapted to cities, especially

in the deserts of the Western United States, where it is commonly observed in urban ponds, but not in

nearby natural areas with high odonate diversity, such as national parks. Many cohorts of *P. longipennis*

develop over the course of a year, but bi-voltinism has also been observed in this species (Wissinger,

1988).

Pantala flavescens (Libellulidae)

P. flavescens, the wandering glider, is one of two species in *Pantala.* It is among the most well-studied

species of Anisoptera, and is the most widely distributed species (Russell et al., 1998). It makes long

transcontinental and trans-oceanic migrations (Rowe, 2004; Ware et al., 2022), and is frequently found in

warmer regions, particularly in the tropics, but is found in temperate regions in the summer (May, 2013).

Though it is not surprising—given the migratory nature of the species—evidence suggests widespread gene

flow between populations across the globe, and it may be appropriate to refer to many, if not all, of the

migratory sub-populations as panmictic (Ware et al., 2022). An smc++ (Liu and Hansen, 2017) analysis,

using individuals from Guangdong Province, China found that the effective population size of *P. flavescens*

has varied between 1 million and 10 million individuals over the last million years, with declines associated

with the onset of the agricultural and industrial period (H. Liu et al., 2022).

Sympetrum striolatum (Charpentier, 1840; Libellulidae)

S. striolatum is a palearctic dragonfly (John Horne, 2012) that is found in Europe and other areas in the

Mediterranean (Borkenstein and Jödicke, 2022), with bodies of salt-water likely serving as a major barrier

to gene flow within this species (Parkes et al., 2009). A demographic model of *S. striolatum*, generated

from a reference assembly of an individual from Wytham Woods, Oxfordshire, UK (Crowley et al., 2023)

estimated a relative stable effective population size around 1 million individuals throughout most of the

past 10 million years (which may predate the divergence times of this species), with declines to an effective

population size of around 100,000 occurring in the last 100,000 years (Tolman, Bruchim, et al., 2023).

Tanypteryx hageni (Petaluridae)

T. hageni is one of only two species in *Tanypteryx* and is the only Petaluridae to inhabit the fens of Western

North America, ranging from Southern California, USA, to Southern British Columbia, Canada (Turner,

1970). *T. hageni* burrows in its nymphal stage, and takes five years to emerge as an adult (Turner, 1970). A

recent behavioral analysis demonstrated that *T. hageni* nymphs regularly leave their burrows for extended

periods of time during both day and night, navigating a terrestrial environment.

Uropetala carovei (Petaluridae)

U. carovei is one of two species of the genus *Uropetala*, and is found in the lowlands of both the Northern

and Southern islands of New Zealand (Rowe, 1981). Their lifespan ranges from five to six years, depending

upon the length of the growing seasons (L. S. Wolfe, 1952). Like *T. hageni*, *U. carovei* burrows in its

nymphal stage (Green, 1977). It has been suggested that *U. carovei* leaves its burrow to forage (G.

Theischinger and I. Endersby, 2009), but it is unclear if *U. carovei* is as active outside of its burrow as *T.*

hageni.

Materials and Methods

Specimen collection

We sequenced the genome of *U. carovei*, which diverged from the lineage leading to *T. hageni* in the late

Jurassic (Ware et al., 2014a; Tolman et al., 2024), to represent the previously unsampled Gondwanan clade

of Petaluridae. *U. carovei* was collected as a nymph near Aorere goldfields in New Zealand and was

injected with and stored in 100% ethanol prior to DNA extraction and sequencing. To expand the taxon

coverage of the Libellulidae we sequenced the genome of *Pachydiplax longipennis*. An adult specimen was

collected from Parkcenter Park Pond in Boise, Idaho, and was stored in 100% ethanol prior to sequencing.

Sequencing and Genome Size Estimation

High molecular weight DNA was extracted from both the *U. carovei* and *P. longipennis* specimens using

the Qiagen Genomic-tip kit. High molecular weight DNA was sheared to 18 kbp using a Diagenode

Megaruptor and separate libraries were prepared for each specimen using the PacBio HiFi SMRTbell

Express Template Kit 2.0. Each library was sequenced on a single PacBio Revio SMRT cell in CCS mode

at the BYU DNA sequencing center.

Genome Assembly and QC

For both *U. carovei* and *P. longipennis* we used HiFi reads to estimate genome size with Genomescope 2.0

(Ranallo-Benavidez et al., 2020). We first generated read histograms with a histogram length of 21 using

Jellyfish (Marçais and Kingsford, 2011), and ran Genomescope 2.0 with default settings. We generated

both whole genome assemblies with hifiasm v.0.16.0 (Cheng et al., 2021).

We used taxon-annotated GC-coverage plots generated with BlobTools v1.1.1 (Laetsch and

Blaxter, 2017) to screen for contamination. We mapped all HiFi reads against the final assembly for each

species using minimap2 v2.1 (Li, 2018) and sorted the resulting bam files with samtools v1.13 (Danecek et

al., 2021). We then assigned taxonomy with megablast (Shiryev et al., 2007) using the parameters: task

megablast and -e-value 1e-25. We calculated coverage using the blobtools function map2cov, created the

blobtools database for each species using the command blobdb, and generated the blobplot with the

command "blobtools plot". After examining the blobplots we removed one contig from the assembly of *P.*

longipennis which was assigned to the phylum tracheophyta and removed all contigs from the assembly of

U. carovei with coverage above 100x, or below 10x, which did not blast to Arthropoda.

Additional contaminants were automatically removed from the assembly of *U. carovei* as a part of

the NCBI genome upload (supplementary table 1). No additional contaminants were removed from the

assembly of *P. longipennis*.

We generated all contiguity stats for both assemblies with the command line tool assembly-stats

(v0.1.4) (Trizna, 2020). We assessed the completeness of both assemblies with BUSCO (v.4.1.4)(Manni et

al., 2021) using the Insecta ODB10 database, in genome mode with the flag –long to retrain the initial

BUSCO results for a more accurate identification of BUSCO genes.

Whole genome alignment

Given the contiguity of the assembly of *P. longipennis*, we calculated synteny between the draft genome

assembly and the chromosome length genome assembly of *S. striolatum* to identify any complete

chromosomes. We identified potentially homologous genes using BLASTp (Camacho et al., 2009),

restricting output to 5 hits per gene and filtering for hits with an e-value below 1e-5. We combined the

BLAST output and gff annotation files for *P. longipennis* and *S. striolatum* and used MCscanX (Wang et

al., 2012) to identify areas of synteny between the genomes with a minimum of 5 genes. We visualized the

syntenic relationships in a Circos (v.0.69-18)(Krzywinski et al., 2009) plot to identify chromosome-length

contigs.

To assemble the remaining contigs into chromosomes, we aligned the chromosome length

assemblies of *Pantala flavescens*, *Sympetrum striolatum*, *Tanypteryx hageni* and the Platycnemididae

Platycnemis pennipes with the assembly of *P. longipennis* using progressive cactus v.2.6.7(Armstrong et

al., 2020). We then used ragout v.2.3(Kolmogorov et al., 2014; Kolmogorov et al., 2018), with the option --

solid-scaffolds, to generate a scaffolded assembly from the alignment. As the karyotype of *U. carovei* is

currently unclear (Kuznetsova and Golub, 2020), it would not be possible to assess the quality of the

scaffolding, so we did not attempt to scaffold this genome. This would not impact the results of the gene

family expansion analysis.

Gene family expansion analysis

Data selection

In addition to the genomes of *P. longipennis* and *U. carovei* we included publicly available genomes of

Odonata from GenBank (October 1, 2023). To minimize bias due to differing genome sequencing methods

we only included genomes that had been generated from PacBio high-fidelity technology, which has been

shown to generate the highest quality insect genomes to date (Hotaling et al., 2021). We identified four

Anisopteran genomes sequenced with this technology, comprising two Libellulidae (*Pantala flavescens* (H.

Liu et al., 2022) and *Sympetrum striolatum* (Crowley et al., 2023)) and one Petaluridae (*Tanypteryx hageni*

(Tolman et al., 2023)). We also chose to include the three publicly available genome assemblies of

Zygoptera, the Platycnemididae *Platycnemis pennipes* (Benjamin W. Price and E. Louise Allan, 2023), the

Coenagrionidae *Ischnura elegans* (Price et al., 2022), and the Calopterygidae *Hetaerina americana*

(Grether et al., 2023) to use as an outgroup in the gene family expansion analysis.

Annotation

To ensure consistency in the genome annotations we annotated with genomes of *H. americana*, *P.

pennipes, I. elegans, P. longipennis, S. striolatum, U. carovei,* and *T. hageni* in GALBA(Brůna et al., 2023)

using the annotated protein set of *P. flavescens* (H. Liu et al., 2022). GALBA aligns the reference protein

set to the soft-masked target genome using miniport (Li, 2023) and uses the alignments to train Augustus

(Stanke et al., 2006; Hoff and Stanke, 2019), and has produced more accurate annotations of insect

genomes than common annotation pipelines such as BRAKER2 (Brůna et al., 2021; Brůna et al., 2023)

which use RNA-seq. Of the published genomes of Odonata, the annotated protein set of *P. flavescens*(H.

Liu et al., 2022) is by far the most complete (Tolman et al., 2023), with nearly 99% of BUSCO genes

present (fig 1.A), justifying its use as a reference. Prior to annotation with GALBA, repetitive elements of

all target species were modeled and soft-masked with RepeatModeler2 (v2.0.1) and RepeatMasker (v.4.1.2-

p1)(Flynn et al., 2020). Because the initial GALBA annotations found an unreasonably high (>50,000)

number of genes, we only kept annotated genes with a significant BLAST(Camacho et al., 2009) hit (with

an e-value cutoff of 1e-10) to the annotated protein set of *P. flavescens* (H. Liu et al., 2022). Given the

completeness of this annotated protein set (H. Liu et al., 2022) this filtering step increases the specificity of

GALBA without a drop in sensitivity, and has since been incorporated into the GALBA pipeline (see

https://github.com/Gaius-Augustus/GALBA/releases/tag/v1.0.10). The new filtered annotations are much

more complete than the original genome annotations for the species (Tolman et al., 2023). We generated all

contiguity stats for both assemblies with the command line tool assembly-stats (v0.1.4)(Trizna, 2020). We

assessed the completeness of all annotations with BUSCO (v4.1.4)(Manni et al., 2021) using the Insecta

ODB10 database, in protein mode.

As part of the annotation process, we also identified the mitochondrial genomes from the de novo

assembles of *P. longipennis* and *U. carovei* using mitohifi (v. 3.0.0)(Uliano-Silva et al., 2021). We used the

mitochondrial genome of *T. hageni* (Tolman et al., 2023)] as a reference for *U. carovei* and the

mitochondrial genome of *S. striolatum* (Crowley et al., 2023) as a reference for *P. longipennis*. The

mitochondrial genome sequences have been uploaded, as part of the reference genome sequences, to

GenBank.

Gene family expansion

While the phylogenetic relationships between the families of Odonata are well established(Bybee et al.,

2021; Kohli et al., 2021; Suvorov et al., 2021), the relationships between the three Libelullidae (*P.*

longipennis, *S. striolatum,* and *P. flavescens*) have not previously been studied with genomic data. To

resolve this relationship, we generated a species tree using the amino acid sequences of 1102 BUSCO

Insecta odb10 genes present in all eight species. We calculated a maximum likelihood tree for each gene

with iqtree2(Minh et al., 2020) using the option "-m test" for model selection. We then generated a species

tree using the estimated gene trees with weighted ASTRAL (Zhang and Mirarab, 2022).

We clustered the amino sequences from the longest isoforms of each gene for each species with

Orthofinder(Emms and Kelly, 2019). We then used CAFE5 (Mendes et al., 2021) to generate four different

lambda (gene birth/death rate) models. First, a global lambda for the whole tree, second, eparate lambdas

for Anisoptera and Zygoptera, third separate lambdas for each family, and fourth separate lambdas for each

species.

As models generated with CAFE5 are not likely to converge when using a large number of

parameters(CAFE, 2023), we only tested the base and error models. For each lambda model we ran the

base and error models ten times to check for convergence and used model one, the only model which

converged, for downstream analysis. We then identified significantly expanding and contracting gene

families from the converged model.

Functional annotation

We obtained GO terms for all of the protein sequences (with the exception of *P. flavescens* which

were previously reported (H. Liu et al., 2022) using interproscan (Mariani et al., 2013; Baek et al., 2021;

Hiranuma et al., 2021). We summarized the GO terms of significantly expanding gene families into

TreeMaps with REVIGO (Supek et al., 2011) for expanded gene families in the Libellulidae and

Petaluridae. Given the missing taxon sampling in other areas of the tree, it would not be appropriate to

analyze patterns of gene family expansion on a broader level.

Demographic histories

As the demographic history of a species may be important when studying genome evolution, particularly in

relic lineages (H.-L. Liu et al., 2022; Bentley et al., 2023), we estimated the demographic histories of *U.*

carovei, *P. longipennis* and *T. hageni* (estimates for all other species have previously been described (H.

Liu et al., 2022; Tolman, Bruchim, et al., 2023)). As with the blobtools analysis, HiFi reads were mapped

to the genome assemblies with minimap2 v2.1.1(Li, 2018), and the bam files were sorted with samtools

v1.16.1 (Danecek et al., 2021) and bcftools v1.6 (Danecek et al., 2021), with a minimum depth of 5 and a

maximum depth of 100, were used to make base calls from the sorted bam files. We then used psmc v0.6.5-

r67(Liu and Hansen, 2017) to convert the fastq basecalling output to psmcfa format and estimate the

pairwise sequential markovian coalescent (PSMC) with 100 bootstrap replicates for each species. For all

three species we visualized the PSMC with the programs psmc(Liu and Hansen, 2017) and gnuplot

v5.2(Phillips, 2012) using four different mutation rates (1e-9, 2e-9, 3e-9, and 4e-9), encompassing known

genome-wide mutation rates in Insecta (Liu et al., 2017), and values previously used in demographic

analyses of Odonata (H. Liu et al., 2022; Ethan R. Tolman et al., 2023). As *P. longipennis* cohorts are

typically uni-voltine, but occasionally exhibit bi-voltinism (Wissinger, 1988) , we used a generation time of

0.8 to reflect the occasional bi-voltine generations. *U. carovei* has a generation time of 5-6 years depending

on the length of growth seasons (Winstanley and Rowe, 1980; Baird, 2012) so we used a generation time of

5.5 years. *T. hageni* has a life cycle of 5 years (Turner, 1970; Baird, 2012), and this was the value used in

the coalescent analysis.

Results

Sequencing and Genome Size Estimation

We generated over 6 million high-fidelity reads, totaling more than 93 Gbp, for *U. carovei* (read

N50=15,101; L50=2,488,241; ave=14,547.95) and over 5.8 million reads totaling more than 76 Gbp for *P.*

longipennis (read N50 = 13,394; L50 = 2,369,807; ave= 13,034.64). Using GenomeScope 2.0, we

estimated a genome size of 0.88 Gb (87.4% unique sequence, 1.8% heterozygosity, kcov: 40.5, err:

0.0885%, k:21) for *P. longipennis* (supplementary fig. 1A). and 1.00 Gb (67.5% unique sequence, 1.18%

heterozygosity, kcov: 36.7, err: 0.224%, k:21) for *U. carovei* (supplementary fig. 1B).

Genome Assembly and QC

The initial assembly of *U. carovei* was highly contiguous when compared to other whole genome

assemblies of Odonata (length=1.33Gb, N50=8.67Mb, L50=49, ave=1.72Mb, N=773). The assembly of *P.*

longipennis was among the most contiguous genomes ever assembled without scaffolding (length=1.0Gb,

N50=66.9 1Mb, L50=7, ave=3.71Mb, N=270). The blobplot, which displays the assigned taxonomy,

coverage and GC content of each contig to identify potential contaminants, of *U. carovei* showed potential

contamination (supplementary fig. 2A) of contigs with coverage above 100x, or below 10x. Such contigs

that were not assigned to Arthropoda were removed. Only one contig of *P. longipennis*, which was

assigned by BLAST to tracheophyta (a common laboratory plant) was removed from the assembly

(supplementary fig. 2B), as all other contigs were assigned to Arthropoda. The identification of core

BUSCO orthologues indicate that the genomes were highly complete, with nearly 97 percent of assessed

Insecta genes present in the assembly of *U. carovei* (C:96.9%[S:95.9%,D:1.0%],F:1.0%,M:2.1%,n:1367),

and 98% of the Insecta genes present in *P. longipennis* (C:98.0% [S:95.2%,D:2.8%], F:0.5%, M:1.5%,

n:1367). Both are comparable to other HiFi assemblies of Odonata(Newton et al., 2023 Jun 7; Tolman et

al., 2023).

Whole genome alignment

We generated eight pseudo-chromosomes for the assembly of *P. longipennis* from the cactus whole

genome alignment with *S. striolatum* (supplementary table 2). The remaining four chromosomes still

showed high levels of synteny with a select number of contigs (supplementary table 2, supplementary fig.

3).

Annotation

All our annotations recovered more than 97% of BUSCO Insecta db10 genes (fig. 1A) after filtering. We

initially recovered 114,332 genes for *H. americana*, 103,410 for *P. pennipes*, 66,745 for *I. elegans*, 45,043

for *P. longipennis*, 53,217 for *S. striolatum*, 53,629 for *U. carovei*, and 75,637 for *T. hageni*. After filtering

for genes that had a significant BLAST hit with the proteome of *P. flavescens* we recovered 13,567 genes

for *H. americana*, 15,064 from *P. pennipes*, 15,378 from *I. elegans*, 16,663 from *P. longipennis*, 17,672

from *S. striolatum*, 15,122 from *U. carovei*, and 17,447 from *T. hageni* (fig. 1B).

Figure 1: Gene Family Expansion Analysis

Gene family evolution

Phylogeny

All expected relationships were recovered with posterior probability support of 1 (fig. 1C). *P. flavescens*

was recovered as the outgroup within Libellulidae, with posterior probability support of 1.

Gene birth-death rate

All ten estimates of the global lambda converged at 0.19 (fig. 1C), but no other models converged.

Expansions and contractions

The suborders had differing numbers of significantly expanding (Anisoptera=24, Zygoptera=37) but the

same number of contracting (66) gene families (fig. 1C). The Petaluridae had 73 significantly expanding,

and 282 significantly contracting gene families, compared to 106 significantly expanding and 300

significantly contracting families in the Libellulidae (fig. 1C). Of the species included, *S. striolatum* had the

largest number of significant gene family expansions (281) and *P. flavescens* had the fewest (77). *H.*

americana had the highest number of significantly contracting gene families (627), and *U. carovei* had the

fewest significantly contracting families (87).

Functional annotation

Figure 2: Tree maps of GO terms associated with significantly expanding gene families in Libellulidae and

Petaluridae

Gene Ontology (GO) terms associated with significantly expanding gene families shared amongst the

Libellulidae and Petaluridae were summarized into 26 (Fig. 2A) and 25 (Fig 2B) biological processes

respectively. Within Libellulidae, GO terms associated with gene family expansion were summarized into

37 biological processes in *S. striolatum*, 17 in *P. flavescens*, and 29 processes in *P. longipennis*. In

Petaluridae, GO terms from significantly expanded gene families were summarized into 26 biological

processes in *T. hageni*, and 31 processes in *U. carovei*.

Figure 3: *Demographic histories* of *Pachydiplax longipennis, Uropetala carovei, and Tanypteryx hageni*

To test the assumption that the Petaluridae have long been habitat specialists, and the Libellulidae have

long been generalists, we modeled the historical effective population size of the species considered here, as

habitat specialists are more likely to have a historically smaller effective population size. PSMC models

showed one climb in N_e of *P. longipennis*, starting between 7 and 20 Mya, and peaking from 1-6 Mya (fig.

3A). The peak N_e was estimated to be between 0.5 and 6 million, with declines stabilizing less than 100k

years before present at ~100,000 (supplementary fig. 3A). The PSMC model suggests the N_e of *Uropetala*

carovei fluctuated throughout a period from 25-100 Mya depending on the mutation rate (fig. 3B). In the

scenario with the lowest genome-wide mutation rate, from 100 Mya to 10 Mya, the N_e of *Uropetala*

increases, but continuously decreases until around a million years ago. In the scenario with the fastest

mutation rate, the population starts increasing around 25 Mya and peaks around 2.5 Mya (fig. 3B). Among

the different scenarios the highest peak was between 7 and 25 million individuals (fig. 3B).

All estimates for the change in effective population size (N_e) for *T. hageni* showed an initial drop in

population size starting around 30 million years ago (Mya), until a brief uptick in population size

approximately 3 Mya, followed by another period of decline stabilizing approximately 200,000 years

before present (fig. 3C), as made evident by the tight bootstrap replicates. Regardless of the mutation rate

used in the model, the N_e of *T. hageni* never climbed higher than 2 million individuals, and lower estimates

of the genome-wide mutation rate do not show an effective population size above 1 million (fig. 3C).

Discussion

Our results provide a number of testable hypotheses relating to genome size, gene family expansion, and

the diversification of odonates into the range of ecosystems they currently inhabit. We consider here the

strengths and implications of the technologies used to generate these results, and then some of the major

research areas suggested by our results.

Genome size estimation and quality control

Genome size estimates of Odonata assembled from PacBio high-fidelity sequences have been found to

diverge from genome sizes estimated with Genomescope(Ranallo-Benavidez et al., 2020). Although many

contigs from the *U. carovei* assembly were not assigned to the Arthropoda, it is likely an artifact of an

incomplete NCBI reference database (Tolman et al., 2023). Another Petaluridae, *T. hageni*, showed a

similar profile with contigs not assigned to Arthropoda, even in well supported scaffolds with HiC data,

likely due to poor coverage in NCBI databases (Tolman et al., 2023). This phenomenon has also been

observed in horseshoe crabs, and in the transcriptomes of other Petaluridae (Tolman et al., 2023). With our

current understanding, taxonomically mis-assigned contigs with a similar GC content and coverage to

contigs not assigned to Arthropoda are not a major source of concern (Tolman et al., 2023).

Implications of PacBio Revio Technology in Insect genomics

PacBio high-fidelity sequencing has become the gold standard for insect genomics (Hotaling et al., 2021).

The near chromosome-level assembly of *P. longipennis* presented here, before any scaffolding, further

demonstrates the potential of this sequencing technology. As this technology is further developed,

complete, chromosome-length genome assemblies will be available without any scaffolding.

GALBA as a tool for genome annotation

The genome annotation tool GALBA has been demonstrated to outperform more established pipelines,

such as BRAKER2 (Brůna et al., 2021) when annotating insect genomes, and does not need further

sequencing of transcriptomes (Brůna et al., 2023). While our genome annotations initially had a high

(>50,000) number of genes compared to the annotation of *P. flavescens*, by conservatively retaining only

genes which had a BLAST hit to a protein sequence from *P. flavescent s*(H. Liu et al., 2022), we were able

to generate highly complete annotations with nearly all insecta BUSCO orthologues present (fig 1. A, B),

further supporting the efficacy of this pipeline in insects, especially in the absence of transcriptomic data.

Gene birth/death rate

The global lambda across Odonata (λ=.19) was unusually high compared to other branches of the tree of

life. Estimated lambdas of primates (λ<0.01), birds of paradise (λ<0.01), and the insect orders

Hymenoptera (λ<0.01) (Mendes et al., 2021), and Hemiptera (λ<0.01) (Kwak et al., 2023) are all less than

5% of the global lambda of Odonata (λ=0.19). This is not solely due to the age of Odonata (Anisoptera and

Zygoptera split some 250Mya (Kohli et al., 2021; Suvorov et al., 2021)), as the tree used to study gene

family expansion in Hemiptera was estimated to have a root divergence time older than 500 Mya (Kwak et

al., 2023). An analysis of the synteny between chromosome level genome assemblies of Odonata found a

high degree of conservation in chromosome structure, even between suborders (Tolman, Beatty, et al.,

2023). In *P. flavescens* and *T. hageni* the relative proportion of CDS content was not retained in a linear

model predicting the proportion of synteny in a given partition of the genome assembly, suggesting

divergence in gene sequences, even though the chromosome structure was largely conserved (Tolman,

Beatty, et al., 2023). The high gene turnover rates observed in Odonata could explain this dissociation

between CDS and synteny. Our estimate, made with currently available data, provides evidence that high

levels of ecological diversification and speciation in Libellulidae, and the habitat specialization and low

levels branching of the Petaluridae, could both be supported in systems with high levels of gene turnover.

While more research is necessary to better understand the molecular processes tied to the

evolutionary and ecological histories of odonates, our functional annotations of expanding gene families

can be used to generate testable hypotheses related to the ecology and evolution of Petaluridae and

Libellulidae. Here, we discuss hypotheses derived from our GO analyses that warrant further investigation

in future studies.

As *T. hageni* and *U. carovei* represent ~18% of all Petaluridae, and both major clades within the

family (Ware et al., 2014, Tolman et al., 2024), we believe it is highly likely that gene family expansions

shared between these two species are shared across Petaluridae. However, as the family Gomphidae is

likely sister to Petaluridae (Bybee et al., 2021; Kohli et al., 2021; Suvorov et al., 2021), without

representative genomes for Gomphidae, we cannot determine whether these expanded gene families would

be found in both families, thereby suggesting that they may have originated in a most-recent common

ancestor. Given these limitations, the expanding gene families shared between the two Petaluridae still

suggest candidate genes to explore for genomic adaptations related to the life history attributes of this

family, namely the longevity of these lineages through evolutionary time, the long development times seen

in these species, and the unique semi-terrestrial lifestyle of the nymphs.

Hypothesis 1: older lineages have expanded immune response families. One of the expanded

gene families seen in the petalurids is associated with *immune response* (Fig. 2B). A possibility is that

these are associated with organisms descended from older lineages; another long-lived arthropod lineage,

the mangrove horseshoe crab, also has an expansion of genes related to immune response (Shingate et al.,

2020). The two petalurid species analyzed here separated from their closest relatives approximately 25 and

5 Mya (Tolman et al., 2024). To explore this correlation, data on more "ancient" arthropod lineages are

needed.

Hypothesis 2: terrestrial nymphs have different sensory perception profiles. Across the

family Petaluridae, and in both representative species, we detected an expansion in genes that play a part in

sensory perception of taste (fig. 2B). While all odonate nymphs are aquatic, the nymphs of these petaltail

species have been observed to forage in the terrestrial habitats around their subterranean burrows (which

fill with water, providing an aquatic environment). Odonates are generalist predators in their nymph and

adult stages. Terrestrial foraging could bring these petalurid nymphs in contact with a range of potential

prey very different from those they would encounter in a strictly aquatic environment. Insects generally

utilize taste to avoid toxic chemicals, find suitable food sources, select mates and find oviposition sites

(Rebora et al., 2022). Adult Odonata, in particular, have been observed catching and rejecting toxic prey

upon tasting it (Rebora et al., 2022). It is currently unclear if nymphs exhibit this behavior. With terrestrial

foraging, petalurid nymphs may encounter a greater proportion of toxic prey, such as arachnids (which

have been observed in and around the burrows of *T. hageni*. Further examination of the specific genes

involved in taste perception in petalurids needs to be performed, along with greater documentation of their

foraging behavior and analysis of their diet to test this hypothesis.

Hypothesis 3: terrestrial nymphs have different phototransduction gene families. The

expansion of gene families related to *phototransduction* in *T. hageni* could be due to the differing light

conditions the nymphs face in a terrestrial environment. Transcriptomic data has shown that opsin genes in

particular have undergone a number of expansions and contractions across Odonata, but the reasons for this

evolutionary pattern are currently unclear (Bybee et al., 2022). Transcriptomic work will be needed to

confirm whether expanded gene families related to phototransduction are expressed in the nymphs, as

previously published transcriptomic work including Petaluridae sequenced adults (Suvorov et al., 2017). It

is currently unclear if *U. carovei* is semi-terrestrial to the same extent that *T. hageni*, and it is necessary to

determine the extent of semi-terrestriality across Petaluridae to refine any hypotheses related to this

behavior. As the nymphal behavior of the Petaluridae is better understood, and more genomes and

transcriptomes are sequenced, gene families associated with taste and vision could be a promising avenue

for understanding their interaction with a terrestrial environment as nymphs.

Hypothesis 4: highly tolerant dragonflies have expanded gene families to deal with

desiccation and bacterial exposure. While it is certainly plausible that shared expansions in the three

Libellulidae analyzed here are shared across the family, we will be more certain about gene family

expansions among libellulids when more genomes are available. Several biological processes related to the

shared expansions of these three taxa may be associated with the generalist life-history of libellulids. A

major overrepresented GO term shared between these three species is the *peptidoglycan catabolic process*

(fig. 2A). Peptidoglycan is an essential element in the walls of most bacteria (Vollmer et al., 2008). Several

hypotheses could explain why gene families related to catabolizing peptidoglycan would be present. First:

genes involved in the peptidoglycan catabolic process were upregulated when the Antarctic midge *Belgica*

antarctica was placed under desiccation conditions (Teets et al., 2012). In the case of odonates, expansions

could be adaptive to deal with more arid environments, or in ephemeral aquatic environments. Secondly,

Libellulidae often inhabit warm, eutrophic waters, where harmful bacteria are more likely to be found

(Sharip and Fauzi, 2019). Expansion of gene families involved in the peptidoglycan catabolic process could

enable Libellulidae to tolerate exposure to high levels of bacteria.

Hypothesis 5: highly tolerant dragonflies have expanded gene families to deal with exposure

to free radicals. Another GO term associated with expanded gene families in Libellulidae is *cell redox*

homeostasis (fig. 2A), a dynamic but essential process ensuring balance between oxidizing and reducing

reactions in cells that is also involved in many biological responses (Le Gal et al., 2021). These elements

have been shown to maintain homeostasis in fluid conditions(Le Gal et al., 2021). An ability to maintain

homeostasis would be critical for Libellulidae, as the shallow bodies of water they inhabit can often

undergo rapid changes in temperature and water chemistry. GO terms associated with expanding gene

families of *P. longipennis* could offer some explanation for how this species has been so successful in

colonizing urban environments. One of the most overrepresented GO terms for this species was *response to*

oxidative stress (fig. 2A). Birds in urban or polluted environments have increased antioxidant protection,

and face higher levels of oxidative stress (Salmón et al., 2018), so it is likely that Odonata face these same

pressures. It follows that a greater capacity to respond to oxidative stress in urban environments would

allow *P. longipennis* to outcompete other species that do not have this capacity. Additionally, the expansion

of gene families related to *chromatin remodeling* and *methylation* (fig. 2A) could allow *P. longipennis*

sufficient plasticity to thrive in an urban environment.

Hypothesis 6: highly tolerant dragonflies have expanded gene families to allow for greater

phenotypic plasticity. Expansions related to both *methylation*, *response to ATP*, and *response to stress* in

Libellulidae (fig. 2A) could result in increased phenotypic and molecular plasticity, which are a necessity

for evolutionary stability in fluctuating environments (Lalejini et al., 2021), and could be another molecular

adaptation to fluctuating environments.

Hypothesis 7: migratory dragonflies can more efficiently process carbohydrates. In *P.*

flavescens, the expansion of gene families related to the *carbohydrate metabolic process* are a plausible

mechanism for allowing this species to undertake trans-oceanic and trans-continental migrations in a single

generation (Ware et al., 2022). It is likely that an especially efficient process for obtaining energy from

carbohydrates would be selected for on such long migrations. The genome of closely related non-migratory

species to *P. flavescens* are needed to explore the genomic adaptations to a migratory lifestyle.

We recognize that a limitation of this study is a lack of taxonomic coverage. *T. hageni* and *U.*

carovei represent the two major clades of Petaluridae (Ware et al., 2014b) (Laurasian and Gondwanan) and

~18% of the family. However, the three species of Libellulidae anlayzed here represent only 3 of 29 genera

and 3 of >1500 species. Additionally, our analysis includes only 2 of the 39 families of Anisoptera, 3 of

the 39 Zygoptera, and only 8 of ~6300 species of Odonata(May, 2019), so we acknowledge the birth/death

rates calculated could be influenced by poor taxon sampling, and more genomes are needed to more

accurately model gene birth and death. We have included all of the high-quality odonate genome

assemblies currently available at the time of analysis, offering predictions about birth/death rates in this

order that can be refined as more data becomes available.

Demographic histories

Similar to *I. elegans* and *S. striolatum* (Tolman, Bruchim, et al., 2023) the PSMC model for *P. longipennis*

showed a decline in effective population size $(N)_e$ roughly corresponding with the Pleistocene ice ages. As

the climate warms this species could become more widely distributed, but more research into the ecological

niche of this species is needed to affirm this assumption. The N_e of the two fen-dwelling petalurid species–

U. carovei and *T. hageni*–were estimated to be smaller than *P. longipennis* or the other libellulids through

time (H. Liu et al., 2022; Tolman, Bruchim, et al., 2023). If the Petaluridae have consistently been habitat

specialists through the period covered by our analysis, we would predict that their population size would be

limited by availability of their habitat.

Conclusions

Our findings are the first comprehensive analysis of gene family expansion in Odonata. The results suggest

that gene turnover is much higher in Odonata than in other insect orders, and we identify a number of gene

families that may be associated with the differing evolutionary and ecological histories of the Petaluridae

and the Libellulidae. The role of these gene families, and the relatively high levels of gene turnover will be

an exciting avenue of future research as more genomes from across the Odonata, along with more

transcriptome data, become available.

Author contributions

E.R.T: Methodology, formal analysis, data curation, writing.

C.D.B: Methodology, formal analysis, writing.

P.B.F: Methodology, writing, resources.

R.J.B: Methodology, writing.

O.R.B: Formal analysis, writing.

E.S.D: Formal analysis.

K.M.H: Formal analysis.

D.J: Methodology, resources, writing.

M.K.K Methodology.

J.P: Formal analysis.

S.P: Formal analysis.

K.R: Formal analysis, writing.

M.R: Formal analysis.

J.L.R: Formal analysis.

V.W: Formal analysis, writing.

A.S. Resources, supervision, writing.

J.L.W: Methodology, formal analysis, resources, supervision, writing.

Acknowledgments

We would like to thank Kristin Gnojewski and Cindy Busche for their support of this project. We

acknowledge funding from the Conservation Connection Foundation, the Society of Systematic Biologists

Graduate Student Research Award, the 2023 Sydney Anderson Travel Award from the American Museum

of Natural History, and the National Science Foundation (DEB#2002473).

Declaration of interests

The authors declare no competing interests.

Data availability

577   The genome assemblies and rawreads of U. carovei (PRJNA1043152 and P. longipennis (PRJNA999334)

578   are available on GenBank. RepeatMasker output (10.6084/m9.figshare.24714909), unfiltered annotations

579   (10.6084/m9.figshare.24714918) and filtered proteins sets used for gene family expansion analysis

580   (10.6084/m9.figshare) are all available on figshare.

581
References

Armstrong J, Hickey G, Diekhans M, et al. 2020. Progressive Cactus is a multiple-genome aligner for the
thousand-genome era. Nature. 587(7833):246–251. https://doi.org/10.1038/s41586-020-2871-y.

Baek M, DiMaio F, Anishchenko I, et al. 2021. Accurate prediction of protein structures and interactions
using a three-track neural network. Science. 373(6557):871–876. https://doi.org/10.1126/science.abj8754.

Baird I. 2012. The wetland habitats, biogeography and population dynamics of Petalura gigantea (Odonata:
Petaluridae) in the Blue Mountains of New South Wales [PhD]. The University of Western Sydney.

Baird IRC. 2019. Establishment of larval pits by Tachopteryx thoreyi (Odonata: Petaluridae): habitat
modification by a non-burrowing petalurid. International Journal of Odonatology. 22(2):135–146.
https://doi.org/10.1080/13887890.2019.1636889.

Benjamin W. Price, E. Louise Allan. 2023. The genome sequence of the White-legged damselfly,
Platycnemis pennipes (Pallas, 1771). Available from https://wellcomeopenresearch.org/articles/8-320.

Bentley BP, Carrasco-Valenzuela T, Ramos EKS, et al. 2023. Divergent sensory and immune gene
evolution in sea turtles with contrasting demographic and life histories. Proceedings of the National
Academy of Sciences. 120(7):e2201076120. https://doi.org/10.1073/pnas.2201076120.

Borkenstein A, Jödicke R. 2022. Thermoregulatory behaviour of Sympetrum striolatum at low
temperatures with special reference to the role of direct sunlight (Odonata: Libellulidae). 51:83–109.

Breeschoten T, van der Linden CFH, Ros VID, et al. 2022. Expanding the Menu: Are Polyphagy and Gene
Family Expansions Linked across Lepidoptera? Genome Biology and Evolution. 14(1):evab283.
https://doi.org/10.1093/gbe/evab283.

Brůna T, Hoff KJ, Lomsadze A, et al. 2021. BRAKER2: automatic eukaryotic genome annotation with
GeneMark-EP+ and AUGUSTUS supported by a protein database. NAR Genomics and Bioinformatics.
3(1):lqaa108. https://doi.org/10.1093/nargab/lqaa108.

Brůna T, Li H, Guhlin J, et al. 2023. Galba: genome annotation with miniprot and AUGUSTUS. BMC
Bioinformatics. 24(1):327. https://doi.org/10.1186/s12859-023-05449-z.

Bybee SM, Futahashi R, Renoult JP, et al. 2022. Transcriptomic insights into Odonata ecology and
evolution. In: Cordoba-Aguilar A, Beatty C, Bried J, editors. Dragonflies and Damselflies: Model
Organisms for Ecological and Evolutionary Research. Oxford University Press. p. 0.
https://doi.org/10.1093/oso/9780192898623.003.0003.

Bybee SM, Kalkman VJ, Erickson RJ, et al. 2021. Phylogeny and classification of Odonata using targeted
genomics. Molecular Phylogenetics and Evolution. 160:107115.
https://doi.org/10.1016/j.ympev.2021.107115.

CAFE. 2023. Available from https://github.com/hahnlab/CAFE5.

Camacho C, Coulouris G, Avagyan V, et al. 2009. BLAST+: architecture and applications. BMC
Bioinformatics. 10(1):421. https://doi.org/10.1186/1471-2105-10-421.

Cheng H, Concepcion GT, Feng X, et al. 2021. Haplotype-resolved de novo assembly using phased
assembly graphs with hifiasm. Nat Methods. 18(2):170–175. https://doi.org/10.1038/s41592-020-01056-5.

Christopher D. Beatty, Aaron Goodman, Ethan R. Tolman, et al. In Preparation. The Larval Behavior of
Tanypteryx Hageni.

Crowley LM, Price BW, Allan EL, et al. 2023. The genome sequence of the Common Darter, Sympetrum
striolatum (Charpentier, 1840). Wellcome Open Res. 8:389.
https://doi.org/10.12688/wellcomeopenres.19937.1.

Danecek P, Bonfield JK, Liddle J, et al. 2021. Twelve years of SAMtools and BCFtools. Gigascience.
10(2):giab008. https://doi.org/10.1093/gigascience/giab008.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics.
Genome Biology. 20(1):238. https://doi.org/10.1186/s13059-019-1832-y.

Flynn JM, Hubley R, Goubert C, et al. 2020. RepeatModeler2 for automated genomic discovery of
transposable element families. Proceedings of the National Academy of Sciences. 117(17):9451–9457.
https://doi.org/10.1073/pnas.1921046117.

G. Theischinger, I. Endersby. 2009. Identification Guide to the Australian Odonata. Hurtsville, New South
Wales, Australia: Department of Environment, Climate Change and Water NSW.

Gao H, Li Y, Tian Y, et al. 2023. Gene family expansion analysis and identification of the histone family in
Spodoptera frugiperda. Comparative Biochemistry and Physiology Part D: Genomics and Proteomics.
48:101142. https://doi.org/10.1016/j.cbd.2023.101142.

Green LFB. 1977. Aspects of the respiratory and excretory physiology of the nymph of Uropetala carovei
(Odonata: Petaluridae). New Zealand Journal of Zoology. 4(1):39–43.
https://doi.org/10.1080/03014223.1977.9517934.

Grether GF, Beninde J, Beraut E, et al. 2023. Reference genome for the American rubyspot damselfly,
Hetaerina americana. Journal of Heredity. 114(4):385–394. https://doi.org/10.1093/jhered/esad031.

Guignard Q, Allison JD, Slippers B. 2022. The evolution of insect visual opsin genes with specific
consideration of the influence of ocelli and life history traits. BMC Ecology and Evolution. 22(1):2.
https://doi.org/10.1186/s12862-022-01960-8.

Hazzouri KM, Sudalaimuthuasari N, Kundu B, et al. 2020. The genome of pest Rhynchophorus ferrugineus
reveals gene families important at the plant-beetle interface. Commun Biol. 3(1):1–14.
https://doi.org/10.1038/s42003-020-1060-8.

Hiranuma N, Park H, Baek M, et al. 2021. Improved protein structure refinement guided by deep learning
based accuracy estimation. Nat Commun. 12(1):1340. https://doi.org/10.1038/s41467-021-21511-x.

Hoff KJ, Stanke M. 2019. Predicting Genes in Single Genomes with AUGUSTUS. Curr Protoc
Bioinformatics. 65(1):e57. https://doi.org/10.1002/cpbi.57.

Hotaling S, Sproul JS, Heckenhauer J, et al. 2021. Long Reads Are Revolutionizing 20 Years of Insect
Genome Sequencing. Genome Biology and Evolution. 13(8):evab138. https://doi.org/10.1093/gbe/evab138.

Hu Q-L, Ye Z-X, Zhuo J-C, et al. 2023. A chromosome-level genome assembly of Stenchaetothrips
biformis and comparative genomic analysis highlights distinct host adaptations among thrips. Commun
Biol. 6(1):1–10. https://doi.org/10.1038/s42003-023-05187-1.

John Horne. 2012. Emergence, maturation time and oviposition in the Common Darter Sympetrum
striolatum (Charpentier). Journal of the British Dragonfly Society. 28(2). Available from https://british-
dragonflies.org.uk/wp-content/uploads/2020/11/JBDS_Vol28_2.pdf#page=13.

Kim H, Kim H-W, Lee JH, et al. 2022. Gene family expansions in Antarctic winged midge as a strategy for
adaptation to cold environments. Sci Rep. 12(1):18263. https://doi.org/10.1038/s41598-022-23268-9.

Kohli M, Letsch H, Greve C, et al. 2021. Evolutionary history and divergence times of Odonata
(dragonflies and damselflies) revealed through transcriptomics. iScience. 24(11):103324.
https://doi.org/10.1016/j.isci.2021.103324.

Kolmogorov M, Armstrong J, Raney BJ, et al. 2018. Chromosome assembly of large and complex genomes
using multiple references. Genome Res. 28(11):1720–1732. https://doi.org/10.1101/gr.236273.118.

Kolmogorov M, Raney B, Paten B, et al. 2014. Ragout—a reference-assisted assembly tool for bacterial
genomes. Bioinformatics. 30(12):i302–i309. https://doi.org/10.1093/bioinformatics/btu280.

Krzywinski M, Schein J, Birol I, et al. 2009. Circos: an information aesthetic for comparative genomics.
Genome Res. 19(9):1639–1645. https://doi.org/10.1101/gr.092759.109.

Kuang J-G, Han Z-T, Kang M-H, et al. 2023. Chromosome-level de novo genome assembly of two conifer-
parasitic wasps, Megastigmus duclouxiana and Megastigmus sabinae, reveals genomic imprints of
adaptation to hosts. Molecular Ecology Resources. 23(5):1142–1154. https://doi.org/10.1111/1755-
0998.13785.

Kuznetsova VG, Golub NV. 2020. A checklist of chromosome numbers and a review of karyotype
variation in Odonata of the world. Comp Cytogenet. 14(4):501–540.
https://doi.org/10.3897/CompCytogen.v14i4.57062.

Kwak Y, Argandona JA, Degnan PH, et al. 2023. Chromosomal-level assembly of Bactericera cockerelli
reveals rampant gene family expansions impacting genome structure, function and insect-microbe-plant-
interactions. Molecular Ecology Resources. 23(1):233–252. https://doi.org/10.1111/1755-0998.13693.

L. S. Wolfe. 1952. A study of the genus Uropetala Selys (Order Odonata) from New Zealand. Transactions
of the Royal Society of New Zealand. 80:245–275.

Laetsch DR, Blaxter ML. 2017. BlobTools: Interrogation of genome assemblies. F1000Res. 6:1287.
https://doi.org/10.12688/f1000research.12232.1.

Lalejini A, Ferguson AJ, Grant NA, et al. 2021. Adaptive Phenotypic Plasticity Stabilizes Evolution in
Fluctuating Environments. Frontiers in Ecology and Evolution. 9. Available from
https://www.frontiersin.org/articles/10.3389/fevo.2021.715381.

Le Gal K, Schmidt EE, Sayin VI. 2021. Cellular Redox Homeostasis. Antioxidants (Basel). 10(9):1377.
https://doi.org/10.3390/antiox10091377.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 34(18):3094–3100.
https://doi.org/10.1093/bioinformatics/bty191.

Li H. 2023. Protein-to-genome alignment with miniprot. Bioinformatics. 39(1):btad014.
https://doi.org/10.1093/bioinformatics/btad014.

Lis C, Moore MP, Martin RA. 2020. Warm developmental temperatures induce non-adaptive plasticity in
the intrasexually selected colouration of a dragonfly. Ecological Entomology. 45(3):663–670.
https://doi.org/10.1111/een.12839.

Liu H, Jia Y, Sun X, et al. 2017. Direct Determination of the Mutation Rate in the Bumblebee Reveals
Evidence for Weak Recombination-Associated Mutation and an Approximate Rate Constancy in Insects.
Molecular Biology and Evolution. 34(1):119–130. https://doi.org/10.1093/molbev/msw226.

Liu H, Jiang F, Wang S, et al. 2022. Chromosome-level genome of the globe skimmer dragonfly (Pantala
flavescens). GigaScience. 11:giac009. https://doi.org/10.1093/gigascience/giac009.

Liu H-L, Harris AJ, Wang Z-F, et al. 2022. The genome of the Paleogene relic tree Bretschneidera sinensis:
insights into trade-offs in gene family evolution, demographic history, and adaptive SNPs. DNA Research.
29(1):dsac003. https://doi.org/10.1093/dnares/dsac003.

Liu S, Hansen MM. 2017. PSMC (pairwise sequentially Markovian coalescent) analysis of RAD
(restriction site associated DNA) sequencing data. Molecular Ecology Resources. 17(4):631–641.
https://doi.org/10.1111/1755-0998.12606.

Luo S, Tang M, Frandsen PB, et al. 2018. The genome of an underwater architect, the caddisfly
Stenopsyche tienmushanensis Hwang (Insecta: Trichoptera). GigaScience. 7(12):giy143.
https://doi.org/10.1093/gigascience/giy143.

Manni M, Berkeley MR, Seppey M, et al. 2021. BUSCO Update: Novel and Streamlined Workflows along
with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral
Genomes. Molecular Biology and Evolution. 38(10):4647–4654. https://doi.org/10.1093/molbev/msab199.

Marçais G, Kingsford C. 2011. A fast, lock-free approach for efficient parallel counting of occurrences of
k-mers. Bioinformatics. 27(6):764–770. https://doi.org/10.1093/bioinformatics/btr011.

Mariani V, Biasini M, Barbato A, et al. 2013. lDDT: a local superposition-free score for comparing protein
structures and models using distance difference tests. Bioinformatics. 29(21):2722–2728.
https://doi.org/10.1093/bioinformatics/btt473.

May ML. 2013. A critical overview of progress in studies of migration of dragonflies (Odonata:
Anisoptera), with emphasis on North America. J Insect Conserv. 17(1):1–15.
https://doi.org/10.1007/s10841-012-9540-x.

May ML. 2019. Odonata: Who They Are and What They Have Done for Us Lately: Classification and
Ecosystem Services of Dragonflies. Insects. 10(3):62. https://doi.org/10.3390/insects10030062.

Mendes FK, Vanderpool D, Fulton B, et al. 2021. CAFE 5 models variation in evolutionary rates among
gene families. Bioinformatics. 36(22–23):5516–5518. https://doi.org/10.1093/bioinformatics/btaa1022.

Minh BQ, Schmidt HA, Chernomor O, et al. 2020. IQ-TREE 2: New Models and Efficient Methods for
Phylogenetic Inference in the Genomic Era. Molecular Biology and Evolution. 37(5):1530–1534.
https://doi.org/10.1093/molbev/msaa015.

Newton L, Tolman E, Kohli M, et al. 2023 Jun 7. Evolution of Odonata: Genomic insights. Current
Opinion in Insect Science.:101073. https://doi.org/10.1016/j.cois.2023.101073.

Parkes K, Amos W, Moore N, et al. 2009. Population structure and speciation in the dragonfly Sympetrum
striolatum/nigrescens (Odonata: Libellulidae): An analysis using AFLP markers. European Journal of
Entomology. 106:179–184. https://doi.org/10.14411/eje.2009.021.

Phillips L. 2012. gnuplot Cookbook. Birmingham: Packt Publishing. 220 p.

Price BW, Winter M, Brooks SJ. 2022. The genome sequence of the blue-tailed damselfly, Ischnura
elegans (Vander Linden, 1820). Wellcome Open Res. 7:66.
https://doi.org/10.12688/wellcomeopenres.17691.1.

R F, R K-M, M K, et al. 2015. Extraordinary diversity of visual opsin genes in dragonflies. Proceedings of
the National Academy of Sciences of the United States of America. 112(11).
https://doi.org/10.1073/pnas.1424670112.

Ranallo-Benavidez TR, Jaron KS, Schatz MC. 2020. GenomeScope 2.0 and Smudgeplot for reference-free
profiling of polyploid genomes. Nat Commun. 11(1):1432. https://doi.org/10.1038/s41467-020-14998-3.

Rebora M, Salerno G, Piersanti S. 2022. Odonata perception is more than vision. In: Cordoba-Aguilar A,
Beatty C, Bried J, editors. Dragonflies and Damselflies: Model Organisms for Ecological and Evolutionary
Research. Oxford University Press. p. 0. https://doi.org/10.1093/oso/9780192898623.003.0007.

Rowe RJ. 1981. AN ANNOTATED KEY TO THE NEW ZEALAND ODONATA. 9.

Rowe RJ. 2004. Conservation of Odonata in the South Pacific and Australasia. International Journal of
Odonatology. 7(2):139–147. https://doi.org/10.1080/13887890.2004.9748206.

Russell RW, May ML, Soltesz KL, et al. 1998. Massive Swarm Migrations of Dragonflies (Odonata) in
Eastern North America. amid. 140(2):325–342. https://doi.org/10.1674/0003-
0031(1998)140[0325:MSMODO]2.0.CO;2.

Salmón P, Watson H, Nord A, et al. 2018. Effects of the Urban Environment on Oxidative Stress in Early
Life: Insights from a Cross-fostering Experiment. Integr Comp Biol. 58(5):986–994.
https://doi.org/10.1093/icb/icy099.

Sharip Z, Fauzi M. 2019. Microbial Contamination in Urban Tropical Lentic Waterbodies and Ponds along
Urbanization Gradient. Pertanika Journal of Tropical Agricultural Science. 42.

Shingate P, Ravi V, Prasad A, et al. 2020. Chromosome-level assembly of the horseshoe crab genome
provides insights into its genome evolution. Nature Communications. 11(1):2322.
https://doi.org/10.1038/s41467-020-16180-1.

Shiryev SA, Papadopoulos JS, Schäffer AA, et al. 2007. Improved BLAST searches using longer words for
protein seeding. Bioinformatics. 23(21):2949–2951. https://doi.org/10.1093/bioinformatics/btm479.

Šigutová H, Dolný A, Samways MJ, et al. 2022. Odonata as indicators of pollution, habitat quality, and
landscape disturbance. In: Cordoba-Aguilar A, Beatty C, Bried J, editors. Dragonflies and Damselflies:
Model Organisms for Ecological and Evolutionary Research. Oxford University Press. p. 0.
https://doi.org/10.1093/oso/9780192898623.003.0026.

Stanke M, Tzvetkova A, Morgenstern B. 2006. AUGUSTUS at EGASP: using EST, protein and genomic
alignments for improved gene prediction in the human genome. Genome Biology. 7(1):S11.
https://doi.org/10.1186/gb-2006-7-s1-s11.

Supek F, Bošnjak M, Škunca N, et al. 2011. REVIGO Summarizes and Visualizes Long Lists of Gene
Ontology Terms. Gibas C, editor. PLoS ONE. 6(7):e21800. https://doi.org/10.1371/journal.pone.0021800.

Suvorov A, Jensen NO, Sharkey CR, et al. 2017. Opsins have evolved under the permanent heterozygote
model: insights from phylotranscriptomics of Odonata. Molecular Ecology. 26(5):1306–1322.
https://doi.org/10.1111/mec.13884.

Suvorov A, Scornavacca C, Fujimoto MS, et al. 2021 Jul 29. Deep Ancestral Introgression Shapes
Evolutionary History of Dragonflies and Damselflies. Systematic Biology.:syab063.
https://doi.org/10.1093/sysbio/syab063.

Teets NM, Peyton JT, Colinet H, et al. 2012. Gene expression changes governing extreme dehydration
tolerance in an Antarctic insect. Proceedings of the National Academy of Sciences. 109(50):20744–20749.
https://doi.org/10.1073/pnas.1218661109.

Tennessen K. 2019. Libellulidae. In: Tennessen KJ, editor. Dragonfly Nymphs of North America: An
Identification Guide. Cham: Springer International Publishing. p. 407–576. https://doi.org/10.1007/978-3-
319-97776-8_12.

Tollett VD, Benvenutti EL, Deer LA, et al. 2009. Differential Toxicity to Cd, Pb, and Cu in Dragonfly
Larvae (Insecta: Odonata). Arch Environ Contam Toxicol. 56(1):77–84. https://doi.org/10.1007/s00244-
008-9170-1.

Tolman ER, Beatty CD, Bush J, et al. 2023. Exploring chromosome evolution in 250 million year old
groups of dragonflies and damselflies (Insecta:Odonata). Mol Ecol. 32(21):5785–5797.
https://doi.org/10.1111/mec.17147.

Tolman ER, Beatty CD, Bush J, et al. 2023 Feb 18. A Chromosome-length Assembly of the Black Petaltail
(Tanypteryx hageni) Dragonfly. Genome Biology and Evolution.:evad024.
https://doi.org/10.1093/gbe/evad024.

Tolman ER, Beatty CD, Kohli MK, et al. 2024. A molecular phylogeny of the Petaluridae (Odonata:
Anisoptera): A 160-Million-Year-Old story of drift and extinction. Molecular Phylogenetics and Evolution.
200:108185. https://doi.org/10.1016/j.ympev.2024.108185.

Tolman ER, Bruchim OR, Driever ES, et al. 2023. Changes in effective population size of Odonata in
response to climate change revealed through genomics. IJO. 26:205–211.
https://doi.org/10.48156/1388.2023.1917241.

Trizna M. 2020. assembly_stats 0.1.4. https://doi.org/10.5281/zenodo.3968775.

Turner PE. 1970. Allusive Dragons: Functional Integration and Structural Integrity among Natural
Populations of the Relict Dragonfly Tanyptery hageni. [PhD Thesis]. Berkeley, California: University of
California, Berkeley.

Uliano-Silva M, Nunes JGF, Krasheninnikova K, et al. 2021. marcelauliano/MitoHiFi: mitohifi_v2.0.
https://doi.org/10.5281/zenodo.5205678.

Vollmer W, Blanot D, De Pedro MA. 2008. Peptidoglycan structure and architecture. FEMS Microbiology
Reviews. 32(2):149–167. https://doi.org/10.1111/j.1574-6976.2007.00094.x.

Wang Y, Luan Y, Luo J, et al. 2023. 300 Million years of coral treaders (Insecta: Heteroptera:
Hermatobatidae) back to the ocean in the phylogenetic context of Arthropoda. Proceedings of the Royal
Society B: Biological Sciences. 290(2001):20230855. https://doi.org/10.1098/rspb.2023.0855.

Wang Y, Tang H, DeBarry JD, et al. 2012. MCScanX: a toolkit for detection and evolutionary analysis of
gene synteny and collinearity. Nucleic Acids Res. 40(7):e49. https://doi.org/10.1093/nar/gkr1293.

Ware J, Kohli MK, Mendoza CM, et al. 2022. Evidence for widespread gene flow and migration in the
Globe Skimmer dragonfly Pantala flavescens. International Journal of Odonatology. 25:43–55.

Ware JL, Beatty CD, Herrera MS, et al. 2014a. The petaltail dragonflies (Odonata: Petaluridae): Mesozoic
habitat specialists that survive to the modern day. Journal of Biogeography. 41(7):1291–1300.
https://doi.org/10.1111/jbi.12273.

Ware JL, Beatty CD, Herrera MS, et al. 2014b. The petaltail dragonflies (Odonata: Petaluridae): Mesozoic
habitat specialists that survive to the modern day. Journal of Biogeography. 41(7):1291–1300.
https://doi.org/10.1111/jbi.12273.

Winstanley WJ, Rowe RJ. 1980. The larval habitat of Uropetala carovei carovei (Odonata: Petaluridae) in
the North Island of New Zealand, and the geographical limits of the subspecies. New Zealand Journal of
Zoology. 7(1):127–134. https://doi.org/10.1080/03014223.1980.10423769.

Wissinger SA. 1988. Life History and Size Structure of Larval Dragonfly Populations. Journal of the North
American Benthological Society. 7(1):13–28. https://doi.org/10.2307/1467827.

Zhang C, Mirarab S. 2022. Weighting by Gene Tree Uncertainty Improves Accuracy of Quartet-based
Species Trees. Molecular Biology and Evolution. 39(12):msac215.
https://doi.org/10.1093/molbev/msac215.

828
Figure 1: (A) BUSCO results of the eight utilized Odonata genomes. (B) Number of genes recovered with
GALBA (Brůna et al., 2023) annotation, after filtering out genes that did not have a BLAST match with the
genome annotation of *P. flavescens.* (C) Species tree calculated from BUSCO Insecta genes (Manni et al.,
2021) using Weighted-Astral (Zhang and Mirarab, 2022) with posterior probability at each node. The
global lambda is shown at the root. Bottom right, the two species sequenced for this study, *Pachydiplax*
longipennis (left), and *U. carovei* (right).
*Annotation previously reported (H. Liu et al., 2022)
836

Figure 2. *TreeMaps summarizing rapidly expanding gene families in the (A) Libellulidae and (B) Petaluridae. Shared gene family expansions are represented under "Libellulidae" and "Petaluridae". The enriched biological process terms were summarized with REVIGO [44]. (C) From left to right, an adult of the petalurid T. hageni, an adult of the libellulid Pantala flavescens, (top) a fen habitat inhabited by T. hageni (California, USA) and (bottom) an urban pond where the authors have observed both P. longipennis and P. flavescens (New York, USA)*

845
846 *Figure 3: Demographic histories of A) P. longipennis, B) U. carovei, and C) Tanypteryx hageni estimated*
847 *with the Pairwise Sequential Markovian Coalescent (PSMC) using mutations of 1,2,3, and 4e-9. Bootstrap*
848 *replicates are shown as lighter lines. A generation time of .8 years was used for P. longipennis, and 5 years*
849 *was used for T. hageni and U. carovei. D) The semi-terrestrial nymphs of U. carovei (left) and T. hageni*
850 *(right) in their specialized fen habitats.*