Aberrant methylation and expression of TNXB promotes chondrocyte apoptosis and extracullar matrix degradation in hemophilic arthropathy via AKT signaling

Jiali Chen [1, 2, †], Zeng Qinghe [1, 2, †], Xu Wang [1, 2, †], Rui Xu [3], Weidong Wang [4], Yuliang Huang [4], Qi Sun [5], Wenhua Yuan [1], Pinger Wang [1], Di Chen [6, 7 *], Peijian Tong [8 *], Hongting Jin [1, *]

[1] Institute of Orthopaedics and Traumatology, the First Affiliated Hospital of Zhejiang Chinese Medical University , Zhejiang Provincial Hospital of Chinese Medicine, Hangzhou, Zhejiang, China

[2] The First College of Clinical Medicine, Zhejiang Chinese Medical University, Hangzhou, China

[3] Department of Orthopedics, Affiliated Hospital of Jiangxi University of Traditional Chinese Medicine, Nanchang, Jiangxi, China.

[4] Department of Osteology, The Second Affiliated Hospital of Zhejiang Chinese Medical University, Hangzhou, Zhejiang, China

[5] Department of Orthopaedic Surgery, Fuyang Orthopaedics and Traumatology Affiliated Hospital of Zhejiang Chinese Medical University, Hangzhou, Zhejiang, China

[6] Research Center for Computer-aided Drug Discovery, Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen, 518005, China.

[7] Faculty of Pharmaceutical Sciences, Shenzhen Institute of Advanced Technology,

Chinese Academy of Sciences, Shenzhen, 518055, China.

[8] Department of Orthopaedic Surgery, the First Affiliated Hospital of Zhejiang

Chinese Medical University , Zhejiang Provincial Hospital of Chinese Medicine,

Hangzhou, Zhejiang, China

* Corresponding authors:

Hongting Jin, Institute of Orthopedics and Traumatology, The First Affiliated

Hospital of Zhejiang Chinese Medical University, No.548, Binwen Road, Hangzhou,

Zhejiang Province, 310053, China. Email: hongtingjin@163.com.

Peijian Tong, Institute of Orthopedics and Traumatology, The First Affiliated

Hospital of Zhejiang Chinese Medical University, No.548, Binwen Road, Hangzhou,

Zhejiang Province, 310053, China. Email: tongpeijian@163.com.

Di Chen, Faculty of Pharmaceutical Sciences, Shenzhen Institute of Advanced

Technology, Chinese Academy of Sciences, No.1068, Xueyuan Avenue, Shenzhen,

518000, China. Email: di.chen@siat.ac.cn.

Conflicts of Interest

All authors declare that they have no conflict of interest.

Abstract

Backgroud: Recurrent joint bleeding in patients with hemophilia frequently results in hemophilic arthropathy (HA). Drastic degradation of articular cartilage is a major characteristic of HA, but its pathological mechanisms has not yet been clarified. Here, we conducted a genome-wide DNA methylation study with the goal of identifying critical genes for HA cartilage degeneration.

Methods: DNA was isolated from human osteoarthritis (N = 5) and HA (N = 5) articular cartilages and analyzed using the Infinium Human Methylation 850 BeadChip array. Adeno-associated virus-mediated shRNA and siRNA were used to knock down Tenascin XB (TNXB) in chondrocytes and F8$^{-/-}$ male mice, respectively. Then histopathological analysis, qPCR, Western blotting and immunofluorescence assays were conducted to detected chondrocyte homeostasis and HA progression.

Results: We found that Dnmt1 and Dnmt3a protein levels were increased in cartilages from HA patients compared with OA patients. Genome-wide DNA methylation analysis identified 1228 differentially methylated regions (DMRs) associated with HA. Functional enrichment analyses then revealed that DMR genes (DMGs) were related to extracellular matrix organization. Among these DMGs, TNXB protein expression was down-regulated in human and mouse HA cartilages. Further, the loss of *Tnxb* in F8$^{-/-}$ mouse cartilage provides a disease-promoting role in HA by augmenting cartilage matrix degeneration and subchondral bone loss. *Tnxb* knockdown also promoted chondrocyte apoptosis and inhibited phosphorylation of AKT. Importantly, AKT agonist showed a chondroprotective effect following *Tnxb*

knockdown.

Conclusions: Our study demonstrated that TNXB is a central mediator of cartilage

matrix degradation following joint bleeding, which functions by regulating the

activation of AKT. These mechanistic insights allow targeted development of

potentially new strategies for cartilage protection in HA.

Keywords: Hemophilic arthropathy; Methylation; Tenascin XB; Cartilage

degeneration; Chondrocyte apoptosis.

Introduction

Hemophilia is a X-linked bleeding disorder due to the deficiency of coagulation factor VIII (hemophilia A) and IX (hemophilia B). As a one of the most frequent rare diseases, hemophilia is a persistent, challenging condition for patients, physicians and society [1]. Joint bleeding (hemarthrosis), especially in knee joint, is the most common clinical manifestation in patients with severe hemophilia (i.e., plasma FVIII or FIX < 1 U/dl) [2]. Recent evidence also indicates that hemarthrosis may occur spontaneously in patients with moderate (plasma factor levels of 1– 5 UI/dl) or mild disease (plasma factor levels of >5 UI/dl) [3,4]. Repeated episodes of hemarthrosis eventually result in hemophilic arthropathy (HA), a debilitating and irreversible condition. HA often starts at an early age and characterized by joint impairment, chronic pain, and reduced quality of life [5-10].

Current efforts to prevent the development of HA are mainly focused on management of acute joint bleeding and optimizing prophylactic replacement therapy [2,11,12]. However, the widespread adoption of the modern hematological care has not conferred considerable protection against the development of HA; even in patients with hemophilia receiving intermediate- and high-dose prophylaxis, many still develop HA [13]. Pathological changes of HA begin with the first joint bleedings, either breakthrough or sub-clinical, usually during the first years of life [4,14]. They consist in progressive and irreversible alterations induced by the direct and indirect toxicity of the free blood in the joint space. The joint replacement surgery, such as Total Knee Arthroplasty (TKA), is considered an intervention for alleviating pain and restoring

 Given no disease modifying therapy available to intervene

in the HA perpetuating process, HA remains a persistent problem and challenge for

hemophilia patients. Thus, to understand the pathogenesis of HA and subsequently

explore possible targets for therapy is necessary.

One distinctive pathophysiological manifestation of HA is cartilage degeneration [15].

Cartilage is a rather inert tissue, consisting of matrix proteins and only one cell type,

chondrocytes that are responsible for production of matrix synthesis and rely on

synovial fluid for nutrients as cartilage lacks blood supply [16,17]. Nevertheless,

recurrent intra-articular bleeding creates a 'toxic' environment to cartilage, by

inducing synovial inflammation, hemosiderin accumulation and excessive mechanical

stress [15,18]. As a consequence, dysregulation of metabolism and abnormal apoptosis

occur in chondrocytes, ultimately resulting in deterioration of the cartilage matrix [2].

DNA methylation, a core epigenetic mechanism, is high dynamic and susceptible to

cues from the environment [19,20]. DNA methylation is the addition of a methyl group to

the cytosine residue of DNA molecules, which is catalyzed by DNA methyl

transferase (Dnmt) family of proteins that is comprised of three members: Dnmt1,

Dnmt3a, and Dnmt3b. DNA methylation is mainly located in the gene regulatory

region, usually repressing gene expression by blocking the transcriptional

accessibility of regulatory genomic regions [21]. Recently, multiple studies confirmed

the important role of abnormal DNA methylation in the pathogenesis of arthritis [22,23].

However, the epigenetic mechanisms underlying HA-related cartilage degradation have not been explored.

In this study, we conducted a genome-wide DNA methylation study with the goal of identifying differentially methylated genes (DMGs) and pathways for HA. Further, we knocked down *Tnxb*, a key DMG, in primary mouse chondrocytes and hemophilia A (F8$^{-/-}$) mice. The biological function and underlying mechanisms of TNXB in HA progression were also investigated *in vivo* and *in vitro*. Our findings provide a new mechanism for articular cartilage lesion in HA, which may lead to the discovery of novel therapeutic targets for the treatment of HA.

Materials and Methods

Cartilage specimen collection

Osteoarthritis (OA) is often used as "disease" control to reveal the characteristics in HA [24,25], although the mechanistic and phenotypic are different between HA and OA. In this study, 5 HA and 5 OA knee cartilage specimens were collected from patients undergoing total knee replacement surgery. Knee cartilage tissues were dissected and then rapidly frozen in liquid nitrogen or fixed in 4% paraformaldehyde. All subjects were Chinese Han and their demographic characteristics were listed in Supplementary table 1. The human cartilage samples were obtained from Department of Orthopedic Surgery at The First Affiliated Hospital of Zhejiang Chinese Medical University. This study was approved by the Ethics Committee of the First Affiliated Hospital of Zhejiang Chinese Medical University (2019-ZX-004-02).

Magnetic Resonance Image (MRI)

MRI examination: a GE 3.0 T Signa Excite superconducting MRI machine with a 32-channel body coil, a gradient field of 24 mT/m, and a creep rate of mT/(m.s) as well as a Siemens Verio 3.0 T superconducting magnetic resonance imaging machine with a 16-channel phased array coil were applied. An ultrashort echo time (UTE) pulse sequence (echo time 0.07 ms) was performed in the sagittal plane of the knee joint of the affected limb. Scanning parameters: layer thickness of 0.8 mm, field of view of 320 mm×320 mm, matrix of 160 mm×160 mm, intra-layer resolution of 0.6 mm×0.6 mm, and excitation of 2 times. Scanning software: NooPhase Wrap, Variable Bandwidth, Tailored RF.

Genome-wide DNA methylation profiling

Genomic DNA was prepared from cartilage specimens using QIAamp DNA Blood Mini Kit (QIAGEN, Germany) according to the manufacturer's instructions and was stored at -80 °C until used. Then bisulfite treatment of genomic DNA was performed with the EZ DNA methylation kit (Zymo Research) according to manufacturer's procedure. Genome-wide DNA methylation patterns were evaluated by Infinium Human Methylation 850 K BeadChips (Illumina), which determine the methylation levels of 853,307 CpG sites. R/Bioconductor (version 3.3.3) package ChAMP was used to process the Illumina intensity data (IDAT) files from the chip. DNA methylation level of each CpG site was described as a β value, ranging from zero (representing fully unmethylated) to one (representing fully methylated). Probes that had a detection P value of > 0.01 and those located on the X and Y chromosome were filtered away. We also removed SNP-related probes and all multi-hit probes.

BMIQ (Beta MIxture Quantile dilation) algorithm was used to correct for the Infinium

type I and type II probe bias. Furthermore, the regression models were adjusted for

age and sex. Differentially methylated probes (DMPs) at significance of $P < 0.05$

after the Benjamini & Hochberg correction for multiple testing. Differentially

Methylated Regions (DMRs) were identified by Bumphunter with default settings.

Additionally, we annotated the DMR-related genes (DMGs), and then performed

Gene Ontology (GO) and Kyoto Encyclopedia of Genes (KEGG) pathway analysis by

using the online tool DAVID (http://david.abcc.ncifcrf.gov/). The significant

pathway/ GO term was identified with P value < 0.05.

Mouse

The FVIII target gene knockout hemophilia (F8$^{-/-}$) mouse were purchased from

the Shanghai Laboratory Animal Center of the Chinese Academy of Sciences. All

mouse were kept in the Laboratory Animal Center of Zhejiang University of Chinese

Medicine. The animals were placed in the controlled environmental condition with

relatively constant air humidity and temperature as well as manual control of day and

night replacement for 12h. The experiment was approved by the Experimental Animal

Ethics Committee of Zhejiang University of Traditional Chinese Medicine

(LZ12H27001).

The mouse experimental model of HA

The HA model is constructed according to the description of Hakobyan et al [26].

F8$^{-/-}$ male mouse were induced with 3% isoflurane, and 1% isoflurane was maintained

under anesthesia. The right knee joint capsule of mouse was pierced with a 30g needle

for bleeding modeling, and the left knee of each animal was used as a control group.

Mouse were sacrificed at 4 and 8 weeks after injury, and knee joints were collected

for further experiments.

Intra-articular injection

F8$^{-/-}$ male mouse were induced with 3% isoflurane, and 1% isoflurane was

maintained under anesthesia. The right knee joint capsule of mouse was pierced with

a 30g needle for bleeding modeling. The needle of the insulin syringe was sagittally

inserted into the intercondylar area of the mouse right knee joint, and one dose of 10

μL AAV-Tnxb (GenePharma, concentration: 1×10^{10}/10μL) or AAV-NC was injected.

After 4 weeks, mouse was sacrificed, respectively, and the right knee joints were

collected for further experiments.

Micro-computed tomography (μCT) analysis

Micro-computed tomography (μCT) (Skyscan 1176, Bruker μCT, Kontich,

Gelgium) was used to analyze the knee joints. The area between the proximal tibia

growth plate and the tibial plateau was chosen as the region of interest. The

parameters collected form μCT were percent bone volume (BV/TV, %), Trabecular

thickness (Tb.Th, mm), Trabecular number (Tb.N, 1/mm) and Trabecular

Spacing(Tb.Sp, mm).

Histological analysis

The human cartilage samples were successively fixed in 4% paraformaldehyde

for 5 days, decalcified with 14% EDTA solution for 3 months, and the mouse samples

were successively fixed in 4% paraformaldehyde for 3 days, decalcified with 14%

EDTA solution for 14 days. These samples subsequently embedded in paraffin. Then

3-μm thick sections at the medial compartment of the joints were cut for Alcian blue

hematoxylin/Orange G (ABH) or Toluidine Blue staining to analyze the gross

cartilage structural changes. Histomorphometry analysis was performed through

Osteomeasure software (Decatur, GA).

Immunohistochemistry

The immunohistochemistry was examined to observe the expressions of protein

in cartilage. Briefly, the deparaffinized sections were soaked in 0.3% hydrogen

peroxide to block the activity of endogenous peroxidase, then blocked with normal

goat serum (diluted 1:20) for 20 min at room temperature. Subsequently, the primary

antibodies were added and incubated overnight at 4° C. The next day, the sections

were treated with secondary antibodies for 30 min and positive staining was visible by

using diaminobenzidine solution (Invitrogen, MD, United States). Then

counterstaining was performed with hematoxylin for 5 s. Anti-Col2a1 (Abcam,

ab34712, 1:200), Anti-Mmp13 (Abcam, ab39012, 1:300), Anti-TNXB (Proteintech,

13595-1-AP, 1:50), Anti-Bax (Huabio, ET1603-34, 1:200), Anti-Bcl-2 (Huabio,

ET1702-53, 1:200), Anti-p-Akt1 (Abclonal, AP0140, 1:200), Dnmt1(Abcam，1：200),

Dnmt3a (Abcam，1：200), Dnmt3b (Huabio，1：200) were used in this study.

Cell isolation and culture

Primary mouse chondrocytes were obtained from the femoral head of 2-week-old

C57BL/6 J mouse purchased from the Experimental Animal Center of Zhejiang

Chinese Medical University. Mouse were sacrificed and disinfected with 75% ethyl

alcohol. Specimens were isolated and rinsed by Phosphate Buffer Saline (PBS) 3

times. Then the cartilage tissues were digested with 0.25% collagenase P at 37 °C for

4h. Chondrocytes were cultured in DMEM/F-12 medium containing 10% fetal bovine

serum (FBS) and 1% streptomycin/penicillin in 5% CO_2 at 37 °C for further

experiment.

Tunel assay.

To evaluate the apoptotic cells in the chondrocytes, we performed a Tunel assay

according to the manufacturer's guideline (Beyotime, catalog C1088). Briefly,

following deparaffinage and rehydration, sections were permeabilized with

DNase-free Proteinase K (20 μg/mL) for 15 minutes at 37°C. Subsequently, slides

were treated with Tunel solution and incubated at 37°C for 1 hour in a dark

environment and counterstained with DAPI (1:1000; Solarbio) for 10 minutes. Finally,

TUNEL positive cells were detected by fluorescence microscope.

Transfection of siRNA-*Tnxb*

Chondrocytes were transiently transfected with siRNA targeting *Tnxb*

(GenePharma) or negative□control siRNA (scrambled; GenePharma) using

X□tremeGENE siRNA transfection reagent (Sigma) following the manufacturer's

instructions. All experiments were performed 72 hours after transfection, and the most

effective single siRNA was used for further experiments. After transfection for 48 h,

cells were treated with various concentrations of SC79 (10μM, Selleck) for 24 h.

Cell Immunofluorescence

The cells were fixed with 4% paraformaldehyde and blocked with goat serum for

1 hour. Then, cells were incubated with Col2a1 antibody (1:200; Abcam) and Mmp13

antibody (1:100; Abcam) overnight. The next day, the knee joints samples and

Chondrocytes with goat anti□rabbit antibody (1:1000; Invitrogen) conjugated with

Alexa Fluor 488 for 1 hour. Cells were stained with DAPI (1:1000; Solarbio).

Western blot analysis

Total proteins were extracted by RIPA buffer, and concentration of protein was

determined by bicinchoninic acid (BCA, Beyotime, Shanghai, China). Proteins were

separated on 10% SDS-PAGE gels and transferred to PVDF membranes. The

membranes were then blocked with 5% skim milk for 1 h, and were incubated with

primary antibodies (Col2a1, Abclonal, A1560, 1:1500), (Mmp13, Abclonal, A11755,

1:1000), (TNXB, Abclonal, A2535, 1:1000), (p-Akt1, Abclonal, AP0140, 1:1000),

(Akt1, Huabio, ET1609-47, 1:1000), (Bcl-2, Huabio, ET1702-53, 1:1000), (Bax,

Huabio, ET1603-34, 1:1000), (active-pro caspase-3, Huabio, ET1608-64, 1:1000),

(Cleaved-Caspase9, Abclonal, A22672, 1:1000), (pSmad2, CST, 8828, 1:1000)

overnight at 4°C. The membranes were washed 3 times, and then cultured with the

secondary antibody for 1 h. The immunoreactivity was detected with the ECL

substrate (Thermo Fisher Scientific, United States) on an Image Quant LAS 4000 (EG,

United States). The grey value was calculated by the software of Image J. *β-actin* was

used as an internal control in all western blot analysis.

Quantitative RT-PCR

TRIzol reagent (Invitrogen, United States) was used according to the

manufacturer's protocol to extract total RNA from chondrocytes. The total RNA

quantity and purity was evaluated by using NanoDrop 2000 (Thermo Fisher Scientific,

United States). Reverse transcription was carried out with cDNA Synthesis Kit

(Bmake, Beijing, China). The quantitative real-time-polymerase chain reaction (qPCR)

was conducted with SYBR Premix Ex Taq™ II (Takara, Dalian, China), and

performed on on a QuantStudio™ 7 Flex Real□Time PCR System. All of the PCR

reactions were repeated 3 times for each gene. The data were analyzed by the

2−ΔΔCT method using *β-actin* as the internal control. Supplementary table 2 showed

the primer sequences of target genes used in the current study.

Treatment of DNA Methyltransferase inhibitors

To investigate whether DNA methylation affects the expression of TNXB, we

treated primary mouse chondrocytes with RG108 or 5-Aza-dc (0, 1, 10, and 25 μM)

(RG-108, Beyotime, SD-1137; 5-Aza-2'-deoxycytidine, Beyotime, SD-1047) for 24 h.

The cells were then collected for qPCR to detect the mRNA level of *Tnxb*.

Statistics

All experimental data of this subject were analyzed by GraphPad Prism software

version 8.0, and the statistical results of measurement data were expressed as mean ±

standard deviation. 2-tailed unpaired parametric Student's t test or nonparametric

Mann-Whitney U test or one-way ANOVA with Tukey's multiple comparisons test

was used for comparison between different groups. The difference was statistically

significant with $P < 0.05$.

Results

Articular cartilages of HA patients show severe damage and aberrant elevations

of Dnmt1 and Dnmt3a

In this study, we collected cartilage samples in tibial plateau from OA and HA

patients undergo total knee replacement (TKR) surgery. Consistent with previous

reports, the HA cartilage exhibited severe damage and significant hemosiderin

deposition, compared with that of OA (Figure 1A). MRI analysis revealed the

subchondral bone loss in HA patients (Figure 1B). ABH staining further assessed the

cartilage degeneration, and detected a substantial sulfated glycosaminoglycan (sGAG)

depletion in HA cartilages (Figure 1, C and D). As expected, comparing to OA, HA

cartilages showed prominent decrease in cartilage matrix protein Col2a1 and increase

in expression of Mmp13, the primary enzymes responsible for cartilage degeneration

(Figure 1, E and F). Additionally, obvious increase in Dnmt1 and Dnmt3a protein

levels was also detected in HA cartilages (Figure 1, G and H), whereas no significant

changes were found in Dnmt3b (Supplementary Figure 1). These data confirmed the

severe damage in articular cartilage and subchondral bone of HA patients.

Genome-wide DNA methylation analysis reveals the epigenetic landscape of HA

cartilage

To understand the methylation profile associated with HA cartilage degeneration,

genome-wide DNA methylation alterations were examined in 10 subjects (5 OA

cartilages and 5 HA cartilages) by using 850K ChIP. The methylation levels of whole

genome-wide CpGs were in a classic bimodal distribution where most CpGs were

either slightly or highly methylated (Figure 2A). Of note, global DNA methylation

levels exhibited systematic differences between HA and OA, as illustrated by their

separation pattern in principle component analysis (Figure 2B). Overall, 1288

significant differentially methylated regions (DMRs) (P < 0.05 after the Benjamini

& Hochberg correction for multiple testing) were identified, with a mean fold change

in methylation difference of 0.90 (Figure 2C, and Supplementary Table 3). Compared

with the OA, 69.95% and 30.05% DMRs were hypermethylated and hypomethylated

in HA, respectively (Supplementary Figure 2A). Given the specificity of DMR, we

then determined whether the identified DMRs were preferentially colocalized with

some specific genomic features. The large majority (93.49%) of these DMRs were

located in gene promoters, 5'-UTRs, 3'-UTRs, and exons, with only 6.51%

overlapping intergenic regions (Supplementary Figure 2B).

To determine the biological functions of these significant DMRs, GO and KEGG

enrichment analysis were performed on DMR-related genes (DMGs). GO analysis

revealed enrichment for terms largely related to extracellular matrix (ECM)

organization, skeletal system development (Figure 2D, Supplementary Table 4). The

top two KEGG pathways were ECM-receptor interaction (P value = 5.2×10^{-6}) and

PI3K-Akt signaling pathway (P value = 5.3×10^{-4}) (Figure 2E, Supplementary Table 5).

DMGs presented in these pathways have mostly been proved to play an important role

in cartilage matrix homeostasis, such as *COL1A1*, *COL1A2*, *COMP* [27-29]. Among

them, we found TNXB were significantly low expressed in human HA cartilage

compared to OA cartilage by using IHC staining (Figure 2, F and G). However,

TNXB was not differentially expressed in OA and HA synovial membranes

(Supplementary Figure 2, C and D).

Decreased expression of Tnxb is associated with cartilage degradation by joint bleeding in F8$^{-/-}$ mice

To assess the potential relevance of TNXB in HA *in vivo*, we established a mouse model for HA, in which F8$^{-/-}$ mice received a needle puncture injury to the knee joint [30,31]. This needle puncture induced excessive bleeding and caused severe hemarthrosis (Supplementary Figure 3A). Analysis of cartilage tissue sections by Toluidine blue staining uncovered that joint bleeding remarkably promoted cartilage degeneration at 4 and 8 weeks after injury (Figure 3, A and B). IHC staining further demonstrated lower expression of Col2a1 and increased expression of Mmp13 (Figure 3, C-F). Next, micro-CT was used to assess the influence of hemarthrosis in osteogenesis. An obvious osteopenia was detected in subchondral bone at 4 and 8 weeks post injury (Supplementary Figure 3, B-E). As expected, the above histopathological changes in F8$^{-/-}$ mice following needle puncture were similar with what we observed in Human HA. Meanwhile, decreased expression of Tnxb and increased expressions of Dnmt1 and Dnmt3a were showed in articular cartilages 4 and 8 weeks following injury (Figure 3, G-L). Above results suggest that the suppression of TNXB in cartilage promotes the HA development.

TNXB **knockdown impairs chondrocyte metabolism and aggravates the progression of HA**

To verify the relationship between TNXB and methylation levels, we treated primary mouse chondrocytes with DNA Methyltransferase inhibitors RG108 or 5-Aza-dc, well-known to block DNA methylation [32,33]. After 24 hours treatment,

RG108 or 5-Aza-dc (10, 25 μM) significantly up-regulated the mRNA level of *Tnxb*,

indicating that the DNA methylation inhibits the expression of TNXB in chondrocytes

(Supplementary Figure 4, A and B). To further identify the biological functions of

TNXB in HA, we first knocked it down by specific siRNA in primary mouse

chondrocytes. The efficiency of *Tnxb* knockdown was confirmed by results of qPCR

and Western blot experiments (Figure 4, A-C). When compared with the

siRNA-control group, we found that *Tnxb* knockdown led to a decrease in *Col2a1*

mRNA expression and an increase in *Mmp13* mRNA expression (Figure 4, D and E).

Western blot and immunofluorescence staining analysis also showed a downregulated

expression of Col2a1 protein and upregulated expression of Mmp13 protein (Figure 4,

E-H). Those observations revealed that TNXB positively regulated the metabolism of

chondrocyte ECM.

Subsequently, we explored the effects of TNXB on HA cartilage degradation *in*

vivo, using intra-articular injection of adeno-associated virus (AAV) carrying

Tnxb-specific short hairpin RNA (shRNA) in F8$^{-/-}$ mice. 4 weeks after injection, more

severe cartilage degeneration was observed in *Tnxb*-KD mice, compared with vehicle

group (Figure 4, I and J). In addition, *Tnxb*-KD mice showed decreased expression of

Col2a1 and increased expression of Mmp13 in cartilages (Figure 4, K and L). The

subchondral bone of *Tnxb*-KD mice was significantly less than vehicle group mice

(Supplementary Figure 5, A-E). Collectively, the above results implied that TNXB

knockdown in cartilage accelerated the HA development.

TNXB deficiency sensitizes chondrocytes to apoptosis by inhibiting AKT

activation

Since blood exposure in joint cavity strongly causes chondrocyte apoptosis

(Supplementary Figure 6, A and B), we next explored the functional mechanisms of

TNXB in HA chondrocyte apoptosis. *Tnxb* knockdown induced the apoptosis in

chondrocytes, as evidenced by an increased proportion of Tunel-positive cells (Figure

5, A and B). Meanwhile, protein expression levels of pro-apoptotic Bax and

Cleaved-Caspase3 were significantly increased in *Tnxb*-KD chondrocytes, while

anti-apoptotic Bcl-2 was decreased (Figure 5, C-F). Moreover, AAV-shTnxb injection

also increased the number of Tunel-positive cells in articular cartilages in HA mice

(Figure 5, G and H). IHC staining confirmed that Bcl-2 expression was significantly

reduced in *Tnxb*-KD HA mice accompanied by enhanced expressions of Bax and

Caspase-3 (Figure 5, I-L).

It is well known that the PI3K/Akt signaling pathway regulates apoptosis [34-36], and

our KEGG analysis and IHC staining confirmed the involvement of this pathway in

HA cartilage degeneration (Figure 5, M and N). We, therefore, analyzed PI3K/Akt

pathway following *Tnxb* knockdown. In *Tnxb*-KD chondrocytes, p-Akt protein level

decreased significantly, while Akt level remained unchange (Figure 6, A and B). Next,

we sought to investigate whether Akt activation could attenuate chondrocyte

apoptosis induced by *Tnxb* knockdown. AKT agonist SC79 was used to treated

Tnxb-KD chondrocytes, and then its efficiency was verified by western blotting

(Supplementary Figure 5, C and D). 24 hours after SC79 treatment, the expressions of

Bcl-2 and Col2a1 (Figure 6, A, B and F) were upregulated, while Bax, Mmp13 and

Cleaved-Caspase9 expressions were markedly downregulated in *Tnxb*-KD

chondrocytes (Figure 6, A, C, E and G). These findings indicate that *TNXB*

knockdown induced chondrocyte apoptosis and ECM degradation through regulating

AKT activation.

Discussion

HA, a common manifestation of hemophilia, is typical characterized by dramatic

cartilage degradation [6]. Thus, the goal of our work was to provide a novel insight into

pathogenic mechanisms underlying HA cartilage degeneration. Figure 6 H

summarized our findings. Genome-wide DNA methylation analysis revealed the DNA

methylation changes in cartilages from hemophilia patients and identified *Tnxb* gene

associated with HA. The decreased expression of TNXB protein was also confirmed

in both human and mouse HA cartilages. Further, *Tnxb* knockdown promoted ECM

catabolism and chondrocyte apoptosis, and eventually accelerated the development of

HA in $F8^{-/-}$ mice. Meanwhile, decreased TNXB expression inhibited the activation of

AKT *in vivo* and *in vitro*. Notably, AKT agonist enhanced ECM synthesis and

suppressed apoptosis in *Tnxb*-KD chondrocytes. From these findings, it could be

hypothesized that abnormal methylation and decreased expression of TNXB are

responsible for disruption of cartilage homeostasis in HA.

In hemophlia patients, recurrent joint bleeding creates a toxic environment

including multifaceted processes such as hemosiderin deposition, inflammatory

response and mechanical pressure overload [37-40]. DNA methylation is a dynamic

epigenetic modification in response to environment. Since there are fewer epigenetic

studies on the pathological mechanisms of HA, we performed genome-wide DNA

methylation analysis, we found 1288 significant DMRs associated with HA with

69.95% hypermethylated, which was also supported by a corresponding increase in

Dnmt1 and Dnmt3a protein levels in HA cartilages. These results are in line with

previous reports showing that aberrant elevations of Dnmt1 and Dnmt3a promote

cartilage degeneration in destabilization of medial meniscus and aging models [33,41].

DNA methylation at gene promoter region is one of the important epigenetic

mechanisms in the regulation of gene expression [42,43]. The identified HA-related

DMRs showed high proportion in transcriptional start sites. These data suggest that

DNA methylation is a critical mediator in the pathology of HA. Nevertheless, due to

the severe erosion of HA cartilage, the mRNA levels of DMR-related genes were not

explored in this study. Besides, our sample size for DNA methylation sequencing is

relatively small. Thus, further studies using additional more HA samples will be

conducted to investigate the possible transcriptional alteration of identified genes as

well as their potential roles in HA development.

Furthermore, our enrichment analysis showed that DMR-related genes were

significantly enriched in ECM organization and ECM-receptor interaction terms. It is

consistent with previous reports that ECM acts as an epigenetic informational entity

capable of transducing and integrating intracellular signals via ECM-receptor

interaction [44-46]. DMR-related genes enriched in these terms, many are known to be

critical compositions of cartilage matrix, such as *COMP*, *COL1A1* and *COL1A2* [27-29].

In Articular cartilage, ECM provides nutrients and mechanical support for

chondrocytes [47], and its age-related stiffening epigenetically regulates gene expression

and compromises chondrocyte integrity [41]. Above findings highlight that ECM-related

genes are potential candidates for HA.

Tenascin-X (TNXB), an ECM glycoprotein, is the top differentially methylated

gene in our DNA methylation analysis of HA cartilages. In addition, DNA

methylation of TNXB has also been reported in whole blood from rheumatoid arthritis

patients and in retinal pigment epithelium from patients with age-related macular

degeneration [48,49]. Using IHC staining, we detected the downregulated expression of

TNXB in human and mouse HA cartilages. TNXB belongs to tenascin family, whose

members (TNC, TNR, TNXB, and TNW) share a similar domain pattern: an

N-terminal oligomerization domain, a series of epidermal growth factor-like repeats, a

variable number of fibronectin-type III (FNIII) repeats and a C-terminal

fibrinogen-like domain [50]. Emerging evidence suggests the involvement of TNC in

cartilage development and degeneration in arthritis [51,52]. Here, we prove for the first

time a role of TNXB in cartilage homeostasis *in vivo* and *in vitro*. Specifically, *Tnxb*

knockdown contributed to an increase in Mmp13 expression and decrease in Col2a1

expression in chondrocytes. In agreement with these findings, knockdown of *Tnxb* in

cartilages of F8$^{-/-}$ mice resulted in accelerated HA-like lesions. It has been previously

reported that TNXB blocks the interaction between TGF-β and its receptor in

endothelial cells. Since a potent role of TGF-β in chondrocyte differentiation, we

examined the effect of TNXB on TGF-β signaling, but did not observe significant

alterations in this signaling in *Tnxb*-KD chondrocytes (Supplementary Figure 7, A and

B). The FNIII repeats domain of TNXB undergoes alternative splicing to interact with different ECM proteins and growth factors, and then affects ECM network formation and three-dimensional collagen matrix stiffness [53]. Thus, future study would benefit from exploring the effect of the FNIII repeats domain in HA development to further clarify the specific mechanisms of TNXB.

The process of HA development is accompanied by the chondrocyte apoptosis which in part promotes the ECM degeneration [15,54]. We also observed remarkably increased apoptosis following joint bleeding in F8$^{-/-}$ mice. Interestingly, *Tnxb* knockdown could significantly induce apoptosis in chondrocytes *in vitro*, and even aggravate apoptosis in mouse HA cartilage, by enhancing expression of markers associated pro-apoptosis (Bax, C-Caspase3) and suppressing anti-apoptotic proteins (Bcl-2). In TNXB family, TNC was reported to inhibit human chondrosarcoma cell apoptosis by activation of AKT [55]. PI3K signaling pathway regulated AKT phosphorylation to suppress apoptosis [56,57]. Our KEGG analysis of DMGs showed the potential association of PI3K/AKT signaling with HA. We then detected much lower phosphorylation level of AKT in *Tnxb*-KD chondrocytes and HA models. These results are in line with previous report showing that the expression of p-AKT1 was down-regulated in the articular cartilage of patients with HA compared with that in patients with OA [58]. Notably, we further demonstrated that AKT agonist effectively restored the abnormal Mmp13 expression and apoptosis induced by *Tnxb* knockdown. Recent studies have shown that AKT signaling affected cartilage metabolism by regulating Mmp13 expression [59,60]. These findings indicate that AKT1 activation

mediates the effect of TNXB on HA chondrocytes apoptosis and ECM catabolism.

Thus, therapeutic effect of AKT agonist on HA development deserves further study.

Conclussions

Conclusively, our study provides the first comprehensive methylation profile in

the progression of knee HA, and shows that abnormal expression of TNXB is

important in HA cartilage degeneration by suppressing the activation of AKT1. These

findings provide a new insight into mechanisms responsible for HA disease and

suggest TNXB as a potentially novel therapeutic target for HA treatment.

List of abbreviations

HA: hemophilic arthropathy, DMRs: differentially methylated regions, DMGs:

DMR genes, TNXB: Tenascin XB, DNMT: DNA methyl transferase, ABH: Alcian

blue hematoxylin/Orange G, ECM: extracellular matrix, AAV: adeno-associated

virus.

Study approval

In the current study, the human cartilage samples were obtained from

Department of Orthopedic Surgery at The First Affiliated Hospital of Zhejiang

Chinese Medical University. This study was approved by the Ethics Committee of the

First Affiliated Hospital of Zhejiang Chinese Medical University (2019-ZX-004-02).

And all the animal operating procedures in this study were approved by the

Experimental Animal Ethics Committee of Zhejiang Chinese Medical University

(LZ12H27001).

Data Availability Statement

The data used to provide support for the results of this study can be obtained

from the corresponding authors.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the

corresponding author on reasonable request.

Conflicts of Interest

All authors declare that they have no conflict of interest.

Funding Statement

This research has been partially supported by the Natural Science Foundation of

China (Grant nos. 82274280, 82104891, 82074457), the Natural Science Foundation

of Zhejiang Province (LQ22H270006), the Fund for Distinguished Young Scholars of

Zhejiang Province (LR23H270001).

Author contributions

J.C., Q.Z., D.C., P.T., and H.J. conceptualized the study. Q.Z. conducted the *in*

vivo experiments. J.C., X.W. and R.X. collected the human samples and performed

the *in vitro* assay. W.W., Y.H. and Q.S. analyzed the data. J.C. wrote the original draft

and W.Y., P.W., D.C., P.T., and H.J. revised the manuscript critically for important

intellectual content. D.C., P.T, and H.J. supervised this study. All the authors read and

approved the final version of this manuscript.

Acknowledgments

We appreciate the great help from the Public Platform of Medical Research

Center, Academy of Chinese Medical Science, Zhejiang Chinese Medical University.

Reference:

1. Carulli, C., Innocenti, M., Linari, S., Morfini, M., Castaman, G., and Innocenti, M. (2021).
Joint replacement for the management of haemophilic arthropathy in patients with inhibitors:
A long-term experience at a single Haemophilia centre. Haemophilia *27*, e93-e101.
10.1111/hae.14169.

2. Gualtierotti, R., Solimeno, L.P., and Peyvandi, F. (2021). Hemophilic arthropathy: Current
knowledge and future perspectives. J Thromb Haemost *19*, 2112-2121. 10.1111/jth.15444.

3. Di Minno, M.N., Ambrosino, P., Franchini, M., Coppola, A., and Di Minno, G. (2013).
Arthropathy in patients with moderate hemophilia a: a systematic review of the literature.
Semin Thromb Hemost *39*, 723-731. 10.1055/s-0033-1354422.

4. Wyseure, T., Yang, T., Zhou, J.Y., Cooke, E.J., Wanko, B., Olmer, M., Agashe, R.,
Morodomi, Y., Behrendt, N., Lotz, M., et al. (2019). TAFI deficiency causes maladaptive
vascular remodeling after hemophilic joint bleeding. JCI Insight *4*. 10.1172/jci.insight.128379.

5. Aledort, L.M., Haschmeyer, R.H., and Pettersson, H. (1994). A longitudinal study of
orthopaedic outcomes for severe factor-VIII-deficient haemophiliacs. The Orthopaedic
Outcome Study Group. J Intern Med *236*, 391-399. 10.1111/j.1365-2796.1994.tb00815.x.

6. Bolton-Maggs, P.H., and Pasi, K.J. (2003). Haemophilias A and B. Lancet *361*, 1801-1809.
10.1016/S0140-6736(03)13405-8.

7. Fischer, K., Bom, J.G., Mauser-Bunschoten, E.P., Roosendaal, G., and Berg, H.M. (2005).
Effects of haemophilic arthropathy on health-related quality of life and socio-economic
parameters. Haemophilia *11*, 43-48. 10.1111/j.1365-2516.2005.01065.x.

8. Manco-Johnson, M.J., Abshire, T.C., Shapiro, A.D., Riske, B., Hacker, M.R., Kilcoyne, R.,
Ingram, J.D., Manco-Johnson, M.L., Funk, S., Jacobson, L., et al. (2007). Prophylaxis versus
episodic treatment to prevent joint disease in boys with severe hemophilia. N Engl J Med *357*,
535-544. 10.1056/NEJMoa067659.

9. Oldenburg, J. (2015). Optimal treatment strategies for hemophilia: achievements and
limitations of current prophylactic regimens. Blood *125*, 2038-2044.
10.1182/blood-2015-01-528414.

10. Olivieri, M., Kurnik, K., Pfluger, T., and Bidlingmaier, C. (2012). Identification and
long-term observation of early joint damage by magnetic resonance imaging in clinically
asymptomatic joints in patients with haemophilia A or B despite prophylaxis. Haemophilia *18*,
369-374. 10.1111/j.1365-2516.2011.02682.x.

11. Jackson, S.C., Yang, M., Minuk, L., Sholzberg, M., St-Louis, J., Iorio, A., Card, R., and Poon,
M.C. (2015). Prophylaxis in older Canadian adults with hemophilia A: lessons and more
questions. BMC Hematol *15*, 4. 10.1186/s12878-015-0022-8.

12. Schramm, W., Gringeri, A., Ljung, R., Berger, K., Crispin, A., Bullinger, M., Giangrande,
P.L., Von Mackensen, S., Mantovani, L.G., Nemes, L., et al. (2012). Haemophilia care in
Europe: the ESCHQoL study. Haemophilia *18*, 729-737. 10.1111/j.1365-2516.2012.02847.x.

13. Kleiboer, B., Layer, M.A., Cafuir, L.A., Cuker, A., Escobar, M., Eyster, M.E., Kraut, E.,
Leavitt, A.D., Lentz, S.R., Quon, D., et al. (2022). Postoperative bleeding complications in
patients with hemophilia undergoing major orthopedic surgery: A prospective multicenter
observational study. J Thromb Haemost *20*, 857-865. 10.1111/jth.15654.

14. Wojdasiewicz, P., Poniatowski, L.A., Nauman, P., Mandat, T., Paradowska-Gorycka, A.,

594   Romanowska-Prochnicka, K., Szukiewicz, D., Kotela, A., Kubaszewski, L., Kotela, I., et al.
595   (2018). Cytokines in the pathogenesis of hemophilic arthropathy. Cytokine Growth Factor
596   Rev *39*, 71-91. 10.1016/j.cytogfr.2017.11.003.

597   15.   Pulles, A.E., Mastbergen, S.C., Schutgens, R.E., Lafeber, F.P., and van Vulpen, L.F. (2017).
598   Pathophysiology of hemophilic arthropathy and potential targets for therapy. Pharmacol Res
599   *115*, 192-199. 10.1016/j.phrs.2016.11.032.

600   16.   Fermor, B., Christensen, S.E., Youn, I., Cernanec, J.M., Davies, C.M., and Weinberg, J.B.
601   (2007). Oxygen, nitric oxide and articular cartilage. Eur Cell Mater *13*, 56-65; discussion 65.
602   10.22203/ecm.v013a06.

603   17.   Lafont, J.E. (2010). Lack of oxygen in articular cartilage: consequences for chondrocyte
604   biology. Int J Exp Pathol *91*, 99-106. 10.1111/j.1365-2613.2010.00707.x.

605   18.   van Vulpen, L.F.D., Holstein, K., and Martinoli, C. (2018). Joint disease in haemophilia:
606   Pathophysiology, pain and imaging. Haemophilia *24 Suppl 6*, 44-49. 10.1111/hae.13449.

607   19.   Feil, R., and Fraga, M.F. (2012). Epigenetics and the environment: emerging patterns and
608   implications. Nat Rev Genet *13*, 97-109. 10.1038/nrg3142.

609   20.   Jirtle, R.L., and Skinner, M.K. (2007). Environmental epigenomics and disease susceptibility.
610   Nat Rev Genet *8*, 253-262. 10.1038/nrg2045.

611   21.   Stelzer, Y., Shivalila, C.S., Soldner, F., Markoulaki, S., and Jaenisch, R. (2015). Tracing
612   dynamic changes of DNA methylation at single-cell resolution. Cell *163*, 218-229.
613   10.1016/j.cell.2015.08.046.

614   22.   Wang, W., Yu, Y., Hao, J., Wen, Y., Han, J., Hou, W., Liu, R., Zhao, B., He, A., Li, P., et al.
615   (2017). Genome-wide DNA methylation profiling of articular cartilage reveals significant
616   epigenetic alterations in Kashin-Beck disease and osteoarthritis. Osteoarthritis Cartilage *25*,
617   2127-2133. 10.1016/j.joca.2017.08.002.

618   23.   Zhang, Y., Fukui, N., Yahata, M., Katsuragawa, Y., Tashiro, T., Ikegawa, S., and Lee, M.T.
619   (2016). Genome-wide DNA methylation profile implicates potential cartilage regeneration at
620   the late stage of knee osteoarthritis. Osteoarthritis Cartilage *24*, 835-843.
621   10.1016/j.joca.2015.12.013.

622   24.   Cooke, E.J., Zhou, J.Y., Wyseure, T., Joshi, S., Bhat, V., Durden, D.L., Mosnier, L.O., and
623   von Drygalski, A. (2018). Vascular Permeability and Remodelling Coincide with
624   Inflammatory and Reparative Processes after Joint Bleeding in Factor VIII-Deficient Mice.
625   Thromb Haemost *118*, 1036-1047. 10.1055/s-0038-1641755.

626   25.   Kalebota, N., Salai, G., Peric, P., Hrkac, S., Novak, R., Durmis, K.K., and Grgurevic, L.
627   (2022). ADAMTS-4 as a possible distinguishing indicator between osteoarthritis and
628   haemophilic arthropathy. Haemophilia *28*, 656-662. 10.1111/hae.14569.

629   26.   Hakobyan, N., Enockson, C., Cole, A.A., Sumner, D.R., and Valentino, L.A. (2008).
630   Experimental haemophilic arthropathy in a mouse model of a massive haemarthrosis: gross,
631   radiological and histological changes. Haemophilia *14*, 804-809.
632   10.1111/j.1365-2516.2008.01689.x.

633   27.   Chen, Y.R., Yang, K.C., Lu, D.H., Wu, W.T., Wang, C.C., and Tsai, M.H. (2019). The
634   chondroprotective effect of diosmin on human articular chondrocytes under oxidative stress.
635   Phytother Res *33*, 2378-2386. 10.1002/ptr.6425.

636   28.   Maly, K., Andres Sastre, E., Farrell, E., Meurer, A., and Zaucke, F. (2021). COMP and TSP-4:
637   Functional Roles in Articular Cartilage and Relevance in Osteoarthritis. Int J Mol Sci *22*.

10.3390/ijms22052242.

29.   Zhang, H., Chen, C., Cui, Y., Li, Y., Wang, Z., Mao, X., Dou, P., Li, Y., and Ma, C. (2019).
lnc-SAMD14-4 can regulate expression of the COL1A1 and COL1A2 in human chondrocytes.
PeerJ *7*, e7491. 10.7717/peerj.7491.

30.   Haxaire, C., Hakobyan, N., Pannellini, T., Carballo, C., McIlwain, D., Mak, T.W., Rodeo, S.,
Acharya, S., Li, D., Szymonifka, J., et al. (2018). Blood-induced bone loss in murine
hemophilic arthropathy is prevented by blocking the iRhom2/ADAM17/TNF-alpha pathway.
Blood *132*, 1064-1074. 10.1182/blood-2017-12-820571.

31.   Magisetty, J., Kondreddy, V., Keshava, S., Das, K., Esmon, C.T., Pendurthi, U.R., and Rao,
L.V.M. (2022). Selective inhibition of activated protein C anticoagulant activity protects
against hemophilic arthropathy in mice. Blood *139*, 2830-2841. 10.1182/blood.2021013119.

32.   He, Y., Zheng, Z., Liu, C., Li, W., Zhao, L., Nie, G., and Li, H. (2022). Inhibiting DNA
methylation alleviates cisplatin-induced hearing loss by decreasing oxidative stress-induced
mitochondria-dependent apoptosis via the LRP1-PI3K/AKT pathway. Acta Pharm Sin B *12*,
1305-1321. 10.1016/j.apsb.2021.11.002.

33.   Zhu, X., Chen, F., Lu, K., Wei, A., Jiang, Q., and Cao, W. (2019). PPARgamma preservation
via promoter demethylation alleviates osteoarthritis in mice. Ann Rheum Dis *78*, 1420-1429.
10.1136/annrheumdis-2018-214940.

34.   Cai, C., Min, S., Yan, B., Liu, W., Yang, X., Li, L., Wang, T., and Jin, A. (2019). MiR-27a
promotes the autophagy and apoptosis of IL-1beta treated-articular chondrocytes in
osteoarthritis through PI3K/AKT/mTOR signaling. Aging (Albany NY) *11*, 6371-6384.
10.18632/aging.102194.

35.   Hu, P.F., Chen, W.P., Bao, J.P., and Wu, L.D. (2018). Paeoniflorin inhibits IL-1beta-induced
chondrocyte apoptosis by regulating the Bax/Bcl-2/caspase-3 signaling pathway. Mol Med
Rep *17*, 6194-6200. 10.3892/mmr.2018.8631.

36.   Luo, J., Manning, B.D., and Cantley, L.C. (2003). Targeting the PI3K-Akt pathway in human
cancer: rationale and promise. Cancer Cell *4*, 257-262. 10.1016/s1535-6108(03)00248-4.

37.   Hooiveld, M.J., Roosendaal, G., van den Berg, H.M., Bijlsma, J.W., and Lafeber, F.P. (2003).
Haemoglobin-derived iron-dependent hydroxyl radical formation in blood-induced joint
damage: an in vitro study. Rheumatology (Oxford) *42*, 784-790.
10.1093/rheumatology/keg220.

38.   Srivastava, A. (2015). Inflammation is key to hemophilic arthropathy. Blood *126*, 2175-2176.
10.1182/blood-2015-09-665091.

39.   van Vulpen, L.F., van Meegeren, M.E., Roosendaal, G., Jansen, N.W., van Laar, J.M.,
Schutgens, R.E., Mastbergen, S.C., and Lafeber, F.P. (2015). Biochemical markers of joint
tissue damage increase shortly after a joint bleed; an explorative human and canine in vivo
study. Osteoarthritis Cartilage *23*, 63-69. 10.1016/j.joca.2014.09.008.

40.   Visser, A.W., de Mutsert, R., le Cessie, S., den Heijer, M., Rosendaal, F.R., Kloppenburg, M.,
and Group, N.E.O.S. (2015). The relative contribution of mechanical stress and systemic
processes in different types of osteoarthritis: the NEO study. Ann Rheum Dis *74*, 1842-1847.
10.1136/annrheumdis-2013-205012.

41.   Iijima, H., Gilmer, G., Wang, K., Bean, A.C., He, Y., Lin, H., Tang, W.Y., Lamont, D., Tai,
C., Ito, A., et al. (2023). Age-related matrix stiffening epigenetically regulates alpha-Klotho
expression and compromises chondrocyte integrity. Nat Commun *14*, 18.

10.1038/s41467-022-35359-2.

42.   Kucuk, C., Hu, X., Jiang, B., Klinkebiel, D., Geng, H., Gong, Q., Bouska, A., Iqbal, J.,
Gaulard, P., McKeithan, T.W., and Chan, W.C. (2015). Global promoter methylation analysis
reveals novel candidate tumor suppressor genes in natural killer cell lymphoma. Clin Cancer
Res *21*, 1699-1711. 10.1158/1078-0432.CCR-14-1216.

43.   Kulis, M., and Esteller, M. (2010). DNA methylation and cancer. Adv Genet *70*, 27-56.
10.1016/B978-0-12-380866-0.60002-2.

44.   Jones, F.S., and Jones, P.L. (2000). The tenascin family of ECM glycoproteins: structure,
function, and regulation during embryonic development and tissue remodeling. Dev Dyn *218*,
235-259. 10.1002/(SICI)1097-0177(200006)218:2<235::AID-DVDY2>3.0.CO;2-G.

45.   Kim, S.H., Turnbull, J., and Guimond, S. (2011). Extracellular matrix and cell signalling: the
dynamic cooperation of integrin, proteoglycan and growth factor receptor. J Endocrinol *209*,
139-151. 10.1530/JOE-10-0377.

46.   Alcaraz, L.B., Exposito, J.Y., Chuvin, N., Pommier, R.M., Cluzel, C., Martel, S., Sentis, S.,
Bartholin, L., Lethias, C., and Valcourt, U. (2014). Tenascin-X promotes
epithelial-to-mesenchymal transition by activating latent TGF-beta. J Cell Biol *205*, 409-428.
10.1083/jcb.201308031.

47.   Sutherland, A.J., Converse, G.L., Hopkins, R.A., and Detamore, M.S. (2015). The bioactivity
of cartilage extracellular matrix in articular cartilage regeneration. Adv Healthc Mater *4*,
29-39. 10.1002/adhm.201400165.

48.   Anaparti, V., Agarwal, P., Smolik, I., Mookherjee, N., and El-Gabalawy, H. (2020). Whole
Blood Targeted Bisulfite Sequencing and Differential Methylation in the C6ORF10 Gene of
Patients with Rheumatoid Arthritis. J Rheumatol *47*, 1614-1623. 10.3899/jrheum.190376.

49.   Porter, L.F., Saptarshi, N., Fang, Y., Rathi, S., den Hollander, A.I., de Jong, E.K., Clark, S.J.,
Bishop, P.N., Olsen, T.W., Liloglou, T., et al. (2019). Whole-genome methylation profiling of
the retinal pigment epithelium of individuals with age-related macular degeneration reveals
differential methylation of the SKI, GTF2H4, and TNXB genes. Clin Epigenetics *11*, 6.
10.1186/s13148-019-0608-2.

50.   Tucker, R.P., Drabikowski, K., Hess, J.F., Ferralli, J., Chiquet-Ehrismann, R., and Adams, J.C.
(2006). Phylogenetic analysis of the tenascin gene family: evidence of origin early in the
chordate lineage. BMC Evol Biol *6*, 60. 10.1186/1471-2148-6-60.

51.   Hasegawa, M., Yoshida, T., and Sudo, A. (2018). Role of tenascin-C in articular cartilage.
Mod Rheumatol *28*, 215-220. 10.1080/14397595.2017.1349560.

52.   Hasegawa, M., Yoshida, T., and Sudo, A. (2020). Tenascin-C in Osteoarthritis and
Rheumatoid Arthritis. Front Immunol *11*, 577015. 10.3389/fimmu.2020.577015.

53.   Valcourt, U., Alcaraz, L.B., Exposito, J.Y., Lethias, C., and Bartholin, L. (2015). Tenascin-X:
beyond the architectural function. Cell Adh Migr *9*, 154-165.
10.4161/19336918.2014.994893.

54.   Hooiveld, M., Roosendaal, G., Wenting, M., van den Berg, M., Bijlsma, J., and Lafeber, F.
(2003). Short-term exposure of cartilage to blood results in chondrocyte apoptosis. Am J
Pathol *162*, 943-951. 10.1016/S0002-9440(10)63889-8.

55.   Jang, J.H., and Chung, C.P. (2005). Tenascin-C promotes cell survival by activation of Akt in
human chondrosarcoma cell. Cancer Lett *229*, 101-105. 10.1016/j.canlet.2004.12.012.

56.   Tan, B.S., Tiong, K.H., Choo, H.L., Chung, F.F., Hii, L.W., Tan, S.H., Yap, I.K., Pani, S.,

Khor, N.T., Wong, S.F., et al. (2015). Mutant p53-R273H mediates cancer cell survival and

anoikis resistance through AKT-dependent suppression of BCL2-modifying factor (BMF).

Cell Death Dis *6*, e1826. 10.1038/cddis.2015.191.

57.   Testa, J.R., and Bellacosa, A. (2001). AKT plays a central role in tumorigenesis. Proc Natl

Acad Sci U S A *98*, 10983-10985. 10.1073/pnas.211430998.

58.   Zheng, L., Yao, Y., Luo, D., Han, Z., Zhang, X., Pang, N., Ding, M., Ye, H., Zhu, K., and Yi,

W. (2023). Iron regulates chondrocyte phenotype in haemophilic cartilage through the

PTEN/PI3 K/AKT/FOXO1 pathway. Hematology *28*, 2240585.

10.1080/16078454.2023.2240585.

59.   El Mabrouk, M., Sylvester, J., and Zafarullah, M. (2007). Signaling pathways implicated in

oncostatin M-induced aggrecanase-1 and matrix metalloproteinase-13 expression in human

articular chondrocytes. Biochim Biophys Acta *1773*, 309-320. 10.1016/j.bbamcr.2006.11.018.

60.   Lu, J., Feng, X., Zhang, H., Wei, Y., Yang, Y., Tian, Y., and Bai, L. (2021). Maresin-1

suppresses IL-1beta-induced MMP-13 secretion by activating the PI3K/AKT pathway and

inhibiting the NF-kappaB pathway in synovioblasts of an osteoarthritis rat model with

treadmill exercise. Connect Tissue Res *62*, 508-518. 10.1080/03008207.2020.1780218.

Figure 1. The severe cartilage damage in human HA. (A) Human cartilage samples in the tibial plateau were obtained from patients with OA and HA. Black arrow indicates hemorrhagic ferruginous deposits. **(B)** MRI imaging of the knee joint in patients with HA and OA. Red arrow indicates the wear area. **(C)** Representative images of ABH/OG staining for HA and OA cartilage. **(D)** The degree of cartilage degeneration was quantified according to the OARSI score. **(E)** Representative IHC staining of COL2A1 and MMP13. **(F)** Quantification of the proportion of COL2A1 and MMP13 positive regions in human cartilage. **(G)** Representative IHC staining of DNMT1 and DMMT3A. **(H)** Quantification of the proportion of NMT1 and DMMT3A positive cells in human cartilage. Red arrows indicate positive cells. Scale bar: 100 μm. Data were presented as means ± SD; n = 5 per group. And analyzed by 2-tailed unpaired parametric Student's t test, **$P < 0.01$, ***$P < 0.001$.

Figure 2. Genome-wide DNA methylation profile in HA cartilage and biological functions of DMR-Related genes. (A) Distribution of methylation level density of CpGs. Note: X = degree of methylation; Y = the CpG site density corresponding to the level of methylation. **(B)** Principal component analysis of DNA methylation data. **(C)** Heatmap shows the 700 significant DMRs between HA and OA. **(D)** Enriched GO terms for DMR-related genes. **(E)** KEGG enrichment analysis of DMR-related genes. **(F)** Representative IHC staining of TNXB. Red arrows indicate positive areas. Scale bar: 100 μm. **(G)** Quantification of the proportion of TNXB positive regions in human cartilage. Data were presented as means ± SD; n = 5 per group. And analyzed by 2-tailed unpaired parametric Student's t test, ***$P < 0.001$.

Figure 3. TNXB expression is drastically reduced in HA mouse cartilages. (A) Representative images and **(B)** Quantitative analysis of Toluidine blue staining for knee sections of F8$^{-/-}$ mice at 4 and 8 weeks following injury. Representative IHC staining of **(C)** Col2a1, **(E)** Mmp13, **(G)** Tnxb, **(I)** Dnmt1 and **(K)** Dnmt3a in F8$^{-/-}$ mice at 4 and 8 weeks post injury. Quantification of the proportion of **(D)** Col2a1, **(F)** Mmp13, **(H)** Tnxb, **(J)** Dnmt1 and **(L)** Dnmt3a positive regions. Scale bar: 100 μm. Data were presented as means ± SD; n = 6 mice per group. And analyzed by 2-tailed unpaired parametric Student's t test, ***$P < 0.001$.

Figure 4. knockdown of Tnxb in chondrocytes induces extracellular matrix degradation and accelerates the progression of HA. (A) qPCR and **(B)** Western

blotting analysis for Tnxb in mouse chondrocytes treated with siRNA-NC or

siRNA-*Tnxb*. **(C)** Corresponding quantification analysis of Tnxb protein.

Quantification of mRNA levels for **(D)** *Mmp13* and*Col2a1* in *Tnxb*-KD chondrocytes.

(E) Western blot and **(F)** corresponding quantification analysis for Mmp13 and

Col2a1 protein. **(G)** Representative immunofluorescence images of Col2a1 and

Mmp13 expression in *Tnxb*-KD chondrocytes. **(H)** Quantification of Col2a1 and

Mmp13 fluorescence intensity. Representative images of **(I)** Toluidine blue staining

and **(K)** IHC staining of Mmp13 and Col2a1 for knee sections of F8$^{-/-}$ mice at 4 weeks

after Intra-articular injection of AAV-sh*Tnxb*. **(J)** Quantitative detection of the area of

the tibial cartilage area. **(L)** Quantification of the proportion of Col2a1 and Mmp13

positive regions. Red arrow indicates the wear area. Scale bar: 100 μm. Data were

presented as means ± SD; n ≥ 3 in each group. And analyzed by 2-tailed unpaired

parametric Student's t test, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

**Figure 5. Tnxb knockdown increases apoptosis in chondrocytes and HA
cartilages.** **(A)** Representative images of TUNEL staining in Tnxb-KD chondrocytes

(B) Quantification for TUNEL positive cells. **(C-F)** Western blot and corresponding

quantification analysis for Bax, Cleaved-Caspase3, and Bcl-2. **(G)** TUNEL staining

for apoptosis in articular cartilage from Tnxb-KD HA mice. **(H)** Quantification for

TUNEL positive cells in articular cartilage. Representative IHC staining of Bax **(I)**,

Bcl-2 **(K)**, and p-Akt1 **(M)** in articular cartilage from *Tnxb*-KD HA mice.

Corresponding quantification of the proportion of **(J)** Bax, **(L)** Bcl-2 and **(N)** p-Akt1

positive regions. Red arrows indicate positive cells. Scale bar: 100 μm. Data were

presented as means ± SD; n ≥ 3 in each group. And analyzed by 2-tailed unpaired

parametric Student's t test, **$P < 0.01$, ***$P < 0.001$.

Figure 6. Treatment with AKT agonist decreases apoptosis in TNXB-KD

chondrocytes. (A) Western blot analysis for Col2a1, Mmp13, Bax, Bcl-2, p-Akt1,

Akt1 and Cleaved-Caspase9 in Tnxb-KD chondrocytes treated with SC79 (0 or 10μM)

for 24 hours. **(B-G)** Corresponding quantification of these proteins. **(H)** Schematic of

the role of TNXB in the pathogenesis of HA cartilage degeneration. Data were

presented as means ± SD; n = 4 per group. And analyzed by 2-tailed unpaired

parametric Student's t test or one-way ANOVA with Tukey's multiple comparisons

test, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A)

Col2a1			
Mmp13			
AKT1			
p-AKT1			
Bax			
Bcl-2			
Cleaved-Caspase9			
β-Actin			
siRNA-NC	+	-	-
siRNA-Tnxb	-	+	+
AKT agonist	-	-	+
Vehicle	+	+	-

B) Col2a1

C) Mmp13

D) pAKT1/AKT1

siRNA-NC	+	-	-
siRNA-Tnxb	-	+	+
AKT agonist	-	-	+
Vehicle	+	+	-

E) Bax

F) Bcl-2

G) Cleaved-Caspase9

siRNA-NC	+	-	-
siRNA-Tnxb	-	+	+
AKT agonist	-	-	+
Vehicle	+	+	-

H)

Hemophilic arthropathy

Methylation

TNXB level

Cartilage degeneration

Joint bleeding

TNXB

P AKT

Chondrocyte

MMP13 COL2A1

BCL2 BAX
Cleaved-CASP3

ECM metabolism imbalance

Apoptosis