Longitudinal profiling of the pregnancy plasma proteome through organic solvent precipitation and nano LC-MS/MS

George, A.L[1], Cook E[2], Smith G.C.S[2], Charnock-Jones D.S[2], O'Rahilly S[3], Reimann F[3], Gribble F.M[3], Kay R.G[3*].

1. Laboratory for Biological Mass Spectrometry, Biosciences Institute, Newcastle University, Newcastle-upon-Tyne, U.K.

2. Department of Obstetrics and Gynaecology, University of Cambridge, NIHR Cambridge Biomedical Research Centre, Cambridge, UK

3. Medical Research Council (MRC) Metabolic Diseases Unit, Wellcome-MRC Institute of Metabolic Science, University of Cambridge, Cambridge, UK

* Corresponding author (rgk27@medschl.cam.ac.uk)

Abstract

Proteins secreted from maternal, fetal, and placental tissue are vital for processes like placental development, immunotolerance, and fetal growth, and are associated with pregnancy complications, necessitating predictive biomarkers. In this study, we introduce an acetonitrile-based precipitation coupled with solid-phase extraction that addresses limitations of current low throughput blood-based biomarker discovery workflows. Our method is efficient and cost-effective, and identified 433 protein groups, showing specific tissue associations and enrichment in reproductive tissues such as the placenta, breast, and endometrium. Significant alterations in proteins related to hormonal regulation, immune modulation, and placental development throughout gestation were observed. This approach offers comprehensive insights into the circulating pregnancy proteome, but also provides a scalable solution for larger studies for biomarker discovery in the context of pregnancy complications.

Keywords: Pregnancy, Gestation, Plasma Proteomics, Biomarkers, Mass spectrometry

Introduction

Proteins, including hormones, growth factors, cytokines, cell-adhesion molecules and immunomodulators play a crucial role in establishing and maintaining pregnancy. They promote fetal growth and development and orchestrate adaptations in maternal physiology to support successful gestation (Napso et al. 2018, Newbern and Freemark 2011). Their regulated secretion from fetal-placental and maternal tissue into the bloodstream supports placental development, implantation, immunotolerance and fetal growth (Baud and Berkane 2019). Importantly, perturbations in these physiological processes can lead to conditions such as preeclampsia, fetal growth restriction and preterm birth, which can arise without clear predictive biomarkers, necessitating extensive research for new biomarkers to enable a more personalised approach to pregnancy care (Beernink et al. 2022, Paules et al. 2020, D'Silva, Hyett and Coorssen 2018).

Blood (plasma or serum) samples, collected during pregnancy provide a minimally-invasive and protein-rich source of biomarkers secreted from maternal, fetal and placental tissues (Shiefa et al. 2013). In recent years, these samples have been analysed with a focus on gaining a deeper understanding of the complex molecular adaptations key for healthy pregnancy, as well as to investigate indicators of placental function, fetal, and maternal health associated with pregnancy complications (Gaccioli et al. 2018). Various blood-based biomarkers have been proposed to stratify the risk of different pregnancy complications but have struggled to provide the necessary sensitivity and specificity for early risk prediction (MacDonald et al. 2022, Conde-Agudelo et al. 2013, Thomas et al. 2023).

Advancements in proteomic technology for multiplexed analysis have accelerated capabilities in biomarker discovery. Explorative studies of protein abundance during both healthy and complicated pregnancies have adopted antibody or aptamer-based approaches, enabling sensitive detection of multiple proteins simultaneously (Aghaeepour et al. 2018, Romero et al. 2017, Tarca et al. 2022). For example, Wang et al. explored a panel of 92 inflammatory markers using the proximity extension assay, O-Link platform, to investigate differences between patients with preeclampsia and healthy pregnant controls, as inflammation is reported to be involved in the occurrence and development of preeclampsia. They identified cysteine-cysteine motif chemokine ligand 20 (CCL20) as being significantly elevated (Wang et al. 2022). However, these platforms are limited by their capacity to measure only a predetermined set of proteins.

Liquid chromatography coupled with high resolution mass spectrometry (LC-MS) offers distinct advantages for biomarker discovery and clinical proteomics compared to affinity-based approaches, enabling untargeted analysis with exceptional analytical specificity. Peptide sequences with a single-mass unit difference can be differentiated, allowing quantification of proteoforms and post-translational modifications (Foreman et al. 2022, Katz et al. 2022). Yet, analysis of plasma or serum

samples is analytically challenging due to their high dynamic range of protein concentration, which is exacerbated by the presence of highly abundant proteins such as albumin, masking detection of the more relevant but less abundant putative biomarkers. Existing workflows for undepleted plasma analysis that can measure over 300 abundant circulating proteins in a 3-hour analysis, have not been adopted for identifying biomarkers of pregnancy complications (Geyer et al. 2016). This is because signalling molecules likely to be markers of placental function circulate at low concentrations. Rather, studies of the circulating pregnancy proteome utilise in-gel digestion, extensive peptide fractionation or immunodepletion to remove the most abundant proteins and reduce sample complexity prior to mass spectrometry (Yohannes et al. 2022, Pernemalm et al. 2019). However, these approaches are costly, time consuming and introduce experimental variability (Bader, Albrecht and Mann 2023). To overcome these challenges, a more efficient extraction approach is needed to eliminate sample preparation as a rate-limiting step and facilitate larger studies with appropriate sample sizes for verifying proposed markers.

Organic solvent precipitation is a simple, cost-effective and efficient approach for removing high abundance and high molecular weight proteins (Kay et al. 2008). This method, applicable to animal plasma as demonstrated in murine pregnancy studies (Napso et al. 2021, Lopez-Tello et al. 2023), is capable of enriching for low molecular weight proteins circulating at much lower concentrations. In this study, we employed an acetonitrile-based precipitation method coupled with solid-phase extraction to rapidly prepare plasma for mass spectrometry analysis. By precipitating the most abundant circulating proteins, we measured a plethora of proteins secreted from the placenta and other reproductive tissues, that significantly change in abundance during uncomplicated pregnancy in healthy controls at 12-, 20-, 28- and 36-weeks gestational age. Samples were obtained from the Pregnancy Outcome Prediction Study (POPs) which was a prospective sample collection including approximately 4000 participants from the Cambridge area in the United Kingdom (Pasupathy et al. 2008).

Materials and methods

Study design and sample collection

The POPs study recruited only nulliparous women, upon confirmation of pregnancy after a dating scan. Participants enrolled into the study had blood taken at approximately 12, 20, 28 and 36 weeks gestational age (wkGA). The POPs study was ethically approved by the Cambridge Local Research Ethics Committee 2 and written informed consent obtained from all participants. Both plasma and serum were collected at each visit, with the serum samples used for protein quantitation on the Cobas e411 clinical analyser. The plasma was collected into EDTA tubes, centrifuged at +4ºC, separated into 4 aliquots and then stored at -80ºC until required. Plasma samples at four time points from six

94   healthy volunteers with uncomplicated pregnancy were obtained for this prospective LC-MS/MS
95   study.

Sample preparation

Plasma samples (25 µL) were diluted with 25 µL of 6M guanidine hydrochloride and mixed on an
Eppendorf MixMate for 1 minute at 1500 RPM. Proteins were precipitated with 300 µL of 75% ACN
in water with 50 ng/mL of bovine insulin as an internal standard. The plate was mixed for 1 minute at
1500 RPM, before centrifugation at 3900 x g for 10 minutes at +4ºC. The supernatant was transferred
to an Eppendorf Lo-Bind 96 well plate and evaporated under oxygen free nitrogen (OFN) heated at
40 ºC. The dried sample residue was reconstituted into 200 µL of 0.1% formic acid in water and
transferred to a Waters HLB PRiME µ-elution SPE plate and pushed through carefully under positive
pressure OFN. The wells were washed with 200 µL of 0.1% formic acid in water, then 200 µL of 5%
methanol in 1% acetic acid in water. The proteins were eluted with 2 x 30 µL of 60% methanol in 10%
acetic acid in water into a fresh Lo-Bind plate and the eluant dried under OFN at 40 ºC. To reduce
and alkylate disulphide bonds, 75 µL of 10 mM DTT in 50 mM ammonium bicarbonate was added
and the plate incubated at 60 ºC for 1 hour before addition of 20 µL of 100 mM iodoacetamide in 50
mM ammonium bicarbonate and left in the dark for 30 minutes at room temperature. Samples were
left on the bench at RT in the light for 20 minutes before 10 µL of 100 µg/mL trypsin gold in 50 mM
ammonium bicarbonate was added and incubated overnight at 37 ºC. The digestion was halted by
the addition of 20 µL of 1% formic acid in water and 10 µL was injected for LC-MS/MS analysis.

Liquid chromatography mass spectrometry (LC-MS/MS)

Samples were analysed using a Thermo Fisher Ultimate 3000 nano LC system coupled to a Q
Exactive Plus Orbitrap mass spectrometer (ThermoScientific, San Jose, CA, USA). Tryptic peptides
were loaded onto a 0.3 x 5mm peptide trap column (ThermoFisher Scientific) at 30 µL per minute for
15 minutes before switching in line with a 0.075 × 250 mm nano easy column (ThermoFisher
Scientific) flowing at 300 nL/min. Initial gradient conditions were 2% ACN rising to 50% over 90
minutes, returning to initial conditions after a 20-minute wash at 90% ACN, giving a total run time of
130 minutes. Electrospray analysis was performed using a spray voltage of +1.8 kV, and a full scan
range of 400–1600 *m/z* was performed at a resolution of 75,000 before the top 10 ions of each
spectrum were selected for MS/MS analysis. Existing ions selected for fragmentation were added to
an exclusion list for 30 s.

Data analysis

Acquired mass spectrometry data was searched against the human Uniprot protein database
(downloaded Aug 2021) using PEAKS 8.5, with fixed carbamidomethylation on cysteine residues, and

variable modifications for oxidised methionines and asparagine / glutamine deamidation. Results were
filtered to include only proteins with a 1% false discovery rate (FDR) and at least 1 unique peptide
(Table S1A). Tissue enrichment analysis was performed with TissueEnrich
(https://tissueenrich.gdcb.iastate.edu/). Peptides from proteins related to pregnancy (selected based
on their number of spectral matches, and sensitivity and selectivity for that protein) were targeted for
label free quantitation (manual integration) and their peak areas were calculated using the data raw
files in Quan Browser (ThermoScientific, San Jose, CA, USA)(Table S1B). For global analysis, peak
areas were Log_2 transformed and filtered to include only proteins present in at least 4 of the 6 patient
samples, in at least one time point. Missing values of this filtered dataset were then imputed in
Perseus (version 2.0.1.1) to mimic proteins of low abundance that were below the limit of detection
during some timepoints (width 0.3, downshift 0.1). All data were expressed as a ratio to gestational
age week 12. Gene Ontology enrichment analysis of proteins significantly changing during gestation
was performed using the bioinformatic resource DAVID (Sherman et al. 2022). Protein-protein
interactions were analysed in STRING v12.0 and visualised in Cytoscape v3.9.1.

Automated immunoassay analysis

During the POPs study, maternal serum was analysed for alpha fetoprotein (AFP), chorionic
gonadotropin (hCG), Pregnancy associated plasma protein-A (PAPP-A), placental growth factor
(PlGF), and soluble FMS like Tyrosine kinase 1 (sFlt-1) on a Cobas e411 analyser (Roche
Diagnostics, Mannheim, Germany)(Gaccioli et al. 2017). The serum hCG concentrations of the
samples from the six individuals used in this study were compared against the respective LC-MS/MS
derived values.

Statistical analysis

Statistical analysis was performed in R (v4.0.4). One-way ANOVA repeated measures was used to
investigate differences in protein abundance associated with gestational age, followed by pairwise t-
tests for multiple comparison of gestation weeks (FDR adjusted p-value < 0.05).

Results and discussion

Characterisation of the circulating proteome

To profile the longitudinal plasma proteome during healthy pregnancy, maternal plasma samples were
collected at approximately 12, 20, 28 and 36 wkGA (**Figure 1A**). A central challenge with plasma or
serum-based proteomics is the limitation of achievable sensitivity, as the prevalence of highly
abundant proteins such as albumin dominate the measure of quantified peptides by mass
spectrometry, making the detection of less abundant biomarkers originating from disease tissue much
more difficult (Foreman et al. 2021).   Removal of these highly prevalent proteins is therefore

necessary, but immunodepleting or fractionation approaches require expensive reagents and lengthy workflows, which are less than optimal for a high-throughput workflow to accommodate the large sample sizes that are necessary for clinical biomarker studies. Protein precipitation using organic solvents offers an alternative method for protein enrichment, and can be rapidly performed in 96-well plate format, is cost-effective as there is no need for specialised reagents or antibodies, and reliably detects low-concentration plasma proteins (Kay et al. 2008, Napso et al. 2021). In this study, we diluted plasma with guanidine hydrochloride to dissociate small proteins bound to blood carrier proteins, like insulin-like growth factor 1 (IGF-1) which circulates bound to IGF binding protein 3. Subsequently acetonitrile (75% in water) was used to precipitate high abundance proteins but retain the soluble, less abundant proteins in the supernatant. The supernatant was evaporated and subjected to solid-phase extraction to remove unwanted hydrophobic small molecules, followed by in-solution digestion, and was analysed by label-free single-shot mass spectrometry (**Figure 1B**).

A total of 433 protein groups were identified, exhibiting specific tissue associations, as indicated in **Figure 1C** and **Figure 1D**. The acetonitrile precipitation method effectively depleted albumin, although other abundant plasma proteins such as low molecular weight apolipoproteins (APOA2, APOC3, APOA1, APOC2) that are secreted from the liver and circulate at µg/mL to mg/mL levels were also measured and ranked highest when ordering each identified protein by their average intensities across all samples (**Figure 1C**). Tissue enrichment analysis highlighted many proteins that are significantly enriched in the liver, an expected result, given the liver's major role in the biosynthesis of many plasma proteins. Notably, many of the other identified proteins are significantly enriched in female reproductive tissues, including the placenta, breast, cervix, or endometrium (**Figure 1D**).

For example, 24 proteins with elevated expression in the placenta were identified, thirteen of which are classified as tissue enriched, defined by the Human Protein Atlas to have a four-fold higher expression in the placenta compared to other tissues (**Figure S1A**). They included pregnancy-specific β-1-glycoproteins (PSG11, PSG3 and PSG4) that are uniquely produced by placental syncytiotrophoblast cells and play key roles in immunomodulation (Zhou et al. 1997). A recent mass spectrometry-based plasma proteomics study proposed 16 candidate biomarkers for ectopic pregnancy prediction, which included PSG11 and PSG3, but their workflow consisted of a two-step depletion of abundant plasma proteins with IGY-14 and super mix immune-depletion columns, followed by in-gel fractionation (6 fractions) to achieve such proteomic depth (Beer et al. 2023).

Compared to other methods, our single-shot analysis, with minimal sample handling, offers sufficient sensitivity to detect low abundance proteins known to circulate in maternal blood at low ng/mL range such as leptin and tachykinin-3. Leptin, for instance, is produced by both maternal and fetal adipose tissue, but also by the placenta to support implantation, human chorionic gonadotrophin production, placental growth, amino acid uptake, and mitogenesis (Taylor et al. 2015). Perturbations in leptin

levels have been correlated with various reproduction and gestation disorders such as polycystic ovary syndrome, recurrent miscarriage, gestational diabetes mellitus, pre-eclampsia, and intrauterine growth restriction (Pérez-Pérez et al. 2018). The neuropeptide, tachykinin-3 (TAC3), has enriched expression in the outer syncytiotrophoblast of the placenta and was among the lowest ranking proteins in the abundance plot of circulating proteins (**Figure 1C**). During normal pregnancy, it circulates at 0.29–0.61 nmol/L and is elevated in women suffering with preeclampsia (Lowry and Woods 2018, Pérez-Pérez et al. 2018). Identifying these low abundance proteins alongside abundant apolipoproteins and immunoglobulins, demonstrates the method's ability to measure proteins over a wide dynamic range.

Plotting the frequency of protein molecular weight demonstrates that precipitation using 75% acetonitrile effectively removes large molecular weight proteins and enriches those with a molecular weight of less than 60kDa (**Figure 1E**). In another study, a different ratio of blood to the organic solvent (1:2 ratio, 100% acetonitrile, v/v) was employed which identified mainly 2.5 kDa proteins, along with some in the range of 2.5 to 10 kDa, emphasising the adaptability of the extraction approach depending on the proteins of interest (Merrell et al. 2004).

Figure 1. *Characterisation of the circulating proteome during pregnancy.* ***A)*** *Sample collection time points of the six subjects.* ***B)*** *Sample preparation workflow to isolate the circulating proteome. Plasma proteins are diluted with guanidine hydrochloride before precipitation with acetonitrile and solid phase extraction. Extracted proteins are digested with trypsin and peptides analysed by LC-MS/MS.* ***C)*** *Ranked average abundance plot of all proteins in maternal plasma demonstrating depletion of abundant circulating proteins, enriching for less abundant signalling proteins.* ***D)*** *Tissue enrichment*

 analysis of all proteins detected revealed a significant enrichment in placental (blue), breast (pink)
and uterine (green) proteins. ***E)*** *Histogram showing frequency in molecular weight of the identified*
433 plasma proteins.

Peak areas of peptides from proteins with known roles in pregnancy were manually integrated and plotted by gestational age (**Figure 2**). This highlighted differences in the low molecular weight proteome and bioactive peptide profiles between individuals. For example, cystatin (13kDa), trefoil factor (7kDa), insulin-like growth factor 1 (IGF-1) (8kDa) and insulin-like growth factor 2 (IGF-2) are all low molecular weight proteins that regulate or support processes such as immune modulation, implantation, or fetal growth. Plasma levels of cystatin-C during normal pregnancy are described as stable during the first trimesters of pregnancy, but increase significantly during the final trimester (Schmitz et al. 2022). These patterns were also identified in our dataset, as cystatin-C increased over pregnancy in all participants (**Figure 2A**). Cystatin-C plasma levels have been explored as a marker of renal function in preeclampsia (Guo et al. 2012). Trefoil factors are peptides secreted from mucosal surfaces for protection and repair, primarily in the gastrointestinal tract. During pregnancy, circulating levels of TFF3 are reported to be 47 times higher than postpartum (Samson et al. 2008), and were significantly elevated in our dataset (**Figure 2B**). Growth factors, insulin-like growth factor 1 (IGF-1) and insulin-like growth factor 2 (IGF-2) that play important roles in cell growth, differentiation, and development, are involved in the development of the placenta and fetal tissues during pregnancy (Kaur et al. 2021) and were significantly higher compared to week 12 (**Figure 2 C and D**).

Corticoliberin, also known as corticotropin releasing hormone (CRF) is a peptide hormone secreted into the blood at very low levels by the hypothalamus but is also highly expressed in the placenta. Placental CRF secretion increases exponentially, peaking at birth and maternal plasma CRF levels increase accordingly from the second trimester. CRF levels in maternal plasma in the first and second trimester have been measured to remain within non-pregnant range (20pg/mL at 28 weeks) but increase significantly in the final trimester to 1320 pg/mL at 40 weeks triggering parturition (Wolfe et al. 1988, Zannas and Chrousos 2015), and high maternal levels have been associated with preterm labour (Smith 1998). In our dataset the peptide was detected in four of the six volunteers (**Figure 2E**). CRF has been identified previously by LC-MS approaches in placental tissue using solid phase extraction, in-gel digestion or a molecular-weight cutoff filter (Lin et al. 2019), but is reported only once before to be detected in plasma using an LC-MS method that included immunodepletion and extensive fractionation (Pernemalm et al. 2019).

GDF15, a member of the TGF-beta family, exhibits high expression in the placenta, and in healthy pregnancy circulating levels rise rapidly in maternal blood during the first trimester and continue to elevate until term, as shown with our dataset (**Figure 2F**). Importantly, immunoassay measurement of GDF15 in human plasma and serum has been reported to underestimate levels in individuals who

carry a histidine (H) to aspartate (D) variant at position 202 in the propeptide (Karusheva et al. 2022).
However, mass spectrometry can distinguish between these variants and can therefore be utilised to
gain an understanding of the origin of proteins that circulate in maternal blood. By exploiting the
specificity of mass spectrometry, the stoichiometry of the H and D peptides was measured to
determine the relative levels of maternal or fetal derived GDF15 in the maternal circulation, which
confirmed circulating GDF15 in human pregnancy is predominantly of fetal origin (Fejzo et al. 2023).

Prolactin (PRL) is a growth hormone which acts primarily on the mammary gland to stimulate lactation.
Additonally, it plays important roles in metabolism and immune system regulation (Ladyman et al.
2021). In healthy non-pregnant women, circulating prolactin levels are low but rise during gestation
(**Figure 2G**). Moreover, alpha-lactalbumin (LALBA), a human milk protein, that is expressed in the
glandular and ductal cells the breast. As a result of breast development during pregnancy, LALBA
levels rise significantly and due to tissue leakage, its presence can be detected in the blood at
increasing levels. Studies have indicated a significant increase until the mid-trimester, after which
concentrations remained stable until term (Martin et al. 1980), a trend reflected in our data for five of
the participants **(Figure 2H).**

Figure 2. Peak area of peptides from pregnancy-associated proteins relative to week 12 gestational
age. Numbers correspond to the amino acid positions of the monitored peptide in the primary amino
*acid sequence. **A)** cystatin-C (CST3), **B)** trefoil factor 3 (TFF3), **C)** insulin-like growth factor I (IGF-I),*
***D)** insulin-like growth factor II (IGF-II), **E)** corticoliberin (CRF), **F)** growth-differentiation factor 15*
*(GDF15), **G)** prolactin (PRL), **H)** alpha-lactalbumin (LALBA).*

Changes in the circulating proteome associated with gestational age.

For unbiased assessment of the method for measuring significant changes in protein abundance during pregnancy, we compared protein group abundance at each of the four time points by one-way repeated measures ANOVA. As the expression of proteins can change considerably over gestational weeks and circulating concentrations can vary hundred-fold, missing values were imputed to mimic proteins of low abundance that were below the limit of detection during some timepoints, resulting in comparison of 231 protein groups. This identified 16 proteins with significant changes associated with gestational age (**Figure 3A**)(**Table S1C and Table S1D)**. These protein groups exhibited distinct profiles as either increasing (red) or decreasing (blue) during pregnancy. Human chorionic gonadotropin hormone (hCG) is a glycoprotein consisting of two non-covalently linked subunits of alpha and beta (hormone specific subunit). While the hormone is produced by trophoblast cells at high levels very early on during pregnancy (and indeed is used as the measured target of point of care pregnancy tests), it peaks at week 10 before gradually declining and stabilising until term. Isoforms of the hormone with identical amino acid sequences exist (hCG, sulfated hCG, hyperglycosylated hCG, hCG free ß-subunit and hyperglycosylated hCG free ß-subunit) and are produced at different concentrations, by different cells and with different functions. Different immunoassays vary in their ability to recognise these isoforms forms, although there is no evidence necessitating accuracy of its measurement to diagnose pregnancy or non-pregnancy (Cole 2010). In our study, we detected beta subunits, CGB5, CGB7 and CGB2, as being highest in maternal plasma at week 12 of gestation, decreasing by week 20, then remaining constant during weeks 28 and 36 (**Figure 3A)**. These measurements align with known secretion patterns during healthy pregnancy. Relative quantification for hCG over the course of pregnancy measured by LC-MS/MS showed good correlation ($R^2 = 0.76$) with measurements from the Roche Cobas e411, which is a clinical analyser used in many hospital laboratories (**Figure 3B**).

Functional annotation through gene ontology (GO) enrichment analysis of the significantly changing proteins highlighted the importance of hormones in regulating gestation, as the molecular function, hormonal activity, was most significantly enriched (6.17E-07) (**Table S1E**). In addition to CGB5, CGB7 and CGB2, the growth factor placental lactogen significantly changed (CSH1 and CSH2). Placental lactogen, also known as chorionic somatotropin, is a peptide hormone produced by syncytiotrophoblast cells during human pregnancy. This growth hormone primarily circulates in the maternal blood, and is detectable from six weeks of gestation (Handwerger 1991, Sibiak et al. 2020). As seen in **Figure 3C**, levels increase throughout gestation, with levels thought to reflect total placental mass (Handwerger 1991).

Additionally, we noted changes in the endometrial protein, glycodelin (GD) also known as progestogen-associated endometrial protein (PAEP), which progressively declined in maternal

plasma over the course of gestation (**Figure 3A)**. Studies have shown concentrations in maternal serum to increase in the case of premature rupture of membranes (Wang et al. 2018). This low molecular weight protein has been detected previously in maternal serum using LC-MS and was identified in the aforementioned Beer at al. study of ectopic pregnancy, but a lengthy depletion and enrichment protocol was previously adopted, and verification was limited because of the moderate sample size (Beer et al. 2023).

We also observed significant changes in the interleukin-27 subunit beta (EBI3) (48kDa) between weeks 12 and 36 of gestation, reflecting the cytokine's role in regulating maternal immune tolerance and angiogenesis during pregnancy (Coulomb-L'Herminé et al. 2007). This protein has also been identified as a potential biomarker for preeclampsia, where the circulating plasma levels were higher in patients with preeclampsia (~1 ng/mL) compared to healthy women (~800 pg/mL) (Jahantigh et al. 2018).

Kisspeptin is a peptide of approximately 6 kDa that stimulates release of hypothalamic gonadotropin-releasing hormone and has a role in regulating trophoblast invasion. It is expressed in various tissues but is primarily secreted by the early syncytiotrophoblasts, increasing levels in the maternal circulation thousand-fold during healthy pregnancy. Kisspeptin has been proposed as a biomarker for placental function, as it increases progressively over pregnancy. Our data mirrored this secretion pattern, as levels increased with gestational age. Low circulating levels are associated with placental dysfunction, which is linked to complications such as miscarriage, hypertensive diseases of pregnancy, fetal growth restriction, and gestational diabetes (GDM) (Abbara et al. 2022).

In addition to placental-derived proteins, we observed changes in low molecular weight proteins with critical roles in supporting metabolic and nutritional adaptations during pregnancy. Proteins responsible for lipid transport and maintaining triglyceride homeostasis (APOL1, APOE) gradually increased with gestational age and were highest at week 36. Studies have suggested that levels of APOA2, APOC1, APOC3, and APOE are lower in women with gestational diabetes mellitus (Ramanjaneya et al. 2021).

Figure 3. *Changes in the circulating proteome associated with gestational age. **A)** Hierarchical*
clustering of proteins with significant changes in abundance associated with gestational age (ANOVA
p < 0.05). Increased abundance is represented in the heat map in red and those decreased are
*represented in blue. Gestational age is shown. **B)** Correlation between hCGbeta levels measured*
*relatively by LC-MS/MS compared with immunoassay quantification in all 24 samples. **C)** Chord*
diagram presenting gene ontology enriched terms in proteins that significantly change with gestational
age.

While our plasma preparation method effectively addresses the challenge of highly abundant proteins
that can dominate proteomic analyses, it predominantly enriches the low molecular weight proteome.
Nonetheless, the proteins identified hold significant clinical relevance, playing crucial roles in
biological processes vital for pregnancy, including implantation, placental development, and immune
modulation, which are known to be perturbed during pregnancy complications. We also demonstrate
the accuracy and quantitative performance of our method, as findings correlate well with
immunoassay data and expected secretion patterns of known pregnancy-related proteins during
normal pregnancy. In future applications, investigating these protein profiles during pregnancy
complications holds promising clinical value, as many proteins identified have previously been
proposed as biomarkers of pregnancy complication. Coupling this rapid extraction workflow with
recent advances in mass spectrometry, such as faster scanning instruments capable of performing
data independent acquisition experiments, supporting shorter analytical columns and gradients will
significantly reduce run-times without compromising quantitative performance. This updated
technology could accelerate biomarker discovery and verification in the context of pregnancy
complications by enabling the analysis of larger sample cohorts for pregnancy-related clinical studies.

Authorship contributions

360 ALG extracted, analysed the samples and performed the bioinformatics analyses. ALG and RGK
361 prepared the manuscript, and all authors reviewed the manuscript. EC, GCSS, and DSCJ designed
362 and led the POPs study and supplied the samples. SOR, FMG, FR and RGK guarantee the study.

Acknowledgments

The Prediction of Pregnancy Outcome Prediction Study (POPS) was supported by the Medical
Research Council (United Kingdom; G1100221) and the National Institute for Health Research (NIHR)
Cambridge Biomedical Research Centre (Women's Health theme). The instrumentation used for the
analysis was funded by the MRC Enhancing UK Clinical Research grant (MR/M009041/1). Research
in the FMG/FR group is funded by MRC (MC_UU_00014/3) and Wellcome (220271/Z/20/Z), and the
LC-MS peptidomics facility receives core support from MRC [MC_UU_00014/5] and Wellcome
[100574/Z/12/Z]).

Abbreviations

373	ANOVA	Analysis of variance
374	DAVID	The Database for Annotation, Visualization and Integrated Discovery
375	DTT	Dithiothreitol

376	GDM	Gestational Diabetes Mellitus
377	GO	Gene Ontology
378	OFN	Oxygen Free Nitrogen
379	PRIDE	PRoteomics IDEntificaitons
380	RT	Room Temperature
381	wkGA	Week of Gestational Age

Supporting information

*Figure S1. TissueEnrich analysis of detected proteins in maternal plasma. **Table S1A.** Raw data of protein peak areas exported from PEAKS search. **Table S1B.** Manual integration of peptides from protein with known pregnancy-related functions. **Table S1C**. Proteins significantly associated with gestational age. **Table S1D.** Pairwise comparison between weeks of gestational age. **Table S1E**. Gene ontology enrichment analysis of significantly changing proteins. **Table S1F.** LC-MS/MS file information, showing subject number and sampling time.*

References

Aghaeepour, N., B. Lehallier, Q. Baca, E. A. Ganio, R. J. Wong, M. S. Ghaemi, A. Culos, Y. Y. El-Sayed, Y. J. Blumenfeld, M. L. Druzin, V. D. Winn, R. S. Gibbs, R. Tibshirani, G. M. Shaw, D. K. Stevenson, B. Gaudilliere & M. S. Angst (2018) A proteomic clock of human pregnancy. *Am J Obstet Gynecol,* 218**,** 347.e1-347.e14.

Bader, J. M., V. Albrecht & M. Mann (2023) MS-Based Proteomics of Body Fluids: The End of the Beginning. *Mol Cell Proteomics,* 22**,** 100577.

Baud, O. & N. Berkane (2019) Hormonal Changes Associated With Intra-Uterine Growth Restriction: Impact on the Developing Brain and Future Neurodevelopment. *Front Endocrinol (Lausanne),* 10**,** 179.

Beer, L. A., X. Yin, J. Ding, S. Senapati, M. D. Sammel, K. T. Barnhart, Q. Liu, D. W. Speicher & A. R. Goldman (2023) Identification and verification of plasma protein biomarkers that accurately identify an ectopic pregnancy. *Clin Proteomics,* 20**,** 37.

Beernink, R. H. J., E. F. Zwertbroek, J. H. N. Schuitemaker, T. Cremers & S. A. Scherjon (2022) First trimester serum biomarker discovery study for early onset, preterm onset and preeclampsia at term. *Placenta,* 128**,** 39-48.

Cole, L. A. (2010) Biological functions of hCG and hCG-related molecules. *Reprod Biol Endocrinol,* 8**,** 102.

Conde-Agudelo, A., A. T. Papageorghiou, S. H. Kennedy & J. Villar (2013) Novel biomarkers for predicting intrauterine growth restriction: a systematic review and meta-analysis. *Bjog,* 120**,** 681-94.

Coulomb-L'Herminé, A., F. Larousserie, S. Pflanz, E. Bardel, R. A. Kastelein & O. Devergne (2007) Expression of interleukin-27 by human trophoblast cells. *Placenta,* 28**,** 1133-40.

D'Silva, A. M., J. A. Hyett & J. R. Coorssen (2018) Proteomic analysis of first trimester maternal serum to identify candidate biomarkers potentially predictive of spontaneous preterm birth. *J Proteomics,* 178**,** 31-42.

Fejzo, M., N. Rocha, I. Cimino, S. M. Lockhart, C. Petry, R. G. Kay, K. Burling, P. Barker, A. L. George, N. Yasara, A. Premawardhena, S. Gong, E. Cook, K. Rainbow, D. J. Withers, V. Cortessis, P. M. Mullin, K. W. MacGibbon, E. Jin, A. Kam, A. Campbell, O. Polasek, G. Tzoneva, F. M. Gribble, G. Yeo, B. Lam, V. Saudek, I. A. Hughes, K. K. Ong, J. Perry, A. Sutton Cole, M. Baumgarten, P. Welsh, N. Sattar, G. Smith,

D. S. Charnock Jones, A. P. Coll, C. L. Meek, S. Mettananda, C. Hayward, N. Mancuso & S. O'Rahilly
(2023) Fetally-encoded GDF15 and maternal GDF15 sensitivity are major determinants of nausea and
vomiting in human pregnancy. *bioRxiv*.
Foreman, R. E., A. L. George, F. Reimann, F. M. Gribble & R. G. Kay (2021) Peptidomics: A Review of Clinical
Applications and Methodologies. *J Proteome Res,* 20**,** 3782-3797.
Foreman, R. E., C. L. Meek, G. P. Roberts, A. L. George, F. Reimann, F. M. Gribble & R. G. Kay (2022) LC-MS/MS
based detection of circulating proinsulin derived peptides in patients with altered pancreatic beta cell
function. *J Chromatogr B Analyt Technol Biomed Life Sci,* 1211**,** 123482.
Gaccioli, F., I. Aye, U. Sovio, D. S. Charnock-Jones & G. C. S. Smith (2018) Screening for fetal growth restriction
using fetal biometry combined with maternal biomarkers. *Am J Obstet Gynecol,* 218**,** S725-s737.
Gaccioli, F., S. Lager, U. Sovio, D. S. Charnock-Jones & G. C. S. Smith (2017) The pregnancy outcome prediction
(POP) study: Investigating the relationship between serial prenatal ultrasonography, biomarkers,
placental phenotype and adverse pregnancy outcomes. *Placenta,* 59**,** S17-S25.
Geyer, P. E., N. A. Kulak, G. Pichler, L. M. Holdt, D. Teupser & M. Mann (2016) Plasma Proteome Profiling to
Assess Human Health and Disease. *Cell Syst,* 2**,** 185-95.
Guo, H. X., C. H. Wang, Z. Q. Li, S. P. Gong, Z. Q. Zhou, L. Z. Leng & M. Zhong (2012) The application of serum
cystatin C in estimating the renal function in women with preeclampsia. *Reprod Sci,* 19**,** 712-7.
Handwerger, S. (1991) Clinical counterpoint: the physiology of placental lactogen in human pregnancy. *Endocr*
Rev, 12**,** 329-36.
Jahantigh, D., M. Mousavi, F. Forghani, M. R. Javan, S. Movahedinia & M. Rezaei (2018) Association between
maternal circulating IL-27 levels and preeclampsia. *Cytokine,* 102**,** 163-167.
Karusheva, Y., M. Ratcliff, A. Mörseburg, P. Barker, A. Melvin, N. Sattar, K. Burling, A. Backmark, R. Roth, L.
Jermutus, E. Guiu-Jurado, M. Blüher, P. Welsh, M. Hyvönen & S. O'Rahilly (2022) The Common H202D
Variant in GDF-15 Does Not Affect Its Bioactivity but Can Significantly Interfere with Measurement of
Its Circulating Levels. *J Appl Lab Med,* 7**,** 1388-1400.
Katz, D. H., J. M. Robbins, S. Deng, U. A. Tahir, A. G. Bick, A. Pampana, Z. Yu, D. Ngo, M. D. Benson, Z. Z. Chen,
D. E. Cruz, D. Shen, Y. Gao, C. Bouchard, M. A. Sarzynski, A. Correa, P. Natarajan, J. G. Wilson & R. E.
Gerszten (2022) Proteomic profiling platforms head to head: Leveraging genetics and clinical traits to
compare aptamer- and antibody-based methods. *Sci Adv,* 8**,** eabm5164.
Kaur, H., B. S. Muhlhausler, C. T. Roberts & K. L. Gatford (2021) The growth hormone-insulin like growth factor
axis in pregnancy. *J Endocrinol*.
Kay, R., C. Barton, L. Ratcliffe, B. Matharoo-Ball, P. Brown, J. Roberts, P. Teale & C. Creaser (2008) Enrichment
of low molecular weight serum proteins using acetonitrile precipitation for mass spectrometry based
proteomic analysis. *Rapid Commun Mass Spectrom,* 22**,** 3255-60.
Ladyman, S. R., C. M. Larsen, R. S. Taylor, D. R. Grattan & L. M. E. McCowan (2021) Case-control study of
prolactin and placental lactogen in SGA pregnancies. *Reprod Fertil,* 2**,** 244-250.
Lin, Z., Y. Zhang, H. Pan, P. Hao, S. Li, Y. He, H. Yang, S. Liu & Y. Ren (2019) Alternative Strategy To Explore
Missing Proteins with Low Molecular Weight. *J Proteome Res,* 18**,** 4180-4188.
Lopez-Tello, J., H. E. J. Yong, I. Sandovici, G. K. C. Dowsett, E. R. Christoforou, E. Salazar-Petres, R. Boyland, T.
Napso, G. S. H. Yeo, B. Y. H. Lam, M. Constancia & A. N. Sferruzzi-Perri (2023) Fetal manipulation of
maternal metabolism is a critical function of the imprinted Igf2 gene. *Cell Metab,* 35**,** 1195-1208.e6.
Lowry, P. & R. Woods (2018) The placenta controls the physiology of pregnancy by increasing the half-life in
blood and receptor activity of its secreted peptide hormones. *J Mol Endocrinol,* 60**,** R23-r30.
MacDonald, T. M., S. P. Walker, N. J. Hannan, S. Tong & T. J. Kaitu'u-Lino (2022) Clinical tools and biomarkers
to predict preeclampsia. *EBioMedicine,* 75**,** 103780.
Martin, R. H., M. R. Glass, C. Chapman, G. D. Wilson & K. L. Woods (1980) Human alpha-lactalbumin and
hormonal factors in pregnancy and lactation. *Clin Endocrinol (Oxf),* 13**,** 223-30.
Merrell, K., K. Southwick, S. W. Graves, M. S. Esplin, N. E. Lewis & C. D. Thulin (2004) Analysis of low-abundance,
low-molecular-weight serum proteins using mass spectrometry. *J Biomol Tech,* 15**,** 238-48.
Napso, T., H. E. J. Yong, J. Lopez-Tello & A. N. Sferruzzi-Perri (2018) The Role of Placental Hormones in
Mediating Maternal Adaptations to Support Pregnancy and Lactation. *Front Physiol,* 9**,** 1091.

Napso, T., X. Zhao, M. I. Lligoña, I. Sandovici, R. G. Kay, A. L. George, F. M. Gribble, F. Reimann, C. L. Meek, R.
S. Hamilton & A. N. Sferruzzi-Perri (2021) Placental secretome characterization identifies candidates
for pregnancy complications. *Commun Biol,* 4**,** 701.
Newbern, D. & M. Freemark (2011) Placental hormones and the control of maternal metabolism and fetal
growth. *Curr Opin Endocrinol Diabetes Obes,* 18**,** 409-16.
Pasupathy, D., A. Dacey, E. Cook, D. S. Charnock-Jones, I. R. White & G. C. Smith (2008) Study protocol. A
prospective cohort study of unselected primiparous women: the pregnancy outcome prediction study.
BMC Pregnancy Childbirth, 8**,** 51.
Paules, C., L. Youssef, J. Miranda, F. Crovetto, J. M. Estanyol, G. Fernandez, F. Crispi & E. Gratacós (2020)
Maternal proteomic profiling reveals alterations in lipid metabolism in late-onset fetal growth
restriction. *Sci Rep,* 10**,** 21033.
Pérez-Pérez, A., A. Toro, T. Vilariño-García, J. Maymó, P. Guadix, J. L. Dueñas, M. Fernández-Sánchez, C. Varone
& V. Sánchez-Margalet (2018) Leptin action in normal and pathological pregnancies. *J Cell Mol Med,*
22**,** 716-727.
Pernemalm, M., A. Sandberg, Y. Zhu, J. Boekel, D. Tamburro, J. M. Schwenk, A. Björk, M. Wahren-Herlenius, H.
Åmark, C. G. Östenson, M. Westgren & J. Lehtiö (2019) In-depth human plasma proteome analysis
captures tissue proteins and transfer of protein variants across the placenta. *Elife,* 8.
Ramanjaneya, M., A. E. Butler, M. Bashir, I. Bettahi, A. S. M. Moin, L. Ahmed, M. A. Elrayess, S. C. Hunt, S. L.
Atkin & A. B. Abou-Samra (2021) apoA2 correlates to gestational age with decreased apolipoproteins
A2, C1, C3 and E in gestational diabetes. *BMJ Open Diabetes Res Care,* 9.
Romero, R., O. Erez, E. Maymon, P. Chaemsaithong, Z. Xu, P. Pacora, T. Chaiworapongsa, B. Done, S. S. Hassan
& A. L. Tarca (2017) The maternal plasma proteome changes as a function of gestational age in normal
pregnancy: a longitudinal study. *Am J Obstet Gynecol,* 217**,** 67.e1-67.e21.
Samson, M. H., E. M. Vestergaard, N. Milman, S. S. Poulsen & E. Nexo (2008) Circulating serum trefoil factors
increase dramatically during pregnancy. *Scand J Clin Lab Invest,* 68**,** 369-74.
Schmitz, E. M., M. L. Langelaan, M. J. de Groot, T. A. Ermens & A. M. Joosen (2022) Trimester-specific reference
intervals for cystatin C and neutrophil gelatinase-associated lipocalin during pregnancy. *Ann Clin
Biochem,* 59**,** 199-204.
Sherman, B. T., M. Hao, J. Qiu, X. Jiao, M. W. Baseler, H. C. Lane, T. Imamichi & W. Chang (2022) DAVID: a web
server for functional enrichment analysis and functional annotation of gene lists (2021 update).
Nucleic Acids Res, 50**,** W216-21.
Shiefa, S., M. Amargandhi, J. Bhupendra, S. Moulali & T. Kristine (2013) First Trimester Maternal Serum
Screening Using Biochemical Markers PAPP-A and Free β-hCG for Down Syndrome, Patau Syndrome
and Edward Syndrome. *Indian J Clin Biochem,* 28**,** 3-12.
Sibiak, R., M. Jankowski, P. Gutaj, P. Mozdziak, B. Kempisty & E. Wender-Ożegowska (2020) Placental Lactogen
as a Marker of Maternal Obesity, Diabetes, and Fetal Growth Abnormalities: Current Knowledge and
Clinical Perspectives. *J Clin Med,* 9.
Smith, R. (1998) Alterations in the hypothalamic pituitary adrenal axis during pregnancy and the placental
clock that determines the length of parturition. *J Reprod Immunol,* 39**,** 215-20.
Tarca, A. L., R. Romero, G. Bhatti, F. Gotsch, B. Done, D. W. Gudicha, D. M. Gallo, E. Jung, R. Pique-Regi, S. M.
Berry, T. Chaiworapongsa & N. Gomez-Lopez (2022) Human Plasma Proteome During Normal
Pregnancy. *J Proteome Res,* 21**,** 2687-2702.
Taylor, B. D., R. B. Ness, J. Olsen, D. M. Hougaard, K. Skogstrand, J. M. Roberts & C. L. Haggerty (2015) Serum
leptin measured in early pregnancy is higher in women with preeclampsia compared with
normotensive pregnant women. *Hypertension,* 65**,** 594-9.
Thomas, G., A. Syngelaki, K. Hamed, A. Perez-Montaño, A. Panigassi, R. Tuytten & K. H. Nicolaides (2023)
Preterm preeclampsia screening using biomarkers: combining phenotypic classifiers into robust
prediction models. *Am J Obstet Gynecol MFM,* 5**,** 101110.
Wang, X., K. C. Yip, A. He, J. Tang, S. Liu, R. Yan, Q. Zhang & R. Li (2022) Plasma Olink Proteomics Identifies
CCL20 as a Novel Predictive and Diagnostic Inflammatory Marker for Preeclampsia. *J Proteome Res,*
21**,** 2998-3006.

Wang, Y., H. Luo, G. Che, Y. Li, J. Gao, Q. Yang, B. Zhou, L. Gao, T. Wang, Y. Liang & L. Zhang (2018) Placental protein 14 as a potential biomarker for diagnosis of preterm premature rupture of membranes. *Mol Med Rep,* 18**,** 113-122.

Wolfe, C. D., S. P. Patel, E. A. Campbell, E. A. Linton, J. Anderson, P. J. Lowry & M. T. Jones (1988) Plasma corticotrophin-releasing factor (CRF) in normal pregnancy. *Br J Obstet Gynaecol,* 95**,** 997-1002.

Yohannes, E., D. L. Ippolito, J. R. Damicis, E. M. Dornisch, K. M. Leonard, P. G. Napolitano & N. Ieronimakis (2022) Longitudinal Proteomic Analysis of Plasma across Healthy Pregnancies Reveals Indicators of Gestational Age. *Int J Mol Sci,* 23.

Zannas, A. S. & G. P. Chrousos (2015) Glucocorticoid signaling drives epigenetic and transcription factors to induce key regulators of human parturition. *Sci Signal,* 8**,** fs19.

Zhou, G. Q., V. Baranov, W. Zimmermann, F. Grunert, B. Erhard, L. Mincheva-Nilsson, S. Hammarström & J. Thompson (1997) Highly specific monoclonal antibody demonstrates that pregnancy-specific glycoprotein (PSG) is limited to syncytiotrophoblast in human early and term placenta. *Placenta,* 18**,** 491-501.

535 **Table of Content (TOC)**