Interspecific relationships in a suboptimal habitat, an unexpected ally for the survival of the threatened Tenerife speckled lizard (*Gallotia intermedia*).

Gonzalo Albaladejo-Robles[1,2,5,6], Alejandro Escánez[2,3,4], Alicia V. Perera-Castro[5]

1. Research Department, Natural History Museum, Cromwell Road, London SW7 5BD, UK

2. Centre for Biodiversity and Environment Research, Department of Genetics, Evolution and Environment, University College London, London, UK.

3. Department of Ecology and Animal Biology, Campus As Lagoas-Marcosende, University of Vigo, Vigo, Spain.

4. MARE – Marine and Environmental Sciences Centre, ARNET – Aquatic Research Network, Regional Agency for the Development of Research, Technology and Innovation (ARDITI), Madeira, Portugal.

5. Asociación Tonina. C/Dr. Antonio González nº 5, 4º piso. La Laguna. S/C de Tenerife. Spain.

6. Departamento de Botánica, Ecología y Fisiología Vegetal, Universidad de La Laguna, 38206 La Laguna, Spain.

Keywords: Marine subsidies, Seagulls, Stable isotopes, Diet, protector species, Conservation, Reptiles

Abstract

Anthropogenic driven environmental changes are pushing species to the limits of their habitats. More often species are restricted to relic or suboptimal habitats that present the minimum requirements to sustain species populations. In this scenario of acerated environmental change and biodiversity loss, is fundamental to understand why species are able to survive in such suboptimal conditions. In this study, we conduct an isotopic trophic analysis along with a behavioural experiment to show how novel ecological interactions allow an endangered species to maintain stable populations on suboptimal habitats. We show how the Tenerife speckled lizard (*Gallotia intermedia*), a critically endangered endemic reptile from Tenerife Island (Canary Islands), is able to maintain stable populations in relic habitats thanks to its interactions with the yellow-legged gull (*Larus michahellis*) colony. The stable isotope analysis revealed that *G. intermedia* rely direct or indirectly on marine subsidies for its diet and that the foraging area of this reptile is likely to be restricted to the area occupied by a breeding colony of *L. michahellis*. Furthermore, an antipredator behaviour analysis showed that *L. michahellis* displayed a strong anti-predator or mobbing respond against cats (*Felis catus*), one of the main threats for *G. intermedia*, thus potentially providing some protection against predators for the reptiles inhabiting the seabird colony. Our results show how unusual and poorly studied biotic interactions can provide a valuable resources and conditions for the conservation of a critically endangered species inhabiting a suboptimal or relict habitat.

Introduction

Biodiversity is facing a human driven crisis. The last Living Planet Report showed an overall decrease of 68% in the abundance of monitored populations of vertebrates at a global scale (Dirzo et al., 2014; WWF, 2020). Globally, the main anthropogenic threats for biodiversity are habitat change, followed by overexploitation of species, climate change, pollution, and biological invasions (Brondizio et al., 2019; Dirzo et al., 2014). In many cases, these and other stressors are pushing species to the brink of extinction, with many species and populations forced to inhabit the edge of their distributions or refugial habitats, persisting over time in the face of biophysical disturbances (Keppel et al., 2012; Sedell et al., 1990). In many cases, these refuge habitats are considered suboptimal or marginal habitats because they do not provide the best habitat but the required environmental conditions (abiotic and a biotic context) for the species to reach a minimum fitness to fulfil their life histories, and in the case of refugee species this depictures their realized or contemporary niche (Brown & Carnaval, 2019; Kerley et al., 2020; Scheele et al., 2017). However, these marginal habitats may not be the most favourable and represent the least affected or most resilient habitat to the drivers of decline (Kerley et al., 2020). In contrast, populations located in these marginal environments may be less tolerant to increasing threats impacts (Scheele et al., 2017), which can turn these refuges into potential ecological traps (Robertson & Hutto, 2006). Even though the importance of this refugees is well recognized for biodiversity conservation (Braidwood et al., 2018; Keppel et al., 2015; Lea et al., 2016; Pavey et al., 2017; Selwood & Zimmer, 2020), it is still unclear, in most cases, which are the mechanisms and processes that allow species and populations to persist in these habitats, especially those involving biotic conditions (Gaston & Gaston, 2003).

Refugee species are those considered to have lost access to their past optimal habitats and are, therefore, currently confined to suboptimal habitats. Usually these species follow three 'rules of thumb' 1) have suffered a severe historical range contraction, 2) maintain slow population growth rates and/or low densities despite conservation efforts, and 3) appear to make use of anomalous resources (e.g., dietary shifts) in the current occupied habitat compared to historical records and that of close relatives (Kerley et al., 2012). Many threatened faunal species are recognized as

refugee species (e.g., Mediterranean monk seal, European bison, kakapo or giant panda) in their current distributional areas (Kerley et al., 2020).

Biotic conditions or biotic environment, such as interspecific interactions (i.e., mutualism, commensalism, predation, parasitism, competition, etc.), in addition to availability and dynamic of resources can generate positive interactions and boost species distribution and abundance in otherwise unsuitable areas (Godsoe et al., 2017; Soberón, 2007). Despite this, the relevance of these biotic interactions on shaping the large-scale distribution of a species is still a matter of debate (Godsoe et al., 2017; Peterson et al., 2011; Wiens, 2011). However, their effects at local scales on species ranges seems to be clear when barriers to its dispersal and abiotic gradients are weak (Davis et al., 1998; Guisan & Thuiller, 2005; Louthan et al., 2015; Wisz et al., 2013).

As a case of study on the importance of interspecific interactions in refugee species living on a suboptimal habitat, we focus our work on the endemic Canarian giant lizards which are currently represented by four species of the same genus: *Gallotia simonyi* from El Hierro, *Gallotia bravoana*, from La Gomera and *Gallotia intermedia* from Tenerife. The Giant lizards of the Canaries are a good example of species widely distributed in the past, that have seen their natural distribution drastically reduced (Gonzalez et al., 2014; Palacios-García et al., 2021). Human activities, such as habitat destruction and especially the introduction of invasive alien species (IAS) such as cats and rats, have driven all three species to the brink of (Afonso & Mateo, 2009; Bonnaud et al., 2011; García Márquez et al., 1997; Medina & Nogales, 2009). Nowadays these "giants" are restricted to small populations refugees in the most inaccessible and almost barren coastal cliffs of their respective islands (Nogales et al., 2001; J. C. Rando et al., 2000; J.-C. Rando et al., 2004). Due to all those factors, all the giant lizards from the Canary Islands are classify as Critically Endangered (CR) by the IUCN Red-list of threatened species and protected under the Spanish and international legislation. From these species, only the Tenerife speckled lizard, *G. intermedia*, presents two natural populations in its native range, Teno massif and Guaza cliffs. This makes this species a candidate of refugee species and a perfect species to study which are the conditions that allow these group of "giant" lizards to maintain their population in suboptimal habitats.

Here we focus on the *G. intermedia* population located in Guaza mountain cliffs, which was discover in 2003 (J.-C. Rando et al., 2004). Soon after its discovery this species was classified as Critically endangered by the IUCN (Mateo et al., 2019), and Endangered by the Spanish legislation (BOC, 2010; BOE, 2011). This population is restricted to inaccessible platforms and some ravines along the coastal cliffs of Guaza Mountain, located in the south-west of Tenerife Island, where it shares space with one of the largest resident colonies of yellow-legged gull (*Larus michahellis*) of the island (Figure 1) (Albaladejo et al., 2015; J.-C. Rando et al., 2004, p. 20). Although this population was described more than two decades from the elaboration of this manuscript, little is known about the biotic interactions between *G. intermedia* and its immediate surrounding environment. As stated before, Guaza cliffs are poor in vegetation cover. This alone suppose a challenge for *G. intermedia*. All *Gallotia* genus lizards are mostly herbivorous, feeding on fruits, leaves, and flowers of a wide variety of plants but also include arthropods and other small animals, including carrion (Fariña & Martín, 2013; Machado, 1985; Pérez-Mellado et al., 1999; Rica, 1981). However, the presence of a stable colony *L. michahellis* creates a potential source of food for *G. intermedia*. Previous studies report observations of *G. simonyi* and *G. intermedia* feeding on the remains leave by *L. michahellis* (Mateo, 2006; Siverio & Felipe, 2009). Furthermore, some studies show how seabird-derived resources are an important energy income for a variety of reptilian populations in areas with low primary productivity (Barret W.B. Anderson, D. A. Wait, L.L. Grismer, G.A. Polis & M.D. Rose et al., 2005; Kenny et al., 2017; Spiller et al., 2010). Therefore, we hypothesized that, in the absence of a substantial vegetation cover, *G. intermedia* is using seabird-derived resources left in the area by the permanent colony of *L. michahellis* as a complementary source of energy. Additionally, we believe that the interaction with *L. michahellis* goes beyond marine subsidies, actin also as indirect "bodyguards" or "protector species" for the reptiles (Gameiro et al., 2022). The abrupt and hard topography of Guaza coastal cliffs serve as a natural barrier for feral cats. Feral cats are considered to be the main responsible for the decline of several giant endemic lizards (genus Gallotia) in the Canaries, such as Gallotia simonyi (García Márquez et al., 1997), Gallotia intermedia (Albaladejo et al., 2015; Hernández et al., 2000; J.-C. Rando et al., 2004) and Gallotia gomerana (Nogales et al., 2001; J. C. Rando et al., 2000). The relatively low permeability of Guaza cliffs to this predator allows *G. intermedia* to inhabit the steepest profiles of these habitats with a lower risk of

predation. However, as stated before, *G. intermedia* also use the numerous platforms and some of the ravines present along the coastline. These areas are more permeable to cats, and so evidence of feral cat attacks and predation over *G. intermedia* have been recorded in those areas (Albaladejo et al., 2015; Flores Ravelo & Rando Reyes, 2021). These same ravines and platforms are used by *L. michahellis* colony for breeding and resting. The yellow-legged gull, as many other bird species, present an aggressive anti-predator behaviour specially during the breeding season (Clode et al., 2000; Frixione & Salvadeo, 2021; Gameiro et al., 2022; Guidos et al., 2023). Previous studies showed how seabird breeding colonies, due to its aggressive response to predators, work as facilitators for breeding of less aggressive and/or solitary species (Erwin, 1988; Gameiro et al., 2022; Quinn & Ueta, 2008; D. S. Richardson & Bolen, 1999). We hypothesize that *L. michahellis* anti-predator behaviour could also work in favour of *G. Intermedia*, providing populations inhabiting within the breeding grounds of this species some indirect protection against common predators. All these factors might explain, at least in part, the actual constrained distribution and conservation status of *G. intermedia* and to a larger extent the distribution of the rest of giant reptiles of the Canary Islands.

To evaluate our hypothesis, we conducted two distinct yet complementary experiments. In order to assess whether the populations of *G. intermedia* residing along the coastal cliffs of Guaza are trophically reliant on marine subsidies from the yellow-legged gull colony and the vegetation situated on the platforms and ravines both under and beyond the influence of the seagulls, we conducted a stable isotope analysis (SIA). In a second experiment we test whether the colony of *L. michahellis* was effective at keeping feral cats out of this species breeding area. To test the anti-predator behaviour of *L. michahellis* towards feral cats, we placed a cat decoy at random locations inside the breeding area of *L. michahellis* and recorded the behaviour of the animals. Finally, based on our results, as well as on the characteristics and conservation status of *G. intermedia* and the rest of the denoted, giant lizards of the Canary Islands, we propose these species to be considered refugee species.

2. Material and methods

2.1. Study system

Nowadays, the Tenerife speckled lizard naturally occurs in two different populations 30 km from each other, one in the Teno Massif and another one in Guaza Mountain (J.-C. Rando et al., 2004). In this study we focused in the later. The coastal cliffs of Guaza Mountain, located in the southern-west side of Tenerife, are part of an eroded volcanic dome which is considered a natural monument protected by the Canarian autonomous government and are included into the European Natura 2000 Network due to its unique geological features and the presence of protected birds' species under the European Birds Directive (Directive 2009/147/EC) (Figure 1). This volcanic dome is located between two of the biggest touristic localities of the island, Las Americas and El Palm-Mar. At a large scale, the vegetation is mainly form by a coastal shrubland with a plant association described as *Ceropegio fuscae-Euphorbietum balsamiferae salsoletosum divaricatae*, dominated by *Euphorbia balsamifera* (del Arco Aguilar et al., 2010). Guaza Mountain is located in the arid infra-mediterranean bioclimatic region (del Arco Aguilar et al., 2010), which is characterized by seasonal and scarce precipitation and constant high temperatures. *G. intermedia* inhabits the thin strip of land that contains the coastal cliffs-area and the transition area between those cliffs and the plateaus (J.-C. Rando et al., 2004; Ginoves et al., 2005) (Figure 1). To obtain an accurate description of the vegetation community in the study area, we conducted a randomized stratified sampling of the vegetation (Elzinga et al., 2001) for each of the topographic features present within the distribution area of the speckled lizard; these features included plateaus, ravines, and cliff platforms (Figure 1, details Supplementary Materials S1). We found differences between the plants communities present at the different topographic features; (Figure S.1); Plateaus were mainly dominated by *Euphorbia balsamifera*, *Limonium pectinatum* and *Frankenia ericifolia* ravines were characterized by *Plocama pendula* and *E. balsamifera* and the cliff platforms were mostly populated by *Patellifolia patellaris* (Table S1-S4).

2.3. Sampling

To measure the δ^{15}N and δ^{13}C composition of *G. intermedia* collected tissue samples of in 4 different localities within its distribution area (2 ravines and 2 platforms) (Figure 1). Reptiles were trapped using pit fall traps of 30x40x50 cm baited with raw tomato and placed using a regular grid of approximately 5 m along the area of the sampling

localities. Traps were placed during the early hours of the morning and let active for 7-8 hours for every trapping session. To avoid animals to suffer for excessive heat, we partially cover the entrance of the traps and check them every 30 to 45 minutes, depending on the heat. Each sampling locality was trapped for five consecutive days using 15-23 traps depending on the size of the sampling area. In addition to the pit falls, we used long-rods and lassos to capture the lizards that were more reluctant to fall into the baited traps. Once capture, reptiles were placed in a textile bag and weighted using a spring valance with a range of 300 g and a 0.1 g precision. After this, we recorded some other morphological important metrics such as snout ventral length or SVL (measured ventrally from the tip of the snout to the start of the cloaca), and pilleum width and length using precision callipers and rulers. Tissue samples of *G. intermedia* were collected by cutting, using surgical scissors, a fragment of 2-3 mm from the tip of the tail. *G. intermedia*, as well as the rest of *Gallotia* lizards, use the tail as a fat reservoir as well as a defensive decoy against predators and competitors. For these reasons tail autotomy is frequent amongst lacertids, and this procedure cause no harm to the individuals (García-Muñoz et al., 2011; Lin & Ji, 2005). Additionally, almost 90% of the capture lizards presented sights of partial or total tail regeneration (Table S5). This sampling method is less invasive than others and allowed us to obtain samples of relatively young tissue from a wide variety of cell types (skin, muscle, bone, etc). Handling of the individuals was carried out in the field, to reduce the interaction time between the animals and the researchers, and under strict sanitary conditions. Once all the samples and metrics were taken, we marked the animals with a temporal identification number (ID in Table S5), using xylene free nail paint (Boone & Larue, 1999) to avoid re-sampling, and released them in the vicinity of its capture point.

Since we were interested in the role of *L. michaaellis* on shaping the habitat use and diet of *G. intermedia*, we collected tissue samples from yellow-legged seagull nestlings. Tissue sample from chicks were collected by cutting, using a surgical scissor, a small section of the vane from the 2nd and 4th primary feather of each wing (Cherel et al., 2000; Han et al., 2021). We decide to extract the samples from the younglings from various reasons; first it simplified the sampling and handling of animals (adults of *L. michaaellis* are especially aggressive and cunning, making the direct trapping of the individuals difficult, especially in the harsh terrain of the cliffs, which limits both our movements and the placing of traps); 2) chicks are fast growing

individuals that depend fully on their parents for food. Therefore, we expected the isotopic signal of chick's feathers to be a strong proxy of the potential marine subsidies' seagulls bring to the breeding area (Cherel et al., 2000). Finally, leaves samples of the dominant plant species in each habitat (platfomrs, ravines and plateus) were collected by triplicate (Table S6). In the case of the Cactaceae *O. dillenii* only its fruit was sampled.

2.4. Sample processing

Once collected, the tail tips samples of *G. intermedia* as well plants and seagulls' feathers samples were rinsed thoroughly in distilled water to remove any surface contaminants before dried in an oven for 48 h at 45ºC, and then ground in a fine powder before isotopic analyses. Then, 0.3 ± 0.05 mg subsamples of dry powder were weighed in tin capsules for stable isotope analyses. All the tin cups were analysed by combustion at 900ºC in a Flash 1112 Elemental analyzer coupled to an isotope ratio mass spectrometer Delta C (ThermoFinnigan, ThermoFisher Scientific) in the lab facilities of the Centres Científics i Tecnològics de la Universitat de Barcelona (www.ccit.ub.edu) in Barcelona, Spain. The abundance of stable isotopes was expressed using the δ notation, where the relative variations of stable isotope ratios are expressed as parts per thousand (‰) deviations from predefined reference scales [atmospheric nitrogen for $\delta^{15}N$ and Vienna Pee Dee Belemnite (V-PDB) calcium carbonate for $\delta^{13}C$]. expressed as $\delta^y X = [(R_{sample} - R_{standard}) / (R_{standard})]$, where X is the element, y is the atomic mass of the stable isotope, and R_{sample} is the ratio of heavy to light isotopes ($^{13}C/^{12}C$ or $^{15}N/^{14}N$) in the sample and $R_{standard}$ is the ratio of the heavy isotope to light isotope in both working reference standards. These analyses were performed with an isotope ratio mass spectrometer. Salt of ammonium sulphate $(NH_4)_2SO_4$ (IAEAN1, $\delta^{15}N$= +0.4‰, and IAEAN2, $\delta^{15}N$ =+20.3 ‰), L-glutamic acid (USGS40, $\delta^{15}N$ = -4.5 ‰), and potassium nitrate, KNO_3 (IAEA-NO-3, $\delta^{15}N$ = +4.7 ‰) and caffeine (IAEA 600, δ15N = +1.1 ‰. For carbon, secondary isotopic reference materials of known $^{13}C/^{13}C$ ratios were used to a precision of 0.2 ‰, and these were namely: polyethylene (IAEACH7, $\delta^{13}C$ = −32.15‰), sucrose (IAEA CH6, $\delta^{13}C$ = −10.4‰), L-glutamic acid (USGS40, $\delta^{13}C$ = −26.4‰), and caffeine (IAEA600, $\delta^{13}C$ = −27.77‰). These isotopic reference materials were used to recalibrate the system once every 15 samples and were analysed in order to compensate any drift over time.

2.5. Stable isotopes statistical analysis

Since stable isotopes estimated values turned to be not normally distributed (Shapiro-wilk test p-value>0.05 in all cases), we used Kruskal-Wallis rank sum test along with a pairwise Wilcoxon rank-sum test to assess the effects of habitat type and *G. intermedia* on the values of $\delta^{15}N$ and $\delta^{13}C$. In addition, same tests were implemented to testing differences on isotopic values among plants samples from different sampling habitats and carbon metabolic pathways (C_3, C_4/CAM).

A Bayesian stable isotope mixing model was implemented to assess the relative proportional contribution of food resources derived from the three habitats sampled (plateau, platform, and ravine), as well as the contribution of seabird-derived resources, using the package MixSIAR version 3.1.10, (Stock et al., 2018) in R version 3.5.3 (R Core Team, 2021). Both the $\delta^{15}N$ and $\delta^{13}C$ values of potential food sources were incorporated into the model, as was the corresponding isotope values of Tenerife speckled lizards. Given that there is no previous information on the diet of *G. intermedia*, we ran the model using uninformative priors that assumed a generalist diet and weighted forage items groups equally ($\alpha = 1, 1, 1, 1$).

When analysing $\delta^{13}C$ and $\delta^{15}N$ in consumer tissue, there is often an offset between the diet and the tissue, termed diet-to-tissue trophic discrimination factor (TDF) or trophic enrichment factor (TEF). These TDFs may vary depending on diet source, growth rates, tissue type, or species (Caut et al., 2008). For accurate diet proportion estimations through the Bayesian isotopic mixing model, it is necessary to use adequate TDFs for each consumer species analysed. Because there are no published trophic discrimination factors for *G. intermedia* or for any other species of the genus *Gallotia*, we used TDFs determined for poikilotherm tissue ($\delta^{13}C$ 0.4 ± 0.14 ‰ and $\delta^{15}N$ 2.3 ± 0.2 ‰) according with (McCutchan Jr et al., 2003) and applied in other lizards' diet studies (Murray et al., 2016).

We estimated the proportion of trophic resources from the three habitats to lizards' diet, grouping into four forage item groups by the isotopic values of plants by habitat and plant metabolic pathway, since C_3 or CAM/C_4 plants differ significantly in their $\delta^{13}C$ values, as well as the seabird-derived resources. To estimate the proportion of the later we use as forage item group the isotopic values of seagull chicks by

subtracting beforehand the trophic discrimination factor (TDF) calculated on feathers of *L. michaellis* when feeding on marine preys according to Ramos et al. (2009). All the isotopic values of the different forage item groups, as well species their species composition is shown in Table S6. A Markov chain Monte Carlo (MCMC) sampling by running three replicate chains, each with 100,000 iterations, with a burn-in of 50,000 iterations and thin by 50 was used as setup in MixSIAR. Diagnostic tests (Gelmin–Rubin, Heidelberger–Welch and Geweke) and trace plots were examined for model convergence.

2.6. Body-guard effect sampling and analysis

To test the potential role of the yellow-legged seagulls in the protection of *G. intermedia* from predators we placed a cat decoy in 15 different locations along *L. michaaellis* breeding colony. For the cat decoy, we used a staffed feral cat fixed to a wooden plank in the stand position (Figure 1 and S3.c). This decoy was placed at random in the breeding colony by one of the members of the team while other researcher was strategically hide from the seabirds within direct sighting line to the decoy. Once the decoy was placed, we waited until the yellow-legged seagulls detect it, usually marked by multiple alarm songs by hovering birds, and then recorded the number and type of aggressive interaction against the cat decoy (similar to Gameiro et al., 2022). These interactions were classified into three basic types; 1) dives, in which seagulls perform a rapid "diving" flight towards the decoy and try to hit it with its legs; 2) Faecal attack, in which the birds defecated in mid-flight trying to hit the decoy; and 3) ground attacks, in this instance the animals try to land behind the decoy, out of its potential sight, and directly peck at the tail and rear legs of the decoy (Figure 1). This type of behaviour has been previously described in the literature as aggressive or territorial defensive behaviours (Frixione & Salvadeo, 2021; Larsen, 1991; Minias et al., 2020). To reduce our disturbance on the breeding colony, and to avoid potential behavioural changes on the individuals (Gaynor et al., 2018), we retired the decoy after two minutes of first aggressive interaction. Similarly, if the decoy was detected but no attack was recorded during the first 3 minutes after its detection, we also retired the decoy. Since we hypothesized that the anti-predator behaviour is greater within the breeding area of *L. michahellis*, once the decoy was hide, we counted the number of young chicks located in a radius of 5 m from where the decoy was placed.

Time was controlled using digital chronometers and attacks were recorded using a combination of binoculars and hidden wide-angle video cameras located in the vicinity of the decoy (GoPro Hero3 model CHDHN-301). The feral cat decoy used for this experiment presented the usual stripped brownish coloration common amongst wild-born feral cats (Figure S3.c). This animal was accidentally run over and killed on the island of Gran Canaria in 1985. It was later donated and prepared for the collection of the Zoology Department of the University of La Laguna, where it was returned after the experiment.

To test whether the number, type, and probability of aggressions towards the decoy was related with the density of birds within the breeding colony, we fitted a generalized linear models (GLMs) with a Poison and Binomial distributions using counts of the different attacks (Dives; Faecal-attack; ground attack; and total number of aggressions) and it's binomial form (1= attack, 0= no attack) as response variables, using the number of yellow-legged chicks as the explanatory variable. Seagull nests, and therefore chicks, are not uniformity distributed along the breeding colony. Terrain features such as slope can determine where nests and animals are located (Momberg et al., 2023) (see also supplementary materials S2). Due to this we added terrain ruggedness, calculated as the mean of the absolute elevation difference between a cell and its 8 surrounding neighbouring cells (Wilson et al., 2007), into the models as a co-variable.

All models were fitted by maximum likelihood, using an adaptative Gauss-Hermite quadrature approximation, implemented in the *lme4* R package version 1.1-2368 (Bates et al., 2015). To satisfy the assumptions of linear relationships of the residuals and to reconcile the different scales of the variables, terrain ruggedness was standardized by subtracting the mean and dividing by the standard deviation. Model performance was measure using the Nagelkerke's r^2 (Nagelkerke & others, 1991) implemented in the R package *performance* version 0.9.1 (Lüdecke et al., 2021). The statistical significance of the model effects was calculated using the Type II Wald Chi-square test, implemented in the *car* R package version 3.1-0 (Fox & Weisberg, 2019). All analysis, data manipulation, and spatial pre-processing was carried out using the R statistical software version 4.3.1 (R Core Team, 2021).

3. Results

3.1. Isotopic characterization of the Tenerife speckled lizard, habitat, and trophic resources.

In total, we analysed the $\delta^{15}N$ and $\delta^{13}C$ of 79 plant samples from 12 different species from 10 families, 10 feather samples from gull chicks (*L. michaellis*) and a total of 28 samples of Tenerife speckled lizards (*G. intermedia*) from Guaza cliffs. Lizards' samples were archived between July and September from two different habitats, (n=7) from isolated platforms, and (n=21) samples from ravines. Values of $\delta^{15}N$ for *G. intermedia* ranged from 11.53‰ to 15.94‰ and values of $\delta^{13}C$ ranged from -21.55‰ to -18.41‰, and their mean values expressed both as (mean ±SD) were 14.03 ± 1.01‰ for $\delta^{15}N$ and -19.76 ± 0.9‰ for $\delta^{13}C$, (Figure 2.a). Stable isotope ratios of $\delta^{15}N$ from platforms lizards ranged widely from 11.93‰ to 14.81‰ (mean ± SD = 13.5 ± 0.99 ‰) and from 11.53‰ to 15.95‰ (mean ±SD = 14.24 ± 0.97 ‰) in ravines lizards, while $\delta^{13}C$ values from platforms specimens ranged from -20.45‰ to -18.41‰ (mean ± SD = -19.1 ± 0.83‰) and from -21.55‰ to -18.78‰ (mean ± SD = -20.01 ± 0.83 ‰), for ravine habitat specimens. Lizards inhabiting ravines showed slightly higher values of $\delta^{13}C$ those of platforms (Kruskal-Wallis χ^2= 5.3265, df=1, p=0.021). There were no statistically significant differences in the values of $\delta^{15}N$ between reptiles inhabiting different habitats (χ^2= 2.61, df=1, p=0.1056).

Vegetation samples were classified as either C_3 ($\delta^{13}C$ values within −22‰ to −30‰ range) or C_4/CAM ($\delta^{13}C$ values in the −10‰ to −14‰ range). In this manner, $\delta^{13}C$ values of individual C3 plant species ranged from −28.12‰ to −19.16‰ and from −17.40‰ to −12.33‰ for C_4/CAM plants. $\delta^{15}N$ values for C3 plants ranged from 4.14‰ to 19.15‰ and from 9.14‰ to 20.11‰ for C_4/CAM plants (Table S7). Seagull chick feather samples ranged in $\delta^{13}C$ values from -18.92 ‰ to -20.03‰ and from 9.89 to 11.47 for $\delta^{15}N$ values (Table S7).

The composition of $\delta^{15}N$ and $\delta^{13}C$ across the four foraging items groups formed by plants metabolic pathway and habitat were statistically significant different in both isotopes ratios values ($\delta^{15}N$ χ^2=59.269, df=3, p<0.001; and $\delta^{13}C$ χ^2= 52.315, df=3, p<0.001) (Figure 2.a-c). Pairwise comparisons revealed statistically significant differences among $\delta^{15}N$ values in platforms C3 and C_4/CAM plants and ravine-plateu

plants, as well from seagulls-derived resources. Platform plants showed in general higher $\delta^{15}N$ values (15.89 ± 2.20‰) than ravine/plateu plants (8.57 ± 4.25‰), even than seagull-derived resources (10.49 ± 0.45‰) (Table S7). Whereas for $\delta^{13}C$ values only C_3 plants showed no significant differences (p>0.05), independently of their habitat, being statistically different from C_4/CAM plants (-14.45 ± 1.49‰) which presented the highest values among plants following by seagulls-derived resources (-19.53 ± 0.37‰) (p<0.001), see Table S7 for full numerical values of pairwise Wilcoxon rank-sum test.The MixSIAR model estimated that the mean (±SD) contribution of seabird-derived resources dominated the dietary composition of the Guaza mountain population of *Gallotia intermedia* (62.2 ± 14.3%), following by C_3 plants of platforms (15 ± 0.42%) and C_3 plant from ravines and plateu habitats (12 ± 10.1%) (Table 2, Figure. 2.a).

3.2. Bodyguard effect

We recorded a total of 165 aggressive interactions between *L. michahellis* and the feral cat decoy across 14 different locations within the breeding colony; 135 dives; 27 faecal attacks; and 3 ground or direct attacks (the later ones were not included in the analysis due to its low frequency). The frequency of dives (Poisson family models), as well as the total number of attacks were higher in the areas with a higher number of *L. michahellis* chicks (Type II Wald Chisquare test; χ^2 = 77.130, d.f = 1, P < 0.001; and χ^2 = 64.397, d.f = 1, P < 0.001, respectively) (Figures 3.a and 4.a). Model predictions for the frequency of dives and total number of attacks showed that these are mainly restricted to the limits of the yellow-legged seagull breading colony, being strongest in the core areas of the breeding colony and fading away towards their edges (Figures 3.c and 4.c). However, Fecal attacks were independent of the number of chicks (χ^2 =0.073, d.f = 1, P=0.785), and the ruggedness of the terrain (χ^2 =1.027, d.f = 1, P=0.31). Model predictions showed that faecal attacks were restricted to *L. michahellis* breeding area but randomly distributed within its limits.

In terms of *L. michahellis* attack probability (Binomial family models), models predicted an independent distribution of the probability of an attack taking place (dives and fecal attacks) in relation with the number of seagull chicks (χ^2 =0, d.f =1, P=0.999). The same pattern was observed for dives and fecal attacks separately (χ^2 =2.307, d.f =1, P=0.128 and χ^2 =0.17, d.f =1, P=0.676 respectively).

4. Discussion

In this study, we showed how direct and indirect ecological interactions allow a critically endangered reptile to inhabit and maintain relatively stable populations in a suboptimal habitat. Through two different experiments, a dietary stable isotopes analysis, and an ecological behaviour study, we showed how the Tenerife speckled lizards interact with a yellow-legged seagull colony obtaining food, and protection against alien predator species. The stable isotope analysis showed that revealed that *G. intermedia* was mainly feeding in the area under the direct influence of seagulls' guano and taking advantage of the seagull-derived food resources that they provide directly or indirectly. Additionally, the behaviour experiment showed that, *L. michahellis* displayed an aggressive antipredator behaviour against predators within the limits of its breeding area with an increase on the frequency of aggressive behaviours correlated with the abundance of seagull's chicks.

The effects of seabird-derived resources on lizards' populations have been found to positively affect different parameters such as reproduction, body size, density and intraspecific antagonistic behaviour (Larsen, 1991; Pafilis et al., 2009, 2011; K. M. Richardson et al., 2019; Stadler et al., 2023). The patterns of stable nitrogen and carbon isotopes found on *G. intermedia* back those past results. Furthermore, the patterns of stable carbon and nitrogen found on the different trophic resources as well as on *G. intermedia,* showed us how the latter is using its environment (Cherel et al., 2000; Grundler et al., 2017; Han et al., 2021). The vegetation in areas under the direct influence of *L. michahellis* colony show a completely different isotopic signal. The high values of $\delta^{15}N$ of the plants located nearby or directly inside the colony show a clear influence of nitrogen derived from guano, which exhibited $\delta^{15}N$ values ~7‰ higher than those on the areas out of seagull colony (Figure 2.a-b, Table S.7). The vegetation from platforms were as expected showed the highest $\delta^{15}N$ values, even higher than those of the seagull chicks (mainly piscivores) and lizards. It is well known that seabird colonies, through the deposition of guano, rich in nitrogen and phosphorous compounds, induce profound changes in soil properties, including eutrophication, salinization, acidification and nutrient imbalances, modifying the plant communities that live there (Anderson & Polis, 1999; Ellis, 2005; Wainright et al., 1998; Wait et al., 2005). As consequence, plants enriched in $\delta^{15}N$ improve their quality and quantity and

significantly increase the $\delta^{15}N$ content in all components of the food webs of coastal and island ecosystems through multi-trophic cascading effects (Barrett et al., 2005; Fukami et al., 2006). In the case of *G. intermedia*, its diet seems to depend on approximately 25% of plants that live in the area influenced by the seagull colony. This limited contribution may be indicative of the sparse vegetation cover in these areas, where C_4/CAM plants such as *Portulaca* spp., *M. crystallynum* and *O. dillenii* predominate, along with C_3 plants such as *P. patelleris*, *L. itrincatum* and *K. neriifolia*. Many of these plants have an annual cycle, appearing only during the wet season. In contrast, the prickly pear *O. dillenii* (cactacea) produce ripe fruits nearly year-round, which are consumed by *G. intermedia* (pers. obs.). However, the distribution of *O. dillenii* on the Guaza cliff platforms is limited to a few locations. Conversely, the prickly pear is a non-native invasive species in the Canary Islands, leading to calls for its control and eradication in protected areas (García-Gallo et al., 2009). Given our findings, the eradication of this species in areas inhabited by the Tenerife speckled lizard could potentially jeopardize its survival. Therefore, we recommend considering the trophic and other ecological relationships that may exist between naturalized invasive plant species and threatened endemic species before implementing eradication plans.

The MixSIAR model suggests that the diet of *G. intermedia* is primarily comprised of food items derived from seagulls (~62%). These resources displayed $\delta^{13}C$ levels of approximately -18‰, which are markedly different from the $\delta^{13}C$ levels of C3 plants (~ -24‰) and C4/CAM plants (~ -14‰). These levels indicate that the carbon is sourced from marine and epipelagic food sources in the area, such as epipelagic fishes like *Trachurus picturatus* and *Scomber colias*, which have $\delta^{13}C$ values ranging between -19‰ and -20‰ (Romero et al., 2021). In a similar manner, the $\delta^{13}C$ values of *G. intermedia* (~ -20 ‰) are indicative of this marine epipelagic $\delta^{13}C$ signature. This suggests that most of the foraging activity of *G. intermedia* is taking place in the areas under the direct influence of the yellow-legged seagulls, despite the majority of speckled lizards being caught in ravines, where apparently vegetation cover and diversity can offer a more diverse range of foods resources. Seagull-derived resources may encompass a wide range of foods, including remains brought directly by the parents to the chicks (such as epipelagic fish and cephalopods), which may be regurgitated and left outside the nest, as well as remains of chick's or adult seagulls

carcasses, eggshells, seagull droppings and scavenger insects that feed on these remains and move out of the colony area dispersing this marine subsidy through the food chain. Future dietary studies with trophic markers such as those used here can provide greater resolution if more effort is made to sample all possible food sources used by this species throughout the year, obtain food consumption data derived from fecal analysis (priors), and generate species-specific or genus specific for *Gallotia* spp. TDFs to improve the fit of Bayesian isotopic mixing models.

We cannot discount the possibility that ravine areas serve as dispersal zones for juvenile lizards originating from the cliff. Ontogenetic habitats shift in lizards, where juveniles actively avoid the presence of adults due to competition, cannibalism or select other habitats according to its different life traits requirements have been documented in multiple lizard species (Delaney & Warner, 2017; Jenssen et al., 1989; McMann & Paterson, 2003; San-Jose et al., 2016; Stamps, 1983). In the context of *G. intermedia*, the forced removal of smaller individuals from cliff regions, where they benefit from protection and food supplied by gulls, could heighten their susceptibility to predation and diminish the fitness of these juveniles in ravines. This may ultimately lead to these ravines becoming sink sites for the population. Unfortunately, the low number of captures in our study does not allow us to investigate this question, so future research should be carried out in order to further investigate the possible habitat selection of *G. intermedia* by age and sex and the structuring of their populations.

Additionally, the results from the anti-predator behaviour experiment further back the potential positive interaction between the breeding colony of *L. michahellis* and *G. intermedia*. Although the probability of attack was randomly distributed withing the colony, the frequency of attacks was strongly correlated with the abundance of the seagull's chicks. Similar mobbing behaviours have been recorded in other seabirds' colonies (Clode et al., 2000; Larsen, 1991), and in some cases it has been reported that other species use this indirect protection on their advantage (Gameiro et al., 2022; Guidos et al., 2023). For example, Monk parakeets (*Myiopsitta monachus*), an IAS built their nest under those of white storks (*Ciconia ciconia*) to reduce predation risk which indirectly offers protection against predators (Hernández-Brito et al., 2020). The anti-predator behaviour of *L. michahellis* wasn't limited only to our cat decoy. During the months of fieldwork, we observed aggressive behaviour towards a wide range of

species, from kestrels to Eleonora falcons, egrets, and biologists (see Supplementary Materials Figure S3.a-c and Figures S4). Again, these types of interactions can help us explain why *G. intermedia*, a critically endangered island species, is able to maintain its populations in an area under feral cats' pressure and surrounded by huge touristic developments. The scale and reach of the effects of feral and domestic cats on the populations of *G. intermedia* are still unknown, however the limited number of studies on the subject have shown that the Tenerife speckled lizard is a relatively common prey for the cats that roams the coastal areas of Guaza Mountain (Albaladejo et al., 2015; Flores Ravelo & Rando Reyes, 2021). Although *L. michahellis* can help to mitigate the damage, at least during its breeding season, previous monitoring of *G. intermedia* populations on the protected area of Guaza Mountain have shown a potential population decline of more than 32% over the past 18 years (Albaladejo et al., 2015). Therefore, active conservation actions that take into consideration the complex interactions between the species are fundamental to push endangered species away from extinction.

Although we focused just on one species and a set of small populations, our findings are of interest for the conservation and study of other critically endangered lizards' species that share similar characteristics and requirements with *G. intermedia*. This is the case, for example, of *Gallotia bravoana* (Mateo et al., 2009a) and *Gallotia simonyi* (Mateo et al., 2009b). These endemism's of La Gomera and El Hierro islands, respectively, are also severely affected by feral cats (Afonso & Mateo, 2009; García Márquez et al., 1997) and their natural populations are restricted to isolated and harsh cliffs (Afonso & Mateo, 2009; Albaladejo et al., 2015; Rica, 1981) occupied by different breeding populations of seabirds (e.g., marsh and shearwaters, seagulls). Understanding which were the conditions that allowed this species to maintain wild populations in such harsh conditions, can aid the reintroduction success of future populations by not only looking for the right geographic conditions (isolation and difficult access to invasive species) but also at the adequate biotic interactions (Bogoni et al., 2019; Renzi et al., 2019). Furthermore, populations of the giant lizard of Gran Canaria (*Gallotia stehlini*) are being decimated by the introduction of the California Kingsnake (*Lampropeltis californiae*) (Piquet & López-Darias, 2021) and so it is likely that in the near future this species also will face the same challenges that the rest of endemic giant reptiles of the Canary Islands. In this context, refugees can be identified

and preserved in advance, and individuals can be introduced in these suboptimal habitats to preserve the genetic diversity of the species (Braidwood et al., 2018) while other conservation actions are undergoing to try to halt the expansion of the California Kingsnake to the rest of the island (Piquet et al., 2021).

At a global scale, habitat destruction and invasive alien species are one of the main causes of biodiversity loss (Dirzo et al., 2014; WWF, 2020). On this scenario we should expect more species to be pushed into suboptimal or relict habitats. As we have shown with this work, understanding which elements allow species to survive in such conditions is of paramount importance for the correct planning and management of conservation actions. Therefore, a focus on disentangled the hidden biotic and abiotic interactions that allow species to inhabit these suboptimal habitats can help us to better prioritize habitat conservation, reintroduction areas, and long-term species management. For this, it is fundamental that basic ecological studies are conducted alongside biodiversity conservation (Brussard, 1991; Fontúrbel et al., 2018).

Overall, we found that populations of a critically endangered species inhabiting a relict or suboptimal habitat rely on indirect ecological interactions to maintain their populations. Our results suggest not only an important interconnectivity between the yellow-legged seagull colony and the Tenerife Speckled lizard, but also that the actual realize niche of *G. intermedia* is defined by this interaction. Our observations further back the importance of basic ecological research to understand the mechanism behind the distribution and survival of endangered species, and the importance of this knowledge for the correct application of conservation actions. The results of this study can further help understand the patterns of distribution of other endemic and endangered reptiles of the Canary Islands as well as to design long-term conservation strategies. At a global, since we expect more species to be pushed to suboptimal habitats, we suggest that hidden biotic interactions as the one we describe here might be of paramount importance for the conservation of biodiversity and the planning of conservation actions.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank all the people who helped us during the fieldwork (M. Gil-Velasco, J. Marrero, N. Álvarez-Quintero). Also, to Drs. D. Hernández-Teixenor and Miquel Lombarte for their help with the SIA analyses. To Dr. A. Martín who provided the essential logistical support that made this research possible, as well as lending us the decoy cat "Fermín" for the experiments. We would also like to thank the staff of As. Tonina for the management of the project.

Funding

This research has been funded by The Mohamed bin Zayed Species Conservation Fund within the 2016 call, project number 152511662. A. E. has been funded by the Actions of the Ministry of Universities under the application 33.50.460A.752 and by the European Union Next Generation EU/PRTR through a Margarita Salas postdoctoral contract of the University of Vigo.

Author contributions

G. A. R.: conceptualization, formal analysis, investigation, methodology, software, writing the original draft, writing the review and editing A. E.: conceptualization, formal analysis, funding acquisition, investigation, methodology, project administration, writing the original draft, review and editing. A. V. P. C.: formal analysis, investigation.

Data Availability

Data will be made available on request.

Bibliography

Afonso, O. M., & Mateo, J. A. (2009). *Depredación por gatos de Lagartos Gigantes de La Gomera, Gallotia bravoana (Sauria; Lacertidae)*.

Albaladejo, G., Martín, A., Sämann, J., & Pfau, B. (2015). Gallotia intermedia Barbadillo, Lacomba, Pérez-Mellado, Sancho & López-Jurado, 1999 – Die Rieseneidechse von Teneriffa. *Die Eidechse*, *26*, 33–48.

Anderson, W. B., & Polis, G. A. (1999). Nutrient fluxes from water to land: Seabirds affect plant nutrient status on Gulf of California islands. *Oecologia*, *118*, 324–332.

Barret W.B. Anderson, D. A. Wait, L.L. Grismer, G.A. Polis & M.D. Rose, K., Barrett, K., Anderson, W. B., Wait, D. A., Grismer, L. L., Polis, G. A., & Rose, M. D. (2005). Marine subsidies alter the diet and abundance of insular and coastal lizard populations. *Oikos*, *109*(1), 145–153. https://doi.org/10.1111/j.0030-1299.2005.13728.x

Barrett, K., Anderson, W. B., Wait, D. A., Grismer, L. L., Polis, G. A., & Rose, M. D. (2005). Marine subsidies alter the diet and abundance of insular and coastal lizard populations. *Oikos*, *109*(1), 145–153. https://doi.org/10.1111/j.0030-1299.2005.13728.x

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

BOC, 2010. LEY 4/2010, de 4 de junio, del Catálogo Canario de Especies Protegidas. Boletín Oficial de Canarias, Nº 112, 15200-15225.

BOE, 2011. Real Decreto 139/2011, de 4 de febrero, para el desarrollo del Listado de Especies Silvestres en Régimen de Protección Especial y del Catálogo Español de Especies Amenazadas. Boletín Oficial del Estado, Nº 46, 20912-20951.

Bogoni, J. A., Navarro, A. B., Graipel, M. E., & Peroni, N. (2019). Modeling the frugivory of a plant with inconstant productivity and solid interaction with relictual vertebrate biota. *Ecological Modelling*, *408*, 108728. https://doi.org/10.1016/j.ecolmodel.2019.108728

Bonnaud, E., Medina, F. M., Vidal, E., Nogales, M., Tershy, B., Zavaleta, E., Donlan, C. J., Keitt, B., Le Corre, M., & Horwath, S. V. (2011). The diet of feral cats on islands: A review and a call for more studies. *Biological Invasions*, *13*(3), 581–603. https://doi.org/10.1007/s10530-010-9851-3

Boone, J. L., & Larue, E. A. (1999). Effects of marking Uta stansburiana (Sauria: Phrynosomatidae) with xylene-based paint. *Herpetological Review*, *30*(1), 33.

Braidwood, D. W., Taggart, M. A., Smith, M., & Andersen, R. (2018). Translocations, conservation, and climate change: Use of restoration sites as protorefuges and protorefugia. *Restoration Ecology*, *26*(1), 20–28. https://doi.org/10.1111/rec.12642

Brondizio, E. S., Settele, J., Diaz, S., & Ngo, H. T. (2019). *IPBES(2019): Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.*

Brown, J. L., & Carnaval, A. C. (2019). A tale of two niches: Methods, concepts, and evolution. *Frontiers of Biogeography*, *11*(4). https://doi.org/10.21425/F5FBG44158

Brussard, P. F. (1991). The role of ecology in biological conservation. *Ecological Applications*, *1*(1), 6–12.

Caut, S., Angulo, E., & Courchamp, F. (2008). Discrimination factors (Δ15N and Δ13C) in an omnivorous consumer: Effect of diet isotopic ratio. *Functional Ecology*, *22*(2), 255–263.

Cherel, Y., Hobson, K. A., & Weimerskirch, H. (2000). Using stable-isotope analysis of feathers to distinguish moulting and breeding origins of seabirds. *Oecologia*, *122*(2), 155–162. https://doi.org/10.1007/PL00008843

Clode, D., Birks, J. D. S., & Macdonald, D. W. (2000). The influence of risk and vulnerability on predator mobbing by terns (Sterna spp.) and gulls (Larus spp.). *Journal of Zoology*, *252*(1), 53–59. https://doi.org/10.1111/j.1469-7998.2000.tb00819.x

Davis, A. J., Lawton, J. H., Shorrocks, B., & Jenkinson, L. S. (1998). Individualistic species responses invalidate simple physiological models of community dynamics under global environmental change. *Journal of Animal Ecology*, *67*(4), 600–612. https://doi.org/10.1046/j.1365-2656.1998.00223.x

del Arco Aguilar, M. J., González-González, R., Garzón-Machado, V., & Pizarro-Hernández, B. (2010). Actual and potential natural vegetation on the Canary Islands

and its conservation status. *Biodiversity and Conservation*, *19*(11), 3089–3140. https://doi.org/10.1007/s10531-010-9881-2

Delaney, D. M., & Warner, D. A. (2017). Effects of age-and sex-specific density on behaviour and survival in a territorial lizard (Anolis sagrei). *Animal Behaviour*, *129*, 31–41.

Dirzo, R., Young, H. S., Galetti, M., Ceballos, G., Isaac, N. J. B., & Collen, B. (2014). Defaunation in the Anthropocene. *Science*, *345*(6195), 401–406. https://doi.org/10.1126/science.1251817

Ellis, J. C. (2005). Marine birds on land: A review of plant biomass, species richness, and community composition in seabird colonies. *Plant Ecology*, *181*, 227–241.

Elzinga, C. L., Salzer, D. W., Willoughby, J. W., & Gibbs, J. P. (2001). *Monitoring plant and animal populations: A handbook for field biologists*. John Wiley & Sons.

Erwin, R. M. (1988). Correlates of Nest-Defense Behavior of Common Terns (Correlaciones en la Defenza del Nido por Parte de Sterna hirundo). *Journal of Field Ornithology*, *59*(2), 135–142.

Fariña, B., & Martín, A. (2013). Ornitofagia de Gallotia intermedia en Guaza (Tenerife, Islas Canarias). *Boletín de Asociación Española de Herpetologia*, *24*(1), 43–45.

Flores Ravelo, A. J., & Rando Reyes, J. C. (2021). Trophic ecology of cats (Felis catus) in montaña de Guaza: Implications for the conservation of the critically endangered giant lizard of Tenerife (Gallotia intermedia). *Scientia Insularum. Revista de Ciencias Naturales En Islas*, *4*, 63–80. https://doi.org/10.25145/j.SI.2021.04.04

Fontúrbel, F. E., Lara, A., Lobos, D., & Little, C. (2018). The cascade impacts of climate change could threaten key ecological interactions. *Ecosphere*, *9*(12), e02485. https://doi.org/10.1002/ecs2.2485

Fox, J., & Weisberg, S. (2019). *An R Companion to Applied Regression*. Sage. https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Frixione, M. G., & Salvadeo, C. (2021). Drones, Gulls and Urbanity: Interaction between New Technologies and Human Subsidized Species in Coastal Areas. *Drones*, *5*(2), 30. https://doi.org/10.3390/drones5020030

Fukami, T., Wardle, D. A., Bellingham, P. J., Mulder, C. P. H., Towns, D. R., Yeates, G. W., Bonner, K. I., Durrett, M. S., Grant-Hoffman, M. N., & Williamson, W. M. (2006). Above- and below-ground impacts of introduced predators in seabird-dominated island ecosystems. *Ecology Letters*, *9*(12), 1299–1307. https://doi.org/10.1111/j.1461-0248.2006.00983.x

Gameiro, J., Franco, A. M. A., Catry, T., Palmeirim, J. M., & Catry, I. (2022). Antipredator benefits of heterospecific colonial breeding for a predominantly solitary bird. *Animal Behaviour, 189,* 101–111. https://doi.org/10.1016/j.anbehav.2022.05.005

García Gallo, A., Rodríguez Delgado, O. y GesPlan S.A.U., 2009. Opuntia dillenii (Ker-Gawl.) Haw. En: Ojeda Land, E. y Rodríguez Luengo, J. L. (Eds.), 2022. Compendio de fichas de la Base de especies introducidas en Canarias (2008-2011). Viceconsejería de Lucha contra el Cambio Climático. Dirección General de Lucha contra el Cambio Climático y Medio Ambiente. Gobierno de Canarias. 599-603.

García Márquez, M., López-Jurado, L. F., & Mateo, J. A. (1997). Predación de Gallotia simonyi por gatos cimarrones. *Boletín de la Asociación Herpetológica Española [ISSN 1130-6939], v. 8, p. 20-22.* https://accedacris.ulpgc.es/jspui/handle/10553/18285

García-Muñoz, E., Ceacero, F., Pedrajas, L., Kaliontzopoulou, A., & Carretero, M. Á. (2011). Tail tip removal for tissue sampling has no short-term effects on microhabitat selection by Podarcis bocagei, but induced autotomy does. *Acta Herpetologica*, *6*(2), 223–227.

Gaston, K. J., & Gaston, K. J. (2003). *The Structure and Dynamics of Geographic Ranges.* Oxford University Press.

Gaynor, K. M., Hojnowski, C. E., Carter, N. H., & Brashares, J. S. (2018). The influence of human disturbance on wildlife nocturnality. *Science, 360*(6394), 1232–1235. https://doi.org/10.1126/science.aar7121

Ginovés, J., A. Betoret & A. Martín (2005): Estudio de la población del lagarto gigante de Tenerife (Gallotia intermedia) en el Espacio Natural Protegido de la montana de Guaza. – Departamento de Biología Animal (Zoologia) Universidad de La Laguna. Unpublished report

Godsoe, W., Franklin, J., & Blanchet, F. G. (2017). Effects of biotic interactions on modeled species' distribution can be masked by environmental gradients. *Ecology and Evolution*, *7*(2), 654–664. https://doi.org/10.1002/ece3.2657

Gonzalez, E. G., Cerón-Souza, I., Mateo, J. A., & Zardoya, R. (2014). Island survivors: Population genetic structure and demography of the critically endangered giant lizard of La Gomera, Gallotia bravoana. *BMC Genetics*, *15*(1), 121. https://doi.org/10.1186/s12863-014-0121-8

Grundler, M. R., Pianka, E. R., Pelegrin, N., Cowan, M. A., & Rabosky, D. L. (2017). Stable isotope ecology of a hyper-diverse community of scincid lizards from arid Australia. *PLOS ONE*, *12*(2), e0172879. https://doi.org/10.1371/journal.pone.0172879

Guidos, S., van Dijk, J., Systad, G., & Landa, A. (2023). Colony-nesting gulls restrict activity levels of a native top carnivore during the breeding season. *Remote Sensing in Ecology and Conservation*, *9*(4), 527–539. https://doi.org/10.1002/rse2.326

Guisan, A., & Thuiller, W. (2005). Predicting species distribution: Offering more than simple habitat models. *Ecology Letters*, *8*(9), 993–1009. https://doi.org/10.1111/j.1461-0248.2005.00792.x

Han, Z., Wang, H.-T., Kardynal, K. J., Hobson, K. A., Shang, W.-P., Zhang, L.-S., Jiang, Y., & Jiguet, F. (2021). Stable isotopes (δ2H) in feathers identify non-breeding origins of the endangered Jankowski's Bunting. *Journal of Ornithology*, *162*(4), 987–995. https://doi.org/10.1007/s10336-021-01897-7

Hernández, E., Nogales, M. ;, & Martín, A. (2000). Discovery of a New Lizard in the Canary Islands, with a Multivariate Analysis of Gallotia (Reptilia: Lacertidae). *Herpetologica, 56*(1), 63–76.

Hernández-Brito, D., Blanco, G., Tella, J. L., & Carrete, M. (2020). A protective nesting association with native species counteracts biotic resistance for the spread of an invasive parakeet from urban into rural habitats. *Frontiers in Zoology, 17*(1), 13. https://doi.org/10.1186/s12983-020-00360-2

Jenssen, T. A., Marcellini, D. L., Buhlmann, K. A., & Goforth, P. H. (1989). Differential infanticide by adult curly-tailed lizards, Leiocephalus schreibersi. *Animal Behaviour*, *38*(6), 1054–1061.

Kenny, H. V., Wright, A. N., Piovia-Scott, J., Yang, L. H., Spiller, D. A., & Schoener, T. W. (2017). Marine subsidies change short-term foraging activity and habitat utilization of terrestrial lizards. *Ecology and Evolution*, *7*(24), 10701–10709. https://doi.org/10.1002/ece3.3560

Keppel, G., Mokany, K., Wardell-Johnson, G. W., Phillips, B. L., Welbergen, J. A., & Reside, A. E. (2015). The capacity of refugia for conservation planning under climate change. *Frontiers in Ecology and the Environment*, *13*(2), 106–112. https://doi.org/10.1890/140055

Keppel, G., Van Niel, K. P., Wardell-Johnson, G. W., Yates, C. J., Byrne, M., Mucina, L., Schut, A. G. T., Hopper, S. D., & Franklin, S. E. (2012). Refugia: Identifying and understanding safe havens for biodiversity under climate change. *Global Ecology and Biogeography*, *21*(4), 393–404. https://doi.org/10.1111/j.1466-8238.2011.00686.x

Kerley, G. I. H., Kowalczyk, R., & Cromsigt, J. P. G. M. (2012). Conservation implications of the refugee species concept and the European bison: King of the forest or refugee in a marginal habitat? *Ecography*, *35*(6), 519–529. https://doi.org/10.1111/j.1600-0587.2011.07146.x

Kerley, G. I. H., te Beest, M., Cromsigt, J. P. G. M., Pauly, D., & Shultz, S. (2020). The Protected Area Paradox and refugee species: The giant panda and baselines shifted towards conserving species in marginal habitats. *Conservation Science and Practice*, *2*(6), e203. https://doi.org/10.1111/csp2.203

Larsen, T. (1991). Anti-predator behaviour and mating systems in waders: Aggressive nest defence selects for monogamy. *Animal Behaviour*, *41*(6), 1057–1062. https://doi.org/10.1016/S0003-3472(05)80643-8

Lea, J. M., Kerley, G. I., Hrabar, H., Barry, T. J., & Shultz, S. (2016). Recognition and management of ecological refugees: A case study of the Cape mountain zebra. *Biological Conservation*, *203*, 207–215. https://doi.org/10.1016/j.biocon.2016.09.017

Lin, Z.-H., & Ji, X. (2005). Partial tail loss has no severe effects on energy stores and locomotor performance in a lacertid lizard, Takydromus septentrionalis. *Journal of Comparative Physiology B*, *175*(8), 567–573. https://doi.org/10.1007/s00360-005-0017-z

Louthan, A. M., Doak, D. F., & Angert, A. L. (2015). Where and When do Species Interactions Set Range Limits? *Trends in Ecology & Evolution*, *30*(12), 780–792. https://doi.org/10.1016/j.tree.2015.09.011

Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P., & Makowski, D. (2021). performance: An R Package for Assessment, Comparison and Testing of Statistical Models. *Journal of Open Source Software*, *6*(60), 3139. https://doi.org/10.21105/joss.03139

Machado, A. (1985). New data on the Hierro giant lizard and the lizard of Salmor (Canary Islands). *Bonn. Zool. Beitr.*, *36*, 429–470.

Mateo, J. A. (2006). *El lagarto gigante de La Gomera*. Cabildo Insular de La Gomera.

Mateo, J. A. (2009). Lagarto gigante de La Palma – Gallotia auaritae. En: Enciclopedia Virtual de los Vertebrados Españoles. Carrascal, L. M., Salvador, A. (Eds.). Museo Nacional de Ciencias Naturales, Madrid. http://www.vertebradosibericos.org/

Mateo, J.A., Miras, Pérez-Mellado, V., Martínez-Solano, I. (2009a). *Gallotia bravoana*. The IUCN Red List of Threatened Species 2009: e.T61502A12493034. http://dx.doi.org/10.2305/IUCN.UK.2009.RLTS.T61502A12493034.en

Mateo, J.A., Miras, Pérez-Mellado, V., Martínez-Solano, I. (2009b). *Gallotia simonyi*. The IUCN Red List of Threatened Species 2009: e.T8881A12935900. http://dx.doi.org/10.2305/IUCN.UK.2009.RLTS.T8881A12935900.en

McCutchan Jr, J. H., Lewis Jr, W. M., Kendall, C., & McGrath, C. C. (2003). Variation in trophic shift for stable isotope ratios of carbon, nitrogen, and sulfur. *Oikos*, *102*(2), 378–390.

McMann, S., & Paterson, A. V. (2003). The relationship between location and displays in a territorial lizard. *Journal of Herpetology*, 414–416.

Medina, F. M., & Nogales, M. (2009). A review on the impacts of feral cats (Felis silvestriscatus) in the Canary Islands: Implications for the conservation of its endangered fauna. *Biodiversity and Conservation*, *18*(4), 829–846. https://doi.org/10.1007/s10531-008-9503-4

Minias, P., Gach, K., Włodarczyk, R., Bartos, M., Drzewińska-Chańko, J., Rembowski, M., Jakubas, D., & Janiszewski, T. (2020). Colony size as a predictor of breeding behaviour in a common waterbird. *PLOS ONE*, *15*(11), e0241602. https://doi.org/10.1371/journal.pone.0241602

Momberg, M., Ryan, P. G., Hedding, D. W., Schoombie, J., Goddard, K. A., Craig, K. J., & Le Roux, P. C. (2023). Factors determining nest-site selection of surface-nesting seabirds: A case study on the world's largest pelagic bird, the Wandering Albatross (Diomedea exulans). *Ibis*, *165*(1), 190–203. https://doi.org/10.1111/ibi.13111

Murray, I. W., Lease, H. M., Hetem, R. S., Mitchell, D., Fuller, A., & Woodborne, S. (2016). Stable isotope analysis of diet confirms niche separation of two sympatric species of Namib Desert lizard. *Integrative Zoology*, *11*(1), 60–75.

Nagelkerke, N. J. & others. (1991). A note on a general definition of the coefficient of determination. *Biometrika*, *78*(3), 691–692.

Nogales, M., Rando, J. C., Valido, A., & Martín, A. (2001). Discovery of a Living Giant Lizard, Genus Gallotia (Reptilia: Lacertidae), from la Gomera, Canary Islands. *Herpetologica*, *57*(2), 169–179.

Pafilis, P., Foufopoulos, J., Sagonas, K., Runemark, A., Svensson, E., & Valakos, E. D. (2011). Reproductive biology of insular reptiles: Marine subsidies modulate expression of the "island syndrome." *Copeia*, *2011*(4), 545–552.

Pafilis, P., Meiri, S., Foufopoulos, J., & Valakos, E. (2009). Intraspecific competition and high food availability are associated with insular gigantism in a lizard. *Naturwissenschaften*, *96*(9), 1107–1113.

Palacios-García, S., Cruzado-Caballero, P., Casillas, R., & Ruiz, C. C. (2021). Quaternary biodiversity of the giant fossil endemic lizards from the island of El Hierro (Canary Islands, Spain). *Quaternary Science Reviews*, *262*, 106961. https://doi.org/10.1016/j.quascirev.2021.106961

Pavey, C. R., Addison, J., Brandle, R., Dickman, C. R., McDonald, P. J., Moseby, K. E., & Young, L. I. (2017). The role of refuges in the persistence of Australian dryland mammals. *Biological Reviews*, *92*(2), 647–664. https://doi.org/10.1111/brv.12247

Pérez-Mellado, V., Romero-Beviá, M., Ortega, F., Martín-García, S., Perera, A., López-Vicente, M., & Galache, C. (1999). El uso de los recursos tróficos en Gallotia simonyi (Sauria, Lacertidae) de la isla de El Hierro (Islas Canarias). *Monogr. Herpetol*, *4*, 63–83.

Peterson, A. T., Soberón, J., Pearson, R. G., Anderson, R. P., Martínez-Meyer, E., Nakamura, M., & Araújo, M. B. (2011). *Ecological Niches and Geographic Distributions (MPB-49)*. Princeton University Press. http://www.jstor.org/stable/j.ctt7stnh

Piquet, J. C., & López-Darias, M. (2021). Invasive snake causes massive reduction of all endemic herpetofauna on Gran Canaria. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1964), 20211939. https://doi.org/10.1098/rspb.2021.1939

Piquet, J. C., Warren, D. L., Saavedra Bolaños, J. F., Sánchez Rivero, J. M., Gallo-Barneto, R., Cabrera-Pérez, M. Á., Fisher, R. N., Fisher, S. R., Rochester, C. J., Hinds, B., Nogales, M., & López-Darias, M. (2021). Could climate change benefit invasive snakes? Modelling the potential distribution of the California Kingsnake in the Canary Islands. *Journal of Environmental Management*, *294*, 112917. https://doi.org/10.1016/j.jenvman.2021.112917

Quinn, J. L., & Ueta, M. (2008). Protective nesting associations in birds. *Ibis*, *150*(s1), 146–167. https://doi.org/10.1111/j.1474-919X.2008.00823.x

R Core Team. (2021). R: A Language and Environment for Statistical Computing. *R Foundation for Statistical Computing*. https://www.r-project.org/

Ramos, R., Ramírez, F., Sanpera, C., Jover, L., & Ruiz, X. (2009). Feeding ecology of yellow-legged gulls Larus michahellis in the western Mediterranean: A comparative assessment using conventional and isotopic methods. *Marine Ecology Progress Series*, *377*, 289–297.

Rando, J. C., Valido, A., Nogales, M., & Martín, A. (2000). Lagarto Gigante de La Gomera: Un fósil que vuelva a la vida. *Quercus*, *171*(January), 10–16.

Rando, J.-C., Betoret, A., Martin, A., & Barone, R. (2004). Discovery of a new population of the spotted lizard of the Canary Islands. *Oryx*, *38*(2), 131. https://doi.org/10.1017/S0030605304000225

Renzi, J. J., He, Q., & Silliman, B. R. (2019). Harnessing Positive Species Interactions to Enhance Coastal Wetland Restoration. *Frontiers in Ecology and Evolution*, *7*. https://www.frontiersin.org/articles/10.3389/fevo.2019.00131

Rica, J. P. M. (1981). Primeros datos sobre la población de lagarto negro (Gallotia simonyi simonyi STEIND.) de la Isla de Hierro1. *Amphibia-Reptilia*, *2*(4), 369–380. https://doi.org/10.1163/156853882X00293

Richardson, D. S., & Bolen, G. M. (1999). A nesting association between semi-colonial Bullock's orioles and yellow-billed magpies: Evidence for the predator protection hypothesis. *Behavioral Ecology and Sociobiology*, *46*(6), 373–380. https://doi.org/10.1007/s002650050632

Richardson, K. M., Iverson, J. B., & Kurle, C. M. (2019). Marine subsidies likely cause gigantism of iguanas in the Bahamas. *Oecologia*, *189*(4), 1005–1015. https://doi.org/10.1007/s00442-019-04366-4

Robertson, B. A., & Hutto, R. L. (2006). A framework for understanding ecological traps and an evaluation of existing evidence. *Ecology*, *87*(5), 1075–1085. https://doi.org/10.1890/0012-9658(2006)87[1075:affuet]2.0.co;2

Romero, J., Vieira, C., Garrido, S., Hermida, M., Catry, P., Faria, G., & Granadeiro, J. P. (2021). Diet and trophic position of two mackerel species in the archipelago of Madeira, Portugal. *Journal of Fish Biology*, *99*(3), 831–843. https://doi.org/10.1111/jfb.14766

San-Jose, L. M., Peñalver-Alcázar, M., Huyghe, K., Breedveld, M. C., & Fitze, P. S. (2016). Inter-class competition in stage-structured populations: Effects of adult density on life-history traits of adult and juvenile common lizards. *Oecologia*, *182*(4), 1063–1074.

Scheele, B. C., Foster, C. N., Banks, S. C., & Lindenmayer, D. B. (2017). Niche Contractions in Declining Species: Mechanisms and Consequences. *Trends in Ecology & Evolution*, *32*(5), 346–355. https://doi.org/10.1016/j.tree.2017.02.013

Sedell, J. R., Reeves, G. H., Hauer, F. R., Stanford, J. A., & Hawkins, C. P. (1990). Role of refugia in recovery from disturbances: Modern fragmented and disconnected river systems. *Environmental Management*, *14*(5), 711–724. https://doi.org/10.1007/BF02394720

Selwood, K. E., & Zimmer, H. C. (2020). Refuges for biodiversity conservation: A review of the evidence. *Biological Conservation*, *245*, 108502. https://doi.org/10.1016/j.biocon.2020.108502

Siverio, F., & Felipe, P. (2009). Comensalismo entre lagarto gigante de El Hierro (Gallotia simonyi) y gaviota patiamarilla (Larus michahellis) en el roque Chico de Salmor. *Bol. Asociación Herpetológica Española*, *20*, 40–44.

Soberón, J. (2007). Grinnellian and Eltonian niches and geographic distributions of species. *Ecology Letters*, *10*(12), 1115–1123. https://doi.org/10.1111/j.1461-0248.2007.01107.x

Spiller, D. A., Piovia-Scott, J., Wright, A. N., Yang, L. H., Takimoto, G., Schoener, T. W., & Iwata, T. (2010). Marine subsidies have multiple effects on coastal food webs. *Ecology*, *91*(5), 1424–1434. https://doi.org/10.1890/09-0715.1

Stadler, S. R., Brock, K. M., Bednekoff, P. A., & Foufopoulos, J. (2023). More and bigger lizards reside on islands with more resources. *Journal of Zoology*, *319*(3), 163–174. https://doi.org/10.1111/jzo.13036

Stamps, J. A. (1983). The relationship between ontogenetic habitat shifts, competition and predator avoidance in a juvenile lizard (Anolis aeneus). *Behavioral Ecology and Sociobiology*, *12*, 19–33.

Stock, B. C., Jackson, A. L., Ward, E. J., Parnell, A. C., Phillips, D. L., & Semmens, B. X. (2018). Analyzing mixing systems using a new generation of Bayesian tracer mixing models. *PeerJ*, *6*, e5096. https://doi.org/10.7717/peerj.5096

Wainright, S., Haney, J., Kerr, C., Golovkin, A., & Flint, M. (1998). Utilization of nitrogen derived from seabird guano by terrestrial and marine plants at St. Paul, Pribilof Islands, Bering Sea, Alaska. *Marine Biology*, *131*, 63–71.

Wait, D., Aubrey, D., & Anderson, W. (2005). Seabird guano influences on desert islands: Soil chemistry and herbaceous species richness and productivity. *Journal of Arid Environments*, *60*(4), 681–695.

Wiens, J. J. (2011). The niche, biogeography and species interactions. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *366*(1576), 2336–2350. https://doi.org/10.1098/rstb.2011.0059

Wilson, M. F. J., O'Connell, B., Brown, C., Guinan, J. C., & Grehan, A. J. (2007). Multiscale Terrain Analysis of Multibeam Bathymetry Data for Habitat Mapping on the Continental Slope. *Marine Geodesy*, *30*(1–2), 3–35. https://doi.org/10.1080/01490410701295962

Wisz, M. S., Pottier, J., Kissling, W. D., Pellissier, L., Lenoir, J., Damgaard, C. F., Dormann, C. F., Forchhammer, M. C., Grytnes, J.-A., Guisan, A., Heikkinen, R. K., Høye, T. T., Kühn, I., Luoto, M., Maiorano, L., Nilsson, M.-C., Normand, S., Öckinger,

E., Schmidt, N. M., … Svenning, J.-C. (2013). The role of biotic interactions in shaping distributions and realised assemblages of species: Implications for species distribution modelling. *Biological Reviews*, *88*(1), 15–30. https://doi.org/10.1111/j.1469-185X.2012.00235.x

WWF. (2020). *Living Planet Report 2020 Bending the curve of biodiversity loss* (R. E. A. Almond, M. Grooten, & T. Petersen, Eds.). WWF.

Figure 1. Location and distribution of *Gallotia intermedia* in the Natural Reserve of Guaza Mountain. The red-shaded area describes the distribution of the breeding colony of *Larus michahellis* estimated using on-site observations and high-definition satellite multispectral images (see Supplementary materials S2 for more details). Gridded area marks the described distribution of *G. intermedia* according to Ginoves et al., (2005). Yellow and purple shaded areas mark the sampling localities for isotopes as well as the area used to carried out the anti-predator behaviour experiments. **Panel 1** describes in detail the different types of behaviour considered as aggressions (**a**=dives and **b**= fecal attacks) and the distances from the decoy at which seagulls chicks were counted (**c**).

Figure 2. Values of δ¹⁵N and δ¹³C for each species (**a**) and habitat sampled (**b-c**) during the study of the diet of *Gallotia intermedia*. Points on panel a represent the raw values obtained from the tissue

samples of *G. intermedia* and the rest of potential trophic resources. Species are differentiated by colours whereas the distinct geometrical shapes represent the different foraging groups. The mean ± the standard deviation $\delta^{15}N$ and $\delta^{13}C$ calculated by the MixSiar analysis taking into consideration the Trophic Discrimination Factors (oversized grey shapes and lines of each foraging group) are presented along with the raw $\delta^{15}N$ and $\delta^{13}C$ for the different species. Panel b and c represents the total values of $\delta^{15}N$ and $\delta^{13}C$ for the 4 types of Foraging groups included in the analysis. Lower case letters along the boxplot represent pairwise statistically significant groups.

Figure 3. Results of the anti-predator behaviour analysis. Partial responses of the Poisson generalized linear models describing the relationship between number of seagull chicks (**a**) and standardize topographic ruggedness (**b**) with the total number of recorded attacks (Dives + Fecal attacks). Data points are presented along with predicted responses and 95% confidence intervals. Spatial predictions of the number of *Larus michahellis* attacks (**c**), dotted red lines represent the approximate distribution of *G. intermedia* according to Ginoves et al., (2005).

Figure 4. Results of the anti-predator behaviour analysis. Partial responses of the Poisson generalized linear models describing the relationship between number of seagull chicks (**a**) and standardize topographic ruggedness (**b**) with the number of recorded dives attacks. Data points are presented along with predicted responses and 95% confidence intervals. Spatial predictions of the number of *Larus michahellis* attacks (**c**), dotted red lines represent the approximate distribution of G. intermedia according to Ginoves et al., (2005).

Table 1: Mean (± standard deviation) values ($\delta^{15}N$ and $\delta^{13}C$) of foraging item groups used in the mixing model MixSIAR to calculate diet contributions of *Gallotia intermedia*. n: number of samples analysed. Group 1: *E. balsamifera*, *F. cretica*, *K. neriifolia*, *L. intricatum*, *N. glauca*, *P. patellaris*. Group 2: *M. crystallinum*, *O. dillenii*, *Portulaca* spp. Group 3: *Argyranthemum* spp., *E. balsamifera*, *F. ericifolia*, *K. neriifolia*, *P. pendula*. Group 4: *L. michaellis* chicks' values after TDF subtraction.

Forage item group	Mean ± $\delta^{13}C$‰	Mean ± $\delta^{15}N$‰	n
Group 1: Platform C_3	-24.06 ± 2.51	15.82 ± 1.75	36
Group 2: Platform C_4/CAM	-14.46 ± 1.49	16.07 ± 3.29	13
Group 3: Plateu/Ravine C_3	-23.71 ± 2.27	8.57 ± 3.01	30
Group 4: Seabirds-derived resources	-19.5 ± 0.37	10.5 ± 0.45	10

Table 2: Proportional contribution of each forage item group to the diet of *Gallotia intermedia* estimated using MixSIAR models. Mean ± SD, 50% quantiles (median) and 95% Bayesian credible intervals (CI).

Forage item group	Mean ± SD	50%	95% CI
Group 1: Platform C_3	15 ± 0.42	15.2	5.8-22.7
Group 2: Platform C_4/CAM	10.9 ± 6.6	9.6	1.6-27.6
Group 3: Plateu/Ravine C_3	12 ± 10.1	9.2	0.06-38.7
Group 4: Seabirds-derived resources	62.2 ± 14.3	65.6	24.8-80.5

Supplementary materials

S1. Characterization of the vegetation in the habitat of *G. intermedia*

The vegetation study was conducted during two sampling sessions at the end of the dry season in early March 2016. Stratified random sampling of vegetation was carry out for each of the considered topological features found within the potential distribution area of *G. Intermedia* (Fig 1 main text); cliff platforms; ravines; and plateau. We carried out this sampling by placing five-line intercept transects of five meters long each. The initial position as well as the direction of the transects was randomly selected by throwing a 30 cm wooden ruler into the air. The landing point was set as the starting point of the transect whereas the orientation of the ruler marked the direction of the transect. Vegetation cover was measured along each line intercept transect by noting the point along the tape where the canopy of each plant species begins and the point at which it ends. The percentage of cover for each plant species in the line transect was then calculated as the proportion of transect covered by each plant species (Elzinga et al., 2001).

To characterize and detect potential differences in plant communities between different topological features, we applied a multivariate analysis approach. Species cover data was log (x+1) transformed and used to construct a similarity matrix using the Bray Curtis dissimilarity index. The resulting matrix was then employed to perform a cluster and non-metric multidimensional scaling (nMDS) analyses. Variation in plant community composition between topological features was subsequently tested using a one-way analysis of similarities (ANOSIM). Similarities between plant communities were measured using the ANOSIM R statistic, calculated using 999 Monte Carlo permutations. The R value returned by this test represents the degree of dissimilarity between the plant communities found in the topographic features aforementioned. ANOSIM R values close to "1" suggests dissimilarity between groups while an R value close to "0" suggests an even distribution of high and low ranks within and between groups (Clarke, 1993). In addition, to explore the relative contribution of plant species to the characterization of communities, we applied a similarity percentage analysis (SIMPER). This analysis test which species contribute to the difference between communities and how much of that difference can be explained by each species. All

these analyses were carried out using the R package *Vegan* version 2.6-4 (Oksanen et al., 2022).

Results:

The results from the nMDS, as well as the hierarchical clustering of the Bray-Curtis dissimilarity matrix, suggest that the topography of the environment is a key element in the delimitation and configuration of plant communities along the distribution area of *G. intermedia* in the Natural Reserve of Guaza Mountain (Figure S1 and Table S1-4). This was further confirmed by the results to the ANOSIM analysis which confirm a strong difference between plant communities cording to the topographic features (R-value=0.87; p-value=0.001; with 999 permutations). The SIMPER analysis showed that *Patellifolia patellaris*, *Euphorbia balsamifera*, and *Argyranthemum foeniculaceum* where the main contributors to differentiate the platform from the plateaus and ravines plant communities (Table S1). In the case of plateaus and ravines, *Arguranthemum foeniculaceum*, *Neochamaelea pulvurulenta*, and *Plocama pendula* were the species contributing to the differentiation between those plant communities (Table S1).

Table S1. Aggregated abundances for the plant species detected during the stratified sampling of the 4 different localities. The most abundant species for each locality are marked in bold, the total vegetation cover for each locality are showed at the bottom of the table.

	Ravine	Plateau	Platforms	
Species	**BNAME**	**CNAME**	**NAME**	**NICOTIANA**
Aizoon canariense	0	0	0.0056	0
Argyranthemum foeniculaceum	0.186	0.0032	0	0.012
Astydamia latifolia	0	0	0	0.024
Aeonium sp	0.008	0	0	0
Cenchrus cilliaris	0.04	0	0.02	0.024
Euphorbia balsamifera	0.184	**0.3288**	0.174	0.084
Fagonia cretica	0	0.004	0.01	0
Frankenia ericifolia	0	0.0908	0	0
Kleinia nerifolia	0.044	0.06	0.022	0
Limonium pectinatum	0.008	0.144	0	0
Neochamaelea pulvurulenta	0.112	0	0	0
Opuntia dillenii	0.0528	0.096	0	0
Patellifolia patellaris	0	0	**0.2148**	**0.586**
Plocama pendula	**0.2**	0	0	0
Astydamia latifolia	0	0	0.0092	0
Zygophyllum fontanesii	0	0	0.0572	0
Reseda scoparia	0	0.02	0	0
Schizogyne sericea	0.008	0.056	0	0
Volutaria canariensis	0.0016	0	0	0
Total vegetal cover	0.8444	0.8028	0.5128	0.73

Table S2. ANOSIM results for the plant communities inhabiting **platforms** and **plateaus** habitats along the distribution area of *G. intermedia*. The average contribution of species to the differentiation between plateaus and platforms plant communities (**Average**), is presented along with the standard deviation of these contributions (**sd**); the relative abundance of species (**Abund**); the cumulative contribution of species (**c.sum**); and the p-values (*p*) of the ANOSIM permutations, in this case 999. Species with no abundance or contribution to the communities have been removed from the table.

Species	Average	sd	Abund	c.sum	*p*
Patellifolia patellaris	0.183	0.052	5.008	0.209	0.001
Euphorbia balsamífera	0.128	0.094	1.949	0.355	0.017
Limonium pectinatum	0.101	0.087	0.000	0.471	0.004
Frankenia ericifolia	0.092	0.081	0.000	0.576	0.001
Opuntia dillenii	0.068	0.086	0.000	0.654	0.041
Schizogyne sericea	0.055	0.071	0.000	0.717	0.011
Reseda scoparia	0.049	0.062	0.000	0.772	0.001
Mesembryanthemum nodiflorum	0.042	0.057	1.288	0.820	0.496
Kleinia nerifolia	0.040	0.069	0.403	0.866	0.421
Argyranthemum foeniculaceum	0.027	0.045	0.343	0.896	1.000
Cenchrus cilliaris	0.025	0.052	0.804	0.926	0.895
Fagonia cretica	0.025	0.042	0.326	0.954	0.187
Astydamia latifolia	0.021	0.044	0.676	0.979	0.576
Zygophyllum fontanesii	0.011	0.033	0.318	0.991	0.677
Aizoon canariense	0.008	0.024	0.271	1.000	0.702

Table S3. ANOSIM results for the plant communities inhabiting **platforms** and **ravines** habitats along the distribution area of *G. intermedia*. The average contribution of species to the differentiation between plateaus and platforms plant communities (**Average**), is presented along with the standard deviation of these contributions (**sd**); the relative abundance of species (**Abund**); the cumulative contribution of species (**c.sum**); and the p-values (*p*) of the ANOSIM permutations, in this case 999. Species with no abundance or contribution to the communities have been removed from the table.

Species	Average	sd	Abund	c.sum	p
Patellifolia patellaris	0.172	0.057	5.008	0.190	0.001
Argyranthemum foeniculaceum	0.137	0.067	0.343	0.341	0.001
Neochamaelea pulvurulenta	0.095	0.059	0.000	0.447	0.001
Plocama pendula	0.092	0.078	0.000	0.549	0.001
Euphorbia balsamífera	0.088	0.079	1.949	0.647	0.873
Opuntia dillenii	0.061	0.064	0.000	0.714	0.193
Cenchrus cilliaris	0.054	0.085	0.804	0.774	0.180
Mesembryanthemum nodiflorum	0.039	0.055	1.288	0.817	0.581
Kleinia nerifolia	0.035	0.059	0.403	0.856	0.604
Limonium pectinatum	0.030	0.038	0.000	0.889	0.931
Astydamia latifolia	0.020	0.042	0.676	0.912	0.635
Aeonium sp	0.020	0.041	0.000	0.934	0.002
Schizogyne sericea	0.019	0.038	0.000	0.954	0.884
Volutaria canariensis	0.015	0.030	0.000	0.970	0.002
Zygophyllum fontanesii	0.010	0.031	0.318	0.982	0.734
Fagonia cretica	0.009	0.028	0.326	0.992	0.829
Aizoon canariense	0.008	0.023	0.271	1.000	0.737

Table S4. ANOSIM results for the plant communities inhabiting **plateaus** and **ravines** habitats along the distribution area of *G. intermedia*. The average contribution of species to the differentiation between plateaus and platforms plant communities (**Average**), is presented along with the standard deviation of these contributions (**sd**); the relative abundance of species (**Abund**); the cumulative contribution of species (**c.sum**); and the p-values (*p*) of the ANOSIM permutations, in this case 999. Species with no abundance or contribution to the communities have been removed from the table.

Species	Average	sd	Abund	c.sum	p
Argyranthemum foeniculaceum	0.108	0.042	0.439	0.145	0.011
Neochamaelea pulvurulenta	0.080	0.048	0.000	0.252	0.012
Plocama pendula	0.077	0.065	0.000	0.356	0.019
Euphorbia balsamífera	0.074	0.071	5.042	0.456	0.953
Limonium pectinatum	0.074	0.055	2.871	0.555	0.088
Frankenia ericifolia	0.071	0.062	2.530	0.650	0.019
Opuntia dillenii	0.063	0.058	1.918	0.735	0.266
Schizogyne sericea	0.046	0.052	1.568	0.796	0.169
Kleinia nerifolia	0.040	0.060	1.003	0.850	0.496
Reseda scoparia	0.037	0.047	1.296	0.900	0.133
Cenchrus cilliaris	0.032	0.066	0.000	0.944	0.649
Aeonium sp	0.017	0.034	0.000	0.966	0.484
Fagonia cretica	0.014	0.029	0.480	0.985	0.777
Volutaria canariensis	0.011	0.023	0.000	1.000	0.430

Figure S1. Non-metric multidimensional scaling (NMDS) of plant community composition for the different topographic features described within the distribution area of G. intermedia in the natural area of Guaza Mountain (**a**) and hierarchical cluster of the vegetation communities (**b**). For the NMDS (a), each colour-filled point represents the random stratified sampling vegetation units, whereas the hollow circles represent the different species. Ellipses represent ordination 95% confidence intervals of the standard deviations. For the dendrogram (**b**), each leaf of the dendrogram represent a monitoring unit of the stratified random sampling of vegetation, or a row of the Bray Curtis dissimilarity matrix. Coloured boxes group sampling units according to their topographic features. Both in **a** and **b**, the different topographic features are marked with colours (red=Platform, blue=Ravine, and yellow=Plateau).

S2. Delimitation of the breeding colony of yellow legged seagulls (*Larus michahellis*) in the Natural Reserve of Guaza Mountain:

To spatially extrapolate the results of the body-guard effect experiment to the rest of the distribution of *G. intermedia* that overlaps with *L. michahellis* breeding colony we first need to make a precise delimitation of such colony. Additionally, we need to predict the likely density of seagulls along its breeding area. For this we combined on site detection and delimitation along with remote sensing data and statistical modelling. To delimit the potential breeding and resting area of *L. michahellis*, we delimited 2 reference areas on which we confirmed the long-term presence of these animals. These reference areas were later projected onto a multi-channel high-definition aerial orthophotography (Figure S2.a-b). Using the range of pixel values from the Red, Green, and Blue (RGB) channels, we perform an initial filtering of the image, selecting only the pixels that presented values on the three different channels for the areas we have confirmed the presence of breeding seagulls (Figure S2.b). Based on this initial filtering we selected other two areas in which the satellite image confirms a strong concentration of seabird's guano (Figure S2.b). Once we defined these areas of permanent presence of *L. michahellis*, we used a 2m resolution Digital Elevation Model (DEM) to extract relevant topographic features; Slope, calculated using the difference in height and orientation of the 4 adjacent pixels (Fleming & Hoffer, 1979; Ritter, 1987); Aspect, calculated using the (Fleming & Hoffer, 1979; Ritter, 1987); Ruggedness, calculated as the average angle of the 4 neighbouring pixels with respect to the horizontal plane (Fleming & Hoffer, 1979; Ritter, 1987); and distance to the sea, calculated as the minimum Euclidean distance between each pixel centroid and the sea (elevation equal to 0 in our DEM). Similar than with the RBG channels previously described, we used the range of values from these terrain features to filter the spatial raster and select the areas more likely to be occupied by *L. michahellis* (Figure S2.C). To account for uncertainty and to remove gaps within the area classified as likely occupied by yellow legged gulls, we added a 15 m buffer around it. Using the hight definition orthophotography we confirmed that the selected areas matched the distribution of *L. michahellis* in the Natural Area of Guaza Mountain (Figure S2.e).

Figure S2. Workflow for the spatial delimitation of *Larus michahellis* breeding colony in the natural protected area of Guaza Mountain. The process implies the combination of field observations (counting of nest and seagull chicks around the sampling localities) and high-definition satellite data of areas occupied by *L. michahellis* (**a-b**). Once the areas of known presence of the species are delimited, in situ information about the gull's density is mixed with precise digital topography variables to model the predicted density of *L. michahellis* chicks along the coastal cliffs (**c-e**).

To predict the number of young chicks across the potential distribution area of *L.michahellis*, first we run a generalised linear model (GLM) with a Poisson distribution. For this model we used the number of yellow legged seagulls' chicks around the points used for the body-guard effect experiment (see main text) as the response variable. For the explanatory variables we used the derived terrain features from the DEM, and averaged the values of all the pixels that fall into the 5 m radius of our counting points (Figure S2.c). To account for the differences in units between the different terrain variables we standardize the data by subtracting the mean and dividing the values by the standard deviation of the variables. The model showed that slope have a positive effect on the number of chicks and whereas terrain aspect, distance to the sea, and terrain ruggedness had a negative effect. We use this fitted model to predict the potential number of chicks along the distribution area of *L.*

michahellis breeding colony (Table S5). We use this new layer to make further predictions (see main text).

Table S5. Results from the GLM Poisson-family used to estimate the number of *Larus michahellis* chicks along the coastal cliffs of the natural protected area of Guaza Mountain. Term estimates, standard deviation and z-values from the models are presented along with the Wald-chi type II test performed to test the statistical significance of the terms on the response variable. Statistically significant terms are marked in bold.

GLM family poisson model				Wald-χ2 type II test		
Term	Estimated	sd	z-value	χ2	d.f	p-value
intercept	-10.64	4.52	-2.35			
Slope	2.26	1.96	1.15	1.59	1	0.2
Aspect	-0.62	1.41	-0.43	0.18	1	0.66
Sea Dist	**-7.7**	**3.57**	**-2.15**	**9.66**	**1**	**0.001**
Ruggedness	-2.37	1.79	-1.32	2.12	1	0.14
Nagelkerke's R2	0.945					

High resolution orthophotography as well as the high-definition DEM were directly download from the open repositories of the Spanish National Geographic Institute or 'Instituto Geografico Nacional' (IGN, https://www.ign.es/web/ign/portal) with the references 'PNOA_MA_OF_REGC95_HU28_H50' and 'MDT02_REG95_HU28' respectively. All the spatial analysis and terrain features calculations were performed in R using the Terra version 1.5-34 (Hijmans, 2022) and *raster* version 3.5-21 (Hijmans, 2022) packages. The GLM were fitted using the R-packages *stats* version 4.3.1 (R Core Team, 2021) and *car* version 3.1-2 (Fox & Weisberg, 2019).

S3. Biometric data and exploratory analysis

Table S6. Biometric data of Gallotia intermedia by locality. The biometric data includes; Snout to Vent distance (SVL), measure ventrally in mm from the tip of the snout to the start of the cloaca; Pilleum length (Length-P), measure in mm following the total length of the pilleum scale complex; Pilleum width (widt-P), measure in mm along the widest section of the pilleum; Head length (length-H), mm from the back of the neck to the tip of the snout; Head width (width-H), mm across the widest portion of the head, normally around the cheeks; Total length, distance in mm from the tip of the snout to the tip of the tail; Age, classification according to the SVL of the animal (Adult= SVL>140; SubAdult= 120<SVL<140; Immature 100 < SVL <120; and Juvenile = SVL<100); Tail regeneration (tail-reg), binomial variable that take the value

"yes" if the animals shows signs of recent or past tail regeneration, and "no" otherwise; Sex, sex of the animals (male, female, ind= adult individual that was not possible to determine their sex, immature= the animal was young at it was not possible to determine their sex by morphological traits).

Locality: Andén de los Ñames (platfomr)

ID	Weigth	Total-length	Age	SVL	widht-P	lenth-P	length-H	width-H	tail-reg	SEX
1	80	368	SubAdult	132	13.9	30.1	30.1	18.1	yes	female
2	145	412	Adult	148	17.7	36.9	38.5	22.7	yes	male
3	21	337	Juvenile	96	10.3	23.2	22.9	13.3	yes	ind
7	88	364	SubAdult	132.5	15	31.8	33.4	18.9	yes	female
8	92	384	SubAdult	137	13.9	29.9	32.2	17.5	yes	female
11	40	250	Immature	106	11.8	25.5	27.6	14.4	yes	ind
13	63	411	SubAdult	124	12.9	28.1	30.2	17.4	yes	ind
14	25	297	Juvenile	89	10.1	21.6	23.1	12	yes	ind

Locality: Barranco del Cabezón (ravine)

ID	Weigth	Total-length	Age	SVL	widht-P	lenth-P	length-H	width-H	tail-reg	SEX
15	56	387	SubAdult	123	13	30	31.7	17.4	yes	male
16	49	348	Immature	116	11.9	26	28.6	14.7	no	immature
17	26	295	Juvenile	95	10.5	22.5	25	13.2	no	immature
18	96	350	SubAdult	132	14.1	31.4	34.1	18.9	yes	female
19	96	362	SubAdult	139	14.9	32.7	35.6	20.2	yes	female
20	86	380	Adult	141	14.3	31.9	34.3	19.3	yes	female
21	NA	382	Immature	115	11.9	27.1	29.9	15.3	yes	female
22	81	429	Adult	141	15.9	34	36.8	20.7	yes	male
23	45	338	Immature	117	12.4	28.4	31	16.3	yes	ind
24	123	471	Adult	154	17.5	39	40.8	23.5	yes	ind
25	92	381	Adult	155	15.3	33.3	36.5	20.5	yes	male
26	25	337	Immature	101	10.1	23.4	21.2	12.9	yes	ind

Locality: Foso caimanes (ravine)

ID	Weigth	Total-length	Age	SVL	widht-P	lenth-P	length-H	width-H	tail-reg	SEX
4	118	475	SubAdult	139.5	18	37.4	36.75	23.5	no	male
5	126	462	Adult	163	16.9	34.7	34.6	22.6	yes	female
6	117	409	Adult	146	16	34.7	36.5	22.6	yes	female
9	43	357	Immature	108.5	11.7	25.8	27	15.1	yes	female
10	169	460	Adult	166	22	44.2	44	27.3	yes	male
12	70	375	SubAdult	128	13.7	30.3	31.7	17.7	yes	female

Locality: Foso de la Nicotiana (platfomr)

ID	Weigth	Total-length	Age	SVL	widht-P	lenth-P	length-H	width-H	tail-reg	SEX
27	68	410	SubAdult	132	13.2	29.3	32.4	16.7	yes	female
28	109	410	Adult	145	16.9	35.3	38.7	22.3	yes	male

S4. Stable Isotopes Values

Table S6. $\delta^{15}N$ and $\delta^{13}C$ stable isotope values of vegetation collected at the Guaza Mountain at the cliff platforms, ravines and plateau habitats.

Species	$\delta^{15}N$	$\delta^{13}C$	Foraging group	Family	Metabolic
Argyranthemum spp.	4.14	-27.68	Ravine/Plateu	Asteraceae	C_3
Argyranthemum spp.	6.05	-24.27	Ravine/Plateu	Asteraceae	C_3
Argyranthemum spp.	7.14	-22.65	Ravine/Plateu	Asteraceae	C_3
Argyranthemum spp.	7.26	-26.36	Ravine/Plateu	Asteraceae	C_3
Argyranthemum spp.	7.90	-24.15	Ravine/Plateu	Asteraceae	C_3
Argyranthemum spp.	8.28	-27.00	Ravine/Plateu	Asteraceae	C_3
E. balsamifera	5.69	-25.37	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	5.90	-25.73	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	7.18	-26.06	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	7.41	-25.82	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	7.71	-26.49	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	9.32	-24.58	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	9.58	-21.63	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	9.88	-21.74	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	11.24	-20.98	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	11.38	-20.95	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	12.86	-21.30	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	12.96	-19.89	Platform	Euphorbiaceae	C_3
E. balsamifera	13.56	-21.96	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	13.95	-21.30	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	14.19	-23.49	Platform	Euphorbiaceae	C_3
E. balsamifera	14.37	-21.04	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	14.83	-23.23	Platform	Euphorbiaceae	C_3
E. balsamifera	14.91	-21.23	Platform	Euphorbiaceae	C_3
E. balsamifera	15.09	-19.16	Platform	Euphorbiaceae	C_3
E. balsamifera	15.40	-21.05	Ravine/Plateu	Euphorbiaceae	C_3
E. balsamifera	15.71	-20.45	Platform	Euphorbiaceae	C_3
E. balsamifera	17.71	-23.54	Platform	Euphorbiaceae	C_3
F. cretica	13.66	-26.27	Platform	Zygophyllaceae	C_3
F. cretica	15.21	-24.95	Platform	Zygophyllaceae	C_3
F. cretica	15.25	-27.14	Platform	Zygophyllaceae	C_3
F. cretica	15.39	-27.41	Platform	Zygophyllaceae	C_3
F. cretica	15.42	-28.12	Platform	Zygophyllaceae	C_3
F. ericifolia	7.65	-22.33	Ravine/Plateu	Frankeniaceae	C_3

F. ericifolia	7.79	-23.02	Ravine/Plateu	Frankeniaceae	C$_3$
F. ericifolia	8.33	-24.02	Ravine/Plateu	Frankeniaceae	C$_3$
K. nerrifolia	5.64	-22.96	Ravine/Plateu	Asteraceae	C$_3$
K. nerrifolia	6.11	-20.94	Ravine/Plateu	Asteraceae	C$_3$
K. nerrifolia	6.15	-21.40	Ravine/Plateu	Asteraceae	C$_3$
K. nerrifolia	12.63	-20.62	Platform	Asteraceae	C$_3$
K. nerrifolia	15.43	-21.13	Platform	Asteraceae	C$_3$
K. nerrifolia	15.59	-20.24	Platform	Asteraceae	C$_3$
L. intricatum	13.94	-23.82	Platform	Solanaceae	C$_3$
L. intricatum	15.07	-25.70	Platform	Solanaceae	C$_3$
L. intricatum	16.10	-26.17	Platform	Solanaceae	C$_3$
L. intricatum	16.54	-25.09	Platform	Solanaceae	C$_3$
L. intricatum	17.57	-26.11	Platform	Solanaceae	C$_3$
L. intricatum	18.26	-27.18	Platform	Solanaceae	C$_3$
L. intricatum	18.51	-26.28	Platform	Solanaceae	C$_3$
L. intricatum	19.15	-27.75	Platform	Solanaceae	C$_3$
M. crystallinum	14.49	-16.74	Platform	Aizoaceae	C$_4$/CAM
M. crystallinum	16.52	-17.40	Platform	Aizoaceae	C$_4$/CAM
M. crystallinum	17.61	-15.47	Platform	Aizoaceae	C$_4$/CAM
M. crystallinum	17.63	-14.97	Platform	Aizoaceae	C$_4$/CAM
M. crystallinum	18.16	-14.59	Platform	Aizoaceae	C$_4$/CAM
M. crystallinum	19.40	-14.00	Platform	Aizoaceae	C$_4$/CAM
N. glauca	15.44	-23.93	Platform	Solanaceae	C$_3$
O. dillenii	9.14	-12.58	Platform	Cactaceae	C$_4$/CAM
O. dillenii	11.41	-12.81	Platform	Cactaceae	C$_4$/CAM
O. dillenii	13.36	-12.34	Platform	Cactaceae	C$_4$/CAM
O. dillenii	14.87	-13.99	Platform	Cactaceae	C$_4$/CAM
P. patellaris	14.05	-23.41	Platform	Chenopodiaceae	C$_3$
P. patellaris	14.26	-22.56	Platform	Chenopodiaceae	C$_3$
P. patellaris	14.34	-26.67	Platform	Chenopodiaceae	C$_3$
P. patellaris	14.47	-25.41	Platform	Chenopodiaceae	C$_3$
P. patellaris	14.59	-27.22	Platform	Chenopodiaceae	C$_3$
P. patellaris	15.48	-25.56	Platform	Chenopodiaceae	C$_3$
P. patellaris	16.63	-24.21	Platform	Chenopodiaceae	C$_3$
P. patellaris	17.67	-22.78	Platform	Chenopodiaceae	C$_3$
P. patellaris	17.72	-22.75	Platform	Chenopodiaceae	C$_3$
P. patellaris	18.57	-21.62	Platform	Chenopodiaceae	C$_3$
P. patellaris	18.60	-22.91	Platform	Chenopodiaceae	C$_3$
P. patellaris	18.75	-22.24	Platform	Chenopodiaceae	C$_3$
P. pendula	4.38	-25.39	Ravine/Plateu	Rubiaceae	C$_3$

P. pendula	6.82	-26.17	Ravine/Plateu	Rubiaceae	C_3
P. pendula	8.11	-27.07	Ravine/Plateu	Rubiaceae	C_3
Portulaca spp.	16.69	-14.79	Platform	Portulacaceae	C_4/CAM
Portulaca spp.	19.49	-14.00	Platform	Portulacaceae	C_4/CAM
Portulaca spp.	20.12	-14.24	Platform	Portulacaceae	C_4/CAM

5 Table S7: *p*-values of pairwise comparisons of $\delta^{13}C$‰ and $\delta^{15}N$‰ values among forage item groups
6 using a Wilcoxon rank-sum test. *: Statistically significant values $p < 0.05$.

$\delta^{15}N$ values	Platform C_3	Platform C_4/CAM	Plateu/Ravine C_3
Group 1: Platform C_3	-		
Group 2: Platform C_4/CAM	0.51	-	
Group 3: Plateu/Ravine C_3	9.5e-14*	3.7e-07*	-
Group 4: Seabirds-derived resources	3.4e-10*	0.00033*	0.00847*
$\delta^{13}C$ values			
Group 1: Platform C_3	-		
Group 2: Platform C_4/CAM	4.60e-11*	-	
Group 3: Plateu/Ravine C_3	0.53	1.60e-10*	-
Group 4: Seabirds-derived resources	4.70e-08*	9.60e-07*	1.30e-09*

S5. Media

Figure S3. Exampled of recorded dive attack on the cat decoy (**a-c**) and cat decoy displayed in an open area nearby Guaza Mountain *L. michahellis* ground breeding colony (**d**).

Figure S4. Exampled of recorded mobbing behaviour of *L. michahellis* over an Eleonoras falcon over the breeding colony (**a-c**) and against one of the researchers while sampling into the breeding colony (**d**).

Bibliography

Clarke, K. R. (1993). Non-parametric multivariate analyses of changes in community structure. *Austral Ecology*, *18*(1), 117–143. https://doi.org/10.1111/j.1442-9993.1993.tb00438.x

Elzinga, C. L., Salzer, D. W., Willoughby, J. W., & Gibbs, J. P. (2001). *Monitoring plant and animal populations: A handbook for field biologists*. John Wiley & Sons.

Fleming, M. D., & Hoffer, R. M. (1979). *Machine Processing of Landsat MSS Data and DMA Topographic Data for Forest Cover Type Mapping*.

Fox, J., & Weisberg, S. (2019). *An R Companion to Applied Regression*. Sage. https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard, D., Carvalho, G., Chirico, M., Caceres, M. D., Durand, S., … Weedon, J. (2022). *vegan: Community Ecology Package*. https://CRAN.R-project.org/package=vegan

R Core Team. (2021). R: A Language and Environment for Statistical Computing. *R Foundation for Statistical Computing*. https://www.r-project.org/

Ritter, P. (1987). *A Vector-Based Slope and Aspect Generation Algorithm*.