Limited potexvirus diversity in eastern Gulf of Mexico seagrass meadows

1.1 Author names

Shen Jean Lim[1*] (https://orcid.org/0000-0003-4578-5318)

Karyna Rosario[1] (https://orcid.org/0000-0001-9847-4113)

Meredith E. Kernbach[1] (https://orcid.org/0000-0001-8161-057X)

Anthony J. Gross[1] (https://orcid.org/0000-0001-6392-5121)

Bradley T. Furman[2] (https://orcid.org/0000-0002-9284-8388)

Mya Breitbart[1] (https://orcid.org/0000-0003-3210-2899)

1.2 Affiliation(s)

[1] College of Marine Science, University of South Florida, St Petersburg, FL, USA

[2] Florida Fish and Wildlife Conservation Commission, Florida Fish and Wildlife Research Institute, Saint Petersburg, FL, USA

1.3 Corresponding author and email address

Shen Jean Lim (jeanlim@usf.edu)

1.4 Keywords

Potexvirus, primer, PCR, seagrass, marine, plant, *Alphaflexivirdae*

1.5 Repositories:

All sequence data have been deposited into NCBI GenBank under the accession numbers OR827689-OR827705, OR854648, OR863396, OR879052-OR879056, and PP430548-PP430571.

2. Abstract

Turtlegrass virus X, which infects the seagrass *Thalassia testudinum*, is the only potexvirus known to infect marine flowering plants. We investigated potexvirus distribution in seagrasses using a degenerate reverse transcription polymerase chain reaction (RT-PCR) assay originally designed to capture potexvirus diversity in terrestrial plants. The assay, which implements Potex-5 and Potex-2RC primers, successfully amplified a 584 nt RNA-dependent RNA polymerase (RdRp) fragment from TVX-infected seagrasses. Following validation, we screened 74 opportunistically collected, apparently healthy seagrass samples for potexviruses using this RT-PCR assay. The survey examined the host species *T. testudinum*, *Halodule wrightii*, *Halophila stipulacea*, *Syringodium filiforme*, *Ruppia maritima*, and *Zostera marina*. Potexvirus PCR products were successfully generated only from *T. testudinum* samples and phylogenetic analysis of sequenced PCR products revealed five distinct TVX sequence variants. Although the RT-PCR assay revealed limited potexvirus diversity in seagrasses, the expanded geographic distribution of TVX shown here emphasizes the importance of future studies to investigate *T. testudinum* populations across its native range and understand how the observed fine-scale genetic diversity affects host-virus interactions.

3. Impact statement

Potexviruses are widespread in terrestrial plants; however, the recent discovery of TVX in the seagrass *Thalassia testudinum* extends their host range to marine flowering plants. Here we use existing Potex-5 and Potex-2RC degenerate primers to explore potexvirus infections in several seagrass species. TVX sequence variants were detected in *T. testudinum* collected from the eastern Gulf of Mexico, uncovering previously unknown genetic diversity of this poorly understood virus.

4. Data summary

All sequence data are available in NCBI GenBank under the accession numbers OR827692-OR827705, OR854648, OR863396, OR879052-OR879056, and PP430548-PP430571. The authors confirm all supporting data, code and protocols have been provided within the article or through supplementary data files.

5. Introduction

Potexviruses are positive-sense single-stranded RNA (ssRNA) viruses from the family *Alphaflexiviridae* that infect many agronomically important flowering plants (angiosperms), including potato, tobacco, and tomato (1). The type member of this genus is *Potato virus X* (PVX), from which the genus name *Potexvirus* is derived (2). Potexviruses have a broad

geographic range that reflects the distribution of their hosts and the prevalence of global
commerce (2, 3). Although each potexvirus species has a limited natural host range, some
species can infect a wide range of host plants in experimental settings (1). Potexviruses are
transmitted mechanically through contaminated agricultural equipment or plant-to-plant
contact (1-3). Transmission via seed and aphid vectors and non-specific transmission through
chewing insects has been reported but is less common (1-3). Most potexviruses cause
persistent, asymptomatic infections or mild mosaic disease, while some cause dwarf,
necrotic, or ringspot symptoms in their natural hosts (1, 2). Notably, PVX can cause severe
symptoms when its plant host is co-infected with potyviruses (4-6).

In addition to infecting terrestrial plants, potexviruses are also capable of infecting
seagrasses, the only marine angiosperms. A novel potexvirus species, *Turtlegrass virus X*
(TVX), was discovered in apparently healthy *Thalassia testudinum* seagrass samples collected
from Terra Ceia Aquatic Preserve, Tampa Bay, Florida, USA (7). The identification of TVX
indicates that an important aspect of potexvirus diversity has been overlooked by focusing
only on terrestrial plants. Degenerate primers used in reverse transcriptase polymerase
chain reaction (RT-PCR) assays, such as Potex-5 and Potex-2RC (8), have proven useful in
detecting diverse potexviruses, but have not been tested in aquatic habitats. This study
applied the Potex-5 and Potex-2RC primer pair in a RT-PCR assay to explore potexvirus
diversity in samples from six seagrass host species.

6. Methods

Primer validation. Potex-5/Potex-2RC were tested on cDNA samples each synthesized from 1
μg RNA extracted from *Portulaca* sp. infected with *Alternanthera mosaic virus* (AltMV) and
Opuntia sp. with a mixed infection of potexviruses, as well as cDNA pooled from *T.*
testudinum samples known to contain TVX, based on the TVX-specific RT-PCR assay
described by Van Bogaert *et al.* 2019 (7). The PCR reaction contained 0.48 μM of each
primer, 2 μL cDNA template, 1 μL GC enhancer, and 1X AmpliTaq Gold™ 360 Master Mix
(Applied Biosystems™, Waltham, MA, USA) in a 25 μL reaction volume. PCR was performed
under the following conditions: initial denaturation at 95°C for 10 minutes, 40 cycles of
denaturation at 95°C for 30 seconds, annealing at 51.5°C (as published in van der Vlugt and
Berendsen (8)) for 30 seconds, extension at 72°C for 1 minute, followed by elongation at
72°C for 10 minutes and cooling at 11°C. The PCR product was visualized following gel
electrophoresis on a 1% (wt/vol) agarose gel stained with ethidium bromide. All PCR
reactions, except for the no template control, yielded visible bands. All PCR products were
purified using the Zymoclean Gel DNA Recovery Kit (Irvine, CA, USA), quantified using the
Qubit™ DNA high sensitivity (HS) assay (Invitrogen™, Waltham, MA, USA), and Sanger
sequenced bidirectionally by Eurofins Genomics (Louisville, KY, USA).

RT-PCR. Total RNA extraction was performed on 30-100 mg of leaves from multiple shoots
pooled by seagrass species and collection site (**Table S1**) using Zymo Research's (ZR) Quick-
RNA™ Plant Miniprep kit. Each pooled seagrass sample was homogenized in a BashingBead
Lysis Tube containing 2 mm ceramic beads and 800 µL RNA lysis buffer (provided in the kit)
for 5 minutes at maximum speed using a Fisherbrand™ Bead Mill 4 Homogenizer (Fisher
Scientific, Waltham, MA, USA). Tissue homogenates were centrifuged at maximum speed
(21,130 _g_) for 1 minute and RNA was extracted from the total volume (~800 µL) of the
supernatant. To ensure successful RNA extraction, after each round of extraction, a random
subset of RNA samples was quantified using the Qubit™ RNA high sensitivity (HS) assay
(Invitrogen™). From each sample, cDNA was synthesized from 8 µL RNA using the
SuperScript™ IV First-Strand Synthesis System (Invitrogen™) and following manufacturer's
instructions for random hexamers. PCR amplification was performed on each cDNA sample,
followed by gel electrophoresis, PCR product purification, DNA quantification, and Sanger
sequencing, using the methods described above.

Amplicon sequence analysis. Amplicon sequences were first compared with sequences from
NCBI's nucleotide (nt) and non-redundant protein (nr) collections (9) and the TVX genome
(7) using the megablast and/or blastx programs on NCBI's BLAST® server (10). Sequence
reads from each sample were then mapped to the Potex-5/Potex-2RC amplicon region (with
forward and reverse primer sequences removed) in the TVX genome (7) using the "Map
Sanger Reads to Reference" function implemented in Unipro UGENE v48.1 (11), with a
trimming quality threshold of 20 and mapping minimum similarity of 70%.
Electropherograms of mapped reads were manually inspected to check for mixed infection
of potexviruses, to remove primer sequences and low-quality bases, and to resolve
ambiguous bases.

Sequences from 42 Potex-5/Potex-2RC PCR products amplified from seagrass and terrestrial
plant samples described above were aligned with corresponding amplicon regions (excluding
primer sequences) extracted from the reference genomes of TVX (NC_040644) (7), _Bamboo_
mosaic virus (NC_001642) (12), _Foxtail mosaic virus_ (NC_001483) (13), AltMV (OR607766),
PVX (NC_011620) (14), and _Lolium latent virus_ (NC_010434) (15) retrieved from the NCBI
Virus database (16). Multiple sequence alignment was performed using the L-INS-I and --
adjustdirectionaccurately options in MAFFT v7.5.08 (17), then trimmed to 514 nt to remove
leading and trailing gaps. Pairwise sequence identities were calculated from the trimmed
alignment using Clustal Omega v1.2.3 (18) with the --distmat-out, --percent-id, and --full

options. For phylogenetic analysis, Molecular Evolutionary Genetics Analysis (MEGA) 11 (19)
identified the best model for the trimmed alignment to be the Hasegawa-Kishino-Yano
model (20) with a discrete Gamma distribution (HKY+G). Using this substitution model, a
maximum likelihood tree with 1,000 bootstrap replicates was constructed from the
alignment using MEGA11 (19). The tree was condensed using MEGA11 (19) to eliminate
interior branches with <50% bootstrap support. The final tree was visualized together with
the heatmap of pairwise sequence identity using the ggtree R package v3.6.2 (21).

7. Results

The Potex-5/Potex-2RC primer pair successfully amplified TVX from *T. testudinum* samples
collected from Terra Ceia Aquatic Preserve, AltMV infecting *Portulaca* sp., and an
uncharacterized potexvirus infecting *Opuntia* sp. (8). A RT-PCR assay using Potex-5/Potex-2RC
was subsequently performed on seagrass leaves opportunistically collected from separate
sampling efforts, including 1) *T. testudinum*, *H. wrightii*, *S. filiforme*, and *R. maritima* from
Tampa Bay on Jan 26[th] or Feb 4[th], 2022; 2) *T. testudinum* from the Dry Tortugas National
Park, Florida, USA collected between May 16[th]-20[th], 2022; 3) *T. testudinum* from a systematic
seagrass survey at Terra Ceia Aquatic Preserve on August 1[st], 2022; and 4) *T. testudinum*, *H.*
wrightii, and *S. filiforme* samples collected from Tampa Bay seagrass site S3T8 (Lassing Park)
on October 3[rd], 2023 (**Figure 1** and **Table S1**). Seagrass samples (**Table S1**) were also received
from other research groups, including *T. testudinum* from Panacea (Florida, USA), *Zostera*
marina from York River (Virginia, USA), West Falmouth Harbor (Massachusetts, USA), Sitka
(Alaska, USA), New Zealand, and Kalmar (Sweden), and *Halophila stipulacea* from Jobos Bay
National Estuarine Research Reserve (Puerto Rico, USA). PCR products were successfully
amplified and sequenced from 40 *T. testudinum* samples from Florida sites, including Terra
Ceia Aquatic Preserve, Tampa Bay seagrass sites S1T5 and S3T8 (Lassing Park), Panacea
located in the Florida Panhandle, and Florida Keys sites including Bush Key, Garden Key,
Marquesas Key, and Key West (**Figure 1**). Based on our analysis of Sanger sequencing traces,
no evidence of mixed potexvirus infection was found in these *T. testudinum* samples. Aside
from *T. testudinum*, no other seagrass species produced PCR products.

Figure 1. Map showing locations of seagrass sites in (A) Florida, (B) Tampa Bay, and (C) Terra
Ceia Aquatic Preserve where samples were collected for RT-PCR. Purple markers in (A) and
(B) denote sites where TVX products were amplified from *T. testudinum* with the Potex-
5/Potex-2RC primers, whereas samples from sites with orange markers tested negative.
Marker shapes in (C) represent potexvirus subclades found in Terra Ceia Aquatic Preserve
sampling sites and correspond to **Figure 2** (negative samples not shown). Metadata of
seagrass samples is described in **Table S1**.

Phylogenetic analysis placed all Potex-5/Potex-2RC amplicons from seagrasses in the same
clade as the RdRp sequence fragment from the TVX genome (7) with 100% bootstrap
confidence (**Figure 2A**). Sequences within this clade shared 74.4% to 98.6% nucleotide
sequence identities to each other (**Figure 2B**). TVX sequences from Terra Ceia Aquatic
Preserve formed three subclades. Average pairwise nucleotide sequence identities were
95.8±1.4% within each subclade and 79.3±2.9% between subclades (**Figure 2**). Two Terra
Ceia Aquatic Preserve subclades, including the subclade containing the amplicon region from
the TVX genome, were sister groups to each other, sharing 84.6±1% average pairwise
nucleotide sequence identities (**Figure 2**). Other TVX sequences from Key West, Marquesas
Key, and Panacea formed a subclade with 99.6±0.2% average pairwise nucleotide sequence
identities, while sequences from Garden Key and Bush Key formed another subclade with
89.3±0.9% average pairwise nucleotide sequence identities (**Figure 2**). Potexviruses that
infect terrestrial grasses (*Foxtail mosaic virus* and *Bamboo mosaic virus*) formed sister clades
to the TVX-containing clade (**Figure 2A**).

A   Maximum likelihood tree with 1,000 bootstrap replicates

B   Pairwise nucleotide sequence identities

Figure 2. (A**)** Bootstrap maximum likelihood tree and (B) heatmap of pairwise nucleotide sequence identities of TVX nucleotide sequences
amplified from seagrasses collected from Terra Ceia Aquatic Preserve (purple text) and other sites in the eastern Gulf of Mexico (green text), in
relation to potexvirus sequences amplified from terrestrial plants (brown text) and potexvirus RdRp sequences from reference genomes (black
text). The NCBI accession number for each sequence in (A) is indicated in square brackets. The tree in (A) was constructed by MEGA11 from a
514 nt conserved region in the multiple sequence alignment. Sizes of tree nodes are proportionate to bootstrap values indicating the
percentage of trees, based on 1,000 replicates, in which sequences within a node were clustered together. Only bootstrap values >70% are
labelled on the tree nodes. *Lolium latent virus* was used as an outgroup for the tree. Marker shapes in (A) and (B) preceding sequences in the
Terra Ceia Aquatic Preserve subclades correspond to marker shapes in **Figure 1C**. Abbreviations: TVX, *Turtlegrass virus X*; AltMV, *Alternanthera*
mosaic virus; PVX, *Potato virus X*; LolV, *Lolium latent virus*; FoMV, *Foxtail mosaic virus*; BaMV, *Bamboo mosaic virus*.

8. Discussion

In this study, we validated the ability of the potexvirus degenerate primers, Potex-5/Potex-2RC, published by van der Vlugt and Berendsen (8), to amplify TVX-infected seagrass samples and potexvirus-infected terrestrial plant samples. Potex-5/Potex-2RC were used in a RT-PCR survey to test six seagrass species for potexviruses. Among the seagrass species examined, potexvirus infection was only detected in *T. testudinum*, the species from which TVX was first identified. The criteria published by the International Committee on Taxonomy of Viruses (ICTV) demarcates potexviruses into species based on host range, absence of cross-protection in infected plants, serology, and sequence identity thresholds of 72% for nucleotide sequences and 80% for amino acid sequences (https://ictv.global/report/chapter/alphaflexiviridae/alphaflexiviridae/potexvirus). Potexviruses amplified in this study are specific to *T. testudinum* and shared >72% sequence identity to each other and to the RdRp sequence fragment from the TVX genome (7). Therefore, we considered identified potexviruses the same species as TVX. Note that this study only analyzed a short and conserved fragment of the RdRp sequence and these pairwise identities may change when considering complete genomes. Nevertheless, our results suggest limited potexvirus diversity in marine angiosperms. RT-PCR analysis revealed *T. testudinum* as a natural host for TVX and confirmed TVX infection in eastern Gulf of Mexico seagrass meadows, including those in northwest (Panacea), west-central (Tampa Bay), and southwest (Florida Keys) Florida. Previous phylogenetic analyses of the PVX coat protein and genomic sequences identified phylogroups that reflect intraspecies diversity (22, 23). Here, three distinct TVX phylogroups, sharing >95% intra-group and <80% inter-group amplicon sequence identity, were observed within a single seagrass meadow at Terra Ceia Aquatic Preserve, along with two additional phylogroups in other Tampa Bay sites, namely the Florida Keys and Panacea.

Among all sites sampled in this study, the most extensive TVX survey was conducted within the Terra Ceia Aquatic Preserve. Systematic RT-PCR surveys at other sites will be useful in elucidating potexvirus diversity in seagrasses. The sampling scheme in this study was limited to *T. testudinum* populations along the eastern Gulf of Mexico. However, *T. testudinum* has a widespread geographic range across the Western Atlantic Ocean with high genetic diversity that is generally partitioned into two clusters corresponding to the Gulf of Mexico or the Caribbean phylogeographic regions (24). Although seagrasses can reproduce asexually, some *T. testudinum* populations, such as those in Tampa Bay (25-27) and Florida Bay (28, 29), undergo sexual reproduction and exhibit high genotypic diversity. Currently, the relationship between *T. testudinum* genotype and TVX diversity is unknown and should be further investigated. Future potexvirus surveys covering the total distribution range of *T. testudinum*, the related species *T. hemprichii*, and other seagrass species also have the potential to

229 uncover relationships between potexvirus diversity, host diversity, and their phylogeographic
230 distributions.

9. Author statements

9.1 Author contributions

SJL: Data curation, Formal analysis, Investigation, Methodology, Software, Supervision,
Visualization, Writing - original draft

KRC: Conceptualization, Formal analysis, Funding acquisition, Writing – review & editing.

MEK: Investigation, Writing – review & editing

AJG: Investigation, Writing - review & editing

BTF: Funding acquisition, Investigation, Writing - review & editing

MB: Conceptualization, Formal analysis, Funding acquisition, Investigation, Methodology,
Project administration, Resources, Supervision, Validation, Writing - review & editing.

9.2 Conflicts of interest

243 The author(s) declare that there are no conflicts of interest.

9.3 Funding information

245 This project was supported by award OCE-2219547 from the National Science Foundation to
246 MB, KRC, and BTF and a Proposal Enhancement Grant from the University of South Florida
247 Internal Awards Program to MB. This research was made possible by support from the
248 Florida Institute of Oceanography for collection of the Dry Tortugas samples.

9.4 Ethical approval

250 N/A

9.5 Consent for publication

252 N/A

9.6 Acknowledgements

Fieldwork was conducted under permit no. FKNMS-2022-026 issued by the Florida Keys
National Marine Sanctuary, permit no. DRTO-2022-SCI-0004 issued through the U.S.
Department of the Interior National Park Service and permit no. SAL-19-1172A-SR issued
through the Florida Fish and Wildlife Conservation Commission. We acknowledge the
contributions of Luis Alvarado-Marchena, Victoria Congdon, Keith Keel, Isabella Ritchie,

Taylor Little, Jessye Kirkham, and Mike Wheeler in field work and sample collection at these

sites. Isabella Ritchie provided cDNAs from *T. testudinum* samples from Terra Ceia Aquatic

Preserve. Salvador P Lopez, Jr and Scott Adkins from the Agricultural Research Service,

United States Horticultural Research Laboratory, U.S. Department of Agriculture (USDA), Fort

Pierce, FL, USA provided cDNAs from potexvirus-infected plant samples. We thank Maya

Groner, Ian Hewson, Karin Holmfeldt, Natalia Lopez Figueroa, Carolyn Malmstrom, Jordan

Rede, Cypress Rudloe, and Jack Rudloe for providing seagrass samples from other sites.

266 10. References

1.   Ryu KH, Hong JS. Potexvirus. In: Mahy BWJ, Van Regenmortel MHV, editors. Encyclopedia of
Virology (Third Edition). Oxford: Academic Press; 2008. p. 310-3.
2.   Verchot J. *Potato virus X*: A global potato-infecting virus and type member of the *Potexvirus*
genus. Mol Plant Pathol. 2022;23(3):315-20.
3.   AbouHaidar MG, Gellatly D. Potexviruses. In: Granoff A, Webster RG, editors. Encyclopedia of
Virology (Second Edition). Oxford: Elsevier; 1999. p. 1364-8.
4.   Rochow WF, Ross AF. Virus multiplication in plants doubly infected by potato viruses X and Y.
Virology. 1955;1(1):10-27.
5.   Pacheco R, Garcia-Marcos A, Barajas D, Martianez J, Tenllado F. PVX-potyvirus synergistic
infections differentially alter microRNA accumulation in *Nicotiana benthamiana*. Virus Res.
2012;165(2):231-5.
6.   Pollari M, Sipari N, Poque S, Himanen K, Makinen K. Effects of poty-potexvirus synergism on
growth, photosynthesis and metabolite status of *Nicotiana benthamiana*. Viruses. 2022;15(1):121.
7.   Van Bogaert N, Rosario K, Furman BT, Hall MO, Greco AM, Breitbart M. Discovery of a novel
potexvirus in the seagrass *Thalassia testudinum* from Tampa Bay, Florida. Limnol Oceanogr Lett.
2019;4(1):1-8.
8.   van der Vlugt RAA, Berendsen M. Development of a general potexvirus detection method.
Eur J Plant Path. 2002;108(4):367-71.
9.   Sayers EW, Beck J, Bolton EE, Bourexis D, Brister JR, Canese K, et al. Database resources of
the National Center for Biotechnology Information. Nucleic Acids Res. 2021;49(D1):D10-D7.
10.   Johnson M, Zaretskaya I, Raytselis Y, Merezhuk Y, McGinnis S, Madden TL. NCBI BLAST: a
better web interface. Nucleic Acids Res. 2008;36(Web Server issue):W5-9.
11.   Okonechnikov K, Golosova O, Fursov M, team U. Unipro UGENE: a unified bioinformatics
toolkit. Bioinformatics. 2012;28(8):1166-7.
12.   Lin NS, Lin BY, Lo NW, Hu CC, Chow TY, Hsu YH. Nucleotide sequence of the genomic RNA of
bamboo mosaic potexvirus. J Gen Virol. 1994;75 (9):2513-8.
13.   Bancroft JB, Rouleau M, Johnston R, Prins L, Mackie GA. The entire nucleotide sequence of
Foxtail mosaic virus RNA. J Gen Virol. 1991;72 (9):2173-81.
14.   Huisman MJ, Linthorst HJ, Bol JF, Cornelissen JC. The complete nucleotide sequence of *Potato*
virus X and its homologies at the amino acid level with various plus-stranded RNA viruses. J Gen
Virol. 1988;69 (Pt 8):1789-98.
15.   Vaira AM, Maroon-Lango CJ, Hammond J. Molecular characterization of *Lolium latent virus*,
proposed type member of a new genus in the family Flexiviridae. Arch Virol. 2008;153(7):1263-70.
16.   Brister JR, Ako-Adjei D, Bao Y, Blinkova O. NCBI viral genomes resource. Nucleic Acids Res.
2015;43(Database issue):D571-7.
17.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol. 2013;30(4):772-80.

18.   Sievers F, Higgins DG. Clustal Omega for making accurate alignments of many protein
sequences. Protein Sci. 2018;27(1):135-45.
19.   Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis Version
11. Mol Biol Evol. 2021;38(7):3022-7.
20.   Hasegawa M, Kishino H, Yano T. Dating of the human-ape splitting by a molecular clock of
mitochondrial DNA. J Mol Evol. 1985;22(2):160-74.
21.   Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. GGTREE: an R package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. MEE. 2017;8(1):28-
36.
22.   Cox BA, Jones RA. Genetic variability in the coat protein gene of *Potato virus X* and the
current relationship between phylogenetic placement and resistance groupings. Arch Virol.
2010;155(8):1349-56.
23.   Fuentes S, Gibbs AJ, Hajizadeh M, Perez A, Adams IP, Fribourg CE, et al. The phylogeography
of *Potato Virus X* shows the fingerprints of its human vector. Viruses. 2021;13(4).
24.   van Dijk K-j, Bricker E, van Tussenbroek BI, Waycott M. Range-wide population genetic
structure of the Caribbean marine angiosperm *Thalassia testudinum*. Ecol Evol. 2018;8(18):9478-90.
25.   Grey WF, Moffler MD. Flowering of the seagrass *Thalassia testudinum* (Hydrocharitaceae) in
the Tampa Bay, Florida area. Aquat Bot. 1978;5:251-9.
26.   Durako MJ, Moffler MD. Factors affecting the reproductive ecology of *Thalassia testudinum*
(Hydrocharitaceae). Aquat Bot. 1987;27(1):79-95.
27.   Campanella JJ, Bologna PAX, Carvalho M, Smalley JV, Elakhrass M, Meredith RW, et al. Clonal
diversity and connectedness of turtle grass (*Thalassia testudinum*) populations in a UNESCO
Biosphere Reserve. Aquat Bot. 2015;123:76-82.
28.   Bricker E, Waycott M, Calladine A, Zieman JC. High connectivity across environmental
gradients and implications for phenotypic plasticity in a marine plant. Mar Ecol Prog Ser.
2011;423:57-67.
29.   Peñalver MM, Durako M, Furman BT, Hall M. Multiple stressors result in reduced
reproductive effort by *Thalassia testudinum* in Florida Bay, USA. Mar Ecol Prog Ser. 2020;647.

Table S1. Metadata of seagrass samples used in the RT-PCR survey. Samples successfully amplified by Potex-5/Potex-2RC primers are
highlighted in bold and the NCBI accession numbers of their amplicon sequences are listed in the last column.

Sample ID	State, Country	Region	Collection Date	Site Description	Latitude	Longitude	Species	Accession
S1T5 Tt	**FL, USA**	**Upper Tampa Bay**	**4-Feb-22**	**S1T5 - Old Tampa Bay**	**27.95788**	**-82.54663**	***Thalassia testudinum***	**OR827692**
S1T5 Sf	FL, USA	Upper Tampa Bay	4-Feb-22	S1T5 - Old Tampa Bay	27.95788	-82.54663	*Syringodium filiforme*	
S1T5 Rm	FL, USA	Upper Tampa Bay	4-Feb-22	S1T5 - Old Tampa Bay	27.95788	-82.54663	*Ruppia maritima*	
S1T16 Tt	FL, USA	Upper Tampa Bay	4-Feb-22	S1T16 - Old Tampa Bay	27.88228	-82.62015	*Thalassia testudinum*	
S1T16 Sf	FL, USA	Upper Tampa Bay	4-Feb-22	S1T16 - Old Tampa Bay	27.88228	-82.62015	*Syringodium filiforme*	
S1T16 Rm	FL, USA	Upper Tampa Bay	4-Feb-22	S1T16 - Old Tampa Bay	27.88228	-82.62015	*Ruppia maritima*	
S1T16 Hw	FL, USA	Upper Tampa Bay	4-Feb-22	S2T2 - Hillsborough Bay	27.82347	-82.39305	*Halodule wrightii*	
S1T16 Rm	FL, USA	Upper Tampa Bay	4-Feb-22	S2T2 - Hillsborough Bay	27.82347	-82.39305	*Ruppia maritima*	
S3T8 Tt 2021	FL, USA	Upper Tampa Bay	26-Jan-22	S3T8 - Lassing Park	27.75327	-82.63035	*Thalassia testudinum*	
S3T8 Sf 2021	FL, USA	Upper Tampa Bay	26-Jan-22	S3T8 - Lassing Park	27.75327	-82.63035	*Syringodium filiforme*	
S3T8 Rm 2021	FL, USA	Upper Tampa Bay	26-Jan-22	S3T8 - Lassing Park	27.75327	-82.63035	*Ruppia maritima*	
S3T8 Tt 2023	**FL, USA**	**Upper Tampa Bay**	**3-Oct-23**	**S3T8 - Lassing Park**	**27.75327**	**-82.63035**	***Thalassia testudinum***	**OR879052**
S3T8 Sf 2023	FL, USA	Upper Tampa Bay	26-Jan-22	S3T8 - Lassing Park	27.75327	-82.63035	*Syringodium filiforme*	
S3T8 Rm 2023	FL, USA	Upper Tampa Bay	26-Jan-22	S3T8 - Lassing Park	27.75327	-82.63035	*Ruppia maritima*	
S3T13 Tt	FL, USA	Upper Tampa Bay	4-Feb-22	S3T13 - Apollo Beach	27.7566	-82.44758	*Thalassia testudinum*	
S3T13 Hw	FL, USA	Upper Tampa Bay	4-Feb-22	S3T13 - Apollo Beach	27.7566	-82.44758	*Halodule wrightii*	
S4T5 Tt	FL, USA	Lower Tampa Bay	26-Jan-22	S4T5 - Terra Ceia Bay	27.54417	-82.60615	*Thalassia testudinum*	
S4T5 Sf	FL, USA	Lower Tampa Bay	26-Jan-22	S4T5 - Terra Ceia Bay	27.54417	-82.60615	*Syringodium filiforme*	

S4T5 Hw	FL, USA	Lower Tampa Bay	26-Jan-22	S4T5 - Terra Ceia Bay	27.54417	-82.60615	*Halodule wrightii*	
S4T5 Rm	FL, USA	Lower Tampa Bay	26-Jan-22	S4T5 - Terra Ceia Bay	27.54417	-82.60615	*Ruppia maritima*	
S4T10 Tt	FL, USA	Lower Tampa Bay	26-Jan-22	S4T10 - Lower Tampa Bay	27.52134	-82.65712	*Thalassia testudinum*	
S4T10 Sf	FL, USA	Lower Tampa Bay	26-Jan-22	S4T10 - Lower Tampa Bay	27.52134	-82.65712	*Syringodium filiforme*	
S4T10 Hw	FL, USA	Lower Tampa Bay	26-Jan-22	S4T10 - Lower Tampa Bay	27.52134	-82.65712	*Halodule wrightii*	
S4T10 Rm	FL, USA	Lower Tampa Bay	26-Jan-22	S4T10 - Lower Tampa Bay	27.52134	-82.65712	*Ruppia maritima*	
BCB11 Tt	FL, USA	Lower Tampa Bay	26-Jan-22	BCB11 - Boca Ciega Bay	27.72243	-82.69762	*Thalassia testudinum*	
BCB11 Sw	FL, USA	Lower Tampa Bay	26-Jan-22	BCB11 - Boca Ciega Bay	27.72243	-82.69762	*Syringodium filiforme*	
BCB11 Hw	FL, USA	Lower Tampa Bay	26-Jan-22	BCB11 - Boca Ciega Bay	27.72243	-82.69762	*Halodule wrightii*	
27	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58404412**	**-82.61651603**	***Thalassia testudinum***	**OR827699**
28	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58404505**	**-82.61643497**	***Thalassia testudinum***	**PP430550**
29	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58404597**	**-82.61635392**	***Thalassia testudinum***	**OR827693**
37	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58397095**	**-82.61659599**	***Thalassia testudinum***	**OR827703**
48	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.5838997**	**-82.61651385**	***Thalassia testudinum***	**OR827694**
49	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58390062**	**-82.61643288**	***Thalassia testudinum***	**PP430551**
50	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58390163**	**-82.61635182**	***Thalassia testudinum***	**PP430552**
51	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58390255**	**-82.61627077**	***Thalassia testudinum***	**PP430553**
52	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58390347**	**-82.6161898**	***Thalassia testudinum***	**PP430554**
53	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.5839044**	**-82.61610875**	***Thalassia testudinum***	**OR827695**
54	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.5839054**	**-82.61602778**	***Thalassia testudinum***	**PP430555**

62	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.5838303	-82.61626977	*Thalassia testudinum*	PP430556
64	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58383223	-82.61610774	*Thalassia testudinum*	OR827702
66	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58383407	-82.61594564	*Thalassia testudinum*	OR827697
67	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58383508	-82.61586467	*Thalassia testudinum*	PP430557
74	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58375813	-82.61626868	*Thalassia testudinum*	PP430558
75	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58375905	-82.61618771	*Thalassia testudinum*	PP430559
77	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58376098	-82.6160256	*Thalassia testudinum*	PP430560
78	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.5837619	-82.61594463	*Thalassia testudinum*	PP430561
79	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58376283	-82.61586358	*Thalassia testudinum*	PP430562
80	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58376383	-82.61578253	*Thalassia testudinum*	PP430563
84	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58368689	-82.61618662	*Thalassia testudinum*	PP430564
88	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58369066	-82.61586249	*Thalassia testudinum*	PP430565
93	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58361564	-82.61610456	*Thalassia testudinum*	OR827698
96	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58361849	-82.61586148	*Thalassia testudinum*	PP430566
97	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58361941	-82.61578043	*Thalassia testudinum*	OR827705
101	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58354439	-82.61602242	*Thalassia testudinum*	OR879054
105	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58354817	-82.61569837	*Thalassia testudinum*	OR827704
119	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58340475	-82.61561522	*Thalassia testudinum*	PP430548
128	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58325941	-82.6156941	*Thalassia testudinum*	OR879053
132	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58318816	-82.61561204	*Thalassia testudinum*	PP430549

A1	**FL, USA**	**Dry Tortugas National Park**	**20-May-22**	**Key West**	**24.581857**	**-81.799379**	***Thalassia testudinum***	**PP430567**
A2	**FL, USA**	**Dry Tortugas National Park**	**20-May-22**	**Key West**	**24.581865**	**-81.799295**	***Thalassia testudinum***	**PP430568**
L1	FL, USA	Dry Tortugas National Park	19-May-22	Marquesas Key	24.560963	-82.125589	*Thalassia testudinum*	
L2	**FL, USA**	**Dry Tortugas National Park**	**19-May-22**	**Marquesas Key**	**24.560981**	**-82.12555**	***Thalassia testudinum***	**PP430569**
Q1	FL, USA	Dry Tortugas National Park	17-May-22	Garden Key	24.626714	-82.874075	*Thalassia testudinum*	**PP430570**
Q2	FL, USA	Dry Tortugas National Park	17-May-22	Garden Key	24.626673	-82.874162	*Thalassia testudinum*	**PP430571**
R	**FL, USA**	**Dry Tortugas National Park**	**16-May-22**	**Bush Key**	**24.626357**	**-82.868817**	***Thalassia testudinum***	**OR879056**
Y	FL, USA	Dry Tortugas National Park	18-May-22	Loggerhead Key	24.635007	-82.921341	*Thalassia testudinum*	
Panacea	**FL, USA**	**Florida Panhandle**	**10-May-23**	**Panacea**	**29.99911**	**-84.35056**	***Thalassia testudinum***	**OR879055**
MAIRIM	Puerto Rico	Jobos Bay National Estuarine Research Reserve	1-Feb-23	MAIRIM Site	Not available	Not available	*Halophila stipulacea*	
CPL	Puerto Rico	Jobos Bay National Estuarine Research Reserve	1-Feb-23	Cayo Puerca Lulu	Not available	Not available	*Halophila stipulacea*	
Z1	VA, USA	York River	2-Aug-17	York River State Park	Not available	Not available	*Zostera marina*	
Z2	MA, USA	West Falmouth Harbor	Not available	West Falmouth Harbor	Not available	Not available	*Zostera marina*	
Z3	AK, USA	Sitka	Not available	Sitka	Not available	Not available	*Zostera marina*	
Z4	New Zealand	Not available	Dec-22	New Zealand	Not available	Not available	*Zostera marina*	
Z5	Kalmar, Sweden	Not available	Not available	Kalmar	Not available	Not available	*Zostera marina*	

A Maximum likelihood tree with 1,000 bootstrap replicates

B Pairwise nucleotide sequence identities