Limited potexvirus diversity in seagrasses

1.1 Author names

Shen Jean Lim[1*] (https://orcid.org/0000-0003-4578-5318)

Karyna Rosario[1] (https://orcid.org/0000-0001-9847-4113)

Meredith E. Kernbach[1] (https://orcid.org/0000-0001-8161-057X)

Anthony J. Gross[1] (https://orcid.org/0000-0001-6392-5121)

Bradley T. Furman[2] (https://orcid.org/0000-0002-9284-8388)

Mya Breitbart[1] (https://orcid.org/0000-0003-3210-2899)

1.2 Affiliation(s)

[1] College of Marine Science, University of South Florida, St Petersburg, FL, USA

[2] Florida Fish and Wildlife Conservation Commission, Florida Fish and Wildlife Research Institute, Saint Petersburg, FL, USA

1.3 Corresponding author and email address

Shen Jean Lim (jeanlim@usf.edu)

1.4 Keywords

Potexvirus, primer, PCR, seagrass, marine, plant, *Alphaflexivirdae*

1.5 Repositories:

All sequence data have been deposited into NCBI GenBank under the accession numbers OR827689-OR827705, OR854648, OR863396, OR879052-OR879056, and PP430548-PP430571.

2. Abstract

Turtlegrass virus X, which infects the seagrass *Thalassia testudinum*, is the only potexvirus known to infect marine flowering plants. We investigated potexvirus distribution in seagrasses using a degenerate reverse transcription polymerase chain reaction (RT-PCR) assay originally designed to capture potexvirus diversity in terrestrial plants. The assay, which implements Potex-5 and Potex-2RC primers, successfully amplified a 584 nt RNA-dependent RNA polymerase (RdRp) fragment from TVX-infected seagrasses. Following validation, we screened 74 opportunistically collected, apparently healthy seagrass samples for potexviruses using this RT-PCR assay. The survey examined the host species *T. testudinum*, *Halodule wrightii*, *Halophila stipulacea*, *Syringodium filiforme*, *Ruppia maritima*, and *Zostera marina*. Potexvirus PCR products were successfully generated only from *T. testudinum* samples and phylogenetic analysis of sequenced PCR products revealed five distinct TVX sequence variants. Although the RT-PCR assay revealed limited potexvirus diversity in seagrasses, the expanded geographic distribution of TVX shown here emphasizes the importance of future studies to investigate *T. testudinum* populations across its native range and understand how the observed fine-scale genetic diversity affects host-virus interactions.

3. Impact statement

Potexviruses are widespread in terrestrial plants; however, the recent discovery of TVX in the seagrass *Thalassia testudinum* extends their host range to marine flowering plants. Here we use existing Potex-5 and Potex-2RC degenerate primers to explore potexvirus infections in several seagrass species. TVX sequence variants were detected in *T. testudinum* collected from the eastern Gulf of Mexico, uncovering previously unknown genetic diversity of this poorly understood virus.

4. Data summary

All sequence data are available in NCBI GenBank under the accession numbers OR827692-OR827705, OR854648, OR863396, OR879052-OR879056, and PP430548-PP430571. The authors confirm all supporting data, code and protocols have been provided within the article or through supplementary data files.

5. Introduction

Potexviruses are positive-sense single-stranded RNA (ssRNA) viruses from the family *Alphaflexiviridae* that infect many agronomically important flowering plants (angiosperms), including potato, tobacco, and tomato (1). The type member of this genus is *Potato virus X* (PVX), from which the genus name *Potexvirus* is derived (2). Potexviruses have a broad

geographic range that reflects the distribution of their hosts and the prevalence of global
commerce (2, 3). Although each potexvirus species has a limited natural host range, some
species can infect a wide range of host plants in experimental settings (1). Potexviruses are
transmitted mechanically through contaminated agricultural equipment or plant-to-plant
contact (1-3). Transmission via seed and aphid vectors and non-specific transmission through
chewing insects has been reported but is less common (1-3). Most potexviruses cause
persistent, asymptomatic infections or mild mosaic disease, while some cause dwarf,
necrotic, or ringspot symptoms in their natural hosts (1, 2). Notably, PVX can cause severe
symptoms when its plant host is co-infected with potyviruses (4-6).

In addition to infecting terrestrial plants, potexviruses are also capable of infecting
seagrasses, the only marine angiosperms. A novel potexvirus species, *Turtlegrass virus X*
(TVX), was discovered in apparently healthy *Thalassia testudinum* seagrass samples collected
from Terra Ceia Aquatic Preserve, Tampa Bay, Florida, USA (7). The identification of TVX
indicates that an important aspect of potexvirus diversity has been overlooked by focusing
only on terrestrial plants. Degenerate primers used in reverse transcriptase polymerase
chain reaction (RT-PCR) assays, such as Potex-5 and Potex-2RC (8), have proven useful in
detecting diverse potexviruses, but have not been tested in aquatic habitats. This study
applied the Potex-5 and Potex-2RC primer pair in a RT-PCR assay to explore potexvirus
diversity in samples from six seagrass host species.

75 6. Methods

Primer validation. Potex-5/Potex-2RC were tested on cDNA samples each synthesized from 1
µg RNA extracted from *Portulaca* sp. infected with *Alternanthera mosaic virus* (AltMV) and
Opuntia sp. with a mixed infection of potexviruses, as well as cDNA pooled from *T.*
testudinum samples known to contain TVX, based on the TVX-specific RT-PCR assay
described by Van Bogaert *et al.* 2019 (7). The PCR reaction contained 0.48 µM of each
primer, 2 µL cDNA template, 1 µL GC enhancer, and 1X AmpliTaq Gold™ 360 Master Mix
(Applied Biosystems™, Waltham, MA, USA) in a 25 µL reaction volume. PCR was performed
under the following conditions: initial denaturation at 95°C for 10 minutes, 40 cycles of
denaturation at 95°C for 30 seconds, annealing at 51.5°C (as published in van der Vlugt and
Berendsen (8)) for 30 seconds, extension at 72°C for 1 minute, followed by elongation at
72°C for 10 minutes and cooling at 11°C. The PCR product was visualized following gel
electrophoresis on a 1% (wt/vol) agarose gel stained with ethidium bromide. All PCR
reactions, except for the no template control, yielded visible bands. All PCR products were
purified using the Zymoclean Gel DNA Recovery Kit (Irvine, CA, USA), quantified using the
Qubit™ DNA high sensitivity (HS) assay (Invitrogen™, Waltham, MA, USA), and Sanger
sequenced bidirectionally by Eurofins Genomics (Louisville, KY, USA).

RT-PCR. Total RNA extraction was performed on 30-100 mg of leaves from multiple shoots

pooled by seagrass species and collection site (**Table S1**) using Zymo Research's (ZR) Quick-

RNA™ Plant Miniprep kit. Each pooled seagrass sample was homogenized in a BashingBead

Lysis Tube containing 2 mm ceramic beads and 800 µL RNA lysis buffer (provided in the kit)

for 5 minutes at maximum speed using a Fisherbrand™ Bead Mill 4 Homogenizer (Fisher

Scientific, Waltham, MA, USA). Tissue homogenates were centrifuged at maximum speed

(21,130 *g*) for 1 minute and RNA was extracted from the total volume (~800 µL) of the

supernatant. To ensure successful RNA extraction, after each round of extraction, a random

subset of RNA samples was quantified using the Qubit™ RNA high sensitivity (HS) assay

(Invitrogen™). From each sample, cDNA was synthesized from 8 µL RNA using the

SuperScript™ IV First-Strand Synthesis System (Invitrogen™) and following manufacturer's

instructions for random hexamers. PCR amplification was performed on each cDNA sample,

followed by gel electrophoresis, PCR product purification, DNA quantification, and Sanger

sequencing, using the methods described above.

Amplicon sequence analysis. Amplicon sequences were first compared with sequences from

NCBI's nucleotide (nt) and non-redundant protein (nr) collections (9) and the TVX genome

(7) using the megablast and/or blastx programs on NCBI's BLAST® server (10). Sequence

reads from each sample were then mapped to the Potex-5/Potex-2RC amplicon region (with

forward and reverse primer sequences removed) in the TVX genome (7) using the "Map

Sanger Reads to Reference" function implemented in Unipro UGENE v48.1 (11), with a

trimming quality threshold of 20 and mapping minimum similarity of 70%.

Electropherograms of mapped reads were manually inspected to remove primer sequences

and low-quality bases and to resolve ambiguous bases.

Sequences from 42 Potex-5/Potex-2RC PCR products amplified from seagrass and terrestrial

plant samples described above were aligned with corresponding amplicon regions (excluding

primer sequences) extracted from the reference genomes of TVX (NC_040644) (7), *Bamboo*

mosaic virus (NC_001642) (12), *Foxtail mosaic virus* (NC_001483) (13), AltMV (OR607766),

PVX (NC_011620) (14), and *Lolium latent virus* (NC_010434) (15) retrieved from the NCBI

Virus database (16). Multiple sequence alignment was performed using the L-INS-I and --

adjustdirectionaccurately options in MAFFT v7.5.08 (17), then trimmed to 514 nt to remove

leading and trailing gaps. Pairwise sequence identities were calculated from the trimmed

alignment using Clustal Omega v1.2.3 (18) with the --distmat-out, --percent-id, and --full

options. The pairwise sequence identity matrix was clustered, based on Euclidean distances,

using the hierarchical clustering with complete linkage method implemented in the

heatmap.2 function of the gplots R package (https://github.com/talgalili/gplots).

For phylogenetic analysis, Molecular Evolutionary Genetics Analysis (MEGA) 11 (19)
identified the best model for the trimmed alignment to be the Hasegawa-Kishino-Yano
model (20) with a discrete Gamma distribution (HKY+G). Using this substitution model, a
maximum likelihood tree with 1,000 bootstrap replicates was constructed from the
alignment using MEGA11 (19). The tree was condensed using MEGA11 (19) to eliminate
interior branches with <50% bootstrap support. The final tree was visualized and annotated
using FigTree v1.4.4 (http://tree.bio.ed.ac.uk/software/Figtree/).

7. Results

The Potex-5/Potex-2RC primer pair successfully amplified TVX from *T. testudinum* samples
collected from Terra Ceia Aquatic Preserve, AltMV infecting *Portulaca* sp., and an
uncharacterized potexvirus infecting *Opuntia* sp. (8). A RT-PCR assay using Potex-5/Potex-2RC
was subsequently performed on seagrass leaves opportunistically collected from separate
sampling efforts, including 1) *T. testudinum*, *H. wrightii*, *S. filiforme*, and *R. maritima* from
Tampa Bay on Jan 26th or Feb 4th, 2022; 2) *T. testudinum* from the Dry Tortugas National
Park, Florida, USA collected between May 16th-20th, 2022; 3) *T. testudinum* from a systematic
seagrass survey at Terra Ceia Aquatic Preserve on August 1st, 2022; and 4) *T. testudinum*, *H.
wrightii*, and *S. filiforme* samples collected from Tampa Bay seagrass site S3T8 (Lassing Park)
on October 3rd, 2023 (**Figure 1** and **Table S1**). Seagrass samples (**Table S1**) were also received
from other research groups, including *T. testudinum* from Panacea (Florida, USA), *Zostera
marina* from York River (Virginia, USA), West Falmouth Harbor (Massachusetts, USA), Sitka
(Alaska, USA), New Zealand, and Kalmar (Sweden), and *Halophila stipulacea* from Jobos Bay
National Estuarine Research Reserve (Puerto Rico, USA). PCR products were successfully
amplified and sequenced from 40 *T. testudinum* samples from Florida sites, including Terra
Ceia Aquatic Preserve, Tampa Bay seagrass sites S1T5 and S3T8 (Lassing Park), Panacea
located in the Florida Panhandle, and Florida Keys sites including Bush Key, Garden Key,
Marquesas Key, and Key West (**Figure 1**). Aside from *T. testudinum*, no other seagrass species
produced PCR products.

Figure 1. Map showing locations of seagrass sites in (A) Florida, (B) Tampa Bay, and (C) Terra

Ceia Aquatic Preserve where samples were collected for RT-PCR. Purple markers in (A) and

(B) denote sites where TVX products were amplified from *T. testudinum* with the Potex-

5/Potex-2RC primers, whereas samples from sites with orange markers tested negative.

Marker shapes in (C) represent potexvirus subclades found in Terra Ceia Aquatic Preserve

sampling sites and correspond to **Figure 2** and **Figure 3** (negative samples not shown).

Metadata of seagrass samples are described in **Table S1**.

Phylogenetic analysis placed all Potex-5/Potex-2RC amplicons from seagrasses in the same

clade as the RdRp sequence fragment from the TVX genome (7) with 100% bootstrap

confidence (**Figure 2**). TVX sequences from Terra Ceia Aquatic Preserve formed three

subclades. Average pairwise nucleotide sequence identities were 95.8±1.4% within each

subclade and 79.3±2.9% between each subclade (**Figure 3**). Two Terra Ceia Aquatic Preserve

subclades, including the subclade containing the amplicon region from the TVX genome,

were sister groups to each other (**Figure 2**), sharing 84.6±1% average pairwise nucleotide

sequence identities (**Figure 3**). Other TVX sequences from Key West, Marquesas Key, and

Panacea formed a subclade with 99.6±0.2% average pairwise nucleotide sequence identities,

while sequences from Garden Key and Bush Key formed another subclade with 89.3±0.9%%

average pairwise nucleotide sequence identities (**Figure 2** and **Figure 3**). Potexviruses that

infect terrestrial grasses (*Foxtail mosaic virus* and *Bamboo mosaic virus*) formed sister clades

to the TVX-containing clade (**Figure 2**).

73.7% ■ Terra Ceia Aquatic Preserve and *Turtlegrass virus X* [NC_040644]

84.1%

98.9% ▲ Terra Ceia Aquatic Preserve

S1T5 Old Tampa Bay [OR827692]

51.4%

S3T8 Lassing Park [OR879052]

94% Key West, Marquesas Key [PP430567-PP430569] and Panacea [OR879055]

100%

95.9% ♦ Terra Ceia Aquatic Preserve

83.8% Garden Key [PP430570-PP430571] and Bush Key [OR879056]

68.1%

Foxtail mosaic virus [NC_001483]

Bamboo mosaic virus [NC_001642]

97%

Potato virus X [NC_011620]

53.3% Uncharacterized potexvirus [OR863396]

Alternanthera mosaic virus [OR854648]

99.9%

Alternanthera mosaic virus isolate DSMZ PV-0849 [OR607766]

Lolium latent virus [NC_010434]

2.0

Figure 2. Condensed bootstrap consensus tree of TVX nucleotide sequences amplified from seagrasses (purple text), in relation to potexvirus

sequences amplified from terrestrial plants (brown text) and potexvirus RdRp sequences from reference genomes (black text). The NCBI

accession number for each sequence is indicated in square brackets. Accession numbers for sequences from Terra Ceia Aquatic Preserve

samples are listed in **Figure 3**. The tree was constructed by MEGA11 from a 514 nt conserved region in the multiple sequence alignment.

Branch lengths indicate the number of substitutions per site. Bootstrap values on tree nodes indicate the percentage of trees, based on 1,000

replicates, in which sequences within a node were clustered together. Shapes beside bootstrap values of Terra Ceia Aquatic Preserve subclades

correspond to shapes in **Figure 1C**. *Lolium latent virus* was used as an outgroup.

Figure 3. Histogram (upper right corner), heatmap, and dendrogram (hierarchical clustering)

of pairwise nucleotide sequence identities between TVX sequences amplified from

seagrasses (purple text), potexvirus sequences amplified from terrestrial plants (brown text),

and potexvirus RdRp sequences from reference genomes (black text). The NCBI accession

number for each sequence is indicated in square brackets. Marker shapes preceding Terra

Ceia Aquatic Preserve seagrass sample names correspond to shapes in **Figure 1C**.

Abbreviations: TVX, *Turtlegrass virus X*; AltMV, *Alternanthera mosaic virus*; PVX, *Potato virus*

X; LolV, *Lolium latent virus*; FoMV, *Foxtail mosaic virus*; BaMV, *Bamboo mosaic virus*.

8. Discussion

In this study, we validated the ability of the potexvirus degenerate primers, Potex-5/Potex-2RC, published by van der Vlugt and Berendsen (8), to amplify TVX-infected seagrass samples and potexvirus-infected terrestrial plant samples. Potex-5/Potex-2RC were used in a RT-PCR survey to test six seagrass species for potexviruses. Among the seagrass species examined, potexvirus infection was only detected in *T. testudinum*, the species from which TVX was identified. Notably, the only potexviruses identified in this study were related to TVX, suggesting limited potexvirus diversity in marine angiosperms. RT-PCR analysis revealed *T. testudinum* as a natural host for TVX and confirmed TVX infection in eastern Gulf of Mexico seagrass meadows, including those in the northwest (Panacea), west-central (Tampa Bay), and southwest (Florida Keys) Florida. Three distinct TVX sequence variants were observed within a single seagrass meadow at Terra Ceia Aquatic Preserve, along with two additional TVX sequence variants in other Tampa Bay sites, Florida Keys, and Panacea.

Among all sites sampled in this study, the most extensive TVX survey was conducted within the Terra Ceia Aquatic Preserve. Systematic RT-PCR surveys at other sites will be useful in elucidating potexvirus diversity in seagrasses and in resolving potexvirus sequence variants into species or strains. The sampling scheme in this study was limited to *T. testudinum* populations along the eastern Gulf of Mexico. However, *T. testudinum* has a widespread geographic range across the Western Atlantic Ocean, with high genetic diversity that is generally partitioned into two clusters corresponding to the Gulf of Mexico or the Caribbean phylogeographic regions (21). Future potexvirus surveys covering the total distribution range of *T. testudinum*, the related species *T. hemprichii*, and other seagrass species have the potential to uncover relationships between potexvirus diversity, host diversity, and their phylogeographic distributions.

9. Author statements

9.1   Author contributions

SJL: Data curation, Formal analysis, Investigation, Methodology, Software, Supervision, Visualization, Writing - original draft

KRC: Conceptualization, Formal analysis, Funding acquisition, Writing – review & editing.

MEK: Investigation, Writing – review & editing

AJG: Investigation, Writing - review & editing

BTF: Funding acquisition, Investigation, Writing - review & editing

MB: Conceptualization, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Resources, Supervision, Validation, Writing - review & editing.

9.2 Conflicts of interest

The author(s) declare that there are no conflicts of interest.

9.3 Funding information

This project was supported by award OCE-2219547 from the National Science Foundation to MB, KRC, and BTF and a Proposal Enhancement Grant from the University of South Florida Internal Awards Program to MB. This research was made possible by support from the Florida Institute of Oceanography for collection of the Dry Tortugas samples.

9.4 Ethical approval

N/A

9.5 Consent for publication

N/A

9.6 Acknowledgements

Fieldwork was conducted under permit no. FKNMS-2022-026 issued by the Florida Keys National Marine Sanctuary, permit no. DRTO-2022-SCI-0004 issued through the U.S. Department of the Interior National Park Service and permit no. SAL-19-1172A-SR issued through the Florida Fish and Wildlife Conservation Commission. We acknowledge the contributions of Luis Alvarado-Marchena, Victoria Congdon, Keith Keel, Isabella Ritchie, Taylor Little, Jessye Kirkham, and Mike Wheeler in field work and sample collection at these sites. Isabella Ritchie provided cDNAs from *T. testudinum* samples from Terra Ceia Aquatic Preserve. Salvador P Lopez, Jr and Scott Adkins from the Agricultural Research Service, United States Horticultural Research Laboratory, U.S. Department of Agriculture (USDA), Fort Pierce, FL, USA provided cDNAs from potexvirus-infected plant samples. We thank Maya Groner, Ian Hewson, Karin Holmfeldt, Natalia Lopez Figueroa, Carolyn Malmstrom, Jordan Rede, Cypress Rudloe, and Jack Rudloe for providing seagrass samples from other sites.

10. References

1.   Ryu KH, Hong JS. Potexvirus. In: Mahy BWJ, Van Regenmortel MHV, editors. Encyclopedia of Virology (Third Edition). Oxford: Academic Press; 2008. p. 310-3.

2.   Verchot J. *Potato virus X*: A global potato-infecting virus and type member of the *Potexvirus* genus. Mol Plant Pathol. 2022;23(3):315-20.

3.   AbouHaidar MG, Gellatly D. Potexviruses. In: Granoff A, Webster RG, editors. Encyclopedia of Virology (Second Edition). Oxford: Elsevier; 1999. p. 1364-8.

4.   Rochow WF, Ross AF. Virus multiplication in plants doubly infected by potato viruses X and Y. Virology. 1955;1(1):10-27.

268   5.   Pacheco R, Garcia-Marcos A, Barajas D, Martianez J, Tenllado F. PVX-potyvirus synergistic
269   infections differentially alter microRNA accumulation in *Nicotiana benthamiana*. Virus Res.
270   2012;165(2):231-5.
271
272   6.   Pollari M, Sipari N, Poque S, Himanen K, Makinen K. Effects of poty-potexvirus synergism on
273   growth, photosynthesis and metabolite status of *Nicotiana benthamiana*. Viruses. 2022;15(1):121.
274
275   7.   Van Bogaert N, Rosario K, Furman BT, Hall MO, Greco AM, Breitbart M. Discovery of a novel
276   potexvirus in the seagrass *Thalassia testudinum* from Tampa Bay, Florida. Limnol Oceanogr Lett.
277   2019;4(1):1-8.
278
279   8.   van der Vlugt RAA, Berendsen M. Development of a general potexvirus detection method.
280   European Journal of Plant Pathology. 2002;108(4):367-71.
281
282   9.   Sayers EW, Beck J, Bolton EE, Bourexis D, Brister JR, Canese K, et al. Database resources of
283   the National Center for Biotechnology Information. Nucleic Acids Res. 2021;49(D1):D10-D7.
284
285   10.   Johnson M, Zaretskaya I, Raytselis Y, Merezhuk Y, McGinnis S, Madden TL. NCBI BLAST: a
286   better web interface. Nucleic Acids Res. 2008;36(Web Server issue):W5-9.
287
288   11.   Okonechnikov K, Golosova O, Fursov M, team U. Unipro UGENE: a unified bioinformatics
289   toolkit. Bioinformatics. 2012;28(8):1166-7.
290
291   12.   Lin NS, Lin BY, Lo NW, Hu CC, Chow TY, Hsu YH. Nucleotide sequence of the genomic RNA of
292   bamboo mosaic potexvirus. J Gen Virol. 1994;75 (9):2513-8.
293
294   13.   Bancroft JB, Rouleau M, Johnston R, Prins L, Mackie GA. The entire nucleotide sequence of
295   *Foxtail mosaic virus* RNA. J Gen Virol. 1991;72 (9):2173-81.
296
297   14.   Huisman MJ, Linthorst HJ, Bol JF, Cornelissen JC. The complete nucleotide sequence of *Potato
298   virus X* and its homologies at the amino acid level with various plus-stranded RNA viruses. J Gen
299   Virol. 1988;69 (Pt 8):1789-98.
300
301   15.   Vaira AM, Maroon-Lango CJ, Hammond J. Molecular characterization of *Lolium latent virus*,
302   proposed type member of a new genus in the family Flexiviridae. Arch Virol. 2008;153(7):1263-70.
303
304   16.   Brister JR, Ako-Adjei D, Bao Y, Blinkova O. NCBI viral genomes resource. Nucleic Acids Res.
305   2015;43(Database issue):D571-7.
306
307   17.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
308   improvements in performance and usability. Mol Biol Evol. 2013;30(4):772-80.
309
310   18.   Sievers F, Higgins DG. Clustal Omega for making accurate alignments of many protein
311   sequences. Protein Sci. 2018;27(1):135-45.
312
313   19.   Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis Version
314   11. Mol Biol Evol. 2021;38(7):3022-7.
315

316   20.   Hasegawa M, Kishino H, Yano T. Dating of the human-ape splitting by a molecular clock of
317   mitochondrial DNA. J Mol Evol. 1985;22(2):160-74.
318
319   21.   van Dijk K-j, Bricker E, van Tussenbroek BI, Waycott M. Range-wide population genetic
320   structure of the Caribbean marine angiosperm *Thalassia testudinum*. Ecol Evol. 2018;8(18):9478-90.
321

Sample ID	State, Country	Region	Collection Date	Site Description	Latitude	Longitude	Species	Accession
S1T5 Tt	**FL, USA**	**Upper Tampa Bay**	**4-Feb-22**	**S1T5 - Old Tampa Bay**	**27.95788**	**-82.54663**	***Thalassia testudinum***	**OR827692**
S1T5 Sf	FL, USA	Upper Tampa Bay	4-Feb-22	S1T5 - Old Tampa Bay	27.95788	-82.54663	*Syringodium filiforme*	
S1T5 Rm	FL, USA	Upper Tampa Bay	4-Feb-22	S1T5 - Old Tampa Bay	27.95788	-82.54663	*Ruppia maritima*	
S1T16 Tt	FL, USA	Upper Tampa Bay	4-Feb-22	S1T16 - Old Tampa Bay	27.88228	-82.62015	*Thalassia testudinum*	
S1T16 Sf	FL, USA	Upper Tampa Bay	4-Feb-22	S1T16 - Old Tampa Bay	27.88228	-82.62015	*Syringodium filiforme*	
S1T16 Rm	FL, USA	Upper Tampa Bay	4-Feb-22	S1T16 - Old Tampa Bay	27.88228	-82.62015	*Ruppia maritima*	
S1T16 Hw	FL, USA	Upper Tampa Bay	4-Feb-22	S2T2 - Hillsborough Bay	27.82347	-82.39305	*Halodule wrightii*	
S1T16 Rm	FL, USA	Upper Tampa Bay	4-Feb-22	S2T2 - Hillsborough Bay	27.82347	-82.39305	*Ruppia maritima*	
S3T8 Tt 2021	FL, USA	Upper Tampa Bay	26-Jan-22	S3T8 - Lassing Park	27.75327	-82.63035	*Thalassia testudinum*	
S3T8 Sf 2021	FL, USA	Upper Tampa Bay	26-Jan-22	S3T8 - Lassing Park	27.75327	-82.63035	*Syringodium filiforme*	
S3T8 Rm 2021	FL, USA	Upper Tampa Bay	26-Jan-22	S3T8 - Lassing Park	27.75327	-82.63035	*Ruppia maritima*	
S3T8 Tt 2023	**FL, USA**	**Upper Tampa Bay**	**3-Oct-23**	**S3T8 - Lassing Park**	**27.75327**	**-82.63035**	***Thalassia testudinum***	**OR879052**
S3T8 Sf 2023	FL, USA	Upper Tampa Bay	26-Jan-22	S3T8 - Lassing Park	27.75327	-82.63035	*Syringodium filiforme*	
S3T8 Rm 2023	FL, USA	Upper Tampa Bay	26-Jan-22	S3T8 - Lassing Park	27.75327	-82.63035	*Ruppia maritima*	
S3T13 Tt	FL, USA	Upper Tampa Bay	4-Feb-22	S3T13 - Apollo Beach	27.7566	-82.44758	*Thalassia testudinum*	
S3T13 Hw	FL, USA	Upper Tampa Bay	4-Feb-22	S3T13 - Apollo Beach	27.7566	-82.44758	*Halodule wrightii*	
S4T5 Tt	FL, USA	Lower Tampa Bay	26-Jan-22	S4T5 - Terra Ceia Bay	27.54417	-82.60615	*Thalassia testudinum*	
S4T5 Sf	FL, USA	Lower Tampa Bay	26-Jan-22	S4T5 - Terra Ceia Bay	27.54417	-82.60615	*Syringodium filiforme*	

S4T5 Hw	FL, USA	Lower Tampa Bay	26-Jan-22	S4T5 - Terra Ceia Bay	27.54417	-82.60615	*Halodule wrightii*	
S4T5 Rm	FL, USA	Lower Tampa Bay	26-Jan-22	S4T5 - Terra Ceia Bay	27.54417	-82.60615	*Ruppia maritima*	
S4T10 Tt	FL, USA	Lower Tampa Bay	26-Jan-22	S4T10 - Lower Tampa Bay	27.52134	-82.65712	*Thalassia testudinum*	
S4T10 Sf	FL, USA	Lower Tampa Bay	26-Jan-22	S4T10 - Lower Tampa Bay	27.52134	-82.65712	*Syringodium filiforme*	
S4T10 Hw	FL, USA	Lower Tampa Bay	26-Jan-22	S4T10 - Lower Tampa Bay	27.52134	-82.65712	*Halodule wrightii*	
S4T10 Rm	FL, USA	Lower Tampa Bay	26-Jan-22	S4T10 - Lower Tampa Bay	27.52134	-82.65712	*Ruppia maritima*	
BCB11 Tt	FL, USA	Lower Tampa Bay	26-Jan-22	BCB11 - Boca Ciega Bay	27.72243	-82.69762	*Thalassia testudinum*	
BCB11 Sw	FL, USA	Lower Tampa Bay	26-Jan-22	BCB11 - Boca Ciega Bay	27.72243	-82.69762	*Syringodium filiforme*	
BCB11 Hw	FL, USA	Lower Tampa Bay	26-Jan-22	BCB11 - Boca Ciega Bay	27.72243	-82.69762	*Halodule wrightii*	
27	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58404412**	**-82.61651603**	***Thalassia testudinum***	**OR827699**
28	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58404505**	**-82.61643497**	***Thalassia testudinum***	**PP430550**
29	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58404597**	**-82.61635392**	***Thalassia testudinum***	**OR827693**
37	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58397095**	**-82.61659599**	***Thalassia testudinum***	**OR827703**
48	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.5838997**	**-82.61651385**	***Thalassia testudinum***	**OR827694**
49	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58390062**	**-82.61643288**	***Thalassia testudinum***	**PP430551**
50	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58390163**	**-82.61635182**	***Thalassia testudinum***	**PP430552**
51	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58390255**	**-82.61627077**	***Thalassia testudinum***	**PP430553**
52	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.58390347**	**-82.6161898**	***Thalassia testudinum***	**PP430554**
53	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.5839044**	**-82.61610875**	***Thalassia testudinum***	**OR827695**
54	**FL, USA**	**Lower Tampa Bay**	**1-Aug-22**	**Terra Ceia Aquatic Preserve**	**27.5839054**	**-82.61602778**	***Thalassia testudinum***	**PP430555**

62	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.5838303	-82.61626977	*Thalassia testudinum*	PP430556
64	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58383223	-82.61610774	*Thalassia testudinum*	OR827702
66	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58383407	-82.61594564	*Thalassia testudinum*	OR827697
67	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58383508	-82.61586467	*Thalassia testudinum*	PP430557
74	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58375813	-82.61626868	*Thalassia testudinum*	PP430558
75	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58375905	-82.61618771	*Thalassia testudinum*	PP430559
77	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58376098	-82.6160256	*Thalassia testudinum*	PP430560
78	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.5837619	-82.61594463	*Thalassia testudinum*	PP430561
79	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58376283	-82.61586358	*Thalassia testudinum*	PP430562
80	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58376383	-82.61578253	*Thalassia testudinum*	PP430563
84	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58368689	-82.61618662	*Thalassia testudinum*	PP430564
88	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58369066	-82.61586249	*Thalassia testudinum*	PP430565
93	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58361564	-82.61610456	*Thalassia testudinum*	OR827698
96	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58361849	-82.61586148	*Thalassia testudinum*	PP430566
97	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58361941	-82.61578043	*Thalassia testudinum*	OR827705
101	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58354439	-82.61602242	*Thalassia testudinum*	OR879054
105	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58354817	-82.61569837	*Thalassia testudinum*	OR827704
119	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58340475	-82.61561522	*Thalassia testudinum*	PP430548
128	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58325941	-82.6156941	*Thalassia testudinum*	OR879053
132	FL, USA	Lower Tampa Bay	1-Aug-22	Terra Ceia Aquatic Preserve	27.58318816	-82.61561204	*Thalassia testudinum*	PP430549

A1	**FL, USA**	**Dry Tortugas National Park**	**20-May-22**	**Key West**	**24.581857**	**-81.799379**	***Thalassia testudinum***	**PP430567**
A2	**FL, USA**	**Dry Tortugas National Park**	**20-May-22**	**Key West**	**24.581865**	**-81.799295**	***Thalassia testudinum***	**PP430568**
L1	FL, USA	Dry Tortugas National Park	19-May-22	Marquesas Key	24.560963	-82.125589	*Thalassia testudinum*	
L2	**FL, USA**	**Dry Tortugas National Park**	**19-May-22**	**Marquesas Key**	**24.560981**	**-82.12555**	***Thalassia testudinum***	**PP430569**
Q1	FL, USA	Dry Tortugas National Park	17-May-22	Garden Key	24.626714	-82.874075	*Thalassia testudinum*	**PP430570**
Q2	FL, USA	Dry Tortugas National Park	17-May-22	Garden Key	24.626673	-82.874162	*Thalassia testudinum*	**PP430571**
R	**FL, USA**	**Dry Tortugas National Park**	**16-May-22**	**Bush Key**	**24.626357**	**-82.868817**	***Thalassia testudinum***	**OR879056**
Y	FL, USA	Dry Tortugas National Park	18-May-22	Loggerhead Key	24.635007	-82.921341	*Thalassia testudinum*	
Panacea	**FL, USA**	**Florida Panhandle**	**10-May-23**	**Panacea**	**29.99911**	**-84.35056**	***Thalassia testudinum***	**OR879055**
MAIRIM	Puerto Rico	Jobos Bay National Estuarine Research Reserve	1-Feb-23	MAIRIM Site	Not available	Not available	*Halophila stipulacea*	
CPL	Puerto Rico	Jobos Bay National Estuarine Research Reserve	1-Feb-23	Cayo Puerca Lulu	Not available	Not available	*Halophila stipulacea*	
Z1	VA, USA	York River	2-Aug-17	York River State Park	Not available	Not available	*Zostera marina*	
Z2	MA, USA	West Falmouth Harbor	Not available	West Falmouth Harbor	Not available	Not available	*Zostera marina*	
Z3	AK, USA	Sitka	Not available	Sitka	Not available	Not available	*Zostera marina*	
Z4	New Zealand	Not available	Dec-22	New Zealand	Not available	Not available	*Zostera marina*	
Z5	Kalmar, Sweden	Not available	Not available	Kalmar	Not available	Not available	*Zostera marina*	

324