ER-anchored protein sorting controls the fate of two proteasome activators

for intracellular organelle communication during proteotoxic stress

Gautier Langin[1,2], Margot Raffeiner[1], David Biermann[2], Mirita Franz-Wachtel[3], Daniela

Spinti[4], Frederik Börnke[4,5], Boris Macek[3], Suayib Üstün[1,*]

[1] Faculty of Biology & Biotechnology, Ruhr-University Bochum, Bochum, Germany

[2] Center for Plant Molecular Biology (ZMBP), University of Tübingen, Tübingen, Germany

[3] Institute for Cell Biology, Department of Quantitative Proteomics, University of Tübingen,

Tübingen, Germany Method section for MS/MS analysis

[4] Leibniz-Institute of Vegetable and Ornamental Crops (IGZ), Großbeeren, Germany

[5] Institute of Biochemistry and Biology, University of Potsdam, Potsdam, Germany

[*]Correspondence: suayb.uestuen@rub.de

Summary

Proteotoxic stress, characterized by the accumulation of damaged proteins, poses a significant

challenge to cellular homeostasis. To mitigate proteotoxicity eukaryotes employ the proteasome

that is regulated by proteasome activators, e.g. transcription factors that promote gene

expression of proteasome subunits. As proteotoxicity originates in different compartments, cells

need to perceive signals from various locations. Understanding which components integrate

signals to address proteotoxicity is essential to develop strategies to cope with proteotoxicity

but remain elusive. Here, we identify that the proteasome autoregulatory feedback loop acts as

a gatekeeper to facilitate the communication between nucleus and chloroplast. We reveal that

the ER-anchored protein sorting system (ERAPS) controls the proteasomal degradation or

nuclear translocation of proteasome activators NAC53 and NAC78. While both transcription

factors activate the proteasome gene expression, they repress photosynthesis-associated nuclear

genes during proteotoxicity. Collectively, our discoveries provide a novel conceptual

framework in which the proteasome autoregulatory feedback loop coordinates subcellular

proteostasis.

Introduction

Protein homeostasis, hereafter proteostasis, is defined as the synthesis of proteins by translating

genetic information, encoded in form of mRNA, into a sequence of amino acids and their

regulated degradation. This intimate ying-yang balance between newly synthesized proteins

and protein degradation is tightly regulated in all organisms, in order to develop or respond to

several environmental stimuli (Jayaraj et al., 2020; Orosa et al., 2020; Sampaio-Marques and

Ludovico, 2018). Various perturbations can alter this equilibrium, leading to proteotoxic stress

by disruption of normal proteostasis. This alteration can ultimately result in irreversible cellular

damage due to the accumulation of aberrant proteins, an outcome known as proteotoxicity.

Perturbations resulting in proteotoxic stress can range from ageing to pathological diseases such

as Parkinson and Alzheimer as well as infection by pathogens (Jayaraj et al., 2020; Langin et

al., 2023). The latter is a consequence of the onset and maintenance of defense responses during

microbial infections and microbes' ability to manipulate proteostasis for their own benefit

(Langin et al., 2023; Langin et al., 2020).

To mitigate deleterious effects due to excessive proteotoxic stress, cells employ various protein

quality control machineries. Strikingly, the response to proteotoxic stress is often controlled by

intricate regulatory feedback loops influenced by proteotoxic stress itself. One of the major

protein quality control machineries across the tree of life is the ubiquitin-proteasome system

(UPS). The ubiquitin-proteasome system controls proteostasis through selective elimination of

defective proteins as well as short-lived regulatory proteins (Raffeiner et al., 2023). Selective

protein degradation by the proteasome proceeds from the ligation of ubiquitin proteins to a

lysine residue within specific target proteins via the concerted action of E1, E2, and E3

enzymes. The ubiquitinated target is then recognized by the 26S proteasome for degradation.

The 26S proteasome itself is a conserved 2.5 MDa ATP-dependent protease complex composed

of a 20S core protease (CP) capped by 19S regulatory particles (RPs), that assists in ubiquitin

recognition and target protein unfolding. Recent findings highlight that the proteasome is a

highly dynamic structure, as more than 40 subunits participate in forming the 26S proteasome.

Changes in the subunit composition or the association of the 20S CP with other adaptors drive

flexibility of the 26S proteasome which can help to cope with proteotoxic stress (Albert et al.,

2020; Colberg et al., 2020; Peters et al., 2013; Tanaka et al., 2012). To guarantee this high

degree of structural flexibility and to immediately react to proteotoxic stress, all the constituents

of the 26S proteasome are under tight transcriptional control.

All eukaryotic kingdoms possess transcription factors (TFs) required for proteasome genes

transcriptional activation (Marshall and Vierstra, 2019). Yeast utilizes the C2H2 zinc-finger TF

Rpn4, mammals the bZIPs (Basic Leucine Zipper) Nrf1/2 and plants the pair of NACs (NAM,

ATAF and CUC) NAC53/NAC78 as a conserved mechanism to cope with proteotoxicity

although all TFs are evolutionary distinct. In yeast and animal systems, the proteasome

activators Rpn4 and Nrf1/2 have been shown to be UPS targets (Radhakrishnan et al., 2014;

Sha and Goldberg, 2014; Steffen et al., 2010; Ulasov et al., 2022; Xie and Varshavsky, 2001;

Xie, 2001). Rpn4 and Nrf2 are degraded in the cytosol while Nrf1, a tail-anchored (TA) protein,

is subjected to endoplasmic reticulum-associated degradation (ERAD). In the context of

exceeded proteasomal capacity, hereafter proteasomal stress, their degradation is impaired,

leading to their stabilization and subsequent nuclear translocation for proteasome gene

activation (Marshall and Vierstra, 2019). This transcriptional activation triggered by Rpn4 and

Nrf1/2 is mediated through binding of, respectively, the proteasome-associated control element

(PACE, [(A/G)GTGGC)]) or antioxidant response element (ARE, [(A/G)TGACTCAGC]

(Baird et al., 2017; Mannhaupt et al., 1999; Shirozu et al., 2015; Xie and Varshavsky, 2001).

This mechanism constitutes a negative feedback loop which allows systematic maintenance of

proteasome homeostasis. In plants, NAC78 has been shown to mediate transcriptional

activation of proteasome promoters through association with the proteasome regulatory cis

element (PRCE, [TGGGC], (Nguyen et al., 2013). In addition, loss of the NAC53/78 TF pair

rendered plants hypersensitive to proteasome inhibition due to its incapacity to activate

proteasome genes expression (Gladman et al., 2016). Therefore, a similar proteasome

regulatory feedback loop has been proposed in plants (Marshall and Vierstra, 2019); however,

to date, no evidence supports NAC53/78 proteasomal degradation.

Given the importance of proteasomal degradation in various cellular pathways, from cell cycle

to immune response (Langin et al., 2023; Ma and Liu, 2010; Wang et al., 2008; Wang et al.,

2010), it appears that the autoregulatory feedback loop is a central mechanism to precisely

integrate and coordinate stress responses through proteotoxic stress sensing. Indeed, a growing

body of evidence suggests that localized proteotoxic stress in compartments and organelles

causes the activation of respective proteasome-mediated degradation pathways to release the

stress. These mechanisms include ERAD, Endosome/Golgi-associated degradation (EGAD),

mitochondria-associated degradation (MAD) and chloroplast-associated degradation

(CHLORAD) (Christianson and Ye, 2014; Fonseca and Carvalho, 2019; Liao et al., 2020; Ling

et al., 2019). This implies a multi-layered role of the proteasome in governing subcellular

proteostasis. This aspect is particularly relevant for semi-autonomous organellar nuclear-

encoded proteins, as their regulation also involves cytosolic proteasomal degradation prior to

organellar import (Bragoszewski et al., 2017; Hristou et al., 2020). In turn, perturbations in the

turn-over of such proteins can lead to proteasomal stress. Indeed, Rpn4 is activated by

mitochondrial precursor protein accumulation to likely upregulate the proteasome for removal

of precursor proteins (Kramer et al., 2021). While for Nrf1/2 similar mechanisms remain

elusive, Nrf2 is activated upon mitochondrial stress and activates mitochondrial biogenesis and

precursors import (Gureev et al., 2019). This places the evolutionary distinct transcription

factors Rpn4, Nrf1/2 and NAC53/78 into the spotlight: Are they the main communicators

between organelles and the 26S proteasome to cope with proteotoxicity? As such, proteasome

activators could act as gatekeepers of the communication between the nucleus and energy-

producing organelles to maintain subcellular proteostasis. While to date it remains elusive

whether proteasome activators might act as such communicators, it would constitute a direct

way to coordinate 26S proteasome capacity and proteasome substrates to cope with proteotoxic

stress. It is also unknown how different signals from various organelles and compartments are

integrated to respond to overall proteotoxic stress.

Here we demonstrate that the plant proteasome autoregulatory feedback loop, driven by the

proteasome activators NAC53/78, functions as signal integrators by controlling the intracellular

communication between the nucleus and the chloroplast during proteotoxic stress. We show

that NAC53/78 are short-lived proteins regulated by the ERAD machinery through a

mechanism we termed ER-Anchored-Protein Sorting (ERAPS). At steady-state conditions

NAC53/78 are degraded via the HRD1-CDC48 axis. In contrast, proteotoxic stress caused by

pathogenic infection or pharmacological drug treatments stabilizes both transcription factors,

leading to their ERAPS-dependent nuclear translocation and subsequent transcriptional

reprogramming. We identified that NAC53/78 coordinate the transcriptional activation of

proteasome genes and repression of photosynthesis associated nuclear genes (PhANGs)

through association with a conserved PRCE-like cis-element. Strikingly, we have found that

the proteasome autoregulatory feedback loop is essential to maintain normal response to

subcellular proteostasis perturbation. Ultimately, we identified that the transcriptional signature

coordinated by NAC53/78 is a general feature in context of biotic interactions as wells as in

response to various developmental cues. Taken together, our findings provide a new conceptual

framework, in which an autoregulatory feedback loop shapes the response to environmental

stress: the integration of signals from different organelles and the role of proteasome activators

as gatekeepers of organelle communication constitutes a novel mechanism for ensuring the

global surveillance of cellular proteostasis in photosynthetic tissues.

NAC53/78 are central integrators of various proteotoxic stress conditions

Proteotoxicity is triggered by a wide range of physicochemical stresses. The induced proteotoxic stress is often due to a perturbation of local subcellular proteostasis. For instance, damage in energy-producing organelles such as chloroplast and mitochondria often lead to accumulation of proteins that are not imported into these organelles (Hristou et al., 2020; Kramer et al., 2021). How such signals from different cellular compartments are integrated is still not understood. Thus, we hypothesized that proteasome activators might play a role in integrating signals from different locations. To test our hypothesis, we subjected the double mutant *nac53-1 78-1,* known to be hypersensitive towards proteasome inhibition (Gladman et al., 2016), to various treatments previously reported to induce proteotoxic stress through specific mechanisms. We inhibited the proteasome (BTZ, (Gladman et al., 2016) and the ubiquitin-dependent segregase CDC48, a central player of multiple associated-degradation pathways (CB-5083, (Li et al., 2022; Marshall et al., 2019), and induced the ER unfolded protein response (DTT, TM, (Pastor-Cantizano et al., 2020). Additionally, we targeted semi-autonomous organelles such as chloroplasts and mitochondria (Lin, Cml and MV (Li et al., 2022; Wu et al., 2019), and the cytosolic chaperone Hsp90 that is essential for the organellar import of chloroplastic/mitochondrial precursors or proteasomal degradation of these precursors (GDA, (Wu et al., 2019). Ultimately, we used pathogenic infection with the bacterial strain *Pseudomonas syringeae pv. tomato DC3000* (*Pst*), known to impact cellular proteostasis through multiple mechanism notably via direct proteasome targeting (Üstün et al., 2016) (**Figure 1A**).

Consistent with previous studies, BTZ treatment led to *nac53-1 78-1* hypersensitivity resulting in a complete growth defect. Similarly, CDC48 inhibition resulted in a strong plant growth defect (**Figure 1B**), supporting the importance of the proteasome activators in the context of general proteotoxic stress, beyond direct proteasome targeting. Consistently, we observed increased sensitivity of the double mutant to ER stress inducers DTT and TM (**Figure 1B**). Strikingly, all other treatments resulted in *nac53-1 78-1* double mutant increased tolerance compared to WT plants (**Figure 1B**). In accordance with the hypersensitivity towards proteasome inhibition, bacterial infection also increased susceptibility of the *nac53-1 78-1* compared to Col-0 plants (**Figure 1C**).

Figure 1. NAC53/78 are central integrators of various proteotoxic stress conditions.
(A) The role of the 26S Proteasome in degradation of proteins from distinct subcellular compartments. Prime site
of action of the several treatments used in this study: Blunt head arrows indicate enzymatic activity suppression;
pointed head arrows indicate trigger of substrate accumulation.
(B) Fresh weight of seedlings grown under the indicated treatments at 10-14 day after germination (dag). Fresh
weight is expressed as a percentage of Col-0 mock conditions. Boxplots colors refer to the genotype. Statistical
significance is assessed by a Wilcoxon-Mann-Whitney test (p values: n.s. > 0.05; * < 0.05; ** < 0.01). Every
treatment has been repeated at least twice with similar results.
(C) Bacterial density in Log10 Colony-Forming-Unit (CFU) per leaf cm². Statistical significance is assessed by a
Wilcoxon-Mann-Whitney test (p values: n.s. > 0.05; ** < 0.01). Boxplots colors refer to the genotype as in panel
B. The treatment has been repeated three times with consistent results.
(B-C) Representative phenotypic pictures of Col-0 and *nac53-1 78-1* plants under several treatments are included.
181

Taken together this suggests that both proteasome activators are involved in general regulation

of subcellular proteostasis. However, depending on the perturbation site, in the light of

organellar stress, the response appears multi-layered. Overall, it suggests a precise subcellular

regulation of NAC53 and NAC78 to adjust their impact on plant response to stress. But how do

two transcription factors integrate different signals to have a global influence on the proteome

during proteotoxic stress conditions?

NAC53/78 are part of an autoregulatory proteasome feedback loop

To elucidate how both TFs might integrate signals from different compartments and organelles,

we aimed to decipher their regulation through interactome analysis. To this end, we generated

transgenic plants expressing NAC53 or NAC78 tagged with eGFP at the N-terminus (N-ter),

under control of the UBQ10 promoter (**Figure S1A**). Intriguingly, none of the generated lines

displayed visible fluorescence under basal conditions (**Figure 2A**). Only upon proteasome

inhibition with bortezomib (BTZ), GFP fluorescence was detected in nucleus (**Figure 2A**).

These observations were confirmed by immunoblotting using an anti-GFP antibody (**Figure**

2B). Taken together, our results suggest that NAC53/78 are constitutively turned over by the

proteasome implying the presence of an autoregulatory feedback loop.

To circumvent the short-lived feature of both NAC transcription factors in *A. thaliana*, and to proceed with our interactome study, we used *Agrobacterium*-mediated transient expression in *N. benthamiana* epidermal cells. Firstly, pUBQ10::GFP/RFP-NAC53/78 (hereafter referred as sTag-NACs, **Figure S1A**) were transiently expressed in *N. benthamiana* to obtain details on their subcellular localization in basal conditions. We could observe a fluorescence signal in reticulated structures reminiscent of the endoplasmic reticulum (ER) as confirmed by co-localization with an ER marker (mCherry-HDEL, **Figure 2C**). Such localization is consistent with the presence of a predicted transmembrane domain (TM) at the C-terminus of both proteins (**Figure S1A**). Due to this specific localization, we developed a second type of chimeric constructs. We tagged the proteins at both ends including a mCerulean3 (CFP) at the N-ter and a mCherry (RFP) at the C-terminus (C-ter), after the TM. These constructs were expressed under regulation of a tandem *Lotus japonicus UBQ1* promoter to obtain stronger expression (hereafter refers as dTag-NACs; **Figure S1A**; (Binder et al., 2014; Maekawa et al., 2008). Interestingly, we could observe the CFP signal was mainly located in the nucleus while the RFP remained strictly at the ER (**Figure 1D**). Both constructs appeared responsive to BTZ treatment that led to stabilization of both NAC transcription factors in *N. benthamiana* strengthening our findings that NAC53/78 are targets of the proteasome. (**Figure S1B and S1C**).

 To decipher how both TFs might integrate signals from different compartments/organelles and if an autoregulatory feedback loop governs their function, we performed immuno-Precipitation (IP) followed by tandem Mass Spectrometry (MS/MS) with both constructs. We processed a total of 4 IP-MS/MS conditions for both NACs among 2 experiments. We performed IP of sTag-NACs in mock conditions and at 8 hours after bacterial infection, to induce proteoxic stress (**Figure S1D**). In parallel, we performed IP of dTag-NACs enriching for the N-ter or C-ter end of the proteins, which allowed us to characterize the interactome at a subcellular resolution (**Figure S1E**). For our downstream analysis, we focused on the common interactors for both NACs. Out of the 4 conditions we identified 245 unique *A. thaliana* orthologous proteins as interactor candidates (**Figure 2E and Table S1**).

We first analysed the localization profile of NAC53/78 common interactome using the Gene Ontology (GO) annotations for cellular components and the SUBA5 data base (https://suba.live/). The GO analysis displayed enrichment for terms mainly associated with the proteasome complex and plastid components (**Figure S1F**). This supports the importance of the proteasome complex as a central hub in the NAC53/78 interactome. Investigating the

Figure 2. NAC53/78 are part of an autoregulatory proteasome feedback loop.
(A) Confocal microscopy pictures of roots of GFP-NAC53/78 *Arabidopsis thaliana* transgenics at 7dag exposed to mock or BTZ 10µM treatment for 3h. Treatments were repeated at least three times with similar observations.
(B) Immunoblot analysis against GFP on crude extract of seedlings related to panel A. Ponceau staining is used as loading control.
(C) Confocal microscopy pictures of transiently co-expressed sTag-NAC53/78 with prom35S::RFP-HDEL or together in *N. benthamiana* epidermis cells. Pearson index (ρ) represents co-localization index. Scale bars = 10µm
(D) Confocal microscopy pictures of transiently co-expressed sTag-NAC53/78 together or dTag-NAC53/78 alone in *N. benthamiana* epidermis cells. Scale bars = 10µm
(E) Bar plots representing the number of significantly enriched proteins in the 4 IP-MS/MS conditions for NAC53 only, NAC78 only and NAC53+NAC78 (Common).
(F) Cladogram of top 30 gene ontology (GO) terms for biological process (BP) enriched in the list of *A. thaliana* ortholog proteins found from NAC53/78 common interactome.
(G) Protein network of the interacting candidates related to the GO BP terms annotated as "Transport" and "Proteolysis" in panel F.

profile; we found enrichment for only 4 compartments: plastid, peroxisome, golgi, and
endoplasmic reticulum (**Figure S1G**). The latter is in accordance with the observed localization
(**Figure 2C**) and supports the importance of the ER-localization for the regulation of
NAC53/78. To understand how this regulation takes place, we further extended our analysis.
We performed GO analysis for biological processes and revealed that possible candidates were
enriched in proteins related to three main biological process: proteolysis, trafficking and
translation (**Figure 2F**). Based on our previous results we focused on the first two groups. We
extracted the corresponding proteins and performed a protein network analysis (**Figure 2G**).
This analysis highlighted the presence of numerous components of the ubiquitin-proteasome
system (UPS) (**Figure 2G; Figure S1H and S1I**); confirming the strong interplay of NAC53/78
with ubiquitin-dependent proteasomal degradation. Comparison of the protein enrichment level
in the several conditions revealed that association with the proteasome complex components
was strongly reduced by bacterial infection and mainly enriched when targeting the ER-
anchored end of the TFs (C-ter IP) (**Figure S1H**). It suggests that both proteasome regulators
escape proteasomal degradation in the context of proteotoxic stress and that ER is the major
site for their proteasomal removal. The latter was further supported by the presence of several
ERAD components in the interactome (**Figure 2G and S1I**). Additionally, both TFs were
associated with various trafficking components or peptidases (**Figure 2G**). Interestingly, the
affinity with most of these proteins was substantially differing based on the analysed condition
(**Figure S1J and S1K**). This indicates that the NAC53/78 interactome might involve further
regulatory layers dynamically shaped in response to proteotoxic stress.
Taken together our analysis confirmed that both proteasome activators are short-lived proteins
degraded by the proteasome. These findings reveal the presence of an autoregulatory feedback
in plants, similar to what has been found in yeast and animal (Marshall & Vierstra, 2019). In
addition, our interactome of NAC53/78 corroborates their role as integrators of subcellular
proteostasis perturbations.
276

ER-Anchored Protein sorting (ERAPS) regulates the subcellular fate of NAC53 & NAC78
Our analysis of the NAC53/78 interactome revealed that NAC53 and NAC78 associate with
proteins involved in the ubiquitin-proteasome system at the ER. Interestingly, among these
candidates, three belonged to the ER-Associated Degradation (ERAD) machinery, including
the membrane-bound E3 ligase HRD1 and the central component CDC48 (**Figure 2G and S1I**).
Given the ER localization of the TFs (**Figure 2C**) and the strong enrichment of proteasome

 subunits in the C-ter IP-MS/MS condition (**Figure S1I**) we hypothesized that the turnover of
 both NAC TFs are regulated by ERAD.

Figure 3. ER-Anchored Protein Sorting (ERAPS) coordinates the subcellular fate of NAC53 & NAC78.
(A) Confocal microscopy pictures of GFP-NAC53/78 expressing *A. thaliana* roots. 7 dag seedlings were exposed to mock treatment, BTZ 10µM, CB-5083 10µM or LS-102 100µM for 3h. The treatments were repeated at least three times with similar observations. Scale bars = 10µm
(B) Immunoblot against GFP on crude extracts of GFP-NAC53/78 *A. thaliana* adult plants infiltrated with mock treatment, BTZ 10µM, CB-5083 10µM or LS-102 100µM for 6h. Ponceau staining is used as loading control.
(C) Confocal microscopy pictures of *N. benthamiana* transiently co-expressing GFP-NAC53/78 full-length (FL) or deleted for the transmembrane domain (ΔTM) with RFP-HRD1a or RFP-HRD1b. Pearson index (ρ) indicates correlation index at the orange lines. Scale bars = 10µm
(D) Co-IP of RFP-HRD1a/b with GFP-NAC53/78. IP was performed with anti-GFP beads. GFP-HDEL and free-GFP were used as negative control. The experiment was repeated twice with similar results.
(E) Immunoblot against GFP on crude extract of *N. benthamiana* leaves transiently expressing GFP-NAC53/78FL or ΔTM. Ponceau was used as loading control. Boxplots represent the relative ΔTM abundance compared FL.

(F) *In vitro* trans-ubiquitination assays of GST-HRD1a/b against MBP-NAC53/78. Removal of E1 and ATP from
the reaction was used as negative control. The experiment was repeated twice with similar results.
(G) Representation of the ER-Anchored Protein Sorting (ERAPS) mechanism. NAC53/78 are ER-anchored
proteins recognized and ubiquitinated by the HRD1 complex. The CDC48 complex subsequently extracts
NAC53/78 from the ER to mediate ER-Associated Degradation or allowing nuclear translocation in the context of
proteasomal stress to mitigate proteotoxicity.

To investigate this possibility, we took advantage of previously characterized drugs: we treated the GFP-NAC53 and GFP-NAC78 transgenic lines with CB-5083 and LS-102, inhibitors of CDC48 and HRD1 proteins, respectively (Yagishita et al., 2012; Zhou et al., 2015). The effect of CB-5083 has already been characterized in plants (Li et al., 2022; Marshall et al., 2019), while LS-102 has not been characterized in plants to date. Our analysis revealed that LS-102 can efficiently inhibit both *A. thaliana* orthologs AtHRD1a and AtHRD1b *in vitro* (**Figure S2A and S2B**). Similar to BTZ treatment, CDC48 and HRD1 inhibition led to a stabilization of both proteins; visualized by confocal microscopy and through GFP immunoblotting (**Figure 3A and 3B**). Collectively, these results indicate that NAC53/78 are targets of ERAD. We next investigated whether NAC53 and NAC78 physically interact with both HRD1 isoforms. Transient co-expression of GFP-NACs with RFP-HRD1a/b revealed a strong association of NAC TFs with HRD1s at the ER, that was further confirmed by co-IP experiments (**Figure 3C and 3D**). The subcellular association was largely reduced upon deletion of the TM of both NAC TFs (ΔTM, **Figure 3C**). Interestingly, loss of ER association led to a strong stabilization of the proteins, supporting ERAD-dependent turn-over for both TFs (**Figure 3E**). These results support that NAC53/78 are targets of HRD1-mediated ubiquitination for proteasomal degradation.

To investigate whether HRD1s are indeed responsible for NAC ubiquitination, we performed *in vitro* ubiquitination assays using purified GST-HRD1a/b isoforms and MBP-NAC53/78 proteins. The assay revealed trans-ubiquitination of both NAC proteins in the presence of HRD1a and b that appeared in form of higher molecular weight bands of MBP-NAC proteins (**Figure 3E**). We could confirm the presence of ubiquitination marks at different sites via MS/MS analysis of the *in vitro* ubiquitination reactions (**Figure S2C and S2E**). In addition, ubiquitin marks were consistently inhibited by the addition of the inhibitor LS-102 to the *in vitro* ubiquitination assay (**Figure S2F**). The *in vitro* data was corroborated by identification of several ubiquitination sites on NAC53 and 78 in our *in vivo* IP-MS/MS analysis (**Figure S2D and S2E**). We confirmed *in planta* ubiquitination by performing ubiquitin IP on *N. benthamiana* leaves transiently expressing the dTag-NACs constructs (**Figure S2G**). Mutation of identified lysine residues substantially reduced the association with anti-ubiquitin beads *in planta* (**Figure S2H and S2I**). Altogether, our *in vivo* and *in vitro* analysis demonstrate that

NAC53 and NAC78 are directed to proteasomal degradation via ERAD through HRD1 ubiquitination.

During our analysis we could observe that NAC53/78 did not respond similarly to the tested inhibitors. Indeed, upon BTZ or CB-5083 treatment, GFP signal in the transgenic lines displayed different subcellular localization (**Figure 2A**). While BTZ led to a strong nuclear localization; CDC48 inhibition by CB-5083 led to a nuclear localization and cytoplasmic dot-like structures. To rule out the possibility of difference in drug potency, we treated plants with higher concentration of both drugs. While double amount of BTZ resulted in a stronger nuclear localization, an increase in CB-5083 concentration resulted in the accumulation of dot-like structures (**Figure S3A**), which could be potential sites of CDC48/NAC association. Co expression of *A. thaliana* CDC48 isoform (AtCDC48c) with both NACs revealed that CDC48 and NAC53/78 were able to associate in dot-like structures (**Figure S3B**). This indicates that upon CDC48 inhibition in *A. thaliana*, NAC53/78 accumulate at sites of CDC48 action. Given the role of CDC48 in retro-translocation of ER proteins to the cytosol, it could prevent degradation of NAC53/78 but also perturb their subcellular sorting. To test this, we compared the GFP signal intensity in the nucleus and cytoplasm upon BTZ and CB-5083 treatments (**Figure S3C**). Upon CB-5083 treatment the mean GFP intensity per cell was generally higher than upon BTZ (**Figure S3C**). This is consistent with a more diffuse localization of GFP signal. However, we observed that upon BTZ treatment more than 50% of the signal was found in the nucleus compared to 30% upon CB-5083 (**Figure S3D**). In addition, analysing the nucleus/cytoplasm mean intensity ratio revealed that NAC53/78 concentrate 6 times more in the nucleus upon BTZ compared to CB-5083 (**Figure S3E**). This analysis strongly suggests that CDC48 is partially involved in the nuclear translocation of NAC53 and NAC78 in addition to its role in proteasomal degradation.

In summary, we showed that the subcellular fate of NAC53 and 78 and hence the autoregulatory feedback loop is coordinated by the ERAD machinery. The turnover of ER-anchored NACs and their subsequent physiological impact is also the first example of the plant ERAD machinery having a role beyond quality control of aberrant proteins. In parallel this mechanism is responsible for their nuclear translocation (**Figure 3G**). Thus, we propose to refer to this mechanism as ERAPS (ER-Anchored Protein Sorting).

The transcriptional landscape of proteotoxic stress

Having established that ERAPS is controlling the proteasome autoregulatory feedback loop in plants, we wanted to understand why the *nac53-1 78-1* mutant is more tolerant to organelle-directed proteotoxic stress while it is hypersensitive to general proteotoxic stress (**Figure 1B**). To decipher the underlying mechanism, we utilized the fact that *Pseudomonas synringae* infection is causing proteotoxic stress through the suppression of proteasome activity and broad modification of chloroplast proteome (Littlejohn et al., 2021; Üstün et al., 2018; Üstün et al., 2016). We first characterized the involvement of the proteasome autoregulatory feedback loop during bacterial infection. Monitoring mRNA levels for both TFs genes, revealed an induction of *NAC53* and *78* upon infection (**Figure 4A**). In addition, transcriptional activation of proteasome genes was dependent on the presence of *NAC53* and *NAC78* WT alleles (**Figure 4B**). This was also consistent with an impaired protein accumulation of proteasome subunits in *nac53-1 78-1* upon infection (**Figure 4C**). Finally, we observed that bacterial infection was associated with the stabilization of NAC53 and NAC78 in *N. benthamiana* and *A. thaliana* (**Figure 4E and 4F**). Altogether these results demonstrate that bacterial infection displays all the features associated with proteotoxic stress that is dependent on the presence of both proteasome activators. As such, this system is ideal to unveil the function of the proteasome autoregulatory feedback loop in response to environmental perturbation.

To obtain a global idea on what else beyond the proteasome genes might be affected in *nac53-1 78-1* plants we undertook a transcriptomic analysis. We first analysed the transcriptomic profile of *nac53-1 78-1* infected with *Pst*. Interestingly, *nac53-1 78-1* displayed only 7 differentially expressed genes (DEGs) compared to Col-0 in mock (**Figure 4G)**, confirming that NAC53/78 are mainly involved in stress responses (Seo et al., 2008). Comparing the *nac53-1 78-1* to Col-0 in the context of bacterial infection revealed 945 DEGs. Strikingly, most of the genes display a positive Log2FC (**Figure 4H**). This suggests that NAC53/78 are major transcriptional repressors upon bacterial infection induced proteotoxic stress. We could confirm that bacterial infection triggers a general NAC53/78-dependent transcriptional up-regulation of proteasome related genes (**Figure S4A**). These results are in accordance with the involvement of the proteasome regulatory feedback loop during infection. Further investigation of which biological processes are enriched within the DEGs in our *nac53-1 78-1* transcriptome revealed photosynthesis, glucosinolates, cell wall and auxin signalling as major terms (**Figure 4I**). To refine our investigation and to include a broader view of the transcriptome landscape during proteotoxic stress, we included two other datasets into our analysis. Firstly, we generated

 another transcriptome in response to pathogenic bacteria, subjecting the *rpt2a-2* proteasome

Figure 4. The transcriptional landscape of proteotoxic stress

(A) Log2FC mRNA levels of *NAC53* and *NAC78* transcripts after *Pst* infection of adult Col-0 plant leaves. The heatmap represent the mean of 4 biological replicates.

(B) Log2FC mRNA levels of 26S proteasome transcripts after *Pst* infection of Col-0 or *nac53-1 78-1* adult plant leaves. The heatmap represent the mean of 4 biological replicates. The experiment has been repeated 3 times with consistent results.

(C) Immunoblot against multiple 26S proteasome subunits on crude extracts of Col-0 or *nac53-1 78-1* adult plant leaves inoculated with *Pst* or mock solutions for 24h. Ponceau staining is used as loading control.

(D) Relative Log2FC abundance of representative 26S proteasome subunits in Col-0 or *nac53-1 78-1* adult plant leaves. The abundance is calculated relative to the mock conditions. Statistical differences are assessed via a Welch t.test (p values: n.s > 0.05, * < 0.05, ** < 0.01).

(E) Immunoblot against GFP on crude extract of *N. benthamiana* leaves transiently expressing GFP-NAC53/78 after mock treatment or *PstΔHopQ* infection for 8h. Ponceau staining is used as loading control.

(F) Immunoblot against GFP on extract of adult *A. thaliana* transgenics GFP-NAC53/78 subjected to mock treatment or *Pst* infection 8h by vacuum infiltration followed by IP with anti-GFP beads.

(G) Number of differentially expressed genes (DEGs) in the different conditions of the *nac53-1 78-1 Pst* RNAseq
analysis. Total number of DEGs per condition is indicated. Colors indicate the proportion of DEGs up-regulated
and down-regulated. DEGs are considered when |Log2FC| > 1.5. Genot. mock: *nac53-1 78-1* mock vs. Col-0
mock; WT *Pst*: Col-0 *Pst* vs. Col-0 mock; Mut. *Pst*: *nac53-1 78-1 Pst* vs. *nac53-1 78-1* mock; Genot. *Pst*: *nac53-*
1 78-1 Pst vs. Col-0 *Pst*.
(H) Level of differential expression (Log2FC) of the 945 DEGs found in Genot. *Pst* (see panel G) for the several
conditions analyzed.
(I) Cladogram of top 40 GO terms for BP enriched in the list of 945 DEGs. Colors represent broad GO clusters.
(J) Venn diagram representing the overlap between the 945 DEGs and DEGs from *rpt2a-2 Pma* (|Log2FC| > 0.5)
or Col-0 MG132 24h (|Log2FC| > 0.5).
(K) Cladogram of top 30 GO terms for BP enriched in the list of 538 DEGs extracted from the combined analysis
(panel J). For conciseness terms are hidden and grouped in 4 processes.
(L) Protein network of the genes from the 945 DEGs list related to the GO BP terms from panel K. Node colors
refers to the 4 GO clusters from panel K.
431

mutant to *Pseudomonas syringae pv. maculicola* (*Pma*) infection. Contrary to *nac53-1 78-1,*

this genotype displayed 251 DEGs in mock conditions (**Figure S4B**); indicating a basal stress

response, including upregulation of proteasome genes, consistent with previous report (Lee et

al., 2011). Upon infection we observed a large transcriptome variation compared to WT plant;

with 981 DEGs identified specific to *rpt2a-2 Pma* condition (**Figure S4B and S4C**). Analysing

the transcriptomic response of the proteasome complex, we found that most of the genes were

further activated in response to infection in *rpt2a-2* background (**Figure S4D**). Interestingly,

beyond terms associated with the proteasome, GO analysis on the 981 DEGs revealed others

enriched process overlapping in both, *nac53-1 78-1* and *rpt2a-2*, transcriptomes (**Figure 4I and**

S4G). Secondly, we included a previously characterized transcriptome analysis on Col-0 plants

upon MG132 treatments (Gladman et al., 2016). This led to three distinct proteotoxic stress

transcriptomes induced by drugs, infection and/or genetic mutations.

The comparison of all three datasets allowed us to extract a core set of 538 DEGs present in the

nac53-1 78-1 dataset and one or both others (**Figure 4J, Table S2**). Strikingly, GO enrichment

on this subset of genes revealed 4 distinct biological processes involved (**Figure 4K**).

Extracting the genes associated with these processes from the *nac53-1 78-1* 945 DEGs, we

could define the core transcriptional network associated with proteotoxic stress (**Figure 4L**).

This network is comprised of 4 modules: the proteasome genes, a major cluster of

Photosynthesis associated nuclear genes (PhANGs), glucosinolate and auxin signalling

pathways (**Figure 4K and 4L**). Analysing the transcriptional profile of the several clusters we

identified interesting patterns (**Figure S4F**). While the proteasome cluster was present in all

three datasets (**Figure S4C and S4F,** (Gladman et al., 2016), the glucosinolate related cluster

was infection specific and the auxin cluster was found only in *nac53-1 78-1 Pst* and MG132

24h (**Figure S4H**). This further emphasized the specificity and complexity of the proteotoxic

stress response. The photosynthesis cluster was found in significant proportions in all three

datasets with 68 DEGs for *nac53-1 78-1 Pst*, 23DEGs for *rpt2a-2 Pma* and 57DEGs for MG132

24h (**Figure S4H, Table S2**). This suggests that PhANGs transcriptional repression is a
recurrent response to proteotoxic stress. Strikingly, similar to the transcriptional response of
proteasome genes, the differential expression of the other clusters is also happening in opposite
manner in *nac53-1 78-1 Pst* compared to *rpt2a-2 Pma* or MG132 24h (**Figure S4H**). It shows
that the observed transcriptional response is not due to an indirect effect of dysfunctional
proteasome in *nac53-1 78-1* background as our previous results could have suggested.
Altogether our analysis revealed a so far unidentified link between NAC53/78 and the
transcriptional regulation of PhANGs in response to proteotoxic stress.
466
NAC53 and NAC78 coordinate the regulation of proteasome and photosynthesis gene
expression through the same cis-element
To identify whether NAC53/78 mediate a link between proteasome activation and repression
of photosynthesis, we hypothesized NAC53/78 can act as novel repressors of PhANGs.
Previous reports have shown that NAC TFs are involved in transcriptional repression (Hao et
al., 2010; Shih et al., 2014). We therefore speculated PhANGs being direct targets of NAC53
and NAC78. This was also corroborated by the detailed analysis of distinct photosynthetic
processes: While many processes were downregulated in the proteotoxic stress core
transcriptome in the *rpt2a-2 Pma* and MG132 data sets, the absence of NAC53/78 had the
opposite effect, indicating a possible repression via the NAC53/78 module (**Figure S5A**). To
investigate a possible direct repression of PhANGs, we extracted the promoter regions of the 4
gene clusters from our transcriptional network (**Figure 4K and 4L**). Running enrichment
analysis for specific cis-element, we found that Proteasome and PhANGs promoters share a
similar cis-element characterized by the PRCE-like [TGGGC] motif, while the other clusters
did not (**Figure 5A**). This further supported the link between PhANGs regulation and the
proteasome regulatory feedback loop. In line with this, PRCE-like elements appeared
homogeneously distributed among PhANGs (**Figure S5B**). Additionally, analysis of the cis-
element position revealed a high frequency of occurrence and enrichment close to the
transcription starting site, similar to proteasome promoters (**Figure S5B and S5C**). We then
extracted two candidates' regions from LHCa3 and PSAD1 promoters (**Figure 5B**),
respectively members of the light harvesting complex I and photosystem I (**Figure S5A,**
(Leister, 2003). Performing electromobility shift assay (EMSA), we could confirm NAC53 and
NAC78 were able to associates with these elements (**Figure 5C**). Consistently, this association
was dependent on the presence of their respective DNA recognition motif (**Figure 5C**).

Furthermore, we confirmed the TGGGC motif as being the driver of this association, as mutated

Figure 5. NAC53 and NAC78 coordinate the regulation of proteasome and photosynthesis gene expression through the same cis-element.

(A) Top 3 motifs found by STREME software in the promoter regions from the 4 network clusters (Figure 4L). PRCE-like elements from proteasome and PhANGs clusters are highlighted.

(B) 25bp region comprising the PRCE-like elements from LHCa3 and PSAD1 promoters used in this figure. The region corresponds to the probe sequence used in panel C and D. The PRCE is highlighted in red.

(C) Electro-Mobility Shift Assay (EMSA) of MBP-NAC53/78 with LHCa3/PSAD1 probes (see panel B). As a negative control, probes were incubated without MBP-NAC53/78 (-) or with MBP-NAC53/78 deleted for their DNA recognition motif (ΔRM). Experiment was repeated twice with similar results.

(D) EMSA competition assay of MBP-NAC53/78 association with LHCa3/PSAD1 probes. Competitor is applied at a concentration gradient (25X, 50X, 75X and 100X) indicated by the shade of grey. WT and mutated competitors are labelled TGGGC and AAAAA, respectively. Experiment was repeated twice with similar results.

(E) Boxplot representing association of GFP-NAC53/78 with LHCa3/PSAD1 PRCE in adult *A. thaliana* transgenics lines infected with *Pst* or mock for 8h after Chromatin Immunoprecipitation. The association is quantified as % of input and amplification of a region in the CDS is used as negative control.

(F) Immunoblot against mature PhANGs proteins in Col-0 or *nac53-1 78-1* adult plant leaves inoculated with *Pst*
or mock solutions for 24h. Ponceau staining is used as loading control.
(G) Relative Log2FC abundance of PhANGs mature proteins immunoblotted in panel F. The abundance is
calculated relative to the mock conditions. Statistical differences are assessed via a Welch t.test (p values: n.s >
0.05, * < 0.05, ** < 0.01).
(H) Photosystem II (PSII) activity measurement on Col-0 or *nac53-1 78-1* adult plants inoculated with *Pst* or mock
solutions for 24h. Letters indicate the statistical group assessed by pairwise Welch t.test (p value < 0.05). The
experiment was repeated 3 times with consistent results.
(I) Representative pictures of measurements from panel I. Fv/Fm values are represented by false color as indicated
by the color gradient legend.
(J) Total chlorophylls and carotenoids from Col-0 or *nac53-1 78-1* adult plant leaves inoculated with *Pst* or mock
solutions for 24h. Letters indicate the statistical group assessed by pairwise Wilcoxon-Mann-Whitney test (p value
< 0.05). The experiment was repeated 3 times with consistent results.
521

oligos were unable to outcompete the association of NAC53/78 with the probes (**Figure 5D**).

To corroborate our findings *in vivo*, we performed chromatin IP followed by qPCR in the

context of bacterial-induced proteotoxic stress (**Figure 5E**). As expected, we could see that

bacterial infection triggers the association of both TFs to both promoter regions, confirming the

direct targeting of NAC53/78 to PhANGs promoters. To assess the regulatory role of NAC53/78

on PhANGs promoter activity, we used the luciferase reporter system in protoplasts (Yoo et al.,

2007). Our analysis revealed that the PSAD1 promoter activity was consistently repressed by

NAC53 and 78, which did not occur in *nac53-1 78-1* mutant protoplasts (**Figure S5D**). Deletion

of the PRCE motif in PSAD1 abolished NAC78-mediated repression, while led NAC53 to

mediate an increase in promoter activity (**Figure S5E**). In contrast, LHCa3 promoter activity

appeared to be rather induced by NAC53 and unchanged by NAC78 (**Figure S5D**), with

NAC53-mediated activation dependent on the PRCE motif (**Figure S5E**). This difference might

be due to the fact that both TFs are expressed without a TM domain, bypassing ERAPS, in a

rather artificial system. To circumvent this problem, we analysed the gene expression of *LHCa3*

and *PSAD1* in the transgenic NAC53 and 78 lines upon BTZ treatment. Proteasome inhibition

in the transgenic NAC53 and 78 lines partially enhanced the repression of PhANGs, supporting

the notion that NAC53/78 act as repressors of photosynthesis during proteotoxic stress (**Figure**

S5F). In addition, both lines showed an increase in *PBA1* and *RPT2a* mRNA (**Figure S5G**),

confirming that NAC53/78 can act concomitantly as transcriptional repressor and activator.

These results confirmed the importance of the PRCE-like element in PhANGs promoters,

suggesting that NAC53 and NAC78 possess the ability to modulate each other's activity and

highlights the complexity of the NAC53/78-PRCE regulatory module mediating transcriptional

repression and activation on distinct gene cluster.

As NAC53 and NAC78 can associate with the PRCE-like element in PhANGs promoters to

dampen their transcriptional activity, it is likely that they act as novel repressors of

photosynthesis. To confirm this hypothesis, we analysed the protein abundance of LHCa3 and

PSAD1 proteins in the context of bacterial infection (**Figure 5F and 5G**). Immunoblot revealed
that contrary to Col-0, infection induced reduction of photosynthesis-associated proteins
appeared less pronounced in the *nac53-1 78-1* background (**Figure 5F**). Semi-quantitative
analysis on several replicates confirmed this observation (**Figure 5G**). This confirms that the
observed transcriptional repression mediated by the NACs has a major impact on PhANGs
protein levels. To test if this protein pool perturbation would impact photosynthesis, we
monitored PSII activity during infection. We found that *nac53-1 78-1* displayed significantly
less PSII activity reduction compared to Col-0 (**Figure 5H and 5I**). In addition, when
monitoring photosynthetic pigment content in the same context, we observed that total
chlorophyll and carotenoid content reduction during infection occurred to a lesser extent in the
nac53-1 78-1 background (**Figure 5J**). Consistently with these findings, subjecting the
NAC53/78 transgenics to constant proteasome inhibition resulted in strong developmental
defect compared to WT (**Figure S5H and S5I**), suggesting that the role of NAC53/78 on
photosynthesis repression is not restricted to bacterial infection.
Altogether, we demonstrate the unique ability of NAC53 and NAC78 to mediate activation and
repression of target genes through the PRCE transcriptional module. This makes them novel
repressors of photosynthesis during bacterial infection-induced proteotoxic stress. It also might
explain our previous findings that the NAC mutant was more tolerant to organelle-directed
proteotoxic stress, suggesting that photosynthesis repression is a general feature of the
proteasome autoregulatory feedback loop to cope with proteotoxicity.
568
The proteasome autoregulatory feedback loop monitors photosynthesis homeostasis
during stress responses
Our results show that NAC53/78 co-regulate proteasome and PhANGs by recognizing the same
cis-element during bacterial infection induced proteotoxic stress. However, this feature does
not only occur during bacterial infection as the PhANGs repression was also observed when
proteasome was chemically inhibited (**Figure 4L, S5A, S5F and S5G**). Consistent with this,
we have observed that the *nac53-1 78-1* mutant also performed better when we perturbed
organellar proteostasis (**Figure 1B**). In addition, recent evidence suggests that loss of
proteasome function in a chloroplast import deficient mutant increased photosynthetic capacity
due to a lack of chloroplast precursor proteins degradation (Grimmer et al., 2020). Altogether,
this hints towards a general role of the proteasome activators NAC53/78 to co-ordinate
photosynthesis and proteasome homeostasis during proteotoxic stress.
581

To investigate to which extend NAC53/78 were involved in response to chloroplast
perturbation, we analysed the photosynthetic response of *nac53-1 78-1* plants upon Bortezomib
and Lincomycin treatments. Our analysis revealed that the decrease in pigment content was
impaired in *nac53-1 78-1* in comparison to Col-0 plants upon BTZ and Lincomycin treatment
(**Figure 6A**). Consistent with this, repression of PhANGs was less pronounced in the double
mutant (**Figure 6B**) which resulted in a higher photosynthetic activity in *nac53-1 78-1* upon
Lin treatment (**Figure S6A**).

To directly study the involvement of the proteasome autoregulatory feedback coordination of
cellular proteostasis and photosynthesis, we decided to analyse the central ERAPS regulators
HRD1a/b utilizing the inhibitors that were used in the NAC double mutant phenotyping (**Figure
1**). Proteasome inhibition rendered the *hrd1a 1b* more tolerant in comparison to Col-0 (**Figure
6C**), which is consistent with its role in negatively regulating NAC53/78 stability (**Figure 3G**).
In contrast, all other tested drugs or bacterial infection rendered the *hrd1a 1b* mutant more
susceptible (**Figure 6C and 6D**). The reduced growth of *hrd1a 1b* upon chloroplastic
perturbations (Cml and Lin) are consistent with our previous results (**Figure 1B**), as
stabilisation of NACs would lead to increased repression of PhANGs and growth reduction
(**Figure S5A, S5H and S5I**). The impaired performance of *hrd1a 1b* upon CDC48 inhibition,
pathogen infection or ER stress induction might be due to the general function of HRDs in
ERAD (Chen et al., 2016) and thus appear to be pleiotropic. Consistent with this idea, analysis
of the photosystem II activity revealed a stronger reduction in the *hrd1a 1b* double mutant
(**Figure S6B**) upon infection corroborating our findings that ERAPS coordinate proteasome
and photosynthesis through NAC53/78 stability. In addition, these data suggest that organellar
perturbation would lead the NAC53/78 activation. Indeed, subjecting GFP-NACs transgenics
to several organellar perturbators led to stabilization and nuclear localization of both TFs
(**Figure S6C**). Taken together our analysis illustrates the importance of the proteasome
autoregulatory feedback loop in maintaining the fine equilibrium of proteostasis to ensure
normal response to proteotoxic stress.

We found that regardless the genetic mutations impacting the feedback loop, the plants
displayed an increased susceptibility to bacterial infection (*rpt2a-2*, Üstün et al. (2016)). This
finding demonstrates the importance of this mechanism in agronomically important context and
raises the question if the autoregulatory feedback loop is involved in other conditions. To

answer this, we performed a large-scale transcriptomic analysis using the recently developed

Figure 6. The proteasome autoregulatory feedback loop monitors photosynthesis homeostasis during stress responses

(A) Photosynthetic pigment content from Col-0 or *nac53-1 78-1* adult plant leaves infiltrated with Lin 200μM, BTZ 2μM or mock solutions for 24h. Content is expressed in percentage relative to Col-0 mock. Letters indicate the statistical group assessed by pairwise Wilcoxon-Mann-Whitney test (p value < 0.05). The experiment was repeated 3 times with consistent results.

(B) Log2FC mRNA level of the indicated transcripts in Col-0, *nac53-1 78-1* adult plants leaves Lin 200μM, BTZ 2μM or mock solutions for 24h. Log2FC is calculated from Col-0 mock and *PDH1* is used as negative control. Statistical differences are assessed with a Welch t-test (p values: n.s. > 0.05, * < 0.05, ** < 0.01). The experiment was repeated twice with consistent results.

(C) Fresh weight of seedlings grown under the indicated treatments at 10-14 day after germination. The fresh weight is represented as a percentage of Col-0 mock conditions. Letters indicate the statistical groups assessed by pairwise Wilcoxon-Mann-Whitney test (p value < 0.05). Every treatment has been repeated at least twice with consistent results. Representative phenotypic pictures are included.

(D) Bacterial density shown as Log10 Colony-Forming-Unit (CFU) per leaf cm² in Col-0 and *hrd1a hrd1b* plants. Statistical significance is assessed by a Welch t-test (p values: * < 0.05). The experiment has been repeated three times with consistent results.

(E) Correlation heatmap of the 54 26S proteasome subunit genes and the expression of 68 PhANGs from publicly available 1223 RNAseq libraries identified in this study.

Arabidopsis RNAseq data base (Zhang et al., 2020). We first investigated the relationship of

transcript abundance between the 54 genes encoding 26S proteasome subunits and the 68

PhANGs associated with proteotoxic stress (**Figure 4**). We first extracted 165 unique projects

of interest (see Methods, **Table S3**) to perform a correlation analysis between mRNA abundance

of individual genes from both clusters (**Figure 6E**). We could observe that genes within each

cluster displayed a strong positive correlation ($\rho \approx 0.6$ and $\rho \approx 0.5$ for 26S Proteasome and PhANGs, respectively). Strikingly, analysing relationship between the two clusters showed a significant negative correlation ($\rho \approx -0.4$). It demonstrates that both clusters are regulated in opposite manner (**Figure S4H**) and that this is a general feature of stress responses. We therefore refined our analysis to define the spectrum of conditions displaying this transcriptional signature (see Methods). Thus, we extracted 43 projects in which 26S Proteasome and PhANGs appear transcriptionally co-regulated (**Table S3**). From these 43 projects, 35 displayed an opposite regulation for 26S Proteasome and PhANGs mRNA level (**Figure S6G**), confirming that both clusters undergo recurrent contrasting co-regulation. Looking at the conditions related to these datasets revealed that bacterial, oomycete infection or elicitation, led to a negative correlation of proteasome gene expression and PhANGs expression. This is in line with our previous observation using *Pst* infection. Beyond biotic stress, our analysis unveiled several other contexts, such as cold, dark and drought treatments as factors triggering this negative correlation (**Figure S6G**). Furthermore, we found that treatment with trans-chalcone a phytotoxin notably causing shoot bleaching (Diaz-Tielas et al., 2019) also resulted in a similar response. In addition to abiotic stress, nutrient stress and circadian rhythm triggered the same transcriptional signature. It suggests that coordination of PhANGs/Proteasome is not only a feature of plant stress response but also a characteristic of developmental processes which is additionally mirrored in hormone treatments such as SA and Auxin (**Figure S6G**).

Taken together, we found that the role of NAC53/78 in repression of photosynthesis is a general feature of proteotoxic stress at any subcellular level. We discovered that the coordination of proteasome activation and PhANGs repression is a key feature of the *Arabidopsis thaliana* transcriptome in response to multiple biotic, abiotic, and developmental cues. Given the conserved nature of the autoregulatory feedback loop in yeast, plants and animals, our findings bear the potential that proteasome activators are general gatekeepers of the interplay between the proteasome and energy metabolism across the tree of life.

Discussion

Understanding how eukaryotic organisms orchestrate proteostasis in a complex compartmentalized cellular environment is a long-standing question. Although the proteasome can degrade proteins from different compartments and organelles, how these different subcellular signals are integrated upon external cues and coordinated to maintain overall proteostasis was unknown.

675 Here, we have characterized that the proteasome autoregulatory feedback loop, involving the

676 transcriptional NAC53/78-PRCE module, acts as a gatekeeper to facilitate the communication

677 between the triangle chloroplast-proteasome-nucleus during proteotoxic stress (**Figure 7**).

Figure 7. ER-anchored protein sorting (ERAPS) controls the fate of two proteasome activators for intracellular organelle communication during proteotoxic stress.

In steady state conditions, the ER-anchored protein sorting system (ERAPS) promotes the constitutive degradation of NAC53/78 via the 26S proteasome. Meanwhile, PhANGs expression is activated and subsequent chloroplastic import permits the maintenance of active photosynthesis. Upon proteotoxic stress, the chloroplast import machinery is targeted for proteasomal degradation. This leads to an accumulation of PhANGs precursors which are therefore subjected to proteasomal degradation, inducing proteotoxicity. Thus, to avoid proteotoxicity, the ERAPS system facilitates the nuclear translocation of NAC53/78, to activate the production of a new proteasome complex and to repress PhANGs expression to mitigate substrate accumulation.

690 We have shown that the trade-off between proteasome activation and photosynthesis

691 downregulation is activated in response to various environmental and developmental cues

including disease. Using bacterial infection induced proteotoxic stress, we have identified ERAPS as the key regulatory node of this feedback loop and how the latter shapes and maintains subcellular proteostasis in plants. Considering that the autoregulatory feedback loop is mechanistically conserved across kingdoms, similar control mechanisms of subcellular proteostasis are possible in yeast and in animals (Marshall and Vierstra, 2019). While in yeast the main proteasome regulator is the cytosolic TF Rpn4; in animal systems, SKN-1A/Nrf1 is an ER-anchored protein which has also been shown to be subjected to ERAD (Radhakrishnan et al., 2014; Ruvkun and Lehrbach, 2023; Steffen et al., 2010). It suggests further mechanistic similarities between plant and animal systems, despite proteasome activators distinct evolutionary origins, as our study demonstrates the importance of ERAPS for the subcellular fate of NAC53/78, to a similar extend to what was reported for Nrf1 (Ruvkun and Lehrbach, 2023), emphasizing the conserved role ERAPS in regulating the proteasome activators. However, our data might suggest that nuclear translocation of NAC53 and 78 is only partially dependent on CDC48, as both NACs are still present in the nucleus upon CDC48 inhibition. In addition, the phenotypes observed using the *hrd1a hrd1b* mutant and the localization of both TFs upon LS-102 treatment suggest that HRD1-mediating ubiquitination is not strictly required for NAC53/78 nuclear translocation. Thus, it is likely that the ERAPS system involves alternative actors or other regulatory layers to mediate the transport of both TFs into the nucleus. For instance, SKN-1A/Nrf1 TFs were reported to be targeted by the ubiquitin-dependent DDI1/2 aspartyl-proteases for cleavage, downstream of their retro-translocation (Ruvkun and Lehrbach, 2023). Although NAC53/78 are evolutionary distinct from SKN-1A/Nrf1 and do not share the known DDIs cleavage site, we cannot exclude similar activation process due to the presence of an ortholog, DDI1, in *A. thaliana* (Farmer et al., 2010). In addition, a recent report has proposed a second ubiquitination step for Nrf1 in the cytosol which would facilitates DDI2-mediated cleavage (Hu et al., 2023). The presence of cytosolic E3 in NAC53/78 interactome could support similar mechanisms.

The ERAPS-dependent autoregulatory pathway proposed here would permit a rapid integration of proteotoxic signals from various compartments. As such, it would allow the communication between different organelles such as chloroplast or mitochondria with the nucleus, a process referred as retrograde signaling, known as the modulation of nuclear encoded gene expression in response of organellar proteostasis modification (Woodson and Chory, 2012). This communication is essential to respond to sudden changes in the environment. As such, retrograde signaling from mitochondria or chloroplast to the nucleus involves massive transcriptional changes that in turn influence organelle function (Richter et al., 2023). Thus, it

requires tight transcriptional regulation to respond to signals from different organelles. Some
nuclear communicators have been identified to mediate retrograde signaling (Richter et al.,
2023). However, to date, transcription factors GLK1 and 2, are the only known PhANGs
activators (Waters et al., 2009), while putative repressors ABI4 and PTM could not be verified
as retrograde signaling components (Kacprzak et al., 2019; Koussevitzky et al., 2007; Page et
al., 2017; Sun et al., 2011). Here we have identified that plants have evolved a dual-regulatory
system through the proteasome autoregulatory feedback loop: the NAC53/78-PRCE module
acts as a novel retrograde signaling component, repressing PhANGs expression during
prolonged proteotoxic stress. By binding to the same cis-element the NAC53/78 module is able
to activate proteasome genes and repress PhANGs. In this scenario, (i) repression of PhANGs
by NAC53/78 may be a strategy to cope with the excess of precursor accumulation due to
proteasomal stress (**Figure 7**) which is corroborated by our findings that the *nac53-1 78-1*
mutant is more tolerant to photosynthesis stress and HSP90 inhibition which activate
NAC53/78 nuclear localization but also (ii) to counteract the overactivation of PhANGs by
GLK1 that is degraded by the proteasome (Tokumaru et al., 2017). Although, NAC TFs have
been reported to possess a repressor domain (NARD) (Hao et al., 2010), an intriguing question
remains: how are NAC53/78 acting as activators and repressors using the same cis-element?
Future investigations on this are necessary to improve our understanding how NACs activate
and repress gene expression simultaneously or in a concurrent manner to mediate retrograde
signaling.

In line with our findings, similar strategies to cope with the trade-off between proteasome
activation and energy producing organelles have been established in yeast for mitochondrial
associated proteins (Boos et al., 2019; Weidberg and Amon, 2018). In this case proteasome
activator Rpn4 is activated upon accumulation mitochondrial precursor protein accumulation
to turn on a second regulator Pdr3 which triggers the so called mitochondrial compromised
protein import response (mitoCPR) (Kramer et al., 2021). This rather constitutes an indirect
mechanism to transcriptionally repress the mitochondrial oxidative phosphorylation machinery
(OXPHOS). This implies a divergence between yeast and plants on subcellular proteostasis
coordination. It is tempting to propose these differences have emerged due to multicellularity
allowing an improved flexibility depending on the tissue or cell types. This hypothesis is
supported by the lack of the PRCE cis-element in the proteasome gene promoters of the
monocellular algae *Chlamydomonas reinhardtii* (Nguyen et al., 2013). In addition, we could
observe NAC53/78 activation in root cells, which lacks chloroplasts.. This observation further
supports the general function of the proteasome regulatory feedback loop in maintenance of

subcellular proteostasis upon stress, not limited to chloroplastic impairment. To our knowledge, a similar system has not been found in other multicellular organisms such as animals. However, the characterization of OXPHOS gene repression by ATFS-1 during mitochondrial unfolded protein response (Shpilka et al., 2021), the importance SKN-1A/Nrf1 upon oxidative stress and the role of Nrf2/Nrf1 in mitochondrial biogenesis and proteasome activation (Ruvkun and Lehrbach, 2023) strongly suggests a tight link between proteasome and mitochondrial proteostasis in animals. Interestingly, another NAC TF, ANAC017 was characterized to mediate mitrochondria-nucleus retrograde signaling (Ng et al., 2013). As NAC TFs are known to function as homo or hetero-dimers (Welner et al., 2016), it is tempting to speculate that the proteasome autoregulatory feedback loop involves additional NAC in concert with NAC53/78, or other TFs, orchestrating the specificity of the transcriptional response.

Various environmental perturbations impede the balance between protein translation, sorting and degradation resulting in proteotoxicity. This has a global effect on proteostasis, and as mounting evidence highlights a role for the proteasome in clearing organelle-associated proteins it is essential to decipher how signals from different compartments and organelles are integrated to maintain proteostasis and avoid proteotoxicity. As such, our findings here lay the foundation to study whether the role as communicators between organelles during proteotoxic stress is a common feature of proteasome activators in different kingdoms.

Acknowledgments

We thank Yasin Dagdas and Marja Timmermans for critical discussions during the thesis advisory committee meeting. We are thankful to Ophélie Léger and Paul Gouguet for stimulating discussions. We thank Nan Wang for technical advice on the Ch-IP procedure, Robert Morbitzer for technical support on the golden gate molecular cloning. We are grateful to Ana Gabriela Andrade-Galan and Lars Pospisil for technical support on the Photosystem II activity measurement. Stefan Bieker for the help with the computational analysis. We are thankful to Jos Schippers, Yasin Dagdas and Richard Vierstra for sharing published material. This work was supported by an Emmy Noether Fellowship GZ: UE188/2-1 from the Deutsche Forschungsgemeinschaft (DFG; to S.Ü.) and through the collaborative research council 1101 (SFB1101; to G.L.). We thank the confocal microscopy facility of ZMBP that is supported by funds from DFG (INST 37/819-1 FUGG and INST 37/965-1 FUGG) and the confocal microscopy facility of RUB (DFG Project number 523980288, GZ:INST 213/1180-1 FUGG). Macek Lab is supported by DFG funds (INST 37/1050-1 FUGG and INST 37/1255-1 FUGG).

Author Contribution

S.Ü. conceived the project. G.L. performed most experiments, analyzed the data, and performed all computational analysis. M.R. performed the *in vitro* ubiquitination assays. D.B performed the EMSA, assisted with the Luciferase reporter assay and the molecular cloning. D.S. generated the *rpt2a-2 Pma* transcriptome under the supervision of F.B..M.F.W and B.M. performed LC-MS/MS analysis. S.Ü. wrote the manuscript together with G.L..

Experimental model and Subject Details

Arabidopsis thaliana lines and growth conditions

A. thaliana Columbia-0 (Col-0) ecotype was considered as WT. T-DNA insertion mutants for *NAC53* (*nac53-1*, SALK_009578C) and *NAC78* (*nac78-1*, SALK_025098) were obtained from Richard Vierstra and previously described (Gladman et al., 2016). T-DNA insertion mutants for *RPT2a* was previously described (Lee et al., 2011; Üstün et al., 2016). T-DNA insertion mutants for *HRD1a (hrd1a*, SALK_032914) and *HRD1b* (*hrd1b,* SALK_061776) were obtained from Yasin Dagdas and previously described (Su et al., 2011).

promUBQ10::GFP-NAC53 and *promUBQ10*::GFP-NAC78 transgenics lines were obtained by *Agrobacetrium tumefaciens* floral dipping (Clough and Bent, 1998) of Col-0 plants. Positive transformants were selected by BASTA spraying and seeds were harvested individually. The following generations were sawed on ½ Murashige and Skoog (MS) media supplemented with 10µM phosphinotricin for segregation analysis. Two independent lines showing a 75% survival rate for each construct were kept for subsequent experiments.

For experiments on seedlings, seeds were surface sterilized 10min with 1.3% sodium hypochlorite. Seeds were then sown on ½ MS medium supplemented with 1% sucrose and stratified for 2 days in cold room to synchronize germination. Plants were next grown under long day conditions (light/dark cycles: 16h 22°C/8h 20°C, 130µmol.mm².s^{-1} light intensity, 70% relative humidity).

For experiments on adult plants, seeds were sown on soil soaked with water in individual pots. Trays were put directly in the growth chamber under short day conditions (light/dark cycles: 12h 22°C/ 12h 20°C, 90µmol.mm².s^{-1} light intensity, 70% relative humidity). Seven days after germination, excessive number of seedlings were removed from individual pots to obtain one plant per pot. Plants were grown under same conditions until use (typically 21 to 28 more days).

Nicotiana benthamiana growth

N. benthamiana plants were grown under long day conditions (light/dark cycles: 16h/8h, at

21°C and 70% humidity) and typically used 4 weeks after germination for confocal microscopy

experiments and 5 weeks after germination for co-immunoprecipitation experiments.

Bacterial strains

For Pseudomonas syringae pv. *tomato* DC3000 wild-type and *Pseudomonas syringae* pv.

maculicola strains ES4326 were grown on King's B medium containing rifampicin 100µg.mL^{-}

^{1} at 28°C. For floral dipping and transient expression in *Nicotiana benthamiana* leaves,

Agrobacterium tumefaciens C58C1 strain was grown in LB broth high salt (Duchefa L1704)

containing the required antibiotics at 28°C. For molecular cloning, *Escherichia coli* Top10

strain was grown in LB broth high salt (Duchefa L1704) containing the required antibiotics at

37°C. For protein purification, *Escherichia coli* BL21 (DE3) was grown in LB broth high salt

(Duchefa L1704) containing the required antibiotics at 37°C or 16°C.

Method Details

Molecular cloning

For chimeric protein constructs generated in this study, protein coding sequence (CDS) of the

desired genes were amplified from *A. thaliana* cDNA with addition of flanking region for

subsequent cloning process.

For Golden Gate based cloning (Binder et al., 2014), CDS were amplified with addition of

flanking regions including BpiI/BsaI sites (Fw: 5'-GAGAAGACGATACGGGTCTCACACC-

3'; Rv: 5'-GAGAAGACGACAGAGGTCTCACCTT-3'). PCR products were gel purified and

inserted into LI module (Binder et al., 2014) backbone using BpiI/T4 ligase (both Thermo

Scientic) cut-ligation procedure (40 cycles, 37°C 2min/16°C 5min). The resulting product was

transformed into *E. coli* Top10 followed by gentamycin selection. Positive clones were used for

plasmid miniprep (NEB) and verified by sanger sequencing. For generation of expression

constructs, the LI CDS module and desired other modules (e.g., promoter, tag, terminator etc.)

were assembled in the LIIa F 1-2 vector (Binder et al., 2014) using BpiI/T4 ligase (both Thermo

Scientic) cut-ligation procedure (40 cycles, 37°C 2min/16°C 5min). Resulting product was

transformed into *E. coli* Top10 followed by spectinomycin selection. Positive clones were

verified for correct assembly by colony PCR following by enzyme restriction analysis. The

obtained construct was then transformed into *A. tumefaciens* C58C1 for subsequent use.

For GATEWAY™ based cloning, CDS were amplified with addition of attb1/attb2 flanking

regions (attb1-fw 5'-GGGGACAAGTTTGTACAAAAAAGCAGGCTTC-3'; attb2-rv 5'-

GGGGACCACTTTGTACAAGAAAGCTGGGT-3'). PCR products were gel purified and

inserted into pDONR221 via BP clonase™ II enzymatic reaction. Alternatively, GATEWAY™

compatible entry clone was generated by performing a BsaI/T4 ligase cut-ligation from LI

module to pENTR-BsaI (Binder et al., 2014). Resulting product was transformed into *E. coli*

Top10 followed by kanamycin selection. Positive clones were used for plasmid miniprep (NEB)

and verified by sanger sequencing. For generation of expression constructs, pDONR/pENTR

clone was used for LR Clonase™ II enzymatic reaction into the desired destination vectors.

Resulting product was transformed into *E. coli* Top10 followed by spectinomycin or

carbenicillin selection. Positive clones were verified by colony PCR. The obtained construct

was then transformed into *A. tumefaciens* C58C1 or *E. coli* BL21 (DE) for subsequent use.

Seedlings phenotyping

For phenotyping assay, seeds were sown on ½ MS 1% sucrose round petri dish supplemented

with the indicated concentration of drug or a mock treatment.

After 10 to 14 days of growth fresh weight was measured. Seedlings were carefully extracted

from the media with a curved tweezer (Dumont Nr. 7) ensuring that the entire root was taken.

For one replicate, 5 representative seedlings were scaled together on an analytical scale

(resolution 0.0001g). Typically, 5 replicates were measured for subsequent statistical analyses.

Transient expression in *N. benthamiana* leaves

Bacterial suspension was centrifuged for 4min at 4000g after overnight incubation of *A.*

tumefaciens strain carrying the desired expression construct. Supernatant was removed and

pellet was resuspended in 500µl of agroinfiltration buffer (10mM $MgCl_2$, 10mM MES pH 5.7).

OD600 was measured and infiltration solutions were diluted to reach final 0.5 OD600 in

agroinfiltration buffer supplemented with 200µM acetoseryringone. Solutions were incubated

in dark at least 1h prior to infiltration. Infiltrated tissues were used 30h post infiltration for

subsequent experimental procedures.

Protein purification

890 Recombinant proteins were expressed in E. coli BL21(DE3) by IPTG induction. Bacterial

891 solution was centrifuged 20min at max speed and pellet was subjected to sonication after

892 resuspension in 1mL MBP-buffer (20 mM TRIS-HCl pH 7.5, 200mM NaC, 1mM EDTA) or

893 1mL IPP50 (10 mM Tris-HCl, pH 8.0. 150mM NaCl, 0.1% NP40). Sonicated samples were

894 used for purification by affinity chromatography using amylose resin (New England Biolabs)

for MBP-NAC53/78 or glutathione sepharose 4B (cytiva) for GST-HRD1a/b. For recombinant
His-UBA1 and His-UBC8 protein were purified using Ni-Ted resin (Macherey-Nagel).

In vitro ubiquitination assay

In vitro ubiquitination assay was performed as previously described (Leong et al., 2022).
Purified proteins were used for in vitro ubiquitination assays. Each reaction of 30 ml final
volume contained 25 mM Tris–HCl, pH 7.5, 5 mM MgCl2, 50 mM KCl, 2 mM ATP, 0.6 mM
DTT, 2 μg ubiquitin, 200 ng E1 His- AtUBA1, 1.2 μg E2 His-AtUBC8, 2 μg of GST-HRD1a/b,
and 0.3 μg of MBP-NAC53/78. Samples were incubated for 1 h at 30°C, and the reaction was
stopped by adding SDS loading buffer and incubated for 10 min at 68°C. Samples were
separated by SDS–PAGE electrophoresis using 4–15% Mini-PROTEAN® TGX™ Precast
Protein Gels (BioRad) followed by detection of the ubiquitinated substrate by immunoblotting
using anti-MBP (New England Biolabs), anti-GST and anti-ubiquitin (Santa Cruz
Biotechnology) antibodies.

Co-immunoprecipitation

Plant tissue was homogenized in a mortar with liquid nitrogen to keep the sample frozen during
the process. The powder was then transferred into a 50mL centrifuge tube and 1ml of extraction
buffer (10% glycerol, 25 mM Tris pH 7.5, 1 mM EDTA, 150 mM NaCl, 1 mM DTT, 0.5%
Triton X, 1X protease cocktail inhibitor (Sigma)) was added per gram of tissue. Obtained
sample were vortexed for 3min and incubated in cold room on rotator in cold room for 10min
following a 30min centrifugation at 4000g, 4°C. Supernatant was filtered through 3 layers of
miracloth (Merck 475855-1R) in a 15mL centrifuge tube. 50μl of filtrate was sampled as input
at this step, supplemented with 12.5μL laemmli buffer 4X (Biorad) and boiled for 10min at
90°C. Next, 10μl of ChromoTek GFP-Trap® or RFP-Trap Agarose beads per gram of tissue
were added and tubes were incubated 2-3h in cold room. Tubes were then centrifuge at 800g
for 1min; supernatant was carefully removed and pelleted beads were transferred to 1.5mL
microcentrifuge tube in 1mL of washing buffer (10% glycerol, 25 mM Tris pH 7.5, 1 mM
EDTA, 150 mM NaCl, 0.5% Triton X-100, 1X protease cocktail inhibitor (Sigma)). Washing
was perfomed by centrifugation at 800g for 1min using 1mL washing buffer. After last washing,
10μL of Laemmli buffer 2X (Biorad) was added to equal amounts of beads and sample was
boiled 10min at 90°C.

RNA isolation & RT-qPCR

Total RNA isolation was performed using RNeasy Plant Mini Kit (Qiagen 74904). All steps

were performed according to manufacturer instructions. To exclude potential contaminant

DNA, RNA samples were subjected to DNase I treatment (Thermo scientific™) following

provider instructions. For RNA sequencing, integrity and RNA concentration was determined

(2100 Bioanalyzer, Agilent). For RT-qPCR sample analysis was adapted from Langin and Üstün

(2023); cDNA synthesis was performed using LunaScript® RT SuperMix Kit (NEB) following

provider recommendation. qPCR was performed using MESA BLUE qPCR 2X MasterMix

Plus for SYBR® Assay (Eurogentec) using a 2-step reaction protocol for 40cycles with

systematic evaluation of primer melting curve. mRNA level was quantified based on the $\Delta\Delta Ct$

method followed by Log2 transformation.

NanoLC-MS/MS analysis and data processing

For purification, proteins were subjected to a NuPAGE 12% gel (Invitrogen) and in-gel trypsin

digestion was done on Coomassie-stained gel pieces, as described previously (Borchert et al,

2010) with a modification: iodoacetamide was used instead of chloroacetamide for

carbamidomethylation of cysteine residues to avoid formation of lysine modifications isobaric

to two glycine residues left on ubiquitinylated lysine after tryptic digestion. Next, peptides

mixture were desalted using C18 Stage tips and run on an Easy-nLC 1200 system coupled to a

Q Exactive HF-X mass spectrometer (both Thermo Fisher Scientific) as described elsewhere

(Kliza et al., 2017) with some modifications: separation of the peptide mixtures was done using

a 87-min segmented gradient from 10-33-50-90% of HPLC solvent B (80% acetonitrile in 0.1%

formic acid) in HPLC solvent A (0.1% formic acid) at a flow rate of 200 nl/min. The seven most

intense precursor ions were sequentially fragmented in each scan cycle using higher energy

collisional dissociation (HCD) fragmentation. In all measurements, sequenced precursor

masses were excluded from further selection for 30 s. The target values were 105 charges for

MS/MS fragmentation and 3×106 charges for the MS scan.

Acquired MS spectra were processed with MaxQuant software package version 1.5.2.8 with an

integrated Andromeda search engine. For *in vivo* IP samples, database search was performed

against a *N. benthamiana* database containing 74,802 protein entries (Kourelis et al., 2019), the

sequences of eGFP-NAC53/78, mCerulean3-NAC53/78-mCherry and 285 commonly

observed contaminants.

For *in vitro* ubiquitination samples, database search was performed against the Uniprot *E. coli*

database, the sequences of MBP-NAC53/78, GST-HRD1a/b and 285 commonly observed

contaminants.

Endoprotease trypsin was defined as a protease with a maximum of two missed cleavages.

Oxidation of methionine, phosphorylation of serine, threonine, and tyrosine, GlyGly dipeptide

on lysine residues, and N-terminal acetylation were specified as variable modifications.

Carbamidomethylation on cysteine was set as a fixed modification. Initial maximum allowed

mass tolerance was set to 4.5 parts per million (ppm) for precursor ions and 20 ppm for fragment

ions. Peptide, protein, and modification site identifications were reported at a false discovery

rate (FDR) of 0.01, estimated by the target-decoy approach (Elias and Gygi). The iBAQ

(Intensity Based Absolute Quantification) and LFQ (Label-Free Quantification) algorithms

were enabled, as was the "match between runs" option (Schwanhäusser et al., 2011).

Photosynthesis monitoring

For photosynthetic pigment concentration, 2 leaf discs (6mm) were sampled in a 2mL

microcentrifuge tube after the indicated treatments and incubated over-night in 1mL acetone

100% on a rotating wheel in cold room. Tubes were next centrifuged for 3min at 400g. 200µl

of was then pipetted in a transparent 96-well plate and absorbance was measured at 470nm,

646nm and 663nm using a plate reader (Tecan Infinite 200 PRO®). Acetone 100% was used as

blank. Concentration was calculated as following:

Total Chlorophylls ($\mu g.ml^{-1}$): $((12.25*A_{662}) - (2.79*A_{646}) + (21.5*A_{646}) - (5.1*A_{663}))/0.2$

Total Carotenoids ($\mu g.ml^{-1}$): $((1000*A470-1.63*((12.25*A_{662}) - (2.79*A_{646}))$ -

$85.02*((21.5*A_{646}) -(5.1*A_{663})))/221)/0.2$

The equations for concentration calculation and procedure were obtained from Karlsruher

Institut für Technologie (KIT, https://www.jkip.kit.edu/molbio/998.php, "Chlorophyll and

Carotenoid determination in leaves") as an updated version of the original procedure

(Lichtenthaler, 1987).

Values were then then expressed as ng.mm² by dividing them by the total leaf area used for

extraction (typically 56.55mm²).

For photosynthetic efficiency, photosystem II (PSII) activity was measured by quantifying the

maximum photosystem II yield, Fv/Fm via the saturation pulse methods on dark acclimated

plants (Schreiber, 2004). After treatments, plants were acclimated in dark for 15min. Next,

leaves were detached and PSII activity was assessed by Fv/Fm values using the Imaging-PAM

chlorophyll fluorometers: Maxi version v2-46i, for measurements related to Figure 5 and Dual-

PAM100, for measurements related to Figure 6 (Walz GmbH, Effeltrich, Germany).

Bacterial Infection

For bacterial infections, *Pst* overnight liquid culture was centrifuged for 15min at 4000g.
Bacterial pellet was resuspended in 5mL of $MgCl_2$ 10mM prior to OD600 measurement.

For qPCR, RNAseq and western blot analysis; a *Pst* solution at OD600=0.2 was used. Adult
plant leaves were infiltrated with a 1mL syringe (Braun) with the *Pst* infiltration solution or a
control solution (10mM $MgCl_2$) and put back in growth chamber for 24h until sampling or
further analysis.

For IP and Chromatin-IP on *A. thaliana*, *Pst* infiltration was prepared as explained above and
detached whole rosettes of adult plants were vacuum infiltrated 2 times 2 min at < 15mbar. For
mock, control solution (10mM $MgCl_2$) was used similarly. Rosettes were put back in the growth
chamber for 8h on multiple layers of wet tissue.

For assessment of bacterial density (related to Figure 1C and 6D); *Pst* infiltration solution was
prepared in 10mM $MgCl_2$ at an OD600 of 0.0001. Adult plant leaves were infiltrated with a
1mL syringe (Braun) with the *Pst* infiltration solution and put back in growth chamber for 72h
in high humidity conditions. After 3 days, 2 leaf disks per replicates were sampled with a biopsy
punch 6.0mm (Harris Uni-Core) in a 1.5mL microcentrifuge tube. Tissue was homogenized in
200µl 10mM and a serial dilution of factor 10 was done for each replicate in 200µL nano pure
water until 10^{-6} dilution was reached. For bacterial counting, 20µL of several resulting dilutions
(typically 10^{-5} and 10^{-6}) were plated on KB media with rifampicin for each replicate. Growing
colony-forming units were counted and Log10CFU/cm² was calculated for graphical
representation and statistical testing.

1017

Chromatin Immuno-Precipitation
For Chromatin-IP, after Arabidopsis rosettes were vacuum infiltrated for 30min with a fixation
buffer solution (1% formaldehyde, 10mM KH2PO4 pH 7, 50mM NaCl, 0.1M sucrose, 0.01%
Triton X-100) to cross-link protein-DNA. Plant material (typically 1g) was dried and frozen in
liquid nitrogen for each replicate. Tissue was next ground in 5mL nuclei isolation buffer (20mM
Hepes pH8, 250mM sucrose, 1 mM $MgCl_2$, 5mM KCl, 40% Glycerol, 0.25% Triton X-100,
0.1mM PMSF, 0.1% 2-mercaptoethanol, 1µM BTZ). The resulting solution was filtered
through three layers of miracloth paper in a 50mL centrifuge tube following 10min
centrifugation at 4°C, 3000g. Pellet was resuspended in 1mL nuclei isolation buffer and
transferred into a 1.5mL microcentrifuge tube. Tube was centrifuged 5min, at 4°C, 3000g and
the previous step was repeated once. Pellet was resuspended in 200µl M3 buffer (10mM
KH2PO4 pH7, 0.1mM NaCl, 10mM 2-mercaptoethanol) and recentrifuged 5min, 4°C, 3000g.
Nuclei pellet was resuspended in 1mL sonication buffer (10mM KH2PO4 pH7, 0.1mM NaCl,

0.5% Sarkosyl, 10mM EDTA). Sonication was performed with an Active Motif Sonicator (Amp 25%; Pulse: 30sec ON – 30sec OFF; Timer: 300sec). Sonicated sample was centrifuged for 5min, 4°C at max speed. Supernatant was tranferred to a new 1.5mL microcentrifuge tube and 10% (v/v) Triton X-100 was added to neutralize the sarkosyl. 5% (v/v) of this solution was saved in another 1.5mL microcentrifuge as input sample. For IP, 15µL ChromoTek GFP-Trap® were activates in 15µL IP buffer (50mM Hepes pH 7.5, 150mM NaCl, 5mM MgCl$_2$, 10uM ZnSO4, 1% Triton X -100, 0.05% SDS) and added to the tube and incubated for 3h on a rotating wheel at 4°C. Beads were washed for 4 rounds by centrifuging for 30sec, at 1000g and subsequent incubation on rotating wheel in 500µL buffer for 3 min; twice in IP buffer, once in LNDET buffer (0.25 LiCl, 1% NP-40, 1% deoxycholate, 1mM EDTA pH 8, 10mM Tris pH 8) and once in TE buffer (10mM Tris pH 8, 1mM EDTA pH 8). After the last washing step, beads were centrifuged for 30sec at 1000g, resuspended in 200µL EB buffer (50mM Tris pH 8, 10mM EDTA pH 8, 1% SDS) supplemented with 8µL 5M NaCl to reverse cross-linking and incubated at 65°C for at least 6h. At this step input sample was added and treated similarly. For protein digestion, 1µL of Proteinase K (Thermo Scientific) was added and incubated for 1h at 45°C. Next, DNA was purified using a PCR purification kit (NEB) following manufacturer instructions.

For DNA quantification, qPCR was performed using MESA BLUE qPCR 2X MasterMix Plus for SYBR® Assay (Eurogentec) using a 2-step reaction protocol for 40 cycles and each sample was processed in technical triplicates. For quantification, % of input was calculated and used for graphical representation and statistical analysis.

Immunoblot analysis

For immunoblot analysis, sample processing was adapted from (Langin and Üstün, 2023); 2 leaf discs from *N. benthamiana* or *A. thaliana* were sampled using a biopsy punch 6mm (Harris Uni-Core) in 2mL microcentrifuge tube containing 2 glass beads (2mm) for one replicate in the given condition and frozen in liquid nitrogen. Tissue was homogenized using a Tissue Lyzer II (Qiagen), 2 times 30sec, frequency: 25; followed by the addition of 150µL protein extraction buffer (100mM Tris-HCl pH 7.5, 3% SDS, 10mM EDTA) per tube. Tubes were next mixed with the Tissue Lyzer II, 2 times 15sec, frequency: 15. Finally, 50µL laemmli buffer 4X (biorad) was added and samples were boiled for 10min at 95°C. Tubes were centrifuged for 1min at 13000g and 150µL of the supernatant was transferred in a new 1.5mL microcentrifuge tube and kept at -20°C until use.

Western blotting was done by SDS-PAGE using TGX FastCast 10% Acrylamide gels (Biorad)
with 1.0 mm spacer plates according to manufacturer recommendations. SDS-PAGE was
performed in TGS running buffer (25 mM Tris pH 8.3, 192mM glycine, 0.1% SDS) and proteins
were transferred on PVDF membrane 0.2mm (Biorad) using Trans-Blot Turbo system (Biorad)
according to manufacturer recommendations. PVDF membranes were incubated for 1h in
TBST (50mM Tris-HCl pH 7.5, 150mM NaCl, 0.1% Tween20) solution containing 5%
Skimmed-Milk followerd by antibody incubation in TBST 5% Milk for 1h at RT or over-night
at 4°C. Membranes were washed 3 times in TBST for 5min and 1 time in TBS for 5min. HRP
chemiluminescence was detected using Amersham™ ECL Select™ detection reagent and
images were taken using a CCD camera.

1074

Electromobility Shift Assay
For EMSA assays, complementary DNA oligos were synthesized by EuroFins Genomics with
ATTO565 dye linked to the 5' end of minus strand for probes. To generate double stranded
probes, complementary oligos were incubated at 2µM in annealing buffer (25mM HEPES-
KOH pH 7.8, 40mM KCl) at 70°C for 5min and cool down to room temperature.
EMSA reactions were performed in 20µL reaction buffer (25 mM HEPES-KOH pH 7.8, 40
mM KCl, 1 mM DTT, 10% Glycerol) using 1µg purified MBP-NAC53/78 and 50ng DNA probe
for 30min at 25°C. Reactions were subjected to native polyacrylamide migration with TGX
FastCast 7.5% Acrylamide gels (Biorad) in TAE (Tris-base 40 mM pH 8.3, 20mM acetic acid,
1mM EDTA). After migration, probes were imaged using a Typhoon FLA 9000 Gel Scanner.

1085

Luciferase reporter assay in Protoplasts
For promoter activity analysis, *Arabidopsis* mesophyll protoplasts were generated adapted from
Yoo et al., 2007. Leaves from adult plants were cut into strips (0.5mm) and incubated in the
Maceroenzyme R-10 (Duchefa M8002) and Cellulase Onozuka R-10 (Duchefa C8001) solution
for 3h. Solution was subsequently filtered through 3 layers of miracloth paper (Merck 475855-
1R) in a 50mL centrifuge tube. Protoplast suspension was centrifuged for 1min at 200g at 4°C
and supernatant was carefully removed from the protoplast pellet. Pellet was washed twice with
W5 buffer (Yoo et al., 2007) in ice. Protoplasts were resuspended in MMG buffer (Yoo et al.,
2007), 200µL dispatched in 2mL microcentrifuge tube and 5µg of each plasmid were
transfected by addition of 220µL 30% PEG4000 (Sigma-Aldrich 81240) solution for 15min.
880µL of W5 was added to stop the reaction followed by 1min centrifugation at 200g.
Supernatant was removed and transfected protoplast pellet were resuspended in 250µL WI

buffer (Yoo et al., 2007). Tubes were left over night in dark for construct expression prior to measurements.

Luciferase quantification was performed according to Dual-Luciferase® Reporter Assay System from Promega recommendation (see link). Protoplast expressing the desired contructs were centrifuged for 15sec at max speed. Supernatant was removed and protoplast pellet was resuspended in 100µl of 1X passive lysis buffer (see link). The lysed protoplast suspension was transferred into a white 96-well plate (Thermo Scientific™ Nunc™ MicroWell™). Measurement of Firefly Luciferase (FfLuc, reporter constructs) was performed by adding 40µl LARII reagent (see link) and luminescence was measured using a plate reader (Berthold Tristar LB 940). Measurement of Renilla Luciferase (RnLuc, internal control) was performed adding 40µl of Stop & Glo® (see link) and luminescence was measured. Due to an apparent cross-talk between NAC53/78ΔTM expression and the *prom35S*::*RnLuc* constructs (see AtNAM DBD and 35S promoter interaction, (Duval et al., 2002); the results showed Figure S5 are based on the FfLuc luminescence only.

QUANTIFICATION AND STATISTICAL ANALYSIS

For all statistical analysis, R programming language was used in the Rstudio environment.

For non-omics data, the statistical test and associated p value threshold for each analysis are indicated in the figure legends. Data dispersion, medians and quantiles are represented with boxplots and replicates displayed in every figures. For bar plots in Figure S2, number of replicates are indicated in the figure legend, error bars represent the standard deviation from the mean.

RNA sequencing

For *rpt2a-2 Pma* transcriptome, sequencing was performed by ATLAS Biolabs, Berlin. For *nac53-1 78-1 Pst* transcriptome, sequencing was performed by the NGS Competence Center, Tübingen. After quality control, reads were mapped to the *A. thaliana* TAIR10 reference genome. The mapped reads were counted with *htseq-count* for subsequent analysis. Log2FC and false discovery rate (FDR) were determined using R package DEseq2 (Love et al., 2014) with default settings. A gene is considered significantly differentially expressed when its FDR < 0.05. Log2FC threshold is indicated in the figure legends according to the analysis.

IP-MS/MS analysis

For IP-MS/MS analysis in Figure 2 and Figure S2, Log2FC and false discovery rate (FDR) for
each protein group were determined using R package IPinquiry4 (Kuhn et al., 2023). For GFP-
NAC53/78 mock conditions, free-GFP mock was used as control; for GFP-NAC53/78 *Pst*
conditions, free-GFP *Pst* was used as control; for CFP-NAC53/78-RFP N-ter conditions, free-
GFP was used as control; for CFP-NAC53/78-RFP C-ter conditions, mCherry-HDEL was used
as control. For downstream analysis, to each protein group a unique Agi-code was assigned by
blasting every *N. benthamiana* protein present in the protein group and taking the most frequent
Agi-code with the best E-value. An e-value threshold of e-10 was used, in case of absence of
A. thaliana the protein group was not considered for downstream analysis. For heatmap
representation in Figure S1H-K, in case multiple protein groups correspond to the same *A.*
thaliana orthologs the mean Log2FC is represented.
1142

Meta-Transcriptomic analysis
For meta-transcriptomic analysis in Figure 6 and FigureS6, fragments per kilobase per million
counts were retrieved from the data base (http://ipf.sustech.edu.cn/pub/athrna/). Log2FC for
individual genes in every project was calculated based on the control RNAseq libraries as
annotated in the original publication (Zhang et al., 2020). Filtering of relevant projects for
subsequent correlation analysis was done based on the median |Log2FC|. Median|Log2FC| >
0.5 was used for identification of projects with at least one gene cluster transcriptionally
impacted (see Table S3, related to Figure 6H) and median |Log2FC| > 0.3219 was used for
identification of projects in which 26S proteasome and PhANGs clusters were co-
transcriptionally impacted (see Table S3, related to Figure S6G). Pearson correlations were
calculated using base R function. Only correlations with p-value < 0.05 were considered
significant and used for calculation of mean/median.
1155

Omics downstream analysis: Gene ontology, Protein Network, subcellular enrichment &
cis-element analysis
For Gene Ontology enrichment, the list of Agi-code was uploaded into ShinyGO software
(http://bioinformatics.sdstate.edu/go/, (Ge et al., 2020). Cladogram of the indicated number of
top terms for the indicated type of terms was used for graphical representation. Table of
annotations was extracted for further analysis.
For network creation, list of Agi-code was provided to the string database (https://string-
db.org/). Table of interactions was extracted and uploaded to cytoscape software
(https://cytoscape.org/) for network generation, design, and annotation. For purposes of clarity,

network nodes were manually arranged. Edge transparency relate to the strength of interaction
with opaque edges corresponding to strong interactions and transparent edges to weak
interactions. Node color corresponds to manual annotation inferred based on GO annotations
and further curated from the literature.

For subcellular compartment enrichment, the list of Agi-code was uploaded into SUBA5db
(https://suba.live/). Compartment annotation was extracted, and occurrence of every
compartment was counted to estimate the list distribution. For theorical distribution 20 Agi-
code lists of the same length were randomly generated and processed similarly. The mean
distribution of the 20 lists was used as theorical distribution. Enrichment for every compartment
was calculated as a Log2FC of the ratio between the observed proportion and the theorical one.

1175

Cis-Regulatory Element analysis
For cis-regulatory element analysis, promoter sequences of 1000bp upstream of the
transcription starting site were retrieved from plant ensembl database using biomart browser
(https://plants.ensembl.org/biomart/). For identification of novel cis-regulatory elements in
Figure 5A, promoter sequences of the several gene clusters were analyzed using the STREME
software (https://meme-suite.org/meme/tools/streme) with default settings. For mapping and
counting in Figure S5B and S5C, promoter sequences were analyzed in R using regular
expression matching for "tgggc" or "gccca" pattern.

1184

Immunoblot semi-quantitative analysis
Quantification of protein abundance based on immunoblot signals (related to Figure 4D and
5G) was adapted from (Gallo-Oller et al., 2018). Raw image of the immublot signals against
the protein of interest and the corresponding loading control were processed using Fiji software
(https://imagej.net/software/fiji/downloads). Image was converted to 8-bit format. A duplicated
image was used to generate a background image using the function "Substract Background"
with a ball radius value of 50 pixels and the option "create background (don't substract)
activated. The resulting background was then subtracted from the original 8-bit image with the
function "Image Calculator". Individual lanes were quantified by measurement of the peak area
after circumscription with the rectangular ROI selection and "Gels" function. The obtained
values for individual bands were normalized by the corresponding loading control values. For
calculation of Log2FC in abundance, quantification of bands in infected sample were divided
by the corresponding mock sample quantification of the same genotype; the obtained ratio was
next transformed using Log2 function.

Confocal microscopy imaging

Confocal microscopy was done using an inverted Zeiss LSM 880 microscope, an upright Leica SP8 microscope or an inverted Leica Stellaris 8 microscope. For mCerulean3 imaging, excitation was done using a 458nm laser and emission signal was measured from 465nm to 500nm; for eGFP or sGFP imaging, excitation was done using a 488nm laser and emission signal was measured from 510nm to 540nm; for mRFP or mCherry imaging, excitation was done using a 561nm laser and emission signal was measured from 590nm to 630nm; in addition chlorophyll auto fluorescence was measured in the far red wavelength (> 700nm). Images were acquired with a 40X water immersion objective, pinhole was set to 1 airy unit, resolution of acquisition was set to at least to 1024x1024 with a line average of 4.

After acquisition images were processed in Fiji Software. For images obtained from Leica microsystem devices, a gaussian blur (radius 0.75) was applied for a smoother rendering. For clarity purposes, contrast on the individual channels was manually adjusted.

For subcellular quantification of GFP signal (related to Figure S3), all images used were acquired on the Zeiss LSM 880 with a 16bits depth. Prior to imaging seedlings were incubated for 10min in a 10μM propidium iodide (PI) ½ MS solution followed by a brief washing in clear ½ MS. PI fluorescence was excited with the 488nm laser and measured from 590nm to 660nm. For all acquisitions within an experimental repetition the laser power and gain for the GFP channel was kept constant. For segmentation, images were processed in Fiji Software. Each cell was segmented using the polygon ROI selection tool. Each nucleus was segmented in the same order using the oval ROI selection tool. Next, mean intensity and integrated density of the whole cell, the nucleus, and the whole cell after removal of nucleus ROI (considered as cytosolic ROI) was measured. Signal proportion in the nucleus was quantified by the ratio between integrated density of nucleus ROI by whole cell ROI and ratio of signal between nucleus and cytosol was calculated by the ratio between nucleus ROI mean intensity and cytosolic ROI mean intensity.

For calculation of co-localization index in Figure 2C, the image J plugin "co-localization finder" was used and calculated pearson index is provided in the Figure. For calculation in Figure 3C, plot profile of GFP and RFP signals was extracted at the indicated line using Fiji software and the pearson index indicated on the Figure was calculated between the two profile values on R.

RESOURCE AVALAIBILITY

1233 Source code of R and Python3 scripts used for the several computational analysis done in this
1234 study will be put available on https://github.com/Gogz31 or can be directly requested.
1235

DATA AVAILABILITY

1237 The mass spectrometry data from this publication will be made available on the PRIDE archive
1238 (https://www.ebi.ac.uk/pride/archive/) and with the identifier (PXDXXXXX; and
1239 (PXDXXXX). All relevant proteomics data are made available in the supplemental information.
1240

SUPPORTING INFORMATION

Figure S1. Deciphering the details of NAC53/78 interactome. Related to Figure 2.
Figure S2. NAC53/78 are subjected to high level of poly-ubiquitination. Related to Figure 3.
Figure S3. NAC53/78 association with CDC48 is required for their nuclear translocation.
Related to Figure 3.
Figure S4. Transcriptional analysis of proteotoxic stress reveals a trade-off between 26S
proteasome and other biological process. Related to Figure 4.
Figure S5. Characterization of the NAC53/78-PRCE module in PhANGs promoters. Related
to Figure 5.
Figure S6. The proteasome autoregulatory feedback loop transcriptional signature is a systemic
response to environmental cues. Related to Figure 6.
Table S1. Spectral count enrichment of the protein groups identified in NAC53/78 common
interactome. Related to Figure 2.
Table S2. Enrichment of mRNA of 538 DEGs in *nac53-1 78-1* Pst, overlapping with *rpt2a-2*
Pma and/or MG132 24h data sets. Related to Figure 4.
Table S3. List of the several RNAseq projects identified in study in which 26S proteasome
and/or PhANGs are differentially regulated. Related to Figure 6.
1258

FIGURE LEGENDS

Figure 1. NAC53/78 are central integrators of various proteotoxic stress conditions.
(A) The role of the 26S Proteasome in degradation of proteins from distinct subcellular
compartments. Prime site of action of the several treatments used in this study: Blunt head
arrows indicate enzymatic activity suppression; pointed head arrows indicate trigger of
substrate accumulation.
(B) Fresh weight of seedlings grown under the indicated treatments at 10-14 day after
germination (dag). Fresh weight is expressed as a percentage of Col-0 mock conditions.
Boxplots colors refer to the genotype. Statistical significance is assessed by a Wilcoxon-Mann-
Whitney test (p values: n.s. > 0.05; * < 0.05; ** < 0.01). Every treatment has been repeated at
least twice with similar results.
(C) Bacterial density in Log10 Colony-Forming-Unit (CFU) per leaf cm². Statistical
significance is assessed by a Wilcoxon-Mann-Whitney test (p values: n.s. > 0.05; ** < 0.01).
Boxplots colors refer to the genotype as in panel B. The treatment has been repeated three times
with consistent results.
(B-C) Representative phenotypic pictures of Col-0 and *nac53-1 78-1* plants under several
treatments are included.
Figure 2. NAC53/78 are part of an autoregulatory proteasome feedback loop.

(A) Confocal microscopy pictures of roots of GFP-NAC53/78 *Arabidopsis thaliana* transgenics
at 7dag exposed to mock or BTZ 10μM treatment for 3h. Treatments were repeated at least
three times with similar observations.
(B) Immunoblot analysis against GFP on crude extract of seedlings related to panel A. Ponceau
staining is used as loading control.
(C) Confocal microscopy pictures of transiently co-expressed sTag-NAC53/78 with
prom35S::RFP-HDEL or together in *N. benthamiana* epidermis cells. Pearson index (ρ)
represents co-localization index. Scale bars = 10μm
(D) Confocal microscopy pictures of transiently co-expressed sTag-NAC53/78 together or
dTag-NAC53/78 alone in *N. benthamiana* epidermis cells. Scale bars = 10μm
(E) Bar plots representing the number of significantly enriched proteins in the 4 IP-MS/MS
conditions for NAC53 only, NAC78 only and NAC53+NAC78 (Common).
(F) Cladogram of top 30 gene ontology (GO) terms for biological process (BP) enriched in the
list of *A. thaliana* ortholog proteins found from NAC53/78 common interactome.
(G) Protein network of the interacting candidates related to the GO BP terms annotated as
"Transport" and "Proteolysis" in panel F.
Figure S1. Deciphering the details of NAC53/78 interactome. Related to Figure 2
(A) Chimeric NAC53/78 constructs generated in this study. Upstream promoter, fluorescent tag
and protein domain of interests are represented.
(B) Immunoblot against GFP on crude extracts of *N. benthamiana* leaves transiently expressing
sTag-NACs in addition to mock or 6h BTZ (10μM) treatment. Ponceau staining is used as
loading control.
(C) Immunoblot against GFP or RFP on crude extracts of *N. benthamiana* leaves transiently
expressing dTag-NACs in addition to mock or BTZ 10μM for 6h. Ponceau staining is used as
loading control.
(D) Immunoblot against GFP after immunoprecipitation (IP) with anti-GFP agarose beads on
N. benthamiana leaf extracts transiently expressing sTag-NACs after mock or *PstΔHopQ*
infection for 8h. The same samples were used for subsequent MS/MS analysis.
(E) Immunoblot against GFP or RFP after IP with anti-GFP or anti-RFP agarose beads on *N.*
benthamiana leaf extracts transiently expressing dTag-NACs. The same samples were used for
subsequent MS/MS analysis.
(F) Cladogram of top 30 GO terms for cellular components enriched in the list of *A. thaliana*
ortholog proteins found in NAC53/78 common interactome.
(G) Subcellular compartments enrichment in the list of *A. thaliana* proteins from NAC53/78
common interactome based on their SUBA5db annotation. Statistical significance is provided
through a chi-square testing.
(H-K) Heatmap representing the peptide enrichment (Log2FC) of the protein clusters from the
network Figure 2G.
Figure 3. ER-Anchored Protein Sorting (ERAPS) coordinates the subcellular fate of
NAC53 & NAC78.
(A) Confocal microscopy pictures of GFP-NAC53/78 expressing *A. thaliana* roots. 7 dag
seedlings were exposed to mock treatment, BTZ 10μM, CB-5083 10μM or LS-102 100μM for
3h. The treatments were repeated at least three times with similar observations. Scale bars =
10μm
(B) Immunoblot against GFP on crude extracts of GFP-NAC53/78 *A. thaliana* adult plants
infiltrated with mock treatment, BTZ 10μM, CB-5083 10μM or LS-102 100μM for 6h. Ponceau
staining is used as loading control.
(C) Confocal microscopy pictures of *N. benthamiana* transiently co-expressing GFP-
NAC53/78 full-length (FL) or deleted for the transmembrane domain (ΔTM) with RFP-HRD1a
or RFP-HRD1b. Pearson index (ρ) indicates correlation index at the orange lines.

(D) Co-IP of RFP-HRD1a/b with GFP-NAC53/78. IP was performed with anti-GFP beads. GFP-HDEL and free-GFP were used as negative control. The experiment was repeated twice with similar results.

(E) Immunoblot against GFP on crude extract of *N. benthamiana* leaves transiently expressing GFP-NAC53/78FL or ΔTM. Ponceau was used as loading control. Boxplots represent the relative ΔTM abundance compared FL.

(F) *In vitro* trans-ubiquitination assays of GST-HRD1a/b against MBP-NAC53/78. Removal of E1 and ATP from the reaction was used as negative control. The experiment was repeated twice with similar results.

(G) Representation of the ER-Anchored Protein Sorting (ERAPS) mechanism. NAC53/78 are ER-anchored proteins recognized and ubiquitinated by the HRD1 complex. The CDC48 complex subsequently extracts NAC53/78 from the ER to mediate ER-Associated Degradation or allowing nuclear translocation in the context of proteasomal stress to mitigate proteotoxicity.

Figure S2. NAC53/78 are subjected to high level of poly-ubiquitination. Related to Figure 3.

(A) *In vitro* analysis of LS-102 effect on HRD1a/b ubiquitination activity. Removal of E1 and ATP from the reaction was used as negative control.

(B) Quantification of the inhibitory effect of LS-102 (100μM) on HRD1 activity. Bar plots represent the mean of three independent replicates with error bars representing the standard deviation.

(C) Heatmaps representing the mean Log10 intensity of peptides bearing a di-glycine mark on the indicated lysin residue identified by MS/MS analysis of *in vitro* trans-ubiquitination reactions (see Figure 3F).

(D) Heatmaps representing the mean Log10 intensity of peptides bearing a di-glycine mark on the indicated lysin residue identified by MS/MS analysis of IP samples generated for Figure 2.

(E) Alignments of NAC53 and NAC78 peptide sequence regions around the identified lysin residues in panel C and D. Colors represent the residue conservation (green: conserved, shade of red: not conserved). All the identified lysins in one protein are conserved in the other. Underlined residues correspond to the mutated residues generated for the panel H and I.

(F) *In vitro* trans-ubiquitination assays of GST-HRD1a/b against MBP-NAC53/78 in the presence of LS-102 inhibitor. Removal of E1 and ATP from the reaction was used as negative control.

(G) Immunoblot analysis of *N. benthamiana* transiently expressing free-GFP, RFP-HDEL or dTag-NAC53/78 subjected to IP with anti-ubiquitin beads. The experiment was repeated twice with similar results.

(H) Immunoblot analysis of *N. benthamiana* transiently expressing GFP-NAC53/78 mutated for 5 lysins (Mut., see panel E) or not (WT) subjected to IP with anti- ubiquitin beads.

(I) Quantification of the relative ubiquitination levels assessed by the ratio between GFP signal and ubiquitin signal after IP as in panel H. Barplots represent the mean ratio of two replicates, normalized to the ratio of WT constructs with error bars representing the standard deviation.

Figure S3. NAC53/78 association with CDC48 is required for their nuclear translocation. Related to Figure 3.

(A) Confocal microscopy pictures of transgenic GFP-NAC53/78 root seedlings exposed to BTZ 10μM, BTZ 20μM, CB-5083 10μM or CB-5083 20μM for 3h. Scale bars = 10μm

(B) Confocal microscopy pictures of *N. benthamiana* leaves transiently co-expressing AtCDC48c-GFP with RFP-NAC53/78. Co-localization is visible in aggregate-like structures. Scale bars = 1μm

(C) Confocal microscopy representative pictures (used for panel D-E) of transgenic GFP-NAC53/78 roots treated with BTZ 10μM and CB-5083 10μM followed by Propidium Iodide staining. Scale bars = 10μm

(D-F) Boxplot representing the GFP signal quantification after confocal microscopy imaging
of transgenic GFP-NAC53/78 roots exposed to BTZ 10μM or CB-5083 10μM. Black dots
represent one cell. Statistical difference is assessed by a Wilcoxon-Mann-Whitney test.
Figure 4. The transcriptional landscape of proteotoxic stress
(A) Log2FC mRNA levels of *NAC53* and *NAC78* transcripts after *Pst* infection of adult Col-0
plant leaves. The heatmap represent the mean of 4 biological replicates.
(B) Log2FC mRNA levels of 26S proteasome transcripts after *Pst* infection of Col-0 or *nac53-*
1 78-1 adult plant leaves. The heatmap represent the mean of 4 biological replicates. The
experiment has been repeated 3 times with consistent results.
(C) Immunoblot against multiple 26S proteasome subunits on crude extracts of Col-0 or *nac53-*
1 78-1 adult plant leaves inoculated with *Pst* or mock solutions for 24h. Ponceau staining is
used as loading control.
(D) Relative Log2FC abundance of representative 26S proteasome subunits in Col-0 or *nac53-*
1 78-1 adult plant leaves. The abundance is calculated relative to the mock conditions.
Statistical differences are assessed via a Welch t.test (p values: n.s > 0.05, * < 0.05, ** < 0.01).
(E) Immunoblot against GFP on crude extract of *N. benthamiana* leaves transiently expressing
GFP-NAC53/78 after mock treatment or *PstΔHopQ* infection for 8h. Ponceau staining is used
as loading control.
(F) Immunoblot against GFP on extract of adult *A. thaliana* transgenics GFP-NAC53/78
subjected to mock treatment or *Pst* infection 8h by vacuum infiltration followed by IP with anti-
GFP beads.
(G) Number of differentially expressed genes (DEGs) in the different conditions of the *nac53-*
1 78-1 Pst RNAseq analysis. Total number of DEGs per condition is indicated. Colors indicate
the proportion of DEGs up-regulated and down-regulated. DEGs are considered when |Log2FC|
> 1.5. Genot. mock: *nac53-1 78-1* mock vs. Col-0 mock; WT *Pst*: Col-0 *Pst* vs. Col-0 mock;
Mut. *Pst*: *nac53-1 78-1 Pst* vs. *nac53-1 78-1* mock; Genot. *Pst*: *nac53-1 78-1 Pst* vs. Col-0 *Pst*.
(H) Level of differential expression (Log2FC) of the 945 DEGs found in Genot. *Pst* (see panel
G) for the several conditions analyzed.
(I) Cladogram of top 40 GO terms for BP enriched in the list of 945 DEGs. Colors represent
broad GO clusters.
(J) Venn diagram representing the overlap between the 945 DEGs and DEGs from *rpt2a-2 Pma*
(|Log2FC| > 0.5) or Col-0 MG132 24h (|Log2FC| > 0.5).
(K) Cladogram of top 30 GO terms for BP enriched in the list of 538 DEGs extracted from the
combined analysis (panel J). For conciseness terms are hidden and grouped in 4 processes.
(L) Protein network of the genes from the 945 DEGs list related to the GO BP terms from panel
K. Node colors refers to the 4 GO clusters from panel K.
Figure S4. Transcriptional analysis of proteotoxic stress reveals a trade-off between 26S
proteasome and other biological process. Related to Figure 4
(A) mRNA level Log2FC of all 26S proteasome genes and associated interactors/chaperones in
several conditions of the *nac53-1 78-1 Pst* RNAseq.
(B) Number of differentially expressed genes (DEGs) in the different conditions of the *nac53-*
1 78-1 Pst RNAseq analysis. Total number of DEGs per condition is indicated. Colors indicate
the proportion of DEGs up-regulated and down-regulated. DEGs are considered when |Log2FC|
> 1.5. Genot. mock: *rpt2a-2* mock vs. Col-0 mock; WT *Pma*: Col-0 *Pma* vs. Col-0 mock; Mut.
Pma: *rpt2a-2 Pma* vs. *nac53-1 78-1* mock; Genot. *Pma*: *rpt2a-2 Pma* vs. Col-0 *Pma*.
(C) mRNA levels Log2FC of the 981 DEGs found in Genot. *Pma* (see panel D) for the several
conditions analyzed.
(D) mRNA level Log2FC of all 26S proteasome genes and associated interactors/chaperones in
the several conditions of the *rpt2a-2 Pma* RNAseq.
(E) Cladogram of top 40 GO terms for BP enriched in the list of 945 DEGs. Colors represent
the broad GO clusters.

(F) mRNA level Log2FC of the genes present in the 4 network modules from Figure 4L in the
3 Transcriptome used.
Figure 5. NAC53 and NAC78 coordinate the regulation of proteasome and photosynthesis
gene expression through the same cis-element.
(A) Top 3 motifs found by STREME software in the promoter regions from the 4 network
clusters (Figure 4L). PRCE-like elements from proteasome and PhANGs clusters are
highlighted.
(B) 25bp region comprising the PRCE-like elements from LHCa3 and PSAD1 promoters used
in this figure. The region corresponds to the probe sequence used in panel C and D. The PRCE
is highlighted in red.
(C) Electro-Mobility Shift Assay (EMSA) of MBP-NAC53/78 with LHCa3/PSAD1 probes (see
panel B). As a negative control, probes were incubated without MBP-NAC53/78 (-) or with
MBP-NAC53/78 deleted for their DNA recognition motif (ΔRM). Experiment was repeated
twice with similar results.
(D) EMSA competition assay of MBP-NAC53/78 association with LHCa3/PSAD1 probes.
Competitor is applied at a concentration gradient (25X, 50X, 75X and 100X) indicated by the
shade of grey. WT and mutated competitors are labelled TGGGC and AAAAA, respectively.
Experiment was repeated twice with similar results.
(E) Boxplot representing association of GFP-NAC53/78 with LHCa3/PSAD1 PRCE in adult
A. thaliana transgenics lines infected with *Pst* or mock for 8h after Chromatin
Immunoprecipitation. The association is quantified as % of input and amplification of a region
in the CDS is used as negative control.
(F) Immunoblot against mature PhANGs proteins in Col-0 or *nac53-1 78-1* adult plant leaves
inoculated with *Pst* or mock solutions for 24h. Ponceau staining is used as loading control.
(G) Relative Log2FC abundance of PhANGs mature proteins immunoblotted in panel F. The
abundance is calculated relative to the mock conditions. Statistical differences are assessed via
a Welch t.test (p values: n.s > 0.05, * < 0.05, ** < 0.01).
(H) Photosystem II (PSII) activity measurement on Col-0 or *nac53-1 78-1* adult plants
inoculated with *Pst* or mock solutions for 24h. Letters indicate the statistical group assessed by
pairwise Welch t.test (p value < 0.05). The experiment was repeated 3 times with consistent
results.
(I) Representative pictures of measurements from panel I. Fv/Fm values are represented by
false color as indicated by the color gradient legend.
(J) Total chlorophylls and carotenoids from Col-0 or *nac53-1 78-1* adult plant leaves inoculated
with *Pst* or mock solutions for 24h. Letters indicate the statistical group assessed by pairwise
Wilcoxon-Mann-Whitney test (p value < 0.05). The experiment was repeated 3 times with
consistent results.
Figure S5. Characterization of the NAC53/78-PRCE module in PhANGs promoters.
Related to Figure 5.
(A) mRNA level Log2FC of the PhANGs clusters from Figure 4L in the 3 transcriptomes used.
Genes are separated based on their associated process in the photosynthetic pathway.
(B) Mapping of the PRCE-like motifs in PhANGs promoter regions from panel A. Promoter
regions are organized according to the order of the genes from panel A. Purple marks represent
the position of the motifs based on their distance from the transcription starting site (TSS). The
two red arrows highlight LHCa3 and PSAD1 promoter regions.
(C) Histograms representing the number of occurrences of the PRCE-like motifs across the 4
genes cluster from the network Figure 4L. Each bar represents a 20bp region, frequency *f* of
promoters bearing at least one motif is indicated as a percentage.
(D) LHCa3/PSAD1 promoter activity measured by luminescence in Col-0 or *nac53-1 78-1*
protoplasts extract transiently co-expressing LHCa3/PSAD1 reporter constructs and
NAC53/78ΔTM. Log2FC is calculated from the luminescence level of protoplasts expressing

the reporter construct alone. Replicates are a pool of at least two independent experiments.
Letters indicate the statistical group assessed by pairwise Wilcoxon-Mann-Whitney test (p
value < 0.05).
(E) Promoter activity as calculated in panel D for LHCa3/PSAD1 promoters deleted for their
respective DNA regions from Figure 5B. Replicates are a pool of at least two independent
experiments. Letters indicate the statistical group assessed by pairwise Wilcoxon-Mann-
Whitney test (p value < 0.05).
(F-G) Boxplot representing the Log2FC mRNA level the indicated transcripts in Col-0, GFP-
NAC53 or GFP-NAC78 seedlings treated with BTZ 10µM or mock solutions for 6h. Log2FC
is calculated from Col-0 mock. Letters indicate the statistical group assessed by pairwise
Wilcoxon-Mann-Whitney test (p value < 0.05).
(H) Fresh weight of seedlings from indicated genotypes grown under the indicated treatments
at 10 days after germination. Letters indicate the statistical group assessed by pairwise
Wilcoxon-Mann-Whitney test (p value < 0.05). Experiment was repeated 3 times with
consistent results.
(I) Representative phenotypic pictures related to panel D.
Figure 6. The proteasome autoregulatory feedback loop monitors photosynthesis
homeostasis during stress responses
(A) Photosynthetic pigment content from Col-0 or *nac53-1 78-1* adult plant leaves infiltrated
with Lin 200µM, BTZ 2µM or mock solutions for 24h. Content is expressed in percentage
relative to Col-0 mock. Letters indicate the statistical group assessed by pairwise Wilcoxon-
Mann-Whitney test (p value < 0.05). The experiment was repeated 3 times with consistent
results.
(B) Log2FC mRNA level of the indicated transcripts in Col-0, *nac53-1 78-1* adult plants leaves
Lin 200µM, BTZ 2µM or mock solutions for 24h. Log2FC is calculated from Col-0 mock and
PDH1 is used as negative control. Statistical differences are assessed with a Welch t-test (p
values: n.s. > 0.05, * < 0.05, ** < 0.01). The experiment was repeated twice with consistent
results.
(C) Fresh weight of seedlings grown under the indicated treatments at 10-14 day after
germination. The fresh weight is represented as a percentage of Col-0 mock conditions. Letters
indicate the statistical groups assessed by pairwise Wilcoxon-Mann-Whitney test (p value <
0.05). Every treatment has been repeated at least twice with consistent results. Representative
phenotypic pictures are included.
(D) Bacterial density shown as Log10 Colony-Forming-Unit (CFU) per leaf cm² in Col-0 and
hrd1a hrd1b plants. Statistical significance is assessed by a Welch t-test (p values: * < 0.05).
The experiment has been repeated three times with consistent results.
(E) Correlation heatmap of the 54 26S proteasome subunit genes and the expression of 68
PhANGs from publicly available 1223 RNAseq libraries identified in this study.
Figure S6. The proteasome autoregulatory feedback loop transcriptional signature is a
systemic response to environmental cues. Related to Figure 6.
(A) PSII activity measurement on Col-0 and *nac53-1 78-1* treated as indicated in Figure 6A.
Replicates are a pool of 3 independent experiments. Letters indicate the statistical group
assessed by pairwise Wilcoxon-Mann-Whitney test (p value < 0.05).
(B) PSII activity measurement on Col-0 and *hrd1a hrd1b* adult plant leaves inoculated with *Pst*
or mock solutions for 24h. Letters indicate the statistical group assessed by pairwise Welch
t.test (p value < 0.05). The experiment was repeated 2 times with consistent results.
(C) Confocal microscopy pictures of transgenic GFP-NAC53/78 *A. thaliana* roots exposed to
mock treatment, MV 100µM, GDA 100µM and Lin 100µM for 2h. The treatments were
repeated at least three times with similar observations. Scale bars = 10µm
(D) Density functions of the 54 26S proteasome genes and 68 PhANGs transcripts (Log2FC)
in the different transcriptome projects. Log2FC limits are -10 to +10. Treatment types and

relevant condition information are provided in the figure. Plots are separated by treatment
groups and subdivided by experiments.
Figure 7. ER-anchored protein sorting (ERAPS) controls the fate of two proteasome
activators for intracellular organelle communication during proteotoxic stress.
In steady state conditions, the ER-anchored protein sorting system (ERAPS) promotes the
constitutive degradation of NAC53/78 via the 26S proteasome. Meanwhile, PhANGs
expression is activated and subsequent chloroplastic import permits the maintenance of active
photosynthesis. Upon proteotoxic stress, the chloroplast import machinery is targeted for
proteasomal degradation. This leads to an accumulation of PhANGs precursors which are
therefore subjected to proteasomal degradation, inducing proteotoxicity. Thus, to avoid
proteotoxicity, the ERAPS system facilitates the nuclear translocation of NAC53/78, to activate
the production of a new proteasome complex and to repress PhANGs expression to mitigate
substrate accumulation.

1543

REFERENCES

Albert, S., Wietrzynski, W., Lee, C.W., Schaffer, M., Beck, F., Schuller, J.M., Salome, P.A.,
Plitzko, J.M., Baumeister, W., and Engel, B.D. (2020). Direct visualization of degradation
microcompartments at the ER membrane. Proc Natl Acad Sci U S A *117*, 1069-1080.

Baird, L., Tsujita, T., Kobayashi, E.H., Funayama, R., Nagashima, T., Nakayama, K., and
Yamamoto, M. (2017). A Homeostatic Shift Facilitates Endoplasmic Reticulum Proteostasis
through Transcriptional Integration of Proteostatic Stress Response Pathways. Mol Cell Biol
37.

Binder, A., Lambert, J., Morbitzer, R., Popp, C., Ott, T., Lahaye, T., and Parniske, M. (2014). A
modular plasmid assembly kit for multigene expression, gene silencing and silencing rescue in
plants. PLoS One *9*, e88218.

Boos, F., Kramer, L., Groh, C., Jung, F., Haberkant, P., Stein, F., Wollweber, F., Gackstatter, A.,
Zoller, E., van der Laan, M.*, et al.* (2019). Mitochondrial protein-induced stress triggers a global
adaptive transcriptional programme. Nat Cell Biol *21*, 442-451.

Bragoszewski, P., Turek, M., and Chacinska, A. (2017). Control of mitochondrial biogenesis
and function by the ubiquitin-proteasome system. Open Biol *7*.

Chen, Q., Zhong, Y., Wu, Y., Liu, L., Wang, P., Liu, R., Cui, F., Li, Q., Yang, X., Fang, S.*, et al.*
(2016). HRD1-mediated ERAD tuning of ER-bound E2 is conserved between plants and
mammals. Nat Plants *2*, 16094.

Christianson, J.C., and Ye, Y. (2014). Cleaning up in the endoplasmic reticulum: ubiquitin in
charge. Nat Struct Mol Biol *21*, 325-335.

Clough, S.J., and Bent, A.F. (1998). Floral dip: a simplified method for Agrobacterium-
mediated transformation of Arabidopsis thaliana. Plant J *16*, 735-743.

Colberg, L., Cammann, C., Greinacher, A., and Seifert, U. (2020). Structure and function of the
ubiquitin-proteasome system in platelets. J Thromb Haemost *18*, 771-780.

Diaz-Tielas, C., Grana, E., Sanchez-Moreiras, A.M., Reigosa, M.J., Vaughn, J.N., Pan, Z.,
Bajsa-Hirschel, J., Duke, M.V., and Duke, S.O. (2019). Transcriptome responses to the natural
phytotoxin t-chalcone in Arabidopsis thaliana L. Pest Manag Sci *75*, 2490-2504.

Duval, M., Hsieh, T.-F., Kim, S.Y., and Thomas, T.L. (2002). Molecular characterization of
AtNAM: a member of theArabidopsis NAC domain superfamily. Plant Molecular Biology *50*,
237-248.

Farmer, L.M., Book, A.J., Lee, K.H., Lin, Y.L., Fu, H., and Vierstra, R.D. (2010). The RAD23
family provides an essential connection between the 26S proteasome and ubiquitylated proteins
in Arabidopsis. Plant Cell *22*, 124-142.

Fonseca, D., and Carvalho, P. (2019). EGAD! There is an ERAD doppelganger in the Golgi.
EMBO J *38*, e102679.

Gallo-Oller, G., Ordoñez, R., and Dotor, J. (2018). A new background subtraction method for Western blot densitometry band quantification through image analysis software. Journal of Immunological Methods *457*, 1-5.

Ge, S.X., Jung, D., and Yao, R. (2020). ShinyGO: a graphical gene-set enrichment tool for animals and plants. Bioinformatics *36*, 2628-2629.

Gladman, N.P., Marshall, R.S., Lee, K.H., and Vierstra, R.D. (2016). The Proteasome Stress Regulon Is Controlled by a Pair of NAC Transcription Factors in Arabidopsis. Plant Cell *28*, 1279-1296.

Grimmer, J., Helm, S., Dobritzsch, D., Hause, G., Shema, G., Zahedi, R.P., and Baginsky, S. (2020). Mild proteasomal stress improves photosynthetic performance in Arabidopsis chloroplasts. Nat Commun *11*, 1662.

Gureev, A.P., Shaforostova, E.A., and Popov, V.N. (2019). Regulation of Mitochondrial Biogenesis as a Way for Active Longevity: Interaction Between the Nrf2 and PGC-1alpha Signaling Pathways. Front Genet *10*, 435.

Hao, Y.J., Song, Q.X., Chen, H.W., Zou, H.F., Wei, W., Kang, X.S., Ma, B., Zhang, W.K., Zhang, J.S., and Chen, S.Y. (2010). Plant NAC-type transcription factor proteins contain a NARD domain for repression of transcriptional activation. Planta *232*, 1033-1043.

Hristou, A., Grimmer, J., and Baginsky, S. (2020). The Secret Life of Chloroplast Precursor Proteins in the Cytosol. Mol Plant *13*, 1111-1113.

Hu, X., Zou, R., Zhang, Z., Ji, J., Li, J., Huo, X.-Y., Liu, D., Ge, M.-X., Cui, M.-K., Wu, M.-Z.*, et al.* (2023). UBE4A catalyzes NRF1 ubiquitination and facilitates DDI2-mediated NRF1 cleavage. Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms *1866*, 194937.

Jayaraj, G.G., Hipp, M.S., and Hartl, F.U. (2020). Functional Modules of the Proteostasis Network. Cold Spring Harb Perspect Biol *12*.

Kacprzak, S.M., Mochizuki, N., Naranjo, B., Xu, D., Leister, D., Kleine, T., Okamoto, H., and Terry, M.J. (2019). Plastid-to-Nucleus Retrograde Signalling during Chloroplast Biogenesis Does Not Require ABI4. Plant Physiol *179*, 18-23.

Kliza, K., Taumer, C., Pinzuti, I., Franz-Wachtel, M., Kunzelmann, S., Stieglitz, B., Macek, B., and Husnjak, K. (2017). Internally tagged ubiquitin: a tool to identify linear polyubiquitin-modified proteins by mass spectrometry. Nature Methods *14*, 504-512.

Kourelis, J., Kaschani, F., Grosse-Holz, F.M., Homma, F., Kaiser, M., and van der Hoorn, R.A.L. (2019). A homology-guided, genome-based proteome for improved proteomics in the alloploid Nicotiana benthamiana. BMC Genomics *20*, 722.

Koussevitzky, S., Nott, A., Mockler, T.C., Hong, F., Sachetto-Martins, G., Surpin, M., Lim, J., Mittler, R., and Chory, J. (2007). Signals from Chloroplasts Converge to Regulate Nuclear Gene Expression. Science *316*, 715-719.

Kramer, L., Groh, C., and Herrmann, J.M. (2021). The proteasome: friend and foe of mitochondrial biogenesis. FEBS Lett *595*, 1223-1238.

Kuhn, L., Vincent, T., Hammann, P., and Zuber, H. (2023). Exploring Protein Interactome Data with IPinquiry: Statistical Analysis and Data Visualization by Spectral Counts. In Statistical Analysis of Proteomic Data: Methods and Tools, T. Burger, ed. (New York, NY: Springer US), pp. 243-265.

Langin, G., Gonzalez-Fuente, M., and Üstün, S. (2023). The Plant Ubiquitin-Proteasome System as a Target for Microbial Manipulation. Annu Rev Phytopathol *61*, 351-375.

Langin, G., Gouguet, P., and Üstün, S. (2020). Microbial Effector Proteins - A Journey through the Proteolytic Landscape. Trends Microbiol *28*, 523-535.

Langin, G., and Üstün, S. (2023). A Pipeline to Monitor Proteasome Homeostasis in Plants. In Plant Proteostasis: Methods and Protocols, L.M. Lois, and M. Trujillo, eds. (New York, NY: Springer US), pp. 351-363.

Lee, K.H., Minami, A., Marshall, R.S., Book, A.J., Farmer, L.M., Walker, J.M., and Vierstra, R.D. (2011). The RPT2 subunit of the 26S proteasome directs complex assembly, histone

dynamics, and gametophyte and sporophyte development in Arabidopsis. Plant Cell *23*, 4298-4317.

Leister, D.S., A. (2003). From Genes to Photosynthesis in Arabidopsis thaliana. Int Rev Cytol *228*.

Leong, J.X., Raffeiner, M., Spinti, D., Langin, G., Franz-Wachtel, M., Guzman, A.R., Kim, J.G., Pandey, P., Minina, A.E., Macek, B., *et al.* (2022). A bacterial effector counteracts host autophagy by promoting degradation of an autophagy component. Embo J *41*, e110352.

Li, J., Yuan, J., Li, Y., Sun, H., Ma, T., Huai, J., Yang, W., Zhang, W., and Lin, R. (2022). The CDC48 complex mediates ubiquitin-dependent degradation of intra-chloroplast proteins in plants. Cell Rep *39*, 110664.

Liao, P.C., Wolken, D.M.A., Serrano, E., Srivastava, P., and Pon, L.A. (2020). Mitochondria-Associated Degradation Pathway (MAD) Function beyond the Outer Membrane. Cell Rep *32*, 107902.

Lichtenthaler, H.K. (1987). [34] Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. In Methods in Enzymology (Academic Press), pp. 350-382.

Ling, Q., Broad, W., Trosch, R., Topel, M., Demiral Sert, T., Lymperopoulos, P., Baldwin, A., and Jarvis, R.P. (2019). Ubiquitin-dependent chloroplast-associated protein degradation in plants. Science *363*.

Littlejohn, G.R., Breen, S., Smirnoff, N., and Grant, M. (2021). Chloroplast immunity illuminated. New Phytol *229*, 3088-3107.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology *15*, 550.

Ma, M., and Liu, Z.L. (2010). Comparative transcriptome profiling analyses during the lag phase uncover YAP1, PDR1, PDR3, RPN4, and HSF1 as key regulatory genes in genomic adaptation to the lignocellulose derived inhibitor HMF for Saccharomyces cerevisiae. BMC Genomics *11*, 660.

Maekawa, T., Kusakabe, M., Shimoda, Y., Sato, S., Tabata, S., Murooka, Y., and Hayashi, M. (2008). Polyubiquitin Promoter-Based Binary Vectors for Overexpression and Gene Silencing in Lotus japonicus. Molecular Plant-Microbe Interactions *21*, 375-382.

Mannhaupt, G., Schnall, R., Karpov, V., Vetter, I., and Feldmann, H. (1999). Rpn4p acts as a transcription factor by binding to PACE, a nonamer box found upstream of 26S proteasomal and other genes in yeast. FEBS Letters *450*, 27-34.

Marshall, R.S., Hua, Z., Mali, S., McLoughlin, F., and Vierstra, R.D. (2019). ATG8-Binding UIM Proteins Define a New Class of Autophagy Adaptors and Receptors. Cell *177*, 766-781 e724.

Marshall, R.S., and Vierstra, R.D. (2019). Dynamic Regulation of the 26S Proteasome: From Synthesis to Degradation. Front Mol Biosci *6*, 40.

Ng, S., Ivanova, A., Duncan, O., Law, S.R., Van Aken, O., De Clercq, I., Wang, Y., Carrie, C., Xu, L., Kmiec, B., *et al.* (2013). A membrane-bound NAC transcription factor, ANAC017, mediates mitochondrial retrograde signaling in Arabidopsis. Plant Cell *25*, 3450-3471.

Nguyen, H.M., Schippers, J.H., Goni-Ramos, O., Christoph, M.P., Dortay, H., van der Hoorn, R.A., and Mueller-Roeber, B. (2013). An upstream regulator of the 26S proteasome modulates organ size in Arabidopsis thaliana. Plant J *74*, 25-36.

Orosa, B., Üstün, S., Calderon Villalobos, L.I.A., Genschik, P., Gibbs, D., Holdsworth, M.J., Isono, E., Lois, M., Trujillo, M., and Sadanandom, A. (2020). Plant proteostasis - shaping the proteome: a research community aiming to understand molecular mechanisms that control protein abundance. New Phytol *227*, 1028-1033.

Page, M.T., Kacprzak, S.M., Mochizuki, N., Okamoto, H., Smith, A.G., and Terry, M.J. (2017). Seedlings Lacking the PTM Protein Do Not Show a genomes uncoupled (gun) Mutant Phenotype. Plant Physiol *174*, 21-26.

Pastor-Cantizano, N., Ko, D.K., Angelos, E., Pu, Y., and Brandizzi, F. (2020). Functional
Diversification of ER Stress Responses in Arabidopsis. Trends Biochem Sci *45*, 123-136.
Peters, L.Z., Hazan, R., Breker, M., Schuldiner, M., and Ben-Aroya, S. (2013). Formation and
dissociation of proteasome storage granules are regulated by cytosolic pH. J Cell Biol *201*, 663-
671.
Radhakrishnan, S.K., den Besten, W., and Deshaies, R.J. (2014). p97-dependent
retrotranslocation and proteolytic processing govern formation of active Nrf1 upon proteasome
inhibition. Elife *3*, e01856.
Raffeiner, M., Zhu, S., Gonzalez-Fuente, M., and Üstün, S. (2023). Interplay between
autophagy and proteasome during protein turnover. Trends Plant Sci *28*, 698-714.
Richter, A.S., Nagele, T., Grimm, B., Kaufmann, K., Schroda, M., Leister, D., and Kleine, T.
(2023). Retrograde signaling in plants: A critical review focusing on the GUN pathway and
beyond. Plant Commun *4*, 100511.
Ruvkun, G., and Lehrbach, N. (2023). Regulation and Functions of the ER-Associated Nrf1
Transcription Factor. Cold Spring Harb Perspect Biol *15*.
Sampaio-Marques, B., and Ludovico, P. (2018). Linking cellular proteostasis to yeast longevity.
FEMS Yeast Res *18*.
Schreiber, U. (2004). Pulse-Amplitude-Modulation (PAM) Fluorometry and Saturation Pulse
Method: An Overview. In Chlorophyll a Fluorescence: A Signature of Photosynthesis, G.C.
Papageorgiou, and Govindjee, eds. (Dordrecht: Springer Netherlands), pp. 279-319.
Schwanhäusser, B., Busse, D., Li, N., Dittmar, G., Schuchhardt, J., Wolf, J., Chen, W., and
Selbach, M. (2011). Global quantification of mammalian gene expression control. Nature *473*,
337-342.
Seo, P.J., Kim, S.G., and Park, C.M. (2008). Membrane-bound transcription factors in plants.
Trends Plant Sci *13*, 550-556.
Sha, Z., and Goldberg, A.L. (2014). Proteasome-mediated processing of Nrf1 is essential for
coordinate induction of all proteasome subunits and p97. Curr Biol *24*, 1573-1583.
Shih, C.F., Hsu, W.H., Peng, Y.J., and Yang, C.H. (2014). The NAC-like gene ANTHER
INDEHISCENCE FACTOR acts as a repressor that controls anther dehiscence by regulating
genes in the jasmonate biosynthesis pathway in Arabidopsis. J Exp Bot *65*, 621-639.
Shirozu, R., Yashiroda, H., and Murata, S. (2015). Identification of minimum Rpn4-responsive
elements in genes related to proteasome functions. FEBS Lett *589*, 933-940.
Shpilka, T., Du, Y., Yang, Q., Melber, A., Uma Naresh, N., Lavelle, J., Kim, S., Liu, P.,
Weidberg, H., Li, R.*, et al.* (2021). UPR(mt) scales mitochondrial network expansion with
protein synthesis via mitochondrial import in Caenorhabditis elegans. Nat Commun *12*, 479.
Steffen, J., Seeger, M., Koch, A., and Kruger, E. (2010). Proteasomal degradation is
transcriptionally controlled by TCF11 via an ERAD-dependent feedback loop. Mol Cell *40*,
147-158.
Su, W., Liu, Y., Xia, Y., Hong, Z., and Li, J. (2011). Conserved endoplasmic reticulum-
associated degradation system to eliminate mutated receptor-like kinases in Arabidopsis.
Proceedings of the National Academy of Sciences *108*, 870-875.
Sun, X., Feng, P., Xu, X., Guo, H., Ma, J., Chi, W., Lin, R., Lu, C., and Zhang, L. (2011). A
chloroplast envelope-bound PHD transcription factor mediates chloroplast signals to the
nucleus. Nat Commun *2*, 477.
Tanaka, K., Mizushima, T., and Saeki, Y. (2012). The proteasome: molecular machinery and
pathophysiological roles. Biol Chem *393*, 217-234.
Tokumaru, M., Adachi, F., Toda, M., Ito-Inaba, Y., Yazu, F., Hirosawa, Y., Sakakibara, Y., Suiko,
M., Kakizaki, T., and Inaba, T. (2017). Ubiquitin-Proteasome Dependent Regulation of the
GOLDEN2-LIKE 1 Transcription Factor in Response to Plastid Signals. Plant Physiol *173*,
524-535.

Ulasov, A.V., Rosenkranz, A.A., Georgiev, G.P., and Sobolev, A.S. (2022). Nrf2/Keap1/ARE
signaling: Towards specific regulation. Life Sci *291*, 120111.
Üstün, S., Hafrén, A., Liu, Q., Marshall, R.S., Minina, E.A., Bozhkov, P.V., Vierstra, R.D., and
Hofius, D. (2018). Bacteria Exploit Autophagy for Proteasome Degradation and Enhanced
Virulence in Plants. Plant Cell *30*, 668-685.
Üstün, S., Sheikh, A., Gimenez-Ibanez, S., Jones, A., Ntoukakis, V., and Börnke, F. (2016). The
Proteasome Acts as a Hub for Plant Immunity and Is Targeted by Pseudomonas Type III
Effectors. Plant Physiol *172*, 1941-1958.
Wang, X., Xu, H., Ju, D., and Xie, Y. (2008). Disruption of Rpn4-induced proteasome
expression in Saccharomyces cerevisiae reduces cell viability under stressed conditions.
Genetics *180*, 1945-1953.
Wang, X., Yen, J., Kaiser, P., and Huang, L. (2010). Regulation of the 26S proteasome complex
during oxidative stress. Sci Signal *3*, ra88.
Waters, M.T., Wang, P., Korkaric, M., Capper, R.G., Saunders, N.J., and Langdale, J.A. (2009).
GLK transcription factors coordinate expression of the photosynthetic apparatus in
Arabidopsis. Plant Cell *21*, 1109-1128.
Weidberg, H., and Amon, A. (2018). MitoCPR-A surveillance pathway that protects
mitochondria in response to protein import stress. Science *360*.
Welner, D.H., Deeba, F., Lo Leggio, L., and Skriver, K. (2016). NAC Transcription Factors:
From Structure to Function in Stress-Associated Networks. In Plant Transcription Factors, pp.
199-212.
Woodson, J.D., and Chory, J. (2012). Organelle signaling: how stressed chloroplasts
communicate with the nucleus. Curr Biol *22*, R690-692.
Wu, G.Z., Meyer, E.H., Richter, A.S., Schuster, M., Ling, Q., Schottler, M.A., Walther, D.,
Zoschke, R., Grimm, B., Jarvis, R.P.*, et al.* (2019). Control of retrograde signalling by protein
import and cytosolic folding stress. Nat Plants *5*, 525-538.
Xie, Y., and Varshavsky, A. (2001). RPN4 is a ligand, substrate, and transcriptional regulator of
the 26S proteasome: A negative feedback circuit. Proceedings of the National Academy of
Sciences *98*, 3056-3061.
Xie, Y.V., A. (2001). RPN4 is a ligand, substrate, and transcriptional regulator of the 26S
proteasome: A negative feedback circuit. Proc Natl Acad Sci U S A *98*, 3056-3061.
Yagishita, N., Aratani, S., Leach, C., Amano, T., Yamano, Y., Nakatani, K., Nishioka, K., and
Nakajima, T. (2012). RING-finger type E3 ubiquitin ligase inhibitors as novel candidates for
the treatment of rheumatoid arthritis. Int J Mol Med *30*, 1281-1286.
Yoo, S.D., Cho, Y.H., and Sheen, J. (2007). Arabidopsis mesophyll protoplasts: a versatile cell
system for transient gene expression analysis. Nat Protoc *2*, 1565-1572.
Zhang, H., Zhang, F., Yu, Y., Feng, L., Jia, J., Liu, B., Li, B., Guo, H., and Zhai, J. (2020). A
Comprehensive Online Database for Exploring approximately 20,000 Public Arabidopsis
RNA-Seq Libraries. Mol Plant *13*, 1231-1233.
Zhou, H.J., Wang, J., Yao, B., Wong, S., Djakovic, S., Kumar, B., Rice, J., Valle, E., Soriano,
F., Menon, M.K.*, et al.* (2015). Discovery of a First-in-Class, Potent, Selective, and Orally
Bioavailable Inhibitor of the p97 AAA ATPase (CB-5083). J Med Chem *58*, 9480-9497.
1773
1774