1 Assembly processes inferred from eDNA surveys of a

2 pond metacommunity are consistent with known

3 species ecologies

Authors: Wang Cai[1,2], Maximilian Pichler[3], Jeremy Biggs[4,5], Pascale Nicolet[4], Naomi

Ewald[4], Richard A. Griffiths[6], Alex Bush[7], Mathew A. Leibold[8], Florian Hartig[3], Douglas W.

Yu[1,9,10,11]*

Author affiliations:

[1] State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology,
Chinese Academy of Sciences, Kunming, China
[2] Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden,
Chinese Academy of Sciences, Mengla 666303, China
[3] Theoretical Ecology, University of Regensburg, Regensburg, Germany
[4] Freshwater Habitats Trust, First Floor, Bury Knowle House, North Place, Old High Street,
Headington, Oxford, UK
[5] Oxford Brookes University, Headington, Oxford, OX3 0BP, UK
[6] Durrell Institute of Conservation and Ecology, School of Anthropology and Conservation,
University of Kent, UK
[7] Lancaster Environment Centre, Lancaster University, UK
[8] Department of Integrative Biology, University of Texas at Austin, Austin, TX USA
[9] Center for Excellence in Animal Evolution and Genetics, Chinese Academy of Sciences,
Kunming, China
[10] Yunnan Key Laboratory of Biodiversity and Ecological Security of Gaoligong Mountain,
Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, China
[11] School of Biological Sciences, University of East Anglia, Norwich Research Park,
Norwich, UK

*__Correspondence:__ email: dougwyu@mac.com; address: School of Biological Sciences,
Univ. of East Anglia, Norwich, Norfolk NR4 7TJ UK

KEYWORDS: Aquatic eDNA, metabarcoding, novel community data, biodiversity, pond,
joint species distribution model (JSDM), macroecology, *Triturus cristatus*

31 Abstract

Technological progress is enabling ecologists to create repeated, large-scale, structured, and

standardised community surveys. However, it is unclear how best to extract information from

these novel community data. We metabarcoded 48 vertebrate species from their eDNA in 320

ponds in England and applied the 'internal-structure' approach, which uses joint species

distribution models to explain community compositions as the outcome of four
metacommunity assembly processes: environmental filtering, dispersal, species interactions,
and stochasticity. We find that the environment plays an important role in community assembly
and that the inferred environmental preferences of species are consistent with their ecologies.
We also infer negative biotic covariances between fish and amphibians, which is consistent
with predator-prey interactions, and high spatial autocorrelation for the palmate newt, which is
consistent with its hypothesised relictual distribution. Comparing sites in the metacommunity,
environmentally and spatially distinctive sites are better explained by their environmental
covariates and geographic locations, respectively, revealing sites where environmental filtering
and dispersal limitation act more strongly. Furthermore, species belonging to different trait
groups differ in how well environmental covariates, biotic covariances, and geographical
locations explain their distributions. Overall, our results highlight the value of a modern
interpretation of metacommunity ecology that embraces the fact that assembly processes differ
between individual species and sites. We discuss how novel community data make feasible
several study-design improvements that will strengthen the inference of metacommunity
assembly processes from observational data.

Introduction

Metacommunity theory, which explicitly models feedback between local communities and

regional species pools, has been proposed as a unifying theory of spatial community ecology

(Leibold *et al.* 2004; Leibold & Chase 2018). In the framework of this theory, we consider a

set of communities whose local population dynamics are governed by environmental filtering,

species interactions, and ecological drift and that are additionally linked by dispersal. The goal

of the theory is to understand how these four basic assembly processes determine species

compositions in the metacommunity (Vellend 2016). Traditionally, this has been done with

classical community data gathered by human observers, but the fact that modern sensors such

as eDNA deliver community observations that are ideally suited for metacommunity analysis

has created excitement in the field and also makes metacommunity analysis interesting for

molecular ecologists (Hartig *et al.* 2024).

Empirical approaches to studying metacommunities mainly aim at inferring the relative

contributions of the four assembly processes (dispersal, environmental filtering, species

interactions, and drift) from empirical data. Examples of these are analyses based on

community summary statistics, such as ordinations that describe different metacommunities

using centroids and distances, and alpha and beta diversities (Figure 1). Another common

approach to analyse metacommunity data is variation partitioning, where classically,

community composition is explained by metacommunity-level contributions of environmental

and spatial factors (e.g. Cottenie 2005). However, those approaches exhibit limited power to

reveal assembly processes (Guzman *et al.* 2022; Ovaskainen *et al.* 2019), in part because

summary metrics cannot reveal how the four processes differentially affect individual species

and sites. Leibold *et al.* (2022) refer to such metacommunity-level metrics as studying the

'external structure' of metacommunities, because they assume that there is one "average" mix

of assembly processes that is the same across sites and species.

To avoid averaging assembly processes across sites and species, Leibold *et al.* (2022) propose

to study the 'internal structure' of metacommunities, which dissects the importance of different

assembly processes *by each species and site*. Technically, this can be done by using a joint

species distribution model (JSDM) to partition the varying contributions of three model

components (environmental covariates, species covariances, and spatial autocorrelation) to

explaining species presence/absence, for each individual species and site (Figure 1). Among

other things, *this approach allows one to relate environmental differences between sites, and*

trait differences between species, to differences in the variation explained by each component,
thereby generating testable hypotheses that link distribution patterns to metacommunity
assembly processes.

Simulation studies have shown that internal-structure analysis can indeed differentiate
synthetic metacommunities that differ in site environmental distinctiveness and in species niche
breadth, dispersal ability, niche centrality, and the presence or absence of competitive
interactions (Figure 1, Leibold *et al.* 2022; Terry *et al.* 2023). While these simulation results
are encouraging, real metacommunity datasets have more complicated properties, including
detection failures, measurement errors, and model uncertainty; not all species, environmental
covariates, and sites can be included; species interact in multiple ways; and real
metacommunities might be non-stationary, not least because of climate change (Abrego *et al.*
2021; Kadoya *et al.* 2024; Terry *et al.* 2023). Thus, it is important to gain more experience
about the applicability of the internal structure idea to real data.

An ideal empirical metacommunity dataset for inferring internal structure would consist of (1)
many local-community inventories with standardised species presence-absence or abundance
information, (2) over an area that is connected (and large) enough for dispersal (and dispersal
limitation) to operate, (3) with the taxonomic breadth to include interacting guilds such as
predators and prey, and (4) measures of local environmental conditions relevant to the niche
requirements of all these species. An exemplary study is provided by Kadoya et al. (2024) who
applied internal-structure analysis to gillnet survey data covering three countries, 93 fish
species, and 1,853 lakes and found that environmental covariates explained the most variation
in species distributions and lake compositions, highlighting the importance of environmental
filtering. Kadoya et al. then projected the effect of future climate heating on lake species
compositions by running the fitted model with higher values of the degree-days environmental
covariate while using the biotic covariances to simulate the effect of species interactions.

A promising alternative to traditional community observations is eDNA metabarcoding, which
can generate repeated, large-scale, structured, and standardised community surveys (Hartig *et*
al. 2024), but eDNA has rarely been used in metacommunity ecology. To our knowledge, the
only attempt is by Vass *et al.* (2022), who metabarcoded aquatic fungi from four separate inlets
along the Swedish coast and found in an internal-structure analysis that most species and site
variance was attributable to environmental covariates, again highlighting the importance of
environmental filtering. Vass *et al.* also found high biotic covariance values for some species,
which is consistent with additional contributions from species interactions or unmeasured

environmental covariates (Blanchet *et al.* 2020; Dormann *et al.* 2018; Hartig *et al.* 2024;
Poggiato *et al.* 2021; Zurell *et al.* 2018).

Here, we test how well the internal structure of a candidate pond metacommunity surveyed
with eDNA matches expectations derived from external knowledge of species ecologies. Our
survey data come from ponds in the South Midlands of England that were originally sampled
to detect the great crested newt (*Triturus cristatus*), a UK-protected amphibian species that
breeds in ponds (Biggs *et al.* 2015). We metabarcoded the residual eDNA to detect vertebrates,
generating a community matrix of 320 ponds X 48 vertebrate species. Each pond was associated
with 8 environmental covariates and a geographic location, allowing us to fit three data
matrices in a JSDM (Figure 1).

Among other results, we report at the species level that the fitted environmental covariates are
consistent with known fish and amphibian ecologies, that the negative biotic covariances
between fish and amphibians are consistent with expected predator-prey relationship between
the two taxa, and that the high spatial autocorrelation for the palmate newt is consistent with a
hypothesised relictual distribution for this species. At the site level, environmentally distinctive
ponds are better explained by their environmental covariates, and spatially distinctive ponds
are better explained by their geographic locations, revealing sites where environmental filtering
and dispersal limitation act more strongly, respectively.

Methods

Environmental covariates

To quantify land cover around each pond, we used Rowland *et al.*'s (2017) 21 UK land classes.
For each pond, we calculated the proportions of land class within a 500 m radius of its point
location and used principal component analysis in {FactoMineR} v.2.4 (Lê *et al.* 2008) to
extract the top three principal components (accounting for 40% of total variation, Figure S1),
which correlate with the degree of agriculture vs. urban cover, grassland cover, and woodland
cover. Each pond was also scored during sampling for ten standard pond variables used by
surveyors to calculate the pond's Habitat Suitability Index (HSI) for the great crested newt
(ARG-UK 2010), of which we used five (Table S1).

Pond sampling and newt assays

The pond water samples were the result of a single-season, great crested newt survey of 544

ponds in the South Midlands of England, UK in 2017, covering ponds in urban/suburban,

woodland and agricultural areas. The survey used a grid design to generate single-species

presence-absence data as input to a species distribution model that underpins a biodiversity

offset market (Box 3 in Hartig *et al.* 2024). Samples were collected and processed following

Biggs *et al.* (2015) and were stored at ambient temperature until shipped to a commercial lab

(NatureMetrics, Egham, UK), and DNA was extracted using a precipitation protocol (Tréguier

et al. 2014), after which each sample's DNA was cleaned and subjected to 12 separate qPCR

tests. After the qPCR assays, the residual eDNA was stored at -80°C. We loosely estimate that

the cost of sample collection and great crested newt assays, including effort getting permission

to sample private properties, exceeded US$120,000. Pond eDNA collected for great crested

newts thus has several advantages for metacommunity studies in that we have a standardised,

professionally collected, commercially processed, and spatially contiguous set of eDNA

samples, with standardised environmental covariates.

Metabarcoding assays

In 2019, the residual eDNA samples were retrieved and subjected to metabarcoding at

NatureMetrics (PCR) and at Kunming Institute of Zoology (library prep), using a two-step

protocol targeting a 73-110 bp fragment of 12S ribosomal RNA (Riaz *et al.* 2011). In the first

step, 12 replicates per sample were amplified and then pooled to maximise the targeted

sequence (Lahoz-Monfort *et al.* 2016). In the second step, three independent PCR replicates

were performed on the pooled samples using twin-tagging (Yang *et al.* 2021; Zepeda-Mendoza

et al. 2016). The three independent (separately twin-tagged) PCR replicates per sample in the

second step were pooled into three approximately equimolar libraries for bead purification,

library preparation, and sequencing on an Illumina HiSeq platform (PE150) at Novogene

Tianjin, China.

We processed the raw sequence data with the modified DAMe bioinformatics pipeline of Cai

et al. (2021), where we filtered out PCR and sequencing errors by retaining only sequences

present in ≥2 PCR replicates with a minimum copy number of 25 per PCR. After sequence

clustering, we generated a table of 540 ponds by 74 OTUs (operational taxonomic units). We

assigned taxonomies to the OTUs using PROTAX (Axtner *et al.* 2019; Somervuo *et al.* 2017),

setting prior probabilities to 0.90 for a list of expected UK vertebrate species (Harper *et al.*

2018).

Joint species distribution modelling

To fit JSDMs to the observed community data, we converted OTU read counts to presence-

absence data. We retained only OTUs present in ≥5 ponds and only sites with ≥1 targeted OTU

(= vertebrate species present in the United Kingdom), which reduced the number of OTUs from

74 to 48 and the number of ponds from 540 to 320. We assigned species-level taxonomies to

OTUs that received ≥98% PROTAX probability of species assignment, and we classified the

OTUs into 6 trait groups: fish, amphibians, perching birds, waterfowl, mammals, and domestic

species. Domestic species are whose distributions we deemed as determined largely by humans

(Table S2).

We fit our data with two distinct JSDM structures. One model was fit to all the species in our

dataset, terrestrial and aquatic (320 ponds X 48 species), and the other model fit to only the

aquatic species (amphibians and fish, 279 ponds X 15 species; fewer ponds because we

excluded those without aquatic species). *A priori*, aquatic species should be more likely to be

filtered by pond characteristics, which make up 5 of our 8 environmental covariates. Thus, we

expect the aquatic species, especially the fish, to act more like narrow-niche species (Figure

1C, top row), and the terrestrial species to act more like broad-niche species (Figure 1C, bottom

row).

All models were fitted using {sjSDM} v.1.0.6 (Pichler & Hartig 2021) running under R 4.2.2

(R Core Team 2022). We used a binomial likelihood and a multivariate probit link, linear main

effects for the eight environmental covariates, and a DNN (deep neural net) spatial model (trend

surface model). To avoid overfitting, a light elastic net regularisation (Zou & Hastie 2005) was

applied to all regression slopes and weights of the DNN (model fitting details in Appendix S1).

Internal structure of the metacommunity

After fitting the two models (aquatic, terrestrial + aquatic), we used the ANOVA functions

implemented in sjSDM (based on Leibold *et al.* 2022) to partition the variation of each species'

distribution and each site's composition across environmental covariates (E), spatial

autocorrelation (S), and biotic covariances (C), without the shared fractions of the pairwise

components. The results are visualised using ternary plots, where the positions of the species

and sites reveal the relative contributions of E, S, and C: the metacommunity's internal
structure (Figure 1B2).

Leibold *et al*. (2022) found that environmentally more distinctive sites (at the ends of their one
niche axis) received higher E values (implying a greater contribution of environmental
filtering). To test for this result in a natural metacommunity, we regressed the individual pond
absolute E, S, and C R^2 values against pond environmental distinctiveness using quantile
regression (50% quantile) (Fasiolo *et al.* 2021). We also tested the parallel hypothesis that
geographically distinctive sites would have higher absolute S R^2 values (implying a greater
contribution of dispersal limitation). In both cases, we defined distinctiveness as the leading
eigenvector of the corresponding environmental or geographical Euclidean distance matrix.
We note that if predictor variables are collinear, bivariate correlations can be spurious and
partial correlations calculated using multiple regressions should be preferred. However,
environmental and geographic distinctiveness showed practically no collinearity (Figure S2).

Model generality

Our results are the outputs of a complex model that includes a linear environmental structure
with eight environmental covariates and a DNN spatial structure with 30 x 2 layers. Complex
models run a risk of overfitting, so to estimate the risk of overfitting after elastic-net
regularisation, we carried out a 20-fold cross-validation test with stratified multi-label sampling
(Gunopulos *et al.* 2011; Szymański & Kajdanowicz 2017). In each of 20 rounds, 19 folds (304
sites) were used as a training dataset, and the fitted model was used to predict species
compositions in the hold-out fold (16 sites). Model fitting used the same structure,
regularisation strengths, and scaled environmental and spatial covariates as the original full
model. For each round and species, we calculated from the training dataset an explanatory-
performance AUC (Area Under the Curve) metric. We also used each hold-out fold to calculate
a predictive-performance AUC for each species: a measure of model generality. The final
explanatory and predictive AUCs per species are the means over 20 folds. Species with higher
predictive AUCs are those whose fitted models are more general (details in Appendix S1).

Results

Internal structure of the pond metacommunity

When analysing aquatic species only, we find that the fish are bimodally arrayed along the E-
C axis, with four species relatively better explained by environmental covariates (higher E

values), and six species relatively better explained by biotic covariances (higher C values),
which reflect the effects of unknown environmental covariates plus possible species
interactions (Figure 2A).

In contrast, none of the five amphibian species shows a high contribution of either
environmental covariates or biotic covariances, but relative to fishes, amphibians show greater
contributions of dispersal limitation (higher S values), especially the palmate newt (*Lissotriton*
helveticus), whose distribution is mostly explained by dispersal limitation. This species was
detected in 16 ponds, in three separate sections of the survey area (Figure S3), and its high S
value indicates that other ponds in which this species was not detected have environmental
conditions similar to those in which the palmate newt was detected.

Including terrestrial species in the analysis (Figure 2C) increases the relative contribution of
biotic covariance for both fish and amphibians, which could reflect either the contributions of
species interactions with terrestrial species or more unmeasured environmental covariates that
have been revealed by adding the terrestrial species. The terrestrial species themselves also
largely range along the E-C axis, with no clear clustering by trait group. Like the aquatic species,
only one terrestrial species, Mandarin duck (*Aix galericulata*), has a high S value (Figure S3)
and is found in only five ponds. For the two site ternary plots (Figure 2B, D), the general effect
of adding terrestrial species is an increase in the variance accounted for by biotic covariances
(site points shift upwards toward C).

We now examine species and site variation to try to infer some of the assembly processes that
have resulted in these observed internal structures.

Estimated environmental preferences

In the aquatic-only model (Figure 3A), the pond effects for the fish species are in the direction
of greater prevalence in *larger* ponds with *lower* risk of drying and *less* macrophyte cover.
Several of the fish species are known to eat macrophytes, reduce macrophyte cover through
other behaviours, and/or require higher oxygen with less macrophyte cover (Lopes *et al.* 2015;
Maceda-Veiga *et al.* 2017; Stefanoudis *et al.* 2017). In contrast, for the amphibians, pond
effects are in the direction of greater prevalence in *smaller* ponds with *higher* macrophyte cover.
Pond drying risk, water quality, and shade showed essentially no effects on amphibian
prevalence. Most of the effects of land cover on fish species are in the direction of lower
prevalence in areas surrounded by agriculture or grassland. Grassland here is dominated by the
'improved grassland' subtype (Rowland *et al.* 2017), characterised by regular fertilisation,

herbicide application, and drainage (DEFRA & Rural Payments Agency 2022). For amphibians,
the only effects are in the direction of greater prevalence of palmate newt and common frog in
ponds bordered by woodland. All five amphibian species may utilise woodland outside the
breeding season (Mulkeen *et al.* 2017; Oldham *et al.* 2000).

In the aquatic+terrestrial model (Figure 3B), the effects of the environmental covariates on fish
and amphibians remain largely the same as in the aquatic-only model. For the terrestrial species,
most of the significant environmental-covariate effects are shade (% of pond perimeter shaded
by trees), macrophyte cover, and land cover. Shade, which affects many perching birds and the
grey squirrel, is most parsimoniously interpreted as increasing species detectabilities.
Macrophyte cover is positively correlated with three rodent species and the grey heron and
negatively correlated with rabbits and ducks (Anatidae sp.). Land-cover effects are variable
across species, but we note that cows and sheep have higher prevalences in ponds bordered by
('improved') grassland.

Finally, we observe that the Eurasian coot (*Fulica atra*) has higher prevalences at ponds of
higher water quality, which in the HSI scoring system is inferred from the presence of an
"abundant and diverse invertebrate community" (ARG-UK 2010).

Biotic covariances
We visualise the residual biotic covariances in pairwise correlation plots (Figure 4), where
large absolute correlation-coefficient values correlate with high C-values in the internal-
structure ternary plots (Figure 2A and 2C; linear model, aquatic species only, $R^2 = 0.726$, P <
0.001; aquatic + terrestrial, $R^2 = 0.201$, P < 0.001).

In the aquatic species model (Figure 4A, B) and after filtering to the 2.5% most negative and
positive values, the three surviving negative correlations are between the common frog (*Rana*
temporaria) and two omnivorous fish species (*Carassius carassius* and *Cyprinus carpio*) and
between great crested newt (*Triturus cristatus*) and a carnivorous fish (*Esox lucius*). There are
also three surviving positive correlations between fish species, which we conservatively
interpret as indicating unmeasured environmental covariates.

In the aquatic+terrestrial species model (Figure 4C, D) and after filtering to the 2.5% most
negative and positive values, there are four surviving negative correlations between amphibians
and fish. The common frog is negatively correlated with two omnivorous fish species
(*Cyprinus carpio* and *Rutilus rutilus*), and great crested newt (*Triturus cristatus*) is negatively
correlated with two carnivorous/omnivorous fish (*Esox lucius* and *Tinca tinca*). Most of the

surviving positive correlations occur among the fish species and among the bird species, which
we again interpret as unmeasured environmental covariates. Also notable are negative
correlations between several fish species with ring-necked pheasant (*Phasianus colchicus*) and
red fox (*Vulpes vulpes*).

Relating distribution patterns to assembly processes, by site and species group

By site, the absolute R^2 explained by the environment increases significantly with the
environmental distinctiveness of the site (Figure 5A), and the absolute R^2 explained by space
increases significantly with the geographical distinctiveness of the site (Figure 5B). In other
words, environmental filtering appears to be an increasingly more important assembly process
for more environmentally distinctive sites, as predicted by Leibold et al. (2022), and dispersal
limitation appears to be increasingly more important for geographically distinctive sites. Given
this environment effect, we *post-hoc* tested each covariate individually and found that the
absolute R^2 explained by the environment increases only with pond area (Figures 5C, S4),
suggesting that the species compositions of large ponds are determined more strongly by
environmental filtering.

By species group, the absolute R^2 explained by biotic covariances is greatest for fish,
amphibians, and waterfowl, and about equal with the R^2 explained by the environment for the
other three trait groups (Figure 5D), which is consistent with species distributions being
primarily governed by a combination of environmental filtering and (to a lesser extent) species
interactions. Interestingly, the most pond-dependent species (fish, amphibians, waterfowl) are
the least well explained by our environmental covariates, although there are individual
exceptions (Figure 2A, C).

Model generality

For the aquatic-only model, explanatory AUCs are always slightly greater than predictive
AUCs, and explanatory and predictive AUCs are positively correlated (linear model, adjusted
R^2=0.546, P=0.001) (Figure 6A). For the aquatic+terrestrial model, explanatory AUCs are
again still always greater than predictive AUCs, but the correlation weakens considerably
(linear model, adjusted R^2=0.139, P=0.005), and for some terrestrial species, the model makes
worse-than-random predictions (predictive AUCs<0.5) (Figure 6B). The risk of overfitting is
greater for low-predictive-AUC species, so the risk is greater for terrestrial species. Both
models are generally better at predicting fish distributions than amphibian distributions.
Palmate newt (*Lissotriton helveticus*) is an exception, with the overall second-highest

predictive AUC (Figure 6A), which is probably explained by the large contribution of space to

its explained variance. Stickleback (*Gasterosteus aculeatus*) and European perch (*Perca*

fluviatilis) are exceptions for fish, with the overall two lowest predictive AUCs in the aquatic

model (Figure 6A).

Discussion

The goals of our study were to reveal the internal structure of a real metacommunity, which

infers the importance of different assembly processes per species and site. We used a joint

species distribution model to estimate these relative and absolute contributions of

environmental covariates, biotic covariances, and space for explaining spatial variation in pond

compositions. Pondscapes are convenient study systems because (1) each pond is

unambiguously identified as a local community, (2) there is an *a priori* division between likely

pond niche specialists (aquatic species) versus generalists (terrestrial species), (3) aquatic

eDNA metabarcoding can efficiently generate hundreds of local-community inventories, and

(4) the detected species encompass multiple trophic levels, increasing the possibility of

detecting species interactions.

Importance of taxonomic breath

We inferred two internal structures, one for aquatic species only and one for aquatic + terrestrial

species (Figure 2). This gives us our first conclusion, which is that the 'one average mix'

approach to metacommunities indeed loses the useful information revealed in species and site

variation (Leibold *et al.* 2022). In this pond metacommunity, even if one makes the extreme

assumption that all the variation partitioned to biotic covariance is also environmental filtering

(but unmeasured), the species and sites still vary in their inferred degrees of dispersal limitation,

and thus no single metacommunity paradigm (i.e. species-sorting, mass-effect, patch-dynamic,

and neutral communities; Holyoak *et al.* 2005; Shoemaker & Melbourne 2016; Suzuki &

Economo 2021; Thompson *et al.* 2020) can serve as an adequate description.

Influence of environmental effects

Looking at how metacommunity assembly processes vary across sites and species, both of the

internal structures (Figure 2A, C) suggest that environmental filtering is an important

structuring force for many of the species in this pondscape. These findings are supported by

the fact that several of the environmental-covariate coefficient values are consistent with

known biology. For instance, relative to amphibians, fish have a higher prevalence in larger,

more permanent ponds (Figure 3). Macrophyte cover is also lower in ponds with higher
prevalence of fish over amphibians, although the direction of causality might vary by species,
since some fish species can directly reduce macrophyte cover. High macrophyte cover also
appears to increase the prevalence of multiple terrestrial and waterfowl species (Figure 3) On
the other hand, and contrary to its inclusion as a contributor to the suitability of habitats for
newts (ARG-UK 2010), higher water quality is not associated with higher prevalence of any
amphibian species, although it is associated with higher prevalences of Eurasian coot (Figure
3).

If environmental filtering is an important assembly process in this metacommunity, Leibold et
al.'s (2022) corollary is that more environmentally distinctive sites should be more strongly
determined by environmental filtering. This is what we observe (Figure 5A). In fact, this result
holds separately for four of the six trait groups (Figure S5), the exceptions being fish, for which
there is no relationship, and waterfowl, for which *spatial* site R^2 increases with environmental
distinctiveness.

378   Influence on space on the community assembly
Looking at the contribution of space, which can be interpreted as a proxy for dispersal
limitation, we find that spatial effects are dominant for two species, palmate newt and mandarin
duck (Figure 2). The spatial distribution of the 16 palmate newt detections is visibly patchy
(Figure S3), which may be due to the persistence of relictual populations with a historic
distribution associated with woodland (Beebee & Griffiths 2000). There are only five
detections for mandarin duck (Figure S3), so it is unclear whether such a spatial pattern implies
dispersal limitation, but this introduced and free-living species is described as largely sedentary
in the UK (Harris 2024).

In parallel to the environmental distinctiveness test, we also found that dispersal limitation
appears to be stronger in geographically distinctive sites (Figure 5B).

389   Influence of co-distribution on community assembly
With the main exception of common toad (*Bufo bufo*), which is protected from fish predation
by toxins, residual associations between amphibians and fish are mostly negative, and three or
four biotic correlations survive after filtering to the most extreme values (Figure 4). Keeping
in mind the caveat that biotic covariances should not be taken as direct evidence for species
interactions (Blanchet *et al.* 2020; Dormann *et al.* 2018; Hartig *et al.* 2024; Poggiato *et al.* 2021;

Zurell *et al.* 2018), it is plausible that these associations arise from species interactions since
the fish species involved are omnivorous or carnivorous and could thus prey on amphibian
eggs, larvae, or possibly adults.

What can eDNA bring to metacommunity ecology and internal structure analysis?

The large gains from eDNA metabarcoding in efficiency and error homogeneity over
traditional survey methods make it feasible to generate datasets with many samples and many
species, which can strengthen the inference of metacommunity assembly processes. Most
obviously, the large number of species detectable with eDNA increases the probability of
detecting sets of species that are truly interacting, as suggested by the negative correlations
between predatory and omnivorous fish and amphibians from this study (Figure 4). Including
more species could also reveal important but unmeasured environmental covariates via their
pairwise species covariances. For instance, while terrestrial and aquatic species mostly do not
interact directly, some terrestrial species could be proxies for unmeasured land uses that affect
aquatic species, such as increased agricultural runoff. In our models, we included agricultural
and improved-grassland land cover types as environmental covariates, so we saw this effect
directly (Figure 3). The negative covariances of foxes and pheasants with multiple fish species
(Figure 4D) might therefore be revealing other unmeasured land-use covariates (N.B. >35
million pheasants are released annually in the UK, mostly in England, and appear to boost fox
numbers and incentivise land-cover management measures (Sage *et al.* 2020)).

More technically, eDNA sampling makes it more feasible to collect multiple sample replicates,
which would allow combining a JSDM with a detection model to account for observation error
(Diana *et al.* 2022; Doser et al. 2023; Guillera-Arroita *et al.* 2017; Hartig *et al.* 2024; Tobler *et*
al. 2019). Also, only two species in our dataset showed strong signals of dispersal limitation
(Figure 2), but this low number could be because near-neighbour ponds were not sampled in
our dataset, removing the possibility of detecting fine-scale spatial autocorrelation and thereby
possibly reducing the relative importance of dispersal that would support source-sink relations
among closely adjacent ponds. Denser sampling might have detected more evidence of
dispersal limitation. In our case, unfortunately, our dataset used the great crested newt sampling
protocol, which requires only one sample per pond, and the ponds were dispersed across the
landscape because the original use was to fit a species distribution model.

What can JSDMs and internal structure analysis bring to eDNA researchers?

Novel community datasets, including eDNA metabarcoding, are multivariate response datasets;

that is, each species is a response variable, and there are many of them. Before computing

power was widely available, such datasets were generally first reduced to tractable dissimilarity

matrices before visualisation and analysis (e.g. NMDS and constrained ordination), but this

approach loses information and generates artefacts (Warton *et al.* 2012). However, for over a

decade now, it has been possible to analyse multivariate abundance data directly (Warton 2022).

For instance, we have shown here that JSDMs allow the simultaneous analysis of

environmental covariates, biotic covariances, and spatial autocorrelations, and the outputs can

be visualised and interrogated in powerful ways (Hartig *et al.* 2024; Popovic *et al.* 2019; Terry

et al. 2023; van der Veen *et al.* 2022; Warton 2022).

Conclusion

In conclusion, our study demonstrates that the combination of eDNA data and the analytical

approach of exposing the 'internal structure' of metacommunities using JSDMs is a powerful

tool for examining community assembly processes in structured landscapes. For the pond

metacommunity in this study, we could reveal interesting patterns of how the importance of

community assembly processes across sites and species, and relate those differences to spatial,

environmental and trait predictors.

Overall, these results fall in line with other empirical studies (Kadoya *et al.* 2024; Mehner *et*

al. 2021; Vass *et al.* 2022) that support Leibold et al.'s (2022) theoretical argument that JSDMs

can reveal local (site and species-specific) assembly processes from static observational

datasets, thereby making it possible to perform confirmatory tests of established ecological

hypotheses or create new hypotheses from observational data. For instance, the *post-hoc* result

that environmental site R^2 increases with the pond area environmental covariate (Figure S4),

suggests that environmental filtering is more important in larger ponds. This can in principle

be tested by creating ponds of different sizes in the same general area and monitoring species

colonisation.

Our study thus provides a blueprint for ecologists who want to study metacommunity assembly

processes and also for molecular ecologists that want to extract more information from their

eDNA data.

References

Abrego, N., Roslin, T., Huotari, T., Ji, Y., Schmidt, N.M., Wang, J., *et al.* (2021). Accounting for species interactions is necessary for predicting how arctic arthropod communities respond to climate change. *Ecography*, 44, 885–896.

ARG-UK. (2010). *ARG UK Advice Note 5: Great Crested Newt Habitat Suitability Index*. Amphibian and Reptile Groups of the United Kingdom.

Axtner, J., Crampton-Platt, A., Hörig, L.A., Mohamed, A., Xu, C.C.Y., Yu, D.W., *et al.* (2019). An efficient and robust laboratory workflow and tetrapod database for larger scale environmental DNA studies. *GigaScience*, 8, giz029.

Beebee, T.J.C. & Griffiths, R.A. (2000). *Amphibians and reptiles: a natural history of the British Herpetofauna*. The new naturalist. 1. publ. HarperCollins, London.

Biggs, J., Ewald, N., Valentini, A., Gaboriaud, C., Dejean, T., Griffiths, R.A., *et al.* (2015). Using eDNA to develop a national citizen science-based monitoring programme for the great crested newt (Triturus cristatus). *Biological Conservation*, 183, 19–28.

Blanchet, F.G., Cazelles, K. & Gravel, D. (2020). Co-occurrence is not evidence of ecological interactions. *Ecology Letters*, 23, 1050–1063.

Cai, W., Yang, C., Wang, X., Wu, C., Larrieu, L., Lopez-Vaamonde, C., *et al.* (2021). The ecological impact of pest-induced tree dieback on insect biodiversity in Yunnan pine plantations, China. *Forest Ecology and Management*, 491, 119173.

Cottenie, K. (2005). Integrating environmental and spatial processes in ecological community dynamics. *Ecology Letters*, 8, 1175–1182.

DEFRA & Rural Payments Agency. (2022). *Guidance. The SFI improved grassland soils standard*. Available at: https://www.gov.uk/guidance/the-sfi-improved-grassland-soils-standard. Last accessed 19 March 2023.

Diana, A., Matechou, E., Griffin, J., Yu, D., Luo, M., Tosa, M., *et al.* (2022). eDNAPlus: A unifying modelling framework for DNA-based biodiversity monitoring. arXiv.

Dormann, C.F., Bobrowski, M., Dehling, D.M., Harris, D.J., Hartig, F., Lischke, H., *et al.* (2018). Biotic interactions in species distribution modelling: 10 questions to guide interpretation and avoid false conclusions. *Global Ecology and Biogeography*, 27, 1004–1016.

Doser, J. W., Finley, A. O. and Banerjee, S. 2023. Joint species distribution models with imperfect detection for high‐dimensional spatial data. - *Ecology* 104: e4137.

Fasiolo, M., Wood, S.N., Zaffran, M., Nedellec, R. & Goude, Y. (2021). qgam : Bayesian Nonparametric Quantile Regression Modeling in *R. Journal of Statistical Software*, 100.

Guillera-Arroita, G., Lahoz-Monfort, J.J., Rooyen, A.R., Weeks, A.R. & Tingley, R. (2017). Dealing with false-positive and false-negative errors about species occurrence at multiple levels. *Methods in Ecology and Evolution*, 8, 1081–1091.

Gunopulos, D., Hofmann, T., Malerba, D. & Vazirgiannis, M. (Eds.). (2011). *Machine Learning and Knowledge Discovery in Databases: European Conference, ECML PKDD 2011, Athens, Greece, September 5-9, 2011, Proceedings, Part III*. Lecture Notes in Computer Science. Springer Berlin Heidelberg, Berlin, Heidelberg.

Guzman, L.M., Thompson, P.L., Viana, D.S., Vanschoenwinkel, B., Horváth, Z., Ptacnik, R., *et al.* (2022). Accounting for temporal change in multiple biodiversity patterns improves the inference of metacommunity processes. *Ecology*, 103, e3683.

Harper, L.R., Lawson Handley, L., Hahn, C., Boonham, N., Rees, H.C., Gough, K.C., *et al.* (2018). Needle in a haystack? A comparison of eDNA metabarcoding and targeted qPCR for detection of the great crested newt (*Triturus cristatus*). *Ecology and Evolution*, 8, 6330–6341.

Harris, M.S. (2024). Aix galericulata *Mandarin duck. Animal Diversity Web*. Available at:
https://animaldiversity.org/accounts/Aix_galericulata. Last accessed 10 March 2024.
Hartig, F., Abrego, N., Bush, A., Chase, J.M., Guillera-Arroita, G., Leibold, M.A., *et al.* (2024).
Novel community data in ecology – properties and prospects. *Trends in Ecology &*
Evolution, 39, 280–293.
Holyoak, M., Leibold, M.A. & Holt, R.D. (2005). *Metacommunities Spatial Dynamics and*
Ecological Communities. Univ. of Chicago Press, chicago.
Kadoya, T., Nilsson, K.A., Kelly, J., Bartley, T.J., Säterberg, T., Guzzo, M.M., *et al.* (2024).
Common processes drive metacommunity structure in freshwater fish. *Global Ecology*
and Biogeography, e13822.
Lahoz-Monfort, J.J., Guillera-Arroita, G. & Tingley, R. (2016). Statistical approaches to
account for false-positive errors in environmental DNA samples. *Molecular Ecology*
Resources, 16, 673–685.
Lê, S., Josse, J. & Husson, F. (2008). FactoMineR: An R Package for Multivariate Analysis.
Journal of Statistical Software, 25.
Leibold, M.A. & Chase, J.M. (2018). *Metacommunity ecology*. Princeton Univ. Press,
Princeton, NJ.
Leibold, M.A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J.M., Hoopes, M.F., *et al.*
(2004). The metacommunity concept: a framework for multi-scale community ecology:
The metacommunity concept. *Ecology Letters*, 7, 601–613.
Leibold, M.A., Rudolph, F.J., Blanchet, F.G., De Meester, L., Gravel, D., Hartig, F., *et al.*
(2022). The internal structure of metacommunities. *Oikos*, 2022, e08618.
Lopes, T.M., Cunha, E.R., Silva, J.C.B., Behrend, R.D.L. & Gomes, L.C. (2015). Dense
macrophytes influence the horizontal distribution of fish in floodplain lakes.
Environmental Biology of Fishes , 98, 1741–1755.
Maceda-Veiga, A., López, R. & Green, A.J. (2017). Dramatic impact of alien carp Cyprinus
carpio on globally threatened diving ducks and other waterbirds in Mediterranean
shallow lakes. *Biological Conservation*, 212, 74–85.
Mehner, T., Argillier, C., Hesthagen, T., Holmgren, K., Jeppesen, E., Kelly, F., *et al.* (2021).
Model-based decomposition of environmental, spatial and species-interaction effects
on the community structure of common fish species in 772 European lakes. *Global*
Ecology and Biogeogrphy., 30, 1558–1571.
Mulkeen, C.J., Gibson-Brabazon, S., Carlin, C., Williams, C.D., Healy, M.G., Mackey, P., *et*
al. (2017). Habitat suitability assessment of constructed wetlands for the smooth newt
(*Lissotriton vulgaris* [Linnaeus, 1758]): A comparison with natural wetlands.
Ecological Engineering, 106, 532–540.
Oldham, R.S., Keeble, J., Swan, M.J.S. & Jeffcote, M. (2000). Evaluating the suitability of
habitat for the Great Crested Newt (*Triturus cristatus*). *Herpetological Journal*, 10,
143–155.
Ovaskainen, O., Rybicki, J. & Abrego, N. (2019). What can observational data reveal about
metacommunity processes? *Ecography*, 42, 1877–1886.
Pichler, M. & Hartig, F. (2021). A new joint species distribution model for faster and more
accurate inference of species associations from big community data. *Methods in*
Ecology and Evolution, 12, 2159–2173.
Poggiato, G., Münkemüller, T., Bystrova, D., Arbel, J., Clark, J.S. & Thuiller, W. (2021). On
the Interpretations of Joint Modeling in Community Ecology. *Trends in Ecology &*
Evolution, 36, 391–401.
Popovic, G.C., Warton, D.I., Thomson, F.J., Hui, F.K.C. & Moles, A.T. (2019). Untangling
direct species associations from indirect mediator species effects with graphical
models. *Methods in Ecology and Evolution*, 10, 1571–1583.

R Core Team. (2022). R: A language and environment for statistical computing.

Riaz, T., Shehzad, W., Viari, A., Pompanon, F., Taberlet, P. & Coissac, E. (2011). ecoPrimers:
inference of new DNA barcode markers from whole genome sequence analysis. *Nucleic*
Acids Research, 39, e145–e145.

Rowland, C.S., Morton, R.D., Carrasco, L., McShane, G., O'Neil, A.W. & Wood, C.M. (2017).
Land Cover Map 2015 (vector, GB). *NERC Environmental Information Data Centre.*
(Dataset).

Sage, R.B., Hoodless, A.N., Woodburn, M.I.A., Draycott, R.A.H., Madden, J.R. & Sotherton,
N.W. (2020). Summary review and synthesis: effects on habitats and wildlife of the
release and management of pheasants and red-legged partridges on UK lowland shoots.
Wildlife Biology, 2020, wlb.00766.

Shoemaker, L.G. & Melbourne, B.A. (2016). Linking metacommunity paradigms to spatial
coexistence mechanisms. *Ecology*, 97, 2436–2446.

Somervuo, P., Yu, D.W., Xu, C.C.Y., Ji, Y., Hultman, J., Wirta, H., *et al.* (2017). Quantifying
uncertainty of taxonomic placement in DNA barcoding and metabarcoding. *Methods in*
Ecology and Evolution, 8, 398–407.

Stefanoudis, P.V., Sayer, C.D., Greaves, H.M., Davidson, T.A., Robson, H., Almeida, D., *et*
al. (2017). Consequences of fish for cladoceran, water beetle and macrophyte
communities in a farmland pond landscape: implications for conservation. *fal*, 190,
141–156.

Suzuki, Y. & Economo, E.P. (2021). From species sorting to mass effects: spatial network
structure mediates the shift between metacommunity archetypes. *Ecography*, 44, 715–
726.

Szymański, P. & Kajdanowicz, T. (2017). A Network Perspective on Stratification of Multi-
Label Data. Presented at the Proceedings of Machine Learning Research, pp. 22–35.

Terry, J.C.D., Langdon, W. & Rossberg, A.G. (2023). Codistribution as an indicator of whole
metacommunity response to environmental change. *Ecography*, 2023, e06605.

Thompson, P.L., Guzman, L.M., De Meester, L., Horváth, Z., Ptacnik, R., Vanschoenwinkel,
B., *et al.* (2020). A process-based metacommunity framework linking local and
regional scale community ecology. *Ecology Letter*, 23, 1314–1329.

Tobler, M.W., Kéry, M., Hui, F.K.C., Guillera-Arroita, G., Knaus, P. & Sattler, T. (2019). Joint
species distribution models with species correlations and imperfect detection. *Ecology*,
100.

Tréguier, A., Paillisson, J.-M., Dejean, T., Valentini, A., Schlaepfer, M.A. & Roussel, J.-M.
(2014). Environmental DNA surveillance for invertebrate species: advantages and
technical limitations to detect invasive crayfish *Procambarus clarkii* in freshwater
ponds. *Journal of Applied Ecology*, 51, 871–879.

Vass, M., Eriksson, K., Carlsson-Graner, U., Wikner, J. & Andersson, A. (2022). Co-
occurrences enhance our understanding of aquatic fungal metacommunity assembly
and reveal potential host–parasite interactions. *FEMS Microbiology Ecology*, 98,
fiac120.

van der Veen, B., Hui, F.K.C., Hovstad, K.A. & O'Hara, R.B. (2022). Concurrent ordination:
Simultaneous unconstrained and constrained latent variable modelling. *Methods in*
Ecology and Evolution, 14, 683–695.

Vellend, M. (2016). *The theory of ecological communities*. Monographs in population biology.
Princeton University Press, Princeton.

Warton, D.I. (2022). *Eco-Stats: Data Analysis in Ecology: From t-tests to Multivariate*
Abundances. Methods in Statistical Ecology. Springer International Publishing, Cham.

Warton, D.I., Wright, S.T. & Wang, Y. (2012). Distance-based multivariate analyses confound
location and dispersion effects. *Methods in Ecology and Evolution*, 3, 89–101.

Yang, C., Bohmann, K., Wang, X., Cai, W., Wales, N., Ding, Z., *et al.* (2021). Biodiversity
Soup II: A bulk-sample metabarcoding pipeline emphasizing error reduction. *Methods*
in Ecology and Evolution, 12, 1252–1264.
Zepeda-Mendoza, M.L., Bohmann, K., Carmona Baez, A. & Gilbert, M.T.P. (2016). DAMe: a
toolkit for the initial processing of datasets with PCR replicates of double-tagged
amplicons for DNA metabarcoding analyses. *BMC Research Notes*, 9, 255.
Zou, H. & Hastie, T. (2005). Regularization and Variable Selection Via the Elastic Net. *Journal*
of the Royal Statistical Society Series B: Statistical Methodology, 67, 301–320.
Zurell, D., Pollock, L.J. & Thuiller, W. (2018). Do joint species distribution models reliably
detect interspecific interactions from co-occurrence data in homogenous environments?
Ecography, 41, 1812–1819.

Figures and legends

616   **Figure 1. External- versus internal-structure analysis of metacommunities.**

B2. Internal-structure analysis of metacommunities

Leibold et al. (2022) applied a joint species distribution model (JSDM) to synthetic metacommunities and partitioned total variation. JSDMs are multivariate regression models that fit environmental-covariate and (optionally) spatial-autocorrelation terms to individual species distributions and then estimate pairwise species covariances from the residuals. For each species and site, variation in species distributions and compositional variation, respectively, is partitioned into that which can be attributed to environmental covariates (E), spatial autocorrelation (S), and biotic covariances (C, the combined effects of unmeasured environmental covariates + biotic interactions). Unexplained variation measures the contribution of ecological stochasticity.

Biotic covariances

Species distributions that are relatively better explained by negative or positive co-occurrences with other species. These co-occurrences can be caused by correlated responses to the same (but unmeasured) environmental covariates or to dispersal limitation, to mediator species (such as shared predators), or to direct biotic interactions (such as predation).

Environmental covariates

Species distributions that are relatively better explained by environmental covariates, which implies distributions that are determined more by environmental filtering.

Spatial autocorrelation

Species distributions that are relatively better explained by spatial covariates, which implies dispersal limitation (at the spatial scale and resolution being analysed).

C. Relating distribution patterns to assembly processes, by site and species

By species and site, the absolute R^2 values partitioned to environmental covariates (E), spatial autocorrelation (S), and biotic covariances (C) are regressed against species traits, environmental distinctiveness, geographical distinctiveness, and individual environmental covariates. This approach allows one to relate environmental differences between sites, and trait differences between species, to differences in the variation explained by each component, thereby generating testable hypotheses that link distribution patterns to metacommunity assembly processes.

Figure 2. The internal structure of a pond metacommunity. The explained variation in species distributions or site compositions is decomposed,

attributed to the three model components, environmental covariates (E), spatial autocorrelation (S), and biotic covariances (also known as co-

distribution) (C), and visualised in a ternary plot after dividing each component's explained variance by its sum to allow comparison among species.

Top row. Aquatic species only. Bottom row. Aquatic + Terrestrial species. Left column. Each point is a species, point size scales to total $R^2_{McFadden}$

of each species, and the colours code for species trait group. Right column. Each point is a site (pond), and point size scales to total $R^2_{McFadden}$ of

each site.

Figure 3. Estimated environmental effects. Eight environmental covariates were included in the model. The first five covariates from the left are taken from the ten standard pond variables used by surveyors to calculate the Habitat Suitability Index (HSI) of each pond for the great crested newt and are therefore measured at all ponds in our dataset (ARG-UK 2010). The last three covariates describe the dominant land cover class within 500 m of each pond (more details in Table S1). Horizontal bars show the magnitudes, directions, and standard errors of the coefficients of each of the eight environmental covariates for each species. All covariates were normalised before fitting. Significance values are not corrected for multiple comparisons. Colours indicate species trait groups.

Asterisks (*) indicate environmental covariates that have >95% posterior probability of an effect
Significance codes: P < 0.001: ***; P < 0.01: **; P < 0.05: *; P < 0.1: .

634 **Figure 4. Biotic covariances.** A. Aquatic species only, all pairwise covariances. B. Covariances filtered to the 2.5% most negative and positive. C.

635 Aquatic + terrestrial species, all pairwise biotic covariances. D. Covariances filtered to the 2.5% most negative and positive.

Figure 5. Correlation of the importance of assembly processes per site and species to environmental predictors and traits. A-C: Quantile regression, correlating the importance of the three assembly mechanisms (measured by the share of absolute partial R^2 values, each in a different colour) per site against A. environmental distinctiveness. D. association of R^2 shares per species with traits, in this case, species groups. Solid lines indicate significant regressions at P<0.05 significance level (p = 0.006 for the environmental component line), B. geographical distinctiveness (p < 0.001 for the spatial line), and C. pond area (p < 0.001 for the environmental line). Dashed lines are non-significant regressions. Note that the shared fractions were removed from the partial R^2, so the three components will not necessarily sum up to the total R^2 value, which is displayed in Fig. 2.

Figure 6. Predictive versus explanatory performance in two JSDMs. A. Aquatic-species-only model. B. Aquatic + Terrestrial species model. Model performance was assessed using the AUC metric, and the dotted lines indicate mean AUC values (vertical for predictive and horizontal for explanatory). In both models, predictive performance is generally higher for fish than for amphibians, and explanatory performance is generally somewhat greater than the predictive performance, indicating moderate overfitting.