Title

Hybracter: Enabling Scalable, Automated, Complete and Accurate Bacterial Genome

Assemblies

Authors

George Bouras[1,2]*, Ghais Houtak[1,2], Ryan R. Wick[3], Vijini Mallawaarachchi[4], Michael J.

Roach[4,5], Bhavya Papudeshi[4], Lousie M. Judd[3], Anna E. Sheppard[6], Robert A. Edwards[4],

Sarah Vreugde[1,2]

Author institutional affiliations

[1] Adelaide Medical School, Faculty of Health and Medical Sciences, The University of

Adelaide, Adelaide, Australia.

[2] The Department of Surgery - Otolaryngology Head and Neck Surgery, University of

Adelaide and the Basil Hetzel Institute for Translational Health Research, Central Adelaide

Local Health Network, South Australia, Australia.

[3] Department of Microbiology and Immunology, University of Melbourne at the Peter

Doherty Institute for Infection and Immunity, Melbourne, Australia.

[4] Flinders Accelerator for Microbiome Exploration, College of Science and Engineering,

Flinders University, Adelaide, Australia.

[5] Adelaide Centre for Epigenetics and South Australian Immunogenomics Cancer Institute,

The University of Adelaide, Adelaide, Australia.

[6] School of Biological Sciences, The University of Adelaide, Adelaide, Australia.

* Correspondence: george.bouras@adelaide.edu.au;

Abstract

Improvements in the accuracy and availability of long-read sequencing mean that complete

bacterial genomes are now routinely reconstructed using hybrid (i.e. short- and long-reads)

assembly approaches. Complete genomes allow a deeper understanding of bacterial evolution

and genomic variation beyond small nucleotide variants (SNVs). They are also crucial for

identifying plasmids, which often carry medically significant antimicrobial resistance (AMR)

genes. However, small plasmids are often missed or misassembled by long-read assembly

algorithms. Here, we present Hybracter, method for fast, automatic and scalable recovery of

near-perfect complete bacterial genomes using a long-read first assembly approach. We

compared Hybracter to existing automated hybrid assembly tools using a diverse panel of

samples with manually curated ground truth reference genomes. We demonstrate that

Hybracter is more accurate and faster than the existing gold standard automated hybrid

assembler Unicycler. We also show that Hybracter with long-reads only is comparable to

hybrid methods in recovering small plasmids.

Introduction

Reconstructing complete bacterial genomes using *de novo* assembly methods had been

considered too costly and time-consuming to be widely recommended in most cases, even as

recently as 2015 [1]. This was due to the reliance on short-read sequencing technologies, which

does not allow for reconstructing regions with repeats and extremely high GC content [2].

However, since then, advances in long-read sequencing technologies have allowed for the

automatic construction of complete genomes using hybrid assembly approaches. Originally,

this involved starting with a short-read assembly followed by scaffolding the repetitive and

difficult to resolve regions with long-reads [3,4]. This approach was implemented in the

command-line tool Unicycler, which remains the most popular tool for generating complete

bacterial genome assemblies [5]. As long-read sequencing has improved in accuracy and

availability, with the latest Oxford Nanopore Technologies reads recently reaching Q20

(99%+) median accuracy, a long-read first assembly approach supplemented by short-read

polishing has recently been favoured for recovering accurate complete genomes. Long-read-

first approaches provide greater accuracy and contiguity than short-read-first approaches in

difficult regions [6–11]. The current gold standard tool Trycycler even allows for the potential

recovery of perfect genome assemblies [7]. However, Trycycler requires significant microbial

bioinformatics expertise and involves manual decision making, creating a significant barrier

to useability, scalability and automation [12].

Several tools exist that generate automated long-read first genome assemblies, such as

MicroPIPE [13], ASA3P [14], Bactopia [15] and Dragonflye [16]. However, these tools do not consider

factors such as genome reorientation [17] and recent polishing best-practices [18], and often

contain the assembly workflow as a sub-module within a more expansive end-to-end

pipeline. Additionally, none of the existing tools consider the targeted recovery of plasmids.

As long-read assemblers struggle particularly with small plasmids, this leads to incorrectly

recovered or missing plasmids in bacterial assemblies [19].

We introduce Hybracter, a new command-line tool for automated near-perfect long-read-first

complete bacterial genome assembly. It implements a comprehensive and flexible workflow

allowing for long-read assembly polished with long and short-reads ('hybracter hybrid' and

'hybracter hybrid-single') or long-read only assembly polished with long-reads ('hybracter

long' and 'hybracter long-single'). For ease of use and familiarity, Hybracter has been

designed with a command-line interface containing parameters similar to Unicycler.

Additionally, thanks to its Snakemake [20] and Snaketool [21] implementation, Hybracter

seamlessly scales from a single isolate to hundreds or thousands of genomes with high

computational efficiency and supports deployment on HPC clusters and cloud-based

environments.

79   Results

Assembly Workflow

Hybracter implements a long-read-first automated assembly workflow based on current best

practices [12]. The main subcommands available in Hybracter can be found in Table 1 and the

workflow is outlined in Figure 1. Hybracter begins with long-reads for all subcommands, and

with short-reads for polishing for 'Hybracter hybrid' and 'Hybracter hybrid-single'

subcommands.

First, long-read input FASTQs are optionally filtered for quality control with Filtlong [22] and

Porechop_ABI [23], with optional contaminant removal against a host genome using modules

from Trimnami (e.g. if the bacteria has been isolated from a host) [24]. Quality control of short-

read input FASTQs is performed with fastp [25] (Fig 1A). Long-reads are then assembled with

Flye [26]. If at least 1 contig is recovered above the cut-off '-c' chromosome length specified by

the user for the sample, that sample will be denoted as 'complete'. All such contigs will then

be marked as chromosomes and kept for downstream reorientation and polishing. If zero

contigs are above the cut-off chromosome length, the assembly will be denoted as

'incomplete', and all contigs will be kept for downstream polishing (Fig 1B).

For all complete samples, targeted plasmid assembly is then conducted using Plassembler [27]

(Fig 1C). All assemblies are then polished with Medaka, which can be turned off using '--

no_medaka' [28] (Fig 1D). For complete assemblies, the chromosome will be reoriented to

begin with the dnaA chromosomal replication initiator gene using Dnaapler [29]. These

chromosomes are then polished for a second time with Medaka to ensure the sequence around

the original chromosome breakpoint is polished. If the user has provided short-reads using

Hybracter hybrid, the assemblies are then polished with Polypolish [18] followed by pypolca [30]

[31] (Fig 1F). If short-reads are available (Hybracter hybrid), the quality of each assembly

round is scored using ALE [32]. If only long-reads are available (Hybracter long), the mean

coding sequence (CDS) length is calculated for each assembly using Pyrodigal, with larger

mean CDS lengths indicating a better quality assembly [33] [34]. The assembly with the highest

mean CDS length is chosen.

Ultimately, the highest-scoring assembly is chosen as the final assembly (Fig 1G). A final

output assembly FASTA file is created, along with per contig and overall summary statistic

TSV files for each sample, and separate chromosome and plasmid FASTA files for samples

denoted as complete (Fig 1H). Once the final assembly has been chosen for all samples, an

overall 'hybracter_summary.tsv' file is generated. All main output files are explained in more

detail in Table 2. While all these main outputs can be found in the 'FINAL_OUTPUT'

subdirectory, all other intermediate output files are available for users who would like extra

information about their assemblies, including all assembly assessments, comparisons of all

changes introduced by polishing, and Flye and Plassembler output summaries. A full list of

these supplementary outputs can be found in Hybracter's Documentation

(https://hybracter.readthedocs.io/en/latest/output/).

Figure 1: Outline of the Hybracter workflow.

Hybracter
Hybrid or Long-Only
Bacterial Assembly
Pipeline

Long-Reads

Short-Reads
(Hybracter Hybrid Only)

A   **Quality Control**
(Optional)
Long-Reads   Short-Reads

Trimnami (Contamination Removal)   fastp
Filtlong
Porechop-ABI

C Plasmid Assembly   **B Long-Reads Assembly**

Plassembler   Flye
                                     Complete   Incomplete
                                     Chromsome(s)   Contigs

Long-Read Polishing   **Chromosome Reorientation**
(Two Rounds)
                                     Start With *DnaA* Gene
D   Medaka   **E**   Dnaapler

                            Short-Read Polishing   **G Quality**
                                                            Assessment
H   Output   **F**   Polypolish
                                      Pypolca   Run on B,D,E,F
Assembly Statistics
Final Chromosome and Plasmid   ALE or Pyrodigal
Assemblies

123   Table 1.   Summary of the 4 Primary Hybracter Commands

Command	Input	Number of Samples	Description	Workflow Elements Included by Default (From Figure 1)
hybracter hybrid	5 column csv sample sheet specified with '--input' containing: • sample name • long-read FASTQ path, • estimated chromosome length • R1 short-read FASTQ path • R2 short-read FASTQ path	1+	Long-read first assembly followed by long then short-read polishing for multiple isolates. Snakemake implementation ensures efficient use of available resources	A, B, C, D, E, F, G, H
hybracter hybrid-single	• sample name (-s) • long-read FASTQ path (-l) • estimated chromosome length (-c) • R1 short-read FASTQ path (-1) • R2 short-read FASTQ path (-2)	1	Long-read first assembly followed by long then short-read polishing for a single isolate. Similar command line interface to Unicycler	A, B, C, D, E, F, G, H
hybracter long	3 column csv sample sheet specified with '--input' containing: • sample name • long-read FASTQ path, • estimated chromosome length	1+	Long-read first assembly followed by long-read polishing for multiple isolates. Snakemake implementation ensures efficient use of available resources	A (no fastp), B, C, D, E, G, H
hybracter long-single	• sample name (-s) • long-read FASTQ path (-l) • estimated chromosome length (-c)	1	Long-read first assembly followed by long-read polishing on a single isolate.	A (no fastp), B, C, D, E, G, H

126 Table 2.   Description of the Primary Hybracter Output Files

Output File	Description
{sample}_final.fasta	Final assembly FASTA file for the sample. Contains all chromosome(s) and plasmids for complete isolates and all contigs for incomplete isolates.
{final}_chromosome.fasta	Final assembly FASTA file for the chromosomes(s) in a complete sample.
{final}_plasmid.fasta	Final assembly FASTA file for the plasmids in a complete sample.
hybracter_summary.tsv	A TSV file combining the {sample}_summary.tsv files for all samples.
{sample}_summary.tsv	A TSV file containing columns denoting for the sample: - Assembly completeness - Total assembly length - Number of contigs assembled - The polishing round deemed to be most accurate and selected as the final assembly - The length of the longest contig - The estimated coverage of the longest contig - The number of circular plasmids recovered by Plassembler
{sample}_per_contig_stats.tsv	A TSV file containing columns denoting for the sample: - Contig name - Contig Type (chromosome or plasmid) (complete samples only) - Contig length - Contig GC% - Contig circularity (complete samples only)

Tool Selection

Tools were selected for inclusion in Hybracter either based on benchmarking from the literature, or they were specifically developed for inclusion in Hybracter. Flye [26] was chosen as the long-read assembler because it is more accurate than other long-read assemblers with comparable runtimes, such as Raven [35], Redbean [36] and Miniasm [37], while being dramatically faster than the comparably accurate Canu [6,38]. Medaka [28] was chosen as the long-read polisher because of its ability to improve assembly continuity in addition to accuracy [12,39]. The benchmarking results of this study also emphasise that it is particularly good at fixing insertion and deletion (InDel) errors, which cause problematic frameshifts and frequently lead to fractured or truncated gene predictions. Polypolish [18] and POLCA [31] were selected as short-read polishers, as these have been shown to achieve the highest performance with the lowest chance of introducing errors when used in combination [18].

We developed three standalone programs included in Hybracter. These are Dnaapler, Plassembler and Pypolca. Dnaapler was developed to ensure the chromosome(s) identified by Hybracter is reoriented to consistently begin with the dnaA chromosomal replication initiator gene. Full implementation details can be found in the manuscript, with expanded functionality beyond this use case [29]. Plassembler was developed to improve the runtime and accuracy when assembling plasmids in bacterial isolates. Full implementation details can be found in the manuscript for hybrid mode [27]. Hybracter long utilises Plassembler containing a post-publication improvement for long-reads only ('Plassembler long') released in v1.3. Plassembler long assembles plasmids from only long-reads by treating long-reads as both short-reads and long-reads. Plassembler long does this by utilising Unicycler in its pipeline to

create a de Bruijn graph-based assembly, treating the long-reads as unpaired single-end reads,

which are then scaffolded with the same long-read set.

The third tool is Pypolca. Pypolca is a Python re-implementation of the POLCA short-read

genome polisher, created specifically for inclusion in Hybracter and with an almost identical

output format and performance (see Methods). Compared to POLCA, Pypolca features

improved useability with a simplified command line interface and allows the user to specify

an output directory. Furthermore, Pypolca is available on both MacOS and Linux (POLCA is

only available on Linux) and does not require the installation of the entire MaSuRCA genome

assembler toolkit [40]. Pypolca is open-source and freely available on Bioconda, PyPI, and

GitHub (https://github.com/gbouras13/pypolca).

Benchmarking

To test the performance of Hybracter, we used 20 samples with available short- and long-read

sets. These samples represent genomes from a variety of Gram-negative and Gram-positive

bacteria, with most containing plasmids. We chose these samples as they have both real

hybrid read sets and manually curated genome assemblies produced using either Trycycler [7]

or Bact-builder [41]—a consensus-building pipeline based on Trycycler. We tested Hybracter

with both short- and long-reads ('Hybracter hybrid') and long-reads only ('Hybracter long')

against Unicycler and the Dragonflye [16] pipeline both with long-reads only ('Dragonflye

long') and with short-read polishing ('Dragonflye hybrid'). More benchmarking details can

be found in the Methods section.

Chromosome Accuracy Performance

All tools recovered complete circular contigs for each chromosome. SNVs, small InDels

(under 60 bps), and large InDels (over 60 bps) were compared as a measure of assembly

accuracy. To account for differences in genomic size between isolates, SNVs and small InDel

counts were normalised by genome length.

The summary results are presented in Table 3 and visualised in Figure 2. The detailed results

for each tool and sample are presented in Supplementary Table 5. Of the hybrid tools,

Dragonflye hybrid produced the fewest SNVs per 100kbp (median 0.03) followed by

Hybracter hybrid (median 0.16) and Unicycler (median 1.25). Hybracter hybrid produced the

fewest InDels per 100kbp (median 0.05), followed by Unicycler (median 0.28) and

Dragonflye hybrid (median 0.49). Hybracter hybrid also produced the fewest InDels plus

SNVs per 100kbp (median 0.24), followed by Dragonflye hybrid (median 0.74) and

Unicycler (median 1.49). The median InDels plus SNVs per 100kbp rate for Hybracter is

very low, with 0.24 small InDels plus SNVs per 100kbp corresponding to approximately 12

small InDels plus SNVs total for a standard 5MB *E. coli* genome.

Additionally, Hybracter hybrid showed superior performance in terms of large InDels, with a

median of 0 and a total of 59 large InDels across the 20 samples, compared to 1.5 and 91 for

Dragonflye hybrid, and 2.5 and 134 for Unicycler.

Overall, Hybracter hybrid produced the most accurate chromosome assemblies. For eight

isolates described in *Lerminiaux* et al. [9], Hybracter also assembled a perfect chromosome

(Isolates A, B, C, D, E, G, I, L), and another two near-perfect chromosomes (defined as <10

total SNVs plus InDels) (Isolate K and H37R2). Dragonflye hybrid did not assemble any

perfect chromosomes and recovered six near-perfect chromosomes (Isolate B, D, E, G, H, I).

Figure 2: Comparison of the counts of small nucleotide variants (SNVs) and small (<60bp)

insertions and deletions (InDels) per 100kbp (A) and the total number of large (>60bp)

InDels (B) for the Hybrid tools benchmarked (Hybracter hybrid in blue, Dragonflye hybrid,

in orange and Unicycler in green). The counts of SNVs and small InDels per 100kbp (C) and

the total number of large InDels (D) for the long tools benchmarked (Hybracter long in

blue, Dragonflye long in orange) are also shown. All data presented is from the

benchmarking output run with 8 threads.

211
212

213   Table 3. Small (<60bp) InDels, SNVs per 100kbp of sequence and total large (>60bp)

214   InDels of Chromosomes Assemblies for all Benchmarked Isolates.

Tool	Type	Small InDels per 100kbp	SNVs per 100kbp	Small InDels + SNVs per 100kbp	Large InDels
Hybracter hybrid	Hybrid	Median = 0.05 Minimum = 0 Maximum = 12.34	Median = 0.16 Minimum = 0 Maximum = 3.52	Median = 0.24 Minimum = 0 Maximum = 15.79	Total = 59 Median = 0 Minimum = 0 Maximum = 26
Dragonflye hybrid	Hybrid	Median = 0.49 Minimum = 0 Maximum = 11.56	Median = 0.03 Minimum = 0 Maximum = 2.21	Median = 0.74 Minimum = 0 Maximum = 13.41	Total = 91 Median = 1.5 Minimum = 1 Maximum = 29
Unicycler	Hybrid	Median = 0.28 Minimum = 0 Maximum = 9.5	Median = 1.25 Minimum = 0.25 Maximum = 4.13	Median = 1.49 Minimum = 0.43 Maximum = 13.62	Total = 134 Median = 2.5 Minimum = 0 Maximum = 41
Hybracter long	Long	Median = 0.49 Minimum = 0.06 Maximum = 24.82	Median = 1.07 Minimum = 0.07 Maximum = 10.46	Median = 2.08 Minimum = 0.37 Maximum = 35.29	Total = 66 Median = 1 Minimum = 0 Maximum = 27
Dragonflye long	Long	Median = 3.01 Minimum = 1.61 Maximum = 43.8	Median = 0.99 Minimum = 0.33 Maximum = 10.86	Median = 3.81 Minimum = 2.01 Maximum = 53.1	Total = 92 Median = 2 Minimum = 1 Maximum = 30

Similar results were found in the long-read only tool comparison. Dragonflye long produced

slightly fewer SNVs per 100kbp (median = 0.99) than Hybracter long (median 1.07).

However, Hybracter long consistently had fewer small InDels (median 0.49), large InDels

(total 66) and small InDels plus SNVs per 100kbp (median 2.08) than Dragonflye long

(median 3.01, total 92 and median 3.81 respectively). No perfect or near-perfect

chromosomes were assembled by either long-only tool, though Hybracter long did assemble

several chromosomes with fewer than 50 total small InDels plus SNVs (*Lerminiaux* isolates

A, C, D, G, H, L, J, and ATCC BAA-679).

Additionally, long-read only assembly methods had consistently worse performance than

hybrid tools as measured by SNVs and small InDels, suggesting the continuing utility of

short-read polishing for the isolates surveyed.

Plasmid Recovery Performance and Accuracy

Hybracter in both hybrid and long modes was superior at recovering plasmids compared to

the other tools in the same class (Table 4). Hybracter hybrid was able to completely recover

57/59 possible plasmids (the other two were partially recovered), compared to 54/59 for

Unicycler and only 34/59 for Dragonflye hybrid. Hybracter hybrid did not miss a single

plasmid, while Unicycler missed 3/59 (all in Isolate E from *Lerminiaux* et al. [9]) and

Dragonflye hybrid completely missed 9/59. In terms of accuracy, Hybracter hybrid and

Unicycler were similar in terms of SNVs plus small InDels, with medians of 4.15 and 3.83

per 100kbp respectively (Supplementary Table 9), while Hybracter hybrid produced fewer

large InDels than Unicycler (39 vs 51 in total).

Interestingly, Hybracter long showed strong performance at recovering plasmids despite

using only long-reads, completely recovering 54/59 plasmids, completely missing only 2/59.

This performance was far superior to Dragonflye long (35/59 completely recovered, 8/59

missed). In terms of accuracy, both long tools were similar and unsurprisingly less accurate

than the hybrid tools in terms of SNVs plus small InDels (medians of 10.64 per 100kbp for

Hybracter long and 9.22 per 100kbp for Dragonflye long). However, Hybracter long was the

best-performing tool overall as measured by large InDels (total 32), outperforming all hybrid

tools and dragonflye long (total 123). Additionally, all five tools detected an additional

5411bp plasmid in *Lerminiaux* Isolate G not found in the reference sequence and Hybracter

in both hybrid and long modes detected a further 2519bp small plasmid from this genome.

Hybracter hybrid recovers more plasmids than either Unicycler or Dragonflye because it uses

a dedicated plasmid assembler, Plassembler [27]. In addition, Hybracter long, using only long-

reads had an identical complete plasmid recovery rate to Unicycler, which uses both long-

and short-reads (54/59 for both). These results suggest that Hybracter long, by applying

algorithms designed for short-reads on long-reads, largely solves the existing difficulties of

recovering small plasmids from long-reads, at least on the benchmarking dataset of

predominantly R10 Nanopore reads [19,42].

Another interesting result from Hybracter hybrid is that in 6/20 isolates, it assembled

additional non-plasmid contigs, which occurred in only 1/20 isolates for Unicycler. These

contigs are not necessarily an assembly artifact and can provide additional information

regarding the quality control and similarity of short and long-read sets. In Plassembler

implemented within Hybracter hybrid, the existence of such contigs is often indicative of

mismatches between long- and short-read sets [27], suggesting that there is likely some

heterogeneity between long- and short-reads in those six samples.

Table 4. The Total Number of Plasmids Recovered by Each Tool. There were 59 total

reference plasmids in the 20 samples.

Tool	Complete Plasmids Recovered	Total Plasmids Partially Recovered or Misassembled	Total Plasmids Missed	Additional Plasmids Recovered not in Reference	Additional Non-Plasmid Contigs Recovered
Hybracter hybrid	57	2	0	2	6
Unicycler	54	2	3	1	1
Dragonflye hybrid	34	16	9	1	7
Hybracter long	54	3	2	2	3
Dragonflye long	35	16	8	1	5

Runtime Performance Comparison

As shown in Table 5 and Figure 3, median wall-clock times with 8 threads for Dragonflye

hybrid (10m55s) were smaller than Hybracter hybrid (54m23s), which were in turn smaller

than Unicycler (02h03m02s). For the long-only tools, Dragonflye long (9m24s) was faster

than Hybracter long (45m29s). Hybracter long was consistently slightly faster than Hybracter

hybrid (Table 5).

The difference in runtime performance between Hybracter and Dragonflye is predominantly

the result of the included targeted plasmid assembly and the reorientation and assessment

steps in Hybracter that are not included in Dragonflye. Additionally, the results suggest

limited benefits to running Hybacter with more than eight threads. As explained in the

following section, if a user has multiple isolates to assemble, a superior approach is to modify

the configuration file specifying more efficient resource requirements for each job in

Hybracter.

Table 5. Wall-clock Runtime Summary Statistics for Each Tool.

Tool	Type	8 Threads (hh:mm:ss)	16 Threads (hh:mm:ss)
Hybracter hybrid	Hybrid	Median **=** 00:54:23 Minimum = 00:14:22 Maximum = 02:01:37	Median = 00:40:19 Minimum = 00:12:43 Maximum = 01:21:05
Dragonflye hybrid	Hybrid	Median = 00:10:55 Minimum = 00:04:02 Maximum = 00:13:34	Median = 00:07:21 Minimum = 00:03:33 Maximum = 00:09:28
Unicycler	Hybrid	Median = 02:03:02 Minimum = 00:39:09 Maximum = 02:48:16	Median = 01:06:8 Minimum = 00:21:38 Maximum = 01:30m38
Hybracter long	Long	Median **=** 00:45:29 Minimum = 00:10:52 Maximum = 01:23:49	Median **=** 00:34:56 Minimum = 00:09:49 Maximum = 00:59:21
Dragonflye long	Long	Median = 00:09:24 Minimum = 00:03:52 Maximum = 00:13:32	Median **=** 00:07:00 Minimum = 00:03:22 Maximum = 00:08:56

Figure 3: Comparison of wall-clock runtime (in seconds) of Hybracter hybrid, Dragonflye

hybrid, Unicycler, Hybracter long and Dragonflye long when run with 8 and 16 threads.

Parallelisation Allows for Improved Efficiency

Hybracter allows users to specify and customise a configuration file to maximise resource

usage and runtime efficiency. Users can modify the desired threads, memory and time

requirements for each type of job that is run within Hybracter to suit their computational

resources. So that resources are not idle for most users on single sample assemblies, large

jobs such as the Flye and Plassembler assembly steps default to 16 threads and 32 GB of

memory.

To emphasise the efficiency benefits of parallelisation, the 12 *Lerminiaux* et al. isolates were

also assembled using 'hybracter hybrid' with a customised configuration file designed to

improve efficiency on the machine used for benchmarking. Specifically, the configuration

was changed to specify 8 threads and 16 GB of memory allocated to big jobs (assembly,

polishing and assessment) and 4 threads and 8 GB of memory allocated to medium jobs

(reorientation). More details on changing Hybracter's configuration file to suit specific

systems can be found in the documentation

(https://hybracter.readthedocs.io/en/latest/configuration/). We limited the overall 'hybracter

hybrid' run with 32 GB of memory and 16 threads to provide a fair comparison. The overall

'hybracter hybrid' run was then compared to the sum of the 12 'hybracter hybrid-single' runs.

Overall, the 12 isolates took 06h16m53s in the combined run, as opposed to 09h34m08s from

the sum of the 12 'hybracter hybrid-single' and 13h32m51s from the sum of the 12 Unicycler

runs. This inbuilt parallelisation of Hybracter provides significant efficiency benefits if

multiple samples are assembled simultaneously. The performance benefit of Hybracter

afforded by Snakemake integration in parallel computing systems may be variable over

different architectures, but this provides an example case of potential efficiency and

convenience benefits.

Discussion

As long-read sequencing has improved in accuracy with reduced costs, it is now routine to

use a combination of long- and short-reads to generate complete bacterial genomes [3,5] Recent

advances in assembly algorithms and accuracy improvements mean that a long-read first

hybrid assembly should be favoured with short-reads being used after assembly for polishing

[12], as opposed to the short-read first assembly approach (where long-reads are only used for

scaffolding a short-read assembly) utilised by the current automated gold standard Unicycler.

The Unicycler approach is more prone to larger scale InDel errors as well as smaller scale

errors such as those caused by homopolymers or methylation motifs [6,11,43,44]. Additionally, it

should be noted that it is already possible (while perhaps not routine) to generate perfect

hybrid bacterial genome assemblies using manual consensus approaches requiring human

intervention, such as Trycycler [7,45]. While manual approaches such as Trycycler generally

yield superior results to automated approaches, manually assembling many complete

genomes manually is challenging as considerable time, resources and bioinformatics

expertise are required.

The results of this study emphasise that the long-read first hybrid approach consistently

yields superior assemblies than the short-read first hybrid approach and should therefore be

preferred going forward. This study also shows that automated perfect hybrid genome

assemblies are already possible with Hybracter. This study and others [9,46] also confirm that a

long-read first hybrid approach remains preferable to long-read only assembly with Nanopore

reads, as short-reads continue to provide accuracy improvements in polishing steps. However,

it is foreseeable that short-reads will provide little or no accuracy improvements and will not

be needed to polish long-read only assemblies; perfect long-read only assemblies are already

possible, at least with manual intervention using Trycycler [11]. Accordingly, automated perfect

(or near-perfect) bacterial genome assemblies may soon become possible from long-reads

only. Hybracter maintains the flexibility to use long-reads only if desired, allowing users to

turn long-read polishing all-together. This may become increasingly useful as long-read

sequencing continues to improve in accuracy beyond the read sets used in this study, because

long-read polishing can introduce errors and make long-read only assemblies worse with

highly accurate Nanopore and PacBio reads [8,11].

Hybracter was created to bridge the gap from the present to the future of automated perfect

hybrid and long-read-only bacterial genome assemblies. The results of this study show that

Hybracter in hybrid mode is both faster and more accurate than the current gold standard tool

for hybrid assembly Unicycler and is more accurate than Dragonflye in both modes. It should

be noted that if users want fast chromosome only assemblies where accuracy is not essential

(for applications such as species identification or sequence typing), Dragonflye remains a

good option due to its speed.

Hybracter especially excels in recovering complete plasmid genomes compared to other

tools. By incorporating Plassembler, Hybracter recovers more complete plasmid genomes

than Unicycler in hybrid mode. Further, Hybracter long is comparable to Unicycler and

Hybracter hybrid when using long-reads only for plasmid recovery.

The high error rates of long-read sequencing technologies have prevented the application of

assembly approaches designed for highly accurate short-reads, such as constructing de Bruijn

graphs (DBGs) based on strings of a particular length k (k-mers)[47–49]. This resulted in

bioinformaticians initially utilising less efficient algorithms designed with long-reads in

mind, such as utilising overlap graphs in place of DBGs [26,36,38,50,51]. While DBGs have been

used for long-read assembly in some applications [52–54], adoption, especially in microbial

genomics, has been limited.

Although long-read first assembly methods enable complete chromosome and large plasmid

reconstruction, it is well established that long-read only assemblers struggle to assemble

small (<20kbp) plasmids accurately, often leading to missing or multiplicated assemblies[19,27].

These errors may be exacerbated if ligation chemistry based sequencing kits are used[42].

Therefore, hybrid DBG based short-read first assemblies are traditionally recommended for

plasmid recovery [12].

Implemented in our post-publication changes to Plassembler described in this study,

Hybracter solves the problem of small plasmid recovery using long-reads. It achieves this by

implementing a DBG-based assembly approach with Unicycler. The same read set is used

twice, first as unpaired pseudo 'short' reads and then as long-reads; the long-read set

scaffolds a DBG-based assembly based on the same read set. This study demonstrates that

current long-read technologies, such as R10 Nanopore reads, are now accurate enough that

some short-read algorithms are applicable. Our results also suggest that similar DBG-based

algorithmic approaches could be used to enhance the recovery of small replicons in long-read

datasets beyond the use case presented here of plasmids in bacterial isolate assemblies. This

could potentially enhance the recovery of replicons such as bacteriophages [55] or other small

contigs from metagenomes using only long-reads [10,56].

Finally, consistent and resource efficient assemblies that are as accurate as possible in

recovering both plasmids and chromosomes are crucial, particularly for larger studies

investigating plasmid epidemiology and evolution. AMR genes carried on plasmids can have

complicated patterns of transmission involving horizontal transfer between different bacterial

species and lineages, transfer between different plasmid backbones, and integration into and

excision from the bacterial chromosome [57–59] . Accurate plasmid assemblies are crucial in

genomic epidemiology studies investigating transmission of antimicrobial resistant bacteria

within outbreak settings, as well as in a broader One Health context, where hundreds or even

thousands of assemblies may be analysed [60–63]. Hybracter will facilitate the expansion of such

studies, allowing for faster and more accurate automated complete genome assemblies than

existing tools. Additionally, by utilising Snakemake[20] with a Snaketool[21] command line

interface, Hybracter is easily and efficiently parallelised to optimise available resources over

various large-scale computing architectures. Individual jobs (such as each assembly,

reorientation, polishing or assessment step) within Hybracter are automatically sent to

different resources on a high performance computing (HPC) cluster using the HPC's job

scheduling system like Slurm[64]. Hybracter can natively use any Snakemake-supported cloud-

based deployments such as Kubernetes, Google Cloud Life Sciences, Tibanna, and Azure

Batch.

Conclusion

Hybracter is substantially faster than the current gold standard tool Unicycler, assembles

chromosomes more accurately than existing methods, and is superior at recovering complete

plasmid genomes. By applying DBG-based algorithms designed for short-reads on current

generation long-reads, Hybracter long also solves the problem of long-read-only assemblers

entirely missing or duplicating small circular elements such as plasmids. Hybracter is

resource efficient and natively supports deployment on high-performance computer clusters

and cloud environments for massively parallel analyses. We believe Hybracter will prove to

be an extremely useful tool for the automated recovery of complete bacterial genomes from

hybrid and long-read-only sequencing data suitable for massive datasets.

Methods

Benchmarking

To compare Hybracter's functionality and performance, we benchmarked its performance

against other software tools. We focused on the most popular state-of-the-art assembly tools

for automated hybrid and long only bacterial genome assemblies. All code to replicate these

analyses can be found at the repository

(https://github.com/gbouras13/hybracter_benchmarking). All programs and dependency

versions used for benchmarking can be found in Supplementary Table 4. For the hybrid tools,

we chose Unicycler and Dragonflye with both long-read and short-read polishing (denoted

'Dragonflye hybrid'). Dragonflye was chosen as it is a popular long-read first assembly

pipeline [16]. Both tools were run using default parameters. By default, Dragonflye conducts a

long-read assembly with Flye that is polished by Racon [65] followed by Polypolish. For the

long-read only tool, we chose Dragonflye with long-read Racon based polishing only

(denoted 'Dragonflye long').

We used 20 samples for benchmarking, representing genomes from a variety of Gram-

negative and Gram-positive bacteria. We chose these samples as they have real hybrid read

sets in combination with manually curated genome assemblies produced using either

Trycycler or Bact-builder [41]—a consensus-building pipeline based on Trycycler. These

samples came from 4 different studies below. We used the published genomes from these

studies (or the available genomes available from the ATCC) as representatives of the 'ground

truth' for these samples. Where read coverage exceeded 100x samples were subsampled to

approximately 100x coverage of the approximate genome size with Rasusa v0.7.0 [66], as this

better reflects more realistic read depth of real life isolate sequencing. Nanoq v0.10.0 [67] was

used to generate quality control statistics for the subsampled long-read sets. Four isolates did

not have 100x long-read coverage — the entire long-read set was used instead. A full

summary table of the read lengths, quality, Nanopore kit and base-calling models used in

these studies can be found in Supplementary Table 2.

These samples contained varying levels of long-read quality (reflecting improvements in

Oxford Nanopore Technologies long-read technology), with the median Q score of long-read

sets ranging from 12.3 to 18.3. The four studies are:

1. Five ATCC strain isolates (ATCC-10708 *Salmonella enterica*, ATCC-17802 *Vibrio*

paragaemolyticus, ATCC-25922 *Escherichia coli*, ATCC-33560 *Campylobacter*

jejuni and ATCC-BAA-679 *Listeria monocytogenes*) made available as a part of this

study [8]

2.   Twelve diverse carbapenemase-producing Gram-negative bacteria from *Lerminiaux*

et al. [9]

3. *Staphylococcus aureus* JKD6159 sequenced with both R9 and R10 chemistry long-

read sets from *Wick* et al. [45]

4. *Mycobacterium tuberculosis* HR37v from *Chitale* et al. [41]

The full details for each individual isolate used can be found in Supplementary Tables 1 and

2.

Chromosome Accuracy

The assembly accuracy of the chromosomes recovered by each benchmarked tool was

compared using Dnadiff v1.3 packaged with MUMmer v3.23 [68]. Comparisons were

performed on the largest assembled contig (denoted as the chromosome) by each method,

other than for ATCC-17802 *Vibrio parahaemolyticus*, where the two largest contigs were

chosen as it has two chromosomes.

Plasmid Recovery Performance and Accuracy

Plasmid recovery performance for each tool was compared using the following methodology.

Summary statistics are presented considered in Table 4. See Supplementary Table 7 for a full

sample-by-sample analysis. All samples were analysed using the 4-step approach outlined

below using summary length and GC% statistics for all contigs and the output of Dnadiff

v1.3 comparisons generated for each sample and tool combination against the reference

genome plasmids:

1. The number of circularised plasmid contigs recovered for each isolate was compared

to the reference genome. If the tool recovered a circularised contig homologous to that

in the reference, it was denoted as completely recovered. Specifically, a contig was

denoted as completely recovered if it had a genome length within 250bp of the

reference plasmid, a GC% within 0.1% of the reference plasmid and whether the

Total Query Bases covered was within 250bp of the Total Reference Bases from

Dnadiff. For Dragonflye assemblies, some plasmids were duplicated or multiplicated

due to known issues with the long-read first assembly approach for small plasmids

[6,19,42]. Any circularised contigs that were multiplicated compared to the reference

plasmid were therefore denoted as misassembled.

2. For additional circularised contigs not found in the reference recovered, these were

tested for homology with NCBI nt database using the web version of blastn [69]. If there

was a hit to a plasmid, the Plassembler output within Hybracter was checked for

whether the contig had a Mash hit (i.e. a Mash distance of 0.2 or lower) to plasmids in

the PLSDB [70]. If there was a hit, the contig was denoted as an additional recovered

plasmid. There were 2 in total (see Supplementary Table 7 and supplementary data).

3. Plasmids with contigs that were either not circularised but homologous to a reference

plasmid, or circularised but incomplete (failing the genome length and Dnadiff criteria

in 1.) were denoted as partially recovered or misassembled.

4. Reference plasmids without any homologous contigs in the assembly were denoted as

missed.

Additional non-circular contigs that had no homology with reference plasmids and were not

identified as plasmids in step 2 were analysed on a contig-by-contig basis and denoted as

additional non-plasmid contigs (see Supplementary Table 7 for contig-by-contig analysis

details).

Runtime Performance Comparison

To compare the performance of Hybracter, we compared wall-clock runtime consumption on

a machine with an Intel® Core™ i7-10700K CPU @ 3.80 GHz on a machine running Ubuntu

20.04.6 LTS with a total of 16 available threads (8 cores). We ran all tools with 8 and 16

threads and with 32 GB of memory to provide runtime metrics comparable to commonly

available consumer hardware. Hybracter hybrid and long were run with 'hybracter hybrid-

single' and 'hybracter long-single' for each isolate to generate a comparable per sample

runtime for comparison with the other tools. The summary results are available in Table 5

and the detailed results for each specific tool and thread combination are found in

Supplementary Table 8.

Sequencing

DNA extraction was performed with the DNeasy Blood and Tissue kit (Qiagen). Illumina

library preparation was performed using Illumina DNA prep (Illumina Inc.) according to the

manufacturer's instructions. Short-read whole genome sequencing was performed an Illumin

MiSeq with a 250bp PE kit. Oxford Nanopore Technologies library preparation ligation

sequencing library was prepared using the ONT SQK-NBD114-96 kit and the resultant

library was sequenced using an R10.4.1 MinION flow cell (FLO-MIN114) on a MinION

Mk1b device. Data was base-called with Super-Accuracy Basecalling (SUP) using the

basecaller model dna_r10.4.1_e8.2_sup@v3.5.1.

Pypolca Benchmarking

Pypolca v0.2.0 was benchmarked against POLCA (in MaSuRCA v4.1.0) [31] using 18 isolates

described above. These were all 12 *Lerminiaux* et al. isolates, the R10 JKD6159 isolate [45] and

the 5 ATCC samples we sequenced as a part of this study. Benchmarking was conducted on

an Intel® Core™ i7-10700K CPU @ 3.80 GHz on a machine running Ubuntu 20.04.6 LTS.

All short read FASTQs used for benchmarking are identical to those used to benchmark

Hybracter. The assemblies used for polishing were intermediate chromosome assemblies

from Flye v2.9.2 [26] generated within Hybracter. The outputs from Pypolca and POLCA were

compared using Dnadiff v1.3 packaged with MUMmer v3.23 [68] Overall, Pypolca and

POLCA yielded extremely similar results. 16/18 assemblies were identical. ATCC 33560 had

2 Single Nucleotide Polymorphisms (SNPs) between Pypolca and POLCA and *Lerminiaux*

Isolate I also had 2 SNPs.

Data Availability

The subsampled FASTQ files used for benchmarking are publicly available at Zenodo with DOI (https://doi.org/10.5281/zenodo.10158013). All ATCC FASTQ reads sequenced as a part of this study can be found under BioProject PRJNA1042815 with the genomes publicly available from the ATCC. All raw *Lermaniaux* et al. FASTQ read files and genomes (prior to subsampling) can be found in the SRA under BioProject PRJNA1020811. All *Staphylococcus aureus* JKD6159 FASTQ read files and genomes can be found under BioProject PRJNA50759. All *Mycobacterium tuberculosis* H37R2 FASTQ read files and genomes can be found under BioProject PRJNA836783. The complete list of BioSample accession numbers for each benchmarked sample can be found in Supplementary Table 1. The benchmarking assembly output files are publicly available on Zenodo with DOI (https://doi.org/10.5281/zenodo.10158013). All Pypolca benchmarking outputs and code are publicly available on Zenodo with DOI (https://zenodo.org/doi/10.5281/zenodo.10072192).

Code Availability

Hybracter is developed using Python and Snakemake as a command-line software tool for Linux and MacOS systems. Hybracter is freely available under an MIT License on GitHub (https://github.com/gbouras13/hybracter) and the documentation is available at Read the Docs (https://hybracter.readthedocs.io/en/latest/). Hybracter is available to install via PyPI (https://pypi.org/project/hybracter/) and Bioconda (https://anaconda.org/bioconda/hybracter).

573 All code used to benchmark Hybracter, including the reference genomes, is publicly available

574 on GitHub (https://github.com/gbouras13/hybracter_benchmarking) with released DOI

575 (https://zenodo.org/doi/10.5281/zenodo.10157987) available at Zenodo.

Acknowledgements

579 This work was supported with supercomputing resources provided by the Phoenix HPC

580 service at the University of Adelaide. We would particularly like to thank Fabien Voisin for

581 his integral role in maintaining and running Phoenix. We would also like to thank Brad Hart

582 for useful comments in testing Hybracter and Simone Pignotti and Oliver Schwengers for

583 providing helpful comments and GitHub pull requests.

Funding

587 G.H. was supported by The University of Adelaide International Scholarships and a THRF

588 Postgraduate Top-up Scholarship. A.E.S was supported by a University of Adelaide Barbara

589 Kidman Women's Fellowship. R.A.E was supported by an award from the NIH NIDDK

590 RC2DK116713 and an award from the Australian Research Council DP220102915. S.V. was

591 supported by a Passe and Williams Foundation senior fellowship.

References

1. Land, M. *et al.* Insights from 20 years of bacterial genome sequencing. *Funct Integr Genomics* **15**, 141–161 (2015).

2. Goldstein, S., Beka, L., Graf, J. & Klassen, J. L. Evaluation of strategies for the assembly of diverse bacterial genomes using MinION long-read sequencing. *BMC Genomics* **20**, 23 (2019).

3. De Maio, N. *et al.* Comparison of long-read sequencing technologies in the hybrid assembly of complex bacterial genomes. *Microbial Genomics* **5**, e000294 (2019).

4. Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Y. 2017. Completing bacterial genome assemblies with multiplex MinION sequencing. *Microbial Genomics* **3**, e000132.

5. Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. *PLOS Computational Biology* **13**, e1005595 (2017).

6. Wick, R. R. & Holt, K. E. Benchmarking of long-read assemblers for prokaryote whole genome sequencing. Preprint at https://doi.org/10.12688/f1000research.21782.4 (2021).

7. Wick, R. R. *et al.* Trycycler: consensus long-read assemblies for bacterial genomes. *Genome Biology* **22**, 266 (2021).

8. Wick, R. ONT-only accuracy with R10.4.1. *Ryan Wick's bioinformatics blog* https://rrwick.github.io/2023/05/05/ont-only-accuracy-with-r10.4.1.html (2023).

9. Lerminiaux, N., Fakharuddin, K., Mulvey, M. R. & Mataseje, L. Do we still need Illumina sequencing data?: Evaluating Oxford Nanopore Technologies R10.4.1 flow cells and v14 library prep kits for Gram negative bacteria whole genome assemblies.

2023.09.25.559359 Preprint at https://doi.org/10.1101/2023.09.25.559359

(2023).

10. Sereika, M. *et al.* Oxford Nanopore R10.4 long-read sequencing enables the

generation of near-finished bacterial genomes from pure cultures and metagenomes

without short-read or reference polishing. *Nat Methods* **19**, 823–826 (2022).

11. Wick, R. ONT-only accuracy: 5 kHz and Dorado. *Ryan Wick's bioinformatics blog*

https://rrwick.github.io/2023/10/24/ont-only-accuracy-update.html (2023).

12. Wick, R. R., Judd, L. M. & Holt, K. E. Assembling the perfect bacterial genome using

Oxford Nanopore and Illumina sequencing. *PLOS Computational Biology* **19**,

e1010905 (2023).

13. Murigneux, V. *et al.* MicroPIPE: validating an end-to-end workflow for high-quality

complete bacterial genome construction. *BMC Genomics* **22**, 474 (2021).

14. Schwengers, O. *et al.* ASA3P: An automatic and scalable pipeline for the assembly,

annotation and higher-level analysis of closely related bacterial isolates. *PLOS*

Computational Biology **16**, e1007134 (2020).

15. Petit, R. A. & Read, T. D. Bactopia: a Flexible Pipeline for Complete Analysis of

Bacterial Genomes. *mSystems* **5**, 10.1128/msystems.00190-20 (2020).

16. Petit III, R. A. Dragonflye: Assemble bacterial isolate genomes from Nanopore reads.

17. Hunt, M. *et al.* Circlator: automated circularization of genome assemblies using long

sequencing reads. *Genome Biology* **16**, 294 (2015).

18. Wick, R. R. & Holt, K. E. Polypolish: Short-read polishing of long-read bacterial

genome assemblies. *PLOS Computational Biology* **18**, e1009802 (2022).

19. Johnson, J., Soehnlen, M. & Blankenship, H. M. Long read genome assemblers

struggle with small plasmids. *Microbial Genomics* **9**, 001024 (2023).

20. Mölder, F. *et al.* Sustainable data analysis with Snakemake. Preprint at

https://doi.org/10.12688/f1000research.29032.2 (2021).

21. Roach, M. J. *et al.* Ten simple rules and a template for creating workflows-as-

applications. *PLOS Computational Biology* **18**, e1010705 (2022).

22. Wick, R. R. Filtlong. (2018).

23. Bonenfant, Q., Noé, L. & Touzet, H. Porechop_ABI: discovering unknown adapters in

Oxford Nanopore Technology sequencing reads for downstream trimming.

Bioinformatics Advances **3**, vbac085 (2023).

24. Roach, M. J. Trimnami: Trim lots of metagenomics samples all at once. (2023).

25. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics **34**, i884–i890 (2018).

26. Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone reads

using repeat graphs. *Nat Biotechnol* **37**, 540–546 (2019).

27. Bouras, G., Sheppard, A. E., Mallawaarachchi, V. & Vreugde, S. Plassembler: an

automated bacterial plasmid assembly tool. *Bioinformatics* **39**, btad409 (2023).

28. medaka: Sequence correction provided by ONT Research.

29. Bouras, G., Papudeshi, B., Grigson, S., Mallawaarachchi, V. & Roach, M. J. Dnaapler: A

tool to reorient circular microbial genomes. (2023).

30. Bouras, G. & Zimin, A. V. pypolca: Standalone Python reimplementation of the

genome polishing tool POLCA. (2023).

31. Zimin, A. V. & Salzberg, S. L. The genome polishing tool POLCA makes fast and

accurate corrections in genome assemblies. *PLOS Computational Biology* **16**,

e1007981 (2020).

32. Clark, S. C., Egan, R., Frazier, P. I. & Wang, Z. ALE: a generic assembly likelihood

evaluation framework for assessing the accuracy of genome and metagenome

assemblies. *Bioinformatics* **29**, 435–443 (2013).

33. Larralde, M. Pyrodigal: Python bindings and interface to Prodigal, an efficient

method for gene prediction in prokaryotes. *Journal of Open Source Software* **7**, 4296

(2022).

34. Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site

identification. *BMC Bioinformatics* **11**, 119 (2010).

35. Vaser, R. & Šikić, M. Time- and memory-efficient genome assembly with Raven. *Nat*

Comput Sci **1**, 332–336 (2021).

36. Ruan, J. & Li, H. Fast and accurate long-read assembly with wtdbg2. *Nat Methods* **17**,

155–158 (2020).

37. Li, H. Minimap and miniasm: fast mapping and de novo assembly for noisy long

sequences. *Bioinformatics* **32**, 2103–2110 (2016).

38. Koren, S. *et al.* Canu: scalable and accurate long-read assembly via adaptive k-mer

weighting and repeat separation. *Genome Res.* **27**, 722–736 (2017).

39. Zhang, X. *et al.* Benchmarking of long-read sequencing, assemblers and polishers for

yeast genome. *Briefings in Bioinformatics* **23**, bbac146 (2022).

40. Zimin, A. V. *et al.* The MaSuRCA genome assembler. *Bioinformatics* **29**, 2669–2677

(2013).

41. Chitale, P. *et al.* A comprehensive update to the Mycobacterium tuberculosis H37Rv

reference genome. *Nat Commun* **13**, 7068 (2022).

42. Wick, R. R., Judd, L. M., Wyres, K. L. & Holt, K. E. Y. 2021. Recovery of small plasmid

sequences via Oxford Nanopore sequencing. *Microbial Genomics* **7**, 000631 (2021).

43. Wick, R. R., Judd, L. M. & Holt, K. E. Performance of neural network basecalling tools

for Oxford Nanopore sequencing. *Genome Biology* **20**, 129 (2019).

44. Marinus, M. G. & Løbner-Olesen, A. DNA Methylation. *EcoSal Plus* **6**,

10.1128/ecosalplus.ESP-0003–2013 (2014).

45. Wick, R. R., Judd, L. M., Monk, I. R., Seemann, T. & Stinear, T. P. Improved Genome

Sequence of Australian Methicillin-Resistant Staphylococcus aureus Strain JKD6159.

Microbiology Resource Announcements **12**, e01129-22 (2023).

46. Sanderson, N. D. *et al.* Comparison of R9.4.1/Kit10 and R10/Kit12 Oxford Nanopore

flowcells and chemistries in bacterial genome reconstruction. *Microbial Genomics* **9**,

000910 (2023).

47. Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its Applications

to Single-Cell Sequencing. *Journal of Computational Biology* **19**, 455–477 (2012).

48. Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-

node solution for large and complex metagenomics assembly via succinct de Bruijn

graph. *Bioinformatics* **31**, 1674–1676 (2015).

49. Compeau, P. E. C., Pevzner, P. A. & Tesler, G. How to apply de Bruijn graphs to

genome assembly. *Nature Biotechnology* **29**, 987–991 (2011).

50. Wong, J. *et al.* Linear time complexity de novo long read genome assembly with

GoldRush. *Nat Commun* **14**, 2906 (2023).

51. Amarasinghe, S. L. *et al.* Opportunities and challenges in long-read sequencing data

analysis. *Genome Biology* **21**, 30 (2020).

52. Lin, Y. *et al.* Assembly of long error-prone reads using de Bruijn graphs. *Proceedings*

of the National Academy of Sciences **113**, E8396–E8405 (2016).

53. Ekim, B., Berger, B. & Chikhi, R. Minimizer-space de Bruijn graphs: Whole-genome

assembly of long reads in minutes on a personal computer. *Cell Syst* **12**, 958-968.e6

(2021).

54. Bankevich, A., Bzikadze, A. V., Kolmogorov, M., Antipov, D. & Pevzner, P. A. Multiplex

de Bruijn graphs enable genome assembly from long, high-fidelity reads. *Nat*

Biotechnol **40**, 1075–1081 (2022).

55. Mallawaarachchi, V. *et al.* Phables: from fragmented assemblies to high-quality

bacteriophage genomes. *Bioinformatics* **39**, btad586 (2023).

56. Kolmogorov, M. *et al.* metaFlye: scalable long-read metagenome assembly using

repeat graphs. *Nat Methods* **17**, 1103–1110 (2020).

57. Sheppard, A. E. *et al.* Nested Russian Doll-Like Genetic Mobility Drives Rapid

Dissemination of the Carbapenem Resistance Gene blaKPC. *Antimicrobial Agents and*

Chemotherapy **60**, 3767–3778 (2016).

58. Mathers, A. J. *et al.* Klebsiella quasipneumoniae Provides a Window into

Carbapenemase Gene Transfer, Plasmid Rearrangements, and Patient Interactions

with the Hospital Environment. *Antimicrobial Agents and Chemotherapy* **63**,

10.1128/aac.02513-18 (2019).

59. Houtak, G. *et al.* The intra-host evolutionary landscape and pathoadaptation of

persistent Staphylococcus aureus in chronic rhinosinusitis. *Microbial Genomics* **9**,

001128 (2023).

60. Hawkey, J. *et al.* ESBL plasmids in Klebsiella pneumoniae: diversity, transmission

and contribution to infection burden in the hospital setting. *Genome Medicine* **14**, 97

(2022).

61. Roberts, L. W. *et al.* Long-read sequencing reveals genomic diversity and associated

plasmid movement of carbapenemase-producing bacteria in a UK hospital over 6

years. *Microbial Genomics* **9**, 001048 (2023).

62. Matlock, W. *et al.* Enterobacterales plasmid sharing amongst human bloodstream

infections, livestock, wastewater, and waterway niches in Oxfordshire, UK. *eLife* **12**,

e85302 (2023).

63. Lerminiaux, N. *et al.* Plasmid genomic epidemiology of blaKPC carbapenemase-

producing Enterobacterales in Canada, 2010–2021. *Antimicrobial Agents and*

Chemotherapy **0**, e00860-23 (2023).

64. Yoo, A. B., Jette, M. A. & Grondona, M. SLURM: Simple Linux Utility for Resource

Management. in *Job Scheduling Strategies for Parallel Processing* (eds. Feitelson, D.,

Rudolph, L. & Schwiegelshohn, U.) 44–60 (Springer, 2003).

doi:10.1007/10968987_3.

65. Vaser, R., Sović, I., Nagarajan, N. & Šikić, M. Fast and accurate de novo genome

assembly from long uncorrected reads. *Genome Res.* **27**, 737–746 (2017).

66. Hall, M. B. Rasusa: Randomly subsample sequencing reads to a specified coverage.

Journal of Open Source Software **7**, 3941 (2022).

67. Steinig, E. & Coin, L. Nanoq: ultra-fast quality control for nanopore reads. *Journal of*

Open Source Software **7**, 2991 (2022).

68. Kurtz, S. *et al.* Versatile and open software for comparing large genomes. *Genome*

Biology **5**, R12 (2004).

69. Sayers, E. W. *et al.* Database resources of the national center for biotechnology

information. *Nucleic Acids Research* **50**, D20–D26 (2022).

70. Galata, V., Fehlmann, T., Backes, C. & Keller, A. PLSDB: a resource of complete

bacterial plasmids. *Nucleic Acids Research* **47**, D195–D202 (2019).

Hybracter
Hybrid or Long-Only Bacterial Assembly Pipeline

Long-Reads

Short-Reads
(Hybracter Hybrid Only)

A Quality Control (Optional)

Long-Reads

Trimnami (Contamination Removal)
Filtlong
Porechop-ABI

Short-Reads

fastp

C Plasmid Assembly

Plassembler

B Long-Reads Assembly

Flye

Complete Chromsome(s)

Incomplete Contigs

D Long-Read Polishing (Two Rounds)

Medaka

E Chromosome Reorientation

Start With *DnaA* Gene
Dnaapler

F Short-Read Polishing

Polypolish
Pypolca

G Quality Assessment

Run on B,D,E,F

ALE or Pyrodigal

H Output

Assembly Statistics
Final Chromosome and Plasmid Assemblies