Targeting plasmid-encoded proteins that contain immunoglobulin-like domains to combat

antimicrobial resistance.

Alejandro Prieto[1*], Lluïsa Miró[2,3*], Yago Margolles[4], Manuel Bernabeu[1], David Salguero[1], Susana

Merino[1], Joan Tomás[1], Juan Alberto Corbera[5], Anna Pérez-Bosque[2,3], Mário Hüttener[1], Luis Ángel

Fernández[4] and Antonio Juárez[1,6**]

1. Department of Genetics, Microbiology and Statistics, University of Barcelona, Barcelona,
Spain.

2. Department of Biochemistry and Physiology, Universitat de Barcelona, Barcelona, Spain.

3. Institut de Nutrició i Seguretat Alimentària, Universitat de Barcelona, Barcelona, Spain.

4. Department of Microbial Biotechnology, Centro Nacional de Biotecnología, Consejo Superior
de Investigaciones Científicas (CNB-CSIC), Madrid, Spain.

5. Instituto Universitario de Investigaciones Biomédicas y Sanitarias (IUIBS), Facultad de
Veterinaria, Universidad de Las Palmas de Gran Canaria (ULPGC), Campus Universitario de
Arucas, Las Palmas, Spain.

6. Institute for Bioengineering of Catalonia, The Barcelona Institute of Science and Technology,
Barcelona, Spain.

*Both authors contributed equally to this work.

**Corresponding author: Prof. Antonio Juárez, ajuarez@ub.edu

Running Title: Plasmid-encoded Big proteins targeted to combat antimicrobial resistance

Keywords: Antimicrobial resistance; IncHI plasmids; R27; Big proteins; RSP protein; vaccination;
nanobodies; plasmid conjugation.

Abstract

Antimicrobial resistance (AMR) poses a significant threat to human health. Although vaccines have been developed to combat AMR, it has proven challenging to associate specific vaccine antigens with AMR. Bacterial plasmids play a crucial role in the transmission of AMR. Our recent research has identified a group of bacterial plasmids (specifically, IncHI plasmids) that encode large molecular mass proteins containing bacterial immunoglobulin-like domains. These proteins are found on the external surface of the bacterial cells, such as in the flagella or conjugative pili. In this study, we show that these proteins are antigenic and can protect mice from infection caused by an AMR *Salmonella* strain harboring one of these plasmids. Furthermore, we successfully generated nanobodies targeting these proteins, that were shown to interfere with the conjugative transfer of IncHI plasmids. Considering that these proteins are also encoded in other groups of plasmids, such as IncA/C and IncP2, targeting them could be a valuable strategy in combating AMR infections caused by bacteria harboring different groups of AMR plasmids. Since the selected antigens are directly linked to AMR itself, the protective effect extends beyond specific microorganisms to include all those carrying the corresponding resistance plasmids.

Introduction

Despite the availability of antibiotics, bacterial infectious diseases are the second leading cause of death worldwide (Morens et al., 2004; Murray et al., 2022). The gradual increase in the resistance rates of several important bacterial pathogens represents a serious threat to public health (Meyer et al., 2010; Rossolini et al., 2007; Spellberg et al., 2008). In 2019, up to 4,95 million deaths could be associated to antimicrobial resistance (AMR) (Murray et al., 2022).

Plasmids can confer resistance to the major classes of antimicrobials (Carattoli, 2009). Their transmission by horizontal gene transfer (HGT) is largely underlying the dissemination of AMR genes, especially in Gram-negative bacteria (Carattoli, 2013; Thomas and Nielsen, 2005; Wang et al., 2015), but also in Gram-positive bacteria (Vrancianu et al., 2020). Among the various methods utilized for classifying plasmids, grouping them into incompatibility groups (Inc) is a well-established and extensively employed approach (Novick and Richmond, 1965).

The incompatibility group HI includes plasmids that are widespread in the *Enterobacteriaceae*. These plasmids frequently include genetic elements that encode multiple AMR determinants (Phan and Wain, 2008). IncHI-encoded AMR can be present in enterobacteria such as *Salmonella* (Phan and Wain, 2008), *Escherichia coli* (Forde et al., 2018), *Klebsiella pneumoniae* (Villa et al., 2012), and *Citrobacter freundii* (Dolejska et al., 2013). Plasmids of the IncHI2 subgroup predominate in antibiotic-resistant *Salmonella* isolates. In *S. enterica* serotype Typhi, more than 40% of isolates harbor an IncHI plasmid (Holt et al., 2011). Recent reports have shown a novel role of IncHI plasmids in AMR spread. Colistin is a last resort antibiotic for the treatment of severe infections (Lim et al., 2010). Plasmid-mediated resistance, conferred by the mobilized colistin resistance gene (*mcr-1*), has emerged recently. Approximately 20% of global isolates carrying the *mcr-1* gene are associated with an IncHI2 plasmid, and this percentage escalates to 41% when considering exclusively European isolates (Matamoros et al., 2017). Of special concern is the presence of the *mcr-1* resistance determinant in enterobacterial isolates carrying

carbapenem resistance genes, such as *blaNDM* and *blaKPC*. The combination of these AMR

determinants can seriously compromise the treatment of infections caused by virulent strains

harboring these plasmids (Wang et al., 2017; Zheng et al., 2016). An example of this is the recent

report of an AMR clone of the highly virulent *E. coli* ST95 lineage (Forde et al., 2018). *E. coli* ST95

clones cause neonatal meningitis and sepsis. They are usually sensitive to several antibiotics.

This AMR clone harbors an IncHI2 plasmid that carries, among other factors, genes encoding

determinants of resistance to colistin and multiple other antibiotics (including the extended-

spectrum beta-lactamase *bla*$_{CTX-M-1}$ gene cluster). The spread of such an AMR ST95 clone could

pose a threat to human health worldwide (Forde et al., 2018).

The R27 plasmid is the best studied IncHI1 plasmid. It harbors the Tn10 transposon, which

confers resistance to tetracycline. The R27 replicative and conjugative machineries have been

studied in detail (Lawley et al., 2003, 2002) and its complete nucleotide sequence is known

(Sherburne et al., 2000). A newly reported feature of IncHI plasmids is that they code for large

molecular mass proteins that contain bacterial Ig-like (Big) domains (Hüttener et al., 2022, 2019).

Bacterial proteins containing Big domains play different roles in the cell. Some fimbrial subunits,

adhesins, membrane transporters and enzymes contain Big domains (as reviewed in (Bodelón

et al., 2013)). The IncHI1 plasmid R27 encodes the *rsp* and *rsp*2 genes which code respectively

for the 155.4 kDa RSP and 86.75 kDa RSP2 proteins. These proteins interact and target both the

flagella and the conjugative pili to favor conjugation (Hüttener et al., 2022). Big proteins are also

encoded at least by two other groups of plasmids (IncA/C and IncP2) (Hüttener et al., 2022). Due

to their extracellular location, these plasmid-encoded Big proteins could be considered as

valuable antigens to combat antimicrobial resistance.

Functional heavy-chain-only antibodies (HCAbs) were discovered nearly 30 years ago in the

serum of camelids (Hamers-Casterman et al., 1993). HCAbs lack both the light chains and the

first constant CH1 domain within the heavy chain. The variable domain of HCAbs, also termed

VHH for <u>V</u>H of <u>H</u>CAb, represents the smallest antigen-binding unit that occur in nature

(Muyldermans, 2013). The VHH domains can be expressed in recombinant form as highly stable

single-domain Ab fragments of ~14 kDa, also termed Nanobodies (Nbs), which exhibit high

affinity and specificity to its cognate antigens and unique recognition of cryptic epitopes not

recognized by conventional Abs (Muyldermans, 2013). Nbs can be easily produced in bacteria

and other microbial cultures, and used in multiple diagnostic and therapeutic applications,

including bacterial infections (Muyldermans, 2021).

In this study, our objective was to investigate the potential of the recently discovered Big

proteins encoded by IncHI plasmids as targets for combating antibiotic resistance. Our research

focused on two key aspects. First, we examined the efficacy of the RSP protein as a vaccine

antigen in providing protection against infections caused by resistant bacteria carrying an IncHI

plasmid. The results obtained confirmed the efficacy of the RSP protein as a vaccine antigen.

Furthermore, we also generated RSP-specific Nbs and showed that they can interfere with the

conjugative transfer of the R27 plasmid. Hence, the use of antigens that are directly linked to

AMR determinants can be effective in combating infections caused by AMR bacteria.

Results

Ampicillin protects mice from infection with the *S.* Typhimurium SL1344 (WT) strain, but not

from infection with the *S.* Typhimurium SL1344 (pHCM1) strain.

To validate the RSP protein as a vaccine antigen against an antibiotic-resistant *Salmonella* strain,

we employed a mouse infection model. We first established that both *Salmonella* strains (WT

and pHCM1) were capable of infecting mice, and that only the plasmid-bearing strain was

antibiotic resistant. For antibiotic treatment, we specifically selected ampicillin (Amp) due to its

established efficacy in reducing the symptoms of *Salmonella* infection in mice (Butler et al.,

1981; Rossi et al., 2017).

Infection with the *Salmonella* strain SL1344 (WT) reduced the survival rate of the mice by about

40% (Fig. 1A), while the addition of Amp increased the survival rate of these animals. On the

other hand, Amp treatment did not change the survival of mice infected with strain SL1344

(pHCM1).

In antibiotic-naive mice infected with either SL1344 (WT) or SL1344 (pHCM1) strains, gradual

weight loss and clinical signs of disease were observed (Figs. 1B and 1C). However, Amp

treatment only ameliorated these symptoms in mice infected with the SL1344 (WT) strain,

showing no effect on those infected with the SL1344 (pHCM1) strain. Additionally, infection with

both *Salmonella* strains led to an upregulation of proinflammatory cytokines in the spleen (Fig.

1D, 1E, 1F). Amp treatment in mice infected with the SL1344 (WT) strain prevented this increase

in cytokine expression.

Immunization of mice with the RSP protein.

We then purified the RSP protein and employed it as a vaccine antigen to evaluate its efficacy in

protecting mice against infection caused by the *S*. Typhimurium SL1344 strain carrying the

pHCM1 plasmid. To prevent the recurrence of infection by SL1344 clones that spontaneously

cured the pHCM1 plasmid in the absence of Amp, we maintained the antibiotic treatment in the

immunized animals.

The immunization process did not have a significant impact on the body weight progression of

the mice (Fig. 2A). Although the overall concentration of total serum IgG remained unchanged

following immunization (Fig. 2B), there was a substantial increase in the levels of specific IgG

antibodies against the RSP protein (Fig. 2C). In terms of intestinal IgA, immunization resulted in

an elevated concentration of total IgA in both the jejunum content and feces (Fig. 2D and 2H,

respectively), while not affecting its presence in the colon (Fig. 2F). Moreover, immunization led

to enhanced production of specific IgA antibodies against the RSP protein in all three tissue types

(intestinal contents Fig. 2E; colon contents Fig. 2G; and feces Fig. 2I).

Effects of immunization of mice on the infection caused by the _S._ Typhimurium SL1344

(pHCM1) strain.

<u>Survival, body weight evolution and clinical signs</u>

Infection with the SL1344 (pHCM1) strain resulted in a decrease in the survival rate of mice and

an approximate 20% reduction in their body weight after 6 days of infection. Additionally, it led

to an increase in the manifestation of clinical signs associated with infection (Fig. 3). However,

immunization significantly improved the survival rate of mice and reduced the incidence of

clinical signs in infected animals.

<u>Intestinal effects of immunization with the RSP protein and challenge with the SL1344 (pHCM1)</u>

<u>strain</u>

Immunization with the RSP protein increased the secretion of total IgA into the lumen of the

jejunum and colon (Figs. 4A and B), while reducing its concentration in feces (Fig. 4C).

Furthermore, immunization with the RSP protein led to an elevated presence of specific IgA

antibodies against the RSP protein in all three samples (Fig. 4D-F), with even higher levels

observed in infected animals in the case of fecal content. Regarding the expression of

proinflammatory cytokines, infection with the SL1344 (pHCM1) strain resulted in increased

expression of all three cytokines analyzed (Fig. 4G-I) both in immunized and non-immunized

animals. In the case of _Tnf-α_, immunization also increased its expression.

<u>Systemic effects of immunization with the RSP protein and challenge with the SL1344 (pHCM1)</u>

<u>strain</u>

The immunization protocol also resulted in an increase in the concentration of total IgG at the

systemic level (Fig. 5A), while infection with the SL1344 (pHCM1) strain reduced it. Additionally,

immunization also led to an elevated concentration of specific anti-RSP IgG in the serum (Fig.

5B), which remained unaffected after infection with the SL1344 (pHCM1) strain. Following

bacterial challenge, the spleen weight increased nearly threefold in non-immunized mice (Fig.

5C). In contrast, immunized mice exhibited a significantly lower increase in spleen weight.

Furthermore, animals infected with the SL1344 (pHCM1) strain displayed *Salmonella*

colonization in the spleen (Fig. 5D), which was significantly reduced in immunized animals.

Regarding the expression of proinflammatory cytokines in the spleen, *Salmonella* infection

upregulated the expression of all cytokines evaluated (Fig. 5E-G). Immunization partially

mitigated this effect on Il-1β and Tnf-α expression.

Selection of nanobodies with neutralizing activity against the RSP protein.

Given the protective effect against *Salmonella* infection observed in mice immunized with the

RSP protein, we wondered whether we could obtain specific antibodies against RSP that could

directly interfere with its biological activity in the conjugative transfer of IncHI plasmids. We

chose nanobodies as ideal candidates to block RSP activity given their small size, stability and

high affinity to antigens, and their therapeutic potential against infections. Interestingly,

nanobodies can also be expressed on the surface of *E. coli* fused to the initial 654 amino acids

of the enterohemorrhagic *E. coli* intimin protein, termed Neae (Bodelón et al., 2009), comprising

the N-terminal signal peptide, the periplasmic LysM domain, the β-barrel for outer membrane

insertion, and the extracellular Big domains D00 and D0 (Fairman et al., 2012). Expression of

Neae-Nb fusions represent an efficient bacterial display system for their selection of Nbs binding

a target antigen from VHH gene libraries employing *E. coli* cell sorting methods with the labelled

antigen (Salema et al., 2013; Salema and Fernández, 2017). In addition, Neae-Nb fusions

expressed in *E. coli* promote synthetic cell-to-cell adhesions mediated by the specific recognition

of a surface antigen in the target cell (Glass and Riedel-Kruse, 2018; Piñero-Lambea et al., 2015).

In order to obtain Nbs that specifically bind the extracellular domain of the RSP protein, we

immunized two dromedaries with the last 280 residues of the C-terminal domain of the RSP

protein. After immunization, peripheral blood lymphocytes from these animals were isolated

and the VHH gene segments were amplified and cloned in the display vector pNeae2 (Robledo

et al., 2022; Salema et al., 2013), which encodes the Neae intimin fragment under the control of

the IPTG-inducible lacI-Plac promoter region. A library of approximately $1x10^8$ clones was

obtained by electroporation of *E. coli* DH10BT1R strain with the products of the ligated pNeae2-

VHH. Bacteria in library were grown in liquid culture and induced with IPTG. Induced bacteria

binding the C-domain of RSP were selected by incubation with the purified protein labeled with

biotin, following iterative steps of magnetic cell sorting (MACS) and fluorescence-assisted cell

sorting (FACS) (Materials and Methods). After the final sorting step (Fig. 6A), the selected pool

of bacteria was grown on LB-agar plates. 96 colonies were randomly picked from these plates,

grown in liquid, and induced with IPTG to evaluate the Nb display levels on the bacterial surface

and their binding capability to the biotinylated C-domain of the RSP protein by flow cytometry

(Fig. 6B). The VHH sequences from clones with specific binding to purified RSP were determined,

which allowed us to identify 7 different Nbs (Supplementary Fig. 1).

Selection of Nb clones displayed on *E. coli* that agglutinate with the *S*. Typhimurium SL1344

(R27) strain.

For selection of Nbs binding native RSP expressed on *Salmonella* cells, we took advantage of the

specific adhesion observed between *E. coli* cells with displayed Nbs and bacteria having the

target antigen accessible on their surface, which leads to the agglutination of mixed bacterial

cultures (Glass and Riedel-Kruse, 2018; Robledo et al., 2022).

We initially evaluated the agglutination capability of the 7 different clones obtained in the

enrichment protocol of the immune library, from strain *E. coli* DH10BT1R, each carrying a

pNeae2-VHH derivative expressing RSP-specific Nbs, against the *S*. Typhimurium SL1344 (R27)

strain. To achieve this, we cultured the *Salmonella* strain in LB medium to an $OD_{600 nm}$ of 2.

Following centrifugation, cells were resuspended in PBS and mixed with an equal volume of a

culture of the different *E. coli* DH10BT1R clones displaying the different set of RSP-specific Nbs.

Only one of the clones tested (Nb 3) showed ability of agglutinating the *S*. Typhimurium SL1344

(R27) strain (Fig. 7A). Next, we inspected this phenomenon using electron microscopy (Fig. 7B)

The presence of the R27 plasmid dictated the formation of clumps between the SL1344 strain

and *E. coli* DH10BT1R (pNeae2 VHH-RSP #3). In contrast, no clumps or aggregates were observed

when the plasmid-free SL1344 or the SL1344 (R27 Δ*rsp*) strains were used.

To further characterize this clone, the corresponding VHH gene was cloned in the mammalian

expression vector pIgΔCH1 (Casasnovas et al., 2022), which allows the fusion of the selected Nb

domain to the Fc region of human IgG1. The corresponding Nb3-Fc protein fusion was purified

from culture supernatants of transfected Expi293F cells (Fig. 8A), and its binding capacity to the

RSP protein was analyzed by ELISA. The results indicated that this Nb3-Fc presented a high

affinity for binding to the RSP protein, showing an apparent equilibrium dissociation constant

(K_D) in the picomolar range (~44 pM) (Fig. 8B).

Interference of *E. coli* displaying Nb-RSP #3 strain on the conjugation of the R27 plasmid from

the *Salmonella* Typhimurium SL1344 strain.

Our objective was to investigate whether the *E. coli* surface expression of the Nb-RSP #3 would

be able interfere with the conjugation process of the R27 plasmid from the SL1344 *Salmonella*

strain (donor) to the SL1344 *ibpA::lacZ-Km^r* derivative (recipient). To prevent that the Nbs

produced by the *E. coli* DH10BT1R (pNeae2 VHH-RSP #3) were titrated out by an excess of

SL1344 (R27) donor cells, we planned a conjugation experiment with a low number of donor

cells (10^3 cfu) mixed with an abundance of recipient cells (SL1344 *ibpA::lacZ-Km^r*) (10^6 cfu), along

with an excess of cells corresponding to the *E. coli* strain displaying Nb-RSP #3 (10^7 cfu).

Considering the known conjugation frequency of the R27 plasmid in *Salmonella* at 25°C, which

is approximately 10^{-3}, and the low number of donor cells used in our experiment, we chose

specific time intervals of 24, 36 and 48 hours following the mating process to detect

transconjugants within the mating mixture. To accomplish this, the mating mixture was mixed

with PB medium and incubated at 25°C. To confirm that any effect on the conjugation frequency

was solely due to the production of RSP-specific Nbs and not to other non-specific effects of Nb

expression, we utilized as controls both a plasmid-free *E. coli* DH10BT1R strain and the same

strain displaying an unrelated Nb binding human fibrinogen (Salema et al., 2016). The results

obtained (Fig. 9) indicated that the surface display of Nb-RSP #3 by the interfering *E. coli* strain

led to a significant and specific reduction in the conjugation frequency of the R27 plasmid in the

S. Typhimurium SL1344 strain. Interfering *E. coli* strains with no Nb or a control Nb elicited a

nonspecific ~10-fold reduction in the number of transconjugants of R27 plasmid in all time

points, compared to conjugation levels between *Salmonella* SL1344 strains. In contrast,

interfering *E. coli* displaying Nb-RSP #3 reduced the number of transconjugants of R27 up to

~1000-fold at 24h and ~100-fold after 36 and 48h of mating (Fig. 9). These results suggest a

strong inhibitory effect in the conjugation of R27 due to the interaction between bacterial

display Nb-RSP #3 and RSP protein expressed by donor cells. To rule out that the observed

reduction in the conjugation frequency in mixtures containing the *E. coli* strain DH10BT1R

(pNeae2 VHH-RSP #3) and the SL1344 (R27) strains could be due to the specific transfer of the

R27 plasmid to the interferent *E. coli* DH10BT1R (pNeae2 VHH-RSP #3) strain, we also detected

transconjugants of the *E. coli* DH10BT1R (pNeae2 VHH-RSP #3) and *E. coli* DH10BT1R (pNVFib1)

interferent strains by plating them on LB medium containing Tc (R27 plasmid marker) and Cm

(pNeae2 plasmid marker). In both experiments the conjugation frequency was the same (1.5×10^{-2}

transconjugants/donor cells).

Discussion

Vaccination stands as a priority strategy in addressing the challenges posed by AMR (Frost et al.,

2022; Micoli et al., 2021). Vaccines of special interest should target the agents responsible for

significant health care-associated infections, which include multiple resistant Gram-negative

bacteria (Lipsitch and Siber, 2016). Protecting against specific pathogens (i.e., *S. pneumoniae, K.*

pneumoniae) reduced the incidence of infections caused by them, leading to a decrease in

antibiotic usage and hence reducing the proliferation of AMR strains (Buchy et al., 2020).

One of the challenges in vaccine development is the identification of broadly protective antigens

(Troisi et al., 2020). In the context of AMR, current strategies focus on targeting either the most

common antigens among AMR clones of a specific pathogen (Micoli et al., 2021) or specific

virulence determinants (Moriel et al., 2010; Nesta et al., 2012). However, an alternative

approach is to develop a vaccine that directly targets a resistance determinant, which would be

effective not only against a specific microorganism but also against any microorganism

expressing the corresponding resistance determinant. Examples of successful targeting include

penicillin-binding proteins or beta-lactamases, which have demonstrated protection against

infections caused by *Staphylococcus aureus* (Senna et al., 2003), *Neisseria meningitidis*

(Zarantonelli et al., 2006), or *Pseudomonas aeruginosa* (Ciofu et al., 2002).

The novelty of the approach presented in this work is that, instead of targeting an antigen

involved in an antibiotic resistance mechanism, we focus on an antigen associated with the

transmission of resistance determinants.

Big proteins have been previously shown to be good vaccine candidates (Khanum et al., 2022;

Koizumi and Watanabe, 2004). The antigenicity of the RSP protein has been demonstrated by

the presence of higher levels of specific immunoglobulin, IgG in serum and IgA in the colon,

jejunum, and feces. In terms of the protective effect of immunization with the RSP protein in

mice, the results can be interpreted as follows. The secretion of proinflammatory cytokines in

the colon was comparable between immunized and non-immunized animals, as well as the

reduction in body weight. However, clinical symptoms, *Salmonella* counts in the spleen, and the

weight and secretion of proinflammatory cytokines in this organ were significantly reduced in

immunized animals. While immunization with the RSP protein did not provide complete

protection against intestinal infection with the SL1344 strain, it did result in attenuated disease

symptoms and restricted systemic infection. These findings support further research to validate

the RSP protein as a valuable antigen for protection against bacterial infections caused by enteric

bacteria harboring IncHI plasmids.

IncHI plasmids are commonly found in AMR bacteria isolated from clinical samples (Parvez and

Khan, 2018; Zhang et al., 2021). They are also frequently identified in *Salmonella* isolates (Phan

and Wain, 2008). The emergence of AMR *S. typhi* isolates has been reported in developing

countries (Tanmoy et al., 2018). Developing an effective vaccine against this pathogen would be

particularly beneficial for children in endemic countries, including regions such as South Asia,

Southeast Asia, and sub-Saharan Africa, as well as for travelers (World Health Organization,

2019). The RSP protein shows promise as a valuable antigen to be included in such a vaccine,

offering potential protection against AMR *Salmonella* infections associated with IncHI plasmids.

An alternative approach to provide protection against infections caused by virulent AMR strains

carrying IncHI plasmids is to specifically target the RSP protein with antibodies. Monoclonal

antibodies against various bacterial pathogens have demonstrated effectiveness, with some

currently in advanced stages of clinical trials (Cook and Wright, 2022). Nanobodies, a novel class

of antibodies, offer a wide range of clinical applications (Graf et al., 2019; Jovčevska and

Muyldermans, 2020a). The use of nanobodies specifically targeting the RSP protein represents

a promising approach to combat infections caused by pathogenic bacteria harboring IncHI

plasmids. Inhibiting the transfer of antimicrobial resistance plasmids has been identified as a

crucial strategy to combat AMR (Buckner et al., 2018; Graf et al., 2019). Our study demonstrates

that *E. coli* strains producing RSP-specific nanobodies can bind to *Salmonella* cells carrying the

R27 plasmid, resulting in a reduction in the conjugation frequency.

Expressing RSP-specific nanobodies in a probiotic bacterium, such as *E. coli* Nissle 1917, can help

restrict the transfer of IncHI plasmids in natural environments. Furthermore, RSP-specific

antibodies could be utilized to combat infections caused by pathogenic isolates carrying IncHI

plasmids. Nanobodies, known for their heat- and acid-stability and solubility, show potential for

oral administration to control gastrointestinal infections (Yuki et al., 2023). Thus, either purified

RSP-specific nanobodies or probiotic microbial cells expressing these nanobodies could be orally

administered to combat infections caused by AMR virulent strains harboring IncHI plasmids.

The expression of plasmid-encoded Big proteins is not exclusive to IncHI plasmids; other

plasmids such as IncA/C and IncP2 also encode these proteins (Hüttener et al., 2022). IncA/C

plasmids were initially identified in the 1970s among multidrug-resistant *Aeromonas hydrophila*

and *Vibrio* strains that infected cultured fish (Aoki et al., 1971). Since the 1990s, IncA/C plasmids

have gained significant attention due to their ability to mobilize antimicrobial resistance

determinants in enterobacteria and other Gram-negative microorganisms (Call et al., 2010;

Fernández-Alarcón et al., 2011; Fricke et al., 2009; Welch et al., 2007). These plasmids have an

exceptionally wide host range, encompassing Beta-, Gamma-, and Deltaproteobacteria (Suzuki

et al., 2010), and they play a crucial role in the global dissemination of antimicrobial resistance

(Eda et al., 2020; Rozwandowicz et al., 2018). In a recent study, IncA/C plasmids accounted for

50% of all plasmids isolated from clinical *K. pneumoniae* strains harboring the *blaNDM* gene

(Qamar et al., 2021). Notably, the monophasic variant of *S*. Typhimurium, *S. enterica* serovar

4,[5],12:i:-, which has emerged as a global cause of multidrug-resistant salmonellosis,

predominantly harbors IncHI1 and IncA/C resistance plasmids (Ingle et al., 2021).

Therefore, targeting these plasmid-encoded Big proteins through immunotherapies can be a

valuable strategy to combat infections caused by pathogenic microorganisms carrying various

types of resistance plasmids.

Methods

Bacterial strains and growth conditions.

The bacterial strains (see Supplementary Table S1) were routinely grown in Luria-Bertani (LB)

medium (10 g/L NaCl, 10 g/L tryptone and 5 g/L yeast extract) with vigorous shaking at 200 rpm

(Innova 3100; New Brunswick Scientific). The antibiotics used were chloramphenicol (Cm) (25

µg/mL), tetracycline (Tc) (15 µg/mL), carbenicillin (Cb) (100 µg/mL), and kanamycin (Km) (50

µg/mL) (Sigma-Aldrich). *E. coli* strains carrying the plasmid with a VHH gene were grown at 30ºC

on LB-agar plates (1.5% w/v) or in LB liquid medium with the addition of chloramphenicol (Cm,

30 µg/mL) for plasmid selection. To overexpress the nanobodies in the bacterial surface of *E.*

coli DH10BT1R, firstly bacteria were grown overnight (O/N) under statics conditions in LB

medium containing 2% (w/v) of glucose. Then, bacteria were harvested by centrifugation (4000

g for 5 minutes) and grown in the same media supplemented with 0.05 mM isopropylthio-β-D-

galactoside (IPTG) in constant agitation (160 rpm) for 16 hours.

Plasmids.

The different plasmids used in this work are listed in Supplementary Table S1.

Oligonucleotides.

The oligonucleotides used in this work are listed in Supplementary Table S2.

Genetic manipulations.

All enzymes used to perform standard molecular and genetic procedures were used according

to the manufacturer's recommendations. To introduce plasmids into *E. coli, Salmonella* and

Klebsiella, bacterial cells were grown until an $O.D_{600\ nm}$ of 0.8. Cells were then washed several

times with 10% glycerol, and the respective plasmids or DNA were electroporated by using an

Eppendorf gene pulser (Electroporator 2510).

Deletion of the *flhDC* locus was performed in the strain SL1344 by using the λ Red recombination

method, as previously described (Datsenko and Wanner, 2000).

Conjugative transfer of the R27 plasmid.

The R27 plasmid was conjugated as described previously (Hüttener et al., 2019). The mating

frequency was calculated as the number of transconjugants per donor cell.

Purification of the native RSP protein for the immunization assay.

The RSP protein was purified from cell-free supernatant fractions obtained from the *Salmonella* cells lacking the flagella (SL1344 R27 *flhDC::Km*). Cells were grown at 25ºC in LB medium for 16 hours in constant shaking at 200 rpm, then centrifugated at 7000 rpm for 30 minutes at room temperature. Cell pellets were discarded, and the supernatant was filtered through a 0.22 μm filter (Millipore). Supernatants were then ultra-centrifugated at 40000 rpm for 1 hour at 4ºC and carefully aspired to not disturb the protein pellet and discarded. The remaining pellet was solubilized using commercial Phosphate-buffered saline (PBS – Gibco). Proteins were quantified using the Bradford standard method (Pierce).

Purification of the C-terminal domain of the RSP protein.

For the RSP protein production the carboxyl-terminal region of the RSP protein was amplified (280 AA in total). Amplification of that region was achieved by performing PCR using the R27 plasmid as a DNA template and the primers RSP5 BamHI Fw and RSP PstI Rv, together with the Phusion Hot Start II High-fidelity DNA Polymerase (Thermo Fisher Scientific) following the manufacturer's recommendations. The DNA was then purified using the GeneJet PCR Purification Kit (Thermo Fisher Scientific), digested with BamHI and PstI restriction endonucleases according to the manufacturer´s instructions (New England Biolabs), and ligated into the pMAL-p2E vector digested with the same restriction endonucleases using the T4 DNA Ligase (New England Biolabs) according to the manufacturer´s instructions. The resulting plasmid, termed pMAL-RSP#5/7, was Sanger sequenced and transformed into BL21 cells. Cells transformed with pMAL-RSP#5/7 plasmid were grown in LB medium supplemented with carbenicillin at a final concentration of 100 µg/mL at 37°C for 16 hours at constant agitation at 200 rpm. Then, cells were diluted 1:100 in LB with 100 µg/mL of carbenicillin and glucose at 0.45% of the final concentration. Cells were incubated at 37ºC with 200 rpm until reached the O.D$_{600\ nm}$ of 0.5, then IPTG was added at a final concentration of 0.15 mM. Cells were incubated

at 37ºC for 2 hours under constant agitation of 200 rpm for the induction of the recombinant

protein. Cells were then centrifuged at 7500 rpm for 30 minutes at 4°C. The cells pellet was

subsequently resuspended in column buffer (20 mM HEPES (Na) pH 7.5, 200 mM NaCl, 1 mM

EDTA, 1 mM β-mercaptoethanol) plus protease inhibitor (Complete Ultra Tablets, Mini, EDTA-

free, EASYpack, Roche). Cells were then disrupted by French press cell lysis and centrifugated at

12000 rpm for 1 hour at 4°C. Supernatants were filtered through a 0.22 μM filter (Millipore) and

mixed with 250 μl of Amylose resin (New England Biolabs) for 2 hours at 4ºC in an orbital shaker.

The resin was then loaded in chromatography columns and washed twice with the column

buffer. The recombinant RSP C-terminal protein was then eluted using the elution buffer (20

mM HEPES (Na) pH 7.5, 200 mM NaCl, 1 mM EDTA, 1 mM β-mercaptoethanol and 10 mM

maltose). Eluted fractions were collected and then concentrated using Amicon Ultra-15 Ultracel

10K (Millipore) according to the manufacturer´s instructions.

Immunization of dromedaries with the C-terminal fragment of the RSP protein and selection

of clones producing anti-RSP specific nanobodies.

Two different dromedary camels (*Camelus dromedarious*) were immunized with purified RSP

protein (~1.2 mg) mixed with veterinary vaccine adjuvant (GERBU). After 6 rounds of

inoculations by subcutaneous injections, serial dilutions of pre-immune and immune sera were

used in ELISA to confirm the immune response of dromedary after the RSP protein inoculation.

B lymphocytes from peripheral blood were isolated, RNA was extracted and VHH genes were

amplified by RT-PCR using the VHH Sfi and VHH Not oligonucleotides following the protocol

described previously (Salema et al., 2013).

The amplified VHH fragments were digested with SfiI and NotI restriction enzymes and ligated

into the pNeae2 backbone vector digested with the same restriction enzymes, and finally

transformed in *E. coli* DH10BT1R by electroporation. Following this protocol, a library of 8.75x10[7]

clones was obtained.

This library was used for the enrichment of clones that recognized the RSP protein. To this end,

purified RSP protein was mixed with Biotin-NHS (Biotinamidocaproate N-hydroxysuccinimide

ester, Sigma Aldrich) and three rounds of MACS (with RSP concentrations of 100 nM (the first

one) and 50 nM (the other two)) were performed. Thereafter, a FACS selection with a protein

concentration of 50 nM were performed). The positive clones for the RSP binding were selected

for further analysis.

Cloning and purification of the RSP specific nanobody (VHH-RSP #3).

The gene coding for the RSP-specific selected Nb was PCR amplified with oligonucleotides VHH

pIg AgeI and VHH pIg BamHI using the pNeae2 VHH-RSP #3 purified plasmid as a template. This

PCR product was subsequently cloned in AgeI-BamHI sites of the mammalian expression vector

pIgΔCH1 (Casasnovas et al., 2022) derived from pIgγ1HC (Tiller et al., 2008). This vector allows

the expression of the cloned gene located in frame with a IgH signal peptide fused to the human

IgG1 hinge and Fc portion. For the nanobody overexpression and purification, the Expi293™

Expression System Kit (Thermo Fisher Scientific) was used. The mammalian cells transfected with

the purified expression vector (pIgΔCH1 VHH-RSP) were grown 5 days post-transfection. Cells

were then centrifuged, the supernatant was collected and filtered, and the proteins were

purified with an Ig Select or protein A affinity columns (Cytiva), following manufacturer's

instructions.

Enzyme-linked immunosorbent assays to evaluate the specific VHH-RSP #3 binding to the RSP

protein.

ELISA experiments were performed in 96-well plates (Maxisorp, Nunc), which were previously

coated overnight at 4°C with 50 μl/well of the purified RSP protein antigen or BSA at 3 μg/ml in

PBS. Antigen-coated plates were washed with PBS and blocked with 200 μl/well of 3% (w/v)

skimmed milk in PBS at room temperature for 2 h. Afterwards, different dilutions of the VHH-

RSP (prepared in 3% (w/v) skimmed milk in PBS) were added to the wells for 1 h. After the

incubation, the wells were washed three times with PBS, and rabbit anti-human Fc IgG-HRP

(Jackson ImmunoResearch) was added to the plates (diluted 1:5000 in 3% (w/v) skimmed milk

in PBS) to detect the bound Nbs. After 1 h of incubation at room temperature, plates were

washed with PBS and developed with H_2O_2 and o-phenylenediamine (OPD; Merck-Sigma). The

Optical Density (OD) at 490 nm ($OD_{490\ nm}$) of plate wells was determined (iMark, Bio-Rad), and

the values were corrected with the background levels of OD_{490nm} of the wells without antigen

added.

***In vivo* immunization and infection of mice.**

Male and female C57BL/6 mice were purchased from Envigo (Bresso, Italy) and maintained

under stable temperature and humidity conditions with a 12-hour light 12-hour dark cycle and

free access to food and water.

<u>Set up of the infection protocol with the SL1344 (pHCM1) strain.</u>

To induce infection, the *S.* Typhimurium strain SL1344 (pHCM1) was chosen. An assay was

designed to assess whether infection with this strain is similar to infection with the plasmid-free

strain. Four groups of mice were infected with strains SL1344 and SL1344 (pHCM1), and two of

them were treated with 150 mg/kg/day ampicillin (Amp) (Sigma-Aldrich) in drinking water and

two of them were untreated. In addition, a group of uninfected control mice was included as a

reference group. Thus, the groups were: i) Control: uninfected mice; ii) Sal: SL1344-infected mice

not treated with Amp; iii) Sal-pHCM1: SL1344-pHCM1-infected mice not treated with Amp; iv)

Sal WT + Amp: SL1344-infected mice treated with Amp; v) Sal pHCM1 + Amp: SL1344-pHCM1-

infected mice treated with Amp. Three hours before the challenge with the bacterial suspension,

mice received an intraperitoneal injection of cimetidine (50 mg/kg; Sigma-Aldrich) to reduce acid

secretion and improve bacterial survival(Ren et al., 2014). To induce infection, 100 µl of the

bacterial suspension (10^9 cfu of *Salmonella*) was administered by oral gavage. During the

infection period the body weight of animals was monitored daily. Four to six days after infection

(depending on survival), the animals were evaluated for clinical signs of disease and fecal

samples were obtained. They were subsequently euthanized and samples of spleen, colon

mucosa, intestinal lavage and serum were obtained. Samples were obtained and processed as

described previously (Miró et al., 2023).

The parameters analyzed to score the clinical signs were: Coat care (normal, 0; slightly altered,

1; considerably altered, 2; very altered, 3); posture (normal, 0; slightly curved, 1; considerably

curved, 2; very curved, 3); movement (normal, 0; slightly slow, 1; considerably slow, 2; very slow,

3); defecation (normal, 0; soft stools, 1; watery stools, 2; liquid stools, 3); body weight loss (no

weight loss, 0; weight loss less than 5%, 1; weight loss between 5 and 10%, 2; weight loss greater

than 10%, 3).

<u>Immunization with the RSP protein and infection with the *S.* Typhimurium SL1344 (pHCM1)</u>

<u>strain.</u>

Regarding the adjustment of the immunization with the RSP protein as antigen, we proceeded

as previously described (Miró et al., 2023). Briefly, mice were intranasally administered with 3

μg of the RSP protein and 5 μg of cholera toxin (CT; Sigma-Aldrich) as immunological adjuvant.

Immunization was performed 3 times every 2 weeks (at 3, 5 and 7 weeks of age). Non-

immunized mice received CT alone. Mice were distributed randomly into two groups: Non-Imm

(Non-immunized, CT alone); Imm (Immunized mice). During the immunization period the body

weight of animals was monitored weekly. Two weeks after last immunization, animals were

euthanized and samples of feces, intestinal lavage and serum were obtained. Total and RSP-

specific immunoglobulins were determined as explained in the section Immunoglobulin

determination.

In order to evaluate the protective efficacy of immunization, the previously established infection

protocol with the *S.* Typhimurium SL1344 (pHCM1) strain was used. The experiments were

performed using a Latin design with two factors: immunization of mice using the RSP protein as

immunogen; and subsequently infected with the *S.* Typhimurium SL1344 (pHCM1) strain. The

animals were randomly distributed into four groups: i) non-immunized and non-infected mice

(Non-Imm); ii) non-immunized and *Salmonella*-infected animals (Sal); iii) immunized and non-

infected animals with *Salmonella* (Imm); iv) immunized and infected animals (Imm-Sal). The

infection was provoked two weeks after last immunization dose.

Bacterial counts in spleen.

502   Spleen samples were homogenized with sterile PBS (20 mg/mL) using a PRO200 homogenizer

503   (Pro-Scientific, USA) at 18000 g*,* and homogenates were used to count the presence of

504   *Salmonella* cells plated in MacConkey media agar plates.

Immunoglobulin concentration.

Total secreted IgA and anti-RSP-specific IgA were determined in intestinal and colon content and

in feces by sandwich enzyme-linked immunosorbent assay (ELISA) as previously described (Ren

et al., 2014). Briefly, plates were coated with anti-mouse IgA monoclonal antibody (1 µg/mL;

Sigma-Aldrich) in PBS to quantify total IgA concentration or RSP (20 µg/mL) in PBS to measure

RSP-specific IgA concentration. Mouse IgA (Bethyl) was used as a standard for the determination

of total IgA. Goat anti-mouse IgA (Bethyl) conjugated with horseradish peroxidase (HRP) was

used as the detection antibody.

The concentrations of total IgG and specific IgG against the RSP protein in serum were also

determined. Plates were coated with goat anti-mouse IgG (1 µg/mL; Sigma-Aldrich) to quantify

the concentration of total IgG or RSP (10 µg/mL) in PBS to measure the concentration of specific

IgG against RSP. Mouse IgG (Sigma-Aldrich) was used as a standard for the determination of

total IgG. Goat anti-mouse IgG conjugated with HRP (Sigma-Aldrich) was used as detection

antibody.

519 In all cases, o-phenylenediamine (OPD, 0.4 mg/mL; Sigma-Aldrich) was used as HRP substrate,

520 and the color intensity was measured at 492 nm in a microplate reader (Sunrise).

521 **Cytokine expression in spleen and colon.**

522 RNA extraction and reverse transcription were carried out as previously described (Rosell-

523 Cardona et al., 2022). RNA quality and quantity were assessed by spectrophotometry (NanoDrop

524 ND-1000; Thermo Fisher Scientific) and its integrity was determined with an Agilent 2100

525 Bioanalyzer (Agilent Technologies Inc.). In all cases, the RNA integrity was ≥ 9 and the A260/280

526 ratio was between 1.96 and 2.02. Total RNA was reverse-transcribed using an iScript™ cDNA

527 Synthesis Kit (Bio-Rad). For real-time PCR determinations, we used SsoAdvanced™ Universal

528 SYBR® Green Supermix (Bio-Rad). The primers used are described previously (Garcia-Just et al.,

529 2020). Real-time PCR was performed on a MiniOpticon Real-Time PCR System (Bio-Rad). Each

530 PCR run included duplicates of reverse transcript cDNA for each sample and negative controls

531 (reverse transcription-free samples, RNA-free sample). Quantification of the target gene

532 transcripts was conducted using hypoxanthine phosphoribosyl transferase 1 (*hprt1*) gene

533 expression as reference and was performed with the $2^{-\Delta\Delta CT}$ method (Schmittgen and Livak,

534 2008). Product fidelity was confirmed by melting curve analysis.

535 **Immunogold electron microscopy.**

536 Immunogold microscopy experiments were performed as previously described (Hüttener et al.,

537 2019).

538 **Statistical analysis.**

539 The results of mice experiments are presented as mean ± SEM, except for survival curves and

540 disease indices. Survival curves have been compared by Log-rank (Mantel-Cox) test and

541 expressed as a percentage of survival. Disease indices are expressed as median values and

542 quartiles and have been compared by the Kruskal-Wallis test. Continuous data were analyzed

with Levene's test to assess the homogeneity of variance and with the Shapiro-Wilk test to

assess distribution. Homogeneous and normally distributed data were analyzed with Student's

t-test (immunization-tuning experiments) or one-way analysis of variance (ANOVA) (infection-

tuning) followed by Fisher's least significant difference (LSD) post hoc test and two-way ANOVA

(immunization and infection experiments). Student's t-test was also used to determine

statistical significance of R27 conjugation frequency in mating experiments. Data were analyzed

using GraphPad Prism® software v.9.3.1 (GraphPad Software, Inc.). Differences were considered

significant at $P<0.05$.

References

Aoki T, Egusa S, Kimura T, Watanabe T. 1971. Detection of R factors in naturally occurring
Aeromonas salmonicida strains. *Appl Microbiol* **22**:716–717. doi:10.1128/aem.22.4.716-
717.1971

Bodelón G, Marín E, Fernández LA. 2009. Role of Periplasmic Chaperones and BamA
(YaeT/Omp85) in Folding and Secretion of Intimin from Enteropathogenic *Escherichia coli*
Strains. *J Bacteriol* **191**:5169–5179. doi:10.1128/JB.00458-09

Bodelón G, Palomino C, Fernández LÁ. 2013. Immunoglobulin domains in *Escherichia coli* and
other enterobacteria: From pathogenesis to applications in antibody technologies. *FEMS*
Microbiol Rev. doi:10.1111/j.1574-6976.2012.00347.x

Buchy P, Ascioglu S, Buisson Y, Datta S, Nissen M, Tambyah PA, Vong S. 2020. Impact of vaccines
on antimicrobial resistance. *International Journal of Infectious Diseases*.
doi:10.1016/j.ijid.2019.10.005

Buckner MMC, Ciusa ML, Piddock LJV. 2018. Strategies to combat antimicrobial resistance: Anti-
plasmid and plasmid curing. *FEMS Microbiol Rev*. doi:10.1093/femsre/fuy031

Butler T, Shuster CW, Dixon P. 1981. Treatment of experimental *Salmonella* typhimurium
infection with mecillinam and ampicillin. *Antimicrob Agents Chemother* **19**:328–331.
doi:10.1128/AAC.19.2.328

Call DR, Singer RS, Meng D, Broschat SL, Orfe LH, Anderson JM, Herndon DR, Kappmeyer LS,
Daniels JB, Besser TE. 2010. blaCMY-2-positive IncA/C plasmids from *Escherichia coli* and
Salmonella enterica are a distinct component of a larger lineage of plasmids. *Antimicrob*
Agents Chemother **54**:590–596. doi:10.1128/AAC.00055-09

Carattoli A. 2013. Plasmids and the spread of resistance. *International Journal of Medical*
Microbiology. doi:10.1016/j.ijmm.2013.02.001

Carattoli A. 2009. Resistance plasmid families in Enterobacteriaceae. *Antimicrob Agents*
Chemother. doi:10.1128/AAC.01707-08

Casasnovas JM, Margolles Y, Noriega MA, Guzmán M, Arranz R, Melero R, Casanova M, Corbera
JA, Jiménez-de-Oya N, Gastaminza P, Garaigorta U, Saiz JC, Martín-Acebes MÁ, Fernández
LÁ. 2022. Nanobodies Protecting From Lethal SARS-CoV-2 Infection Target Receptor
Binding Epitopes Preserved in Virus Variants Other Than Omicron. *Front Immunol* **13**.
doi:10.3389/fimmu.2022.863831

Ciofu O, Bagge N, Høiby N. 2002. Antibodies against β-lactamase can improve ceftazidime
treatment of lung infection with β-lactam-resistant *Pseudomonas aeruginosa* in a rat
model of chronic lung infection. *APMIS* **110**:881–891. doi:10.1034/j.1600-
0463.2002.1101207.x

Cook MA, Wright GD. 2022. The past, present, and future of antibiotics. *Sci Transl Med* **14**:7793.
doi:10.1126/scitranslmed.abo7793

Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in *Escherichia coli*
K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**:6640–6645.
doi:10.1073/pnas.120163297

Dolejska M, Villa L, Poirel L, Nordmann P, Carattoli A. 2013. Complete sequencing of an IncHI1
plasmid encoding the carbapenemase NDM-1, the ArmA 16s RNA methylase and a
resistance-nodulation-cell division/multidrug efflux pump. *Journal of Antimicrobial*
Chemotherapy **68**:34–39. doi:10.1093/jac/dks357

Eda R, Nakamura M, Takayama Y, Maehana S, Nakano R, Yano H, Kitasato H. 2020. Trends and
molecular characteristics of carbapenemase-producing Enterobacteriaceae in Japanese
hospital from 2006 to 2015. *Journal of Infection and Chemotherapy* **26**:667–671.
doi:10.1016/j.jiac.2020.02.002

Fairman JW, Dautin N, Wojtowicz D, Liu W, Noinaj N, Barnard TJ, Udho E, Przytycka TM, Cherezov
V, Buchanan SK. 2012. Crystal Structures of the Outer Membrane Domain of Intimin and
Invasin from Enterohemorrhagic *E. coli* and Enteropathogenic *Y. pseudotuberculosis*.
Structure **20**:1233–1243. doi:10.1016/j.str.2012.04.011

Fernández-Alarcón C, Singer RS, Johnson TJ. 2011. Comparative genomics of multidrug
resistance-encoding incA/C plasmids from commensal and pathogenic *Escherichia coli*
from multiple animal sources. *PLoS One* **6**. doi:10.1371/journal.pone.0023415

Forde BM, Zowawi HM, Harris PNA, Roberts L, Ibrahim E, Shaikh N, Deshmukh A, Sid Ahmed MA,
Al Maslamani M, Cottrell K, Trembizki E, Sundac L, Yu HH, Li J, Schembri MA, Whiley DM,
Paterson DL, Beatson SA. 2018. Discovery of *mcr-1* -Mediated Colistin Resistance in a Highly
Virulent *Escherichia coli* Lineage. *mSphere* **3**. doi:10.1128/mSphere.00486-18

Fricke WF, Welch TJ, McDermott PF, Mammel MK, LeClerc JE, White DG, Cebula TA, Ravel J.
2009. Comparative genomics of the IncA/C multidrug resistance plasmid family. *J Bacteriol*
191:4750–4757. doi:10.1128/JB.00189-09

Frost I, Balachandran A, Paulin-Deschenaux S, Sati H, Hasso-Agopsowicz M. 2022. The approach
of World Health Organization to articulate the role and assure impact of vaccines against
antimicrobial resistance. *Hum Vaccin Immunother*. doi:10.1080/21645515.2022.2145069

Garcia-Just A, Miró L, Pérez-Bosque A, Amat C, Polo J, Pallàs M, Griñán-Ferré C, Moretó M. 2020.
Dietary spray-dried porcine plasma prevents cognitive decline in senescent mice and
reduces neuroinflammation and oxidative stress. *Journal of Nutrition* **150**:303–311.
doi:10.1093/jn/nxz239

Glass DS, Riedel-Kruse IH. 2018. A Synthetic Bacterial Cell-Cell Adhesion Toolbox for
Programming Multicellular Morphologies and Patterns. *Cell* **174**:649-658.e16.
doi:10.1016/j.cell.2018.06.041

Graf FE, Palm M, Warringer J, Farewell A. 2019. Inhibiting conjugation as a tool in the fight
against antibiotic resistance. *Drug Dev Res* **80**:19–23. doi:10.1002/ddr.21457

Hamers-Casterman C, Atarhouch T, Muyldermans S, Robinson G, Hammers C, Songa EB,
Bendahman N, Hammers R. 1993. Naturally occurring antibodies devoid of light chains.
Nature **363**:446–448. doi:10.1038/363446a0

Holt KE, Phan MD, Baker S, Duy PT, Nga TVT, Nair S, Turner AK, Walsh C, Fanning S, Farrell-Ward
S, Dutta S, Kariuki S, WeillFranç FX, Parkhill J, Dougan G, Wain J. 2011. Emergence of a
globally dominant inchi1 plasmid type associated with multiple drug resistant typhoid.
PLoS Negl Trop Dis **5**. doi:10.1371/journal.pntd.0001245

Hüttener M, Hergueta J, Bernabeu M, Prieto A, Aznar S, Merino S, Tomás J, Juárez A. 2022. Roles
of Proteins Containing Immunoglobulin-Like Domains in the Conjugation of Bacterial
Plasmids. *mSphere* **7**. doi:10.1128/msphere.00978-21

Hüttener M, Prieto A, Aznar S, Bernabeu M, Glaría E, Valledor AF, Paytubi S, Merino S, Tomás J,
Juárez A. 2019. Expression of a novel class of bacterial Ig-like proteins is required for IncHI
plasmid conjugation. *PLoS Genet* **15**. doi:10.1371/journal.pgen.1008399

Ingle DJ, Ambrose RL, Baines SL, Duchene S, Gonçalves da Silva A, Lee DYJ, Jones M, Valcanis M,
Taiaroa G, Ballard SA, Kirk MD, Howden BP, Pearson JS, Williamson DA. 2021. Evolutionary
dynamics of multidrug resistant *Salmonella enterica* serovar 4,[5],12:i:- in Australia. *Nat*
Commun **12**. doi:10.1038/s41467-021-25073-w

Jovčevska I, Muyldermans S. 2020. The Therapeutic Potential of Nanobodies. *BioDrugs* **34**:11–
26. doi:10.1007/s40259-019-00392-z

Khanum S, Carbone V, Gupta SK, Yeung J, Shu D, Wilson T, Parlane NA, Altermann E, Estein SM,
Janssen PH, Wedlock DN, Heiser A. 2022. Mapping immunogenic epitopes of an adhesin-
like protein from Methanobrevibacter ruminantium M1 and comparison of empirical data
with in silico prediction methods. *Sci Rep* **12**:10394. doi:10.1038/s41598-022-14545-8

Koizumi N, Watanabe H. 2004. Leptospiral immunoglobulin-like proteins elicit protective
immunity. *Vaccine* **22**:1545–1552. doi:10.1016/j.vaccine.2003.10.007

Lawley TD, Gilmour MW, Gunton JE, Standeven LJ, Taylor DE. 2002. Functional and mutational
analysis of conjugative transfer region 1 (Tra1) from the IncHI1 plasmid R27. *J Bacteriol*
184:2173–2180. doi:10.1128/JB.184.8.2173-2180.2002

Lawley TD, Gilmour MW, Gunton JE, Tracz DM, Taylor DE. 2003. Functional and mutational
analysis of conjugative transfer region 2 (Tra2) from the IncHI1 plasmid R27. *J Bacteriol*
185:581–591. doi:10.1128/JB.185.2.581-591.2003

Lim LM, Ly N, Anderson D, Yang JC, Macander L, Jarkowski A, Forrest A, Bulitta JB, Tsuji BT. 2010.
Resurgence of colistin: A review of resistance, toxicity, pharmacodynamics, and dosing.
Pharmacotherapy. doi:10.1592/phco.30.12.1279

Lipsitch M, Siber GR. 2016. How can vaccines contribute to solving the antimicrobial resistance
problem? *mBio*. doi:10.1128/mBio.00428-16

Matamoros S, Van Hattem JM, Arcilla MS, Willemse N, Melles DC, Penders J, Vinh TN, Thi Hoa
N, Bootsma MCJ, Van Genderen PJ, Goorhuis A, Grobusch M, Molhoek N, Oude Lashof
AML, Stobberingh EE, Verbrugh HA, De Jong MD, Schultsz C. 2017. Global phylogenetic
analysis of *Escherichia coli* and plasmids carrying the mcr-1 gene indicates bacterial
diversity but plasmid restriction. *Sci Rep* **7**. doi:10.1038/s41598-017-15539-7

Meyer E, Schwab F, Schroeren-Boersch B, Gastmeier P. 2010. Dramatic increase of third-
generation cephalosporin-resistant *E. coli* in German intensive care units: Secular trends in
antibiotic drug use and bacterial resistance, 2001 to 2008. *Crit Care* **14**. doi:10.1186/cc9062

Micoli F, Bagnoli F, Rappuoli R, Serruto D. 2021. The role of vaccines in combatting antimicrobial
resistance. *Nat Rev Microbiol* **19**:287–302. doi:10.1038/s41579-020-00506-3

Miró L, Rosell-Cardona C, Amat C, Polo J, Moretó M, Pérez-Bosque A. 2023. Dietary
supplementation with spray-dried animal plasma improves vaccine protection in aged
mice. *Front Nutr* **10**. doi:10.3389/fnut.2023.1050961

Morens DM, Folkers GK, Fauci AS. 2004. The challenge of emerging and re-emerging infectious
diseases. *Nature* **430**:242–249. doi:10.1038/nature02759

Moriel DG, Bertoldi I, Spagnuolo A, Marchi S, Rosini R, Nesta B, Pastorello I, Corea VAM, Torricelli
G, Cartocci E, Savino S, Scarselli M, Dobrindt U, Hacker J, Tettelin H, Tallon LJ, Sullivan S,
Wieler LH, Ewers C, Pickard D, Dougan G, Fontana MR, Rappuoli R, Pizza M, Serino L. 2010.
Identification of protective and broadly conserved vaccine antigens from the genome of
extraintestinal pathogenic *Escherichia coli*. *Proceedings of the National Academy of*
Sciences **107**:9072–9077. doi:10.1073/pnas.0915077107

Murray CJ, Ikuta KS, Sharara F, Swetschinski L, Robles Aguilar G, Gray A, Han C, Bisignano C, Rao
P, Wool E, Johnson SC, Browne AJ, Chipeta MG, Fell F, Hackett S, Haines-Woodhouse G,
Kashef Hamadani BH, Kumaran EAP, McManigal B, Agarwal R, Akech S, Albertson S, Amuasi
J, Andrews J, Aravkin A, Ashley E, Bailey F, Baker S, Basnyat B, Bekker A, Bender R, Bethou
A, Bielicki J, Boonkasidecha S, Bukosia J, Carvalheiro C, Castañeda-Orjuela C, Chansamouth
V, Chaurasia S, Chiurchiù S, Chowdhury F, Cook AJ, Cooper B, Cressey TR, Criollo-Mora E,
Cunningham M, Darboe S, Day NPJ, De Luca M, Dokova K, Dramowski A, Dunachie SJ,
Eckmanns T, Eibach D, Emami A, Feasey N, Fisher-Pearson N, Forrest K, Garrett D,
Gastmeier P, Giref AZ, Greer RC, Gupta V, Haller S, Haselbeck A, Hay SI, Holm M, Hopkins
S, Iregbu KC, Jacobs J, Jarovsky D, Javanmardi F, Khorana M, Kissoon N, Kobeissi E,
Kostyanev T, Krapp F, Krumkamp R, Kumar A, Kyu HH, Lim C, Limmathurotsakul D, Loftus
MJ, Lunn M, Ma J, Mturi N, Munera-Huertas T, Musicha P, Mussi-Pinhata MM, Nakamura
T, Nanavati R, Nangia S, Newton P, Ngoun C, Novotney A, Nwakanma D, Obiero CW, Olivas-
Martinez A, Olliaro P, Ooko E, Ortiz-Brizuela E, Peleg AY, Perrone C, Plakkal N, Ponce-de-
Leon A, Raad M, Ramdin T, Riddell A, Roberts T, Robotham JV, Roca A, Rudd KE, Russell N,
Schnall J, Scott JAG, Shivamallappa M, Sifuentes-Osornio J, Steenkeste N, Stewardson AJ,
Stoeva T, Tasak N, Thaiprakong A, Thwaites G, Turner C, Turner P, van Doorn HR, Velaphi
S, Vongpradith A, Vu H, Walsh T, Waner S, Wangrangsimakul T, Wozniak T, Zheng P,
Sartorius B, Lopez AD, Stergachis A, Moore C, Dolecek C, Naghavi M. 2022. Global burden
of bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet* **399**:629–
655. doi:10.1016/S0140-6736(21)02724-0

Muyldermans S. 2021. Applications of Nanobodies. *Annu Rev Anim Biosci* **9**:401–421.
doi:10.1146/annurev-animal-021419-083831

Muyldermans S. 2013. Nanobodies: Natural single-domain antibodies. *Annu Rev Biochem*.
doi:10.1146/annurev-biochem-063011-092449

Nesta B, Spraggon G, Alteri C, Moriel DG, Rosini R, Veggi D, Smith S, Bertoldi I, Pastorello I,
Ferlenghi I, Fontana MR, Frankel G, Mobley HLT, Rappuoli R, Pizza M, Serino L, Soriani M.
2012. FdeC, a novel broadly conserved *Escherichia coli* adhesin eliciting protection against
urinary tract infections. *mBio* **3**. doi:10.1128/mBio.00010-12

Novick RP, Richmond MH. 1965. Nature and Interactions of the Genetic Elements Governing
Penicillinase Synthesis in *Staphylococcus aureus*. *J Bacteriol* **90**:467–480.
doi:10.1128/jb.90.2.467-480.1965

Parvez S, Khan AU. 2018. Hospital sewage water: a reservoir for variants of New Delhi metallo-
β-lactamase (NDM)- and extended-spectrum β-lactamase (ESBL)-producing
Enterobacteriaceae. *Int J Antimicrob Agents* **51**:82–88.
doi:10.1016/j.ijantimicag.2017.08.032

Phan M-D, Wain J. 2008. IncHI plasmids, a dynamic link between resistance and pathogenicity.
The Journal of Infection in Developing Countries **2**. doi:10.3855/jidc.221

Piñero-Lambea C, Bodelón G, Fernández-Periáñez R, Cuesta AM, Álvarez-Vallina L, Fernández LÁ.
2015. Programming Controlled Adhesion of *E. coli* to Target Surfaces, Cells, and Tumors
with Synthetic Adhesins. *ACS Synth Biol* **4**:463–473. doi:10.1021/sb500252a

Qamar MU, Ejaz H, Walsh TR, Shah AA, Al Farraj DA, Alkufeidy RM, Alkubaisi NA, Saleem S, Jahan
S. 2021. Clonal relatedness and plasmid profiling of extensively drug-resistant New Delhi
metallo-β-lactamase-producing *Klebsiella pneumoniae* clinical isolates. *Future Microbiol*
16:229–239. doi:10.2217/fmb-2020-0315

Ren W, Yin J, Duan J, Liu G, Zhu X, Chen S, Li T, Wang S, Tang Y, Hardwidge PR. 2014. Mouse
intestinal innate immune responses altered by enterotoxigenic *Escherichia coli* (ETEC)
infection. *Microbes Infect* **16**:954–961. doi:10.1016/j.micinf.2014.09.005

Robledo M, Álvarez B, Cuevas A, González S, Ruano-Gallego D, Fernández LÁ, de la Cruz F. 2022.
Targeted bacterial conjugation mediated by synthetic cell-to-cell adhesions. *Nucleic Acids*
Res **50**:12938–12950. doi:10.1093/nar/gkac1164

Rosell-Cardona C, Amat C, Griñán-Ferré C, Polo J, Pallàs M, Pérez-Bosque A, Moretó M, Miró L.
2022. The Neuroprotective Effects of Spray-Dried Porcine Plasma Supplementation Involve
the Microbiota–Gut–Brain Axis. *Nutrients* **14**. doi:10.3390/nu14112211

Rossi O, Dybowski R, Maskell DJ, Grant AJ, Restif O, Mastroeni P. 2017. Within-host
spatiotemporal dynamics of systemic *Salmonella* infection during and after antimicrobial
treatment. *Journal of Antimicrobial Chemotherapy* **72**:3390–3397. doi:10.1093/jac/dkx294

Rossolini GM, Mantengoli E, Docquier J-D, Musmanno RA, Coratza G. 2007. Epidemiology of
infections caused by multiresistant gram-negatives: ESBLs, MBLs, panresistant strains. *New*
Microbiol **30**:332–9.

Rozwandowicz M, Brouwer MSM, Fischer J, Wagenaar JA, Gonzalez-Zorn B, Guerra B, Mevius DJ,
Hordijk J. 2018. Plasmids carrying antimicrobial resistance genes in Enterobacteriaceae.
Journal of Antimicrobial Chemotherapy **73**:1121–1137. doi:10.1093/jac/dkx488

Salema V, Fernández LÁ. 2017. *Escherichia* coli surface display for the selection of nanobodies.
Microb Biotechnol **10**:1468–1484. doi:10.1111/1751-7915.12819

Salema V, López-Guajardo A, Gutierrez C, Mencía M, Fernández LÁ. 2016. Characterization of
nanobodies binding human fibrinogen selected by *E. coli* display. *J Biotechnol* **234**:58–65.
doi:10.1016/j.jbiotec.2016.07.025

Salema V, Marín E, Martínez-Arteaga R, Ruano-Gallego D, Fraile S, Margolles Y, Teira X, Gutierrez
C, Bodelón G, Fernández LÁ. 2013. Selection of Single Domain Antibodies from Immune
Libraries Displayed on the Surface of *E. coli* Cells with Two β-Domains of Opposite
Topologies. *PLoS One* **8**:e75126. doi:10.1371/journal.pone.0075126

Schmittgen TD, Livak KJ. 2008. Analyzing real-time PCR data by the comparative CT method. *Nat
Protoc* **3**:1101–1108. doi:10.1038/nprot.2008.73

Senna JPM, Roth DM, Oliveira JS, Machado DC, Santos DS. 2003. Protective immune response
against methicillin resistant *Staphylococcus aureus* in a murine model using a DNA vaccine
approach. *Vaccine* **21**:2661–2666. doi:10.1016/S0264-410X(02)00738-7

Sherburne CK, Lawley TD, Gilmour MW, Blattner FR, Burland V, Grotbeck E, Rose DJ, Taylor DE.
2000. The complete DNA sequence and analysis of R27, a large IncHI plasmid from
Salmonella typhi that is temperature sensitive for transfer. *Nucleic Acids Res* **28**:2177–86.
doi:10.1093/nar/28.10.2177

Spellberg B, Guidos R, Gilbert D, Bradley J, Boucher HW, Scheld WM, Bartlett JG, Edwards J.
2008. The epidemic of antibiotic-resistant infections: A call to action for the medical
community from the infectious diseases society of America. *Clinical Infectious Diseases*.
doi:10.1086/524891

Suzuki H, Yano H, Brown CJ, Top EM. 2010. Predicting plasmid promiscuity based on genomic
signature. *J Bacteriol* **192**:6045–6055. doi:10.1128/JB.00277-10

Tanmoy AM, Westeel E, De Bruyne K, Goris J, Rajoharison A, Sajib MSI, Van Belkum A, Saha SK,
Komurian-Pradel F, Endtz HP. 2018. *Salmonella enterica* Serovar Typhi in Bangladesh:
Exploration of Genomic Diversity and Antimicrobial Resistance. doi:10.1128/mBio.00105

Thomas CM, Nielsen KM. 2005. Mechanisms of, and barriers to, horizontal gene transfer
between bacteria. *Nat Rev Microbiol*. doi:10.1038/nrmicro1234

Tiller T, Meffre E, Yurasov S, Tsuiji M, Nussenzweig MC, Wardemann H. 2008. Efficient
generation of monoclonal antibodies from single human B cells by single cell RT-PCR and
expression vector cloning. *J Immunol Methods* **329**:112–124.
doi:10.1016/j.jim.2007.09.017

Troisi M, Andreano E, Sala C, Kabanova A, Rappuoli R. 2020. Vaccines as remedy for antimicrobial
resistance and emerging infections. *Curr Opin Immunol*. doi:10.1016/j.coi.2020.09.003

Villa L, Poirel L, Nordmann P, Carta C, Carattoli A. 2012. Complete sequencing of an IncH plasmid
carrying the bla ndm-1, bla ctx-m-15 and qnrB1 genes. *Journal of Antimicrobial
Chemotherapy* **67**:1645–1650. doi:10.1093/jac/dks114

Vrancianu CO, Popa LI, Bleotu C, Chifiriuc MC. 2020. Targeting Plasmids to Limit Acquisition and
Transmission of Antimicrobial Resistance. *Front Microbiol*. doi:10.3389/fmicb.2020.00761

Wang J, Stephan R, Zurfluh K, Hächler H, Fanning S. 2015. Characterization of the genetic
environment of blaESBL genes, integrons and toxin-antitoxin systems identified on large
transferrable plasmids in multi-drug resistant *Escherichia coli*. *Front Microbiol* **6**.
doi:10.3389/fmicb.2014.00716

Wang Y, Tian GB, Zhang R, Shen Y, Tyrrell JM, Huang X, Zhou H, Lei L, Li HY, Doi Y, Fang Y, Ren H, Zhong LL, Shen Z, Zeng KJ, Wang S, Liu JH, Wu C, Walsh TR, Shen J. 2017. Prevalence, risk factors, outcomes, and molecular epidemiology of mcr-1-positive Enterobacteriaceae in patients and healthy adults from China: an epidemiological and clinical study. *Lancet Infect Dis* **17**:390–399. doi:10.1016/S1473-3099(16)30527-8

Welch TJ, Fricke WF, McDermott PF, White DG, Rosso ML, Rasko DA, Mammel MK, Eppinger M, Rosovitz MJ, Wagner D, Rahalison L, LeClerc JE, Hinshaw JM, Lindler LE, Cebula TA, Carniel E, Ravel J. 2007. Multiple antimicrobial resistance in plague: An emerging public health risk. *PLoS One* **2**. doi:10.1371/journal.pone.0000309

World Health Organization. 2019. Typhoid vaccines: WHO position paper, March 2018 – Recommendations. *Vaccine* **37**:214–216. doi:10.1016/j.vaccine.2018.04.022

Yuki Y, Zuo F, Kurokawa S, Uchida Y, Sato S, Sakon N, Hammarström L, Kiyono H, Marcotte H. 2023. Lactobacilli as a Vector for Delivery of Nanobodies against Norovirus Infection. *Pharmaceutics* **15**. doi:10.3390/pharmaceutics15010063

Zarantonelli ML, Antignac A, Lancellotti M, Guiyoule A, Alonso JM, Taha MK. 2006. Immunogenicity of meningococcal PBP2 during natural infection and protective activity of anti-PBP2 antibodies against meningococcal bacteraemia in mice. *Journal of Antimicrobial Chemotherapy* **57**:924–930. doi:10.1093/jac/dkl066

Zhang Zhiren, Guo H, Li Xiaodong, Li W, Yang G, Ni W, Zhan M, Lu L, Zhang Zhenlin, Li Xiaobin, Zhou Z. 2021. Genetic Diversity and Characteristics of blaNDM-Positive Plasmids in *Escherichia coli*. *Front Microbiol* **12**. doi:10.3389/fmicb.2021.729952

Zheng B, Dong H, Xu H, Lv J, Zhang J, Jiang X, Du Y, Xiao Y, Li L. 2016. Coexistence of MCR-1 and NDM-1 in Clinical *Escherichia coli* Isolates. *Clinical Infectious Diseases*. doi:10.1093/cid/ciw553

Acknowledgments

This work was supported by grants from Fundació "La Marató TV3," Spain (project 201818 10), CERCA Program/Generalitat de Catalunya to AJ, PID2021-124676NB-I00 from the Ministerio de Economía, Industria y Competitividad to SM and JT. Work in the laboratory of LAF is supported by grants PLEC2021-007739 from MCIN/AEI NextGeneration EU/ PRTR and FET Open 965018-BIOCELLPHE from the European Union's Horizon 2020 Future and Emerging Technologies research and innovation program. LM and APB are members of the Institut de Recerca en Nutrició, i Seguretat Alimentària (INSA-UB), which is recognized as a Maria de Maeztu Unit of Excellence and funded by MICIN/AEI/FEDER (CEX2021-001234-M).

Contributions

Genetic manipulations, conjugative transfer of the RSP plasmid and purification of the variants

of the RSP protein were conducted by A. Prieto, M. Hüttener, D. Salguero and M. Bernabeu. *In*

vivo experiments with mice were conducted by L. Miró and A. Pérez-Bosque. Electron

microscopy studies were conducted by S. Merino and J. Tomás. Selection of clones producing

anti-RSP nanobodies and the analysis of the effect of the obtained nanobodies on the

conjugative transfer of the RSP plasmid were performed by Y. Margolles, L.A. Fernández and A.

Prieto. The initial manuscript draft was written by A. Prieto, A. Pérez-Bosque and A. Juárez. A.

Prieto, A. Pérez-Bosque, L.A. Fernández, M. Hüttener, and A. Juárez edited the final version. A.

Juárez was the PI of the project.

Ethics

The dromedary camel immunization protocol followed European Union and Spanish guidelines

of animal experimentation and was approved by the Ethics Committee for Animal

Experimentation from University of Las Palmas de Gran Canaria and the Consejería de

Agricultura, Pesca y Aguas of the Canary Islands Goverment (reference OEBA_ULPGC_20-2023).

Experimentation with mice was conducted at Animal Facility of Faculty of Pharmacy and Food

Science. Experiments were approved by the Ethics Committee of Animal Experimentation of the

University of Barcelona (Ref. 170/20). Animals were handled in strict accordance with the

guidelines of the European Community 86/609/CEE.

Competing interests

The authors declare no competing interests.

Figures

Figure 1. Infection of mice with the *S.* Typhimurium SL1344 and *S.* Typhimurium (pHCM1)

strains. Survival rate **(A)**, body weight evolution **(B)**, disease index **(C)** and proinflammatory

cytokine expression in spleen, *Il-6* **(D)**, *Il-1β* **(E)** and *Tnf-α* **(F)**. Groups treated with ampicillin

(Amp) are indicated with **+Amp**. In panel (A), survival rate of control animals and those infected

with the *S.* Typhimurium SL1344 (WT) strain and treated with Amp, are 100%. Animals infected

with the *S.* Typhimurium (pHCM1) strain have the same survival rate independently of the

treatment with Amp. Results are expressed as mean ± SEM (n= 6-8 animals). Means without a

common letter differ, P<0.05.

Figure 2. Effect of immunization with the RSP protein on body weight evolution (A) and

immunoglobulin concentration (B-I). Open bars represent non-immunized (**No-Imm**) mice;

solid bars represent immunized (**Imm**) mice. Results are expressed as mean ± SEM (n= 6-8

animals). IgA, immunoglobulin A; IgG, immunoglobulin G; nd, non-detected. Means without a

common letter differ, P<0.05.

Figure 3. Survival rate (A), body weight change (B) and clinical signs (C) after challenge with

the SL1344 (pHCM1) strain. Open bars/plots represent non-immunized (**No-Imm**) mice; solid

bars/plots represent immunized (**Imm**) mice. In panel B, results are expressed as mean ± SEM

(n= 12-14 animals). Means without a common letter differ, P<0.05.

Figure 4. Immunoglobulin concentration and cytokine expression in the gastrointestinal tract after RSP immunization and challenge with *Salmonella.* Total IgA concentration in jejunum content **(A)**, in colon content **(B)** and in feces **(C)**. RSP-specific IgA titers in jejunum content **(D)**, in colon content **(E)** and in feces **(F)**. *Il-1β* **(G)**, *Il-6* **(H)** and *Tnf-α* **(I)** expression in colon mucosa. Open bars represent non-immunized (**Non-Imm**) mice; solid bars represent immunized (**Imm**) mice. Results are expressed as means ± SEMs (n= 10–12 animals). Means without a common letter differ, P<0.05. IgA, immunoglobulin A; IgG, immunoglobulin G; Il, interleukin; Int, interaction between both factors; Tnf-α, tumor necrosis factor alpha.

Figure 5. Immunoglobulin concentration, spleen weight, *Salmonella* counts in spleen tissue and cytokine expression in spleen tissue after immunization with the RSP protein and challenge with the SL1344 (pHCM1) strain. Total IgG concentration **(A)** and RSP-specific IgG titers **(B)** in plasma. Spleen weight **(C)** and *Salmonella* counts in spleen tissue **(D)**. *Il-1β* **(E)**, *Il-6* **(F)** and *Tnf-α* **(G)** expression in spleen tissue. Results are expressed as means ± SEMs (n= 12-14 animals). Means without a common letter differ, P<0.05. IgA, immunoglobulin A; IgG, immunoglobulin G; Il, interleukin; nd, non-detected; Tnf-α, tumor necrosis factor alpha.

**Figure 6. Selection of RSP-binding Nbs from immune libraries displayed on the surface of *E.*

coli. (A)** Flow cytometry analysis of the enrichment of bacterial population displaying Nb libraries

generated after the immunization of dromedaries with the C-terminal domain of the RSP

protein. 3 rounds of selection with the RSP protein by magnetic cell sorting (MACS) and one

round of fluorescence activated cell sorting (FACS) were performed, in which bacteria were

incubated with 50 nM of biotin-labeled RSP (or 100 nM of fibrinogen used as a specificity control)

and stained with Streptavidin-APC. **(B)** Flow cytometry of the individual bacterial clones selected

from the immune library. The bacterial surface display of the corresponding nanobody was

897   detected using an anti-c-myc monoclonal antibody. The binding of biotin-labeled antigens to

898   Nbs displayed on bacteria was performed with the incubation of bacterial cells with 50 nM of

899   biotin-labeled RSP.

Figure 7. Analysis of the interaction of the VHH-RSP#3 isolated in the immune library with the native RSP protein at a macro and microscopic level. (A) Assay of the ability of *E. coli* DH10BT1R cells expressing nanobodies on its surface to agglutinate cells of the *S.* Typhimurium SL1344 (R27) strain. Blue arrow marks the aggregation of bacteria after the interaction of *E. coli* cells producing nanobodies directed against the RSP protein (VHH-RSP #3) and *Salmonella* cells harboring the R27 plasmid. (B) Transmission electron microscopy imaging mixtures of *E. coli* DH10BT1R cells expressing nanobodies against the RSP protein on its surface and cells of the *S.* Typhimurium SL1344 strain, harboring or not the R27 plasmid. The studies were performed by labeling the nanobodies producing bacteria (*E. coli* DH10BT1R) with a mouse antibody anti c-myc tag and goat anti-mouse IgG conjugated to 12 nm gold particles. Blue bars represent 1 µm while black bars represent 2 µm.

Figure 8. Characterization of the purified VHH-RSP#3 as a specific binder to the RSP protein.

(A) Purification of Nb3-Fc by affinity column. Coomassie staining of the SDS-PAGE (10%) showing the purified Nb-RSP #3-Fc from transfected mammalian cell culture supernatants. Molecular weight markers are indicated on the left. **(B)** ELISA showing the binding capacity of the purified VHH-RSP#3-Fc to the RSP or BSA protein. The plot represents the OD values at 490 nm obtained with the indicated concentrations of the purified Nb3-Fc after 1 hour of interaction with the corresponding antigen.

Figure 9. Interference of *E. coli* DH10BT1R cells expressing nanobodies on its surface with the

conjugative transfer of the R27 plasmid in *Salmonella*. R27 plasmid was conjugated from a

donor strain (SL1344) to a recipient strain (SL1344 *ibpA::lacZ-Km^r*) with the absence or presence

of the interferent strains *E. coli* DH10BT1R, plasmid-free clone; *E. coli* DH10BT1R (pNVFib1),

clone expressing a fibrinogen-specific VHH; *E. coli* DH10BT1R (pNeae2 VHH-RSP #3), clone

expressing RSP-specific VHH. Means without a common letter differ, P<0.05.