Flower size evolution in the Southwest Pacific

Riccardo Ciarle, Kevin C. Burns and Fabio Mologni

Address: Te Kura Mātauranga Koiora | School of Biological Sciences,

Te Herenga Waka | Victoria University of Wellington, PO Box 600, Wellington, New

Zealand

Email: riccardo.ciarle@vuw.ac.nz

Phone: +64-4-463-8873

Abstract

The island rule is arguably the most controversial topic in evolutionary biogeography. Previous work has focused almost exclusively on vertebrates. Little is known about the island rule in plants, and no previous study has tested for the island rule in flower size. Here, we provide the first test of the island rule in flowers. Using a newly derived dataset from the Southwest Pacific, we tested whether animal-pollinated and wind-pollinated flowers follow the island rule, or whether they exhibit directional size changes (i.e. gigantism or dwarfism). Results indicate that animal-pollinated flowers follow the island rule, while wind-pollinated flowers do not. Instead, they became consistently larger on islands. We conclude that, while *in situ* evolution of flower size is widespread on islands in the Southwest Pacific, species belonging to different pollination syndromes exhibit markedly different evolutionary trajectories.

Keywords

Island rule; flower size; island biogeography; plant evolution; Southwest Pacific

Introduction

Evolutionary size changes are common on islands. After island colonization, some organisms get bigger, such as the Moas of New Zealand and the elephant birds of Madagascar. Paradoxically, other organisms get smaller, such as the pygmy elephants of the Mediterranean islands and the hobbit of Flores (Benitez-Lopez et al., 2021).

Van Valen recognized that the directionality of body size changes was related to body size itself. He termed this phenomenon 'island rule' stating that, after island colonization, small organisms evolve to become larger while large organisms evolve to become smaller (Van Valen, 1973; Lomolino et al., 2012). Whether the island rule provides a general explanation for the evolutionary trajectories of island life has been fiercely debated for decades (Lomolino, 2005; Meiri et al., 2008; Faurby and Svenning, 2016; Itescu et al., 2018; Rebouças et al., 2018; Lokatis and Jeschke, 2018; Benitez-Lopez et al., 2021). To date, the island rule remains one of the most controversial topics in biogeography.

Most work on the island rule focuses on animals, mainly vertebrates (Meiri et al., 2008; Benitez-Lopez et al., 2021; Ciarle et al., 2023; Whittaker et al., 2023). Little is known about the island rule in plants (Burns, 2019). Studies conducted in the Southwest Pacific found evidence for the island rule in plant stature and leaf area but not in seed size (Kavanagh and Burns, 2014; Biddick et al., 2019). Similar evolutionary trajectories might be followed by other plant traits, including flower size. Differences in the size of flowers between island and mainland populations have been investigated in the past (Bernardello et al., 2001; Newstrom and Robertson, 2005; Hetherington-Rauth and Johnson, 2020). However, no previous study has tested for the island rule in flower size.

Two main evolutionary processes may cause flower size to follow the island rule. One is the reduction of pollinator diversity and size range on islands. Considering that size-coupling is an important aspect of plant-pollinator mutualism (Biddick and Burn, 2018), a

reduction in pollinators' diversity and size range could select for reduced flower diversity,

causing animal-pollinated flowers to follow the island rule (Hiraiwa and Ushimaru, 2017;

Burns, 2019). Under this hypothesis, only animal-pollinated flowers should be under

selection. Wind-pollinated flowers should not differ in size when compared to their mainland

relatives.

Alternatively, changes in flower size could be a by-product of selection acting on a

non-flower correlated trait. Since plant traits known to covary with flower size were shown to

evolve predictably on islands (Niklas, 1994; Burns et al., 2012; Biddick et al., 2019; E-

Vojtkó et al., 2022), flower size could follow the island rule simply because of its allometric

relationships. Under this hypothesis, both animal- and wind-pollinated island flowers should

differ from their mainland relatives.

Distinct plant lineages will respond differently to insular selective pressures. For

example, the genetic, morphology and evolutionary history of certain taxonomic families

makes them more likely to evolve generalized floral structures or to break down mechanisms

of self-incompatibility (Goldberg et al., 2010; Minelli, 2018; Cossard et al., 2021). Similarly,

plants with different breeding systems are typically characterized by different flowers (Krizek

and Anderson, 2013; Salisbury et al., 2017) and might respond to insular conditions in a

distinct way. Additionally, evolutionary pressures will vary depending on the abiotic and

biotic characteristics of the islands themselves (MacArthur and Wilson, 1967; Whittaker et

al., 2023). Consequently, to test for the island rule in flower size, it is vital to include multiple

plant lineages with different pollination syndromes, evolutionary histories and breeding

systems spanning multiple archipelagos.

In this study, we used a phylogenetic comparative approach to test whether island

animal-pollinated and wind-pollinated flowers follow the island rule. Specifically, using a

newly derived dataset representing 131 independent colonization events on 10 archipelagos

in the Southwest Pacific, we tried to answer the following questions: I) Do Island animal-

pollinated and wind-pollinated flowers evolve to follow the island rule? II) Do they evolve to

become larger or smaller? III) Do flowers with different pollination syndromes follow

different evolutionary trajectories?

Methods

Study area

We focused on the endemic flora of 10 archipelagos distributed across 25° of latitude (29°04'

S - 54°37' S) around New Zealand (Fig 1). The island groups can roughly be divided into the

sub-tropical archipelagos of Lord Howe, Norfolk, Kermadec, and Three Kings, and the cool-

temperate/subantarctic archipelagos of Chatham, Snares, Antipodes, Auckland, Campbell and

Macquarie. All island systems are oceanic in origin, with faunas and floras derived by

overwater dispersal from New Zealand and Australia or vicariance (Chown et al., 1998;

Rutledge et al., 2003; Pendlebury and Barnes-Keoghan, 2007; Lord, 2015; Fitzgerald, 2020).

We restricted analyses to these island systems so that our set of island taxa would be

derived from a few well-known mainland floras, sharing similar ecological, geological and

evolutionary histories. Chatham, Snares, Antipodes, Auckland, Campbell, Three Kings,

Kermadec and Macquarie were included as most of their endemic species originated from the

New Zealand mainland. Lord Howe and Norfolk islands were included as most of their

endemic species originated from Australia (Supplementary S2). We included endemics from 2

mainland sources to increase the number of taxonomic families present and to account for the

possibility that endemics originating from very different mainlands (i.e. New Zealand and

Australia) would respond differently to insular environments.

Species selection

To test whether island animal- and wind-pollinated flowers follow the island rule, we

compiled a list of island/mainland pairs of taxa. Each mainland taxon is the sister group of

the island taxon so that each pair represents a phylogenetically independent comparison.

We first compiled a list of endemic species and subspecies on each archipelago using

the most recent data available (Oliver, 1948; Godley, 1969; Taylor, 1971; Johnson and

Campbell, 1975; Meurk, 1977; Sykes, 1977; Godley, 1989; Wright, 1983; Green, 1994;

Meurk et al., 1994; Sykes and West, 1996; Amey et al., 2007; De Lange et al., 2011; Lord et

al., 2013; De Lange, 2014; De Salas and Baker, 2015). Because botanical sources varied

widely in age, all species names were cross-checked for synonymy, keeping the New Zealand

Plant Conservation Network (2023) and the Atlas of Living Australia (2023) as taxonomic

references.

We then identified the mainland sister taxon of each island taxon using published

molecular phylogenies. We searched Google Scholar for phylogenies using the species or

genus name and the Boolean operator "AND" along the term "phylogen*" (e.g. *Coprosma*

AND phylogen*). When multiple phylogenies were available, we selected the phylogeny that

used the most comprehensive genetic dataset, used either Bayesian inference or Maximum

likelihood for phylogenetic reconstruction, and showed the strongest support at each node

(supplementary material S3). Species names were standardized following the New Zealand

Plant Conservation Network (2023) and Atlas of Living Australia (2023). When molecular

phylogenies were unavailable, sister taxa were identified based on morphological evidence.

As without phylogenies we could not identify the closest relative with certainty, multiple

mainland species within the genus were selected and their floral traits averaged. If the genus

comprised less than five species on the mainland (e.g. genus *Streblus*), all species were

included as mainland relatives. If the genus comprised more than five species on the

mainland (e.g. genus *Exocarpos*), we included as mainland relatives up to five species based

on morphological affinities. (Supplementary material S3). We excluded island taxa if

mainland taxa could not be identified using either phylogeny or taxonomy.

Trait measurements

Each island/mainland pair was categorized as animal-pollinated or wind-pollinated, and as

monomorphic (including hermaphroditic, monoecious, andromonoecious, and

gynomonoecious species) or dimorphic (including dioecious and gynodioecious species)

using McGlone and Richardson (2023), Atlas of Living Australia (2023), Flora of New South

Wales Online (2023) and the New Zealand Plant Conservation Network (2023).

124   For animal-pollinated flowers, we identified the unit of pollination (i.e. the unit of

pollinator attraction, sensu Faegri and Van der Pijl, 1979) and we characterized the organs

involved in pollinator attraction (i.e. possessing an advertisement function). For radially

symmetric flowers with fused petals and petal lobes (e.g. Plantaginaceae, genus *Veronica*),

we characterized lobe length, lobe width, or corolla diameter. Tube length was collected but

not included in the analysis as it is not a measure of advertisement but of accessibility (Dafni,

1992). For radially symmetric flowers with free petals, we characterized petal length, petal

width or corolla diameter. For zygomorphic flowers, we used the length and width of the

organs most involved in advertisement (e.g. banner and wing of Fabaceae flowers). For the

Myrtaceae family, we characterized both stamen length and petal length, as in this family the

stamens are often involved in pollinator attraction (Vasconcelos et al., 2019). For Asteraceae,

we collected ray floret length, disk diameter or capitulum diameter. For dimorphic taxa, we

always used measurements from male flowers, as we expect the male perianth to be the one

under stronger selective pressures (Barbot et al., 2023) (supplementary material S2).

138   For wind-pollinated flowers, we used the sterile part of flowers. For radially

symmetric flowers with fused petals (e.g. Plantaginaceae, genus *Plantago*), we characterized

lobe length, lobe width, or corolla diameter. For radially symmetric flowers with free petals

(e.g. Arecaceae) we characterized petal length or corolla diameter. For flowers with

undifferentiated sepals and petals (e.g. Juncaceae), we used tepal length. For Poaceae, we

used glume, lemma, palea or lodicule length. For Cyperaceae, we collected glume length. For

dimorphic taxa (e.g. Rubiaceae, genus *Coprosma*) we characterized both male and female

flowers.

For the analysis, we used the same trait for all species within a family (e.g. capitulum

diameter for all Asteraceae). If missing, another flower trait was used in its place. Before

using multiple traits, a Pearson's correlation test was run to assess collinearity, setting 0.7 as a

cut-off (Berry and Feldman, 1985). Only traits above the cut-off value were considered

(Table S3).

We extracted floral traits from the Floras of New Zealand and Australia series (Allan,

1961; Moore and Edgar, 1970; Webb et al., 1988; Edgar et al., 2010) and additional online

sources (New Zealand Plant Conservation Network, 2023; Atlas of Living Australia, 2023;

and New South Wales Flora Online, 2023). We searched the primary literature when data

were missing. When online data was unavailable, we manually measured floral traits from

herbaria in the Botany collection of Te Papa Tongarewa Museum of New Zealand. For each

species trait, 5 to 10 specimens with multiple, well-preserved flowers were sampled. Three to

five measurements were then taken for each specimen and averaged.

Statistical analyses

To test whether wind-pollinated and animal-pollinated island flowers followed the island

rule, we computed a log ratio (LR) for each island/mainland pair as log(mean island flower

size/mean mainland flower size). We then conducted two separate reduced major axis

regressions. We used LR values as dependent variables and average values of mainland

relatives as independent variables. Slopes significantly lower than zero would be evidence for

the island rule. Slopes indistinguishable from zero would denote morphological isometry

(Lomolino et al., 2013). The analysis was then repeated without the LR ratio, with island

flower size as the dependent variable and mainland flower size as the independent variable.

Slopes indistinguishable from zero would denote morphological isometry. Intercepts above or

below zero would be evidence for gigantism or dwarfism respectively (Burns et al., 2012).

Mixed effect models were used to control for potential confounding factors. Because

flower size may differ between plants with different breeding strategies, the breeding system

was included as a fixed factor with two categories: monomorphic and dimorphic. Because

flowers from different mainlands may respond differently to insular systems, mainland source

was included as a fixed factor with two categories: Australia and New Zealand. Because we

expect fully differentiated species to exhibit more pronounced insular trends compared to

partially differentiated species (i.e. subspecies), taxonomic differentiation was included as a

fixed effect with two categories: partially differentiated and fully differentiated. To control

for phylogenetic conservatism, taxonomic families were included as a random effect. To

avoid singularity, only taxonomic families with three or more island/mainland pairs were

included. To control for environmental and ecological differences among islands, islands

were included as a random effect. Since flower size may differ between different flower

morphologies of animal-pollinated species, flower type was included as a fixed effect with

three categories: actinomorphic flowers with fused petals; actinomorphic flowers with free

petals, and zigomorphic flowers.

To test whether animal-pollinated and wind-pollinated island flowers were

consistently larger or smaller than their mainland relatives, we used one-sided t-tests to

determine whether mean LR values differed significantly from zero. Kruskal-Wallis analysis

of variance was used to assess significant differences in LR means among breeding systems,

mainland sources, degree of taxonomic differentiation, taxonomic families (i.e. phylogenetic

conservatism), and islands.

All analyses were conducted in the R environment version 4.3.1 (R Core Team,

2022). Reduced major axis regressions were conducted using the R package *lmodel2*

(Legendre and Oksanen, 2018). Mixed effect models were conducted using the R packages

nlme (Pinhero et al., 2017) and lme4 (Bates et al., 2009). When used as the independent

variable, mainland flower size was log-transformed to avoid heteroskedasticity. *ggplot2* was

used for data visualization (Wickham et al., 2016).

Results

We identified 155 endemic island species, representing 139 independent colonization events.

From these, we could compile island/mainland pairs for 149 species belonging to 43 families

and representing 131 phylogenetically independent events. We identified 92 animal-

pollinated pairs and 40 wind-pollinated pairs. Data for 104 pairs was gathered from online

sources, while data for the remaining 27 pairs was manually measured from herbaria

specimens (Supplementary material S2).

Animal-pollinated flowers followed the island rule, with a slope that was weakly but

significantly lower than 0 (n = -0.17, 95% confidence interval = -0.39, -0.05; p-value = 0.01)

(Fig 2a). Wind-pollinated flowers did not follow the island rule, with a slope that was not

significantly different from 0 (n = -0.01, 95% confidence interval = -0.13, 0.10; p-value =

0.81) (Fig2b). Results did not vary when island flower size was used as the dependent

variable instead of LR (Supplementary material S1). Results remained unchanged after

controlling for breeding system, mainland source, degree of taxonomic differentiation,

phylogenetic conservatism (i.e. taxonomic families), and islands (p-values = 0.02 and 0.81

respectively) (Table S1).

Animal-pollinated flowers were not consistently larger or smaller than mainland

relatives (n = 87, LR mean = -0.005, t = 0.15, p-value = 0.82). Conversely, wind-pollinated

flowers exhibited a trend toward gigantism, being generally larger than mainland relatives (n

= 39, LR mean = 0.060, t = 3.28, p-value = 0.002). For animal-pollinated flowers, the LR

mean did not vary significantly among mainland sources, taxonomic families and degrees of

taxonomic differentiation. However, it did vary among breeding systems and islands (Figure

S1, Table S3). The LR mean of wind-pollinated flowers did not vary significantly between

breeding system, mainland source, degree of taxonomic differentiation, taxonomic families,

and islands (Figure S1, Table S3).

Discussion

We provided the first test for the island rule in flowers and the first evidence of *in-situ*

evolution of flower size on islands. The size of flowers both obeyed and disobeyed the island

rule, and species with different pollination syndromes followed different evolutionary

trajectories. We found significant, albeit weak, support for the island rule in animal-

pollinated flowers. Conversely, wind-pollinated flowers did not follow the island rule but

were consistently larger on islands. This pattern was robust across breeding systems,

mainland sources, degree of taxonomic differentiation, taxonomic families and islands.

Animal-pollinated flowers obey the island rule. Perhaps this is because of reduced

pollinator diversity on islands. As pollinator diversity declines, so should the overall size

range of pollinator species (Burns, 2019). Size coupling is an important aspect of pollination

mutualisms, as smaller flowers tend to be pollinated by smaller pollinators and larger flowers

by larger pollinators (Kuriya et al., 2015; Biddick and Burn, 2018). Consequently, the

reduced size range of island pollinators may in turn reduce the flower-size diversity of

animal-pollinated island flowers.

This explanation faces two key challenges. First, we assume that pollinator size

follows the island rule, but no one has ever tested for the island rule in insect or vertebrate

pollinators in the Southwest Pacific (Burns, 2019). Secondly, plant stature and leaf area were shown to follow the island rule (Biddick et al., 2019). As flower size scales positively with these plant traits (Lambrecht and Dawson, 2007; Gomez et al., 2009; E-Vojtkó et al., 2022; Genty et al., 2023), animal-pollinated flower size may follow the island rule just as a by-product of selection acting of plant stature and/or leaf area. However, trait correlation could explain the evolution of both animal-pollinated and wind-pollinated flowers, in which case the reason why these groups evolve differently on islands becomes puzzling. Biddick et al. (2019) did not account for pollination syndrome, and it is therefore possible that animal-pollinated and wind-pollinated plant species exhibit different patterns in stature and leaf area.

Results for animal-pollinated plants varied significantly among islands and breeding systems (Figure S1). Differences among islands in animal-pollinated plants could be explained by different pollinator communities. For example, Campbell Island hosts various plants with spectacular, large inflorescences (Lord et al., 2013; Buxton et al., 2019). The nocturnal weta (*Notoplectron campbellensis*), a giant flightless cricket, is thought to play an important role in their pollination (Lord et al., 2013). The Antipodes Islands mostly lack plants with large inflorescences (Ciarle and Burns, 2023). Accordingly, they also lack an endemic species of weta (Richards, 1974).

Differences between dimorphic and monomorphic plants in animal-pollinated species were unexpected. Larger flowers in males of dimorphic animal-pollinated island plants may reflect evolution in a depauperate environment where every possibility of attracting pollinators needs to be exploited (Godley, 1979). Additionally, while females allocate more to reproduction in dimorphic animal-pollinated plants, in wind-pollinated species the reproductive costs of males may exceed or match those of females, potentially causing plants with different pollination syndromes to follow different evolutionary trajectories (Harris and Pannell, 2008; Liu et al., 2021; Roddy et al., 2021; but see Midgley, 2022).

Wind-pollinated flowers did not follow the island rule but were consistently larger on

islands. Larger wind-pollinated flowers might be a by-product of selection acting on a

correlated trait. Flower size correlates with other plant traits (Bawa et al., 2019; Roddy et al.,

2021; E-Vojtkó et al., 2022), and thus selection acting on a correlated trait could result in the

evolution of flower size (Niklas, 1994; Burns et al., 2012).

On islands, seeds consistently become larger (Kavanagh and Burns, 2014; Biddick et

al., 2019). Larger seeds might constrain the plant into producing larger fruits and flowers

(Sakai and Sakai, 1995). Conversely, larger seeds may coevolve with smaller flowers. Larger

seeds can lead to increased within-fruit sibling competition. One of the strategies adopted by

plant species to counter sibling competition is to reduce the number of seeds per fruit (Bawa,

2016). While this potentially reduces fitness, it also reduces flower and pollinator investment,

allowing plants to produce small flowers that are wind-pollinated or that rely on generalist

pollinators (Bawa et al., 2019). In other words, as seed size increases, flower size of wind-

pollinated plants might also decline, which is not what we observe. Perhaps insular wind-

pollinated plants allocate resources differently than their mainland counterparts, promoting

consistent positive flower/seed correlations. The reason behind it though remains unclear.

Our analysis encompasses only a portion of the Southwest Pacific. First, the study

does not include any tropical islands. Ecologically, tropical systems might operate differently

than non-tropical ones. For instance, while pollination networks seem to be structurally

similar in tropical and non-tropical areas (Vizentin-Bugoni et al., 2018), many plant traits

follow a latitudinal gradient, including flower morphological diversity (Currie and Paquin,

1987; Chen et al., 2017; Chartier et al., 2021). Additionally, tropical systems are

underrepresented in the literature (Culumber et al., 2019). Secondly, our study focuses on

small, isolated archipelagos with relatively small species pools. As the ecological and

evolutionary dynamics of islands vary markedly with area and isolation (MacArthur and

Wilson, 1967; Whittaker et al., 2023), these results may not apply to larger and less isolated

archipelagos.

In this study, we focused on many island/mainland pairs of species, spanning several

families. However, many of the families included in the analysis were only represented by

one or a few species, making it difficult to evaluate within family trends. Moreover, focusing

on island/mainland pairs implies that we did not account for intraspecific variance. It is

possible that accounting for such variance would alter the observed patterns.

We encourage future work to seek alternative explanations. We currently lack

alternative hypotheses to explain the *in-situ* evolution of island flowers, likely due to most

peculiarities of such flowers being viewed as assembly rules (i.e. a products of biased

colonization/establishment) rather than evolutionary processes (Lord, 2015; Pannell et al.,

2015; Burns, 2019; Hetherington-Rauth and Johnson, 2020). Finally, while our study focused

exclusively on flower size, future research should aim for a multi-trait approach capable of

encompassing the plant island syndrome in its entirety.

In conclusion, we provided the first test for the island rule in flowers, and the first

evidence of *in-situ* evolution of flower size on islands. Results indicate that, while *in situ*

evolution of flower size is widespread on islands in the Southwest Pacific, species belonging

to different pollination syndromes exhibit markedly different evolutionary trajectories. These

results combine with a growing body of literature investigating the island syndrome in plants

(Burns, 2019; Schrader et al., 2021; Delavaux et al., 2021; Meudt et al., 2021; Zizka et al.,

2022; Whittaker et al., 2023), and provide a much-needed insight into the evolution of plants

on islands.

References

Allan, H. H. (1961). *Flora of New Zealand.* New Zealand Government Printer Publications.

Amey, J., Lord, J. M., & de Lange, P. (2007). First record of a vascular plant from the Bounty Islands:

Lepidium oleraceum (nau, Cook's scurvy grass) (Brassicaceae). *New Zealand Journal of*

Botany, 45(1), 87–90.

Atlas of Living Australia. (2023). Atlas of living Australia. https://www.ala.org.au/

Barbot, E., Dufaÿ, M., & De Cauwer, I. (2023). Sex-specific selection patterns in a dioecious insect-

pollinated plant. hal-04083935.

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R. H. B., Singmann, H., Dai, B., Scheipl,

F., Grothendieck, G, Fox, J., Bauer, A., Krivitsky. P. N., Tanaka, E., & Green, P. (2009).

Package 'lme4'. *URL http://lme4. r-forge. r-project. org.*

Bawa, K. S. (2016). Kin selection and the evolution of plant reproductive traits. *Proceedings of the*

Royal Society B: Biological Sciences, *283*(1842), 20160789.

Bawa, K. S., Ingty, T., Revell, L. J., & Shivaprakash, K. N. (2019). Correlated evolution of flower size

and seed number in flowering plants (monocotyledons). *Annals of Botany*, *123*(1), 181-190.

Benítez-López, A., Santini, L., Gallego-Zamorano, J., Milá, B., Walkden, P., Huijbregts, M. A., &

Tobias, J. A. (2021). The island rule explains consistent patterns of body size evolution in

terrestrial vertebrates. *Nature Ecology & Evolution*, *5*(6), 768-786.

Bernardello, G., Anderson, G. J., Stuessy, T. F., & Crawford, D. J. (2001). A survey of floral traits,

breeding systems, floral visitors, and pollination systems of the angiosperms of the Juan

Fernández Islands (Chile). *The Botanical Review*, 67, 255-308.

Berry, W. D., & Feldman, S. (1985). Multiple regression in practice. *SAGE publications*.

Biddick, M., & Burns, K. C. (2018). Phenotypic trait matching predicts the topology of an insular plant–

bird pollination network. *Integrative Zoology*, *13*(3), 339-347.

Biddick, M., Hendriks, A., & Burns, K. C. (2019). Plants obey (and disobey) the island

rule. *Proceedings of the National Academy of Sciences*, 116(36), 17632-17634.

Burns, K. C., Herold, N., & Wallace, B. (2012). Evolutionary size changes in plants of the south-west

Pacific. *Global Ecology and Biogeography*, *21*(8), 819-828.

Burns, K. C. (2019). *Evolution in isolation: the search for an island syndrome in plants*. Cambridge

University Press.

Buxton, M. N., Anderson, B. J., Hoare, R. J., & Lord, J. M. (2019). Are moths the missing pollinators in

Subantarctic New Zealand? *Polar Research*.

Chartier, M., von Balthazar, M., Sontag, S., Löfstrand, S., Palme, T., Jabbour, F., Sauquet, H., &
347   Schönenberger, J. (2021). Global patterns and a latitudinal gradient of flower disparity:
348   perspectives from the angiosperm order Ericales. *New Phytologist*, *230*(2), 821-831.

349 Chen, S. C., Cornwell, W. K., Zhang, H. X., & Moles, A. T. (2017). Plants show more flesh in the
350   tropics: Variation in fruit type along latitudinal and climatic gradients. *Ecography*, *40*(4), 531-
351   538.

352 Chown, S. L., Gremmen, N. J. M., & Gaston, K. J. (1998). Ecological biogeography of Southern
353   Ocean islands: Species-area relationships, human impacts, and conservation. *The American*
354   *Naturalist*, 152(4), 562–575.

355 Ciarle, R., & Burns, K. C. Island biogeography of birds in the South West Pacific: Direct and indirect
356   effects of physical geography and co-occurring vegetation. *Journal of Biogeography*.

357 Ciarle, R., Burns, K. C. & **Mologni, F**. (In press.). *The loss (and gain) of defensive adaptations in*
358   *island plants and animals: A comparative review.* In X. Moreira & L. Abdala-Roberts (Ed.).
359   *Ecology and evolution of plant-herbivore interactions on islands.* Springer.

360 Cossard, G. G., Gerchen, J. F., Li, X., Cuenot, Y., & Pannell, J. R. (2021). The rapid dissolution of
361   dioecy by experimental evolution. *Current Biology*, 31(6), 1277-1283.

362 Culumber, Z. W., Anaya-Rojas, J. M., Booker, W. W., Hooks, A. P., Lange, E. C., Pluer, B., Ramírez-
363   Bullón, N., & Travis, J. (2019). Widespread biases in ecological and evolutionary
364   studies. *BioScience*, *69*(8), 631-640.

365 Currie, D. J., & Paquin, V. (1987). Large-scale biogeographical patterns of species richness of
366   trees. *Nature*, *329*(6137), 326-327.

367 Dafni, A. (1992). *Pollination ecology: a practical approach*. Oxford University Press.

368 Delavaux, C. S., Weigelt, P., Dawson, W., Essl, F., van Kleunen, M., König, C., Pergl. J., Pysek. P.,
369   Stein. A., Winter. M., Taylor. A., Schultz T. A., Whittaker. R. J., Kreft. H., & Bever, J. D.
370   (2021). Mycorrhizal types influence island biogeography of plants. *Communications*
371   *Biology*, *4*(1), 1128.

372 De Lange, P. J., Heenan, P. B., & Rolfe, J. R. (2011). *Checklist of vascular plants recorded from*
373   *Chatham Islands*. Department of Conservation, Wellington Hawke's Bay Conservancy.

374 De Lange, P. J. (2014). The flora of Egeria Rock, Northern Kermadec Island Group. *Wellington*
375   *Botanical Society Bulletin*, 55, 41–49.

De Salas, M. F., & Baker, M. L. (2015). *A census of the vascular plants of Tasmania, including*

Macquarie Island. Tasmanian Herbarium, Tasmanian Museum and Art Gallery.

E-Vojtkó, A., Junker, R. R., de Bello, F., & Götzenberger, L. (2022). Floral and reproductive traits are

an independent dimension within the plant economic spectrum of temperate central

Europe. *New Phytologist*, *236*(5), 1964-1975.

Edgar, E., Connor, H. E., Sykes, W. R., & Dawson, M. I. (2010). Flora of New Zealand: *Gramineae*.

With Contributions by W. R. Sykes and M. I. Dawson: [Illustrations by S. B. Malcolm, K. R.

West and P. A. Brooke, with Photographs by P. N. Johnson]. Manaaki Whenua Press

Faegri, K., & Van der Pijl, L. (1979). A short history of the study of pollination ecology. *The Principles*

of pollination ecology, 1-77.

Faurby, S., & Svenning, J. C. (2016). Resurrection of the island rule: human-driven extinctions have

obscured a basic evolutionary pattern. *The American Naturalist*, *187*(6), 812-820.

Fitzgerald, N. B. (2020). *Vegetation change on subantarctic Macquarie Island (Doctoral dissertation).*

University of Tasmania.

Genty, L., Kazakou, E., Metay, A., Baude, M., Gardarin, A., Michelot-Antalik, A., Leroy. A., Sotillo. A.,

Crouzet. J., & Barkaoui, K. (2023). Flowers of ruderal species are numerous but small, short

and low-rewarding. *Oikos*, 2023(12), e10219.

Godley, E. J. (1969). Additions and corrections to the flora of the Auckland and Campbell Islands.

New Zealand Journal of Botany, 7(4), 336–348.

Godley, E. J. (1979). Flower biology in New Zealand. *New Zealand journal of botany*, *17*(4), 441-466.

Godley, E. J. (1989). The flora of Antipodes Island. *New Zealand Journal of Botany*, 27(4), 531–564.

Goldberg, E. E., Kohn, J. R., Lande, R., Robertson, K. A., Smith, S. A., & Igić, B. (2010). Species

selection maintains self-incompatibility. *Science*, *330*(6003), 493-495.

Gómez, J. M., Abdelaziz, M., Muñoz-Pajares, J., & Perfectti, F. (2009). Heritability and genetic

correlation of corolla shape and size in *Erysimum mediohispanicum*. *Evolution*, *63*(7), 1820-

1831.

Green, P. S. (1994). *Flora of Australia, volume 49, Oceanic Islands 1.* Australian Government

Printing Service.

Harris, M. S., & Pannell, J. R. (2008). Roots, shoots and reproduction: sexual dimorphism in size and
405   costs of reproductive allocation in an annual herb. *Proceedings of the Royal Society B:*
406   *Biological Sciences*, *275*(1651), 2595-2602.

407   Hetherington-Rauth, M. C., & Johnson, M. T. (2020). Floral trait evolution of angiosperms on Pacific
408   Islands. *The American Naturalist*, *196*(1), 87-100.

409   Hiraiwa, M. K., & Ushimaru, A. (2017). Low functional diversity promotes niche changes in natural
410   island pollinator communities. *Proceedings of the Royal Society B: Biological*
411   *Sciences*, *284*(1846), 20162218.

412   Itescu, Y., Schwarz, R., Donihue, C. M., Slavenko, A., Roussos, S. A., Sagonas, K., Valakos. E. D.,
413   Foufopoulos, J., Pafilis, P., & Meiri, S. (2018). Inconsistent patterns of body size evolution in
414   co-occurring island reptiles. *Global Ecology and Biogeography*, *27*(5), 538-550.

415   Johnson, P. N., & Campbell, D. J. (1975). Vascular plants of the Auckland Islands. *New Zealand*
416   *Journal of Botany,* 13(4), 665–720.

417   Kavanagh, P. H., & Burns, K. C. (2014). The repeated evolution of large seeds on
418   islands. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1786), 20140675.

419   Krizek, B. A., & Anderson, J. T. (2013). Control of flower size. *Journal of experimental botany*, *64*(6),
420   1427-1437.

421   Kuriya, S., Hattori, M., Nagano, Y., & Itino, T. (2015). Altitudinal flower size variation correlates with
422   local pollinator size in a bumblebee-pollinated herb, *Prunella vulgaris* (Lamiaceae). *Journal of*
423   *evolutionary biology*, *28*(10), 1761-1769.

424   Lambrecht, S. C., & Dawson, T. E. (2007). Correlated variation of floral and leaf traits along a
425   moisture availability gradient. *Oecologia*, *151*, 574-583.

426   Legendre, P., & Oksanen, M. J. (2018). Package 'lmodel2'. *See https://CRAN. R-project.*
427   *org/package= lmodel2*.

428   Liu, M., Korpelainen, H., & Li, C. (2021). Sexual differences and sex ratios of dioecious plants under
429   stressful environments. *Journal of Plant Ecology*, *14*(5), 920-933.

430   Lokatis, S., & Jeschke, J. M. (2018). The island rule: an assessment of biases and research
431   trends. *Journal of Biogeography*, *45*(2), 289-303.

432   Lomolino, M. V. (2005). Body size evolution in insular vertebrates: generality of the island
433   rule. *Journal of biogeography*, *32*(10), 1683-1699.

Lomolino, M. V., Sax, D. F., Palombo, M. R., & van der Geer, A. A. (2012). Of mice and mammoths:

evaluations of causal explanations for body size evolution in insular mammals. *Journal of*

Biogeography, *39*(5), 842-854.

Lomolino, M. V., van der Geer, A. A., Lyras, G. A., Palombo, M. R., Sax, D. F., & Rozzi, R. (2013). Of

mice and mammoths: generality and antiquity of the island rule. *Journal of*

Biogeography, *40*(8), 1427-1439.

Lord, J. M., Huggins, L., Little, L. M., & Tomlinson, V. R. (2013). Floral biology and flower visitors on

subantarctic Campbell Island. *New Zealand Journal of Botany*, *51*(3), 168-180.

Lord, J. M. (2015). Patterns in floral traits and plant breeding systems on Southern Ocean Islands.

AoB Plants, 7, plv095.

MacArthur, R. H., & Wilson, E. O. (1967). *The theory of island biogeography*. Princeton university

press.

McGlone, M. S., & Richardson, S. J. (2023). Sexual systems in the New Zealand angiosperm

flora. *New Zealand Journal of Botany*, *61*(4), 201-231.

Meiri, S., Cooper, N., & Purvis, A. (2008). The island rule: made to be broken? *Proceedings of the*

Royal Society B: Biological Sciences, *275*(1631), 141-148.

Meudt, H. M., Albach, D. C., Tanentzap, A. J., Igea, J., Newmarch, S. C., Brandt, A. J., Lee. W. G., &

Tate, J. A. (2021). Polyploidy on islands: its emergence and importance for

diversification. *Frontiers in Plant Science*, 12, 637214.

Meurk, C. D. (1977). Alien plants in Campbell Island's changing vegetation. *Mauri Ora*, 5, 93–118.

Meurk, C. D., Foggo, M. N., & Wilson, J. B. (1994). The vegetation of subantarctic Campbell Island.

New Zealand Journal of Ecology, 18(2), 123–168.

Midgley, J. J. (2022). The costs of reproduction in plants cannot differ between the sexes. *Journal of*

Plant Ecology, *15*(6), 1308-1311.

Minelli, A. (2018). *Plant evolutionary developmental biology: the evolvability of the phenotype*.

Cambridge University Press.

Moore, L. B., & Edgar, E. (1970). Flora of New Zealand. In Flora of New Zealand. Volume 2.

Indigenous Tracheophyta. Monocotyledones except Gramineae. Manaaki Whenua Press.

Newstrom, L., & Robertson, A. (2005). Progress in understanding pollination systems in New

Zealand. *New Zealand journal of botany*, *43*(1), 1-59.

New Zealand Plant Conservation Network. (2023). New Zealand Plant Conservation Network.

https://www.nzpcn.org.nz/

NSW Flora Online. (2023). NSW Flora Online. https://plantnet.rbgsyd.nsw.gov.au/floraonline.htm

Niklas, K. J. (1994). *Plant allometry: the scaling of form and process*. University of Chicago Press.

Oliver, W. R. B. (1948). The flora of the Three Kings Islands. *Records of the Auckland Institute and*

Museum, 3(4/5), 211–238.

Pannell, J. R., Auld, J. R., Brandvain, Y., Burd, M., Busch, J. W., Cheptou, P. O., Conner. J. K.,

Goldberg. E. E., Grant. A., Grossenbacher D. L., Hovick. S. M., Igic. B., Kalisz. S., Petanidou.

T., Randle. A. M., Rubio de Casas, R., Pauw. A., Vamosi. J. C., & Winn, A. A. (2015). The

scope of Baker's law. *New Phytologist*, *208*(3), 656-667.

Pendlebury, S., & Barnes-Keoghan, I. P. (2007). Climate and climate change in the sub-Antarctic.

Papers and Proceedings of the Royal Society of Tasmania, 141(1), 67–81.

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Heisterkamp, S., Van Willigen, B., & Maintainer, R.

(2017). Package 'nlme'. *Linear and nonlinear mixed effects models, version*, *3*(1), 274.

R Core Team (2022). R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria.

URL https://www.R-project.org/.

Rebouças, R., da Silva, H. R., & Solé, M. (2018). Frog size on continental islands of the coast of Rio

de Janeiro and the generality of the Island Rule. *PLoS One*, *13*(1), e0190153.

Richards, A. M. (1974). Arthropoda of the subantarctic islands of New Zealand: 7. Orthoptera:

Rhaphidophoridae. *New Zealand Journal of Zoology*, *1*(4), 495-499.

Roddy, A. B., Martínez-Perez, C., Teixido, A. L., Cornelissen, T. G., Olson, M. E., Oliveira, R. S., &

Silveira, F. A. (2021). Towards the flower economics spectrum. *New Phytologist*, *229*(2), 665-

672.

Rutledge, D. T., Wilson, G. W., & Whenua, M. (2003). Climate parameters for New Zealand's

Offshore Islands. *Landcare Research Contract Report*.

Sakai, S., & Sakai, A. (1995). Flower size-dependent variation in seed size: theory and a test. *The*

American Naturalist, *145*(6), 918-934.

Salisbury, P. A., Fripp, Y. J., Gurung, A. M., & Williams, W. M. (2017). Is floral structure a reliable

indicator of breeding system in the Brassicaceae? *Plos one*, *12*(3), e0174176.

Schrader, J., Wright, I. J., Kreft, H., & Westoby, M. (2021). A roadmap to plant functional island

biogeography. *Biological Reviews*, *96*(6), 2851-2870.

Sykes, W. R. (1977). Kermadec Islands flora: an annotated check list. *Manaaki Whenua Press*.

Sykes, W. R., & West, C. J. (1996). New records and other information on the vascular flora of the

Kermadec Islands. *New Zealand Journal of Botany*, 34(4), 447–462.

Taylor, R. H. (1971). Influence of man on vegetation and wildlife of Enderby and Rose Islands,

Auckland Islands. *New Zealand Journal of Botany*, 9(2), 225–268.

Van Valen, L. (1973). Patterns and the balance of nature. *Evolutionary Theory*, *1*, 31-49.

Vasconcelos, T. N., Prenner, G., & Lucas, E. J. (2019). A systematic overview of the floral diversity in

Myrteae (Myrtaceae). *Systematic Botany*, *44*(3), 570-591.

Vizentin-Bugoni, J., Maruyama, P. K., de Souza, C. S., Ollerton, J., Rech, A. R., & Sazima, M. (2018).

Plant-pollinator networks in the tropics: a review. In *Ecological networks in the tropics: an*

integrative overview of species interactions from some of the most species-rich habitats on

earth, pp 73-91. Springer Nature.

Webb, C. J., Sykes, W. R., & Garnock-Jones, P. J. (1988). Flora of New Zealand. Volume IV:

Naturalised pteridophytes, gymosperms, dicotyledons. Botany Division, D.S.I.R.

Whittaker, R. J., Fernández-Palacios, J. M., & Matthews, T. J. (2023). *Island Biogeography: Geo-*

environmental Dynamics, Ecology, Evolution, Human Impact, and Conservation. Oxford

University Press.

Wickham, H., Chang, W., & Wickham, M. H. (2016). Package 'ggplot2'. *Create elegant data*

visualisations using the grammar of graphics. Version, *2*(1), 1-189.

Wright, A. E. (1983). Conservation status of the Three Kings Islands endemic flora in 1982. *Records*

of the Auckland Institute and Museum, 20, 175–184.

Zizka, A., Onstein, R. E., Rozzi, R., Weigelt, P., Kreft, H., Steinbauer, M. J., Bruelheide. H., & Lens, F.

(2022). The evolution of insular woodiness. *Proceedings of the National Academy of*

Sciences, *119*(37), e2208629119.

Acknowledgments

522   We thank Bridget Hatton for helping us navigate through the Kaitiaki Taonga Herbarium

523   Collection at Te Papa Tongarewa Museum of New Zealand. We thank Alessandra Franco for

524   drawing Figure 1.

Ethics

527   This work did not require ethical approval from a human subject or animal welfare

528   committee.

Declaration of AI use

531   We have not used AI-assisted technologies in creating this article.

Data availability Statement

534   Code, plant species lists, flower traits measured, and models used for the analysis are

535   available in the supplementary materials of this article.

Conflict of Interest Declaration

The authors declare no conflict of interest.

Figure legends

Figure 1 Map of the studied archipelagos.

Figure 2 LR (y-axis) varying as a function of log mainland values (x-axis) for a) animal-

pollinated and b) wind-pollinated flowers. Each point denotes an island/mainland pairing. A

horizontal line intercepting the y-axis at 0 denotes morphological isometry. For animal-

pollinated flowers, the slope is significantly smaller than 0, giving support for the island rule.

For wind-pollinated flowers, the slope does not vary significantly from 0, but most points are

above the horizontal line, indicating that island flowers are larger than their mainland relatives.

Fig. 1

Fig 2.