Metal accumulation sites in fruits

Kumbirai Deon Mandebere[1], Utku Deniz[1], Katarina Vogel Mikus[2,3], Cagdas Son[1],

Seckin Eroglu[1*]

1 Department of Biological Sciences, Middle East Technical University, Ankara, Turkey

2 Jozef Stefan, Jamova cesta 39, Ljubljana, Slovenia

3 Biotechnical Faculty, University of Ljubljana, Jamnikarjeva ulica 101, Ljubljana,

Slovenia

Corresponding Author: Seckin Eroglu, erogluseckin@gmail.com

Running title: Metal accumulation sites in fruits

Keywords: fruit, synchrotron, X-ray, metal, iron, MTP8, germination, tomato

Abstract

Global fruit production suffers from pre- and post- harvest losses, part of which are

related to metal deficiencies. Despite fruits being one of the most widely consumed

plant parts, the spatial distribution of metals and their possible physiological impact

remained largely unexplored. In this study, we searched for conserved metal

accumulation sites in fruits of various crops and investigated their physiological function.

By using X-ray and histochemical-based techniques, we found that calcium

accumulated in the outermost hardened tissues, potassium in fleshy tissues, and iron

(Fe) in vascular tissues. Furthermore, seeds accumulated Fe in the fruit-seed juncture,

which remained in the chalazal part of the seed upon dispersal. In *Arabidopsis thaliana*,

seed chalazal Fe formed a distinct circular shape, which appeared diffuse in *metal*

tolerance protein 8 mutants. Seed chalazal Fe appeared also in tomato seeds, albeit

more diffused and at higher concentrations. In tomato, treatment with the Fe immobilizer

desferoxamine delayed germination in wild-type plants but not in mutants with low

chalazal Fe, indicating that chalazal Fe may contribute to germination. This study marks
the first systematic application of X-ray fluorescence and histochemistry to investigate
conserved metal localization sites in mature fleshy fruits. The discovery of metal
hotspots in fruits opens new research avenues to understand fruit physiology, which
may, in the long run, contribute to food security.

Introduction

Fruits are essential to a nutritious human diet. To meet and sustain the dietary needs of
the growing global population, fruit production must increase from one billion tons to
over two billion tons per year by 2050 (Stratton et al., 2021; Willett et al., 2019). A key
challenge in achieving this goal is reducing pre- and post-harvest losses, which can
account for up to half of the production, making fruits the most wasted type of food
(Gustavsson et al., 2011; Wang et al., 2022). Metal deficiencies are significant factors
contributing to such losses, limiting the quality and quantity of fruits (Andresen et al.,
2018; Kaya and Higgs, 2002; Suzuki et al., 2003). The effects of metal deficiencies can
be either direct or indirect. As a direct effect, insufficient Ca levels in apples can result in
bitter pit disease (Song et al., 2018; Torres et al., 2021; Van Goor, 1971). As an indirect
(i.e., systemic) effect, the lack of a particular metal in sink tissues can cause metabolic
remodeling, leading to lower quantity and quality of fruits (Àlvarez-Fernàndez et al.,
2006; Álvarez-Fernández et al., 2003; El-Gioushy et al., 2021). Furthermore, metal
deficiencies may also weaken the defense of fruits against several microbial diseases
(Conway et al., 2001; Morina and Küpper, 2022).

Despite the importance of metals in fruit biology, protocols for the investigation of their
distribution in plant tissues do not always lead to the desired results. The primary
methods for exploring the spatial localization of metals in plant organs such as seeds,
roots, or leaves are X-ray-based techniques (Mijovilovich et al., 2020; Wu and Becker,
2012) and histochemical staining. However, these methods are often impractical for
fleshy fruits with high water content (Lombi et al., 2011; Yadav et al., 2021; Zhao et al.,
2014), which complicates the use of many sample preparation techniques. For example,

fixatives may not effectively replace water, the physical structure of fruits may be

undermined by water loss, and slicing of water-filled soft tissues into uniform sections

may be challenging and require specialized expertise. Consequently, spatial metal

analysis of fruits is currently limited to non-fleshy varieties such as grains or pods and,

with only a few exceptions, does not include juicy, fleshy fruits like tomatoes or oranges

(Trebolazabala et al., 2017). At present, researchers collect samples from various parts

of the fruit and compare their total metal concentrations, resulting in limited spatial

information (Kalcsits, 2016; Ramesh et al., 2021; Saquet et al., 2019). Therefore,

partially due to inefficient protocols, our understanding of how metals localize in fruits

and the physiological functions they perform remains limited.

Here, we systematically investigated metal accumulation sites in freeze-dried fruit

samples using X-ray fluorescence and histochemical staining. This approach identified

metal "hot spots", some of which are surprisingly well-conserved. We further studied

one of these metal hotspots to investigate its possible physiological functions.

Material and Methods

Research material

Fruits of various species were purchased from local markets and bazaars in Turkey

throughout the year (Table S1). For experiments with tomato seeds, if not otherwise

indicated, wild-type *Solanum lycopersicum* cv. Bonner Beste plants and its mutant

chloronerva were used (Herbik et al., 1996). All Arabidopsis lines used in this study

were derived from the Col-0 ecotype. The *mtp8-1* and *mtp8-2* mutants have been

characterized before (Eroglu et al., 2016). The SUC2::GFP line was a donation and has

been published elsewhere (Kalmbach and Helariutta, 2019).

Tabletop XRF

To prepare the fruits for XRF analysis, they were freshly sliced both longitudinally and

cross-sectionally to a thickness of 2-3 mm using a V-blade (Figure S1). The prepared

fruit slices were immediately immersed in liquid nitrogen and kept frozen at -80°C for 24

hours. Frozen samples were then placed in a vacuum freeze dryer for a main drying

phase of 23 hours at -54°C and 0.024 millibars, followed by an additional hour of final

drying at -60°C and 0.011 millibars. Once dried, fruit samples were stored at room

temperature until the analysis.

Freeze-dried fruit samples were analyzed by a micro XRF analyzer (Horiba XGT-9000,

Japan). The benchtop system was equipped with a Rh tube (50 kV, 1 mA) yielding a

polychromatic excitation beam focused to 15 μm by a polycapillary lense. X-ray

fluorescence was detected with a silicon drift diode (SDD) detector. The samples were

mapped with 100-120 μm step size and 200 ms dwell time. The spectra in each pixel

were recorded with the Horiba software, exported as .H5 files, and processed with the

PyMCA 5.8.1 toolkit (Bolte et al., 2011) where full spectra deconvolution and fitting was

performed. Maps were then generated in the RGB correlator of PyMCA on the basis of

the intensities of particular fluorescence K lines (Solé et al., 2007).

Synchrotron XRF

For examining metal enrichment in the vasculature of tomato fruits (*Solanum*

lycopersicum cv. Moneymaker) in high resolution and with minimum treatments, fresh

cut tomato pieces were embedded in the cryo-embedding matrix OCT and immediately

plunged into liquid nitrogen. Fifty μm cross sections were obtained using a Leica

RM2265 semi-automated rotary cryomicrotome and immediately transferred to a

sample holder between two Ultralene foils. The samples were kept in liquid nitrogen

during the transfer and analysis with the Scanning X-ray Microscope (SXM) at ID21 X-

Ray Microscopy Beamline at the European Synchrotron Radiation Facility (Grenoble,

France). The beam size was focused using Kirckpatrick Baez mirror optics to about

1x0.6 (HxV) μm^2. Elemental localization and distribution of Fe were performed at 7.3

keV excitations above the Fe K-edge and 100 ms dwell time. The energy was selected

using a double-crystal monochromator with a Si111 crystal pair. The X-ray fluorescence

signal from the samples was detected using a silicium drift diode detector with an 80

mm^2 active area from SGX Sensortech. The incoming beam intensity was monitored

using a photodiode. The X-ray fluorescence data from each pixel in the maps were fitted

to obtain the final elemental distribution images using the batch processing routines

from PyMCA 5.9.2.

For examining metal accumulation in the chalaza of tomato seeds in high resolution,

two-dimensional SXRF imaging was conducted. Dry seeds were placed on a Kapton

tape. Images were collected at beamline 4-BM of the National Synchrotron Light

Source-II (NSLS-II; Brookhaven National Laboratories, Upton, NY, USA). The elemental

maps were collected using an incident energy of 12 keV, a 7µm step size, and a 100 ms

dwell time. Raw data from the NSLS-II was analyzed using the Larch software package

to produce fitted X-ray fluorescence images. Final SXRF images and intensity profile

plots were generated using the Nikon NIS-elements software.

Histochemical staining of fruits

For preparing the fruit samples for histochemical staining, freeze-dried fruit pieces that

were prepared as described above for the tabletop XRF analyses were used as starting

material. Colorful fruits were subjected to decolorization. Decolorization was conducted

by immersing the pieces into a fixing solution (methanol: chloroform: glacial acetic acid;

6:3:1) in a glass Petri dish and shaking at 90 rpm until the natural color disappeared.

The fixing solution was renewed several times during the shaking. To stop the

decolorization step, the fixative was removed by washing the samples at least three

times with distilled water.

For the histochemical staining of Fe, Perls staining was used as previously described

(Roschzttardtz et al., 2009). Briefly, Perls stain solution was prepared fresh by mixing

4% K-ferrocyanide ($K_4Fe(Cn)_6$) and 4% HCl stock solutions in a 1:1 (v:v) proportion and

vacuum infiltration for one hour at 500 millibars. Next, the staining solution was

removed, and the samples were washed three times with distilled water for 1-2 min

each by slight shaking. The samples were stored in distilled water at 4°C until

observation with a stereo microscope.

For histochemical Fe staining on the thin cross sections of tomato fruits, samples

prepared for Perls staining were first embedded in resin, cut, and then stained. Pieces

were washed three times with 0.1 M phosphate buffer (pH 7.4) and dehydrated in

successive baths of 50%, 70%, 90%, 95%, and 100% ethanol, butanol/ethanol 1:1 (v/v),

and 100% butanol. Then, the pieces were embedded in Technovit 7100 resin (Kulzer,

Tokyo, Japan) according to the manufacturer's instructions and sliced into thin sections

(10 μm). The sections were deposited on glass slides and incubated for 45 min in Perls

stain solution. Perls solution was removed, slides were washed three times and directly

observed with a compound microscope.

Histochemical staining of Arabidopsis seeds

To observe whether Fe also accumulates in the seed chalaza of *Arabidopsis thaliana*,

seeds were prepared as illustrated (Fig. 5). Histochemical staining of Arabidopsis seed

parts was conducted as described above followed by DAB intensification. For DAB

intensification, the stained seed parts were incubated for 30 min in a 0.1 M phosphate

buffer (pH 7.4) solution containing 0.025% (w/v) DAB, 0.005% (v/v) H_2O_2, and 0.005%

(w/v) $CoCl_2$ (Roschzttardtz et al., 2009). The reaction was stopped by rinsing the

samples with distilled water. Images were captured with a compound microscope.

Elemental analysis by ICP-OES

To quantify Fe accumulation in the chalazal part, elemental analysis on seed parts were

conducted. To this end, mature tomato seeds (*Solanum lycopersicum* cv. Candela)

were imbibed for 24 hours and dissected into three parts as shown in Figure 2.

Dissected samples were dried in an oven at 65°C for one week. Dried plant samples

were finely ground with an electric coffee grinder. Approximately 0.3 g of the dried and

ground plant samples were weighed and placed in microwave digestion tubes. On top of

each sample, 2 ml of 30% H_2O_2 and 5 ml of HNO_3 were added, and the digestion was

conducted in a closed-vessel microwave system (MarsExpress; CEM Corp., Matthews,

NC, USA). After cooling down sufficiently, the total sample volume was finalized to 20

ml by adding double-deionized water, and samples were filtered through analytical filter

papers (Macherey-Nagel, Ø125 mm, blue band). Inductively coupled plasma optical

emission spectrometry (ICP-OES) Agilent 5800 ICP-OES (Agilent Technologies, USA),

was used to determine the concentrations of micronutrients in digested plant samples.

The accuracy of the element analyses was validated using certified standard reference

materials obtained from the National Institute of Standards and Technology

(Gaithersburg, MD, USA).

Ferric chelate reductase assay

Ferric chelate reductase activity of endosperm exudates was assessed according to a

method previously described (Grillet et al., 2014). Mature tomato seeds (*Solanum*

lycopersicum cv. Candela) were imbibed in water for 24 h at room temperature and cut

longitudinally. Embryos were discarded, and five endosperms were pooled as one

biological sample (n=3). Isolated endosperms were immersed in distilled water and

allowed to release their exudates for 24 hours. The obtained exudate solution was then

mixed (1:2 ratio) with ferric reductase activity assay solution (0.2 mM CaSO4, 5

mM MES (pH 5.5), 0.2 mM ferrozine, and 0.1 mM Fe-EDTA) on a 12-well plate in the

presence or absence of ascorbate oxidase enzyme (1.5 AOX units ml^{-1}). Following an

incubation of 3 hours, endosperms were quickly dried on tissue paper and weighed. An

800 µl aliquot was taken from the remaining solution and the absorbance was measured

at 562 nm. Experiments were repeated three times with at least three biological

replicates and the result of one representative experiment is shown.

Confocal microscopy

To observe phloem remnants in mature Arabidopsis seeds, the well-known phloem

marker SUC2::GFP was used (Kalmbach and Helariutta, 2019). Arabidopsis seeds

expressing the SUC2 construct were imbibed in distilled water for 24 hours at 4°C. After

imbibition, the seeds were dissected and chalazal+micropylar parts were excised.

These pieces were imaged using a confocal laser scanning microscope (Leica DMI4000

B; Leica Microsystems GmbH, Wetzlar, Germany) equipped with a 488-nm GFP filter.

The images were subsequently overlaid using the Fiji (ImageJ) imaging processing

package.

Germination tests

To determine how Fe availability impacts germination, germination tests were conducted with tomato (*chloronerva* and its wild type Bonner Beste) in the presence and absence of a strong Fe chelator. Seeds were sown on Petri dishes (30 seeds per dish, three replicates) filled with water agar (0.7% agar, buffered at pH 5.6 with MES) with or without desferoxamine (250 µM). Plated tomato seeds were incubated for two days in a refrigerator for stratification and then placed in a growth chamber at 25 °C under continuous light. Germination was scored over time, with a seed considered germinated when the radicle protruded through the testa. The impact of desferoxamine on germination was calculated by subtracting the IC50 values of the best-fit curves of the control from those of the desferoxamine treatment.

Results

Ca, K, and Fe accumulate in specific tissues of the fruit

By introducing an initial freeze-drying step, we determined metal localizations in fruit pieces obtained from various crop species using a bench-top X-ray machine. We opted for fruits from crop species–including tomato, okra, black grape, blueberry, lemon, aubergine, kumquat, squash, and strawberry–due to their size, which is generally larger than those of wild plants. While X-ray mapping revealed metal localizations, it provided only limited resolution. Alternatively, histochemical stains have been used to localize metals, even in cellular compartments (Eroglu et al., 2019). We found that in all fruits K accumulated in their fleshy (water rich) parts (Fig. 1, Fig. S10). We also noticed that Ca preferentially accumulates on the outer skin of the fruits. To identify accumulation sites of Fe, we stained fruit slices with Prussian Blue (also known as Perls staining). Fruits showed either homogeneous staining (Fig. S2) or a more pronounced staining throughout their vascular tissues (blue arrow heads in Fig. 2). This result shows that in fruits, Fe is preferentially stored in the vasculature in a conserved manner. The use of fixatives has been criticized due to the possible remobilization of metals, leading to artifacts and misinterpretation of the results (van der Ent et al., 2018). To address this constraint, we employed synchrotron X-ray mapping, which allowed resolutions sufficient to map vasculature in fruit tissues. Mapping of freshly cut tomato pericarp

under cryo conditions revealed a hollow Fe accumulation pattern, possibly

corresponding to vasculature (Fig. S3). While we were unable to identify Fe-

accumulating cell types, our result suggests that the preferential Fe accumulation in the

vasculature is not an artifact due to sample preparation.

Iron in the chalazal region of tomato seeds facilitates germination

Seeds of various species have been extensively investigated for metal distribution

(Eroglu et al., 2019; Ibeas et al., 2017); however, in tomato seeds, such distribution

patterns remain elusive. In the present survey, we noticed Fe accumulation in the fruit-

seed juncture of tomato slices (red arrowhead in Fig. 2). We examined whether metals

other than Fe are also enriched in this region by mapping seed metal distribution using

synchrotron X-ray fluorescence and ICP-OES. Mn partitioned towards the endosperm

and Zn towards the cells that will develop into apical and root meristems (Fig. 3). Fe

accumulated both in the embryo and the endosperm. While in the embryo, Fe

accumulation was homogeneous, in the endosperm, Fe staining was enriched in

regions close to the germination site. The chalazal endosperm (more accurately,

chalazal and micropylar endosperm, including the associated seed coat) contained

approximately twice as much Fe compared to the remainder of the seed. We further

investigated whether this Fe accumulation is dependent on nicotianamine.

Nicotianamine is a major chelator for metals including Fe in the phloem (Herbik et al.,

1996). Seeds obtained from the nicotianamine-auxotroph mutant *chloronerva* showed

lower concentrations of Fe, which, however, still accumulated preferentially in the

chalaza (Fig. S4). These results indicate that Fe in the chalaza may play a specific role

in tomato seed physiology.

In a germinating seed, endosperm-accumulated nutrients can be remobilized to support

the embryo that is transforming into a seedling (Uraguchi and Fujiwara, 2011). To

validate a possible role for Fe in feeding the germinating embryo, we examined the fate

of the Fe deposits after germination. In seeds of one-week-old seedlings, seed

endosperm tissues were degraded, reminiscent of nutrient remobilization from

endosperm into the embryo (Fig. S5). However, Perls staining showed that these

endosperm remnants preserved the preferential Fe accumulation in the chalaza (Fig.
S5, denoted "After") similar to the control (Fig. S5, denoted "Before"). Metal analysis
showed that the Fe concentration doubled. Since the cultivation media did not contain
nutrients, the observed increase in Fe concentrations was mostly due to loss of mass
through endosperm degradation. These results negate the assumption that Fe in the
chalazal nourishes the germinating embryo.
283
Next, we assessed whether Fe could aid germination by comparing the effect of an Fe
chelator on seeds with contrasting concentrations of Fe. The Fe chelator desferoxamine
has been used to counteract Fe poisoning in humans (Rafati Rahimzadeh et al., 2023)
and to render Fe stores in the plant tissues unavailable for biological processes (Aznar
et al., 2014). Targeting Fe stores in seeds with desferoxamine led to a delay of
approximately 6 hours (Fig. 4). However, the germination of the *chloronerva* mutant,
which contains low concentrations of Fe, was not affected by desferoxamine,
suggesting that Fe accumulation in the seed controls germination speed in tomato.
292
Germination speed is determined by the amount of hydroxyl radicals produced in the
endosperm (Muller et al., 2009; Schweikert et al., 2002). A common pathway in nature
for hydroxyl radical production is through the Fenton chain reaction, which requires Fe,
hydrogen peroxide, and ascorbate (Burkitt and Gilbert, 1990a). Hydrogen peroxide
production in tomato seeds during imbibition was shown before (Morohashi, 2002). We
found that tomato endosperm also produces ascorbate during imbibition (Fig. S6).
Taken together, it can be speculated that Fe aids germination by weakening the
endosperm through the production of hydroxyl radicals.
301
Chalazal Fe accumulation in Arabidopsis seeds follows remnants of vascular
tissues and is dependent on MTP8
Next, we were interested in finding genetic components that mediate chalazal Fe
accumulation. For this purpose, we employed the model plant Arabidopsis to profit from
its genomic resources. We applied Perls staining with DAB amplification (Perls/DAB) to
the cut seeds as depicted in Fig. 5A and investigated the black coloration representing

Fe accumulation. Interestingly, Perls/DAB staining revealed Fe accumulation in the

chalaza (Fig. 5B, top). In Arabidopsis, a chalazal Fe accumulation pattern has not been

reported until now (Eroglu, 2018; Eroglu et al., 2017; Kim et al., 2006). However, unlike

in tomato, this Fe accumulation was not evenly distributed but produced a dashed ring-

like shape (Fig. 5B, bottom). We then screened Arabidopsis metal transporter mutants

for disrupted chalazal Fe accumulation. Two independent alleles of *metal-tolerance-*

protein 8 (*mtp8*) knock-out lines showed a disrupted Fe accumulation pattern. MTP8

has been previously characterized as an Fe and Mn transporter that mediates the

loading of Fe and Mn into the embryo (Eroglu et al., 2017). Our staining showed that in

almost all *mtp8* seeds, the typical sharp Fe localization appeared diffused (Fig. 5C).

Furthermore, in our hands, 11 out of 20 *mtp8* mutant plants showed lighter tones of

black (brownish), and 3 out of 20 *mtp8* mutant plants showed no staining at all (i.e., the

Fe concentration was below the detection limit). This result indicates that preferential Fe

accumulation in the Arabidopsis chalaza is MTP8 dependent. Next, we applied

desferoxamine but found no decrease in germination speed of either wild-type or *mtp8*

Arabidopsis seeds (data not shown).

Then, we investigated whether the ring-like Fe accumulation corresponds to any known

structure in the Arabidopsis seed. A similar structure has recently been identified as the

phloem end of the vasculature of the funiculus (Liu et al., 2025). However, this

observation was made on developing seeds. There have been no reports on vascular

tissue remaining in the mature seeds of Arabidopsis. Due to the similarities between the

shapes (i.e., ring-like) and localizations (i.e., at the end of the chalaza), we hypothesize

that the Fe-accumulating ring corresponds to the remaining vasculature. To test this

hypothesis, we examined the GFP localization in the imbibed mature seeds of

SUC2:GFP, a marker used to reveal the symplastic pathway of the phloem (Kalmbach

and Helariutta, 2019; Otero and Helariutta, 2016). GFP localized exclusively to the end

of the chalaza of the seed, producing a ring-like shape reminiscent of the Fe

accumulation (Fig. 5D). This result indicates that the Fe accumulation in the chalaza

may follow the vasculature remaining in the seed.

With this new finding, we revisited the unique Fe localization around the squash seeds

(Fig. 2F). In most angiosperms, the funiculus attaches to the seed only in the chalaza;

however, in the Cucurbitaceae family, a single vasculature extends throughout the outer

integument of the seed coat from chalaza to the micropyle in a median plane (Johri et

al., 2013) from the rear and the top (Fig. S8A, B). Fe staining patterns matched the

expected vascular bundles (Fig. S8C, D). These data indicate that, at least in

Arabidopsis and squash, Fe followed vascular tissues in the seeds besides the fruits.

Discussion

X-ray and histochemistry-based methods highlighted numerous metal hotspots in

various plant organs (Eroglu et al., 2019, 2016; Kim et al., 2006). In this study, we

examined metal localizations in fruits. Our results show that in fruits, metals including

Ca, K, and Fe accumulate in a tissue-specific way. We further show that in Arabidopsis,

Fe accumulation in the fruit-seed juncture is mediated by MTP8, possibly determining

germination speed in tomato.

Histochemical staining of freeze-dried and decolorized fruits provides tissue-level

information

The histochemical approach to mature fleshy fruits often faces two main obstacles: 1)

too much water, limiting the penetration and activity of chemicals, and 2) too many

pigments, limiting the visual inspection of the stain. Industrially, fruits have been

extensively dried for various purposes. Among drying technologies, vacuum freeze

drying appears to be the golden standard for human consumption (Harper and Tappel,

1957). We observed that once freed from water, the fruit slices behaved like regular

tissue. After vacuum freeze drying, fixatives could decolorize the fruit. This may

represent an advance over the techniques described in the literature. The most

common way to circumvent the natural bright colors of fruits like tomato, has been to

analyze the sample in earlier developmental stages (Saed Taha et al., 2012; Wang et

al., 2014). We propose here that freeze drying, coupled with histochemistry, provides an

efficient and quick method to reveal immobile metal hotspots in fruits. Any finding can

then be later confirmed with more advanced methods that offer higher resolution and

quantification and avoid possible metal re-localization.

Iron follows the vasculature of fruits and seeds

Fe preferentially accumulates in the vascular tissues of several plant parts. Leaves (Xu

et al., 2022), embryos inside the seeds (Kim et al., 2006), and fruits (this study)

accumulate Fe in and around their provascular or vascular strands. In mature fruits,

vascular Fe may represent the most prominent Fe accumulation site (Fig. 3). However,

this accumulation can appear more pronounced than it is, due to clearing off weakly

bound Fe in the rest of the fruit during fixation and decolorization. The physiological

function of Fe in the fruit vasculature is unclear. Fe deficiency in fruit trees has not been

linked with a disease that is specific to fruit vasculature (Àlvarez-Fernàndez et al.,

2006). Alternatively, an immobile Fe reservoir may be a "byproduct" of extensive callose

production. Callose is required to close the xylem connection in fruits (Knipfer et al.,

2015) and may cause the oxidation of a large amount of Fe for its production (Müller et

al., 2015). Another possibility is that the main vasculatures of fruits serve as a sink to

load Fe into the seeds. In this model, the vasculature may represent a reservoir of

excess Fe left over from seed Fe filling, as this process is highly regulated to avoid Fe-

mediated oxidative damage in the seeds (Sun et al., 2021). The remaining Fe can then

be bound to cell walls that can quickly absorb Fe(III) (Xu et al., 2022). The current study

identified the vasculature as the main Fe-accumulating tissue in fruits; however, further

investigations to identify the cells in the vasculature that accumulate Fe and into the

underlying mechanisms of this process are required to substantiate this supposition.

Iron in the seed chalaza facilitates germination

A fraction of Fe-loaded vasculature retains in the seeds upon dispersal from the fruit

(Fig. 3, 5 and Liu et al., 2023). This Fe reservoir may facilitate germination (Fig. 4).

Germination requires softening of the endosperm to allow radical protrusion, a major

process that includes cell wall-cutting hydroxyl radicals (Grillet et al., 2013; Muller et al.,

2009). A well-known source of hydroxyl radicals derives from the reaction of Fe with

H_2O_2, referred to as Fenton reaction (Burkitt and Gilbert, 1990b). Therefore, Fe deposits

in the chalaza may facilitate germination by generating cell wall-attacking hydroxyl

radicals. However, for such a function one would expect Fe to accumulate more

towards the micropylar region where the root protrudes. Fe-enriched chalazae are

preserved between tomato and Arabidopsis (Fig. 5), but only in tomato perturbation of

the Fe deposit by desferoxamine postponed the germination (Fig. 6). This observation

implies that the phenotype may depend on the Fe amount in the chalaza. In line with

this assumption, tomato seeds that accumulate less Fe (Fig. S4 and Fig. 6) or seeds of

species (e.g., Arabidopsis) that accumulate lower concentrations of Fe did not show a

clear germination phenotype when treated with desferoxamine. Further research should

test whether Fe-mobilizing chelators such as nicotianamine increase germination speed

and whether the germination of Arabidopsis Fe over-acccumulating mutants such as

brutus-3 (Hindt et al., 2017) is postponed by desferoxamine.

Micro-XRF mapping is applicable to investigate nutrient deficiency-related fruit diseases

Micro XRF mapping has been successfully used to examine metal localizations in

various plant organs, including leaves, roots, stems, and seeds (Singh et al., 2022). We

detected Ca to be preferentially accumulating in the outer part of fruits, especially in the

citrus genus (Fig. 1, S10). This is in agreement with previous reports where the highest

Ca levels were generally observed in the peel of the fruits (Saure, 2005). Higher Ca in

the outer parts of the fruits is probably caused by cell wall thickening. Ca is a major

component of the cell wall (Thor, 2019) and outer parts of the fruits usually possess

thicker cell walls to maintain integrity (Forlani et al., 2019; Hallett and Sutherland, 2005;

Jeffery et al., 2012). When fruits contain insufficient Ca, they may be more prone to

infections, as Ca deficiency was reported to cause several diseases, including bitter pit

in apples and blossom-end rot in tomatoes, peppers, and watermelons (de Freitas and

Mitcham, 2012). Potassium preferentially accumulates in the fleshy parts of the fruits in

a conserved manner (Fig. 1 and Ahmed et al., 2024). This is not surprising since K is

involved in osmoregulation and sugar accumulation, whereas the fleshy parts of the

fruits often accumulate water and sugar (Kumar et al., 2006).

Imaging element distributions of fruits is associated with certain limitations.
Polychromatic excitation with an X-ray tube (Rh in our case) gave a high background
due to scattering in the energy range of middle z-elements, decreasing the detection
limit for trace elements like Fe, Cu and Zn. We found that in fruits, Fe was under the
detection limit for imaging despite the presence of an Fe peak (Fig. S9). Nevertheless,
our data showed that micro-XRF can be very useful to image distribution of macro-
elements like K and Ca, which can aid in investigating whether specific disease
symptoms overlap with perturbations in metal accumulating sites.
438

Conclusions

Increasing fruit production should target decreasing pre- and post-harvest losses. To
reduce such losses, a better understanding of fruit biology is required. This study
extends the basic techniques to localize metals to mature fleshy fruits, contributing to
preventing yield loss in fruit production and storage in the long run. Fruits preferentially
accumulate Fe in their vascular, K in their fleshy and Ca in their epidermal tissues.
Fruits load Fe into the chalazal part of the seeds, which, in *A. thaliana*, is mediated by
MTP8. Tomato seed chalaza accumulates a large amount of Fe, which may determine
germination speed.
448

Author contributions

450 S.E. designed the study. U.D. and D.M. prepared the fruits and conducted the Perls
451 staining. D.M. conducted the remaining experiments. K.V.M conducted benchtop X-Ray
452 analyses. C.S. conducted confocal microscopy analysis. All authors discussed the data.
453 S.E. wrote the manuscript with contributions from the other authors. All authors read
454 and approved the final version of the manuscript.
455
456

Acknowledgments

We thank Mary Lou Guerinot and Ryan Tappero for synchrotron imaging at X-ray Fluorescence Microprobe (XFM) beamline 4-BM of the National Synchrotron Light Source II, a US Department of Energy (DOE) Office of Science User Facility operated for the DOE Office of Science by Brookhaven National Laboratory under Contract DE-SC0012704. We acknowledge the European Synchrotron Radiation Facility (ESRF) for provision of synchrotron radiation facilities under proposal number LS-3343 for beamline, and we would like to thank Hiram Castillo for assistance and support in using beamline ID21. We also acknowledge the SIMBION (Eurobioimaging) network for benchtop micro-XRF analyses and ARIS for program financing (P1-0212, Plant Biology). We further thank Yka Helariutta for providing SUC2::GFP lines, Petra Bauer for *chloronerva* mutants, Abdulsamet Sakalar for helping with the histochemical staining of fruits, Catherine Curie for help with the germination experiments, and Ahmet Bakirbas for critical reading of the manuscript. This study was supported by the Scientific and Technological Research Council of Turkiye (TUBITAK) under Grant Number 122Z109; European Union under COST action CA19116, Trace metal metabolism in plants (PLANTMETALS), and TENMAK, Ankara, Turkey.

References

Ahmed, I.A.M., Özcan, M.M., AlJuhaimi, F., Albakry, Z., 2024. The Monitoring of Accumulations of Elements in Apple, Pear, and Quince Fruit Parts. Biol Trace Elem Res. https://doi.org/10.1007/s12011-024-04223-3

Àlvarez-Fernàndez, A., Abadía, J., Abadía, A., 2006. Iron deficiency, fruit yield and fruit quality. Iron nutrition in plants and rhizospheric microorganisms 85–101.

Álvarez-Fernández, A., Paniagua, P., Abadía, J., Abadía, A., 2003. Effects of Fe Deficiency Chlorosis on Yield and Fruit Quality in Peach (*Prunus persica* L. Batsch). J. Agric. Food Chem. 51, 5738–5744. https://doi.org/10.1021/jf034402c

Andresen, E., Peiter, E., Küpper, H., 2018. Trace metal metabolism in plants. Journal of experimental botany 69, 909–954.

Aznar, A., Chen, N.W.G., Rigault, M., Riache, N., Joseph, D., Desmaële, D., Mouille, G., Boutet, S., Soubigou-Taconnat, L., Renou, J.-P., Thomine, S., Expert, D., Dellagi, A., 2014. Scavenging Iron: A Novel Mechanism of Plant Immunity Activation by Microbial Siderophores. Plant Physiology 164, 2167–2183. https://doi.org/10.1104/pp.113.233585

Bolte, S., Lanquar, V., Soler, M.-N., Beebo, A., Satiat-Jeunemaître, B., Bouhidel, K., Thomine, S., 2011. Distinct Lytic Vacuolar Compartments are Embedded Inside the Protein Storage Vacuole of Dry and Germinating Arabidopsis thaliana Seeds. Plant Cell Physiol 52, 1142–1152. https://doi.org/10.1093/pcp/pcr065

Burkitt, M.J., Gilbert, B.C., 1990a. Model Studies of the Iron-Catalysed Haber-Weiss Cycle and the Ascorbate-Driven Fenton Reaction. Free Radical Research Communications 10, 265–280. https://doi.org/10.3109/10715769009149895

Burkitt, M.J., Gilbert, B.C., 1990b. Model studies of the iron-catalysed Haber-Weiss cycle and the ascorbate-driven Fenton reaction. Free radical research communications 10, 265–280.

Conway, W.S., Sams, C.E., Hickey, K.D., 2001. Pre-and postharvest Ca treatment of apple fruit and its effect on quality, in: International Symposium on Foliar Nutrition of Perennial Fruit Plants 594. pp. 413–419.

de Freitas, S.T., Mitcham, E.J., 2012. Factors Involved in Fruit Ca Deficiency Disorders, in: Horticultural Reviews. John Wiley & Sons, Ltd, pp. 107–146. https://doi.org/10.1002/9781118351871.ch3

El-Gioushy, S.F., Ding, Z., Bahloul, A.M., Gawish, M.S., Abou El Ghit, H.M., Abdelaziz, A.M., El-Desouky, H.S., Sami, R., Khojah, E., Hashim, T.A., 2021. Foliar application of nano, chelated, and conventional iron forms enhanced growth, nutritional status, fruiting aspects, and fruit quality of Washington navel orange trees (Citrus sinensis L. Osbeck). Plants 10, 2577.

Eroglu, S., 2018. Chapter Four - Metal Transport in the Developing Plant Seed, in: Maurel, C. (Ed.), Advances in Botanical Research, Membrane Transport in Plants. Academic Press, pp. 91–113. https://doi.org/10.1016/bs.abr.2018.09.010

Eroglu, S., Giehl, R.F.H., Meier, B., Takahashi, M., Terada, Y., Ignatyev, K., Andresen, E., Küpper, H., Peiter, E., von Wirén, N., 2017. Metal Tolerance Protein 8 Mediates Manganese Homeostasis and Iron Reallocation during Seed Development and Germination. Plant physiology 174, 1633–1647. https://doi.org/10.1104/pp.16.01646

Eroglu, S., Karaca, N., Vogel-Mikus, K., Kavčič, A., Filiz, E., Tanyolac, B., 2019. The conservation of VIT1-dependent iron distribution in seeds. Frontiers in plant science 10, 907.

Eroglu, S., Meier, B., von Wirén, N., Peiter, E., 2016. The vacuolar manganese transporter MTP8 determines tolerance to iron deficiency-induced chlorosis in Arabidopsis. Plant physiology 170, 1030–1045. https://doi.org/10.1104/pp.15.01194

Forlani, S., Masiero, S., Mizzotti, C., 2019. Fruit ripening: the role of hormones, cell wall modifications, and their relationship with pathogens. Journal of Experimental Botany 70, 2993–3006. https://doi.org/10.1093/jxb/erz112

Grillet, L., Mari, S., Schmidt, W., 2013. Iron in seeds–loading pathways and subcellular localization. Frontiers in plant science 4.

Grillet, L., Ouerdane, L., Flis, P., Hoang, M.T.T., Isaure, M.-P., Lobinski, R., Curie, C., Mari, S., 2014. Ascorbate efflux as a new strategy for iron reduction and transport in plants. Journal of Biological Chemistry 289, 2515–2525.

Gustavsson, J., Cederberg, C., Sonesson, U., Van Otterdijk, R., Meybeck, A., 2011. Global food losses and food waste.

Hallett, I.C., Sutherland, P.W., 2005. Structure and Development of Kiwifruit Skins. International Journal of Plant Sciences 166, 693–704. https://doi.org/10.1086/431232

Harper, J.C., Tappel, A.L., 1957. Freeze-drying of food products. Advances in Food Research 7, 171–234.

Herbik, A., Giritch, A., Horstmann, C., Becker, R., Balzer, H.-J., Baumlein, H., Stephan, U.W., 1996. Iron and copper nutrition-dependent changes in protein expression in a tomato wild type and the nicotianamine-free mutant chloronerva. Plant Physiology 111, 533–540.

Hindt, M.N., Akmakjian, G.Z., Pivarski, K.L., Punshon, T., Baxter, I., Salt, D.E., Guerinot, M.L., 2017. BRUTUS and its paralogs, BTS LIKE1 and BTS LIKE2, encode important negative regulators of the iron deficiency response in Arabidopsis thaliana. Metallomics 9, 876–890. https://doi.org/10.1039/C7MT00152E

Ibeas, M.A., Grant-Grant, S., Navarro, N., Perez, M.F., Roschzttardtz, H., 2017. Dynamic Subcellular Localization of Iron during Embryo Development in Brassicaceae Seeds. Front. Plant Sci. 8, 1–8. https://doi.org/10.3389/fpls.2017.02186

Jeffery, J., Holzenburg, A., King, S., 2012. Physical barriers to carotenoid bioaccessibility. Ultrastructure survey of chromoplast and cell wall morphology in nine carotenoid-containing fruits and vegetables. Journal of the Science of Food and Agriculture 92, 2594–2602. https://doi.org/10.1002/jsfa.5767

Johri, B.M., Ambegaokar, K.B., Srivastava, P.S., 2013. Comparative embryology of angiosperms vol. 1/2. Springer Science & Business Media.

Kalcsits, L.A., 2016. Non-destructive measurement of Ca and P in apple and pear using handheld X-ray fluorescence. Frontiers in plant science 7, 442.

Kalmbach, L., Helariutta, Y., 2019. Sieve Plate Pores in the Phloem and the Unknowns of Their Formation. Plants 8, 25. https://doi.org/10.3390/plants8020025

Kaya, C., Higgs, D., 2002. Response of tomato (*Lycopersicon esculentum* L.) cultivars to foliar application of zinc when grown in sand culture at low zinc. Scientia Horticulturae 93, 53–64. https://doi.org/10.1016/S0304-4238(01)00310-7

Kim, S.A., Punshon, T., Lanzirotti, A., Li, L., Alonso, J.M., Ecker, J.R., Kaplan, J., Guerinot, M.L., 2006. Localization of iron in Arabidopsis seed requires the vacuolar membrane transporter VIT1. Science 314, 1295–1298.

Knipfer, T., Fei, J., Gambetta, G.A., McElrone, A.J., Shackel, K.A., Matthews, M.A., 2015. Water transport properties of the grape pedicel during fruit development: insights into xylem anatomy and function using microtomography. Plant Physiology 168, 1590–1602.

Kumar, A.R., Kumar, N., Kavino, M., 2006. Role of P in fruit crops-a review. Agricultural Reviews 27, 284–291.

Liu, X., Nakajima, K.P., Adhikari, P.B., Wu, X., Zhu, S., Okada, K., Kagenishi, T., Kurotani, K., Ishida, T., Nakamura, M., 2025. Fertilization-dependent phloem end gate regulates seed size. Current Biology.

Liu, X., Nakajima, K.P., Wu, X., Zhu, S., Adhikari, P.B., Okada, K., Kurotani, K., Ishida, T., Nakamura, M., Sato, Y., 2023. Seed size control via phloem end by callose deposition/degradation of β-1, 3-glucanase. bioRxiv 2023–07.

Lombi, E., Scheckel, K.G., Kempson, I.M., 2011. In situ analysis of metal(loid)s in plants: State of the art and artefacts. Environmental and Experimental Botany, Metal(loid) tolerance in plants and lichens 72, 3–17. https://doi.org/10.1016/j.envexpbot.2010.04.005

Mijovilovich, A., Morina, F., Bokhari, S.N., Wolff, T., Küpper, H., 2020. Analysis of trace metal distribution in plants with lab-based microscopic X-ray fluorescence imaging. Plant Methods 16, 82. https://doi.org/10.1186/s13007-020-00621-5

Morina, F., Küpper, H., 2022. Trace metals at the frontline of pathogen defence responses in non-hyperaccumulating plants. Journal of Experimental Botany 73, 6516–6524. https://doi.org/10.1093/jxb/erac316

Morohashi, Y., 2002. Peroxidase activity develops in the micropylar endosperm of tomato seeds prior to radicle protrusion. Journal of Experimental Botany 53, 1643–1650.

Müller, J., Toev, T., Heisters, M., Teller, J., Moore, K.L., Hause, G., Dinesh, D.C., Bürstenbinder, K., Abel, S., 2015. Iron-dependent callose deposition adjusts root meristem maintenance to phosphate availability. Developmental cell 33, 216–230.

Muller, K., Linkies, A., Vreeburg, R.A., Fry, S.C., Krieger-Liszkay, A., Leubner-Metzger, G., 2009. In vivo cell wall loosening by hydroxyl radicals during cress seed germination and elongation growth. Plant physiology 150, 1855–1865.

Otero, S., Helariutta, Y., 2016. Companion cells: a diamond in the rough. Journal of experimental botany erw392.

Rafati Rahimzadeh, Mehrdad, Rafati Rahimzadeh, Mehravar, Kazemi, S., Moghadamnia, A.R., Ghaemi Amiri, M., Moghadamnia, A.A., 2023. Iron; Benefits

or threatens (with emphasis on mechanism and treatment of its poisoning). Hum Exp Toxicol 42, 09603271231192361. https://doi.org/10.1177/09603271231192361

Ramesh, K.V., Paul, V., Pandey, R., 2021. Dynamics of mineral nutrients in tomato (Solanum lycopersicum L.) fruits during ripening: part I—on the plant. Plant Physiol. Rep. 26, 23–37. https://doi.org/10.1007/s40502-020-00546-0

Roschzttardtz, H., Conéjéro, G., Curie, C., Mari, S., 2009. Identification of the Endodermal Vacuole as the Iron Storage Compartment in the Arabidopsis Embryo. Plant Physiol. 151, 1329–1338. https://doi.org/10.1104/pp.109.144444

Saed Taha, R., Ismail, I., Zainal, Z., Abdullah, S.N.A., 2012. The stearoyl-acyl-carrier-protein desaturase promoter (Des) from oil palm confers fruit-specific GUS expression in transgenic tomato. Journal of Plant Physiology 169, 1290–1300. https://doi.org/10.1016/j.jplph.2012.05.001

Saquet, A.A., Streif, J., Almeida, D.P., 2019. Mineral composition and distribution within 'Rocha'pear in relation to internal storage disorders. Postharvest Biology and Technology 158, 111002.

Saure, M.C., 2005. Ca translocation to fleshy fruit: its mechanism and endogenous control. Scientia Horticulturae 105, 65–89. https://doi.org/10.1016/j.scienta.2004.10.003

Schweikert, C., Liszkay, A., Schopfer, P., 2002. Polysaccharide degradation by Fenton reaction- or peroxidase-generated hydroxyl radicals in isolated plant cell walls. Phytochemistry 61, 31–35. https://doi.org/10.1016/S0031-9422(02)00183-8

Singh, Vivek K., Sharma, N., Singh, Virendra K., 2022. Application of X-ray fluorescence spectrometry in plant science: Solutions, threats, and opportunities. X-Ray Spectrometry 51, 304–327. https://doi.org/10.1002/xrs.3260

Solé, V.A., Papillon, E., Cotte, M., Walter, P., Susini, J., 2007. A multiplatform code for the analysis of energy-dispersive X-ray fluorescence spectra. Spectrochimica Acta Part B: Atomic Spectroscopy 62, 63–68.

Song, W., Yi, J., Kurniadinata, O.F., Wang, H., Huang, X., 2018. Linking fruit Ca uptake capacity to fruit growth and pedicel anatomy, a cross-species study. Frontiers in plant science 9, 575.

Stratton, A.E., Finley, J.W., Gustafson, D.I., Mitcham, E.J., Myers, S.S., Naylor, R.L., Otten, J.J., Palm, C.A., 2021. Mitigating sustainability tradeoffs as global fruit and vegetable systems expand to meet dietary recommendations. Environ. Res. Lett. 16, 055010. https://doi.org/10.1088/1748-9326/abe25a

Sun, L., Wei, Y.Q., Wu, K.H., Yan, J.Y., Xu, J.N., Wu, Y.R., Li, G.X., Xu, J.M., Harberd, N.P., Ding, Z.J., 2021. Restriction of iron loading into developing seeds by a YABBY transcription factor safeguards successful reproduction in Arabidopsis. Molecular Plant 14, 1624–1639.

Suzuki, K., Shono, M., Egawa, Y., 2003. Localization of Ca in the pericarp cells of tomato fruits during the development of blossom-end rot. Protoplasma 222, 149–156. https://doi.org/10.1007/s00709-003-0018-2

Thor, K., 2019. Ca—Nutrient and Messenger. Frontiers in Plant Science 10.

Torres, E., Alegre, S., Recasens, I., Asín, L., Lordan, J., 2021. Integral procedure to predict bitter pit in 'Golden Smoothee'apples based on Ca content and symptom induction. Scientia Horticulturae 277, 109829.

Trebolazabala, J., Maguregui, M., Morillas, H., de Diego, A., Madariaga, J.M., 2017. Evaluation of metals distribution in Solanum lycopersicum plants located in a coastal environment using micro-energy dispersive X-ray fluorescence imaging. Microchemical Journal 131, 137–144.

Uraguchi, S., Fujiwara, T., 2011. Significant contribution of boron stored in seeds to initial growth of rice seedlings. Plant Soil 340, 435–442. https://doi.org/10.1007/s11104-010-0614-9

van der Ent, A., Przybyłowicz, W.J., de Jonge, M.D., Harris, H.H., Ryan, C.G., Tylko, G., Paterson, D.J., Barnabas, A.D., Kopittke, P.M., Mesjasz-Przybyłowicz, J., 2018. X-ray elemental mapping techniques for elucidating the ecophysiology of hyperaccumulator plants. New Phytologist 218, 432–452. https://doi.org/10.1111/nph.14810

Van Goor, B.J., 1971. The Effect of Frequent Spraying with Ca Nitrate Solutions on the Mineral Composition and the Occurrence of Bitter Pit of the Apple Cox's Orange Pippin. Journal of Horticultural Science 46, 347–364. https://doi.org/10.1080/00221589.1971.11514415

Wang, M., Xu, Y., Yang, Y., Mu, B., Nikitina, M.A., Xiao, X., 2022. Vis/NIR optical biosensors applications for fruit monitoring. Biosensors and Bioelectronics: X 11, 100197. https://doi.org/10.1016/j.biosx.2022.100197

Wang, S., Lu, G., Hou, Z., Luo, Z., Wang, T., Li, H., Zhang, J., Ye, Z., 2014. Members of the tomato FRUITFULL MADS-box family regulate style abscission and fruit ripening. Journal of Experimental Botany 65, 3005–3014. https://doi.org/10.1093/jxb/eru137

Willett, W., Rockström, J., Loken, B., Springmann, M., Lang, T., Vermeulen, S., Garnett, T., Tilman, D., DeClerck, F., Wood, A., 2019. Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. The lancet 393, 447–492.

Wu, B., Becker, J.S., 2012. Imaging techniques for elements and element species in plant science. Metallomics 4, 403. https://doi.org/10.1039/c2mt00002d

Xu, Z.-R., Cai, M.-L., Yang, Y., You, T.-T., Ma, J.F., Wang, P., Zhao, F.-J., 2022. The ferroxidases LPR1 and LPR2 control iron translocation in the xylem of Arabidopsis plants. Molecular Plant 15, 1962–1975.

Yadav, V., Arif, N., Singh, V.P., Guerriero, G., Berni, R., Shinde, S., Raturi, G., Deshmukh, R., Sandalio, L.M., Chauhan, D.K., Tripathi, D.K., 2021. Histochemical Techniques in Plant Science: More Than Meets the Eye. Plant and Cell Physiology 62, 1509–1527. https://doi.org/10.1093/pcp/pcab022

Zhao, F.-J., Moore, K.L., Lombi, E., Zhu, Y.-G., 2014. Imaging element distribution and speciation in plant cells. Trends in Plant Science 19, 183–192. https://doi.org/10.1016/j.tplants.2013.12.001

Figures and captions

Figure 1: Calcium accumulates in epidermal and K in fleshy parenchyma tissues of fruits. Freeze-dried fruit slices were examined with a bench-top micro-XRF. Left, kumquat; middle, lemon; right, squash. Red rectangles denote the analyzed area. At least two independent pieces were observed for each fruit with comparable results. Scale bar is 5 mm. K: Potassium, Ca: Ca.

Figure 2: Iron accumulates in the fruit vasculature of diverse species and in the fruit-seed juncture in tomato. Cross-sectional or longitudinal cut fruit pieces were freeze-dried and decolorized with fixatives (methanol: chloroform: glacial acetic acid; 6:3:1). After fixation, the pieces were submerged in Perls staining solution for 2 to 16 hours. Blue coloration indicates Fe accumulation, blue arrowheads highlight vasculature stained blue in different fruit species. For this figure, images of fruits with noticeable Fe accumulation in the vascular bundles were chosen. Note that tomato accumulated Fe in the fruit seed juncture, shown by a red arrowhead. A-K; eggplant, cucumber, kiwi, kumquat, haricot, squash, green almond, blueberry, papaya, tomato, and passion fruit. At least five pieces of fruits were investigated, and a representative sample was photographed. Bar scale is 1 cm, unless otherwise indicated.

Figure 3: Metal loading to the fruit-seed juncture is specific to Fe and remains after seed dispersal. A, metal distribution in dry mature tomato seeds. Seeds were placed on Kapton tape and metal distribution was examined by synchrotron XRF. B, metal distribution in dissected seeds parts. Seeds were imbibed for one day and dissected to isolate embryos and lateral or chalazal endosperm. Seed parts were dried, and metal concentrations were determined using ICP-OES.

Figure 4: Iron loading in the fruit-seed juncture may facilitate germination. Left, desferoxamine, a strong chelator of Fe, decreases germination speed. Right, the impact of desferoxamine on germination is abolished in *chloronerva* mutants that contain low Fe concentrations. Tomato seeds were sown on water agar without (Control, circles) or with desferoxamine supplementation (Des, squares), and the germination percentage was calculated over time. The experiment was repeated four times with comparable results. Each treatment consisted of three plates, each containing thirty seeds. Error bars represent SEM. The asterisk indicates that the corresponding mean of the treatment is significantly different from the mean of controls for each time point indicated according to Student's t-test ($P<0.05$).

Figure 5: Iron accumulation in the fruit seed juncture is conserved in Arabidopsis seeds and determined by MTP8. A, sketch outlining the protocol. Seeds were cut along the red dashed line after one day of imbibition, stained with Perls/DAB and placed on a slide for examination under a light microscope. B, Perls/DAB staining of seeds and their unstained controls in side and top view. The ring-like black coloration indicates Fe accumulation. C, ring-like Fe accumulation in wild-type plants (WT) and *mtp8* mutants (*mtp8-1* and *mtp8-2).* These images are close-ups of the chalazal regions of the seeds.

At least twenty seeds were analyzed for each genotype. M: micropylar region, Cha: chalazal region. -Perls/DAB: DAB intensification was conducted without Perls staining and serves as a negative control. D, GFP fluorescence of the phloem marker line SUC2::GFP. Mature seeds were imbibed for one day, cut following the sketch in Fig. 5A, and examined with a confocal laser microscope. At least five different seeds were investigated with comparable results. M: micropylar region, Cha: chalazal region. Bar: 50 µm