Mapping metal accumulation sites in crop fruits revealed a functional

iron reservoir in the tomato seed chalaza

Kumbirai Deon Mandebere[1], Utku Deniz[1], Katarina Vogel Mikus[2,3], Seckin Eroglu[1*]

1 Department of Biological Sciences, Middle East Technical University, Ankara, Turkey

2 Jozef Stefan, Jamova cesta 39, Ljubljana, Slovenia

3 Biotechnical Faculty, University of Ljubljana, Jamnikarjeva ulica 101, Ljubljana,

Slovenia

Corresponding Author: Seckin Eroglu, erogluseckin@gmail.com

Running title: Metal accumulation sites in crop fruits

Keywords: fruit, synchrotron, X-ray, metal, iron, germination, tomato

Abstract

Global fruit production suffers from pre- and post- harvest losses, part of which are

related to metal deficiencies. Despite fruits being one of the most widely consumed

plant parts, the spatial distribution of metals and their physiological significance

remained largely unexplored. In this study, we investigated metal accumulation sites in

fruits of 28 major crops by using X-ray and histochemical-based techniques. We found

that calcium accumulated in the outermost hardened tissues, potassium in sugar

accumulating fleshy tissues, and iron (Fe) in vascular tissues in a conserved manner.

Vascular Fe pattern traced to the seed revealed an Fe reservoir at the fruit-seed

juncture in tomato, which persisted in the seed's chalazal region upon dispersal. To

determine the physiological function of these stored reserves, we manipulated Fe

bioavailability. Two opposely acting chelators, desferoxamine, an Fe immobilizer,

delayed germination, while nicotianamine, a mobilizer, accelerated it in wild-type plants

but not in mutants with low chalazal Fe. Additionally, external Fe supplementation also

increased germination speed in a dose-dependent manner. Collectively, these findings demonstrate that fruit vasculature serves as a critical delivery system to establish seed Fe pools, which are determinants of seed germination speed. This study provides the first comprehensive atlas of metal hotspots in fleshy fruits and links anatomical distribution to a defined physiological mechanism in seed biology.

Introduction

Fruits are essential to a nutritious human diet. To meet and sustain the dietary needs of the growing global population, fruit production must increase from one billion tons to over two billion tons per year by 2050 (Stratton et al., 2021; Willett et al., 2019). A key challenge in achieving this goal is reducing pre- and post-harvest losses, which can account for up to half of the production, making fruits the most wasted type of food (Gustavsson et al., 2011; Wang et al., 2022). Metal deficiencies are significant factors contributing to such losses, limiting the quality and quantity of fruits (Andresen et al., 2018; Kaya and Higgs, 2002; Suzuki et al., 2003). The effects of metal deficiencies can be either direct or indirect. As a direct effect, insufficient calcium levels in apples can result in bitter pit disease (Song et al., 2018; Torres et al., 2021; Van Goor, 1971). As an indirect (i.e., systemic) effect, the lack of a particular metal in sink tissues can cause metabolic remodeling, leading to lower quantity and quality of fruits (Àlvarez-Fernàndez et al., 2006; Álvarez-Fernández et al., 2003; El-Gioushy et al., 2021). Furthermore, metal deficiencies may also weaken the defense of fruits against several microbial diseases (Conway et al., 2001; Morina and Küpper, 2022).Despite the importance of metals in fruit biology, protocols for investigating their distribution in plant tissues often fail to yield the desired results.

The primary methods for exploring the spatial localization of metals in plant organs such as seeds, roots, or leaves are X-ray-based analysis and histochemical staining (Mijovilovich et al., 2020; Wu and Becker, 2012). However, these methods are often impractical for fleshy fruits with high water content (Lombi et al., 2011; Yadav et al., 2021; Zhao et al., 2014). Consequently, spatial metal analysis of fruits is currently

limited to non-fleshy varieties such as grains or pods and, with only a few exceptions, does not include juicy, fleshy fruits like tomatoes or oranges (Trebolazabala et al., 2017). Mapping metals in fruits will also contribute to seed biology because the fruit is the maternal environment that loads the seed.

Fruits use complex strategies to ensure dispersed seeds germinate in favorable conditions. In a recent study, it was shown that seeds are sealed with additional hydrophobic barriers such as lignin and suberin when mother plants are subjected to high heat during the bolting stage (Hyvärinen et al., 2025). Mother plant may determine germination speed by differential loading of metals to the seeds. Plants cultivated in metal-deficient basic soils produced late-germinating seeds (Murgia and Morandini, 2017). Since Fe and copper can impact cell wall integrity through redox reactions (Chen et al., 2023) it was speculated that they may also facilitate germination by weakening the cell walls of the endosperm (Mari et al., 2020); however, such has never been directly tested. Therefore, although there are some indications, a direct link between metals and germination speed has not been established.

Here, we systematically investigated metal accumulation sites in freeze-dried fruit samples using X-ray fluorescence and histochemical staining. This approach identified metal "hot spots", some of which are well-conserved. We further studied one of these metal hotspots to investigate its possible physiological functions.

Material and Methods

Research material

To examine metal localization in fruits, we selected 28 species to represent the major fruit crops consumed globally. Samples were obtained from markets and bazaars to ensure that the findings reflect metal localization patterns in agriculturally relevant, commercially available produce, rather than solely in model organisms grown under artificial laboratory conditions. Fruits were collected in Ankara, Turkey, throughout the

year (Table S1). For experiments with tomato seeds, if not otherwise indicated, wild-

type *Solanum lycopersicum* cv. Bonner Beste plants and its mutant *chloronerva* were

used (Herbik et al., 1996).

Tabletop XRF

To prepare the fruits for XRF analysis, they were freshly sliced both longitudinally and

cross-sectionally to a thickness of 2-3 mm using a V-blade (Fig. S1). The prepared fruit

slices were immediately immersed in liquid nitrogen and kept frozen at -80°C for 24

hours. Frozen samples were then placed in a vacuum freeze dryer for a main drying

phase of 23 hours at -54°C and 0.024 millibars, followed by an additional hour of final

drying at -60°C and 0.011 millibars. Once dried, fruit samples were stored at room

temperature until the analysis.

Freeze-dried fruit samples were analyzed by a micro XRF analyzer (Horiba XGT-9000,

Japan). The benchtop system was equipped with a Rh tube (50 kV, 1 mA) yielding a

polychromatic excitation beam focused to 15 µm by a polycapillary lense. X-ray

fluorescence was detected with a silicon drift diode (SDD) detector. The samples were

mapped with 100-120 µm step size and 200 ms dwell time. The spectra in each pixel

were recorded with the Horiba software, exported as .H5 files, and processed with the

PyMCA 5.8.1 toolkit (Bolte et al., 2011) where full spectra deconvolution and fitting was

performed. Maps were then generated in the RGB correlator of PyMCA on the basis of

the intensities of particular fluorescence K lines (Solé et al., 2007).

Synchrotron XRF

For examining metal enrichment in the vasculature of tomato fruits (*Solanum*

lycopersicum cv. Moneymaker) in high resolution and with minimum treatments, fresh

cut tomato pieces were embedded in the cryo-embedding matrix OCT and immediately

plunged into liquid nitrogen. Fifty µm cross sections were obtained using a Leica

RM2265 semi-automated rotary cryomicrotome and immediately transferred to a

sample holder between two Ultralene foils. The samples were kept in liquid nitrogen

during the transfer and analysis with the Scanning X-ray Microscope (SXM) at ID21 X-

Ray Microscopy Beamline at the European Synchrotron Radiation Facility (Grenoble,

France). The beam size was focused using Kirckpatrick Baez mirror optics to about

1x0.6 (HxV) μm^2. Elemental localization and distribution of Fe were performed at 7.3

keV excitations above the Fe K-edge and 100 ms dwell time. The energy was selected

using a double-crystal monochromator with a Si111 crystal pair. The X-ray fluorescence

signal from the samples was detected using a silicium drift diode detector with an 80

mm^2 active area from SGX Sensortech. The incoming beam intensity was monitored

using a photodiode. The X-ray fluorescence data from each pixel in the maps were fitted

to obtain the final elemental distribution images using the batch processing routines

from PyMCA 5.9.2.

For examining metal accumulation in the chalaza of tomato seeds in high resolution,

two-dimensional SXRF imaging was conducted. Dry seeds were placed on a Kapton

tape. Images were collected at beamline 4-BM of the National Synchrotron Light

Source-II (NSLS-II; Brookhaven National Laboratories, Upton, NY, USA). The elemental

maps were collected using an incident energy of 12 keV, a 7μm step size, and a 100 ms

dwell time. Raw data from the NSLS-II was analyzed using the Larch software package

to produce fitted X-ray fluorescence images. Final SXRF images and intensity profile

plots were generated using the Nikon NIS-elements software.

Histochemical staining of fruits

For preparing the fruit samples for histochemical staining, freeze-dried fruit pieces that

were prepared as described above for the tabletop XRF analyses were used as starting

material. Colorful fruits were subjected to decolorization. Decolorization was conducted

by immersing the pieces into a fixing solution (methanol: chloroform: glacial acetic acid;

6:3:1) in a glass Petri dish and shaking at 90 rpm until the natural color disappeared.

The fixing solution was renewed several times during the shaking. To stop the

decolorization step, the fixative was removed by washing the samples at least three

times with distilled water.

For the histochemical staining of Fe, Perls staining was used as previously described (Roschzttardtz et al., 2009). Briefly, Perls stain solution was prepared fresh by mixing 4% K-ferrocyanide ($K_4Fe(Cn)_6$) and 4% HCl stock solutions in a 1:1 (v:v) proportion and vacuum infiltration for one hour at 500 millibars. Next, the staining solution was removed, and the samples were washed three times with distilled water for 1-2 min each by slight shaking. The samples were stored in distilled water at 4°C until observation with a stereo microscope.

For histochemical Fe staining on the thin cross sections of tomato fruits, samples prepared for Perls staining were first embedded in resin, cut, and then stained. Pieces were washed three times with 0.1 M phosphate buffer (pH 7.4) and dehydrated in successive baths of 50%, 70%, 90%, 95%, and 100% ethanol, butanol/ethanol 1:1 (v/v), and 100% butanol. Then, the pieces were embedded in Technovit 7100 resin (Kulzer, Tokyo, Japan) according to the manufacturer's instructions and sliced into thin sections (10 μm). The sections were deposited on glass slides and incubated for 45 min in Perls stain solution. Perls solution was removed, slides were washed three times and directly observed with a compound microscope.

Elemental analysis by ICP-OES

To quantify Fe accumulation in the chalazal part, elemental analysis on seed parts were conducted. To this end, mature tomato seeds (*Solanum lycopersicum* cv. Candela) were imbibed for 24 hours and dissected into three parts. Dissected samples were dried in an oven at 65°C for one week. Dried plant samples were finely ground with an electric coffee grinder. Approximately 0.3 g of the dried and ground plant samples were weighed and placed in microwave digestion tubes. On top of each sample, 2 ml of 30% H_2O_2 and 5 ml of HNO_3 were added, and the digestion was conducted in a closed-vessel microwave system (MarsExpress; CEM Corp., Matthews, NC, USA). After cooling down sufficiently, the total sample volume was finalized to 20 ml by adding double-deionized water, and samples were filtered through analytical filter papers (Macherey-Nagel, Ø125 mm, blue band). Inductively coupled plasma optical emission spectrometry (ICP-OES) Agilent 5800 ICP-OES (Agilent Technologies, USA), was used

to determine the concentrations of micronutrients in digested plant samples. The

accuracy of the element analyses was validated using certified standard reference

materials obtained from the National Institute of Standards and Technology

(Gaithersburg, MD, USA).

Ferric chelate reductase assay

Ferric chelate reductase activity of endosperm exudates was assessed according to a

method previously described (Grillet et al., 2014). Mature tomato seeds (*Solanum*

lycopersicum cv. Candela) were imbibed in water for 24 h at room temperature and cut

longitudinally. Embryos were discarded, and five endosperms were pooled as one

biological sample (n=3). Isolated endosperms were immersed in distilled water and

allowed to release their exudates for 24 hours. The obtained exudate solution was then

mixed (1:2 ratio) with ferric reductase activity assay solution (0.2 mM CaSO4, 5

mM MES (pH 5.5), 0.2 mM ferrozine, and 0.1 mM Fe-EDTA) on a 12-well plate in the

presence or absence of ascorbate oxidase enzyme (1.5 AOX units ml^{-1}). Following an

incubation of 3 hours, endosperms were quickly dried on tissue paper and weighed. An

800 µl aliquot was taken from the remaining solution and the absorbance was measured

at 562 nm. Experiments were repeated three times with at least three biological

replicates and the result of one representative experiment is shown.

Germination tests

For all germination experiments, plated tomato seeds were incubated for two days in a

refrigerator for stratification and then placed in a growth chamber at 25 °C under

continuous light. Germination was scored over time, with a seed considered germinated

when the radicle protruded through the testa.

To determine how internal Fe availability impacts germination, germination tests were

conducted with tomato (*chloronerva* and its wild type Bonner Beste) in the presence and

absence of a strong Fe chelator. Seeds were sown on Petri dishes (30 seeds per dish,

three replicates) filled with water agar (0.7% agar, buffered at pH 5.6 with MES) with or

without DES (250 µM) or NA (5 µM). Results were analyzed by two-way ANOVA. Upon

finding a significant difference between treatments, Tukey's test was conducted to

identify the time points treatments show significantly different results.

To determine the trends of how metal supplementation impacts germination speed,

$FeCl_2$, $FeCl_3$, Fe-EDTA, Fe-PDMA (Prepared as previously described by Suzuki et al.,

2024), $MnCl_2$, and $CuSO_4$ were added to water agar (0.7% agar, buffered at pH 5.6 with

MES) at 100 µM final concentration of each. 50 seeds for each treatment were counted.

Measurement of hydroxyl radical production

To quantify hydroxyl radical production in the seeds, a radiolabeled benzoyl probe,

BzK_5Me, was used as described by Fry et al. (2002) and Miller and Fry (2004). This

tracer is excluded from the symplast of the cells and reacts specifically with the hydroxyl

radical to form tritiated water, ^3H_2O. ^3H_2O release acts as a proxy for hydroxyl radical

production. At least five seeds were imbibed in a 12-well plate in 2ml solutions on a

shaker in a refrigerator for two days and one day at room temperature, in solutions

containing the probe and the treatments (100 µM $FeCl_2$, 5 µM NA or 100 µM Des).

Solutions were then filtered using Dowex 50 H^+ beads to remove unreacted probe. The

radioactivity of the filtrate was measured without and with evaporation following

resuspension in scintillant buffer. The radioactivity of the evaporated ^3H_2O was

calculated.

Results

Calcium, potassium, and iron accumulate in specific tissues of the fruit

By introducing an initial freeze-drying step, we determined metal localizations in fruit

pieces obtained from various crop species using a bench-top X-ray machine. While X-

ray mapping revealed metal localizations, it provided only limited resolution (Fig. S2).

We examined metal distribution maps to identify conserved metal-enriched sites. We

found that in all fruits, potassium accumulated in their fleshy (water-rich) parts, and

calcium preferentially accumulated on the outer skin of the fruits (Fig. 1, Fig. S2).

Alternatively, histochemical stains have been used to localize metals, even in cellular

compartments (Eroglu et al., 2019). To identify accumulation sites of Fe, we stained fruit

slices with Prussian Blue (also known as Perls staining). Perls staining resulted in

higher resolution compared to the benchtop X-ray (Fig. 2). Although Perls staining

showed homogeneous coloration in some fruits (Fig. S3), it led to a more pronounced

blue color throughout vascular tissues of many others (blue arrow heads in Fig. 2). This

result shows that in fruits, Fe is preferentially stored in the vasculature in a conserved

manner. The use of fixatives has been criticized because they may remobilize metals,

leading to artifacts and misinterpretation of results (van der Ent et al., 2018). To address

this constraint, we employed synchrotron X-ray mapping, which provided sufficient

resolution to map vasculature in fruit tissues. Mapping of freshly cut tomato pericarp

under cryo conditions revealed a hollow Fe accumulation pattern, possibly

corresponding to vasculature (Fig. S4). While we were unable to identify Fe-

accumulating cell types, our result suggests that the preferential Fe accumulation in the

vasculature is not an artifact due to sample preparation.

Iron accumulates in the chalazal region of tomato seeds

Seeds of various species have been extensively investigated for metal distribution

(Eroglu et al., 2019; Ibeas et al., 2017); however, in tomato seeds, such distribution

patterns remain elusive. In the present survey, we noticed Fe accumulation in the fruit-

seed juncture of tomato slices (red arrowhead in Fig. 2). We examined whether metals

other than Fe are also enriched in this region by mapping seed metal distribution using

synchrotron X-ray fluorescence and quantifying them with ICP-OES. Synchrotron X-Ray

fluorescence showed Mn partitions towards the endosperm and Zn towards the cells

that will develop into apical and root meristems (Fig. 3A). Fe accumulated both in the

embryo and the endosperm. Fe accumulation in the embryo was pronounced on the

prevascular strands. Fe accumulation in the endosperm was enriched in regions close

to the germination site, confirming the localization obtained by Perls staining.

Furthermore, signal intensity in the chalaza was comparable to that of the embryo. A

tomato mutant, *chloronerva,* was reported to be devoid of a Fe chelator, NA, and to

contain less Fe (Herbik et al., 1996). In addition to wild-type seeds, we examined

chloronerva seeds to assess the impact of nicotianamine on seed Fe distribution. As

expected, *chloronerva* showed lower signals for all Fe, Mn, and Zn, indicating lower

levels of these metals in the mutant (Fig. 3A, bottom panel). However, except for
predominant Mn accumulation in the chalaza, *chloronerva* showed the same metal
distribution pattern as the wild-type tomato. We also examined spatial metal distribution
in the wild type seeds by analyzing metal concentrations in cut seed parts. This analysis
showed the chalazal endosperm (more accurately, chalazal and micropylar endosperm,
including the associated seed coat) contained approximately twice as much Fe
compared to the remainder of the seed in agreement with the X-Ray fluorescent
measurements and Perls staining. Overall, the mapping of metals in tomato seeds
showed that the chalaza is a primary Fe-enrichment site, and *chloronerva* can serve as
a tool to investigate the physiological function of this store.

In a germinating seed, endosperm-accumulated nutrients can be remobilized to support
the embryo as it transforms into a seedling (Uraguchi and Fujiwara, 2011). To validate a
possible role for Fe in feeding the germinating embryo, we examined the fate of the Fe
deposits after germination. In seeds of one-week-old seedlings, seed endosperm
tissues were degraded, reminiscent of nutrient remobilization from endosperm into the
embryo (Fig. S5). However, Perls staining showed that these endosperm remnants
preserved the preferential Fe accumulation in the chalaza (Fig. S5, denoted "After")
similar to the ungerminated control (Fig. S5, denoted "Before"). Metal analysis showed
that the Fe concentration doubled. Since the cultivation media lacked nutrients, the
observed increase in Fe concentrations was primarily due to mass loss from endosperm
degradation. These results negate the hypothesis that Fe in the chalaza nourishes the
germinating embryo.

Iron is involved in germination
Next, we assessed whether Fe could promote germination by comparing the effects of
Fe chelators on seeds with contrasting Fe concentrations. The Fe chelator DES has
been used to counteract Fe poisoning in humans (Rafati Rahimzadeh et al., 2023) and
to render Fe stores in the plant tissues unavailable for biological processes (Aznar et
al., 2014). Targeting Fe stores in seeds with DES led to a delay of approximately 6
hours (Fig. 4A). Following the same logic, we also added NA to the medium. As

previously mentioned, plants use NA to mobilize metals in the long distant transport

(Takahashi et al., 2003). Although the effect was less striking than that of DES, NA

increased germination speed(Fig. 4A). However, the germination of the *chloronerva*

mutant, which contains low concentrations of Fe, was not affected by either DES or NA

(Fig. 4B), suggesting that seed Fe reserve acts as a determinant for germination speed.

After investigating the effect of Fe mobilization on germination speed, we examined

whether Fe application can promote germination similar to NA. We added various forms

of Fe, both chelated and unchelated, to the medium. Results showed that independent

of the form, Fe increased germination speed (Fig. 4C). To determine whether the early

germination phenotype is specific to Fe, we tested various other micronutrients. None of

them caused an apparent increase in the germination speed (Fig. 4D). Interestingly,

increasing Fe concentrations, even to physiologically toxic levels (200 and 500 µM),

tended to increase the germination speed further (Fig. 4E).

Germination speed is determined by the amount of hydroxyl radicals produced in the

endosperm (Muller et al., 2009; Schweikert et al., 2002). To investigate the mechanism

of Fe-triggered germination, we hypothesized that Fe aids germination by weakening

the endosperm through the production of hydroxyl radicals. A common pathway in

nature for hydroxyl radical production is through the Fenton chain reaction, which

requires Fe, hydrogen peroxide, and ascorbate (Burkitt and Gilbert, 1990a). Since,

hydrogen peroxide production in tomato seeds during imbibition was shown before

(Morohashi, 2002) and we found that tomato endosperm also produces ascorbate

during imbibition (Fig. S6); we predicted that hydroxyl radical production to increase in

the presence of Fe. Comparison of hydroxyl radical production showed that seeds

produce hydroxyl radicals during imbibition, but this production was independent of Fe

availability (Fig. 4F).

334   Discussion

X-ray and histochemistry-based methods highlighted numerous metal hotspots in

various plant organs (Eroglu et al., 2019, 2016; Kim et al., 2006). In this study, we

examined metal localizations in fruits. Our results show that in fruits, metals including

calcium, potassium, and Fe preferentially accumulate in certain tissues. We further
focused on one of the Fe accumulation sites, in the fruit-seed juncture, and found that
this Fe store mobilizes to facilitate seed germination upon seed dispersal.
341
Histochemical staining of freeze-dried and decolorized fruits provides tissue-level
information
The histochemical approach to mature fleshy fruits often faces two main obstacles: 1)
too much water, limiting the penetration and activity of chemicals, and 2) too many
pigments, limiting the visual inspection of the stain. Industrially, fruits have been
extensively dried for various purposes. Among drying technologies, vacuum freeze
drying appears to be the golden standard for human consumption (Harper and Tappel,
1957). We observed that once freed from water, the fruit slices behaved like regular
tissue. After vacuum freeze drying, fixatives could decolorize the fruit. This may
represent an advance over the techniques described in the literature. The most
common way to circumvent the natural bright colors of fruits like tomato, has been to
analyze the sample in earlier developmental stages (Saed Taha et al., 2012; Wang et
al., 2014). We propose here that freeze drying, coupled with histochemistry, provides an
efficient and quick method to reveal immobile metal hotspots in fruits (Fig. 2 and
Fig.S3). Any finding can then be later confirmed with more advanced methods that offer
higher resolution and quantification and avoid possible metal re-localization (Fig. S4).
358
Iron follows the vasculature of fruits and seeds
Fe preferentially accumulates in the vascular tissues of several plant parts. Leaves (Xu
et al., 2022), embryos inside the seeds (Kim et al., 2006 and Fig. S7), and fruits (this
study) accumulate Fe in and around their provascular or vascular strands. In mature
fruits, vascular Fe may represent the most prominent Fe accumulation site (Fig. 3).
However, this accumulation can appear more pronounced than it is, due to clearing off
weakly bound Fe in the rest of the fruit during fixation and decolorization. The
physiological function of Fe in the fruit vasculature is unclear. Fe deficiency in fruit trees
has not been linked with a disease that is specific to fruit vasculature (Àlvarez-
Fernàndez et al., 2006). Alternatively, an immobile Fe reservoir may be a "byproduct" of

extensive callose production. Callose is required to close the xylem connection in fruits

(Knipfer et al., 2015) and may cause the oxidation of a large amount of Fe for its

production (Müller et al., 2015). Another possibility is that the main vasculatures of fruits

serve as a sink to load Fe into the seeds. In this model, the vasculature may represent a

reservoir of excess Fe left over from seed Fe filling, as this process is highly regulated

to prevent Fe-mediated oxidative damage in seeds (Sun et al., 2021). The remaining Fe

can then be bound to cell walls that can quickly absorb Fe(III) (Xu et al., 2022). The

current study identified the vasculature as the main Fe-accumulating tissue in fruits;

however, further investigations to identify the cells within the vasculature that

accumulate Fe and into the underlying mechanisms of this process are required to

substantiate this supposition.

Iron in the seed chalaza facilitates germination

A fraction of Fe-loaded vasculature retains in the seeds upon dispersal from the fruit

(Fig. 3, S5, S7 and Liu et al., 2023). This Fe reservoir facilitates germination (Fig. 4).

Germination requires softening of the endosperm to allow radical protrusion, a major

process that includes cell wall-cutting hydroxyl radicals (Grillet et al., 2013; Muller et al.,

2009). A well-known source of hydroxyl radicals derives from the reaction of Fe with

H_2O_2, referred to as Fenton reaction (Burkitt and Gilbert, 1990b). Therefore, Fe deposits

in the chalaza may facilitate germination by generating cell wall-attacking hydroxyl

radicals. This hypothesis is ruled out, as we failed to link hydroxyl radical production to

Fe availability (Fig. 4F). To elucidate the mechanism, future research should integrate

reverse genetic screens (especially under high Fe supplementation) to identify genes

involved in Fe-triggered germination.

Micro-XRF mapping is applicable to investigate nutrient deficiency-related fruit diseases

Micro-XRF mapping has been successfully used to examine metal distributions in

various plant organs, including leaves, roots, stems, and seeds (Singh et al., 2022). We

detected calcium to be preferentially accumulating in the outer part of fruits, especially

in the citrus genus (Fig. 1, S2). This is in agreement with previous reports where the

highest calcium levels were generally observed in the peel of the fruits (Saure, 2005).

Higher calcium in the outer parts of the fruits is probably caused by cell wall thickening.
Calcium is a major component of the cell wall (Thor, 2019), and outer parts of the fruits
usually possess thicker cell walls to maintain integrity (Forlani et al., 2019; Hallett and
Sutherland, 2005; Jeffery et al., 2012). When fruits contain insufficient calcium they
may be more prone to infections, as calcium deficiency has been reported to cause
several diseases, including bitter pit in apples and blossom-end rot in tomatoes,
peppers, and watermelons (de Freitas and Mitcham, 2012). Potassium preferentially
accumulates in the fleshy parts of the fruits in a conserved manner (Fig. 1 and Ahmed
et al., 2024). This is not surprising since potassium is involved in osmoregulation and
sugar accumulation, whereas the fleshy parts of the fruits often accumulate water and
sugar (Kumar et al., 2006).
411
Imaging element distributions of fruits is associated with certain limitations.
Polychromatic excitation with an X-ray tube (Rh in our case) gave a high background
due to scattering in the energy range of middle z-elements, decreasing the detection
limit for trace elements like Fe, Cu and Zn. We found that in fruits, Fe was under the
detection limit for imaging despite the presence of an Fe peak (Fig. S8). Nevertheless,
our data showed that micro-XRF can be very useful to image distribution of macro-
elements like potassium and calcium which can aid in investigating whether specific
disease symptoms overlap with perturbations in metal accumulating sites.
420

Conclusions

Increasing fruit production should target decreasing pre- and post-harvest losses. To
reduce such losses, a better understanding of fruit biology is required. This study
extends the basic techniques to localize metals to mature fleshy fruits, contributing to
preventing yield loss in fruit production and storage in the long run. Fruits preferentially
accumulate Fe in their vascular, potassium in their fleshy and calcium in their epidermal
tissues. While tomato seeds preferentially accumulate Fe in the fruit-seed juncture,
seed Fe determines germination speed.
429

Author contributions

431   S.E. designed the study. U.D. and D.M. prepared the fruits and conducted the Perls

432   staining. D.M. and S.E. conducted the seed-related experiments. K.V.M conducted

433   benchtop X-Ray analyses. All authors discussed the data. S.E. wrote the manuscript

434   with contributions from the other authors. All authors read and approved the final

435   version of the manuscript.

Acknowledgments

We thank Mary Lou Guerinot and Ryan Tappero for synchrotron imaging at X-ray Fluorescence Microprobe (XFM) beamline 4-BM of the National Synchrotron Light Source II, a US Department of Energy (DOE) Office of Science User Facility operated for the DOE Office of Science by Brookhaven National Laboratory under Contract DE-SC0012704. We acknowledge the European Synchrotron Radiation Facility (ESRF) for provision of synchrotron radiation facilities under proposal number LS-3343 for beamline, and we would like to thank Hiram Castillo for assistance and support in using beamline ID21. We also acknowledge the SIMBION (Eurobioimaging) network for benchtop micro-XRF analyses and ARIS for program financing (P1-0212, Plant Biology). We further thank Petra Bauer for *chloronerva* mutants, Abdulsamet Sakalar for helping with the histochemical staining of fruits, Catherine Curie for help with the germination experiments, Stephen Fry for hydroxyl radical measurements and Ahmet Bakirbas for critical reading of the manuscript. This study was supported by the Scientific and Technological Research Council of Turkiye (TUBITAK) under Grant Number 122Z109; European Union under COST action CA19116, Trace metal metabolism in plants (PLANTMETALS), EMBO scientific exchange grant (10056) and TENMAK, Ankara, Turkey.

References

Ahmed, I.A.M., Özcan, M.M., AlJuhaimi, F., Albakry, Z., 2024. The Monitoring of Accumulations of Elements in Apple, Pear, and Quince Fruit Parts. Biol Trace Elem Res. https://doi.org/10.1007/s12011-024-04223-3

Àlvarez-Fernàndez, A., Abadía, J., Abadía, A., 2006. Iron deficiency, fruit yield and fruit quality. Iron nutrition in plants and rhizospheric microorganisms 85–101.

Álvarez-Fernández, A., Paniagua, P., Abadía, J., Abadía, A., 2003. Effects of Fe Deficiency Chlorosis on Yield and Fruit Quality in Peach (*Prunus persica* L. Batsch). J. Agric. Food Chem. 51, 5738–5744. https://doi.org/10.1021/jf034402c

Andresen, E., Peiter, E., Küpper, H., 2018. Trace metal metabolism in plants. Journal of experimental botany 69, 909–954.

Aznar, A., Chen, N.W.G., Rigault, M., Riache, N., Joseph, D., Desmaële, D., Mouille, G., Boutet, S., Soubigou-Taconnat, L., Renou, J.-P., Thomine, S., Expert, D., Dellagi, A., 2014. Scavenging Iron: A Novel Mechanism of Plant Immunity Activation by Microbial Siderophores. Plant Physiology 164, 2167–2183. https://doi.org/10.1104/pp.113.233585

Bolte, S., Lanquar, V., Soler, M.-N., Beebo, A., Satiat-Jeunemaître, B., Bouhidel, K., Thomine, S., 2011. Distinct Lytic Vacuolar Compartments are Embedded Inside the Protein Storage Vacuole of Dry and Germinating Arabidopsis thaliana Seeds. Plant Cell Physiol 52, 1142–1152. https://doi.org/10.1093/pcp/pcr065

Burkitt, M.J., Gilbert, B.C., 1990a. Model Studies of the Iron-Catalysed Haber-Weiss Cycle and the Ascorbate-Driven Fenton Reaction. Free Radical Research Communications 10, 265–280. https://doi.org/10.3109/10715769009149895

Burkitt, M.J., Gilbert, B.C., 1990b. Model studies of the iron-catalysed Haber-Weiss cycle and the ascorbate-driven Fenton reaction. Free radical research communications 10, 265–280.

Chen, C., Zhang, Y., Cai, J., Qiu, Y., Li, L., Gao, C., Gao, Y., Ke, M., Wu, S., Wei, C., Chen, J., Xu, T., Friml, J., Wang, J., Li, R., Chao, D., Zhang, B., Chen, X., Gao, Z., 2023. Multi-copper oxidases SKU5 and SKS1 coordinate cell wall formation using apoplastic redox-based reactions in roots. Plant Physiol 192, 2243–2260. https://doi.org/10.1093/plphys/kiad207

Conway, W.S., Sams, C.E., Hickey, K.D., 2001. Pre-and postharvest calcium treatment of apple fruit and its effect on quality, in: International Symposium on Foliar Nutrition of Perennial Fruit Plants 594. pp. 413–419.

de Freitas, S.T., Mitcham, E.J., 2012. Factors Involved in Fruit Calcium Deficiency Disorders, in: Horticultural Reviews. John Wiley & Sons, Ltd, pp. 107–146. https://doi.org/10.1002/9781118351871.ch3

El-Gioushy, S.F., Ding, Z., Bahloul, A.M., Gawish, M.S., Abou El Ghit, H.M., Abdelaziz, A.M., El-Desouky, H.S., Sami, R., Khojah, E., Hashim, T.A., 2021. Foliar application of nano, chelated, and conventional iron forms enhanced growth, nutritional status, fruiting

aspects, and fruit quality of Washington navel orange trees (Citrus sinensis L. Osbeck). Plants 10, 2577.

Eroglu, S., 2018. Chapter Four - Metal Transport in the Developing Plant Seed, in: Maurel, C. (Ed.), Advances in Botanical Research, Membrane Transport in Plants. Academic Press, pp. 91–113. https://doi.org/10.1016/bs.abr.2018.09.010

Eroglu, S., Giehl, R.F.H., Meier, B., Takahashi, M., Terada, Y., Ignatyev, K., Andresen, E., Küpper, H., Peiter, E., von Wirén, N., 2017. Metal Tolerance Protein 8 Mediates Manganese Homeostasis and Iron Reallocation during Seed Development and Germination. Plant physiology 174, 1633–1647. https://doi.org/10.1104/pp.16.01646

Eroglu, S., Karaca, N., Vogel-Mikus, K., Kavčič, A., Filiz, E., Tanyolac, B., 2019. The conservation of VIT1-dependent iron distribution in seeds. Frontiers in plant science 10, 907.

Eroglu, S., Meier, B., von Wirén, N., Peiter, E., 2016. The vacuolar manganese transporter MTP8 determines tolerance to iron deficiency-induced chlorosis in Arabidopsis. Plant physiology 170, 1030–1045. https://doi.org/10.1104/pp.15.01194

Forlani, S., Masiero, S., Mizzotti, C., 2019. Fruit ripening: the role of hormones, cell wall modifications, and their relationship with pathogens. Journal of Experimental Botany 70, 2993–3006. https://doi.org/10.1093/jxb/erz112

Fry, S.C., Miller, J.G., Dumville, J.C., 2002. A proposed role for copper ions in cell wall loosening. Plant and Soil 247, 57–67.

Grillet, L., Mari, S., Schmidt, W., 2013. Iron in seeds–loading pathways and subcellular localization. Frontiers in plant science 4.

Grillet, L., Ouerdane, L., Flis, P., Hoang, M.T.T., Isaure, M.-P., Lobinski, R., Curie, C., Mari, S., 2014. Ascorbate efflux as a new strategy for iron reduction and transport in plants. Journal of Biological Chemistry 289, 2515–2525.

Gustavsson, J., Cederberg, C., Sonesson, U., Van Otterdijk, R., Meybeck, A., 2011. Global food losses and food waste.

Hallett, I.C., Sutherland, P.W., 2005. Structure and Development of Kiwifruit Skins. International Journal of Plant Sciences 166, 693–704. https://doi.org/10.1086/431232

Harper, J.C., Tappel, A.L., 1957. Freeze-drying of food products. Advances in Food Research 7, 171–234.

Herbik, A., Giritch, A., Horstmann, C., Becker, R., Balzer, H.-J., Baumlein, H., Stephan, U.W., 1996. Iron and copper nutrition-dependent changes in protein expression in a tomato wild type and the nicotianamine-free mutant chloronerva. Plant Physiology 111, 533–540.

Hindt, M.N., Akmakjian, G.Z., Pivarski, K.L., Punshon, T., Baxter, I., Salt, D.E., Guerinot, M.L., 2017. BRUTUS and its paralogs, BTS LIKE1 and BTS LIKE2, encode important negative regulators of the iron deficiency response in Arabidopsis thaliana. Metallomics 9, 876–890. https://doi.org/10.1039/C7MT00152E

Hyvärinen, L., Fuchs, C., Utz-Pugin, A., Gully, K., Megies, C., Holbein, J., Iwasaki, M., Demonsais, L., Capitão, M.B., Barberon, M., Franke, R.B., Nawrath, C., Loubéry, S., Lopez-Molina, L., 2025. Temperature-dependent polar lignification of a seed coat suberin layer promoting dormancy in Arabidopsis thaliana. Proceedings of the National Academy of Sciences 122, e2413627122. https://doi.org/10.1073/pnas.2413627122

Ibeas, M.A., Grant-Grant, S., Navarro, N., Perez, M.F., Roschzttardtz, H., 2017. Dynamic Subcellular Localization of Iron during Embryo Development in Brassicaceae Seeds. Front. Plant Sci. 8, 1–8. https://doi.org/10.3389/fpls.2017.02186

Jeffery, J., Holzenburg, A., King, S., 2012. Physical barriers to carotenoid bioaccessibility. Ultrastructure survey of chromoplast and cell wall morphology in nine carotenoid-containing fruits and vegetables. Journal of the Science of Food and Agriculture 92, 2594–2602. https://doi.org/10.1002/jsfa.5767

Johri, B.M., Ambegaokar, K.B., Srivastava, P.S., 2013. Comparative embryology of angiosperms vol. 1/2. Springer Science & Business Media.

Kalmbach, L., Helariutta, Y., 2019. Sieve Plate Pores in the Phloem and the Unknowns of Their Formation. Plants 8, 25. https://doi.org/10.3390/plants8020025

Kaya, C., Higgs, D., 2002. Response of tomato (*Lycopersicon esculentum* L.) cultivars to foliar application of zinc when grown in sand culture at low zinc. Scientia Horticulturae 93, 53–64. https://doi.org/10.1016/S0304-4238(01)00310-7

Kim, S.A., Punshon, T., Lanzirotti, A., Li, L., Alonso, J.M., Ecker, J.R., Kaplan, J., Guerinot, M.L., 2006. Localization of iron in Arabidopsis seed requires the vacuolar membrane transporter VIT1. Science 314, 1295–1298.

Knipfer, T., Fei, J., Gambetta, G.A., McElrone, A.J., Shackel, K.A., Matthews, M.A., 2015. Water transport properties of the grape pedicel during fruit development: insights into xylem anatomy and function using microtomography. Plant Physiology 168, 1590–1602.

Kumar, A.R., Kumar, N., Kavino, M., 2006. Role of potassium in fruit crops-a review. Agricultural Reviews 27, 284–291.

Liu, X., Nakajima, K.P., Adhikari, P.B., Wu, X., Zhu, S., Okada, K., Kagenishi, T., Kurotani, K., Ishida, T., Nakamura, M., 2025. Fertilization-dependent phloem end gate regulates seed size. Current Biology.

Liu, X., Nakajima, K.P., Wu, X., Zhu, S., Adhikari, P.B., Okada, K., Kurotani, K., Ishida, T., Nakamura, M., Sato, Y., 2023. Seed size control via phloem end by callose deposition/degradation of β-1, 3-glucanase. bioRxiv 2023–07.

Lombi, E., Scheckel, K.G., Kempson, I.M., 2011. In situ analysis of metal(loid)s in plants: State of the art and artefacts. Environmental and Experimental Botany, Metal(loid) tolerance in plants and lichens 72, 3–17. https://doi.org/10.1016/j.envexpbot.2010.04.005

Mari, S., Bailly, C., Thomine, S., 2020. Handing off iron to the next generation: how does it get into seeds and what for? Biochemical Journal 477, 259–274.

Mijovilovich, A., Morina, F., Bokhari, S.N., Wolff, T., Küpper, H., 2020. Analysis of trace metal distribution in plants with lab-based microscopic X-ray fluorescence imaging. Plant Methods 16, 82. https://doi.org/10.1186/s13007-020-00621-5

Miller, J.G., Fry, S.C., 2004. N-[3H] Benzoylglycylglycylglycine as a probe for hydroxyl radicals. Analytical biochemistry 335, 126–134.

Morina, F., Küpper, H., 2022. Trace metals at the frontline of pathogen defence responses in non-hyperaccumulating plants. Journal of Experimental Botany 73, 6516–6524. https://doi.org/10.1093/jxb/erac316

Morohashi, Y., 2002. Peroxidase activity develops in the micropylar endosperm of tomato seeds prior to radicle protrusion. Journal of Experimental Botany 53, 1643–1650.

Müller, J., Toev, T., Heisters, M., Teller, J., Moore, K.L., Hause, G., Dinesh, D.C., Bürstenbinder, K., Abel, S., 2015. Iron-dependent callose deposition adjusts root meristem maintenance to phosphate availability. Developmental cell 33, 216–230.

Muller, K., Linkies, A., Vreeburg, R.A., Fry, S.C., Krieger-Liszkay, A., Leubner-Metzger, G., 2009. In vivo cell wall loosening by hydroxyl radicals during cress seed germination and elongation growth. Plant physiology 150, 1855–1865.

Murgia, I., Morandini, P., 2017. Iron deficiency prolongs seed dormancy in Arabidopsis plants. Frontiers in Plant Science 8, 2077.

Otero, S., Helariutta, Y., 2016. Companion cells: a diamond in the rough. Journal of experimental botany erw392.

Rafati Rahimzadeh, Mehrdad, Rafati Rahimzadeh, Mehravar, Kazemi, S., Moghadamnia, A.R., Ghaemi Amiri, M., Moghadamnia, A.A., 2023. Iron; Benefits or threatens (with emphasis on mechanism and treatment of its poisoning). Hum Exp Toxicol 42, 09603271231192361. https://doi.org/10.1177/09603271231192361

Roschzttardtz, H., Conéjéro, G., Curie, C., Mari, S., 2009. Identification of the Endodermal Vacuole as the Iron Storage Compartment in the Arabidopsis Embryo. Plant Physiol. 151, 1329–1338. https://doi.org/10.1104/pp.109.144444

Saed Taha, R., Ismail, I., Zainal, Z., Abdullah, S.N.A., 2012. The stearoyl-acyl-carrier-protein desaturase promoter (Des) from oil palm confers fruit-specific GUS expression in transgenic tomato. Journal of Plant Physiology 169, 1290–1300. https://doi.org/10.1016/j.jplph.2012.05.001

Saure, M.C., 2005. Calcium translocation to fleshy fruit: its mechanism and endogenous control. Scientia Horticulturae 105, 65–89. https://doi.org/10.1016/j.scienta.2004.10.003

Schweikert, C., Liszkay, A., Schopfer, P., 2002. Polysaccharide degradation by Fenton reaction- or peroxidase-generated hydroxyl radicals in isolated plant cell walls. Phytochemistry 61, 31–35. https://doi.org/10.1016/S0031-9422(02)00183-8

Singh, Vivek K., Sharma, N., Singh, Virendra K., 2022. Application of X-ray fluorescence spectrometry in plant science: Solutions, threats, and opportunities. X-Ray Spectrometry 51, 304–327. https://doi.org/10.1002/xrs.3260

Solé, V.A., Papillon, E., Cotte, M., Walter, P., Susini, J., 2007. A multiplatform code for the analysis of energy-dispersive X-ray fluorescence spectra. Spectrochimica Acta Part B: Atomic Spectroscopy 62, 63–68.

Song, W., Yi, J., Kurniadinata, O.F., Wang, H., Huang, X., 2018. Linking fruit Ca uptake capacity to fruit growth and pedicel anatomy, a cross-species study. Frontiers in plant science 9, 575.

Stratton, A.E., Finley, J.W., Gustafson, D.I., Mitcham, E.J., Myers, S.S., Naylor, R.L., Otten, J.J., Palm, C.A., 2021. Mitigating sustainability tradeoffs as global fruit and vegetable systems expand to meet dietary recommendations. Environ. Res. Lett. 16, 055010. https://doi.org/10.1088/1748-9326/abe25a

Sun, L., Wei, Y.Q., Wu, K.H., Yan, J.Y., Xu, J.N., Wu, Y.R., Li, G.X., Xu, J.M., Harberd, N.P., Ding, Z.J., 2021. Restriction of iron loading into developing seeds by a YABBY transcription factor safeguards successful reproduction in Arabidopsis. Molecular Plant 14, 1624–1639.

Suzuki, K., Shono, M., Egawa, Y., 2003. Localization of calcium in the pericarp cells of tomato fruits during the development of blossom-end rot. Protoplasma 222, 149–156. https://doi.org/10.1007/s00709-003-0018-2

Suzuki, M., Suzuki, Y., Hosoda, K., Namba, K., Kobayashi, T., 2024. The phytosiderophore analogue proline-2⊡-deoxymugineic acid is more efficient than conventional chelators for improving iron nutrition in maize. Soil Science and Plant Nutrition 70, 435–446. https://doi.org/10.1080/00380768.2024.2385401

Takahashi, M., Terada, Y., Nakai, I., Nakanishi, H., Yoshimura, E., Mori, S., Nishizawa, N.K., 2003. Role of nicotianamine in the intracellular delivery of metals and plant reproductive development. The Plant Cell 15, 1263–1280.

Thor, K., 2019. Calcium—Nutrient and Messenger. Frontiers in Plant Science 10.

Torres, E., Alegre, S., Recasens, I., Asín, L., Lordan, J., 2021. Integral procedure to predict bitter pit in 'Golden Smoothee'apples based on calcium content and symptom induction. Scientia Horticulturae 277, 109829.

Trebolazabala, J., Maguregui, M., Morillas, H., de Diego, A., Madariaga, J.M., 2017. Evaluation of metals distribution in Solanum lycopersicum plants located in a coastal environment using micro-energy dispersive X-ray fluorescence imaging. Microchemical Journal 131, 137–144.

Uraguchi, S., Fujiwara, T., 2011. Significant contribution of boron stored in seeds to initial growth of rice seedlings. Plant Soil 340, 435–442. https://doi.org/10.1007/s11104-010-0614-9

van der Ent, A., Przybyłowicz, W.J., de Jonge, M.D., Harris, H.H., Ryan, C.G., Tylko, G., Paterson, D.J., Barnabas, A.D., Kopittke, P.M., Mesjasz-Przybyłowicz, J., 2018. X-ray elemental mapping techniques for elucidating the ecophysiology of hyperaccumulator plants. New Phytologist 218, 432–452. https://doi.org/10.1111/nph.14810

Van Goor, B.J., 1971. The Effect of Frequent Spraying with Calcium Nitrate Solutions on the Mineral Composition and the Occurrence of Bitter Pit of the Apple Cox's Orange Pippin. Journal of Horticultural Science 46, 347–364. https://doi.org/10.1080/00221589.1971.11514415

Wang, M., Xu, Y., Yang, Y., Mu, B., Nikitina, M.A., Xiao, X., 2022. Vis/NIR optical biosensors applications for fruit monitoring. Biosensors and Bioelectronics: X 11, 100197. https://doi.org/10.1016/j.biosx.2022.100197

Wang, S., Lu, G., Hou, Z., Luo, Z., Wang, T., Li, H., Zhang, J., Ye, Z., 2014. Members of the tomato FRUITFULL MADS-box family regulate style abscission and fruit ripening. Journal of Experimental Botany 65, 3005–3014. https://doi.org/10.1093/jxb/eru137

Willett, W., Rockström, J., Loken, B., Springmann, M., Lang, T., Vermeulen, S., Garnett, T., Tilman, D., DeClerck, F., Wood, A., 2019. Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. The lancet 393, 447–492.

Wu, B., Becker, J.S., 2012. Imaging techniques for elements and element species in plant science. Metallomics 4, 403. https://doi.org/10.1039/c2mt00002d

Xu, Z.-R., Cai, M.-L., Yang, Y., You, T.-T., Ma, J.F., Wang, P., Zhao, F.-J., 2022. The ferroxidases LPR1 and LPR2 control iron translocation in the xylem of Arabidopsis plants. Molecular Plant 15, 1962–1975.

Yadav, V., Arif, N., Singh, V.P., Guerriero, G., Berni, R., Shinde, S., Raturi, G., Deshmukh, R., Sandalio, L.M., Chauhan, D.K., Tripathi, D.K., 2021. Histochemical Techniques in Plant Science: More Than Meets the Eye. Plant and Cell Physiology 62, 1509–1527. https://doi.org/10.1093/pcp/pcab022

Zhao, F.-J., Moore, K.L., Lombi, E., Zhu, Y.-G., 2014. Imaging element distribution and speciation in plant cells. Trends in Plant Science 19, 183–192. https://doi.org/10.1016/j.tplants.2013.12.001

Figures and captions

Figure 1: Calcium accumulates in epidermal and potassium in fleshy parenchyma tissues of fruits. Freeze-dried fruit slices were examined with a bench-top micro-XRF. Left, kumquat; middle, lemon; right, squash. Red rectangles denote the analyzed area. At least two independent pieces were observed for each fruit with comparable results. Scale bar is 5 mm. K: Potassium, Ca: Calcium.

Figure 2: Iron accumulates in the fruit vasculature of diverse species and in the fruit-seed juncture in tomato. Cross-sectional or longitudinal cut fruit pieces were freeze-dried and decolorized with fixatives (methanol: chloroform: glacial acetic acid; 6:3:1). After fixation, the pieces were submerged in Perls staining solution for 2 to 16 hours. Blue coloration indicates Fe accumulation, blue arrowheads highlight vasculature stained blue in different fruit species. For this figure, images of fruits with noticeable Fe accumulation in the vascular bundles were chosen. Note that tomato accumulated Fe in the fruit seed juncture, shown by a red arrowhead. A-K; eggplant, cucumber, kiwi, kumquat, haricot, squash, green almond, blueberry, papaya, tomato, and passion fruit. At least five pieces of fruits were investigated, and a representative sample was photographed. Bar scale is 1 cm, unless otherwise indicated.

Figure 3: Metal loading to the fruit-seed juncture in tomato is specific to Fe and remains after seed dispersal. A, metal distribution in wild-type and mutant dry mature tomato seeds. Two seeds from each genotype were placed on Kapton tape and metal distribution was examined by synchrotron XRF. B, metal distribution in dissected seed parts. Seeds were imbibed for one day and dissected to isolate embryos and lateral or chalazal endosperm. Seed parts were dried, and metal concentrations were determined using ICP-OES. Error bars represent SEM. Different letters indicate that the means of the seed parts for that metal significantly differ from the other according to one-way ANOVA followed by a Tukey's honestly significant difference test ($P<0.05$).

Figure 4: Iron facilitates germination of tomato seeds without inducing hydroxyl radicals. A, impact of chelators desferoxamine (Des)(100 µM) and nicotianamine (NA)(5 µM) on germination speed of wild type seeds. B, impact of the same chelators on germination speed of *chloronerva* mutants that contain low chalazal Fe. For A and B, Each treatment consisted of three plates, each containing at least thirty seeds (n=3). Error bars represent SEM. Different letters at the same time point indicate a significant difference determined by two-way ANOVA, followed by Tukey's test for individual time points ($P<0.05$). The experiment was repeated four times with comparable results. C, impact of Fe form on germination speed. Final concentration of each Fe form was set to 100 µM. D, impact of several micronutrients on germination speed. Final concentration of each micronutrient was set to 100 µM. E, impact of increasing Fe-EDTA concentration on germination speed. For C-E, fifty seeds for each treatment were assessed. F, effect of Fe availability on hydroxyl radical production. Five seeds were pooled as a biological replicate (n=3). Seeds were incubated in water containing a radioactive probe, BzK$_5$Me, which is reactive to hydroxyl radicals in the presence of Fe

(250 µM FeCl$_2$) or Fe chelators (5 µM Na or 100 µM Des). ^{3}H$_2$O release was considered as a proxy for hydroxyl radical production. Error bars represent SEM. The asterisk indicates that the corresponding mean of the treatment or negative control is significantly different from the mean of the control (Seeds imbibed without chelators or Fe) according to Student's t-test ($P<0.05$).