1 Improved sample preparation for fruits allowed histochemistry and X-ray
2 microscopy to reveal conserved iron hotspots

4 Seçkin Eroğlu[1*], Utku Deniz[1], Abdulsamet Sakalar[1], Katarina Vogel-Mikuš[2,3]
5
6 *1 Department of Biological Sciences, Middle East Technical University, Ankara, Turkiye*
7 *2 Institute Jozef Stefan, Jamova cesta 39, Ljubljana, Slovenia*
8 *3 Biotechnical Faculty, University of Ljubljana, Jamnikarjeva ulica 101, Ljubljana, Slovenia*
9
10 Corresponding Author: Seckin Eroglu, erogluseckin@gmail.com

11
12 Running title: Extending histochemical and X-ray-based approaches to the mature fleshy fruits
13
14 Keywords: fruit, stain, X-ray, metal, mature, perls, microscopy, spatial, iron
15
16

Abstract

18
Mature fleshy fruits are the most challenging plant organs to study due to their high water and
flavonoid content. We aim to develop a sample preparation protocol to analyze fleshy fruits
with histochemical and X-ray based methods. By vacuum freeze drying and decolorization with
fixatives, we were able to apply histochemical Perls staining to seamlessly localize iron in fruits.
We screened the fruits to reveal possible conserved iron accumulation sites. This approach
showed iron preferentially accumulated in fruit vascular tissues; at subcellular level, in the cell
walls of vasculature and other tissues, and inside the epidermal cells of polyphenol-rich fruits.
Furthermore, iron accummulated in the endosperm of tomato seeds, indicating a role in
germination. X-ray microscopy of chemically untreated fruit slices showed similar iron
distributions, indicating chemicals used in the protocol did not significantly mobilized the
metals. This has been the first systematic study to extend histochemistry and X-ray fluorescence
to obtain spatial information from mature fleshy fruits. This advancement in methodology may
facilitate fruit research, potentially contributing to food security in the long run.

Introduction

35
Fruits constitute an indispensable component of a nutritious human diet, and to sustain the
dietary needs of our burgeoning global population, fruit production must escalate from one
billion tons to surpass two billion tons by 2050 [1]–[3]. Part of this challenge is to reduce the
pre- and post-harvest losses, which can account for up to half of the production, putting fruits
on the top of most wasted among all kinds of foods [4], [5]. One major factor leading to such

crop losses is metal deficiencies. Metal deficiencies lead to losses by limiting the quality and
quantity of the fruits. These effects can be direct, as in apples, where insufficient calcium levels
can reach the fruit, leading to the bitter pit disease Field [6]–[8]. The effects can also be indirect
(i.e., systemic); for example, starving for a metal can induce remodeling of the metabolism,
leading to lower quantity and quality of fruits [9]–[11]. Metal deficiencies may also reduce
fruits' defense against several microbial diseases [12], [13], which are associated with a
significant part of the post-harvest losses.
48
Despite the well-established importance of metals in fruit biology even the most basic protocols
for their investigation are notably absent. Main methods to investigate spatial localization of
metals in plant organs such as seeds, roots or leaves include X-ray based techniques [14], [15],
histochemical stains and others [16]–[18] prove inconvenient for fleshy fruits due to fruits' very
high water content. This water content prevents the use of many sample preparation
techniques; fixatives may not effectively replace water, water loss may spoil the physical
structure, and water-filled soft tissues might require expertise to cut to get uniform sections. In
addition to being mostly water, fruits are enriched in colorful polyphenols, obscuring the color
of simple histochemical stains. Therefore, spatial metal analysis of fruits is currently limited to
non fleshy fruits -such as grains or pods- and does not extend to juicy and fleshy ones such as
tomatoes or oranges- with only a few exceptions [19]. Consequently, the current practice of
spatially localizing metals in fruits must be improved. Currently, investigators take samples from
different parts of the fruit and compare the total metal concentration in them, providing
extremely low spatial information [20]–[22].
63
Lacking suitable techniques for localizing metals in fleshy fruits may have consequences beyond
the farmers' perspective. From the public's view, consumers sometimes choose one part of the
food over other parts due to its nutritional value. They should know how metal nutrients are
distributed among their everyday fruits, provided that consuming nutrient-rich parts can
alleviate widespread micronutrient deficiency disorders in people [23]. To illustrate the
importance of this choice, how people eat eggs can be given as an example. People doing
exercise often choose to eat the protein-rich white part, while, vitamin-seekers choose the
yellow, and many others choose to eat the membrane beneath the egg shells for therapeutic
effects; consequently, each of these parts is commercialized and available seperately[24], [25].
From the plant physiologists' view, detecting preferential accumulation of metals is the first step
to investigate their biological roles following the principle "form follows function." For example,
the accumulation of metals may indicate the presence of enzymes that use these metals as
cofactors and of metal transporter proteins mediating this accumulation.
77
78 We tested several tissue preparation protocols to allow spatial localization of metals by
79 histochemical staining in fleshy fruits. We aimed to employ such a protocol to seamlessly detect
80 metal distribution at tissue and subcellular levels. This approach allowed us to discover hot
81 spots for metals, some surprisingly widely conserved.
82
83

Method

Fruits were sliced and freeze dried. These fruit slices then used for metal analyses either directly (for X-ray analyses) or after decolorization (Perls and Perls/DAB stainings).

Fruits were purchased from local markets and bazaars in Turkey (Supplementary table 1). Fruits were freshly sliced longitudinally and cross-sectionally with a V-blade in 2-3mm thickness (Supplementary Figure 1). Prepared fruit slices were frozen in -80 °C for 24 hours. Frozen fruit samples were placed in a vacuum freeze dryer for 23 hours of the main drying phase at -54 °C, 0.024 millibar, and an hour of final drying phase at -60 °C, 0.011 millibars. These samples were used directly for X-ray analysis.

For X-Ray analysis, freeze-dried fruit samples were analyzed by an micro XRF analyzer (Horiba, Japan) with step size ranging from 120 to 200 μm with 15μm polycapillary beam.

For decolorization, freeze-dried samples were immersed in a fixing solution (Methanol: Chloroform: Glacial acetic acid; 6:3:1) in a glass petri dish and were shaken at 90 rpm until the fruit's natural color disappeared. The fixing solution was renewed several times as the fruit piece lost color. To stop the decolorization step, the fixative was removed by washing the samples three times with distilled water.

For the histochemical staining of Fe, Perls staining was used as described before [26]. Briefly, Perls stain solution was prepared fresh by mixing 4% K-ferrocyanide ($K_4Fe(Cn)_6$) and 4% HCl stock solutions in a 1:1 (v: v) proportion. Then, it was braught to 37 °C in a standard incubator. Decolorized samples were submerged in the solution and vacuum infiltrated for an hour at 500 millibars. Next, the staining solution was removed, and the samples were washed three times with distilled water for 1-2 min each by slightly shaking. The samples were stored in distilled water at 4°C. The stained samples were observed under a stereoscopic microscope.

To examine the subcellular localization of Fe, Perls/DAB staining was performed on thin sections. For this, freeze-dried fruit slices were embedded in paraffin, and 10 μm sections were obtained. Sections were then deparaffinized by incubation at 65°C for 15 minutes following a xylene treatment for 15 minutes and stained with Perls/DAB according to Roschzttardtz et al. (2009). Briefly, Perl's stain solution was prepared fresh by mixing 4% K-ferrocyanide ($K_4Fe(Cn)_6$) and 4% HCl stock solutions in a 1:1 (v: v) proportion. Then, the stained sections were incubated between 10 and 30 min in a 0.1 M phosphate buffer (pH 7.4) solution containing 0.025% (w/v) DAB (Sigma), 0.005% (v/v) H_2O_2, and 0.005% (w/v) $CoCl_2$. The reaction was stopped by rinsing samples with distilled water. Samples were imaged with a light microscope (Axioskop; Carl Zeiss, Jena, Germany).

RESULTS

For metal localization studies, arguably, the most widely used plant organ can be the seed.
Generally, this is attributed to seeds' low (i.e., around %5-15) water content. We stained tomato
fruit with Perls staining to examine how effective classical histochemical methods are in
localizing metals in fruits. In literature, Perls turn to a bright blue in the presence of Fe and have
been widely used for localizing Fe in various plant organs [27]. Being %95 water, tomato is one
of the most prominent agricultural crop [28], making it a challenging target for metal
localization studies. As expected, lentil seeds that were used as controls were heavily stained by
Perls, producing more or less homogeneously distributed blue color, indicating the stain is
working (Figure 1A). However, tomato fruits show very little or no coloration(Figure 1A). This
indicated either the fruit lacked detectable Fe or Perls could not localize Fe in fresh tomato
fruits. We hypothesized that the high water content of tomato inhibits Perls dye's action, and
the bright red color of tomato fruit further obscures the presence of blue precipitates of Perls.
To remove excess water, we first tried conventional drying methods. Oven drying spoiled the
structure of the tomato(Figure 1B). However, when dried tomato slices were stained by Perls,
we noticed more blue coloration (Figure 1B). Compared to the non-dried and stained tomato
slices, indicating high water content was decreasing staining efficiency. We treated the tomato
fruit with several solvents aiming to remove the colorful polyphenols while preventing the
remobilization of Fe as much as possible. However, treating tomato fruits with solvents led to
loss of rigidity, and fruit slices looked mushy (Figure 1B). We saw a similar trend with other
fruits; when treated with solvents, they lost structural integrity, and developed brown
discoloration (Supplementary Figure 2).
149
We hypothesized that the solvents we used were ineffective due to the high water content of
the tomato fruit. To remove water while maintaining the structural integrity of tomato slices, we
dried the slices using a vacuum freeze drier this time. Vacuum freeze drier kept the structure of
tomato slices(Figure 2A). As expected, when treated with solvents, these dried tomato slices
effectively lost most of their natural color. Among the solvents, we decided to use Methanol:
Chloroform: Glacial acetic acid; 6:3:1 since it effectively removed colors from tomato fruit and at the
same time had been succesfully used as a fixative in Perls staining [29], indicating this mixture should
keep metal remobilization at minimal. When stained by Perls, freeze-dried and decolorized tomato slices
showed intensive blue staining (Figure 2B-G). This blue staining was not just homogeneously distributed
but preferentially localized in the region below the calyx (Figure 2C), in the vasculature (Figure 2D), and
around the seeds (Figure 2E). Seeds of various species had been extensively investigated for metal
distribution before [27], [30]; however, tomato seed remained elusive. To investigate whether Fe staining
around the seed was in the outer or the inner tissues, we made 10 µm cross and longitudinal sections
and stained with Perls. Results showed tomato seeds accumulate Fe in the embryo and in the
endosperm. Embryo Fe was preferentially localized in the vasculature (Figure 2F), in agreement with the
previous data [27]. However, Fe also observed in the endosperm (Figure 2F, G). Endosperm Fe localized
preferentially in the region close to the germination site, including chalazal and micropyler, but not
lateral endosperm. This novel observation may indicate Fe may be involved in the germination of tomato
seeds.
169
Next, we assessed whether the protocol we developed was only effective for localizing metals in
tomato fruits or could be used for other fruits as well. We employed the technique on several
fruits and determined the Fe localizations (Figure 3 and 4). Results showed Fe accumulation

patterns successfully. Many fruits showed more or less homogeneous blue color (Figure 3).
However, we noticed many fruits showed an intense blue color throughout their vasculature
tissues, indicating a preferential Fe accumulatin in this tissue (Figure 3). We also noticed fruits
that were known to accumulate extensive amounts of proanthocyanidins in the fruit skin
showed preferential Fe accumulation in their skin (for example, blueberry in Figure 4H).
178
Perls staining can show Fe localization even in the subcellular level [26]. However, to reach
subcellular resolution, thin sections are required. At this level, total Fe in the tissue is often
insufficient to produce a signal. For such cases, Perls staining had been intensified by DAB [26].
To investigate Fe localization at the subcellular level, we made thin cross-sections of vacuum
freeze-dried fruits slices. Perls staining following DAB intensification revealed the Fe distribution
as black coloration. According to the results, most Fe resided outer parts of the cells, probably
in organelles at the peripheries such as chloroplasts, chromoplasts, and mitochondria of the cell
and in the cell wall (Figure 5). In contrast, Fe was found in inner regions, inside the vacuoles in
the epidermis of blueberry (Figure 5A). In accordance with Perls staining of the slides, vascular
tissues were heavily stained with Perls/DAB (Figure 5B,C).
189
Histochemical methods to localize metals have often been criticized for possible artifacts (i.e.,
remobilization of metals). To verify the Fe hotspots recently identified in the fruits with an
independent technique, we used X-ray-based techniques where tissue preparation was
minimal. Micro XRF mapping of vacuum freeze-dried slices revealed localizations of many
metals in fruit slices cut horizontally or vertically (Figure 6). Next, we assessed whether Micro
XRF can validate tissue level Fe hotspots detected by histochemical methods. Scanning of
vacuum freeze-dried slices of fruits confirmed the Fe hotspots in the vascular tissues of selected
fruits(Figure 7A, B) and in the skin of blueberries (Figure 7C).
198
199
200

201 # Discussion

202
Fruits, especially the fleshy ones, are the fastest deteriorating part of plants upon harvest,
posing a unique challenge to meet the ever increasing global food demand. Fruit spoilage
should be highly related to metals, as metals activate key enzymatic processes such as those
involved in fruit ripening/softening and pathogen defense. To examine metal-involved
physiological processes in fruits, the very first step is to be able to localize the metals in fruit
tissues, preferably, with low cost and fast methods offering high resolution at the same time. In
this study we developed a tissue preparation method that can be applied to even most complex
(i.e., very high water and colorful polyphenol content) fruits and showed that this allowed fruits
to be analyzed by the common metal localization methods, including histochemical staining or
micro X-Ray mapping.
213
Histochemical approach to mature fleshy fruits often face two main obstacles: too much of
water content limiting penetration of chemicals and too much of colors obscuring the chemical

staining (Figure 1A). Although fruits are extensively dried in industry, such tecnologies have
been rarely used for research. Among the drying technologies, vacuum freeze drying offers the
golden standard for human consumption. It is used on freshly cut fruits to increase fruits' shelf
life and simplify their transportation to the market. Freeze drying prevents shrinkage of the
slices, minimizes remobilization of solutes in the tissues [31]. It relies on evaporation directly
from the ice phase, skipping the water phase which is the most prone phase for chemical
reactions and remobilization of solutes. After the evaporation porous tissue remains, perfect for
penetration of chemicals for histochemical approach, despite these advantages, vacuum freeze
drying almost never appears in histochemical methods in fruit research.
225
We observed that once freed from water, the fruit slices behaved as regular tissue (Figure 2).
After vacuum freeze drying, fixatives could decolorize the fruit (Figure 2 and Supplementary
figure 2). Although part of the color was retained(Figure 2A), it did not interfere with the
detection of our stain anymore. In literature, the most common getaround from the natural
bright colors of fruits such as tomato has been to analyze the sample in earlier developmental
stages before the fruit develops colors[32], [33]. This approach is in parallel with histochemistry
applied to seeds. Seed coats develop a dark color due to accumulated proanthocyanins, forcing
researchers to do histochemistry such as GUS staining before the fruit matures or the detection
quality suffers [32], [33]. For fruits, we tested only one type of histochemical stain; however, the
protocol potentially allowed other histochemical methods to be applied.
236
We observed the most prominent Fe accumulation in the vasculature, cell walls, then tissues
around seeds, and sometimes in the skin of the fruits (Figure 4, 5). Regarding the vasculature,
the physiological function of this Fe accumulation is unclear. The physiological functions of
metals have often been studied by imposing their deficiency and investigating the developing
phenotype. For example, Fe deficiency induces chlorosis in the leaves, indicating Fe may be
involved in chlorophyll synthesis. However, Fe deficiency in fruit trees has never been reported
to cause a disease specific to vasculature[9]. However, we cannot rule out a possible
physiological function for this Fe reservoir. Recently, Fe oxidation was linked with the callose
deposition [34], which may decrease xylem functionality in the developing fruits [35]. Another
possibility is that the main vasculatures of fruits may serve as a sink to load Fe to the seeds. In
this model, the vasculature may represent excess Fe reservoir left over from seed Fe filling, as
this process is highly regulated not to induce Fe-mediated oxidative damage in the precious
seeds [36]. The remaining Fe can then be absorbed by the cell walls, as shown cell walls can
quickly absorb Fe(III) [37]. Our study identified vasculature as the main Fe-accumulating tissue;
however, further investigation on which cells in the vasculature accumulate iron is yet to be
investigated.
253
Fe accumulation in the cell walls has been well documented in other organs [37]–[39] but has
not yet been discussed regarding the cell walls of fruits. We propose that the cell walls of fruits
may act as a larger reservoir for metal ions such as Fe due to two main factors. First, among all
the plant cell walls, the cell walls of fruits potentially have the highest concentration of pectin,
the major component of the cell wall that binds Fe [38], [40]. Second, the cell walls of fruits may
be less saturated with cations than other organs' cell walls. Demethylated pectins in cell walls

typically offer free carboxyl groups to absorb positively charged metals, notably calcium, a far
more abundant element than Fe, Mn, and Zn. However, calcium is almost solely transported by
the xylem, which is ineffective in fruits due to low transpiration [35]. Consequently, its low
concentrations commonly cause diseases in fruits such as apples or tomatoes [41], [42]. On the
other hand, all the rest of the cations are phloem mobile, meaning they are transported to the
fruit continuously [43]. Positively charged molecules compete for binding the free carboxyl
residues of pectins at least in the root; for example, in the field, sodium mobilizes Fe from the
cell wall [44] or protons secreted from H+ ATPases mobilize the calcium [45]. We propose that
in fruits, due to high pectin and low calcium concentration, cell walls may act as an unusually
powerful absorbent for metal ions, explaining our observation of conserved Fe accumulation
(Figure 4). Fe and maybe other metals unloaded/leaked from the phloem and xylem may
immediately get stuck in the cell walls of helper cells in the vasculature.

We also detected iron inside the outer cell layers of blueberry or grapes and in the endosperm
of tomato (Figure 5 and 7). These iron spots may represent tannins, as blueberry skin can often
be rich in tannins that bind iron and may be stored in the vacuoles [46]–[48]. In literature, an
increase in fruit polyphenols, including tannins was reported to be correlated with higher Fe
concentrations [49]. Besides, we detected Fe in tomato seed endosperm (Figure 2). In seeds,
main Fe storage is in the embryo, but exceptions such as seeds accumulating Fe in the
endosperm have been reported [27]. However, Fe storage in tomato endosperm shows an
evident preferential accumulation to the sides close to the germination (i.e., micropylar and
chalazal regions). Therefore, we propose that this Fe storage may facilitate germination. For
tomato seeds to germinate, endosperm should first soften to allow radical protrusion, a major
process shown to include cell wall cutting hydroxyl radicals [50], [51]. A well-known source of
hydroxyl radicals is Fe, which reacts with H_2O_2 through the Fenton reaction [52]. Therefore, this
Fe storage in the endosperm may facilitate germination by generating a cell wall attacking
hydroxyl radicals. Further studies should target whether manipulating Fe levels in seeds can
change germination patterns.

Increasing fruit production should target decreasing pre- and post-harvest losses. To reduce the
loss, a better understanding of fruit biology is required. We developed a sample preparation
technique for even the most difficult fruits. By using it, we systematically examined Fe
localizations in the fruits. We determined some Fe hotspots which require further investigation
for their potential physiological function. This study extended the basic and applied research to
mature fleshy fruits, potentially facilitating research to prevent yield loss in the long run.

Figures and captions

Figure 1: The histochemical approach to fruits requires a new protocol for sample preparation. A, Effectiveness of direct staining of seeds compared with fleshy fruits. Lentil seed coat was removed, and lentil embryos were submerged in Perls staining solution for two hours. Cut tomato pieces were immersed directly in Perls solution overnight. B, Common obstacles for histochemical approach on fruits include high water content and bright colors. Left, a tomato slice was dried in an oven; simple drying led to structure loss. Middle, dried tomato ineffectively stained by Perls. The oven-dried tomato slice was incubated in Perls solution for two hours. Right, fixatives decolorized the tomato but led to structure loss. A freshly cut tomato slice was treated with fixatives (Methanol: Chloroform: Glacial acetic acid; 6:3:1) overnight with continuous shaking. Blue coloration shows Fe. Bar 1cm unless otherwise indicated.

Figure 2: A newly developed protocol allowed histochemical methods to apply to tomatoes and revealed preferential Fe accumulation patterns. A, freeze-dried, decolorized tomato slice. B, freeze-dried, decolorized, and stained tomato slices. C-E, close-up pictures of B. F, G; stained thin cross sections of the tomato slice. For A, freshly cut tomato pieces were first freeze-dried and then decolorized using fixatives (Methanol: Chloroform: Glacial acetic acid; 6:3:1). For B, after fixation, pieces were submerged in Perls solution for 2-4 hours. To see the staining inside the seeds, freshly cut tomato pieces were first freeze-dried and then decolorized using fixatives (Methanol: Chloroform: Glacial acetic acid; 6:3:1). For F, G; after fixation, samples were embedded in wax, and 10 µm cuts were prepared on slides, following Perls staining. Blue coloration shows iron. Pink arrows point to blue coloration. Bar 1cm unless otherwise indicated. Cal: calyx detachment site, VB: vascular bundle, M: micropylar region, Cha: chalazal region, C: cotyledon, R: radicle

Figure 3: **The newly developed protocol revealed Fe accumulation in various fruits.** Fruits were either cut cross-sectional or longitudinal with a V- blade (For the V-blade, refer to supplementary figure 1). These pieces were then freeze-dried and decolorized using fixatives (Methanol: Chloroform: Glacial acetic acid; 6:3:1). After fixation, pieces were submerged in Perls' staining solution between 2-16 hours. From A to M; orange, mandarin, lemon, grapefruit, apple, persimmon, honey melon, mango, pumpkin, plantain, nectarin, cape gooseberry, avakado. Note that despite the differences in fruit types, the optimized protocol preserved the structure and decolorized the tissues efficiently. For the complete list of fruits successfully stained, refer to supplementary table 1. Blue coloration shows Fe. Bar 1cm unless otherwise indicated.

Figure 4: Fe accumulation was detected in the main vasculatures of fruits. Fruits were either cut cross-sectional or longitudinal with a V- blade(For the V-blade, refer to supplementary figure 1). These pieces were then freeze-dried and decolorized using fixatives (Methanol: Chloroform: Glacial acetic acid; 6:3:1). After fixation, pieces were submerged in Perls' staining solution between 2-16 hours. For this figure, fruit pictures with apparent Fe accumulation in the vascular bundles were chosen. From A to K; eggplant, cucumber, kiwi, kumquat, haricot, squash, green almond, blueberry, papaya, pepper, passion fruit. Bar 1cm, unless otherwise indicated.

Figure 5: Fe accumulation in fruits could be detected at the subcellular level. A, Fe accumulated inside the epidermal cells of blueberry, most probably bound to polyphenols in the vacuole. Left, control section without staining; middle, Perls/DAB stained section; right, close up of red rectangle of the middle. B, Fe accumulates in cell walls. Left, control section without staining; right, Perls/DAB stained section. C, Fe heavily accumulated in the back of the squash seed embryo, most probably corresponding to the vasculature as Cucurbitacea seeds, including squash, have a coat usually surrounded by a single vein [53]. Left, squash cross section stained with Perls; middle thin section corresponding to the back part of the seed (such as the pink rectangle shown in the left picture); middle, control section without staining; right, Perls/DAB stained section. Blue coloration in C (left) and black coloration in the rest shows Fe accumulation. Pink arrows point to Fe accumulation. VB: Vascular bundle, V: Vasculature, S: Seed

Figure 6: Determining metal localizations on fruit slices by micro XRF mapping. Cross and longitudinal section of Kumquat. The sample was cut, freeze-dried, and analyzed by XRF.

Figure 7: Micro XRF mapping confirmed the histochemical methods on tissue-level Fe hotspots. A, aubergine; B, squash; C, blueberry. Pink arrows point to Fe accumulations. VB: vascular bundle

Graphical abstract, a low-cost, high-throughput protocol to map metal localizations in everyday fruits.

References

[1] W. Willett *et al.*, "Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems," *The lancet*, vol. 393, no. 10170, pp. 447–492, 2019.

[2] A. E. Stratton *et al.*, "Mitigating sustainability tradeoffs as global fruit and vegetable systems expand to meet dietary recommendations," *Environ. Res. Lett.*, vol. 16, no. 5, p. 055010, May 2021, doi: 10.1088/1748-9326/abe25a.

[3] F. A. O. Fruit and V.-Y. D. Essentials, "The International Year of Fruits and Vegetables, 2021," *Background Paper*, 2020.

[4] J. Gustavsson, C. Cederberg, U. Sonesson, R. Van Otterdijk, and A. Meybeck, "Global food losses and food waste." FAO Rome, 2011. Accessed: Nov. 04, 2023. [Online]. Available: https://www.madr.ro/docs/ind-alimentara/risipa_alimentara/presentation_food_waste.pdf

[5] M. Wang, Y. Xu, Y. Yang, B. Mu, M. A. Nikitina, and X. Xiao, "Vis/NIR optical biosensors applications for fruit monitoring," *Biosensors and Bioelectronics: X*, vol. 11, p. 100197, Sep. 2022, doi: 10.1016/j.biosx.2022.100197.

[6] B. J. Van Goor, "The Effect of Frequent Spraying with Calcium Nitrate Solutions on the Mineral Composition and the Occurrence of Bitter Pit of the Apple Cox's Orange Pippin," *Journal of Horticultural Science*, vol. 46, no. 4, pp. 347–364, Jan. 1971, doi: 10.1080/00221589.1971.11514415.

[7] E. Torres, S. Alegre, I. Recasens, L. Asín, and J. Lordan, "Integral procedure to predict bitter pit in 'Golden Smoothee'apples based on calcium content and symptom induction," *Scientia Horticulturae*, vol. 277, p. 109829, 2021.

[8] W. Song, J. Yi, O. F. Kurniadinata, H. Wang, and X. Huang, "Linking fruit Ca uptake capacity to fruit growth and pedicel anatomy, a cross-species study," *Frontiers in plant science*, vol. 9, p. 575, 2018.

[9] A. Àlvarez-Fernàndez, J. Abadía, and A. Abadía, "Iron deficiency, fruit yield and fruit quality," *Iron nutrition in plants and rhizospheric microorganisms*, pp. 85–101, 2006.

[10] A. Álvarez-Fernández, P. Paniagua, J. Abadía, and A. Abadía, "Effects of Fe Deficiency Chlorosis on Yield and Fruit Quality in Peach (*Prunus persica* L. Batsch)," *J. Agric. Food Chem.*, vol. 51, no. 19, pp. 5738–5744, Sep. 2003, doi: 10.1021/jf034402c.

[11] S. F. El-Gioushy *et al.*, "Foliar application of nano, chelated, and conventional iron forms enhanced growth, nutritional status, fruiting aspects, and fruit quality of Washington navel orange trees (Citrus sinensis L. Osbeck)," *Plants*, vol. 10, no. 12, p. 2577, 2021.

[12] F. Morina and H. Küpper, "Trace metals at the frontline of pathogen defence responses in non-hyperaccumulating plants," *Journal of Experimental Botany*, vol. 73, no. 19, pp. 6516–6524, Nov. 2022, doi: 10.1093/jxb/erac316.

[13] W. S. Conway, C. E. Sams, and K. D. Hickey, "Pre-and postharvest calcium treatment of apple fruit and its effect on quality," in *International Symposium on Foliar Nutrition of Perennial Fruit Plants 594*, 2001, pp. 413–419. Accessed: Nov. 30, 2023. [Online]. Available: https://www.actahort.org/books/594/594_53.htm

[14] B. Wu and J. S. Becker, "Imaging techniques for elements and element species in plant science," *Metallomics*, vol. 4, no. 5, p. 403, 2012, doi: 10.1039/c2mt00002d.

[15] A. Mijovilovich, F. Morina, S. N. Bokhari, T. Wolff, and H. Küpper, "Analysis of trace metal distribution in plants with lab-based microscopic X-ray fluorescence imaging," *Plant Methods*, vol. 16, no. 1, p. 82, Dec. 2020, doi: 10.1186/s13007-020-00621-5.

[16] F.-J. Zhao, K. L. Moore, E. Lombi, and Y.-G. Zhu, "Imaging element distribution and speciation in plant cells," *Trends in Plant Science*, vol. 19, no. 3, pp. 183–192, 2014, doi: 10.1016/j.tplants.2013.12.001.

[17] E. Lombi, K. G. Scheckel, and I. M. Kempson, "In situ analysis of metal(loid)s in plants: State of the art and artefacts," *Environmental and Experimental Botany*, vol. 72, no. 1, pp. 3–17, Aug. 2011, doi: 10.1016/j.envexpbot.2010.04.005.

[18] V. Yadav *et al.*, "Histochemical Techniques in Plant Science: More Than Meets the Eye," *Plant and Cell Physiology*, vol. 62, no. 10, pp. 1509–1527, Oct. 2021, doi: 10.1093/pcp/pcab022.

[19] J. Trebolazabala, M. Maguregui, H. Morillas, A. de Diego, and J. M. Madariaga, "Evaluation of metals distribution in Solanum lycopersicum plants located in a coastal environment using micro-energy dispersive X-ray fluorescence imaging," *Microchemical Journal*, vol. 131, pp. 137–144, 2017.

[20] A. A. Saquet, J. Streif, and D. P. Almeida, "Mineral composition and distribution within 'Rocha'pear in relation to internal storage disorders," *Postharvest Biology and Technology*, vol. 158, p. 111002, 2019.

[21] L. A. Kalcsits, "Non-destructive measurement of calcium and potassium in apple and pear using handheld X-ray fluorescence," *Frontiers in plant science*, vol. 7, p. 442, 2016.

[22] K. V. Ramesh, V. Paul, and R. Pandey, "Dynamics of mineral nutrients in tomato (Solanum lycopersicum L.) fruits during ripening: part I—on the plant," *Plant Physiol. Rep.*, vol. 26, no. 1, pp. 23–37, Mar. 2021, doi: 10.1007/s40502-020-00546-0.

[23] R. L. Bailey, K. P. West Jr, and R. E. Black, "The epidemiology of global micronutrient deficiencies," *Annals of nutrition and metabolism*, vol. 66, no. Suppl. 2, pp. 22–33, 2015.

[24] A. M. King'Ori, "A review of the uses of poultry eggshells and shell membranes," *International Journal of Poultry Science*, vol. 10, no. 11, pp. 908–912, 2011.

[25] C. MM Cordeiro and M. T Hincke, "Recent patents on eggshell: shell and membrane applications," *Recent patents on food, nutrition & agriculture*, vol. 3, no. 1, pp. 1–8, 2011.

[26] H. Roschzttardtz, G. Conéjéro, C. Curie, and S. Mari, "Identification of the Endodermal Vacuole as the Iron Storage Compartment in the Arabidopsis Embryo," *Plant Physiol.*, vol. 151, no. 3, pp. 1329–1338, Nov. 2009, doi: 10.1104/pp.109.144444.

[27] S. Eroglu, N. Karaca, K. Vogel-Mikus, A. Kavčič, E. Filiz, and B. Tanyolac, "The conservation of VIT1-dependent iron distribution in seeds," *Frontiers in plant science*, vol. 10, p. 907, 2019.

[28] L. Zhiguo, L. Pingping, and L. Jizhan, "Effect of mechanical damage on mass loss and water content in tomato fruits," *International Agrophysics*, vol. 25, no. 1, 2011, Accessed: Nov. 19, 2023. [Online]. Available: http://agro.icm.edu.pl/agro/element/bwmeta1.element.agro-010e2c1d-4048-4d1b-a146-a171a44147a3

[29] T. Brumbarova and R. Ivanov, "Perls staining for histochemical detection of iron in plant samples," *Bio-protocol*, vol. 4, no. 18, pp. e1245–e1245, 2014.

[30] M. A. Ibeas, S. Grant-Grant, N. Navarro, M. F. Perez, and H. Roschzttardtz, "Dynamic Subcellular Localization of Iron during Embryo Development in Brassicaceae Seeds," *Front. Plant Sci.*, vol. 8, no. 2186, pp. 1–8, 2017, doi: 10.3389/fpls.2017.02186.

[31] J. C. Harper and A. L. Tappel, "Freeze-drying of food products," *Advances in Food Research*, vol. 7, pp. 171–234, 1957.

[32] R. Saed Taha, I. Ismail, Z. Zainal, and S. N. A. Abdullah, "The stearoyl-acyl-carrier-protein desaturase promoter (Des) from oil palm confers fruit-specific GUS expression in transgenic tomato," *Journal of Plant Physiology*, vol. 169, no. 13, pp. 1290–1300, Sep. 2012, doi: 10.1016/j.jplph.2012.05.001.

[33] S. Wang *et al.*, "Members of the tomato FRUITFULL MADS-box family regulate style abscission and fruit ripening," *Journal of Experimental Botany*, vol. 65, no. 12, pp. 3005–3014, Jul. 2014, doi: 10.1093/jxb/eru137.

[34] J. Müller *et al.*, "Iron-dependent callose deposition adjusts root meristem maintenance to phosphate availability," *Developmental cell*, vol. 33, no. 2, pp. 216–230, 2015.

[35] T. Knipfer, J. Fei, G. A. Gambetta, A. J. McElrone, K. A. Shackel, and M. A. Matthews, "Water transport properties of the grape pedicel during fruit development: insights into xylem anatomy and function using microtomography," *Plant Physiology*, vol. 168, no. 4, pp. 1590–1602, 2015.

[36] L. Sun *et al.*, "Restriction of iron loading into developing seeds by a YABBY transcription factor safeguards successful reproduction in Arabidopsis," *Molecular Plant*, vol. 14, no. 10, pp. 1624–1639, 2021.

[37] Z.-R. Xu *et al.*, "The ferroxidases LPR1 and LPR2 control iron translocation in the xylem of Arabidopsis plants," *Molecular Plant*, vol. 15, no. 12, pp. 1962–1975, 2022.

[38] J.-S. Peng *et al.*, "Galactosylation of rhamnogalacturonan-II for cell wall pectin biosynthesis is critical for root apoplastic iron reallocation in Arabidopsis," *Molecular Plant*, vol. 14, no. 10, pp. 1640–1651, 2021.

[39] S. Clemens, "(Not) an elementary question: How to mobilize the iron in the wall?," *Molecular Plant*, vol. 14, no. 10, pp. 1610–1611, 2021.

[40] A. G. J. Voragen, G.-J. Coenen, R. P. Verhoef, and H. A. Schols, "Pectin, a versatile polysaccharide present in plant cell walls," *Struct Chem*, vol. 20, no. 2, pp. 263–275, Apr. 2009, doi: 10.1007/s11224-009-9442-z.

[41] K. Suzuki, M. Shono, and Y. Egawa, "Localization of calcium in the pericarp cells of tomato fruits during the development of blossom-end rot," *Protoplasma*, vol. 222, no. 3–4, pp. 149–156, Dec. 2003, doi: 10.1007/s00709-003-0018-2.

[42] B. Hocking, S. D. Tyerman, R. A. Burton, and M. Gilliham, "Fruit calcium: transport and physiology," *Frontiers in plant science*, vol. 7, p. 569, 2016.

[43] P. J. White and G. Ding, "Chapter 3 - Long-distance transport in the xylem and phloem⌷⌷This chapter is a revision of the third edition chapter by P.J. White, pp. 49–70. DOI: https://doi.org/10.1016/B978-0-12-384905-2.00003-0. © Elsevier Ltd.," in *Marschner's Mineral Nutrition of Plants (Fourth Edition)*, Z. Rengel, I. Cakmak, and P. J. White, Eds., San Diego: Academic Press, 2023, pp. 73–104. doi: 10.1016/B978-0-12-819773-8.00002-2.

[44] X. F. Zhu, Q. Wu, L. Zheng, and R. F. Shen, "NaCl alleviates iron deficiency through facilitating root cell wall iron reutilization and its translocation to the shoot in Arabidopsis

thaliana," *Plant Soil*, vol. 417, no. 1, pp. 155–167, Aug. 2017, doi: 10.1007/s11104-017-3248-3.

[45] S. S. Virk and R. E. Cleland, "Calcium and the mechanical properties of soybean hypocotyl cell walls: Possible role of calcium and protons in cell-wall loosening," *Planta*, vol. 176, no. 1, pp. 60–67, 1988, doi: 10.1007/BF00392480.

[46] M. He, H. Tian, X. Luo, X. Qi, and X. Chen, "Molecular Progress in Research on Fruit Astringency," *Molecules*, vol. 20, no. 1, Art. no. 1, Jan. 2015, doi: 10.3390/molecules20011434.

[47] S. Gagné, C. Saucier, and L. Gény, "Composition and Cellular Localization of Tannins in Cabernet Sauvignon Skins during Growth," *J. Agric. Food Chem.*, vol. 54, no. 25, pp. 9465–9471, Dec. 2006, doi: 10.1021/jf061946g.

[48] L. Michel, Á. Peña, C. Pastenes, P. Berríos, A. D. Rombolà, and J. I. Covarrubias, "Sustainable Strategies to Prevent Iron Deficiency, Improve Yield and Berry Composition in Blueberry (Vaccinium spp.)," *Frontiers in Plant Science*, vol. 10, 2019, Accessed: Dec. 02, 2023. [Online]. Available: https://www.frontiersin.org/articles/10.3389/fpls.2019.00255

[49] C. C. Pereira, E. do Nascimento da Silva, A. O. de Souza, M. A. Vieira, A. S. Ribeiro, and S. Cadore, "Evaluation of the bioaccessibility of minerals from blackberries, raspberries, blueberries and strawberries," *Journal of Food Composition and Analysis*, vol. 68, pp. 73–78, May 2018, doi: 10.1016/j.jfca.2016.12.001.

[50] K. Muller, A. Linkies, R. A. Vreeburg, S. C. Fry, A. Krieger-Liszkay, and G. Leubner-Metzger, "In vivo cell wall loosening by hydroxyl radicals during cress seed germination and elongation growth," *Plant physiology*, vol. 150, no. 4, pp. 1855–1865, 2009.

[51] L. Grillet, S. Mari, and W. Schmidt, "Iron in seeds–loading pathways and subcellular localization," *Frontiers in plant science*, vol. 4, 2013, Accessed: Feb. 26, 2014. [Online]. Available: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3877777/

[52] M. J. Burkitt and B. C. Gilbert, "Model studies of the iron-catalysed Haber-Weiss cycle and the ascorbate-driven Fenton reaction," *Free radical research communications*, vol. 10, no. 4–5, pp. 265–280, 1990.

[53] D. Singh and A. S. R. Dathan, "18. Seed Coat Anatomy Of The Cucurbitaceae," in *Biology and Utilization of the Cucurbitaceae*, C. Jeffrey, Ed., Ithaca, NY: Cornell University Press, 2019, pp. 225–238. doi: 10.7591/9781501745447-020.

Figures and captions

Figure 1: The histochemical approach to fruits requires a new protocol for sample preparation. A, Effectiveness of direct staining of seeds compared with fleshy fruits. Lentil seed coat was removed, and lentil embryos were submerged in Perls staining solution for two hours. Cut tomato pieces were immersed directly in Perls solution overnight. B, Common obstacles for histochemical approach on fruits include high water content and bright colors. Left, a tomato slice was dried in an oven; simple drying led to structure loss. Middle, dried tomato ineffectively stained by Perls. The oven-dried tomato slice was incubated in Perls solution for two hours. Right, fixatives decolorized the tomato but led to structure loss. A freshly cut tomato slice was treated with fixatives (Methanol: Chloroform: Glacial acetic acid; 6:3:1) overnight with continuous shaking. Blue coloration shows Fe. Bar 1cm unless otherwise indicated.

Figure 2: A newly developed protocol allowed histochemical methods to apply to tomatoes and revealed preferential Fe accumulation patterns. A, freeze-dried, decolorized tomato slice. B, freeze-dried, decolorized, and stained tomato slices. C-E, close-up pictures of B. F, G; stained thin cross sections of the tomato slice. For A, freshly cut tomato pieces were first freeze-dried and then decolorized using fixatives (Methanol: Chloroform: Glacial acetic acid; 6:3:1). For B, after fixation, pieces were submerged in Perls solution for 2-4 hours. To see the staining inside the seeds, freshly cut tomato pieces were first freeze-dried and then decolorized using fixatives (Methanol: Chloroform: Glacial acetic acid; 6:3:1). For F, G; after fixation, samples were embedded in wax, and 10 μm cuts were prepared on slides, following Perls staining. Blue coloration shows iron. Pink arrows point to blue coloration. Bar 1cm unless otherwise indicated. Cal: calyx detachment site, VB: vascular bundle, M: micropylar region, Cha: chalazal region, C: cotyledon, R: radicle

Figure 3: **The newly developed protocol revealed Fe accumulation in various fruits.** Fruits were either cut cross-sectional or longitudinal with a V- blade (For the V-blade, refer to supplementary figure 1). These pieces were then freeze-dried and decolorized using fixatives (Methanol: Chloroform: Glacial acetic acid; 6:3:1). After fixation, pieces were submerged in Perls' staining solution between 2-16 hours. From A to M; orange, mandarin, lemon, grapefruit, apple, persimmon, honey melon, mango, pumpkin, plantain, nectarin, cape gooseberry, avakado. Note that despite the differences in fruit types, the optimized protocol preserved the structure and decolorized the tissues efficiently. For the complete list of fruits successfully stained, refer to supplementary table 1. Blue coloration shows Fe. Bar 1cm unless otherwise indicated.

Figure 4: Fe accumulation was detected in the main vasculatures of fruits. Fruits were either cut cross-sectional or longitudinal with a V- blade(For the V-blade, refer to supplementary figure 1). These pieces were then freeze-dried and decolorized using fixatives (Methanol: Chloroform: Glacial acetic acid; 6:3:1). After fixation, pieces were submerged in Perls' staining solution between 2-16 hours. For this figure, fruit pictures with apparent Fe accumulation in the vascular bundles were chosen. From A to K; eggplant, cucumber, kiwi, kumquat, haricot, squash, green almond, blueberry, papaya, pepper, passion fruit. Bar 1cm, unless otherwise indicated.

Figure 5: Fe accumulation in fruits could be detected at the subcellular level. A, Fe accumulated inside the epidermal cells of blueberry, most probably bound to polyphenols in the vacuole. Left, control section without staining; middle, Perls/DAB stained section; right, close up of red rectangle of the middle. B, Fe accumulates in cell walls. Left, control section without staining; right, Perls/DAB stained section. C, Fe heavily accumulated in the back of the squash seed embryo, most probably corresponding to the vasculature as Cucurbitacea seeds, including squash, have a coat usually surrounded by a single vein [53]. Left, squash cross section stained with Perls; middle thin section corresponding to the back part of the seed (such as the pink rectangle shown in the left picture); middle, control section without staining; right, Perls/DAB stained section. Blue coloration in C (left) and black coloration in the rest shows Fe accumulation. Pink arrows point to Fe accumulation. VB: Vascular bundle, V: Vasculature, S: Seed

Figure 6: Determining metal localizations on fruit slices by micro XRF mapping. Cross and longitudinal section of Kumquat. The sample was cut, freeze-dried, and analyzed by XRF.

Figure 7: Micro XRF mapping confirmed the histochemical methods on tissue-level Fe hotspots. A, aubergine; B, squash; C, blueberry. Pink arrows point to Fe accumulations. VB: vascular bundle

Graphical abstract, a low-cost, high-throughput protocol to map metal localizations in everyday fruits.