Java sparrow song conforms to Menzerath's Law but not Zipf's Law of Abbreviation

Rebecca N Lewis[1,2]*, Anthony Kwong[1,3], Masayo Soma[4], Selvino R de Kort[5], R Tucker Gilman[1]

[1]Department of Earth and Environmental Sciences, University of Manchester, Manchester, UK

[2]Chester Zoo, Upton-by-Chester, Chester, UK

[3]Department of Mathematics, University of Manchester, Manchester, UK

[4]Department of Biology, Faculty of Science, Hokkaido University, Hokkaido, Japan

[5]Department of Natural Sciences, Manchester Metropolitan University, Manchester, United Kingdom

*Corresponding author: r.lewis@chesterzoo.org

Abstract

Linguistic laws, which describe common statistical patterns in human language, are often studied in animal communication systems. Despite good evidence for adherence to these laws in non-human primate communication, there is still a paucity of data on other taxa. Birdsong is often used as a model for human language development due to similarities in vocal learning processes. Investigating whether birdsong also adheres to linguistic laws may provide further value or help determine the limitations of such comparisons. Here, we looked for patterns consistent with Menzerath's law, the tendency for longer sequences to be made up of smaller constituent parts, and Zipf's Law of Abbreviation, the tendency for more frequently used elements to be shorter in length, in the songs of Java sparrows (*Padda oryzivora*). We found that songs conformed to Menzerath's law at both the population and individual level; longer sequences were made up of shorter notes. This pattern was achieved through note type selection rather than the shortening of existing note types. Adherence to Menzerath's law may reflect species-specific constraints on song durations. We found no evidence to support adherence to Zipf's Law of Abbreviation, either at population or individual level; more frequently used notes were not shorter in length. However, as repertoire size in Java sparrows is small, we expect our power to detect Zipf's Law of Abbreviation to be weak. Whilst there may be

some limitations in applying linguistic laws derived from human language to birdsong, especially Zipf's

Law of Abbreviation, these laws may still provide insights into animal and human communication

systems.

Keywords

Animal communication; Birdsong; Java Sparrow; Language; Linguistic Laws; Menzerath's Law; *Padda*

oryzivora; Zipf's Law of Abbreviation

Introduction

Birdsong and spoken language share many similarities (Berwick et al., 2011; Bolhuis et al., 2010;

Hyland Bruno et al., 2021). Both are made up of acoustic sequences, and there is overlap in the

behavioural, neural, and genetic bases of vocal learning and production in songbirds and humans

(Berwick et al., 2011, 2012; Bolhuis et al., 2010; Hyland Bruno et al., 2021). Due to these similarities,

birdsong is often used in comparative studies examining the evolution and development of human

language (e.g., Berwick et al., 2011). However, there are also differences between birdsong and

human language. In particular, the close relationship between structure and meaning in human

language is thought not to exist in other animals (Bolhuis et al., 2010). Testing the limitations of the

parallels between human and animal communication may provide insight in the universal laws

affecting acoustic communication.

Linguistic laws describe common statistical patterns in human language, where they have been

studied extensively (Semple et al., 2022). More recently, patterns consistent with linguistic laws have

also been explored in animal communication systems (Semple et al., 2022). Two laws that often

receive attention are Menzerath's law, the tendency for longer sequences to be made up of smaller

constituent parts, and Zipf's Law of Abbreviation (ZLA), the tendency for more frequently used

elements to be shorter in length. In human languages, length is often equated with the number of

letters in a word, whereas for animal communication the duration (e.g., in seconds) of the element is

usually measured (REFS?). These laws are especially well studied in non-human primates (Bezerra et

al., 2011; Clink et al., 2020; Clink and Lau, 2020; Gustison et al., 2016; Huang et al., 2020; Semple et

al., 2010; Valente et al., 2021; Watson et al., 2020) and some other mammal taxa (Demartsev et al.,

2019; Ferrer-i-Cancho et al., 2013; Youngblood, 2025). Studies examining the presence of linguistic

laws in bird communication are still relatively limited (Favaro et al., 2020a; Ferrer-i-Cancho et al.,

2013; Hailman et al., 1985; James et al., 2021a; Wascher & Youngblood, 2025; Youngblood, 2024).

Adherence to Menzerath's law has been widely reported in non-human primates (Clink et al., 2020a;

Gustison et al., 2016; Heesen et al., 2019; Huang et al., 2020a; Valente et al., 2021). Despite this,

adherence to the law is not universal, and may depend on the subset of vocalizations examined; in

mountain gorillas, Menzerath's law was apparent when single-unit sequences were included in

analyses, but not when they were excluded (Watson et al., 2020). Patterns consistent with

Menzerath's law have also been found in a range of songbird species, both at the repeat phrase and

whole song level (James et al., 2021b; Youngblood, 2024). However, the mechanisms by which these

patterns arise have not been explored in detail in songbird species with complex vocalizations. In the

African penguin (*Spheniscuc demersus*), patterns consistent with Menzerath's law are achieved

through a combination of note choice and reducing the length of specific note types (Favaro et al.,

2020b). However, the vocal sequences of penguins are relatively simple compared to birdsong, so

determining if the same mechanisms operate in more complex avian vocal sequences could be

beneficial for comparative studies examining the evolution of language.

ZLA has been reported in a range of mammal taxa (Demartsev et al., 2019; Ferrer-i-Cancho &

Hernández-Fernández, 2013; Semple et al., 2010b; Valente et al., 2021; Youngblood, 2025), but is

not universally found across studies (Bezerra et al., 2011; Clink et al., 2020b; Youngblood, 2025). As

with Menzerath's law, adherence to ZLA may depend on the subset of the repertoire studied; in

common marmosets (*Callithrix jacchus*), long-distance calls did not adhere to ZLA, whereas short-

range signals were consistent with the expected pattern (Ferrer-i-Cancho & Hernández-Fernández,

2013). Adherence to ZLA may also depend on the unit of analysis. Across gibbon species, ZLA has

been reported at the note level, with shorter duration notes being used more frequently in loud

morning calls for the cao vit (*Nomascus nasutus*) and western black-crested (*Nomascus concolor*)

gibbons (Huang et al., 2020b), but not at the phrase level in male solos of the Mueller's gibbon

(*Hylobates muelleri*) (Clink et al., 2020b). ZLA is not commonly studied in birds, although a small

number of species have been examined to date (Favaro et al., 2020b; Ferrer-i-Cancho et al., 2013;

Gilman et al., 2025; Hailman et al., 1985; Youngblood, 2024). In African penguin calls, shorter

duration note types were used more frequently than longer ones, a pattern consistent with ZLA

(Favaro et al., 2020b). However, these calls are relatively simple and contain only a few note types.

In the common raven (*Corvus corax*), researchers found no significant evidence for ZLA (Ferrer-i-

Cancho et al., 2013). More recently, patterns consistent with ZLA have been reported in house finch

(*Haemorhous mexicanus*) songs (Youngblood, 2024). Gilman et al. (2025) found only weak evidence

for ZLA in individual populations of seven bird species, but a synthetic analysis across species

supported the presence of ZLA in birdsong generally. There is, therefore, a need to study more

populations of birds with complex note repertoires to fully understand ZLA in birdsong.

Here, we looked for patterns consistent with Menzerath's law and ZLA in Java sparrow (*Padda*

oryzivora) song (Lewis et al 2023). Each individual male Java sparrow produces a single song type

that is repeatable but varies across renditions (Ota & Soma, 2014). This variability in note usage both

among and within individuals makes the Java sparrow a good candidate for examining linguistic laws

in birdsong. Our goals were i) to assess whether Java sparrow song adheres to Menzerath's law, and,

if so, to determine whether birds achieve this by choosing shorter notes or by producing shorter

versions of the same notes in longer songs; and ii) to determine whether Java sparrow song adheres

to ZLA at the individual and population levels.

Methods

Study species

The Java sparrow (*Padda oryzivora*) is an oscine passerine and member of the family Estrildidae

(Restall, 1996). Songs in this species are used in courtship but not for territory defence. Each male

learns to produce a single song with 2-8 note types from his social father (Lewis et al., 2021, 2023;

Soma, 2011) and produces his song with some variation among renditions (Ota & Soma, 2014).

Learning fidelity is high, with sons' songs resembling their fathers' songs in a range of spectral and

temporal parameters (Lewis et al., 2021, 2023).

Dataset

To investigate the presence of linguistic laws in Java sparrow song, we used the dataset from Lewis

et al. (2023), which contains 22,966 notes across 676 undirected songs from 73 individuals (see

Figure 1 for examples). Birds in this dataset belonged to 10 social lineages. Each social lineage

included a single founder, all the males that learned their songs from that founder (i.e., his social

sons), the social sons of those birds, and so on for up to five generations. Thus, songs within social

lineages were not independent. Some birds (n=6) were reared in the presence of subtutors (an

additional adult male that was not their social or genetic father). Spectrogram inspection suggests

that subtutored birds generally produced songs that resembled their social fathers' song. Therefore,

subtutored individuals were classed as belonging to their social fathers' lineage for this study. Songs

were defined as series of notes with inter-note intervals <1 s (Kagawa & Soma, 2013; Lewis et al.,

2021). Songs were manually segmented into individual notes using the software Koe (Fukuzawa et

al., 2020) and classified into 16 note types, with high inter-observer repeatability (Lewis et al., 2021).

Java sparrow songs frequently begin with a single note or small sequence of notes repeated several

times with uncharacteristically long gaps (>0.25 s but <1 s) between repetitions (Lewis et al. 2023).

These introductory sequences may represent periods of preparation prior to main songs (as in zebra

finches e.g., (Kalra et al., 2021; Rajan & Doupe, 2013; Rao et al., 2019)), but may also be important

in song learning and development (Kalra et al., 2021). Therefore, it is difficult to determine whether

these notes should be counted as part of the main song sequence. As introductory notes contribute

to sequence length (i.e., the number of notes in a song), the presence of introductory notes may

affect our analysis of linguistic laws, and in particular, Menzerath's law. To account for the presence

of introductory notes, all analyses concerning Menzerath's law were conducted twice, once with and

once without introductory notes included, to determine if patterns were similar in both cases

(Appendix 1). As in Lewis et al. 2023, we considered the bird to have begun its main song as soon as

it began a sequence of 5 or more notes with inter-note intervals of <0.25 s, and we considered any

notes before the beginning of the main song to be introductions. The full dataset included 20911

notes when song introductions were excluded. Results were qualitatively similar for analyses with and

without introductory notes, with minor differences indicated in the 'Results' section.

Figure 1: Example spectrograms of songs from 8 individual Java sparrows from different social

lineages. Number of notes and the average log-transformed durations of notes for each song are

indicated next to spectrograms. Spectrograms were created using Seewave (Sueur et al., 2008)

(sample rate = 44.1kHz, window length = 512, overlap = 90%).

Data Analysis

Menzerath's Law

We examined adherence to Menzerath's law by asking i) whether longer songs (i.e., those with more

notes) are made up of shorter duration notes in the population as a whole. This is how Menzerath's

law has been studied in other animals (Gustison et al., 2016; Valente et al., 2021). Java sparrow

songs are composed of different note types, and note duration depends on both the note type and

the singer (Lewis et al., 2021). To examine how Menzerath's law might arise in Java sparrow song,

we asked ii) whether notes of the same type have shorter durations when they appear in longer

songs, and iii) whether longer songs are composed of shorter note types. Menzerath's law could arise

at the population level due to differences in song production among birds even if the same law does

not hold within individual birds. For example, we would see this if some birds produce only long songs

composed of short duration notes and other birds produce only short songs composed of long

duration notes. Therefore, we conducted a second set of analyses to ask whether Menzerath's law

holds within individuals. In particular, we asked, iv) whether individual Java sparrows sing shorter

duration notes when they produce longer songs, v) whether individuals use shorter duration versions

of the same note types in longer songs, and vi) whether individuals use shorter duration note types in

longer songs. To our knowledge, Menzerath's Law has not previously been studied at the level of

individuals in birdsong, although the importance of studying Menzerath's law at the individual level

has been advocated for human language (Grzybek & Stadlober, 2007).

To answer the above questions, we fit a series of linear mixed effect models using the lme4 (Bates et

al., 2015) and lmerTest packages (Kuznetsovs et al., 2017) in R (ver 4.1.3) (R Core Team, 2022). To

address question i) we regressed the log-transformed duration of individual notes on the length of

the song (defined as the number of notes in the song) that contained them, and on their log-

transformed position in the song. A significant negative relationship between the note duration and

song length would be evidence for Menzerath's law, and a significant relationship between note

duration and position in the song would indicate that note duration changed systematically over the

course of each song (i.e., notes later in the song are shorter or longer in duration than those at the

beginning). Note duration was log-transformed to stabilize variance, as longer notes are inherently

more variable than shorter notes. Position in the song was log-transformed, as we expected the

effect of position in the song to increase less than linearly with the position, and models with log-

transformed position provided lower AICc values. AICc values were not consistently lower for models

with log-transformed song length. Therefore, we fit models to untransformed song length fot

consistency with other recent work (Youngblood, 2024). To account for the non-independence of

note durations within songs, individuals and social lineages, we included random effects of song,

individual bird, and social lineage in the model. The random effects of song included a random

intercept, to account for the possibility that some songs might have longer notes than others, and a

random slope of position in song, to account for the possibility that note length might change

differently over the course of different songs. The random effects of bird and social lineage included

intercepts, slopes of log-transformed position in the song, and slopes of song length (number of

notes in the song). The random intercept accounts for the possibility that different birds or social

lineages might sing notes of different durations, the random slope of log-transformed position

accounts for the possibility that note duration may change differently over the course of songs in

different individuals or lineages, and the random effect of song length accounts for the possibility that

song length might affect note length differently in different individuals or lineages. There was no

random slope of song length within songs because each song has only one length. The random

slopes of song length within individuals and lineages allows that the strength of Menzerath's law may

differ among individuals and lineages.

Adherence to Menzerath's law at the population level could be achieved in two ways; birds might sing

shorter duration versions of the same note types in longer songs (question ii), or birds might use

shorter duration note types in longer songs (question iii). To disentangle these two possibilities, we

conducted two regressions identical to the one just described, but with different response variables.

To address question ii) the response variable was the log-transformed duration for each note in the

dataset, mean centred within its note type. Thus, differences in note duration due to the note types

are removed, and any patterns that remain must be due to differences in how notes of each type are

produced. To address question iii) we replaced each log-transformed note duration in the dataset

with the mean log-transformed duration for its note type in the whole dataset. Thus, variability within

note types is removed, and any pattern that remains must be due to the note types that birds use in

their songs.

Java sparrow songs might adhere to Menzerath's law across the whole population even if individual

birds do not use shorter duration notes in longer songs. To investigate patterns associated with

Menzerath's law within individual birds, we conducted a second set of regressions. In these analyses,

we mean-centred the song length within each bird and used the individual mean-centred song length,

rather than raw song length as a predictor. Any relationship between individual mean-centred song

length and note duration in these analyses cannot be due to differences in song length among

individuals, because those have been removed from the predictor, and therefore must be because

individual birds use shorter notes in longer songs. Analyses iv to vi are analogous to analyses i to iii,

respectively.

Zipf's Law of Abbreviation (ZLA)

We used Kendall's rank correlation (τ) test to assess the significance of the relationship between the

mean log-transformed duration of note types and their frequency of use in our study population.

Kendall's τ has been used to study ZLA in human languages (Bentz & Ferrer-i-Cancho, 2016) and

other animal communication systems (Ferrer-i-Cancho et al., 2013; Semple et al., 2010a). Typically, a

representative sample of the language (for humans) or vocalizations (for animals) is examined to

determine whether the length of individual units is negatively correlated with their frequency of use.

However, this simple approach may not transfer directly to birdsong. Individual birds may use highly

divergent or even non-overlapping sets of note types, making it difficult to create truly representative

samples of note type use in the population. Even if representative samples can be obtained,

differences in the use of note types among individuals could create patterns that look like ZLA at the

population level but do not arise by the Zipfian mechanism. Zipf (1935) hypothesised that individuals

shorten vocal units that they use more frequently to make communication more efficient. If this is

true, we should see a negative relationship between the duration and frequency of use of vocal units

within as well as among birds. However, if many birds sing only short note types and a few sing only

long note types, then the resulting population-level pattern may be consistent with ZLA even if no

individual bird uses short note types more frequently than long ones. Another challenge to studying

ZLA in birdsong is that birdsong can be highly stereotyped, and is often learned from other members

of the population. This means that observations of note type use within and among individuals are

not independent, and treating them as independent is likely to result in false positive inferences about

adherence to ZLA in populations.

To determine whether our population of Java sparrows adhered to ZLA, we developed a novel test

appropriate for bird populations. Briefly, we computed Kendall's τ for note duration and frequency of

use within each individual bird, and then computed the mean τ over all birds in the population, with

each bird weighted for the number of note types in its repertoire. This population-mean τ is a

measure of the strength of ZLA in the population and serves as a test statistic. To determine whether

the population-mean τ was larger than we would expect in a random sample from a population

without ZLA, we compared it to a null distribution that we computed while maintaining the observed

song structure and note similarities among birds. We implemented this method using the R package

ZLAvian, which we present in full elsewhere (Gilman et al., 2025).

Ethical Note

The dataset used in this study was initially collated for use in Lewis et al. (2021) and Lewis et al.

(2023). The dataset contains information extracted from archival recordings from a laboratory

population of Java sparrows (Padda oryzivora) housed at Hokkaido University. Archive data was

collected between 2011 and 2019 under the following approvals: Hokkaido University (Permit

numbers: 11-0028, 16-0020) and the University of Manchester AWERB (Permit number: D042).

Data Availability Statement

Data and code associated with this paper can be found at FigShare

https://doi.org/10.48420/24763299. Data and code from Lewis et al. (2023) can be found at

Figshare: http://doi.org/ 10.48420/22550149.

Results

Menzerath's law

We found evidence to support the presence of Menzerath's law in Java sparrow songs when

considering all songs in the dataset (i.e., the population level models i - iii); across birds, as song

length (number of notes) increased, notes within the song were shorter in duration (β = -0.004,

p=0.025; Table 1; Figure 2). Results were qualitatively similar when considering songs with

introductory notes removed, with a trend towards shorter duration notes in long songs (i.e., songs

with more notes) (β = -0.003, p=0.084; Appendix 1). At the population level, we did not find that

notes significantly changed in duration across the course of the song, although the direction of effect

was towards notes being longer when they appeared later in songs (with introductory notes β =

0.058, p=0.121 (Table 1); without introductory notes β = 0.042, p=0.263 (Appendix 1)). When

examining the potential drivers of Menzerath's law in Java sparrow songs, we found that birds did not

reduce the durations of note types in long songs (with introductory notes p=0.988 (Table 1); without

introductory notes p=0.850 (Appendix 1)). For example, note type 'A' in a longer songs was not

shorter in duration than note type 'A' in a shorter songs. There was also no change over the course of

the song (with introductory notes p=0.973 (Table 1); without introductory notes p=0.949 (Appendix

1)); note type 'A' later in a song was not shorter in duration than note type 'A' earlier in the song.

However, we did find evidence that birds used note types with shorter average durations more

frequently in longer songs (with introductory notes β = -0.004, p=0.002 (Table 1); without

introductory notes β = -0.003, p=0.003 (Appendix 1)). In addition, birds used note types with longer

average durations later in songs (β = 0.056, p=0.015; Table 1). Results were qualitatively similar

when considering songs with introductory notes removed (β = 0.038, p=0.051; Appendix 1). Overall,

these results suggest that adherence to Menzerath's law is driven by preferential use of short

duration note types, rather than by shortening the durations of note types in long songs.

Table 1: Results of GLMMs examining the effect of song length (number of notes) and note position

in songs on note duration. All durations were log-transformed.

Population level			
Question	*Measurement*	*Song length*	*Position in song (logged)*
Are longer songs made up of shorter notes?	Log note duration (log(s))	-0.004 p=0.025	0.058 p=0.121
Do longer songs contain shorter versions of the same note types?	Deviation from note type mean (log(s))	-6.24 x 10^{-6} p=0.988	-0.0007 p=0.973
Are longer songs made up of shorter note types?	Average logged duration of note type (log(s)	-0.004 p=0.002	0.056 p=0.015
Individual level			
Question	*Measurement*	*Song length*	*Position in song (logged)*
Are longer songs made up of shorter notes?	Log note duration (log(s)	-0.003 p=0.059	0.055 p=0.144
Do longer songs contain shorter versions of the same note types?	Deviation from note type mean (log(s))	-8.41 x 10^{-6} p=0.985	-0.0003 p=0.989
Are longer songs made up of shorter note types?	Average logged duration of note type (log(s))	-0.003 p=0.009	0.052 p=0.026

Figure 2: Average log-transformed note duration within a song (ms) plotted against song length

(number of notes). Each shape and colour combination represents a single individual. Individuals are

coloured based on their social lineage, which relates to song phenotype. Regression lines are

calculated for each social lineage and coloured accordingly.

We found weak evidence for Menzerath's law at the individual level (i.e., models iv-vi); individuals

used shorter duration notes when they sang songs with more notes (with introductory notes β = -

0.003, p=0.059 (Table 1); without introductory notes β = -0.002, p=0.084 (Appendix 1)). Again,

there was no evidence that note duration became shorter towards the end of the song, and our

results suggest a trend in the opposite direction (with introductory notes β = 0.055, p=0.144 (Table

1); without introductory notes β = 0.037, p=0.326 (Appendix 1)). The pattern in individuals was

driven by the same factors as the population level findings; birds used shorter duration note types in

longer songs (β = -0.003, p=0.009 (Table 1); without introductory notes β = -0.002, p=0.013

(Appendix 1)) and note types with longer average duration later in songs (with introductory notes β

= 0.052, p=0.026 (Table 1); without introductory notes β = 0.033, p=0.084 (Appendix 1)). There

was no evidence that birds changed the duration of individual note types based on song length (with

introductory notes p=0.984 (Table 1); without introductory notes p=0.899 (Appendix 1)) or that

notes of a given type were shorter in duration later in songs (with introductory notes p=0.986 (Table

1); without introductory notes p=0.943 (Appendix 1)).

Zipf's law of abbreviation (ZLA)

We did not find evidence for ZLA at either the population or individual levels. There was no evidence

that shorter duration note types were used more frequently than longer duration note types across all

songs in the population (Kendall's tau; τ=-0.150, p=0.209) or by individual birds (τ=-0.037,

p=0.431). The results are qualitatively similar if we exclude note types used by only one bird

(population level: τ=-0.282, p=0.090; individual level: τ=-0.030, p=0.441) or if we exclude note

types used fewer than 5 times in total (population level: τ=-0.181, p=0.174; individual level: τ=-

0.044, p=0.419).

Discussion

We found evidence that Java sparrow songs adhered to one of the two linguistic laws we studied.

Longer songs (i.e., those with more notes) were made up of shorter duration notes across all songs

in the population and within the songs of individual birds (Menzerath's law). This was due to birds

using shorter duration note types more frequently in longer songs, rather than shortening the

duration individual note types in longer songs. We did not find strong evidence to suggest that note

duration changed as the song progressed, although a trend towards longer duration notes later in the

song was apparent. However, we found that birds included longer duration note types towards the

end of their songs both across the population and within individuals.

Adherence to Menzerath's law at the level of repeat phrases has been reported previously in Java

sparrows (James et al., 2021b), and the same pattern is apparent in this study across full songs. In

our study population, songs often contained notes that were not part of repeat phrases, meaning that

whole song analysis was more appropriate. Adherence to Menzerath's law may be due in part to

constraints on overall song duration, which may relate to energetic costs, female preferences, or

species recognition. In Java sparrows, song length and song tempo are positively correlated; birds

that produce songs with more notes sing them at faster rates (Lewis et al, 2023). The current study

advances this result by showing that this pattern holds not just for song tempo, but for also for the

duration of the notes themselves. At the population level, the social learning of birdsong can create

patterns consistent with Menzerath's law. Each male Java sparrow sings a single song type (Kagawa

& Soma, 2013; Lewis et al., 2021; Soma, 2011), and the note types used and the order of note types

in each male's song are learned from his social father (Lewis et al., 2021; Soma, 2011). Therefore,

fathers and sons sing similar songs (Lewis et al., 2021; Soma, 2011). This means that correlations

between song length and note duration may persist within lineages, with some singing fewer, longer

notes and others singing more, faster notes. Our analyses controlled for non-independence of song

characteristics within birds and social lineages, so learning alone cannot account for the patterns

observed. This is further supported by the presence of Menzerath's law at the individual level; within

an individual, longer songs were also made up of shorter duration notes.

Conformation to Menzerath's law has been reported in a range of other passerine species (James et

al., 2021b), indicating that this pattern may be widespread across the order. However, the methods

by which Menzerath's law arises in birdsong have not been examined in detail. For this population,

notes within songs were classified into discrete categories, meaning that we could investigate how

patterns consistent with Menzerath's law were achieved in more detail. Longer songs consisted of

note types with shorter average durations, and there was no evidence that the duration of individual

note types was compressed in longer songs. Our results indicate that shortening of sequence

components in Java sparrows is achieved through note selection alone, rather than through changes

to note parameters. Note selection also contributes to adherence to Menzerath's law in the African

penguin (Favaro et al., 2020b), although changes to note parameters also play a role. Mediation of

Menzerath's law through note selection was apparent at both the population and individual level in

Java sparrows. As with overall adherence to Menzerath's law, this suggests that the pattern is not

driven solely by a few social lineages with long songs made up of short duration notes, but that

individuals also use this mechanism on a by-song basis.

It has been suggested that breathing and performance constraints may result in a negative

relationship between sequence position and element duration, i.e., elements towards the end of the

sequence are shorter (Gustison et al., 2016). However, we did not find this pattern in Java sparrows.

At the population and individual level, we did not find a significant correlation between note position

and note duration, although in both cases the direction of effect was towards longer duration notes

later in songs. However, at both the population and individual level birds were significantly more likely

to use longer duration note types later in their songs. A similar positive correlation between note

position and duration has been found in a number of non-human primates, with note duration

positively associated with position in the vocal sequence (Clink & Lau, 2020; Huang et al., 2020b;

Valente et al., 2021). The presence of longer duration notes later in the song, where a decline in

performance may be expected due to energetic constraints, could be an indicator of male quality. In

Java sparrows, songs are used during courtship and mate selection. Therefore, indicators of quality,

such as the positioning of long duration note types near the ends of songs may be preferred by

females. However, it may be that short, fast sections, especially those close to theoretical

performance limits (Kagawa & Soma, 2013; Podos, 1997), are more energetically costly, and the use

of longer duration notes later in the sequence is indicative of performance constraints in this species.

Across a range of species, singing performance declined after sustained singing (Sierro et al., 2023),

suggesting that performance limits could play a role in the expression of linguistic laws in this and

other bird species. Faithful song learning in the Java sparrow, where birds closely imitate their tutors'

song (Lewis et al., 2021; Soma, 2011), could contribute to the maintenance of songs with longer

duration notes at the end of the sequence across generations regardless of performance constraints,

especially in songs that are not near to performance limits.

We found no evidence that Java sparrow songs adhered to ZLA, either at the population or individual

level. Shorter duration note types were not used significantly more frequently than longer duration

note types across the population as a whole or within individual birds' repertoires. The presence of

ZLA has not been well studied in birds. Early studies in the black-capped chickadee found that call

bouts with smaller numbers of calls were more frequent than those with more calls (Hailman et al.,

1985), but not that calls with fewer notes were more common than those with more notes (Hailman

et al., 1985). The frequency of specific note types was not examined. This raises an interesting

question regarding the appropriate level of analysis for comparing the acoustic sequences of animals

with human language, as it is difficult to equate the various levels of organization (Semple et al.,

2022). In Java sparrows, note types are used in different combinations by different individuals,

indicating that notes may be the recombinant unit in songs and are an appropriate unit for analysis. A

similar approach was used to examine ZLA in the calls of African penguins, with the proportions of

individual notes being examined (Favaro et al., 2020b). Penguins' ecstatic display calls did adhere to

ZLA, with shorter syllables making up greater proportions of the calls. However, the calls contained

only three syllable types, one of which was inspiratory, making comparisons between these calls and

Java sparrow songs difficult. More recently, studies have explored adherence to ZLA in complex bird

vocalizations i.e., songs. Whilst some studies report adherence to ZLA (youngblood), a study of ZLA

across 11 bird populations found only weak concordance with ZLA for any individual population

(Gilman et al., 2025). Although a synthetic analysis found evidence for ZLA more broadly, the

negative concordance between duration and frequency of use was several times weaker than that

reported between word length and frequency of use in written human languages (Gilman et al.,

2025). If patterns associated with ZLA are weak in birdsong, then species, including the Java

sparrow, may not use enough individual note types to infer statistical significance. Indeed, while

evidence for ZLA in Java sparrows was not significant, the estimated concordance between note

duration and frequency of use was -0.037, which is within the range of values reported by Gilman et

al. (2025) for the populations they studied. It may be necessary to study songs of many different

populations and species to build a more general understanding of ZLA across birds, and this study

contributes to that effort.

If ZLA is weaker in birdsong than in human languages, as our results suggest, it may be due to a

range of phenomena that select against compression of vocal signals in birdsong (Ferrer-i-Cancho et

al., 2013). Java sparrow songs are used in courtship and mate choice, and so may be under different

selective pressures compared to other parts of the vocal repertoire. In this case, compression and

signal efficiency might not be an advantage, as songs can be honest indicators used to advertise male

quality (Gil & Gahr, 2002). This may act against compression in several ways. If longer duration notes

(or the use of multiple long notes) are signals of quality, longer duration notes may be

overrepresented in songs due to their effect on female preference. Similarly, certain note types with

particularly attractive qualities, such as 'sexy syllables' in canaries (Leitner et al., 2001; Vallet &

Kreutzer, 1995), may be over-represented. Faithful social learning of song features, as is apparent in

Java sparrows (Lewis et al., 2021, 2023), may also act against compression of vocal signals by

preventing compression of note types and changes in the proportion of different note types within

songs. Unlike human language, the individual 'meaning' of notes and songs, or how these are

perceived by the receiver, is not known. It is possible that signal compression affects perception by

the receiver. In the closely related Bengalese finch, female-directed songs are more consistent in

terms of acoustic structure of individual notes and sequence structure (Sakata et al., 2008). This

suggests that even small changes to notes within songs may be salient to females. Java sparrows

produce a range of call types in addition to songs (Restall, 1996). Calls often have specific meaning

(Marler, 2004), and so this subset of the repertoire may exhibit different patterns relating to

compression. In the common marmoset, long distance calls, which may be more affected by

attenuation and signal degradation, did not conform to ZLA, whereas short range calls were

consistent with the expected pattern (Ferrer-i-Cancho & Hernández-Fernández, 2013).

Overall, we find that Java sparrow songs adhere to Menzerath's law - longer sequences are made up

of shorter constituent parts – but we found no strong evidence for ZLA – more frequently used note

types do not have significantly shorter durations. It is likely that these laws reflect different

components and constraints on birdsong. Adherence to Menzerath's law may relate to a constraint on

song duration, where songs with more notes necessarily have a faster tempo. However, ZLA is more

closely related to meaning and information transfer, which is difficult to attribute to birdsong. It has

been suggested that some aspects of birdsong may be more suited to comparisons with human

music, rather than language, due to their repetitive nature, aspects of performance, and lack of

semantic meaning (Rothenberg et al., 2014). The case of meaning and information transfer may be a

case where comparisons to music are more appropriate than comparisons to language. Our findings

suggest that although there may be some limitations in applying linguistic laws designed for human

language to birdsong, they may nonetheless provide insight into communication universals and

warrant further investigation.

Author Contributions

Conceptualization: RL, AK, TG; Data curation: RL, MS; Formal analysis: RL, AK, TG; Funding

acquisition; RL, MS, TG; Investigation – RL, MS, SdK, TG; Methodology: RL, AK, TG; Resources: MS;

Supervision: MS, TG; Visualization: RL; Writing – initial draft: RL; Writing – reviewing and editing: RL,

AK, MS, SdK, TG.

Acknowledgements

RL's work on this project was funded by the Natural Environment Research Council (NERC) EAO

Doctoral Training Partnership (grant NE/L002469/1) supported by the Chester Zoo Conservation

Scholar and Fellow Scheme. RL also received funding through the JSPS Summer Programme. AK's

work on this project was funded by the Engineering and Physical Sciences Research Council (EPSRC)

Doctoral Training Partnership (grant number V520299). This project was also funded by the Daiwa

Anglo-Japanese Foundation and JSPS Grants-in-Aid for Young Scientists (grant 23680027 and

16H06177). We thank Nagisa Matsuda, Ken Otsuka, Hiroko Kagawa, and Nao Ota for archival song

recordings. We also thank CD Durrant for computational support. Finally, we thank Patrycja

Strycharczuk for useful discussions and feedback.

References

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

Bentz, C., & Ferrer-i-Cancho, R. (2016). Zipf's Law of Abbreviation as a Language Universal. *Proceedings of the Leiden Workshop on Capturing Phylogenetic Algorithms for Linguistics*, 1–4.

Berwick, R. C., Beckers, G. J. L., Okanoya, K., & Bolhuis, J. J. (2012). A bird's eye view of human language evolution. *Frontiers in Evolutionary Neuroscience*, *4*(APR), 1–25. https://doi.org/10.3389/fnevo.2012.00005

Berwick, R. C., Okanoya, K., Beckers, G. J. L., & Bolhuis, J. J. (2011). Songs to syntax: The linguistics of birdsong. *Trends in Cognitive Sciences*, *15*(3), 113–121. https://doi.org/10.1016/j.tics.2011.01.002

Bezerra, B. M., Souto, A. S., Radford, A. N., & Jones, G. (2011). Brevity is not always a virtue in primate communication. *Biology Letters*, *7*(1), 23–25. https://doi.org/10.1098/rsbl.2010.0455

Bolhuis, J. J., Okanoya, K., & Scharff, C. (2010). Twitter evolution: converging mechanisms in birdsong and human speech. *Nature Reviews Neuroscience*, *11*(11), 747–759. https://doi.org/10.1038/nrn2931

Clink, D. J., Ahmad, A. H., & Klinck, H. (2020a). Brevity is not a universal in animal communication: Evidence for compression depends on the unit of analysis in small ape vocalizations. *Royal Society Open Science*, *7*(4). https://doi.org/10.1098/rsos.200151

Clink, D. J., Ahmad, A. H., & Klinck, H. (2020b). Brevity is not a universal in animal communication: Evidence for compression depends on the unit of analysis in small ape vocalizations. *Royal Society Open Science*, *7*(4). https://doi.org/10.1098/rsos.200151

Clink, D. J., & Lau, A. R. (2020). Adherence to Menzerath's Law is the exception (not the rule) in three duetting primate species: Linguistic rules in duetting primates. *Royal Society Open Science*, *7*(11). https://doi.org/10.1098/rsos.201557

Demartsev, V., Gordon, N., Barocas, A., Bar-Ziv, E., Ilany, T., Goll, Y., Ilany, A., & Geffen, E. (2019). The "Law of Brevity" in animal communication: Sex-specific signaling optimization is determined by call amplitude rather than duration. *Evolution Letters*, *3*(6), 623–634. https://doi.org/10.1002/evl3.147

Favaro, L., Gamba, M., Cresta, E., Fumagalli, E., Bandoli, F., Pilenga, C., Isaja, V., Mathevon, N., & Reby, D. (2020a). Do penguins' vocal sequences conform to linguistic laws? *Biology Letters*, *16*(2). https://doi.org/10.1098/rsbl.2019.0589

Favaro, L., Gamba, M., Cresta, E., Fumagalli, E., Bandoli, F., Pilenga, C., Isaja, V., Mathevon, N., & Reby, D. (2020b). Do penguins' vocal sequences conform to linguistic laws? *Biology Letters*, *16*(2). https://doi.org/10.1098/rsbl.2019.0589

Ferrer-i-Cancho, R., & Hernández-Fernández, A. (2013). The Failure of the Law of Brevity in Two New World Primates. Statistical Caveats. *Glottotheory*, *4*(1). https://doi.org/10.1524/glot.2013.0004

Ferrer-i-Cancho, R., Hernández-Fernández, A., Lusseau, D., Agoramoorthy, G., Hsu, M. J., & Semple, S. (2013a). Compression as a Universal Principle of Animal Behavior. *Cognitive Science*, *37*(8), 1565–1578. https://doi.org/10.1111/cogs.12061

Ferrer-i-Cancho, R., Hernández-Fernández, A., Lusseau, D., Agoramoorthy, G., Hsu, M. J., & Semple, S. (2013b). Compression as a Universal Principle of Animal Behavior. *Cognitive Science*, *37*(8), 1565–1578. https://doi.org/10.1111/cogs.12061

Fukuzawa, Y., Webb, W. H., Pawley, M. D. M., Roper, M. M., Marsland, S., Brunton, D. H., & Gilman, A. (2020). Koe: Web-based software to classify acoustic units and analyse sequence structure in animal vocalizations. *Methods in Ecology and Evolution*, *11*(3), 431–441. https://doi.org/10.1111/2041-210X.13336

Gil, D., & Gahr, M. (2002). The honesty of bird song: Multiple constraints for multiple traits. *Trends in Ecology and Evolution*, *17*(3), 133–141. https://doi.org/10.1016/S0169-5347(02)02410-2

Gilman, R. T., Durrant, C. D., Malpas, L., & Lewis, R. N. (2025). Does Zipf's law of abbreviation shape birdsong? *PLOS Computational Biology*, *21*(8 August), 1–17. https://doi.org/10.1371/journal.pcbi.1013228

Grzybek, P., & Stadlober, E. (2007). Do we have problems with Arens' law? S new look at the sentence-word relation. In P. Grzybek & W. de Gruyter (Eds.), *Exact Methods in the Study of Language and Text: Dedicated to Gabriel Altmann on the Occasion of his 75th Birthday* (pp. 205–218). https://doi.org/10.1515/9783110894219

Gustison, M. L., Semple, S., Ferrer-I-Cancho, R., & Bergman, T. J. (2016). Gelada vocal sequences follow Menzerath's linguistic law. *Proceedings of the National Academy of Sciences of the United States of America*, *113*(19), E2750–E2758. https://doi.org/10.1073/pnas.1522072113

Hailman, J. P., Ficken, M. S., & Ficken, R. W. (1985). The 'chick-a-dee' calls of Parus atricapillus: A recombinant system of animal communication compared with written English. *Semiotica*, *56*(3–4), 191–224. https://doi.org/10.1515/semi.1985.56.3-4.191

Heesen, R., Hobaiter, C., Ferrer-I-Cancho, R., & Semple, S. (2019). Linguistic laws in chimpanzee gestural communication. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1896). https://doi.org/10.1098/rspb.2018.2900

Huang, M., Ma, H., Ma, C., Garber, P. A., & Fan, P. (2020a). Male gibbon loud morning calls conform to Zipf's law of brevity and Menzerath's law: insights into the origin of human language. *Animal Behaviour*, *160*, 145–155. https://doi.org/10.1016/j.anbehav.2019.11.017

Huang, M., Ma, H., Ma, C., Garber, P. A., & Fan, P. (2020b). Male gibbon loud morning calls conform to Zipf's law of brevity and Menzerath's law: insights into the origin of human language. *Animal Behaviour*, *160*, 145–155. https://doi.org/10.1016/j.anbehav.2019.11.017

Hyland Bruno, J., Jarvis, E. D., Liberman, M., & Tchernichovski, O. (2021). Birdsong Learning and Culture: Analogies with Human Spoken Language. *Annual Review of Linguistics*, *7*, 449–472. https://doi.org/10.1146/annurev-linguistics-090420-121034

James, L. S., Mori, C., Wada, K., & Sakata, J. T. (2021a). Phylogeny and mechanisms of shared hierarchical patterns in birdsong. *Current Biology*, *31*(13), 1–13. https://doi.org/10.1016/j.cub.2021.04.015

James, L. S., Mori, C., Wada, K., & Sakata, J. T. (2021b). Phylogeny and mechanisms of shared hierarchical patterns in birdsong. *Current Biology*, *31*(13), 1–13. https://doi.org/10.1016/j.cub.2021.04.015

Kagawa, H., & Soma, M. (2013). Song performance and elaboration as potential indicators of male quality in Java sparrows. *Behavioural Processes*, *99*, 138–144. https://doi.org/10.1016/j.beproc.2013.07.012

Kalra, S., Yawatkar, V., James, L. S., Sakata, J. T., & Rajan, R. (2021). Introductory gestures
before songbird vocal displays are shaped by learning and biological predispositions.
Proceedings of the Royal Society B: Biological Sciences, *288*(1943), 1–9.
https://doi.org/10.1098/rspb.2020.2796
Kuznetsovs, A., Brockhoff, P., & Christensen, R. (2017). lmerTest Package: Tests in Linear
Mixed Effects Models. *Journal of Statistical Software*, *82*(13), 1–26.
Leitner, S., Voigt, C., & Gahr, M. (2001). Seasonal Changes in the Song Pattern of the Non-
Domesticated Island Canary (Serinus canaria), a Field Study. *Behaviour*, *138*(7), 885–
904.
Lewis, R. N., Kwong, A., Soma, M., de Kort, S. R., & Gilman, R. T. (2023). Inheritance of
temporal song features in Java sparrows. *Animal Behaviour*, *206*, 61–74.
https://doi.org/10.1016/j.anbehav.2023.09.012
Lewis, R. N., Soma, M., de Kort, S. R., & Gilman, R. T. (2021). Like Father Like Son: Cultural
and Genetic Contributions to Song Inheritance in an Estrildid Finch. *Frontiers in
Psychology*, *12*(June). https://doi.org/10.3389/fpsyg.2021.654198
Marler, P. (2004). Bird calls: a cornucopia for communication. In *Nature's Music: the Science
of Birdsong* (pp. 132–177).
Ota, N., & Soma, M. (2014). Age-dependent song changes in a closed-ended vocal learner:
Elevation of song performance after song crystallization. *Journal of Avian Biology*, *45*(6),
566–573. https://doi.org/10.1111/jav.00383
Podos, J. (1997). A performance constraint on the evolution of trilled vocalizations in a
songbird family (Passeriformes: Emberizidae). *Evolution*, *51*(2), 537–551.
https://doi.org/10.1111/j.1558-5646.1997.tb02441.x
R Core Team. (2022). *R: A language and environment for statistical computing*. R Foundation
for Statistical Comuting.
Rajan, R., & Doupe, A. J. (2013). Behavioral and neural signatures of readiness to initiate a
learned motor sequence. *Current Biology*, *23*(1), 87–93.
https://doi.org/10.1016/j.cub.2012.11.040
Rao, D., Kojima, S., & Rajan, R. (2019). Sensory feedback independent pre-song vocalizations
correlate with time to song initiation. *Journal of Experimental Biology*, *222*(7).
https://doi.org/10.1242/jeb.199042
Restall, R. (1996). *Munias and mannikins*. Pica Press.
Rothenberg, D., Roeske, T. C., Voss, H. U., Naguib, M., & Tchernichovski, O. (2014).
Investigation of musicality in birdsong. *Hearing Research*, *308*, 71–83.
https://doi.org/10.1016/j.heares.2013.08.016
Sakata, J. T., Hampton, C. M., & Brainard, M. S. (2008). Social modulation of sequence and
syllable variability in adult birdsong. *Journal of Neurophysiology*, *99*(4), 1700–1711.
https://doi.org/10.1152/jn.01296.2007
Semple, S., Ferrer-i-Cancho, R., & Gustison, M. L. (2022). Linguistic laws in biology. *Trends in
Ecology and Evolution*, *37*(1), 53–66. https://doi.org/10.1016/j.tree.2021.08.012
Semple, S., Hsu, M. J., & Agoramoorthy, G. (2010a). Efficiency of coding in macaque vocal
communication. *Biology Letters*, *6*(4), 469–471. https://doi.org/10.1098/rsbl.2009.1062
Semple, S., Hsu, M. J., & Agoramoorthy, G. (2010b). Efficiency of coding in macaque vocal
communication. *Biology Letters*, *6*(4), 469–471. https://doi.org/10.1098/rsbl.2009.1062
Sierro, J., de Kort, S. R., & Hartley, I. R. (2023). A limit to sustained performance constrains
trill length in birdsong. *IScience*, *26*(11), 108206.
https://doi.org/10.1016/j.isci.2023.108206

Soma, M. (2011). Social Factors in Song Learning: A Review of Estrildid Finch Research.
Ornithological Science, *10*(2), 89–100. https://doi.org/10.2326/osj.10.89
Sueur, J., Aubin, T., & Simonis, C. (2008). Seewave, a free modular tool for sound analysis and
synthesis. *Bioacoustics*, *18*(2), 213–226.
https://doi.org/10.1080/09524622.2008.9753600
Valente, D., De Gregorio, C., Favaro, L., Friard, O., Miaretsoa, L., Raimondi, T., Ratsimbazafy,
J., Torti, V., Zanoli, A., Giacoma, C., & Gamba, M. (2021). Linguistic laws of brevity:
conformity in Indri indri. *Animal Cognition*, *0123456789*.
https://doi.org/10.1007/s10071-021-01495-3
Vallet, E., & Kreutzer, M. (1995). Female canaries are sexually responsive to special song
phrases. *Animal Behaviour*, *49*(6), 1603–1610. https://doi.org/10.1016/0003-
3472(95)90082-9
Wascher, C. A. F., & Youngblood, M. (2025). Vocal efficiency in crows. *Animal Cognition*,
28(1). https://doi.org/10.1007/s10071-025-01985-8
Watson, S. K., Heesen, R., Hedwig, D., Robbins, M. M., & Townsend, S. W. (2020). An
exploration of Menzerath's law in wild mountain gorilla vocal sequences: Menzerath's
Law in mountain gorillas. *Biology Letters*, *16*(10).
https://doi.org/10.1098/rsbl.2020.0380rsbl20200380
Youngblood, M. (2024). Language-like efficiency and structure in house finch song.
Proceedings of the Royal Society B, *291*, 20240250.
https://doi.org/10.1098/rspb.2024.0250/1438264/rspb.2024.0250.pdf
Youngblood, M. (2025). Language-like efficiency in whale communication. *Science Advances*,
11, eads6014. https://doi.org/10.1126/sciadv.ads6014
Zipf, G. K. (1935). *The psycho-biology of language: An introduction to dynamic philology*.
Houghton Mifflin Company.

674   Appendix 1: Results of GLMM examining the effect of song length and note position in songs (with

675   introductory notes removed) on duration. All durations were logged.

Population level			
Question	*Measurement*	*Song length*	*Position in song (logged)*
Are longer songs made up of shorter notes?	Log note duration (log(s))	-0.003 p=0.092	0.042 p=0.263
Do longer songs contain shorter versions of the same note types?	Deviation from note type mean (log(s))	0.00007 p=0.850	0.0016 p=0.949
Are longer songs made up of shorter note types?	Average logged duration of note type (log(s)	-0.003 p=0.003	0.038 p=0.051
Individual level			
Question	*Measurement*	*Mean centred song length*	*Position in song (logged)*
Are longer songs made up of shorter notes?	Log note duration (log(s)	-0.002 p=0.084	0.037 p=0.326
Do longer songs contain shorter versions of the same note types?	Deviation from note type mean (log(s))	-5.59×10^{-5} p=0.899	-0.002 p=0.943
Are longer songs made up of shorter note types?	Average logged duration of note type (log(s))	-0.002 p=0.013	0.033 p=0.084

677