Java sparrow song conforms to Mezerath's Law but not Zipf's

Law of Abbreviation

Rebecca N Lewis[1,2*], Anthony Kwong[1,3], Masayo Soma[4], Selvino R de Kort[5], R Tucker Gilman[1]

[1]Department of Earth and Environmental Sciences, University of Manchester, Manchester, UK

[2]Chester Zoo, Upton-by-Chester, Chester, UK

[3]Department of Mathematics, University of Manchester, Manchester, UK

[4]Department of Biology, Faculty of Science, Hokkaido University, Hokkaido, Japan

[5]Department of Natural Sciences, Manchester Metropolitan University, Manchester, United Kingdom

*Corresponding author: rebecca.lewis-3@manchester.ac.uk

Abstract

Linguistic laws, which describe common statistical patterns in human language, are often studied in

animal communication systems. Despite good evidence for the existence of these laws in non-human

primates, data is still limited for other taxa. Birdsong is often used as a model for human language

development due to similarities in vocal learning and production. Therefore, it is useful to explore

features of human language in birdsong to determine the value and limitations of such comparisons.

Here, we looked for patterns consistent with Menzerath's law, the tendency for longer sequences to

be made up of smaller constituent parts, and Zipf's Law of Abbreviation, the tendency for more

frequently used elements to be shorter in length, in the songs of Java sparrows (*Padda oryzivora*).

We found that songs conformed to Menzerath's law at both the population and individual level; longer

sequences were made up of shorter notes, and this pattern was achieved through note selection,

rather than shortening existing notes. This adherence to Menzerath's law may reflect species-specific

constraints on song durations. However, we found no evidence to support adherence to Zipf's Law of

Abbreviation, either at population or individual level; more frequently used notes were not shorter in

length. As they are used in courtship, songs may experience different selection pressure to other

parts of the vocal repertoire, which could act against compression. Whilst there may be some

limitations in applying linguistic laws designed for human language to birdsong, especially Zipf's Law

of Abbreviation, these laws may still provide insights into animal and human communication systems.

Keywords

33 Animal communication; Birdsong; Java Sparrow; Language; Linguistic Laws; Menzerath's Law; *Padda*

34 *oryzivora*; Zipf's Law of Abbreviation

Introduction

Birdsong and spoken language share many similarities (Berwick et al., 2011; Bolhuis et al., 2010;

Hyland Bruno et al., 2021). Both are made up of acoustic sequences, and there is overlap in the

behavioural, neural, and genetic bases of vocal learning and production in songbirds and humans

(Berwick et al., 2012, 2011; Bolhuis et al., 2010; Hyland Bruno et al., 2021). However, there are also

differences. In particular the close relationship between structure and meaning in human language is

thought not to exist in other animals (Bolhuis et al., 2010). Due to the similarities, birdsong is often

used in comparative studies examining the evolution and development of human language (e.g.,

Berwick et al., 2011). Therefore, it is useful to explore features of human language in animal

communication systems, especially birdsong, to help determine the value and limitations of these

comparisons.

Linguistic laws describe common statistical patterns in human language, where they have been

studied extensively (Semple et al., 2022). More recently, patterns consistent with linguistic laws have

also been explored in animal communication systems (Semple et al., 2022). Two laws that often

receive attention are Menzerath's law, the tendency for longer sequences to be made up of smaller

constituent parts, and Zipf's Law of Abbreviation (ZLA), the tendency for more frequently used

elements to be shorter in length. These laws are especially well studied in non-human primates (e.g.,

Bezerra et al., 2011; Clink et al., 2020; Clink & Lau, 2020; Gustison et al., 2016; Huang et al., 2020;

Semple et al., 2010; Valente et al., 2021; Watson et al., 2020) and some other mammal taxa (e.g.,

Demartsev et al. 2019; Ferrer-i-Cancho et al. 2013). Studies examining the presence of linguistic laws

in bird communications are still relatively limited (Favaro et al., 2020; Ferrer-i-Cancho et al., 2013;

Hailman et al., 1985; James et al., 2021).

Adherence to Menzerath's law has been widely reported in non-human primates (Clink et al., 2020;

Gustison et al., 2016; Heesen et al., 2019; Huang et al., 2020; Valente et al., 2021). Despite this,

adherence to the law is not universal, and may depend on the subset of vocalizations examined; in

mountain gorillas, Menzerath's law was apparent when single-unit sequences were included in

analyses, but not when they were excluded (Watson et al., 2020). Patterns consistent with

Menzerath's law have also been found in a range of songbird species, both at the repeat phrase and

whole song level (James et al., 2021). However, the mechanisms by which these patterns arise have

not been explored in detail in songbird species with complex vocalizations. In the African penguin

(*Spheniscuc demersus*), patterns consistent with Menzerath's law are achieved through a combination

of note choice and reducing the length of specific note types (Favaro et al., 2020). However, the

vocal sequences of penguins are relatively simple compared to birdsong, so determining if the same

mechanisms operate in more complex avian vocal sequences could be beneficial for comparative

studies examining the evolution of language.

ZLA has been reported in a range of mammal taxa (e.g., Demartsev et al., 2019; Ferrer-i-Cancho &

Hernández-Fernández, 2013; Semple et al., 2010; Valente et al., 2021), but is not universally found

across studies (e.g., Bezerra et al., 2011; Clink et al., 2020). As with Menzerath's law, adherence to

ZLA may depend on the subset of the repertoire studied; in common marmosets (*Callithrix jacchus*),

long-distance calls did not adhere to ZLA, whereas short-range signals were consistent with the

expected pattern (Ferrer-i-Cancho and Hernández-Fernández, 2013). Adherence to ZLA may also

depend on the unit of analysis. Across gibbon species, ZLA has been reported at the note level, with

shorter notes being used more frequently in loud morning calls for the cao vit (*Nomascus nasutus*)

and western black-crested (*Nomascus concolor*) gibbons (Huang et al., 2020), but not at the phrase

level in male solos of the Mueller's gibbon (*Hylobates muelleri*) (Clink et al., 2020). ZLA is not

commonly studied in birds, although a small number of species have been examined to date (Favaro

et al., 2020; Ferrer-i-Cancho et al., 2013; Hailman et al., 1985). In the African penguin, shorter note

types were used more frequently than longer ones, a pattern consistent with ZLA (Favaro et al.,

2020). However, these calls are relatively simple and contain only a few note types. In the common

raven (*Corvus corax*), patterns consistent with ZLA were not found for call usage; shorter call types

were not more frequently used in the vocal repertoire (Ferrer-i-Cancho et al., 2013). To our

knowledge, ZLA has not been examined in the songs of species with complex note repertoires.

Here, we looked for patterns consistent with Menzerath's law and ZLA in a published dataset of Java

sparrow (*Padda oryzivora*) song (Lewis et al 2023). Each individual male Java sparrow produces a

single song type that is repeatable but varies across renditions (Ota and Soma, 2014). This variability

in note usage both among and within individuals makes the Java sparrow a good candidate for

examining linguistic laws in birdsong. Our goals were i) to assess whether Java sparrow song adheres

to Menzerath's law, and, if so, to determine whether birds achieve this by choosing shorter notes or

by producing shorter versions of the same notes in longer songs; and ii) to determine whether Java

sparrow song adheres to ZLA at the individual and population levels.

Methods

Study species

The Java sparrow (*Padda oryzivora*) is an oscine passerine and member of the family Estrildidae

(Restall, 1996). Songs in this species are used in courtship but not for territory defense. Each male

learns to produce a single song with 2-8 note types (Ota and Soma, 2014) from his social father

(Lewis et al., 2023, 2021; Soma, 2011). Learning fidelity is high, with sons' songs resembling their

fathers' songs in a range of measures (Lewis et al., 2023, 2021)). Each male sings one stereotyped

song (Ota and Soma, 2014) with some variation between renditions.

Dataset

To investigate the presence of linguistic laws in Java sparrow song, we used the dataset from Lewis

et al. (2023), which contains 676 undirected songs from 73 individuals. Birds in this dataset belonged

to 10 social lineages. Each social lineage included a single founder, all of the males that learned their

songs from that founder (i.e., his social sons), the social sons of those birds, and so on for up to five

generations. Thus, songs within social lineages were not independent. Some birds (n=6) were reared

in the presence of subtutors (an additional adult male that was not their social or genetic father).

Spectrogram inspection suggests that subtutored birds generally produced songs that resembled their

social fathers' song. Therefore, subtutored individuals were classed as belonging to their social

fathers' lineage for this study. Songs were defined as series of notes with inter-note intervals <1 s

(Kagawa and Soma, 2013; Lewis et al., 2021). Songs were manually segmented into individual notes

using Koe (Fukuzawa et al., 2020) and classified into 16 note types, with high inter-observer

repeatability (Lewis et al., 2021). As in Lewis et al. (2023), we cleaned 4 additional notes that were

included in the dataset but separated by an interval >1 s. The dataset contained a total of 22,966

notes.

Java sparrow songs frequently contained a series of introductory notes, which are spectrally similar to

other notes in the song but indicated by note repetition and variable, uncharacteristically long gap

durations (>0.25s but <1s) (Lewis et al. 2023). As introductory notes contribute to sequence length,

the presence of introductory notes may, therefore, affect our analysis of linguistic laws, in particular,

Menzerath's law. To account for the presence of introductory notes, all analyses concerning

Menzerath's law were conducted twice, with and without introductory notes included, to determine if

patterns were similar in the two situations (Appendix 1). We identified introductory notes as in Lewis

et al. 2023, reducing the inter-note interval for inclusion in the song to 0.25s until five consecutive

notes with inter-note interval of <0.25s had been identified. The main song was considered to start at

the first note in the series of 5. The full dataset included 20911 notes when song introductions were

excluded. Results were qualitatively similar for analyses with and without introductory notes, and any

differences are indicated in the 'Results' section.

Data Analysis

Menzerath's Law

We examined adherence to Menzerath's law by asking i) whether longer songs are made up of

shorter notes in the population as a whole. This is how Menzerath's law has been studied in other

animals (Gustison et al., 2016; Valente et al., 2021). Java sparrow songs are composed of different

note types, and note duration depends on both the note type and the singer (Lewis et al., 2021). To

examine how Menzerath's law might arise in Java sparrow song, we asked ii) whether notes of the

same type are shorter when they appear in longer songs, and iii) whether longer songs are composed

of shorter note types. Menzerath's law could arise at the population level even if it does not hold

within individual birds due to individual differences in song production. For example, we would see

this if some birds produce only long songs composed of short notes and other birds produce only

short songs composed of long notes. Therefore, we conducted a second set of analyses to ask

whether Menzerath's law holds within individuals. In particular, we asked, iv) whether individual Java

sparrows sing shorter notes when they produce longer songs, v) whether individuals use shorter

versions of the same note types in longer songs, and vi) whether individuals use shorter note types in

longer songs.

To answer the above questions, we fit a series of linear mixed effect models using the lme4 (Bates et

al., 2015) and lmerTest packages (Kuznetsovs et al., 2017) in R (ver 4.1.3) (R Core Team, 2022). To

address question i) we regressed the logged duration of individual notes on the length of the song

that contained them and on their logged position in the song. A significant negative relationship

between the note duration and song length would be evidence for Menzerath's law, and a significant

relationship between note duration and position in the song would indicate that note duration

changed systematically over the course of each song (i.e., notes later in the song are shorter or

longer than those at the beginning). To account for the non-independence of note durations within

songs, individuals and social lineages, we included random effects of song, individual bird, and social

lineage in the model. The random effects of song included a random intercept, to account for the

possibility that some songs might have longer notes than others, and a random slope of position in

song, to account for the possibility that note length might change differently over the course of

different songs. The random effects of bird and social lineage included intercepts, slopes of logged

position in the song, and slopes of song length (number of notes in the song). The random intercept

accounts for the possibility that different birds or social lineages might sing notes of different

durations, the random slope of logged position accounts for the possibility that note duration may

change differently over the course of songs in different individuals or lineages, and the random effect

of song length accounts for the possibility that song length might affect note length differently in

different individuals or lineages. There was no random slope of song length within songs because

each song has only one length. The random slopes of song length within individuals and lineages

allows that the strength of Menzerath's law may differ among individuals and lineages.

Adherence to Menzerath's law at the population level could be achieved in two ways; birds might sing

shorter versions of the same note types in longer songs (question ii), or birds might use shorter note

types in longer songs (question iii). To disentangle these two possibilities, we conducted two

regressions identical to the one just described, but with different response variables. To address

question ii) the response variable was the logged duration for each note in the dataset, mean centred

within its note type. Thus, differences in note duration due to the note types are removed, and any

patterns that remain must be due to differences in how notes of each type are produced. To address

question iii) we replaced each logged note duration in the dataset with the mean logged duration for

its note type in the whole dataset. Thus, variability within note types is removed, and any pattern

that remains must be due to the note types that birds use in their songs.

Java sparrow songs might adhere to Menzerath's law across the whole population even if individual

birds do not use shorter notes in longer songs. To investigate patterns associated with Menzerath's

law within individual birds, we conducted a second set of regressions. In these analyses, we mean-

centred the song length within each bird and used the individual mean-centred song length, rather

than raw song length as a predictor. Any relationship between individual mean-centred song length

and note duration in these analyses cannot be due to differences in song length among individuals,

because those have been removed from the predictor, and therefore must be because individual birds

use shorter notes in longer songs. Analyses iv to vi are analogous to analyses i to iii, respectively.

Zipf's Law of Abbreviation (ZLA)

We used Kendall's rank correlation test to assess the significance of the relationship between the

mean logged duration of each note type and its frequency of use in our study population. This

approach has been used to study ZLA in human languages (Bentz and Ferrer-i-Cancho, 2016) and

other animal communication systems (Ferrer-i-Cancho et al., 2013; Semple et al., 2010). Typically, a

representative sample of the language (for humans) or vocalizations (for animals) is examined to

determine whether length of the individual types is negatively correlated with their frequency of use.

Whilst this approach is analogous to that used in other studies, it may not be the most appropriate

way to study ZLA in birdsong. Individual birds may use highly divergent or even non-overlapping sets

of note types, making it difficult to create truly representative samples of note type use in the

population. Even if representative samples can be obtained, differences in the use of note types

among individuals could create patterns that look like ZLA at the population level even if no individual

bird uses short note types more frequently than long ones. For example, this would arise if many

birds sing only short note types and a few sing only long note types. While this pattern would be

consistent with ZLA, the mechanism would not be. Zipf (1935) hypothesised that individuals shorten

vocal units that they use more frequently to make communication more efficient, and, if this is true,

we should see a negative relationship between the duration and frequency of use of vocal units

within as well as among individuals. Another challenge to studying ZLA in birdsong is that birdsong

can be highly stereotyped, and is often learned from other members of the population. This means

that observations of note type use within and among individuals are not independent, and treating

them as independent is likely to result in false positive inferences about adherence to ZLA in

populations.

With these challenges in mind, we developed a test to assess whether birdsong in a given population

adheres to ZLA. Consider a data set of annotated birdsongs in which notes have been classified to

type. Let d_{ijk} be the logged duration of the k^{th} instance of note type i produced by bird j, and let d_{ij} be

the mean logged duration of note type i produced by bird j. We study logged durations because the

unexplained variability in note duration increases with the mean, and logging standardises this

variability. Let n_{ij} be the number of times bird j produces note type i. ZLA predicts a negative

concordance between d_{ij} and n_{ij} within individual birds.

We assess concordance using Kendall's τ. For each bird j, we compute τ_j, the concordance between

d_{ij} and n_{ij}, and we compute v_j, the variance in our estimate of τ_j, where $v_j \approx \frac{2(2c_j+5)}{9c_j(c_j-1)}$ and c_j is the

number of note types produced by bird j (Valz and McLeod, 1990). We then compute τ_{obs}, the

observed population mean concordance with each τ_j weighted by its inverse variance: $\tau_{obs} =$

$\sum \tau_j v_j^{-1} / \sum v_j^{-1}$. This τ_{obs} assigns more weight to birds that use more note types because their

concordances are better estimates of the expected concordance for the population. This τ_{obs} is our

test statistic. This is convenient because τ_{obs} has a clear biological interpretation: if we randomly

choose one bird from the population and we randomly choose two different note types produced by

that bird, $(1 + \tau_{obs})/2$ is the probability that the note type with the longer duration will also be

performed more frequently.

We construct a null distribution for τ_{obs} while maintaining the observed similarities in song structure

and note similarity among birds. This accounts conservatively for the social learning of birdsong. For

each note type i we compute d_i, the mean logged duration of the note type with each bird weighted

equally. Thus, $d_i = b^{-1} \sum_{j=1}^{b} d_{ij}$ where b is the number of birds in the data set. Then, we compute ε_{ij},

the deviance of the logged duration of each note type i when produced by bird j from the population

mean logged duration for that note type. Thus, $\varepsilon_{ij} = d_{ij} - d_i$. If song structures are constrained but

the duration of notes is independent of their frequency of use, then we can create a null distribution

by permuting the population mean logged durations among the note types in the data set. Thus, for

each note i we select a randomised duration r_i from the set of ds without replacement. Then, we

compute r_{ij}, a randomised mean logged duration for note i as it might have been produced by bird j,

as either $r_{ij} = r_j + \varepsilon_{ij}$ for all i and j or $r_{ij} = r_j - \varepsilon_{ij}$ for all i and j, with equal probability. By either

adding all deviances or subtracting all deviances from the randomised means, we account for

variability in the durations of note types among birds while controlling for the fact that differences

from the population mean duration may be learned with the song structure (Lewis et al., 2021).

Finally, we compute the concordance between r_{ij} and n_{ij} for each bird in the randomised data, and we

average across all birds with each concordance weighted by its inverse variance. This is analogous to

our computation of τ_{obs} from the observed data, and results in one possible value of the test statistic

that we might obtain under the null hypothesis. We repeat this process 10^4 times to produce a null

distribution for τ_{obs}. The p-value for a negative concordance between d_{ij} and n_{ij} within individual birds

is the proportion of observations in the null distribution that are less than or equal to τ_{obs}. This

quantifies the strength of evidence for ZLA in the population.

Ethics statement

The dataset used in this study was initially collated for use in Lewis et al. (2021) and Lewis et al.

(2023) The dataset contains information extracted from archival recordings from a laboratory

population of Java sparrows (Padda oryzivora) housed at Hokkaido University. Archive data was

collected between 2011 and 2019 under the following approvals: Hokkaido University (Permit

numbers: 11-0028, 16-0020) and the University of Manchester AWERB (Permit number: D042).

Data Availability Statement

Data and code associated with this paper can be found at FigShare

https://doi.org/10.48420/24763299. Data and code from the original Lewis et al. (2021) paper can be

found at FigShare: https://doi. org/10.48420/14555247, and from the Lewis et al. (2023) paper at

Figshare: http://doi.org/ 10.48420/22550149.

Results

Menzerath's law

We found evidence to support the presence of Menzerath's law in Java sparrow songs when considering all songs in the dataset (i.e., the population level models i - iii); across birds, as song length (number of notes) increased, notes within the song were shorter (β = -0.004, p=0.025; Table 1; Figure 1). Results were qualitatively similar when considering songs with introductory notes removed, with a trend towards shorter notes in longer songs (β = -0.003, p=0.084; Appendix 1). Although not significant, our results suggest that note length differed across the course of the song, with notes tending to be longer when appearing later in the song (with introductory notes β = 0.058, p=0.121 (Table 1); without introductory notes β = 0.042, p=0.263 (Appendix 1)). When examining the potential drivers of Menzerath's law in Java sparrow songs, we found that birds did not reduce the duration of notes within types in longer songs (with introductory notes p=0.988 (Table 1); without introductory notes p=0.850 (Appendix 1)) i.e., note type 'A' in a long song was not shorter than note type 'A' in a short song. There was also no change over the course of the song (with introductory notes p=0.973 (Table 1); without introductory notes p=0.949 (Appendix 1)); note type 'A' later in the song was not shorter than note type 'A' earlier in the song. However, we did find evidence that birds used note types with shorter average durations in longer songs (with introductory notes β = -0.004, p=0.002 (Table 1); without introductory notes β = -0.003, p=0.003 (Appendix 1)). In addition, birds used note types with longer average durations later in songs (β = 0.056, p=0.015; Table 1). Results were qualitatively similar when considering songs with introductory notes removed, with a trend towards the use of longer note classes later in the song (β = 0.038, p=0.051; Appendix 1). Overall, these results suggest that adherence to Menzerath's law is driven by preferential use of short note types, rather than by shortening note types in long songs.

Table 1: Results of GLMMs examining the effect of song length and note position in songs on note duration. All durations were logged.

Population level			
Question	*Measurement*	*Song length*	*Position in song (logged)*
Are longer songs made up of shorter notes?	Log note duration (log(s))	-0.004 p=0.025	0.058 p=0.121
Do longer songs contain shorter versions of the same note types?	Deviation from note type mean (log(s))	-6.24×10^{-6} p=0.988	-0.0007 p=0.973
Are longer songs made up of shorter note types?	Average logged duration of note type (log(s)	-0.004 p=0.002	0.056 p=0.015
Individual level			
Question	*Measurement*	*Song length*	*Position in song (logged)*
Are longer songs made up of shorter notes?	Log note duration (log(s)	-0.003 p=0.059	0.055 p=0.144
Do longer songs contain shorter versions of the same note types?	Deviation from note type mean (log(s))	-8.41×10^{-6} p=0.985	-0.0003 p=0.989
Are longer songs made up of shorter note types?	Average logged duration of note type (log(s))	-0.003 p=0.009	0.052 p=0.026

Figure 1: Average logged note duration within a song (ms) plotted against song length (number of notes). Each shape and colour combination represents a single individual. Individuals are coloured based on their social lineage, which relates to song phenotype. Regression lines are calculated for each social lineage and coloured accordingly.

We found weak evidence for Menzerath's law at the individual level (i.e., models iv-vi); individuals used shorter notes when they sang longer songs (with introductory notes β = -0.003, p=0.060 (Table 1); without introductory notes β = -0.002, p=0.084 (Appendix 1)). Again, there was no evidence that note duration became shorter towards the end of the song, and our results suggest a trend in the opposite direction (with introductory notes β = 0.055, p=0.144 (Table 1); without introductory notes β = 0.037, p=0.326 (Appendix 1)). The pattern in individuals was driven by the same factors as the population level findings; birds used shorter note types in longer songs (β = -0.003, p=0.009 (Table 1); without introductory notes β = -0.002, p=0.013 (Appendix 1)) and note types with longer

average duration later in songs (with introductory notes β = 0.052, p=0.026 (Table 1); without

introductory notes β = 0.033, p=0.084 (Appendix 1)). There was no evidence that birds changed the

length of individual note types based on song length (with introductory notes p=0.984 (Table 1);

without introductory notes p=0.899 (Appendix 1)) or that notes of a given type were shorter later in

songs (with introductory notes p=0.986 (Table 1); without introductory notes p=0.943 (Appendix 1)).

Zipf's law of abbreviation (ZLA)

We did not find evidence for ZLA at either the population or individual levels. Shorter note types were

not used more frequently than longer note types across all songs in the population (Kendall's tau; τ=-

0.150, p=0.209), and individual birds did not use shorter note types more frequently than longer note

types (τ=-0.037, p=0.431). The results are qualitatively similar if we exclude note types used by only

one bird (population level: τ=-0.282, p=0.090; individual level: τ=-0.030, p=0.441) or if we exclude

note types used fewer than 5 times in total (population level: τ=-0.181, p=0.174; individual level:

τ=-0.044, p=0.419).

Discussion

Java sparrow songs adhered to one of the two linguistic laws tested. We found evidence to support

the presence of Menzerath's law in Java sparrow songs; longer sequences were made up of shorter

notes across all songs in the population and across songs within individuals. This is due to birds using

shorter note types more frequently in longer songs, rather than shortening individual note types in

longer songs. We did not find strong evidence to suggest that note duration changed as the song

progressed, although a trend towards longer notes later in the song was apparent. However, we

found that birds included longer note types towards the end of their songs both across the population

and within individuals. We did not find evidence for Zipf's Law of Abbreviation (ZLA) in Java sparrow

songs; shorter note types were not used more frequently in the population as a whole or by individual

birds.

Adherence to Menzerath's law at the level of repeat phrases has been reported previously in Java

sparrows (James et al., 2021), and the same pattern is apparent in this study across full songs. In our

study population, songs often contained notes that were not part of repeat phrases, meaning that

whole song analysis was more appropriate. Adherence to Menzerath's law may be due in part to

constraints on overall song duration, which may relate to energetic costs, female preferences, or

species recognition. In Java sparrows, song length and song tempo are negatively correlated; birds

that produce songs with more notes sing them at faster rates (Lewis et al, 2023). The current study

advances this result by showing that this pattern holds not just for song tempo, but for also for the

duration of the notes themselves. At the population level, the social learning of birdsong can create

patterns consistent with Menzerath's law. Each male Java sparrow sings a single song type (Kagawa

and Soma, 2013; Lewis et al., 2021; Soma, 2011), and the note types used and the order of note

types in each male's song are learned from his social father (Lewis et al., 2021; Soma, 2011).

Therefore, fathers and sons sing similar songs (Lewis et al., 2021; Soma, 2011). This means that

correlations between song length and note duration may persist within lineages, with some singing

fewer, longer notes and others singing more, faster notes. Our analyses controlled for non-

independence of song characteristics within birds and social lineages, so learning alone cannot

account for the patterns observed. This is further supported by the presence of Menzerath's law at

the individual level; within an individual, longer songs were also made up of shorter notes.

Conformation to Menzerath's law has been reported in a range of other passerine species (James et

al., 2021), indicating that this pattern may be widespread across the order. However, the methods by

which Menzerath's law arises in birdsong have not been examined in detail. For this population, notes

within songs were classified into discrete categories, meaning that we could investigate how patterns

consistent with Menzerath's law were achieved in more detail. Longer songs consisted of note types

with shorter average durations, and there was no evidence that individual note types were

compressed in longer songs. Our results indicate that shortening of sequence components in Java

sparrows is achieved through note selection alone, rather than through changes to note parameters.

Note selection also contributes to adherence to Menzerath's law in the African penguin (Favaro et al.,

2020), although changes to note parameters also plays a role. Mediation of Menzerath's law through

note selection was apparent at both the population and individual level in Java sparrows. As with

overall adherence to Menzerath's law, this suggests that the pattern is not driven solely by a few

social lineages with long songs made up of short notes, but that individuals also use this mechanism

on a by-song basis.

It has been suggested that breathing and performance constraints may result in a negative

relationship between sequence position and element duration, i.e., elements towards the end of the

sequence are shorter (Gustison et al., 2016). However, we did not find this pattern in Java sparrows.

At both the population and individual level, there was a trend towards longer notes later in

sequences, and birds were significantly more likely to use longer note types later in their songs. A

similar positive correlation between note position and duration has been found in a number of non-

human primates, with note length positively associated with position in the vocal sequence (Clink and

Lau, 2020; Huang et al., 2020; Valente et al., 2021). The presence of longer notes later in the song,

where a decline in performance may be expected due to energetic constraints, could be an indicator

of male quality. In Java sparrows, songs are used during courtship and mate selection. Therefore,

indicators of quality, such as the positioning of long note types near the ends of songs may be

preferred by females. However, it may be that short, fast sections, especially those close to

theoretical performance limits (Kagawa and Soma, 2013; Podos, 1997), are more energetically costly,

and the use of longer notes later in the sequence is indicative of performance constraints in this

species. Faithful song learning in the Java sparrow, where birds closely imitate their tutor's song

(Lewis et al., 2021; Soma, 2011), could contribute to the maintenance of songs with longer notes at

the end of the sequence across generations regardless of performance constraints, especially in songs

that are not near to performance limits.

We found no evidence that Java sparrow songs adhered to ZLA, either at the population or individual

level. Shorter note types were not more commonly used across the population as a whole, or within

individual birds' repertoires. The presence of ZLA has not been well studied in birdsong. Early studies

in the black-capped chickadee found that call bouts with a smaller number of calls were more

common than those with more calls (Hailman et al., 1985), but not that calls with fewer notes were

more common than those with more notes (Hailman et al., 1985). The frequency of specific note

types was not examined. This raises an interesting question regarding the appropriate level of

analysis for comparing the acoustic sequences of animals with human language, as it is difficult to

equate the various levels of organization (Semple et al., 2022). In Java sparrows, note types are used

in different combinations by different individuals, indicating that notes may be the recombinant unit in

songs and are an appropriate unit for analysis. A similar approach was used to examine ZLA in the

calls of African penguins, with the proportions of individual notes being examined (Favaro et al.,

2020). Penguins' ecstatic display calls did adhere to ZLA, with shorter syllables making up greater

proportions of the calls. However, the calls contained only three syllable types, one of which was

inspiratory, making comparisons between these calls and Java sparrow songs difficult.

ZLA is generally more challenging to study than Menzerath's law due to the need for classification of

notes or calls into discrete categories. It is not possible to know how birds perceive differences

between specific notes. As such, classification by human observers or computational methods may

not accurately represent the true categorization. Both human and computational classification

methods may be more likely to divide longer note types into spurious subclasses simply because long

note types have more features that can be observed. This means that statistical tests for patterns

consistent with ZLA in birdsong may tend to produce false positives. We did not find evidence for ZLA

in Java sparrows, so a tendency towards false positives was not a problem in our study.

A range of phenomena may select against compression of vocal signals (Ferrer-i-Cancho et al., 2013).

Java sparrow songs are used in courtship and mate choice, and so may be under different selective

pressures compared to other parts of the vocal repertoire. In this case, compression and signal

efficiency might not be an advantage, as songs can be honest indicators used to advertise male

quality (Gil and Gahr, 2002). This may act against compression in several ways. If longer notes (or

the use of multiple long notes) are signals of quality, longer notes may be overrepresented in songs

due to their effect on female preference. Similarly, certain note types with particularly attractive

qualities, such as 'sexy syllables' in canaries (Leitner et al., 2001; Vallet and Kreutzer, 1995), may be

over-represented. Faithful social learning of song features, as is apparent in Java sparrows (Lewis et

al., 2023, 2021), may also act against compression of vocal signals by preventing compression of

note types and changes in the proportion of different note types within songs. Unlike human

language, the individual 'meaning' of notes and songs, or how these are perceived by the receiver is

not known. It is possible that signal compression affects perception by the receiver. In the closely

related Bengalese finch, female-directed songs are more consistent in terms of acoustic structure of

individual notes and sequence structure (Sakata et al., 2008). This suggests that even small changes

to notes within songs may be salient to females. Java sparrows produce a range of call types in

addition to songs (Restall, 1996). Calls often have specific meaning (Marler, 2004), and so this subset

of the repertoire may exhibit different patterns relating to compression. In the common marmoset,

long distance calls, which may be more affected by attenuation and signal degradation, did not

conform to ZLA, whereas short range calls were consistent with the expected pattern (Ferrer-i-

Cancho and Hernández-Fernández, 2013).

Overall, we find that Java sparrow songs adhere to Menzerath's law - longer sequences are made up

of shorter constituent parts. However, they do not correspond with ZLA – more frequently used note

types do not have shorter durations. It is likely that these laws reflect different components and

constraints on birdsong. Adherence to Menzerath's law may relate to a constraint on song duration,

where songs with more notes necessarily have a faster tempo. However, ZLA is more closely related

to meaning and information transfer, which is difficult to attribute to birdsong. It has been suggested

that some aspects of birdsong may be more suited to comparisons with human music, rather than

language, due to their repetitive nature, aspects of performance, and lack of semantic meaning

(Rothenberg et al., 2014). The case of meaning and information transfer may be a case where

comparisons to music are more appropriate than comparisons to language. Our findings suggest that

although there may be some limitations in applying linguistic laws designed for human language to

birdsong, they may nonetheless provide insight into communication universals and warrant further

investigation.

Author Contributions

RL was responsible for the initial idea and RL, AK, and TG conceptualised the study. The original

dataset was collated by RL and MS. RL, MS, SdK, and TG contributed to data extraction. RL, AK and

TG conducted the data analysis. The initial draft was written by RL, with all authors contributing to

reviewing and editing the manuscript.

Acknowledgements

RL's work on this project was funded by the Natural Environment Research Council (NERC) EAO

Doctoral Training Partnership (grant NE/L002469/1) supported by the Chester Zoo Conservation

Scholar and Fellow Scheme. AK's work on this project was funded by the Engineering and Physical

Sciences Research Council (EPSRC) Doctoral Training Partnership (grant number V520299). This

project was also funded by the Daiwa Anglo-Japanese Foundation and JSPS Grants-in-Aid for Young

Scientists (grant 23680027 and 16H06177). We thank Nagisa Matsuda, Ken Otsuka, Hiroko Kagawa,

and Nao Ota for archival song recordings. We also thank CD Durrant for computational support.

Finally, we thank Patrycja Strycharczuk for useful discussions and feedback.

References

Bates, D., Mächler, M., Bolker, B.M., Walker, S.C., 2015. Fitting linear mixed-effects models using

lme4. J. Stat. Softw. 67. https://doi.org/10.18637/jss.v067.i01

Bentz, C., Ferrer-i-Cancho, R., 2016. Zipf's Law of Abbreviation as a Language Universal. Proc. Leiden

Work. Capturing Phylogenetic Algorithms Linguist. 1–4.

Berwick, R.C., Beckers, G.J.L., Okanoya, K., Bolhuis, J.J., 2012. A bird's eye view of human language

evolution. Front. Evol. Neurosci. 4, 1–25. https://doi.org/10.3389/fnevo.2012.00005

Berwick, R.C., Okanoya, K., Beckers, G.J.L., Bolhuis, J.J., 2011. Songs to syntax: The linguistics of

birdsong. Trends Cogn. Sci. 15, 113–121. https://doi.org/10.1016/j.tics.2011.01.002

Bezerra, B.M., Souto, A.S., Radford, A.N., Jones, G., 2011. Brevity is not always a virtue in primate

communication. Biol. Lett. 7, 23–25. https://doi.org/10.1098/rsbl.2010.0455

Bolhuis, J.J., Okanoya, K., Scharff, C., 2010. Twitter evolution: converging mechanisms in birdsong

and human speech. Nat. Rev. Neurosci. 11, 747–759. https://doi.org/10.1038/nrn2931

Clink, D.J., Ahmad, A.H., Klinck, H., 2020. Brevity is not a universal in animal communication:

Evidence for compression depends on the unit of analysis in small ape vocalizations. R. Soc.

Open Sci. 7. https://doi.org/10.1098/rsos.200151

Clink, D.J., Lau, A.R., 2020. Adherence to Menzerath's Law is the exception (not the rule) in three

duetting primate species: Linguistic rules in duetting primates. R. Soc. Open Sci. 7.

https://doi.org/10.1098/rsos.201557

Demartsev, V., Gordon, N., Barocas, A., Bar-Ziv, E., Ilany, T., Goll, Y., Ilany, A., Geffen, E., 2019. The

"Law of Brevity" in animal communication: Sex-specific signaling optimization is determined by

call amplitude rather than duration. Evol. Lett. 3, 623–634. https://doi.org/10.1002/evl3.147

Favaro, L., Gamba, M., Cresta, E., Fumagalli, E., Bandoli, F., Pilenga, C., Isaja, V., Mathevon, N.,

Reby, D., 2020. Do penguins' vocal sequences conform to linguistic laws? Biol. Lett. 16.

https://doi.org/10.1098/rsbl.2019.0589

Ferrer-i-Cancho, R., Hernández-Fernández, A., 2013. The Failure of the Law of Brevity in Two New

World Primates. Statistical Caveats. Glottotheory 4. https://doi.org/10.1524/glot.2013.0004

Ferrer-i-Cancho, R., Hernández-Fernández, A., Lusseau, D., Agoramoorthy, G., Hsu, M.J., Semple, S.,

2013. Compression as a Universal Principle of Animal Behavior. Cogn. Sci. 37, 1565–1578.

https://doi.org/10.1111/cogs.12061

Fukuzawa, Y., Webb, W.H., Pawley, M.D.M., Roper, M.M., Marsland, S., Brunton, D.H., Gilman, A.,

2020. Koe: Web-based software to classify acoustic units and analyse sequence structure in

animal vocalizations. Methods Ecol. Evol. 11, 431–441. https://doi.org/10.1111/2041-

210X.13336

Gil, D., Gahr, M., 2002. The honesty of bird song: Multiple constraints for multiple traits. Trends Ecol.

Evol. 17, 133–141. https://doi.org/10.1016/S0169-5347(02)02410-2

Gustison, M.L., Semple, S., Ferrer-I-Cancho, R., Bergman, T.J., 2016. Gelada vocal sequences follow

Menzerath's linguistic law. Proc. Natl. Acad. Sci. U. S. A. 113, E2750–E2758.

https://doi.org/10.1073/pnas.1522072113

Hailman, J.P., Ficken, M.S., Ficken, R.W., 1985. The 'chick-a-dee' calls of Parus atricapillus: A

recombinant system of animal communication compared with written English. Semiotica 56,

191–224. https://doi.org/10.1515/semi.1985.56.3-4.191

Heesen, R., Hobaiter, C., Ferrer-I-Cancho, R., Semple, S., 2019. Linguistic laws in chimpanzee

gestural communication. Proc. R. Soc. B Biol. Sci. 286. https://doi.org/10.1098/rspb.2018.2900

Huang, M., Ma, H., Ma, C., Garber, P.A., Fan, P., 2020. Male gibbon loud morning calls conform to

Zipf's law of brevity and Menzerath's law: insights into the origin of human language. Anim.

Behav. 160, 145–155. https://doi.org/10.1016/j.anbehav.2019.11.017

Hyland Bruno, J., Jarvis, E.D., Liberman, M., Tchernichovski, O., 2021. Birdsong Learning and Culture:

Analogies with Human Spoken Language. Annu. Rev. Linguist. 7, 449–472.

https://doi.org/10.1146/annurev-linguistics-090420-121034

James, L.S., Mori, C., Wada, K., Sakata, J.T., 2021. Phylogeny and mechanisms of shared hierarchical

patterns in birdsong. Curr. Biol. 31, 1–13. https://doi.org/10.1016/j.cub.2021.04.015

Kagawa, H., Soma, M., 2013. Song performance and elaboration as potential indicators of male

quality in Java sparrows. Behav. Processes 99, 138–144.

https://doi.org/10.1016/j.beproc.2013.07.012

Kuznetsovs, A., Brockhoff, P., Christensen, R., 2017. lmerTest Package: Tests in Linear Mixed Effects

Models. J. Stat. Softw. 82, 1–26.

Leitner, S., Voigt, C., Gahr, M., 2001. Seasonal Changes in the Song Pattern of the Non-Domesticated

Island Canary (Serinus canaria), a Field Study. Behaviour 138, 885–904.

Lewis, R.N., Kwong, A., Soma, M., de Kort, S.R., Gilman, R.T., 2023. Inheritance of temporal song

features in Java sparrows. Anim. Behav. 206, 61–74.

https://doi.org/10.1016/j.anbehav.2023.09.012

Lewis, R.N., Soma, M., de Kort, S.R., Gilman, R.T., 2021. Like Father Like Son: Cultural and Genetic

Contributions to Song Inheritance in an Estrildid Finch. Front. Psychol. 12.

https://doi.org/10.3389/fpsyg.2021.654198

Marler, P., 2004. Bird calls: a cornucopia for communication, in: Nature's Music: The Science of

Birdsong. pp. 132–177.

Ota, N., Soma, M., 2014. Age-dependent song changes in a closed-ended vocal learner: Elevation of

song performance after song crystallization. J. Avian Biol. 45, 566–573.

https://doi.org/10.1111/jav.00383

Podos, J., 1997. A performance constraint on the evolution of trilled vocalizations in a songbird family

(Passeriformes: Emberizidae). Evolution (N. Y). 51, 537–551. https://doi.org/10.1111/j.1558-

5646.1997.tb02441.x

R Core Team, 2022. R: A language and environment for statistical computing.

Restall, R., 1996. Munias and mannikins. Pica Press, Sussex.

Rothenberg, D., Roeske, T.C., Voss, H.U., Naguib, M., Tchernichovski, O., 2014. Investigation of

musicality in birdsong. Hear. Res. 308, 71–83. https://doi.org/10.1016/j.heares.2013.08.016

Sakata, J.T., Hampton, C.M., Brainard, M.S., 2008. Social modulation of sequence and syllable

variability in adult birdsong. J. Neurophysiol. 99, 1700–1711.

https://doi.org/10.1152/jn.01296.2007

Semple, S., Ferrer-i-Cancho, R., Gustison, M.L., 2022. Linguistic laws in biology. Trends Ecol. Evol. 37,

53–66. https://doi.org/10.1016/j.tree.2021.08.012

Semple, S., Hsu, M.J., Agoramoorthy, G., 2010. Efficiency of coding in macaque vocal communication.

Biol. Lett. 6, 469–471. https://doi.org/10.1098/rsbl.2009.1062

Soma, M., 2011. Social Factors in Song Learning: A Review of Estrildid Finch Research. Ornithol. Sci.

10, 89–100. https://doi.org/10.2326/osj.10.89

Valente, D., De Gregorio, C., Favaro, L., Friard, O., Miaretsoa, L., Raimondi, T., Ratsimbazafy, J.,

Torti, V., Zanoli, A., Giacoma, C., Gamba, M., 2021. Linguistic laws of brevity: conformity in

Indri indri. Anim. Cogn. https://doi.org/10.1007/s10071-021-01495-3

Vallet, E., Kreutzer, M., 1995. Female canaries are sexually responsive to special song phrases. Anim.

Behav. 49, 1603–1610. https://doi.org/10.1016/0003-3472(95)90082-9

Valz, P.D., McLeod, A.I., 1990. A Simplified Derivation of the Bariance of Kendall's Rank Correlation

Coefficient. Am. Stat. 44, 39–40.

Watson, S.K., Heesen, R., Hedwig, D., Robbins, M.M., Townsend, S.W., 2020. An exploration of

Menzerath's law in wild mountain gorilla vocal sequences: Menzerath's Law in mountain gorillas.

Biol. Lett. 16. https://doi.org/10.1098/rsbl.2020.0380rsbl20200380

Zipf, G.K., 1935. The psycho-biology of language: An introduction to dynamic philology. Houghton

Mifflin Company, Boston.

Appendix 1: Results of GLMM examining the effect of song length and note position in songs (with

introductory notes removed) on duration. All durations were logged.

Population level			
Question	*Measurement*	*Song length*	*Position in song (logged)*
Are longer songs made up of shorter notes?	Log note duration (log(s))	-0.003 p=0.092	0.042 p=0.263
Do longer songs contain shorter versions of the same note types?	Deviation from note type mean (log(s))	0.00007 p=0.850	0.0016 p=0.949
Are longer songs made up of shorter note types?	Average logged duration of note type (log(s)	-0.003 p=0.003	0.038 p=0.051
Individual level			
Question	*Measurement*	*Mean centred song length*	*Position in song (logged)*
Are longer songs made up of shorter notes?	Log note duration (log(s)	-0.002 p=0.084	0.037 p=0.326
Do longer songs contain shorter versions of the same note types?	Deviation from note type mean (log(s))	-5.59×10^{-5} p=0.899	-0.002 p=0.943
Are longer songs made up of shorter note types?	Average logged duration of note type (log(s))	-0.002 p=0.013	0.033 p=0.084

595