Human Trophoblast Stem Cells Restrict Human Cytomegalovirus Replication

Tyler B. Rollman[1], Zachary W. Berkebile[1], Hiroaki Okae[2], Vivian J. Bardwell[3], Micah D.

Gearhart[4], and Craig J. Bierle[1#]

[1.] Division of Pediatric Infectious Diseases, Department of Pediatrics, University of Minnesota,

Minneapolis, MN.

[2.] Department of Informative Genetics, Tohoku University Graduate School of Medicine, Sendai,

Japan.

[3.] Developmental Biology Center, Department of Genetics, Cell Biology and Development and

the Masonic Cancer Center, University of Minnesota, Minneapolis, MN

[4.] Department of Obstetrics, Gynecology and Women's Health, University of Minnesota,

Minneapolis, MN.

Running Head: Trophoblast Stem Cells Restrict HCMV

#Address correspondence to Craig J Bierle, cjbierle@umn.edu

Abstract word count: 186

Text word count: 10,143

Abstract.

Placental infection plays a central role in the pathogenesis of congenital human cytomegalovirus

(HCMV) infections and is a cause of fetal growth restriction and pregnancy loss. HCMV can

replicate in some trophoblast cell types, but it remains unclear how the virus evades antiviral

immunity in the placenta and how infection compromises placental development and function.

Human trophoblast stem cells (TSCs) can be differentiated into extravillous trophoblasts (EVTs),

syncytiotrophoblasts (STBs), and organoids, and this study assessed the utility of TSCs as a

model of HCMV infection in the first trimester placenta. HCMV was found to non-productively

infect TSCs, EVTs, and STBs. Immunofluorescence assays and flow cytometry experiments

further revealed that infected TSCs frequently only express immediate early viral gene products.

Similarly, RNA-sequencing found that viral gene expression in TSCs does not follow the kinetic

patterns observed during lytic infection in fibroblasts. Canonical antiviral responses were largely

not observed in HCMV-infected TSCs and TSC-derived trophoblasts. Rather, infection

dysregulated factors involved in cell identity, differentiation, and WNT signaling. Thus, while

HCMV does not replicate in TSCs, infection may perturb trophoblast differentiation in ways that

could interfere with placental function.

Importance.

Placental infection plays a central role in HCMV pathogenesis during pregnancy, but the

species-specificity of HCMV and the limited availability and lifespan of primary trophoblasts

have been persistent barriers to understanding how infection impacts this vital organ. Human

TSCs represent a new approach to modeling viral infection early in placental development. This

study reveals that TSCs, like other stem cell types, restrict HCMV replication. However, infection

perturbs the expression of genes involved in differentiation and cell fate determination, pointing

to a mechanism by which HCMV could cause placental injury.

Introduction.

Human cytomegalovirus (HCMV) is a highly prevalent beta-herpesvirus and a leading infectious cause of birth defects and adverse pregnancy outcomes. Recent surveys estimate the global seroprevalence of HCMV to be 83% across all demographics, and in the United States 65% of women of childbearing age are latently infected with HCMV (1). While HCMV infection rarely presents clinically in healthy individuals, the virus is an opportunistic pathogen that can cause severe illness in the immunocompromised and can infect the placenta and developing fetus. Congenital CMV (cCMV) infections occur in an estimated 0.66% of all live births in the United States and 10% of cCMV infections present as symptomatic at birth (2). Roughly 20% of children born with either symptomatic or asymptomatic cCMV develop sensorineural disabilities, such as neurosensory hearing loss, chorioretinitis leading to vision loss, microcephaly, and intellectual disability (3). Placental HCMV infections can cause a broad spectrum of injuries even in the absence of detectable viral transmission to the fetus. Pathologic findings in infected placentas include maldevelopment of chorionic villi, diffuse villitis, and areas of necrosis and calcification (4, 5). HCMV is a major contributor to idiopathic intrauterine growth restriction and an under recognized cause of fetal demise (6, 7). The timing of placental infection and virus-induced placental damage correlate with fetal injury (8-11). While placental infection can cause growth restriction or fetal demise after maternal infection at any time in pregnancy, the highest rates of HCMV-associated growth restriction and placental malformations are observed when infection occurs during the third trimester (12).

The placenta is formed by blastocyst-derived trophoblasts and these cells play a central role in antiviral immunity at the maternal-fetal interface. Trophoblast stem cells (TSCs) originate in the trophectoderm and small populations of TSCs are thought to be maintained throughout pregnancy. TSCs differentiate into proliferative cytotrophoblasts (CTBs), which are located both in cytotrophoblastic columns that anchor the placenta to the decidua and on the fetal side of chorionic villi. CTBs terminally-differentiate into extravillous trophoblasts (EVTs) and

syncytiotrophoblasts (STBs). EVTs invade and remodel the maternal decidua, playing a key role

in regulating maternal blood flow into the placenta. STBs form a non-proliferative, continuous

syncytium that separates maternal blood from fetal circulation and release antiviral cytokines,

type III interferons, and exosomes that contain antiviral microRNAs (miRNAs) (13). Type III

interferon is constitutively expressed by trophoblasts during mid and late gestation, regulating

the expression of select interferon stimulated genes (ISG) (14-17). The chromosome 19 miRNA

Cluster (C19MC) encodes primate-specific miRNAs, which are expressed exclusively in the

placenta, and have broad antiviral activity (14). These and other antiviral properties of

trophoblasts make the placenta an effective barrier against the vertical transmission of most

pathogens.

HCMV can infect and replicate in some trophoblast cell types despite their intrinsic

immune defenses. In infected placenta, HCMV is most frequently detected by

immunohistochemistry in CTBs and EVTs (5, 18). Experimental infection studies in explanted

placenta and primary trophoblasts have revealed that HCMV can productively infect CTBs and

EVTs but not STBs (8, 19, 20). How HCMV infection affects placental function remains unclear.

Proliferative trophoblasts have an impaired capacity to differentiate when infected, which may

interfere with the establishment and maintenance of the maternal-fetal interface and potentially

facilitate viral dissemination from maternal circulation to the fetus (8, 19, 21). Inflammation

triggered by viral infection can also cause placental dysfunction (22). In murine models of

congenital Zika syndrome, fetal wastage may be driven by type I interferon signaling triggered

by viral infection (23). Maternal polyinosinic:polycytidylic acid (Poly I:C) treatment, which triggers

innate immune responses by stimulating pattern recognition receptors and directly activates

intrinsic antiviral pathways such as protein kinase R, can affect placental and fetal development

in mice and rats [reviewed in (24)]. Elevated concentrations of proinflammatory cytokines are

observed in amniotic fluid and maternal blood in cases of cCMV (25-27). *In vitro* studies in

tissue explants and organoids have revealed that HCMV infection triggers distinct patterns of

cytokine production from decidua and placenta (28-30).

Several barriers have prevented a greater understanding of HCMV pathogenesis in the

placenta. The cytomegaloviruses are species-specific and cannot replicate in animals other than

their natural host. Tissue explants and primary trophoblasts are prepared from tissue that is

collected either after elective pregnancy terminations or birth. While term placentas are typically

discarded and can be collected for research purposes with relative ease, the use and collection

of tissue from other times in gestation is more highly regulated if possible at all. Gene

expression studies have focused on placentas aged up to 20 weeks or full-term; other data from

the late second and third trimester comes from placentas that were delivered prematurely (31).

Trophoblast cell lines such as BeWo and JEG3 have abnormal antiviral properties relative to

primary trophoblasts that limit their utility as infection models (32, 33).

Methods to culture and maintain undifferentiated human trophoblast stem cells (TSCs)

were recently developed (34). TSCs were first derived from blastocysts and villous tissue from

first-trimester placentas, but cell lines were subsequently established from naïve embryonic or

induced pluripotent stem cells and older placentas (30, 35). Mutant, clonal TSC lines can be

generated by CRISPR/Cas9 mutagenesis and TSCs can be differentiated into EVT and STB

monocultures or placental organoids (30, 34-37). Given the potential utility of TSCs as an

experimental model of HCMV infection in the first trimester placenta, we investigated the

capacity of HCMV to infect and replicate in TSCs and TSC-derived EVT and STB. While all

three cell types can be infected by HCMV, very little, if any, virus was produced from infected

trophoblasts. Further investigation revealed that HCMV replication was restricted at some point

after entry. TSC growth was not affected by high multiplicity HCMV infections, and the virus was

not maintained in the cells upon passage. While these findings may suggest that TSCs could

function as an infection-resistant reservoir of proliferative cells, gene expression profiling did

note significant changes in the transcription of factors related to cell fate and development,

suggesting that infection may affect the capacity of TSCs to differentiate.

Results.

HCMV infects but does not replicate in TSCs and TSC-derived trophoblasts.

To assess whether HCMV could replicate in TSCs and TSC-derived trophoblasts,

female (CT27) and male (CT29) TSCs were grown in media that either maintained the cells in

an undifferentiated state (self-renewal media) or stimulated differentiation into EVTs or STBs

(34, 36). The cells were infected with TB40/Ewt-mCherry (HCMV mCherry), a clinical-like strain

that maintains the broad tropism of wild-type HCMV, at a multiplicity of infection (MOI) of 0.1

(38). HCMV-permissive retinal pigment epithelial cells (ARPE-19) were infected in parallel.

mCherry-expressing cells were observed when all three cell types were infected (**Fig. 1A**).

When cell-associated and extracellular virus was titered, HCMV replication peaked at 3 days

post infection (dpi) in ARPE-19 cells (**Fig. 1B**). Titers from infected TSCs and STBs were near

the limit of detection throughout the experiment. Peak titers from infected EVTs were higher,

near $1X10^3$ plaque forming units (PFU)/ml. However, this concentration of virus could be

detected at 1 dpi, likely before HCMV could have completed a single cycle of replication, and

may be residual input virus.

HCMV mCherry is a TB40/E Bac4-based recombinant virus where non-excisable

bacterial sequences and the *mCherry* expression cassette have replaced the viral genes *US1*

through *US6* (39). To assess whether this gene replacement affects HCMV mCherry replication

in trophoblasts, the cells were infected with the clinical isolate 181031E (**Fig. 1B**) (40). 181031E

was isolated from the urine of a congenitally infected neonate and propagated exclusively on

ARPE-19 cells to generate high-titer stocks after five passages. As with HCMV mCherry, very

little 181031E replication was observed in trophoblasts relative to ARPE-19 cells. Taken

together, the inability of HCMV to replicate in TSCs and TSC-derived cells does not appear to

be a strain-specific defect.

TSC, ETV, and STB are propagated in media that includes several factors that are not

routinely used when HCMV-permissive ARPE-19 cells and fibroblasts are cultured, including the

histone deacetylase inhibitor valproic acid, the TGF-β inhibitor A83-01, and the Rho-associated

protein kinase inhibitor Y27632. To test whether HCMV replication was inhibited by one or more

of these factors, ARPE-19 cells were grown in TSC self-renewal media for 3 passages and

infected with HCMV mCherry at an MOI of 0.1. While HCMV titers were lower in ARPE-19 cells

grown in TSC media relative cells grown under standard conditions, the virus could still replicate

(**Fig. 1C**). TSCs grow rapidly and must be passaged every three to four days even when

infected with HCMV at a high multiplicity. Thus, TSC overgrowth constrained the duration of our

infection studies. To assess whether HCMV would replicate in TSCs over longer time scales

and if the viral genome was maintained when cells were passaged, TSCs were infected with

HCMV mCherry at a MOI of 1 and passaged a total of six times, splitting 1:10 or 1:15 every

three to four days. DNA was extracted from the cells after each passage and the abundance of

host and viral genomes was quantified using droplet digital PCR (ddPCR) reactions specific to

NRAS and HCMV *UL55* (41). The relative abundance of viral genomes exceeded that of host

genome for the first two passages but rapidly decreased after subsequent passages; HCMV

was undetectable and apparently cleared by the fifth passage (**Fig. 1D)**. Together, our

observations suggest that TSCs and their derivatives do not support robust HCMV replication.

TSCs restrict HCMV replication after entry.

Given that mCherry expression was evidence that TSCs, EVTs, and STBs were being

infected by HCMV (**Fig. 1A**), we next assessed whether other viral proteins were produced in

infected trophoblasts. ARPE-19 cells and TSCs were infected with HCMV-mCherry at a MOI of

1 and immunostained with antibodies specific to the immediate early and late viral gene

products IE1/2 and pp65 (**Fig. 2**). In ARPE-19 infected cells, high rates of IE1/2-positivity were

detected at 24 hpi and virtually all infected cells expressed pp65 and mCherry by 72 hpi. The

proportion of infected TSCs that expressed any virally-encoded protein was comparably lower;

an estimated 10-20% of the cells expressed IE1/2. While the number of infected TSCs that

expressed pp65 and mCherry increased as time progressed, at most 30% of IE1/2 expressing

cells also expressed the other viral proteins.

To better quantify the differences in viral protein expression between cell types, TSCs

and ARPE-19 cells were infected with HCMV3F. HCMV3F encodes three fluorescent reporters

with clearly distinguishable emission spectra that are linked by 2A peptides to HCMV proteins

with immediate early (mNeonGreen-P2A-IE1/IE2), early (mTagBFP2-P2A-UL112/113), and late

(mCherry-T2A-UL48A [small capsid protein]) expression kinetics (42, 43). TSCs and ARPE-19

cells were infected at low and high infectious doses (MOIs of approximately 0.05 and 0.5). Cells

were analyzed by flow cytometry at 12, 24, 48, and 96 hpi. To measure viral protein expression

over time, we first quantified mNeonGreen fluorescence in single cells. Using this reporter of

immediate early protein expression to identify infected cells, mTagBFP2 and mCherry

fluorescence was measured in mNeonGreen[+] cells to determine whether infected cells also

expressed early and late genes (**Fig. S1**).

In ARPE-19 cells, the abundance of mNeonGreen[+] cells correlated with the initial

infectious dose and increased over time (**Fig. 3**). At later time points, this reflects viral spread

through the monolayer. The proportion of mNeonGreen[+] cells that also expressed mTagBFP2

and/or mCherry also increased as time progressed (**Fig. 4**). Consistent with the results IE1/2

immunostaining, the proportion of TSCs that expressed mNeonGreen was 10-20% of the

number of ARPE-19 cells that expressed the immediate early reporter at either infectious dose.

The number of infected, mNeonGreen-expressing TSCs that also expressed either mTagBFP2

or mCherry was low even at late times post-infection. Thus, TSCs can be infected by HCMV,

but these infections are largely unproductive and rarely progress past immediate early gene

expression.

HCMV gene expression is dysregulated in TSCs and TSC-derived cells.

RNA sequencing (RNA-Seq) was used to study host and viral gene expression in mock-

and HCMV-infected TSC and TSC-derived trophoblasts (**Fig. 5A**). TSCs, EVTs, and STBs were

mock- or HCMV mCherry-infected at a MOI of 1 and RNA was extracted for analysis at 24, 48,

and 72 hpi. For TSCs, cells were dissociated and fluorescence activated cell sorting (FACS)

was used to separate the infected cells into populations with low or high mCherry fluorescence

prior to RNA extraction (**Fig. S2**). RNA was extracted from EVTs and STBs directly as the two

cell types uniformly expressed mCherry after infection. Reads were aligned to a concatenated

reference sequence made up of the human and HCMV genomes (GRCh38 and TB40e-Bac4

[NCBI EF999921.1]).

The abundance of viral transcripts in infected trophoblasts was compared with RNA-seq

data from a recent study of HCMV-infected fibroblasts (HFF1) and CD14[+] monocytes, which

model lytic and latent infections, respectively (44). First, the proportion of viral reads among all

reads was compared across the five cell types (**Fig. 5B**). In infected fibroblasts, viral reads

accounted for approximately 50% of all reads at 72 hpi. In contrast, very low levels of viral

transcription was noted in latently infected monocytes and TSCs that expressed low levels of

mCherry. TSCs that expressed high levels of mCherry and infected EVTs and STBs had an

intermediate level of viral transcription, where the abundance of HCMV transcripts increased

over time but peaked at a level that was 10-fold lower than was observed in fibroblasts.

Viral gene expression was next compared by temporal class (44). Most HCMV

transcripts were assigned to one of seven temporal classes (TCs): immediate early (IE) genes

(TC1), true early genes (TC2), delayed early genes (TC3), early-late genes (TC4), replication

independent late genes (TC5), translation independent late genes (TC6), and true late genes

(TC7). The proportional abundance of each TC was determined over time (**Fig. 5C**), and **Figure**

S   3 illustrates the expression of individual viral transcripts in a heat map. During lytic infection,

IE transcripts were the most abundant before 24 hpi, early transcripts (TC2 and TC3) were the

most abundant at 24 hpi, and late transcripts (TC5, TC6, and TC7) predominate at later times

(44). Low-level expression of viral transcripts from all seven TCs was observed in latently-

infected monocytes, where comparably little change in the relative abundance of the TCs

occurred over time (44). Viral gene expression in infected EVTs and STBs was the most like

lytic infection in fibroblasts. In these cell types, early transcripts (TC2 and TC3) were the most

abundant at 24 hpi and late transcripts (TC6 and TC7) increased in abundance at 48 and 72

hpi. Two patterns of viral gene expression were noted in TSCs. There was no evidence of

temporally coordinated viral gene expression in infected TSCs that expressed low levels of

mCherry. In cells that expressed high levels of mCherry, high levels of early transcripts (TC2

and TC3) persisted until 72 hpi, when many late genes were not expressed to their expected

level. Thus, RNA-Seq revealed an unusual pattern of viral gene expression in HCMV-infected

TSCs, which is consistent with the defects in early and late protein expression that were

observed by immunofluorescence and flow cytometry experiments.

HCMV does not trigger a canonical antiviral response from infected trophoblasts.

To assess how HCMV infection affects host transcription, gene expression in mock- and

HCMV-infected cells was compared. Principal component analysis (PCA) revealed that the

samples of TSCs and EVTs clustered according to when the cells had been collected (**Fig. 6A**).

There was also separation between populations of mock- and HCMV-infected EVTs and STBs.

Across all three cell types and time points, infection upregulated more transcripts than it

downregulated (≥2fold, $p < 0.05$). Differential gene expression analysis revealed that HCMV

infection had the greatest effect on transcription in STBs, where nearly 3000 genes were

differentially regulated by 24 hpi (**Fig. 6B**). Infection had the least effect on TSCs that expressed

low levels of mCherry; at most 394 transcripts were affected by HCMV in these cells. EVTs and

TSCs that expressed high levels of mCherry had an intermediate phenotype: about 300

transcripts were differentially regulated in infected cells at 24 hpi and the number of differentially

expressed transcripts increased at later times post-infection. Gene set enrichment analysis was

used to illuminate host pathways that were affected by infection (45). The molecular functions

and biological processes that were most highly enriched among infection-regulated transcripts

were largely related to developmental processes, metabolic regulation, and cellular

morphogenesis (**Fig. 6C**). Terms like "response to virus (GO:0009615)" that include transcripts

that are canonically associated with an antiviral response were significantly enriched in some

cases, but the absolute number of genes that were differentially regulated was low. Pathways

related to inflammatory signaling and immune activation were enriched in some cell types and

times.

We next examined the expression of interferons and cytokines, interferon stimulated

genes (ISGs), and factors involved in trophoblast differentiation or cell type specificity in infected

and uninfected trophoblasts (**Fig. 7**). Primary trophoblasts constitutively express type III IFN,

which helps to establish and maintain an antiviral environment in the placenta and decidua,

while antiviral type I IFN responses have been found to cause placental dysfunction in some

experimental systems (15, 23, 46). Type III IFN transcripts were detected at low levels in many

samples, but HCMV infection did not significantly alter IFN expression. Gene ontogeny terms

related to cytokine activity (e.g. GO:0005125, GO:0034097) were enriched in some infected

trophoblasts. *IL1A, IL6,* and *IL11* were upregulated at every time point post-infection in EVTs

and STBs. *IL11* transcription was also upregulated in TSCs that expressed high levels of

mCherry, but this effect was only significant at 72 hpi. *TNF* was also significantly upregulated in

these infected TSCs at 48 hpi.

Interferons (IFNs) and pattern recognition receptors like RIG-I regulate the expression of

hundreds of antiviral effectors, known as interferon stimulated genes (ISGs). Using an inclusive

list that included 1905 different transcripts, we examined which ISGs were expressed by

trophoblasts and how they were regulated by HCMV infection (47, 48). In TSCs, ISG expression

was temporally regulated, and more transcripts were affected at late times post-infection and in

cells that expressed high levels of viral mCherry. While some HCMV infection regulated ISGs

encoded well known antiviral effectors like *IFIT5*, *MX1* and *OAS3,* many of the other transcripts

have functions related to cell survival and proliferation. For example, *PRAME, NFIX,* and *A2M*

are upregulated by infection and have known anti-apoptotic activities (49-51). *TRIB2* and

CCNA1 play key roles in stem cells self-renewal (52, 53). The greatest number of ISGs were

differentially regulated in STBs (347, 519, and 505 at 24 hpi, 48 hpi, and 72 hpi respectively).

Differentiated trophoblasts were like TSCs in that infection predominantly upregulated anti-

apoptotic transcripts and factors that positively regulate cellular proliferation; *TRIB2* and *PRAME*

were highly upregulated at all time points in infected STBs.

The expression of cell type-specific transcripts was analyzed to assess whether HCMV

infection affected trophoblast differentiation. A recent analysis of differentiation trajectories in

TSCs identified transcripts that could be used as markers of undifferentiated TSCs (*TEAD4*,

YAP1, *TP63*, *ITGA6* and *ELF5*), differentiating EVTs (*ITGA5*, *ITGA2*, and *NOTCH1*), mature

EVTs (*ITGA1*, *NOTCH2*, and *HLA-G*), and mature STBs (*ERVFRD-1*, *ERVV-1*, and *SDC1*) (**Fig.**

7) (54). All TSCs expressed some EVT and STB markers at 72 hpi. TSCs have been noted to

express differentiation markers when grown in three-dimensional cultures, and the expression of

these transcripts may reflect the spontaneous differentiation of some cells as cultures

propagated in self-renewal media overgrow (54). HCMV infection significantly reduced the

expression of *ERVFRD-1*, *ERVV-1, and HLA-G* in TSCs at 72 hpi. Biomarkers for both

differentiating and mature cells were detected in all EVT samples and these transcripts were not

differentially regulated by HCMV infection. However, transcripts associated with stemness, such

as *ELF5* and *TEAD4,* were significantly upregulated in HCMV-infected EVTs. TSC markers

were also upregulated in HCMV-infected STBs. Relatedly, Wingless/Integrated (WNT) signaling

regulates the proliferation and maintenance of TSCs *in vitro*, and the removal of CHIR99021, a

WNT activator, from self-renewal media triggers the differentiation of TSCs into EVTs (34, 54).

Transcripts involved in WNT signaling pathways were among the transcripts that were most

highly upregulated by HCMV infection. For example, *FZD9, NRARP, NKD1, WNT3, WNT5B,*

were upregulated in infected TSCs. HCMV infection also perturbed WNT signaling in

differentiated EVTs and STBs, where transcripts including *WNT3, WNT7B,* and *WNT9A* were

upregulated. In summary, HCMV infection had more significant effects on host gene expression

in differentiated EVTs and STBs than in TSCs. Infection did not appear to affect interferon

transcription, but some inflammatory cytokines were upregulated by infection. While TSCs

propagate after being infected with HCMV, transcripts involved in development and cell

differentiation are dysregulated by infection; whether these changes in gene expression affect

TSC function remains unclear.

Discussion.

TSCs are a recently developed and robustly characterized *in vitro* model of first-trimester

human trophoblasts (34, 37). In this study, we used TSCs to model HCMV infection in the

placenta, studying replication dynamics and the effects of infection on gene expression in

undifferentiated and differentiated trophoblasts. We observed that TSCs and TSC-derived EVTs

and STBs can be infected by HCMV but found little evidence of productive replication in any of

the three cell types. High MOI HCMV infections have no apparent effect on the proliferation of

TSCs and viral genomes are not maintained in the cells beyond the first several passages.

Analyses of viral gene expression and protein production by immunofluorescence microscopy,

flow cytometry, and RNA-Seq suggests that replication is inhibited after entry. Relative to lytic

infection in fibroblasts, viral gene expression was notably dysregulated in TSCs. Infection also

had muted effects on host transcription in TSCs when compared to similarly infected EVTs and

STBs.

Our results were unexpected given decades of prior research that includes pathologic

studies of HCMV-infected placentas and experimental infections done in tissue explants and

primary trophoblasts. The broad consensus has been that HCMV infects and replicates in CTBs

and EVTs but not STBs (5, 8, 18-20, 55-57). Chorion-derived trophoblast progenitor cells are

likely the most similar cell type to TSCs that were previously used to study HCMV infection (58).

HCMV replicates in these trophoblast progenitor cells, interfering with their ability to differentiate

(19, 21). However, recent work has found that trophoblasts derived from the smooth chorion

may be committed to a different cell fate than villous trophoblasts (59). Isolated reports have

suggested that HCMV can replicate in STBs *in vitro* (60, 61). However, contaminating maternal

and/or fetal epithelial cells can outgrow primary trophoblasts, and it is imperative to confirm that

the intended cell type is being studied throughout an experiment when cells have been isolated

from placenta (62). The CT27 and CT29 cell lines are advantageous in that the TSCs have

been robustly characterized and meet many established criteria for first trimester trophoblast

identity (34, 37, 63).

In this study, TSCs were infected with either a HCMV clinical isolate or one of two

TB40E-based "clinical-like" laboratory strains. All viral stocks that were used were prepared by

passaging virus a minimum number of times in epithelial (ARPE-19) cells and were confirmed to

infect fibroblasts and epithelial cells with comparable efficiency. This was necessary because

HCMV rapidly adapts during propagation *in vitro*, and commonly used laboratory strains (e.g.

AD169 and Towne) and other viruses propagated extensively in fibroblasts can lack a

glycoprotein complex that is required for HCMV's naturally broad cell tropism (64). As the

expression of virally encoded proteins confirmed that TSCs can be infected by HCMV, the cells

likely either express one or more restriction factors that inhibit HCMV replication or are deficient

in host dependency factors that the virus requires to replicate. HCMV can latently infect

embryonic and induced pluripotent stem cells and can reactivate when the cells differentiate

(65, 66). This suggests that HCMV may be generally unable to establish lytic infections in stem

cells. TSCs may not be broadly restrictive to viruses as Zika virus can replicate in the cells (67).

Our findings are similar to observations made in HCMV-infected trophoblast organoids.

Derived either directly from human placenta or from stem cells, trophoblast organoids are a

three-dimensional model of the first trimester placenta (35, 68). Trophoblast organoids are

largely refractory to HCMV infection but decidual organoids can be infected by the HCMV

strains AD169r and TB40E (30). Like findings in two-dimensional TSC culture, trophoblast

organoids can be readily infected by Zika virus (35, 67). Interestingly, ZIKV replicates less well

in TSC-derived EVT and STB than it does in TSCs (67). Thus, the broad antiviral properties of

primary trophoblasts may develop as the cells differentiate or as the placenta matures (15, 33,

69). Our gene expression profiling did not find strong evidence of type I or III interferon

expression in TSCs nor or that these genes to be regulated by infection.

Furthermore, while HCMV did upregulate some proinflammatory cytokines and ISGs in infected

TSCs, EVTs, and STBs, the classical antiviral response to infection was more muted than is

observed in other cell types.

HCMV infection early in gestation can cause placenta maldevelopment and early

pregnancy loss and understanding how infection affects trophoblast differentiation has been an

area of intense research focus (4). While HCMV infection did not effect the ability of TSCs to

grow *in vitro*, we did not test whether infected cells could differentiate normally. Others have

found that infected CTBs and chorionic villi have a diminished capacity to differentiate and

invade matrigel (8, 21, 55). In trophoblast progenitor cells, proteins involved in cell cycle

progression, pluripotency, and early differentiation are dysregulated after HCMV infection (19,

21). The HCMV-encoded cytokine cmvIL-10, an ortholog of human IL-10 that the virus

expresses to manipulate the immune response to infection, impairs the invasiveness of infected

trophoblasts (55, 70). WNT signaling also plays an important role in HCMV infection and

trophoblast differentiation. HCMV sequesters and degrades beta-catenin in infected fibroblasts

and SGHPL-4 cells, an EVT cell line, disrupting cellular function and potentially impairing EVT

invasion during placental development (71). Notably, we found that *IL11*, which regulates

trophoblast invasion, was upregulated by infection in most of the conditions we examined (72,

73). Given that RNA-Seq of TSCs, EVTs, and STBs revealed that the expression of cell type

biomarkers is perturbed by HCMV infection and that WNT signaling is dysregulated, further

research into how HCMV affects the capacity of TSCs to differentiate is merited.

Materials and Methods.

Cells and viruses.

ARPE-19 epithelial cells (CRL-2302) and HFF-1 human foreskin fibroblasts (SCRC-

1041) were purchased from the ATCC (SCRC-1041) and propagated according to their

instructions. CT27 and CT29 TSCs were maintained in an undifferentiated state and

differentiated into EVT or STB as previously described (34). Briefly, cell culture plastics were

coated with iMatrix-511 (ReproCELL MP892-011) recombinant laminin dissolved in TSC self-

renewal media (0.05 µg/cm^2) for at least 15 minutes before cells were seeded. TSC self-renewal

medium contained DMEM/F12 (ThermoFisher 11320033) supplemented with 1% KnockOut

Serum Replacement (ThermoFisher 10828010), 0.6% Penicillin-Streptomycin (ThermoFisher

15140122), 0.15% BSA (Millipore Sigma A1470), 1% Insulin-Transferrin-Selenium-

Ethanolamine (ITS -X) (ThermoFisher 51500056), 0.2 mM L-ascorbic acid (Millipore Sigma,

A8960), 25 ng/ml EGF (R&D Systems, 236EG200), 2 µM CHIR99021 (Tocris Bioscience 4423),

5 µM A83-01 (Tocris Bioscience 2939), 0.8 mM valproic acid (Millipore Sigma 676380), and 2.5

µM Y-27632 (Tocris Bioscience 1254). TSCs were dissociated by washing the cells with dPBS

and incubating with a 1:1 mixture of dPBS and TrypLE Express Enzyme (ThermoFisher

12604013). TSCs were passaged every 2 to 3 days (upon reaching ~80% confluence) at a ratio

of 1:10 to 1:15.

For EVT and STB differentiations, cell culture vessels were coated with either 1 µg/ml

(EVT) or 2.5 µg/ml (STB) of mouse Collagen IV (Corning™ 354233). For EVT differentiation,

TSCs were seeded at a density of 7.8X10^3 cells/cm^2 in DMEM/F12 supplemented with 0.1 mM

2-mercaptoethanol (Gibco 21985-023), 0.5% Penicillin-Streptomycin (Gibco 15140-122), 0.3%

BSA (Millipore Sigma A1470-25G), 1% ITS-X supplement (Gibco 51500-056), 100 ng/ml NRG1

(Cell Signaling Technology 5218SC), 7.5 µM A83-01 (Tocris Bioscience 2939/10), 2.5 µM

Y27632 (Tocris Bioscience 1254/10), and 4% KnockOut Serum Replacement (ThermoFisher

10828-010), and 2% Matrigel (Corning 354234). At 3 days post-differentiation, EVT media was

added that was formulated as described except in that it contained Matrigel at 0.5% and did not

contain NRG. For STB differentiation, TSCs were seeded at a density of $2.1X10^4$ cells/cm^2 in

DMEM/F12 supplemented with 0.1 mM 2-mercaptoethanol, 0.5% penicillin-streptomycin, 0.3%

BSA, 1% ITS-X supplement, 2.5 µM Y27632, 2 µM forskolin (Millipore Sigma F6886), and 4%

KSR. STB media was replaced after 3 days.

TB40/Ewt-mCherry (HCMV mCherry) was a gift from Dr. Wade Bresnahan (University of

Minnesota) (38). Infectious virus was reconstituted from the BAC by transfection into human

fibroblasts by Dr. Bresnahan and passaged three times in ARPE-19 cells to generate the viral

stocks used in this project. The clinical isolate 181031E was isolated from the urine of a

congenitally infected child and passaged exclusively through ARPE-19 cells five times to

generate the high titer stocks used in this project (41). The HCMV3F BAC was a gift from Dr.

Luka Čičin-Šain (Helmholtz-Zentrum für Infektionsforschung) (42). Infectious virus was

reconstituted by transfecting ARPE-19 cells with BAC DNA and Lipofectamine 3000

(ThermoFisher L3000001); high titer stocks generated after three passages in ARPE-19 cells

were used for this project. The titers of HCMV stocks were calculated by $TCID_{50}$ assays on both

HFF-1 and ARPE-19 cells to ensure that broad tropism was maintained.

Assaying HCMV infection and replication in trophoblasts.

ARPE-19s, TSCs, EVTs, and STBs were seeded to 12-well plates as described above 3

days before infection. Each well was infected with $2.5X10^4$ PFU of HCMV for an approximate

MOI of 0.1. Fluorescent micrographs of HCMV mCherry-infected cells were taken using an

EVOS Cell Imaging System. At 1, 2, 3, and 4 dpi, cells were scraped into the media, collected,

and flash frozen. To calculate HCMV titer, these samples were freeze-thawed a total of three

times, serially diluted, and used to infect HFF-1 monolayers in plaque assays as previously

described (74).

To assess the effects of TSC self-renewal media on HCMV replication, ARPE-19 cells

were adapted to propagate in TSC self-renewal medium for 3 passages. These adapted and

control cells were seeded onto a 12-well plate and infected with $2.5X10^4$ PFU of HCMV mCherry

for an approximate MOI of 0.1. Cells were scraped into the media, collected, and flash frozen at

24, 48, and 72 hpi. To calculate HCMV titer, samples were freeze-thawed three times, serially

diluted, and used to infect HFF-1 monolayers in plaque assays as previously described (74).

To assess the fate of HCMV-infected TSCs upon passage, TSCs were infected with

HCMV mCherry at a MOI of 1. The cells were passaged every 3-4 days at a ratio of 1:10 to 1:15

a total of six times. After each passage, DNA was extracted from a portion of the remaining cells

using the DNeasy Blood and Tissue kit (Qiagen). The relative abundance of human and viral

genomes was assessed by droplet digital PCR using primers and probe specific to human

HCMV *UL55* (FAM) and *NRAS* (Hex) as previously described (41, 75).

Immunofluorescence assay.

TSCs and ARPE-19 cells were seeded to an ibiTreat 4-well μ-Slide (ibidi 80426) at a

density of $2X10^3$ cells/cm^2 and grown overnight. The cells were infected with HCMV mCherry at

a MOI of 1. After a 2 hour absorption, the cells were washed with dPBS and fresh media was

added. Cells were immunostained at 24, 48, and 72 hpi. Media was aspirated and the cells were

fixed with 4% paraformaldehyde (Sigma Aldrich P6148-1KG), permeabilized with 0.2% Triton X-

100 (Sigma Aldrich 11332481001), and blocked with a 10% normal donkey serum (Sigma

Millipore D9663-10). The cells were stained with mouse α-IE-1 (Ms. X pp72 mAb, Sigma 8B1.2)

and rabbit α-pp65 (Biorbyt orb10511) primary antibodies at 1:200 dilutions, then washed with

blocking solution. Donkey α-mouse AF488 secondary antibody (Jackson ImmunoResearch 715-

545-150) and donkey α-rabbit Cy5 secondary antibody (Jackson ImmunoResearch 711-175-

152) were used at a dilution of 1:250. After a final wash, the cells were mounted and nuclei

stained using Mounting Medium with DAPI (ibidi 50011). Fluorescent micrographs were taken

with a Nikon Ti-E Inverted Deconvolution Microscope and image analyses done with NIS-

Elements Basic Research (version 4.60).

Flow cytometry.

Fluorescent protein expression vectors were cloned by PCR amplifying mNeonGreen

(pCJB139 F: 5' CACTAGTCCAGTGTGGTGGAATTGCCCTTACCATGGTGAGCAAGGGCGAG,

R: 5' TCGAGCGGCCGCCACTGTGCTGGACTTGTACAGCTCGTCCATGC) and mTagBFP2

(pCJB138 F: 5'

CACTAGTCCAGTGTGGTGGAATTGCCCTTACCATGGGACCTAAGAAGAAGAGAAAGG, R: 5'

TCGAGCGGCCGCCACTGTGCTGGAATTAAGCTTGTGCCCCAGTTTGC) from the HCMV3F

BAC. The resulting amplicons were cloned into pKTS786 (a pcDNA 3.1/V5-His based vector)

that had been linearized with BstX1 with an In-Fusion HD cloning kit (Takara 638909) (76).

These vectors were transfected into ARPE-19 cells with Lipofectamine 3000 to use as single-

stain controls for flow cytometry. ARPE-19 cells were infected with HCMV mCherry at a MOI of

0.1 and used as a single stain mCherry control.

TSCs and ARPE-19 cells were seeded at densities of 1×10^4 cells/cm^2. Twenty-four

hours later, the cells were infected with HCMV3F at 1×10^3 or 1×10^4 PFU/cm^2 for approximate

MOIs of 0.05 and 0.5. After adsorbing for 2 hours, the infected cells were washed with dPBS

and cell type-specific media added. Infected cells and single-stain controls were harvested by

dissociation for flow cytometry at 12, 24, 48, 72, and 96 hpi. The dissociated cells were

resuspended in a fixative buffer (0.5% BSA, 0.01% sodium azide in sterile dPBS). Cells were

analyzed using a BD LSRFortessa X-20 Cell Analyzer with the same gating voltages for all time

points. One hundred thousand events were recorded for each time point, and were gated in the

following order: cells, single cells, single stain positive (for each fluorophore), and

BFP+/mCherry+ cells within the mNeonGreen+ cell population. Flow analysis was done using

FlowJo (version 10.8.0).

RNA-Sequencing and bioinformatics analyses

For RNA-seq of HCMV-infected TSCs, 2×10^5 CT27 or CT29 cells were seeded on T25

flasks and infected two days later with 7.5×10^5 PFU of HCMV mCherry (approximate MOI of 1).

Mock- and HCMV infected cells were trypsinized and collected at 1, 2, and 3 dpi, and stained

with Ghost Dye Violet 450 (Tonbo Bioscences) for florescence-activated cell sorting with a

FACSAria II cytometer (BD Biosciences). Cells were sorted directly into 2-Mercaptoethanol-

containing RLT buffer and RNA was extracted using a RNeasy Mini kit (Qiagen). The cell

populations used for RNA sequencing are summarized in **Table S1**. Sequencing libraries were

prepared by the University of Minnesota Genomics Center using the SMARTer Pico Mammalian

V2 kit (Takara). For RNA-Seq from EVTs and STBs, CT27 and CT29 were seeded into 6-well

plates ($7.5X10^4$ and $2X10^5$ cells/well, respectively) and differentiated for 3 days as described

above. At 3 days post-differentiation, each well was infected at an approximate MOI of 1 with

$2X10^5$ PFU of HCMV mCherry. RNA was isolated by directly lysing the cells with RLT buffer and

using RNeasy Mini kits and used to prepare TruSeq Stranded mRNA (Illumina) sequencing

libraries. Libraries were pooled and sequenced on an Illumina NovaSeq (S4 Flow Cell, 2x150-

bp PE reads). A sequencing depth of >40M reads was obtained for TSC samples and >20M

reads for EVT and STB samples; mean quality scores for all sequencing libraries were >Q30.

Sequencing data generated in this project was analyzed along with publically-available

data from HCMV TB40E-GFP-infected fibroblasts and CD14+ monocytes (Gene Expression

OmnibusGSE193467) (44). Sequencing reads were trimmed using Trimmomatic at default

parameters (77) and aligned to a concatenated reference file that contained the human

(GRCh38) and TB40e-BAC4 (NCBI EF999921.1) genomes with STAR version 2.7.2a using the

default parameters (78, 79). Gene count tables were prepared with featureCounts (80). Gene

expression analysis of host transcripts was done using PartekFlow (version 11.0.23.1004). After

filtering out transcripts with fewer than 20 gene counts, transcript abundance was normalized

using the median ratio method, and DESeq2 was used to identify differentially regulated

transcripts when mock- and HCMV-infected samples were compared on a time and cell-type

basis (81). Principal component analysis was used to dimensionally reduce and visualize data

from each cell type. Differentially expressed genes were found to be statistically significant

when they were upregulated or downregulated 2-fold or more, and DESeq2 calculated p-values

were 0.05 or less. The expression of select genes of interest was visualized by heat map. To

analyze the pattern of HCMV gene expression, viral reads were filtered from the host. The

proportional abundance of each transcript in each sample was calculated by dividing its gene

count by the total number of viral gene counts. RNA-Seq data have been deposited in NCBI's

Gene Expression Omnibus and are accessible through GEO Series accession number

GSE248772 (82).

Acknowledgments.

This project was supported by the Eunice Kennedy Shriver National Institute of Child Health and

Human Development (R21HD087496 and R01HD109252 to CJB) and the Minnesota Masonic

Charities (Masonic Cross-Departmental Grant in Children's Health Research to CJB and VJB).

TB40/Ewt-mCherry and HCMV3F were generously shared by Dr. Wade Bresnahan (University

of Minnesota) and Luka Čičin-Šain (Helmholtz Centre for Infection Research). This work was

supported by the resources and staff at the University of Minnesota University Imaging Centers

(SCR_020997), the Genomics Center (https://genomics.umn.edu) and the Flow Cytometry

Resource.

Figure 1. HCMV does not efficiently replicate in TSCs or TSC-derived trophoblasts. TSCs,

TSC-derived EVTs and STBs, and ARPE-19 cells were infected with either HCMV mCherry or

the HCMV clinical isolate 181031E at a MOI of 0.1. **(A)** Representative phase contrast and

fluorescent micrographs show mCherry expression by HCMV mCherry-infected ARPE-19 cells

and trophoblasts (CT29) at 2 dpi (Scale bar = 50 µm). **(B)** Cells were scraped and pooled with

supernatant daily. HCMV production was measured by plaque assay on fibroblasts. (**C**) ARPE-

19 cells were adapted to grow on TSC self-renewal medium for three passages and infected

with HCMV. Virus production was measured as described above and compared with cells that

had been grown under standard conditions. (**D**) HCMV mCherry-infected CT29s were passaged

1:10 to 1:15 six times. After each passage, the abundance of *UL55* and *NRAS* was quantified

by droplet digital PCR. (ND: none detected).

Figure 2. Fewer HCMV mCherry-infected TSCs express pp65 and mCherry than express

IE1/2. ARPE-19 cells and TSCs were infected with HCMV mCherry at a MOI of 1 and

immunostained with α-HCMV IE1/2 and pp65 at 24 and 72 hpi. Nuclei were stained with DAPI

and the expression of virally-encoded mCherry was also assessed. The merged images show

data from the DAPI, IE1/2, and pp65 channels and exclude mCherry data.

Figure 3. Fewer HCMV3F-infected TSCs express mNeonGreen than similarly-infected

ARPE-19 cells. ARPE-19 cells and TSCs were infected at low and high MOIs with HCMV3F.

Cells were harvested at 12, 24, 48, and 96 hpi and fluorescent protein expression measured by

flow cytometry. mNeonGreen fluorescence is shown in the histograms, and the percent

mNeonGreen-expressing cells in each population and time point is shown.

Figure 4. The expression of reporter proteins with early and late expression kinetics are suppressed in HCMV3F-infected TSCs. ARPE-19 cells and TSCs were infected with HCMV3F and the expression of virally-encoded fluorescent proteins was measured by flow cytometry. **(A)** Gating strategy used to quantify mTagBFP2 and mCherry fluorescence in mNeonGreen+ cells. Representative data from ARPE-19 low MOI and TSC high MOI infections at 24 and 90 hpi is shown. **(B)** The proportion of mTagBFP2 and mCherry-expressing cells, as determined by quad gating, is shown over time.

Figure 5. **Infected TSCs and TSC-derived trophoblasts have distinct viral gene expression profiles when compared to infected fibroblasts or monocytes.** TSC, EVT, and STB were infected with HCMV mCherry and RNA was extracted and sequenced at the indicated time points. Viral gene expression was compared with previously-published data from infected fibroblasts (lytic infection) and monocytes (latent expression) (44). (**A**) Summary of sample collection for RNA-sequencing. TSC-derived EVT and STB were grown in differentiation media for 3 days, mock- or HCMV mCherry-infected, and RNA was extracted at 1, 2, and 3 dpi. TSCs were infected similarly but sorted into populations with low or high expression of mCherry before RNA extraction. Figure created with the Biorender. (**B**) Illumina sequencing reads were mapped to the host and viral genomes. The proportion of reads that mapped to the viral genome at each time point is shown. (**C**) Viral reads were classified by temporal class and the abundance of each temporal class, expressed as a proportion of all viral reads, is shown over time.

Figure 6. HCMV infection dysregulates more transcripts in differentiated trophoblasts

than in TSCs. (A) PCA plots illustrate dimensionally-reduced gene expression in mock- and

HCMV-infected TSCs, ETVs, and STBs. **(B)** Table summarizing the number of differentially

regulated host transcripts in each cell type at 24, 48, and 72 hpi (≥2 fold, $p < 0.05$). (**C**) Gene set

enrichment analysis was used to identify GO terms that were enriched among transcripts that

were differentially expressed after infection. The results for select terms are illustrated by a

bubble chart.

Figure 7. HCMV infection dysregulates transcripts involved in cell differentiation and

inflammatory responses. Host gene expression was normalized using Z-scoring across all

samples and organized by time point, infection status, and cell type. Representative transcripts

encoding type I, II, and II interferons, IL-1 and IL-6 family cytokines, trophoblast biomarkers, and

regulators of WNT signaling were assembled and grouped.

Figure S1. Gating strategy used to analyze HCMV3F reporter protein expression.

Representative flow cytometry data is shown from the 24 hpi ARPE-19 Low MOI sample. Single

cells were identified using forward and side scatter. After gating for mNeonGreen-expressing

cells, the expression of virally encoded mBFP2 and mCherry was measured.

Figure S2. Gating strategy for fluorescent activated cell sorting prior to TSC RNA-Seq.

TSCs were mock- or HCMV mCherry-infected. At 24, 48, and 72 hpi cells were dissociated and

divided into populations that expressed high or low levels of mCherry using fluorescent

activated cell sorting (FACS). Representative data from mock- and HCMV- infected TSC (CT29)

cells collected at 72 hpi is shown. Cells were identified by forward scatter area versus side

scatter area. Single cells were identified by using forward and side scatter heights first then

secondarily gating by forward and side scatter widths. Single cells were gated by mCherry

expression on the X-axis and Ghost Violet 450 on the Y-axis to sort live and dead cells, well as

mCherry expressing cells. Three populations of cells were collected for RNA extraction and

sequencing at each time point (summarized in Table S1): mock-infected mCherry[(-)] cells,

infected mCherry[(high)] cells, and pools of infected mCherry[(-)] and mCherry[(low)] cells.

Figure S3. Heat map illustrating normalized HCMV gene expression. The relative

expression of HCMV gene products was normalized by dividing the abundance of the

transcripts by the total number of viral reads in that sample and scaled using a standard Z-scale

across all genes. Viral transcripts were organized by temporal class, cell type, and time post-

infection. Similarly processed, previously published data from HCMV-infected HFF-1 and

CD14+ Monocytes is shown to illustrate gene expression patterns during lytic and latent

infections (44).

References.

1. Zuhair M, Smit GSA, Wallis G, Jabbar F, Smith C, Devleesschauwer B, Griffiths P. 2019. Estimation of the worldwide seroprevalence of cytomegalovirus: A systematic review and meta-analysis. Rev Med Virol 29:e2034.

2. Ssentongo P, Hehnly C, Birungi P, Roach MA, Spady J, Fronterre C, Wang M, Murray-Kolb LE, Al-Shaar L, Chinchilli VM, Broach JR, Ericson JE, Schiff SJ. 2021. Congenital Cytomegalovirus Infection Burden and Epidemiologic Risk Factors in Countries With Universal Screening: A Systematic Review and Meta-analysis. JAMA Network Open 4:e2120736-e2120736.

3. Diseases NCfIaR. 2018. CMV Fact Sheet for Healthcare Providers.

4. Gomes Pinto Garcia A, Ferreira Fonseca E, Lucia De Souza Marques R, Yamaki Lobato Y. 1989. Placental morphology in cytomegalovirus infection. Placenta 10:1-18.

5. Mühlemann K, Miller RK, Metlay L, Menegus MA. 1992. Cytomegalovirus infection of the human placenta: An immunocytochemical study. Human Pathology 23:1234-1237.

6. Pereira L, Petitt M, Fong A, Tsuge M, Tabata T, Fang-Hoonver J, Maidji E, Zydek M, Zhou Y, Inoue N, Loghavi S, Pepkowitz S, Kauvar LM, Ogunyemi D. 2014. Intrauterine growth restriction caused by underlying congenital cytomegalovirus infection. The Journal of infectious diseases 209:1573-1584.

7. Tabata T, Petitt M, Fang-Hoover J, Zydek M, Pereira L. 2016. Persistent Cytomegalovirus Infection in Amniotic Membranes of the Human Placenta. The American Journal of Pathology 186:2970-2970.

8. Fisher S, Genbacev O, Maidji E, Pereira L. 2000. Human cytomegalovirus infection of placental cytotrophoblasts in vitro and in utero: implications for transmission and pathogenesis. J Virol 74:6808-20.

9.   Fowler KB, Stagno S, Pass RF, Britt WJ, Boll TJ, Alford CA. 1992. The outcome of

congenital cytomegalovirus infection in relation to maternal antibody status. N Engl J

Med 326:663-7.

10. Lazzarotto T, Varani S, Guerra B, Nicolosi A, Lanari M, Landini MP. 2000. Prenatal

indicators of congenital cytomegalovirus infection. J Pediatr 137:90-5.

11. Pass RF, Fowler KB, Boppana SB, Britt WJ, Stagno S. 2006. Congenital

cytomegalovirus infection following first trimester maternal infection: symptoms at birth

and outcome. J Clin Virol 35:216-20.

12. Faure-Bardon V, Magny JF, Parodi M, Couderc S, Garcia P, Maillotte AM, Benard M,

Pinquier D, Astruc D, Patural H, Pladys P, Parat S, Guillois B, Garenne A, Bussieres L,

Guilleminot T, Stirnemann J, Ghout I, Ville Y, Leruez-Ville M. 2019. Sequelae of

Congenital Cytomegalovirus Following Maternal Primary Infections Are Limited to Those

Acquired in the First Trimester of Pregnancy. Clin Infect Dis 69:1526-1532.

13. Megli CJ, Coyne CB. 2021. Infections at the maternal–fetal interface: an overview of

pathogenesis and defence. Nature Reviews Microbiology 2021 doi:10.1038/s41579-021-

00610-y:1-16.

14. Delorme-Axford E, Donker RB, Mouillet JF, Chu T, Bayer A, Ouyang Y, Wang T, Stolz

DB, Sarkar SN, Morelli AE, Sadovsky Y, Coyne CB. 2013. Human placental trophoblasts

confer viral resistance to recipient cells. Proceedings of the National Academy of

Sciences of the United States of America 110:12048-12053.

15. Bayer A, Lennemann NJ, Ouyang Y, Bramley JC, Morosky S, Marques ETDA, Cherry S,

Sadovsky Y, Coyne CB. 2016. Type III Interferons Produced by Human Placental

Trophoblasts Confer Protection against Zika Virus Infection. Cell Host and Microbe

19:705-712.

16. Corry J, Arora N, Good CA, Sadovsky Y, Coyne CB. 2017. Organotypic models of type

III interferon-mediated protection from Zika virus infections at the maternal–fetal

interface. Proceedings of the National Academy of Sciences of the United States of
America 114:9433-9438.

17. Bayer A, Delorme-Axford E, Sleigher C, Frey TK, Trobaugh DW, Klimstra WB, Emert-
Sedlak LA, Smithgall TE, Kinchington PR, Vadia S, Seveau S, Boyle JP, Coyne CB,
Sadovsky Y. 2015. Human trophoblasts confer resistance to viruses implicated in
perinatal infection. American Journal of Obstetrics & Gynecology 212:71.e1-71.e8.

18. Sinzger C, Muntefering H, Loning T, Stoss H, Plachter B, Jahn G. 1993. Cell types
infected in human cytomegalovirus placentitis identified by immunohistochemical double
staining. Virchows Arch A Pathol Anat Histopathol 423:249-56.

19. Zydek M, Petitt M, Fang-Hoover J, Adler B, Kauvar LM, Pereira L, Tabata T. 2014.
HCMV infection of human trophoblast progenitor cells of the placenta is neutralized by a
human monoclonal antibody to glycoprotein B and not by antibodies to the pentamer
complex. Viruses 6:1346-64.

20. Genbacev O, Larocque N, Ona K, Prakobphol A, Garrido-Gomez T, Kapidzic M,
Bárcena A, Gormley M, Fisher SJ. 2016. Integrin α4-positive human trophoblast
progenitors: functional characterization and transcriptional regulation. Hum Reprod
31:1300-14.

21. Tabata T, Petitt M, Zydek M, Fang-Hoover J, Larocque N, Tsuge M, Gormley M, Kauvar
LM, Pereira L. 2015. Human cytomegalovirus infection interferes with the maintenance
and differentiation of trophoblast progenitor cells of the human placenta. J Virol 89:5134-
47.

22. Megli CJ, Coyne CB. 2022. Infections at the maternal-fetal interface: an overview of
pathogenesis and defence. Nat Rev Microbiol 20:67-82.

23. Yockey LJ, Jurado KA, Arora N, Millet A, Rakib T, Milano KM, Hastings AK, Fikrig E,
Kong Y, Horvath TL, Weatherbee S, Kliman HJ, Coyne CB, Iwasaki A. 2018. Type I
interferons instigate fetal demise after Zika virus infection. Sci Immunol 3.

24. Bao M, Hofsink N, Plosch T. 2022. LPS versus Poly I:C model: comparison of long-term

effects of bacterial and viral maternal immune activation on the offspring. Am J Physiol

Regul Integr Comp Physiol 322:R99-R111.

25. Scott GM, Chow SS, Craig ME, Pang CN, Hall B, Wilkins MR, Jones CA, Lloyd AR,

Rawlinson WD. 2012. Cytomegalovirus infection during pregnancy with maternofetal

transmission induces a proinflammatory cytokine bias in placenta and amniotic fluid. J

Infect Dis 205:1305-10.

26. Bourgon N, Fitzgerald W, Aschard H, Magny JF, Guilleminot T, Stirnemann J, Romero

R, Ville Y, Margolis L, Leruez-Ville M. 2022. Cytokine Profiling of Amniotic Fluid from

Congenital Cytomegalovirus Infection. Viruses 14.

27. Gervasi MT, Romero R, Bracalente G, Chaiworapongsa T, Erez O, Dong Z, Hassan SS,

Yeo L, Yoon BH, Mor G, Barzon L, Franchin E, Militello V, Palu G. 2012. Viral invasion

of the amniotic cavity (VIAC) in the midtrimester of pregnancy. J Matern Fetal Neonatal

Med 25:2002-13.

28. Weisblum Y, Panet A, Zakay-Rones Z, Vitenshtein A, Haimov-Kochman R, Goldman-

Wohl D, Oiknine-Djian E, Yamin R, Meir K, Amsalem H, Imbar T, Mandelboim O, Yagel

S, Wolf DG. 2015. Human cytomegalovirus induces a distinct innate immune response

in the maternal-fetal interface. Virology 485:289-96.

29. Weisblum Y, Oiknine-Djian E, Vorontsov OM, Haimov-Kochman R, Zakay-Rones Z, Meir

K, Shveiky D, Elgavish S, Nevo Y, Roseman M, Bronstein M, Stockheim D, From I,

Eisenberg I, Lewkowicz AA, Yagel S, Panet A, Wolf DG. 2017. Zika Virus Infects Early-

and Midgestation Human Maternal Decidual Tissues, Inducing Distinct Innate Tissue

Responses in the Maternal-Fetal Interface. J Virol 91:e01905-16.

30. Yang L, Semmes EC, Ovies C, Megli C, Permar S, Gilner JB, Coyne CB. 2022. Innate

immune signaling in trophoblast and decidua organoids defines differential antiviral

defenses at the maternal-fetal interface. Elife 11:e79794.

31.   Soncin F, Khater M, To C, Pizzo D, Farah O, Wakeland A, Arul Nambi Rajan K, Nelson

712   KK, Chang CW, Moretto-Zita M, Natale DR, Laurent LC, Parast MM. 2018. Comparative

713   analysis of mouse and human placentae across gestation reveals species-specific

714   regulators of placental development. Development 145:dev156273.

715   32.   Bayer A, Lennemann NJ, Ouyang Y, Bramley JC, Morosky S, Marques ET, Jr., Cherry

716   S, Sadovsky Y, Coyne CB. 2016. Type III Interferons Produced by Human Placental

717   Trophoblasts Confer Protection against Zika Virus Infection. Cell Host Microbe 19:705-

718   12.

719   33.   Delorme-Axford E, Donker RB, Mouillet JF, Chu T, Bayer A, Ouyang Y, Wang T, Stolz

720   DB, Sarkar SN, Morelli AE, Sadovsky Y, Coyne CB. 2013. Human placental trophoblasts

721   confer viral resistance to recipient cells. Proc Natl Acad Sci U S A 110:12048-53.

722   34.   Okae H, Toh H, Sato T, Hiura H, Takahashi S, Shirane K, Kabayama Y, Suyama M,

723   Sasaki H, Arima T. 2018. Derivation of Human Trophoblast Stem Cells. Cell Stem Cell

724   22:50-63 e6.

725   35.   Karvas RM, Khan SA, Verma S, Yin Y, Kulkarni D, Dong C, Park KM, Chew B, Sane E,

726   Fischer LA, Kumar D, Ma L, Boon ACM, Dietmann S, Mysorekar IU, Theunissen TW.

727   2022. Stem-cell-derived trophoblast organoids model human placental development and

728   susceptibility to emerging pathogens. Cell Stem Cell 29:810-825 e8.

729   36.   Takahashi S, Okae H, Kobayashi N, Kitamura A, Kumada K, Yaegashi N, Arima T.

730   2019. Loss of p57(KIP2) expression confers resistance to contact inhibition in human

731   androgenetic trophoblast stem cells. Proc Natl Acad Sci U S A 116:26606-26613.

732   37.   Sheridan MA, Zhao X, Fernando RC, Gardner L, Perez-Garcia V, Li Q, Marsh SGE,

733   Hamilton R, Moffett A, Turco MY. 2021. Characterization of primary models of human

734   trophoblast. Development 148.

735   38.   O'Connor CM, Shenk T. 2011. Human cytomegalovirus pUS27 G protein-coupled

736   receptor homologue is required for efficient spread by the extracellular route but not for

737   direct cell-to-cell spread. J Virol 85:3700-7.

738   39.   Sinzger C, Hahn G, Digel M, Katona R, Sampaio KL, Messerle M, Hengel H,

739   Koszinowski U, Brune W, Adler B. 2008. Cloning and sequencing of a highly productive,

740   endotheliotropic virus strain derived from human cytomegalovirus TB40/E. J Gen Virol

741   89:359-368.

742   40.   Putri DS, Berkebile ZW, Mustafa HJ, Fernandez-Alarcon C, Abrahante JE, Schleiss MR,

743   Bierle CJ. 2020. Cytomegalovirus infection elicits a conserved chemokine response from

744   human and guinea pig amnion cells. Virology 548:93-100.

745   41.   Sedlak RH, Cook L, Cheng A, Magaret A, Jerome KR. 2014. Clinical utility of droplet

746   digital PCR for human cytomegalovirus. J Clin Microbiol 52:2844-8.

747   42.   Rand U, Kubsch T, Kasmapour B, Cicin-Sain L. 2020. A Novel Triple-Fluorescent HCMV

748   Strain Reveals Gene Expression Dynamics and Anti-Herpesviral Drug Mechanisms.

749   Front Cell Infect Microbiol 10:536150.

750   43.   Ryan MD, Drew J. 1994. Foot-and-mouth disease virus 2A oligopeptide mediated

751   cleavage of an artificial polyprotein. EMBO J 13:928-33.

752   44.   Rozman B, Nachshon A, Levi Samia R, Lavi M, Schwartz M, Stern-Ginossar N. 2022.

753   Temporal dynamics of HCMV gene expression in lytic and latent infections. Cell Rep

754   39:110653.

755   45.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K,

756   Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese

757   JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. 2000. Gene ontology: tool for

758   the unification of biology. The Gene Ontology Consortium. Nat Genet 25:25-9.

759   46.   Lazear HM, Schoggins JW, Diamond MS. 2019. Shared and Distinct Functions of Type I

760   and Type III Interferons. Immunity 50:907-923.

761   47.   Roesch F, OhAinle M, Emerman M. 2018. A CRISPR screen for factors regulating

762   SAMHD1 degradation identifies IFITMs as potent inhibitors of lentiviral particle delivery.

763   Retrovirology 15:26.

764   48.   OhAinle M, Helms L, Vermeire J, Roesch F, Humes D, Basom R, Delrow JJ, Overbaugh

765   J, Emerman M. 2018. A virus-packageable CRISPR screen identifies host factors

766   mediating interferon inhibition of HIV. Elife 7.

767   49.   Xu Y, Yue Q, Wei H, Pan G. 2015. PRAME induces apoptosis and inhibits proliferation

768   of leukemic cells in vitro and in vivo. Int J Clin Exp Pathol 8:14549-55.

769   50.   Walker M, Li Y, Morales-Hernandez A, Qi Q, Parupalli C, Brown S, Christian C,

770   Clements WK, Cheng Y, McKinney-Freeman S. 2023. An NFIX-mediated regulatory

771   network governs the balance of hematopoietic stem and progenitor cells during

772   hematopoiesis. Blood Adv 7:4677-4689.

773   51.   Vandooren J, Itoh Y. 2021. Alpha-2-Macroglobulin in Inflammation, Immunity and

774   Infections. Front Immunol 12:803244.

775   52.   Fang Y, Zekiy AO, Ghaedrahmati F, Timoshin A, Farzaneh M, Anbiyaiee A, Khoshnam

776   SE. 2021. Tribbles homolog 2 (Trib2), a pseudo serine/threonine kinase in tumorigenesis

777   and stem cell fate decisions. Cell Commun Signal 19:41.

778   53.   McLenachan S, Menchón C, Raya A, Consiglio A, Edel MJ. 2012. Cyclin A1 is essential

779   for setting the pluripotent state and reducing tumorigenicity of induced pluripotent stem

780   cells. Stem Cells Dev 21:2891-9.

781   54.   Shannon MJ, Baltayeva J, Castellana B, Wachter J, McNeill GL, Yoon JS, Treissman J,

782   Le HT, Lavoie PM, Beristain AG. 2022. Cell trajectory modeling identifies a primitive

783   trophoblast state defined by BCAM enrichment. Development 149.

784   55.   Yamamoto-Tabata T, McDonagh S, Chang HT, Fisher S, Pereira L. 2004. Human

785   cytomegalovirus interleukin-10 downregulates metalloproteinase activity and impairs

endothelial cell migration and placental cytotrophoblast invasiveness in vitro. J Virol

78:2831-40.

56.   Gabrielli L, Losi L, Varani S, Lazzarotto T, Eusebi V, Landini MP. 2001. Complete

replication of human cytomegalovirus in explants of first trimester human placenta. J

Med Virol 64:499-504.

57.   Halwachs-Baumann G, Wilders-Truschnig M, Desoye G, Hahn T, Kiesel L, Klingel K,

Rieger P, Jahn G, Sinzger C. 1998. Human trophoblast cells are permissive to the

complete replicative cycle of human cytomegalovirus. J Virol 72:7598-602.

58.   Genbacev O, Donne M, Kapidzic M, Gormley M, Lamb J, Gilmore J, Larocque N,

Goldfien G, Zdravkovic T, McMaster MT, Fisher SJ. 2011. Establishment of human

trophoblast progenitor cell lines from the chorion. Stem Cells 29:1427-36.

59.   Marsh B, Zhou Y, Kapidzic M, Fisher S, Blelloch R. 2022. Regionally distinct trophoblast

regulate barrier function and invasion in the human placenta. Elife 11.

60.   Schleiss MR, Aronow BJ, Handwerger S. 2007. Cytomegalovirus infection of human

syncytiotrophoblast cells strongly interferes with expression of genes involved in

placental differentiation and tissue integrity. Pediatr Res 61:565-71.

61.   Hemmings DG, Kilani R, Nykiforuk C, Preiksaitis J, Guilbert LJ. 1998. Permissive

cytomegalovirus infection of primary villous term and first trimester trophoblasts. J Virol

72:4970-9.

62.   Turco MY, Moffett A. 2019. Development of the human placenta. Development 146.

63.   Lee CQ, Gardner L, Turco M, Zhao N, Murray MJ, Coleman N, Rossant J, Hemberger

M, Moffett A. 2016. What Is Trophoblast? A Combination of Criteria Define Human First-

Trimester Trophoblast. Stem Cell Reports 6:257-72.

64.   Wilkinson GW, Davison AJ, Tomasec P, Fielding CA, Aicheler R, Murrell I, Seirafian S,

Wang EC, Weekes M, Lehner PJ, Wilkie GS, Stanton RJ. 2015. Human

cytomegalovirus: taking the strain. Med Microbiol Immunol 204:273-84.

65. D'Aiuto L, Di Maio R, Heath B, Raimondi G, Milosevic J, Watson AM, Bamne M, Parks

WT, Yang L, Lin B, Miki T, Mich-Basso JD, Arav-Boger R, Sibille E, Sabunciyan S,

Yolken R, Nimgaonkar V. 2012. Human Induced Pluripotent Stem Cell-Derived Models

to Investigate Human Cytomegalovirus Infection in Neural Cells. PLoS ONE 7:49700-

49700.

66. Penkert RR, Kalejta RF. 2013. Human Embryonic Stem Cell Lines Model Experimental

Human Cytomegalovirus Latency. mBio 4.

67. Wu H, Huang XY, Sun MX, Wang Y, Zhou HY, Tian Y, He B, Li K, Li DY, Wu AP, Wang

H, Qin CF. 2023. Zika virus targets human trophoblast stem cells and prevents

syncytialization in placental trophoblast organoids. Nat Commun 14:5541.

68. Turco MY, Gardner L, Kay RG, Hamilton RS, Prater M, Hollinshead MS, McWhinnie A,

Esposito L, Fernando R, Skelton H, Reimann F, Gribble FM, Sharkey A, Marsh SGE,

O'Rahilly S, Hemberger M, Burton GJ, Moffett A. 2018. Trophoblast organoids as a

model for maternal-fetal interactions during human placentation. Nature 564:263-267.

69. Bayer A, Delorme-Axford E, Sleigher C, Frey TK, Trobaugh DW, Klimstra WB, Emert-

Sedlak LA, Smithgall TE, Kinchington PR, Vadia S, Seveau S, Boyle JP, Coyne CB,

Sadovsky Y. 2015. Human trophoblasts confer resistance to viruses implicated in

perinatal infection. Am J Obstet Gynecol 212:71 e1-71 e8.

70. Tabata T, McDonagh S, Kawakatsu H, Pereira L. 2007. Cytotrophoblasts infected with a

pathogenic human cytomegalovirus strain dysregulate cell-matrix and cell-cell adhesion

molecules: a quantitative analysis. Placenta 28:527-37.

71. Angelova M, Zwezdaryk K, Ferris M, Shan B, Morris CA, Sullivan DE. 2012. Human

cytomegalovirus infection dysregulates the canonical Wnt/β-catenin signaling pathway.

PLoS Pathog 8:e1002959.

72. Winship AL, Koga K, Menkhorst E, Van Sinderen M, Rainczuk K, Nagai M, Cuman C,

Yap J, Zhang JG, Simmons D, Young MJ, Dimitriadis E. 2015. Interleukin-11 alters

placentation and causes preeclampsia features in mice. Proc Natl Acad Sci U S A

112:15928-33.

73.   Paiva P, Salamonsen LA, Manuelpillai U, Dimitriadis E. 2009. Interleukin 11 inhibits

human trophoblast invasion indicating a likely role in the decidual restraint of trophoblast

invasion during placentation. Biol Reprod 80:302-10.

74.   Britt WJ. 2010. Human cytomegalovirus: propagation, quantification, and storage. Curr

Protoc Microbiol Chapter 14:Unit 14E 3.

75.   Meyer L, Sharon B, Huang TC, Meyer AC, Gravel KE, Schimmenti LA, Swanson EC,

Herd HE, Hernandez-Alvarado N, Coverstone KR, McCann M, Schleiss MR. 2017.

Analysis of archived newborn dried blood spots (DBS) identifies congenital

cytomegalovirus as a major cause of unexplained pediatric sensorineural hearing loss.

Am J Otolaryngol 38:565-570.

76.   Bierle CJ, Schleiss MR, Geballe AP. 2012. Antagonism of the protein kinase R pathway

by the guinea pig cytomegalovirus US22-family gene gp145. Virology 433:157-66.

77.   Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30:2114-20.

78.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,

Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15-21.

79.   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol 37:907-

915.

80.   Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics 30:923-30.

81.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biol 15:550.

82.   Edgar R, Domrachev M, Lash AE. 2002. Gene Expression Omnibus: NCBI gene

expression and hybridization array data repository. Nucleic Acids Res 30:207-10.

HPI	% mNeonGreen
12	2.83
24	9.63
48	15.40
96	24.00

HPI	% mNeonGreen
12	14.40
24	25.30
48	47.50
96	84.80

HPI	% mNeonGreen
12	0.56
24	1.54
48	1.20
96	1.54

HPI	% mNeonGreen
12	1.92
24	4.85
48	3.86
96	6.84

A.

TSC

PCA (56.19%)

PC2 (16.52%)

PC1 (39.67%)

Cell Line
• CT27
● CT29

HPI
● 24
● 48
● 72

Infection Status
● Mock
■ HCMV mCherry High/Infected
✚ HCMV mCherry Low

EVT

PCA (52.49%)

PC2 (24.84%)

PC1 (27.65%)

STB

PCA (58.57%)

PC2 (19.84%)

PCA (38.73%)

B.

Upregulated				
HPI	TSC mCherry High	TSC mCherry Low	EVT	STB
24	282	60	278	1909
48	664	111	1001	3064
72	1355	317	1683	2793

Downregulated				
HPI	TSC mCherry High	TSC mCherry Low	EVT	STB
24	7	11	17	993
48	42	25	84	1773
72	401	77	201	1566

C.

Enrichment score
● 10
● 30
● 60

p-value
1
0.05
0

Developmental process (GO:0032502)
Cell differentiation (GO:0030154)
Immune Response (GO:0006955)
Cytokine activity (GO:0005125)
Response to cytokine (GO:0034097)
Response to virus (GO:0009615)
WNT Signaling pathway (GO:0016055)

mCherry Low 24HPI
mCherry Low 48HPI
mCherry Low 72HPI
mCherry High 24HPI
mCherry High 48HPI
mCherry High 72HPI
EVT 24HPI
EVT 48HPI
EVT 72HPI
STB 24HPI
STB 48HPI
STB 72HPI