**Cytochrome P450 *CYP72A219* is Involved in Evolution of Metabolic Resistance to**

**Tembotrione in *Amaranthus palmeri* Populations**

Carlos Alberto Gonsiorkiewicz Rigon[1], Anita Küpper[2], Crystal Sparks[1], Jacob Montgomery[1], Falco

Peter[2], Simon Schepp[2], Alejandro Perez-Jones[3], Patrick J. Tranel[4], Roland Beffa[5], Franck E. Dayan[1],

Todd A. Gaines[1]

[1] Colorado State University, Department of Agriultural Biology, Fort Collins, CO 80523, USA

[2] Bayer AG, Division CropScience, Weed Control Research, 65926 Frankfurt, Germany

[3] Weed Control Platform Lead, Bayer CropScience, St Louis, MO 63141, USA

[4] University of Illinois, Department of Crop Sciences, Urbana, IL 61801, USA

[5] Senior Scientist Consultant, 65835, Liederbach, Germany

**Corresponding author:** Todd A. Gaines, todd.gaines@colostate.edu

Rigon 0000-0001-8565-7383

Küpper 0000-0001-5122-1514

Sparks

Montgomery

Peter

Schepp

Perez-Jones 0009-0002-3705-5331

Tranel  0000-0003-0666-4564

Beffa 0000-0003-3109-388X

Dayan 0000-0001-6964-2499

Gaines 0000-0003-1485-7665

## SUMMARY

- Evolution of metabolic herbicide resistance is a major issue for weed management. Few genes and regulatory mechanisms have been identified, particularly in dicotyledonous weed species. We identified putative causal genes and regulatory mechanism for tembotrione-resistance in *Amaranthus palmeri*.

- Cytochrome P450 candidate genes were identified through RNA-seq analysis. We validated their functions using heterologous expression in *S. cerevisae*. Promoters of the candidate P450 genes were analyzed. We performed QTL mapping to identify genomic regions associated with resistance.

- CYP72A219_4284 deactivated tembotrione. This gene had increased expression in other *A. palmeri* populations resistant to multiple herbicides, including-tembotrione. Resistant plants exhibited polymorphisms in the promoter of CYP72A219_4284. We identified QTLs linked to herbicide resistance, including one on chromosome 4 approximately 3 Mb away from CYP72A219_4284.

- CYP72A219_4284 is involved in tembotrione resistance in *A. palmeri*. Increased expression of this gene could be due to *cis*-regulation in the promoter, as well as *trans*-regulation from transcription factors. The elucidation of regulatory genes is crucial for developing innovative weed management approaches and target-based novel molecules.

Key Words: metabolism, monooxygenase, herbicide, hydroxylation, oxidation.

## INTRODUCTION

Evolution of herbicide resistance results from the selection pressure imposed by the frequent and consistent use of herbicides over time (Gaines *et al.*, 2020). The emergence and spread of herbicide-resistant weeds poses a major challenge to sustainable agriculture, as they can diminish crop yields and increase the costs associated with weed control (Oerke, 2006, Chauhan, 2020). Herbicide resistance in plants can be selected through different mechanisms, primarily classified as target-site (TSR) and non-target-site resistance (NTSR) (Gaines *et al.*, 2020, Rigon *et al.*, 2020). Between these two groups, NTSR, specifically metabolic resistance, is a critical issue due to the potential interaction of a single metabolic resistance mechanism with herbicides spanning multiple modes of action (Rigon *et al.*, 2020).

Plants possess a diverse array of enzymes involved in the detoxification of xenobiotics. Within these enzymes, cytochrome P450 enzymes (P450) play a key role in the initial step of the detoxification process (Rigon *et al.*, 2020). P450s associated with herbicide metabolism have been identified from various crops (Siminszky *et al.*, 1999, Pan *et al.*, 2006, Kato *et al.*, 2020, Brazier-Hicks *et al.*, 2022, Pan *et al.*, 2022) and implicated in evolved herbicide resistance in weeds. Notable examples from weeds include *CYP81A12*, *CYP81A21*, *CYP81A15*, and *CYP81A14* from *Echinochloa phyllopogon* (Dimaano *et al.*, 2020); *CYP81A69* and *CYP81A70* from *Cynodon dactylon* (Zheng *et al.*, 2022); *CYP81A10v7* from *Lolium rigidum* (Han *et al.*, 2021) and *CYP77B34* from *Descurainia sophia* (Shen *et al.*, 2022). Nearly all of the identified genes from weeds were from grass species (Rigon *et al.*, 2020).

*Amaranthus palmeri* (Palmer amaranth) is a highly invasive and troublesome weed that poses a significant threat to agricultural crops and ecosystems. Native to the southwestern United States, this annual dicot weed has become a pervasive problem across various regions due to its

aggressive growth, adaptability, and resistance to herbicides (Ward *et al.*, 2013, Roberts &

Florentine, 2022). Its dioecious nature, with separate male and female plants, allows for efficient

cross-pollination, leading to the exchange of genetic material and the emergence of novel traits,

including herbicide resistance (Ward *et al.*, 2013). Populations have evolved resistance to

herbicide spanning eight different modes of action, including synthetic auxins (Shyam *et al.*,

2022) and herbicide targeting acetolactate synthase (ALS) (Berger et al., 2016), 5-

enolpyruvylshikimate-3-phosphate synthase (EPSPS) (Gaines *et al.*, 2010), protoporphyrinogen

oxidase (PPO) (Montgomery et al., 2021), photosystem II (PSII) (Jhala et al., 2014), 4-

hydroxyphenylpyruvate dioxygenase (HPPD) (Küpper *et al.*, 2018), very long chain fatty acid

elongase (VLCFAE) (Hwang *et al.*, 2023), and tubulin biosynthesis (González-Torralva &

Norsworthy, 2021).

*Amaranthus palmeri* has evolved herbicide resistance through two main mechanisms, TSR

and NTSR. The most commonly observed cases of resistance involve TSR to ALS- and EPSPS-

inhibitors (Gaines *et al.*, 2010, Palmieri *et al.*, 2022). However, there are increasing resistance

cases due to metabolic-based mechanisms (Küpper *et al.*, 2018, Shyam *et al.*, 2022), and further

analysis is necessary to identify the specific genes involved and their regulation. RNA-seq

analysis plays a crucial role in deciphering the genetic basis of herbicide resistance, especially if

it is a multigenic trait (Gaines et al., 2014, Kohlhase et al., 2019). Additionally, with the

availability of complete genomes for weed species (Montgomery et al., 2023), such as *A.*

*palmeri*, conducting Quantitative Trait Locus (QTL) mapping experiments becomes a valuable

approach for determining the genetic architecture of resistance (Murphy *et al.*, 2021). By

combining comprehensive gene expression data obtained from RNA-seq analysis with QTL

mapping, researchers can gain valuable insights into the genetic factors underlying resistance and

other important characteristics in weeds.

This research aimed to identify and validate metabolic genes associated with the resistance

to tembotrione (an HPPD inhibitor) within a particular population of *A. palmeri*. We

subsequently sought to determine whether an identified gene, *CYP72A19*, was also associated

with resistance in other suspected tembotrione-resistant *A. palmeri* populations. The research

also sequenced the promoter and a run a QTL analysis to investigate regulation of *CYP72A219*.

**MATERIALS AND METHODS**

**Plant material for QTL mapping and RNA-seq analysis**

The resistant (NER) and susceptible (NES) *A. palmeri* populations were collected from

fields in Shickley, Nebraska in 2011. NER is resistant to atrazine and the HPPD inhibitors

tembotrione, mesotrione, and topramezone (Jhala *et al.*, 2014, Küpper *et al.*, 2018). The

enhanced metabolism of tembotrione to hydroxy-tembotrione was identified as the mechanism of

herbicide resistance in the NER population (Küpper *et al.*, 2018). To minimize the impact of

genetic differences unrelated to metabolic herbicide resistance, QTL mapping and RNA-seq was

performed on pseudo-F2 plants generated from controlled pairings of NER and NES parents. The

approaches for the generation of the pseudo-F2 plants are in Methods **S1** and Fig. **S1**.

**RNA-sequencing analysis**

Descriptions of plant population, herbicide application, plant tissue collection, and library

preparation are in Methods **S1**. Raw reads were pre-processed by removal of library adapter

sequences, removal of low quality reads, and assessing the quality control using fastp (Chen *et*

*al.*, 2018). Read alignment of the 108 libraries to the available male genome of *A. palmeri* (v1.1,

id55760) (Montgomery *et al.*, 2020) was performed using Hisat2.2.1 (Kim *et al.*, 2019). Most of

the reads (>77%) aligned concordantly once, while >7% aligned concordantly more than one

time and about 14% of reads aligned not concordantly. Mapped reads were assigned to gene

features using featureCounts (Liao *et al.*, 2014).

**P450 gene sequences and phylogenetic tree**

Consensus sequences of the candidate P450 genes were extracted from the transcriptome

and aligned to the reference *A. palmeri* genome (Montgomery *et al.*, 2020) to analyze single

nucleotide polymorphisms (SNPs). PCR was performed to amplify the coding sequences in four

each of sensitive (S) and resistant (R) plants from the pseudo-F2 population that were used for

the RNA-seq experiment. cDNA synthesis was performed using ProtoScript® II First Strand

cDNA synthesis kit with 1 µg of RNA. Protein sequences from cytochrome P450 known to

metabolize herbicides in different plant species were obtained from NCBI. Multiple protein

alignment was performed using ClustalO and used for tree construction with Neighbor-joining

method using Geneious Prime® 2023.0.1.

**Quantitative reverse transcription-PCR (RT-qPCR) validation**

NES and NER populations were grown, and herbicide applied at 91 g a.i. ha$^{-1}$ as

previously described. Four plants of each population were used for gene expression analysis

validation. Youngest leaf tissue was collected at 0, 3, 6, and 12 h after herbicide application

(HAT) in 2 mL Eppendorf tubes and placed in liquid nitrogen. The tubes were kept at -80 ℃ for

further analysis. Tissue was ground with 3 mm stainless steel beads in a TissueLyser (Qiagen)

with intensity of 30 for 1 min. RNA was isolated using Direct-zol RNA Miniprep from Zymo

Research. cDNA synthesis was performed using ProtoScript® II First Strand cDNA synthesis kit

using 1 µg of RNA and purified with DNase I.

Relative gene expression was analyzed on a T100 Thermal Cycler (BioRad), using

SsoAdvanced™ universal SYBR® Green supermix. Reaction mixtures consisted of 10 µL of

SsoAdvanced universal SYBR® Green supermix ($2\times$), 2.5 µL of forward and reverse primers at

10 µM, and 5 µL of cDNA (1:20 dilution). Thermocycler conditions consisted of an initial step

of 30 sec at 95°C followed by 35 cycles of 5s at 94°C and 30 s at 60°C. Melt curve analysis was

added using 65°C with 0.5°C increments of 5 s/step.

The reference genes used were *18S rRNA* and *Actin7*. These two genes had the best gene

stability as assessed by the NormFinder algorithm (Andersen et al., 2004) (Tab. **S1**). Primers for

the reference genes were designed based on conserved regions after alignment of *18S*

*(FJ669720.1), actin* (HQ656028.1), and *TUB* (XM_010693569.3) from *Beta vulgaris* with the

reference genome from *A. palmeri* (CoGe – v1.1, id55760). The candidate genes tested based on

RNA-seq results were *CYP72A219*_4284, *CYP72A219*_4285, *CYP72A219*_7285, and *CYP81E8*.

Another *CYP72A-like* gene, hereafter named as *CYP72A219*_4286 was used as a cytochrome

P450 not associated with herbicide resistance based on RNA-seq data. Primers for candidate

genes were designed on regions based on consensus sequences from the transcriptome and

Sanger sequencing of the genes and are listed in Tab. **S2**.

**Heterologous expression of CYPs**

The genes of interest were transformed and expressed in yeast for further investigation of

their role in tembotrione metabolism. Coding sequences of the candidate genes were synthesized,

optimized for yeast transformation, and inserted in the pUC-WG/ampv vector by GENEWIZ

(Azenta Life Sciences). The gene sequences used were *CYP72A219*_4284 allele 1 (GenBank

accession number: OR596705) and 2 (OR596706), *CYP72A219_4285* (OR596707),

*CYP72A219_7285* allele 1 (OR596708) and 2 (OR596709), and *CYP81E8* allele 1 (OR596710)

and 2 (OR596711). As a positive control, the gene *CYP81A9* from corn, known as *Nsf1*, was

used. This gene was identified as the major resistance locus in corn for the herbicides

nicosulfuron and mesotrione (Williams & Pataky, 2008). Restriction sites for BamHI and EcoRI

were added the the 5' and 3' ends, respectively. Kozak sequence (AAAAAATCT) was added at

the 5' end as a protein translation initiation site. Restriction enzyme reactions were performed to

cut the optimized sequence from the vector using 1 μL EcoRI, 1 μL BamHI, 500 ng of the

vector, with incubation at 37°C for 2 h. Reaction products were run on 1% agarose gel

electrophoresis and the gene band was isolated and purified using a DNA Gel Purification Kit

from New England Biolabs®.

The pYES2 yeast expression vector was used for recombinant expression. It contains

the *URA3* gene for selection in yeast and 2μ origin for high-copy maintenance and GAL1

promoter to express protein (Thermo Fisher Scientific). The ligation reaction was performed

using LigaFast™ Rapid DNA Ligation System from Promega. The reaction consisted of 50 ng

digested pYES2 plasmid vector, 5 μL 2× Rapid Ligation Buffer, 1 μL T4 DNA ligase (3 Weiss

unit/μL), 50 ng of gene, and purified water up to 10 μL. The reaction was incubated overnight at

4°C. The product from the reaction was used to transform *E. coli* using the One Shot® TOP 10

kit (Invitrogen). The transformed cells were plated with ampicillin (100 μg/mL) and incubated at

37°C overnight. Single colonies were selected, and gene insertion was confirmed by colony PCR

using high-fidelity PrimerStar HS DNA polymerase. The gene sequence was confirmed by

Sanger sequencing. Plasmids were isolated using ZymoPure II Plasmid Miniprep Kit.

**Yeast transformation and tembotrione incubation**

Yeast transformation protocol is described in Methods **S2**. Single colonies containing

empty pYES2 or the candidate CYP genes were incubated in 15 mL 2% raffinose medium and

allowed to grow for 2-3 d at 30 °C. Cell density was measured spectrophotometrically until the

OD600 in 20 mL of induction medium was 5. The volume was removed and pelleted at 1,500 $\times$

g for 15 min at 4 °C. The cells were resuspended with 1 mL of induction medium containing 2%

galactose and inoculated into 20 mL of the same medium. Tembotrione at 1500 μM diluted in

EtOH was applied right after the cells were incubated in the induction medium. The yeast cells

were incubated at 30°C at 200 rpm. Twenty-four hours after herbicide application, 5 mL of the

medium were collected in 15 mL falcon tubes and cells were pelleted at 1,500 $\times$ g for 5 min.

The supernatant was collected, and cleaned by adding 5 mL of 5% acetic acid acetonitrile +

QuEChERS, vortexed for 30 s, and centrifuged for 15 min at 2,000 $\times$ g. The supernatant was

collected, filtered through a nylon 13 mm by 0.2 μm (Econofltr Nyln, Agilent Technologies),

and injected in the LC-MS/MS.

**LC-MS/MS protocol**

LC-MS/MS system consisted of a Nexera X2 UPLC with 2 LC-30AD pumps, a SIL-30AC

MP autosampler, a DGU-20A5 Prominence degasser, a CTO-30A column oven, and SPD-M30A

diode array detector coupled to an 8040 quadrupole mass-spectrometer. For tembotrione, the MS

was in negative mode with an MRM optimized for 439.1>226.05 and set for 100 ms dwell time

with a Q1 pre-bias of 11.0V, a collision energy of 11.0V and a Q3 pre-bias of 14.0V. For

hydroxy-tembotrione, the MS was in negative mode with an MRM optimized for 455.1>419.05

and set for 100 ms dwell time with a Q1 pre-bias of 11.0V, a collision energy of 11.0V and a Q3

pre-bias of 14.0V. The samples were chromatographed on a 100$\times$4.6 mm Phenomenex Kinetex

2.6 μm biphenyl column maintained at 40°C. Solvent A consisted of water with 0.1% formic

acid and solvent B was acetonitrile with 0.1% formic acid. The solvent program started at 80% B

and increased to 100% B in 3.5 min and maintained at 100% for 2 min. The solvent was returned

to 80% B and maintained there for 3 min before the next injection. The flow rate was set at 0.4

mL/min and samples were analyzed as 1 μL injection volumes.

*CYP72A219_4284* and *CYP81E8* **promoter amplification**

Pseudo-F2 plants were grown in the greenhouse and the youngest leaf tissue was collected

when the plants reached four to five true leaf stage. Tembotrione was applied at 77 g ha$^{-1}$ rate.

Five each of R and S plants from a pseudo-F2 population were chosen for promoter

amplification. DNA extraction was performed using hexadecyltrimethylammonium bromide

(CTAB) method (Doyle & Doyle, 1990). DNA quantification was carried out using Nanodrop

2000c (Thermo Fisher Scienfitic). Primers were designed by Primer3Plus

(https://www.bioinformatics.nl/) using as reference the available draft genome of *A. palmeri*

(Montgomery *et al.*, 2020). The primer sequences used are listed in Tab. **S2**. The methods used

to amplify the gene promoter are available in Methos **S3**.

**Involvement of *CYP72A219_4284* in different tembotrione-resistant *A. palmeri***

**populations**

Seeds from *A. palmeri* populations were collected from agricultural fields in 2019 in the

United States with suspected herbicide resistance to mesotrione and tembotrione. Whole-plant

dose response, $^{14}$C-tembotrione metabolism, P450 gene expression and gene copy number were

evaluated with the hypothesis that the same candidate P450 gene is involved in the resistance

mechanism in these additional populations. The methods are described in Methods **S4**.

**Mapping of tembotrione resistance in NER *A. palmeri* population**

*Library preparation and QTL identification*

Dose-response and segregation analysis methods are described in Methods **S5**. Plant tissue

was collected before herbicide application for DNA extraction from single leaves following a

modified CTAB method. DNA samples were assessed for quality using Qubit. Double-digest

restriction site-associated DNA sequencing (ddRADseq) libraries were generated with ApeKI

and sequenced using NovaSeq S4 with 150 bp paired-end reads (Illumina). The average yield

was around 7.4 M reads per library and the mean quality scores were over Q30 for all libraries.

The sequencing was performed at the University of Minnesota Genomics Center.

The ddRAD-seq libraries were trimmed using trimmomatic v0.36 (Bolger et al., 2014),

aligned to the reference male genome of *A. palmeri* (scaffold file - v1.1, id55760) (Montgomery

*et al.*, 2020) with Burrows-Wheeler Aligner (Li & Durbin, 2009) and variants called using

GATK 4.2.0 (Poplin *et al.*, 2018). Samples were computationally binned by shoot fresh mass.

The variant sites were separated by SNPs and insertion/indels and hard filter was performed as

following for SNPs - $QD < 2.0$, $QUAL < 30.0$, $SOR > 4.0$, $FS > 20.0$, $MQ < 50.0$ and

insertion/indels $QD < 2.0$, $QUAL < 30.0$, $FS > 200.0$. The variant sites were filtered based on

depth (at least five reads) and only variants that were homozygous were selected. R/qtl2 package

was used to find QTLs (Broman et al., 2019). The analysis was performed separately for two

separate crosses (A and B) and combined A + B. QTL intervals were calculated using Bayes

credible intervals with the function bayes_int(). To determine the threshold for identifying

potential QTL, 1000 permutation tests were conducted at a confidence level of 95%. The critical

F value obtained from this analysis was used as the criterion to declare putative QTL. The

functional analysis of the genes present in QTLs was conducted using DAVID Bioinformatics

Resources v6.8 (Sherman et al., 2022).

Statistical analysis for all the experiments is described in Methods **S6**.

**RESULTS**

**Differentially expressed genes (DEGs)**

A total of 20,846 genes were analyzed for differential expression out of the initial set of

29,758 genes assigned by featureCounts. Gene-wise dispersion curve was fitted using the

DESeq2 model (Fig. **S2**). There was favorable dispersion pattern, with decreasing dispersion as

the mean expression levels increased, indicating a good fit of the DESeq2 model to the analyzed

data. The principal component analysis (PCA) plot revealed that PCA1 and PCA2 accounted for

35 and 21% of the variation in the data, respectively (Fig. **S3**). PCA1 separated cross A from

cross B, while PCA2 exhibited separation of gene clusters based on HAT within each cross.

When clustering all 72 analyzed transcriptome samples, two major clusters emerged, with

samples from different crosses grouped together, indicating similarities in gene expression

response (Fig. **S4**).

DEGs were observed in different comparisons. In the contrast comparison of R versus S at

constitutive (0 HAT), 6 HAT, and 12 HAT after herbicide application, R had 37, 2, and 9 up-

regulated genes and 3, 2, and 4 down-regulated genes, respectively (Fig. **1a**). Among these

genes, only two genes were commonly up-regulated (Fig. **1a**). For the analysis of the genes that

were responsive to herbicide application for each phenotype, 39, 530, and 106 genes were up-

regulated in S and 6, 185, and 160 were up-regulated in R for the comparisons of time 6 vs 0

HAT, 12 vs 0 HAT, and 12 vs 6 HAT, respectively (Fig. **1a**). The heatmap illustrates the

expression of DEGs from these comparisons and indicates a separation in clusters of resistant

and susceptible plants (Fig. **1a**).

Several genes associated with the detoxification of xenobiotic substances were consistently

identified, suggesting an enhanced genetic metabolism in plants with resistance. These genes

included four cytochrome P450 genes, four glutathione-S-transferase genes, seven

glycosyltransferase genes, a disease resistance protein, and a detoxification 27 gene (Tab. **S3**).

Additionally, two MADS-box transcription factors of types 23 and 27 were found to be up-

regulated and arise hypothesis of gene regulation in resistant plants. The resistance mechanism in

the resistant population was characterized by increased herbicide metabolism including an initial

hydroxylation of the tembotrione molecule (Küpper *et al.*, 2018); therefore, among all

differentially expressed genes (DEGs), our focus was on the four cytochrome P450 genes that

were consistently up-regulated in the R plants. These genes had the genomic IDs

MAKER_29886, MAKER_25717, MAKER_10107, and MAKER_25718, and exhibited fold

changes of 18.3, 11.8, 6.2, and 5.4, respectively (Fig**. 2b-d** and Tab. **S3**). None of these genes

had differential expression at the 6 HAT time point compared to the susceptible plants (Fig. **2c**),

and only one gene (MAKER_29886) exhibited a reactivation at the 12 HAT (Fig. **2c**). This

indicates that the susceptible plants responded to the herbicide treatment (Fig. **2c**), where an

increased transcription of these genes was observed at 6 and 12 HAT. However, the resistant

plants constitutively expressed these cytochrome P450 genes at a higher level in the absence of

herbicide treatment.

Constitutively expressed P450s, namely MAKER_25717, MAKER_25718, and

MAKER_29886, were designated *CYP72A219* due to their significant similarity to *CYP72A219*

from *Spinacia oleracea* (NCBI: XM_056843192.1), with identities of 76.2, 74.3, and 76.3%

respectively. Similarly, the gene ID MAKER_10107 was assigned the name *CYP81E8* due to its

high similarity (73.3%) to *CYP81E8* from *Chenopodium quinoa* (NCBI: XP_021724107.1).

Hereafter, the genes MAKER_25717, MAKER_25718, MAKER_29886, and MAKER_10107

will be referred to as *CYP72A219_4284*, *CYP72A219_4285*, *CYP72A219_7285*, and *CYP81E8*

respectively. The number after CYP gene name indicates the gene number ID in the reference

genome of *A. palmeri*.

**SNPs in cytochrome P450 candidate**

There were no consistent SNPs or amino acid changes between S or R in the P450 genes.

*CYP72A219_4284* has a coding sequence of 1544 bp and 517 amino acids. CYP72A219_4285

has a coding sequence of 1578 bp and 526 amino acids. Both phenotypes share the same two

alleles for these genes. *CYP72A219_7285* has a length of 1394 bp and 464 amino acids.

Susceptible plants have single alleles and R plants have a second allele with amino acid

substitutions of Thr307Ser, Phe388Leu, and Ile407Val. The former two *CYP72A219* (4284 and

4285) are localized on chromosome four and the latter is localized in chromosome 16.

*CYP72A219_4284* shares identity with *CYP72A219_4285* and *CYP72A219_7285* of 76.1 and

55.1%, respectively. Two alleles of *CYP81E8* were present in both S and R samples. The gene

has a length of 1470 bp and 1479 bp in each allele. The short allele has a deletion of 18 bp in

position 35, and the longest allele has a deletion of 9 bp in position 37 when aligned to the draft

genome. In the available *A. palmeri* genome, this specific region of the gene exhibits a

triplication of PPS amino acids, which is likely due to incorrect assembly. All cytochrome P450

possess a conserved cluster of proline in the membrane hinge. Similarly, the sequenced

cytochrome P450 exhibits this conserved region.

In the phylogenetic tree analysis, *CYP72A219* genes clustered with *CYP72A31* from *Oryza*

*sativa*, which can metabolize bispyribac-sodium and bensulfuron-methyl (Saika et al., 2014).

*CYP81E8* was clustered with other P450 of family 81 from grasses but with some significant

distances between them (Fig. **S5**). This gene has a low identity when aligned with the other

P450s. *ApCYP81E8* shares identity of 40.5%, 41.0%, and 39% with *ZmCYP81A9* (Genbank:

EU955910.1), *LrCYP81A10v7* (Genbank: MK629521.1), and *EpCYP81A12*

(Genbank:AB818460.1), respectively, known to metabolize herbicides in grasses (Brazier *et al.*,

2002, Iwakami *et al.*, 2014, Han *et al.*, 2021, Brazier-Hicks *et al.*, 2022).

**P450 qPCR validation**

*CYP72A219_4284*, *CYP72A219_4285*, and *CYP72A219_7285* were upregulated in NER

plants at 3, 6, and 12 HAT, respectively (Fig. **S6**). Among the *CYP72A219* genes,

*CYP72A219_4284* had a higher up-regulation at 3 and 6 HAT. Additionally, *CYP81E8*, located

on chromosome 4, had up-regulation specifically at the 6 HAT (Fig. **S6**). These observations

deviate from those obtained in the RNA-seq analysis, which indicated that these genes are

constitutively expressed in resistant plants (Fig. **2**). However, this discrepancy arises due to the

utilization of the parental populations NES and NER for gene validation, which exhibit

significant variation in their response to tembotrione. Nevertheless, despite the disparity, the

consistency of these findings indicates that the up-regulation of these genes in NER plants differs

from that in NES plants.

*CYP72A219* **metabolizes tembotrione**

Following a 24 h incubation period with tembotrione, some transformed yeast treatments

converted tembotrione into hydroxy-tembotrione. The chromatogram of the empty vector pYES2

revealed solely the presence of the parent tembotrione peak at a retention time of 2.9 min (Fig.

**3**). Notably, the *Nsf1* gene exhibited a pronounced affinity for tembotrione, producing hydroxy-

tembotrione peak at a retention time of 2.7 min and a dihydroxy tembotrione peak at 2.5 min

(Fig. **3**). Among all the candidate P450 genes tested, only *CYP72A219_4284* converted the

parent tembotrione into hydroxy-tembotrione. Both alleles of this gene in either the WAT11 or

WAT21 strain metabolized the herbicide. Conversely, *CYP72A219_4285*, *CYP72A219_7285*,

and *CYP81E8* did not exhibit any ability to metabolize tembotrione. Based on these results, it

can be concluded that only *CYP72A219_4284* possesses the capacity to recognize the

tembotrione molecule and effectively metabolize it.

**CYPs promoter analysis**

The DNA segment amplified upstream *CYP81E8* was 1,250 base pairs for both the

sensitive and resistant plants (Fig. **4**). Pairwise sequence alignment between two samples

revealed 100% identity between the promoters of S and R (Fig. **4a** and Notes **S1**). For the

promoter of the *CYP72A219_4284* gene on chromosome four (MAKER-25717), the primers

amplified regions ranging from 1991 to 2004 base pairs for S plants and from 1747 to 1768 base

pairs for R plants. Pairwise alignment of *CYP72A219_4284* promoter revealed sequence

differences between S and R, with identity of 76.8% (Notes **S1**). Resistant plants had unique

insertions, such as a cytosine at position 45 bp, an 8 bp insertion at position 458 bp, an 8 bp

insertion at position 1247 bp, and a 31 bp insertion at position 1291 bp upstream. Sensitive

plants, on the other hand, had unique insertions, including a 13 bp insertion at position 82, a 10

bp insertion at position 309 bp, and a 277 bp insertion at position 746 bp upstream of the gene

(Fig. **4b** and Notes **S1**).

A total of fifteen significant binding-site motifs were identified in the promoter region of

*CYP81E8*. Interestingly, all motifs were found in both S and R promoters (Fig. **S7**). In the case

of the *CYP72A219_4284* promoter, a total of twenty-five motifs were discovered. Among these,

two motifs were specific to the S promoter, two were specific to the R promoter, and twenty-one

motifs were shared between S and R (Fig. **S7** and **S8**). Notably, the R promoter exhibited

replication of four motifs. Some motifs were identified as a binding site for transcription factors

such as MYB88 (AT2G02820), suppressor factor AIF1 (AT3G05800), and abiotic stress-

responsive transcription factor DREB (AT1G77200) based on GOMo analysis (Fig. **S8**).

Furthermore, the motif GTTATTTAGTAACTTWKYGTD, which serves as a binding site for

the MYB gene transcription factor, was triplicated in the R promoter (Fig. **S8**).

**Whole-plant dose response, metabolism, and *CYP* expression in different herbicide**

**resistant *A. palmeri***

Based on the whole-plant dose response (Fig. **S9**), the confirmation of *A. palmeri*

populations with suspected tembotrione resistance revealed distinct variations, with resistance

index ranging from 2.4 to 7.9 (Tab. **S4** and **S5**). Tembotrione metabolism was assessed in the *A.*

*palmeri* populations using $^{14}$C herbicide (Fig. **S10**). The method used in the present study was

different from the previous study (Küpper *et al.*, 2018), hence, retention times are slightly

different. Parental tembotrione had a retention time of 66.9 min and its major metabolites had

retention times of 47.3 (M1), 50.1 (M2), 56.3 (M3), 59.0 (M4), and 67.7 min (M5) (Fig. S11 and

S12). NER had higher % area peak of hydroxy-tembotrione (M3/M4), the main metabolite

associated with herbicide resistance (Küpper *et al.*, 2018), than the sensitive population at 12

HAT (Fig **S10**). The other resistant populations were similar to NER, except for WR2019-044

and WR2019-273 (Fig. **S10**). The faster hydroxy-tembotrione production in most populations

suggests that enhanced metabolism capacity is the main mechanism of HPPD-resistance of these

populations. *CYP72A219_4284* was upregulated in most resistant populations, either before or 6

HAT (Fig. **S1**). The populations WR2019-034, -137, -140, -199, and -274 had a significantly

higher relative expression compared to the sensitive population, ranging from 100 to 657.8-fold

increase. In contrast, the expression of *CYP81E8* was not constitutively higher or responsive to

tembotrione treatment in *A. palmeri* populations, except in WR2019-140 (Fig. **S1**). The gene

copy number of *CYP72A219_4284* was similar across all resistant populations and with that of

sensitive plants (Fig. **S14**). The higher gene expression of *CYP72A219_4284*, constitutively and

after tembotrione application in resistant plants, suggests its crucial role in herbicide resistance in

the newly characterized tembotrione-resistant populations.

**Degree of dominance and QTL identification**

The $ED_{50}$ of NES and NER were 7.9 and 37.3 g, respectively (Tab. **1**), lower than

previously reported for these populations (Küpper *et al.*, 2018). However, the resistance index

(RI) between these two populations was similar, 4 compared with 3.3 from the previous study

(Küpper *et al.*, 2018). Both F1 populations were intermediate between NES and NER in response

to tembotrione (Fig. **5**). The degree of dominance was calculated and indicated a co-dominant or

semi-dominant trait (cross A = 0.42 and cross B = 0.67) based on the formula proposed by

Bourguet and Raymond (1998). Pseudo-F2 populations, resulting from the cross between A and

B, exhibited survival rates of 11.4% and 8.8%, respectively. These survival rates are similar to

the parental homozygous line NER (12.4%), which was cultivated alongside the pseudo-F2

plants.

Frequency distribution of fresh weight after herbicide treatment on pseudo-F2 plants had

two distinct peaks for each cross, with one peak each indicating phenotypic similarity with each

parental population. A normal distribution of R plants had higher biomass (red bars) and the

same was identified for S samples (Fig. **6a**). The variant calling analysis identified over 3.4

million SNP sites in the *A. palmeri* genome. After filtering, a total of 4,404 SNP variants were

detected throughout the entire *A. palmeri* genome. Genome scanning in cross A did not detect

any QTLs surpassing the LOD threshold of 6.4 after conducting a permutation test. However,

when examining pseudo-F2 plants from cross B, significant QTLs with high effects were found

in multiple scaffolds overpassing LOD threshold of 6.7. The scaffolds include scaffold 10 with

LOD of 11.6, scaffold 81 with LOD of 10.1, scaffold 6 with LOD of 9.5, and scaffold 14 with

LOD of 8. Each QTL was accompanied by other significant QTLs nearby (Fig. **6b**, Tab. **S6**). For

instance, in scaffold 10, there are three QTL peaks, the biggest peak with a LOD score of 11.6 at

position 11,877,263, and two others peaks with LOD scores of 8.4 and 7.9 at positions

10,336,852 and 10,337,656, respectively. The same happened in scaffolds 81, 6, and 14 with

nearby peaks. Combining the datasets from cross A and cross B increased the sample size and

enhanced the statistical power for the analysis. As a result, the genome scan revealed a more

robust QTL effect on scaffold 10, precisely at the same genomic location 11,877,263. The QTL

exhibited a higher LOD score of 12.2, indicating a stronger and more significant association with

the resistance trait (Tab. **S6**). Additional QTLs were found on scaffolds 6, and 14, but with lower

effects.

QTL located in scaffold 10 accounted for 23% of the phenotypic variation observed in the

data for cross B. Additionally, the QTLs in scaffold 81, 6, and 14 explained 20, 19, and 16% of

the phenotypic variation, respectively. However, when the data from both cross A and cross B

were combined, the overall phenotypic variation explained by the QTLs slightly decreased. This

can be attributed to the absence of any QTL in cross A. Specifically, for the combined data, the

QTL in scaffold 10 explained 15% of the phenotypic variation, while the QTLs in scaffold 6 and

14 accounted for 10% each (Tab. **S6**).

To conduct further analysis, functional annotation was performed on the combined dataset

due to its larger number of genes and sample size. The QTLs found in scaffolds 10, 6, and 14 are

localized on chromosomes four, eight, and fifteen of *A. palmeri*, respectively, and encompass 82,

196, and 25 genes, respectively (Supplemental Information Tab. **S6**). The QTL on scaffold

10/chromosome 4 does not contain the *CYP72A219_4284* gene but is located 3 Mb away.

Functional annotation of these genes indicates a predominant molecular function of protein

binding (GO:0005515), RNA binding (GO:0003723), and mRNA binding (GO:0003729),

consistent with proteins involved in regulating other proteins or molecules through selective

protein interactions (Hudson & Ortlund, 2014) (Fig. **S15**). The important TFs that were identified

include *DREB2A*, *WRKY* and GATA16 within the QTLs. The QTLs also include nine zinc finger

proteins, four ubiquitin proteases, three F-box proteins, three ABC transporters, and three

cytochrome P450 enzymes.

For each QTL, multiple markers were identified. In the case of the QTL in scaffold 10, a

total of 54 markers were found, while for the QTL in scaffold 6, 94 markers were identified.

However, for the QTL in scaffold 14, only a single marker was detected within a 1 Mb range

upstream and downstream of the QTL region. The markers associated with scaffolds 10 and 6

displayed a consistent pattern with the evaluated phenotype, where the RR genotype exhibited

higher biomasses, while the SS genotype had lower biomasses (Fig**. 7**). In contrast, the single

marker found in scaffold 14 did not demonstrate the same consistency as the markers identified

in the other QTLs (Fig**. 7i**).

**DISCUSSION**

The evolution of metabolic herbicide resistance mechanisms in weeds represents a

significant challenge in weed control practices (Rigon *et al.*, 2020). These mechanisms enable

weeds to metabolize a diverse array of herbicides from various chemical families. Identifying the

genes responsible for metabolic resistance and comprehending their regulation holds immense

potential for developing improved herbicides and mitigating herbicide resistance. In our current

investigation, four consistently up-regulated P450 genes were identified in HPPD-inhibitor

resistant *A. palmeri*. These four genes did not have any polymorphisms in the coding sequences

between R and S plants. Among these genes, only one, named *CYP72A219_4284*, deactivated

the herbicide tembotrione when tested in a heterologous system. Furthermore, our study revealed

that this specific gene is upregulated in other HPPD-inhibitor resistant *A. palmeri* populations

collected from various fields across the US, supporting its involvement in conferring herbicide

resistance in this globally significant species.

In the past, metabolic herbicide resistance in dicot weeds was relatively uncommon, with

most documented cases of metabolic resistance and identified P450 genes occurring in

monocots. However, recent studies have reported instances of metabolic resistance in dicots such

as *A. tuberculatus* in 2017 (Kaundun et al., 2017), *A. palmeri* (Küpper *et al.*, 2018) in 2018, and

*Raphanus raphanistrum* in 2020 (Lu *et al.*, 2019). These weed species exhibited resistance to

HPPD inhibitors. Furthermore, in 2020, metabolic resistance to 2,4-D was identified in *A.*

*tuberculatus* (de Figueiredo *et al.*, 2022) . The identification of P450 genes capable of

metabolizing herbicides in dicots is relatively limited. For example, *DsCYP77B34* from

*Descurainia sophia*, a broadleaf weed found in Asian wheat fields, conferred resistance to ALS-,

PPO-, VLCFA-, and PSII-inhibitors when expressed in *Arabidopsis thaliana* (Shen *et al.*, 2022).

Arabidopsis lines expressing *R. raphanistrum* genes *RrCYP704C1* or *RrCYP709B1* were

resistant to HPPD-inhibitors such as mesotrione *(RrCYP704C1)*, tembotrione *(RrCYP709B1)*,

and isoxaflutole *(RrCYP709B1)* (Lu *et al.*, 2023). In *A. palmeri*, *CYP81E8* was reported as the

primary gene responsible for metabolic resistance in a HPPD-inhibitor resistant population from

Kansas, USA (KCTR) (Aarthy *et al.*, 2022). This study found that the population exhibited

enhanced metabolism of tembotrione, and *CYP81E8* expression, measured by qPCR, was

implicated as the likely cause of tembotrione resistance. However, our study indicates that

although *CYP81E8* is up-regulated, it is not the causal gene for tembotrione resistance in the

NER population. Instead, we have identified *CYP72A219_4284* as the gene capable of

converting tembotrione to hydroxy-tembotrione. This finding represents the first study which

mechanistically validated the gene responsible for herbicide metabolism in *A. palmeri*.

*CYP81E8* has been hypothesized as the gene responsible for 2,4-D metabolic resistance in

*A. tuberculatus* (Giacomini *et al.*, 2020). This specific cytochrome P450 gene co-segregated with

2,4-D resistance based on RNA-seq analysis conducted on F2 individuals. However, the gene did

not co-segregate with HPPD resistance, indicating that the genetic association was specific to

2,4-D resistance. Mechanistic gene validation for function in 2,4-D metabolism was not

performed (Giacomini *et al.*, 2020). Interestingly, *CYP81E8* alleles from *A. tuberculatus* and *A.*

*palmeri* exhibit high sequence similarity, greater than 92%. A similar behavior of *CYP81E8* was

observed in this study, with higher expression levels in resistant plants of *A. palmeri*; however,

when tested in a heterologous system, this gene did not inactivate tembotrione. Based on this

finding, we hypothesized that the NER population might have resistance to 2,4-D due to up-

regulation of *CYP81E8*. The NER population had a resistance index ranging from 2 to 2.5, but

complete control was achieved when 500 a.e. g ha$^{-1}$ of 2,4-D was applied (data not shown). This

suggests that the NER population may indeed be evolving resistance to 2,4-D; however,

resistance to tembotrione is not attributed to *CYP81E8*, but rather to *CYP72A219_4284*.

Previous research has highlighted *CYP72A219* as a gene involved in herbicide resistance in

weeds. A study indicated that among other metabolic genes, *CYP72A219* had constitutive up-

regulation in glufosinate-resistant *A. palmeri* plants, with eight times higher expression levels in

resistant plants; however, the study did not perform functional gene validation (Salas-Perez *et*

*al.*, 2018). Furthermore, *BsCYP72A219*, along with *BsCYP81Q32* from *Beckmannia syzigachne*,

were up-regulated in mesosulfuron-methyl-resistant plants (Wang *et al.*, 2023). These genes

were validated in transgenic rice, where only *CYP81Q32* conferred resistance to ALS-inhibitors.

Notably, up-regulation of two genes with only one gene having validated metabolic function for

herbicide resistance, resembled the observed pattern in the present study. Additionally, an

alignment of *CYP81E8* from *A. palmeri* revealed a 92.7% identity with *CYP81Q32-like* from

*Amaranthus tricolor* (XP_057543510.1), a recently sequenced *Amaranthus* species (Wang *et al.*,

2022). This finding indicates a very close relationship between *CYP81E8* and *CYP81Q32* and

may even represent the same gene, designated with different names. The pattern of co-

segregation and up-regulation of these genes in different weed species suggests the presence of

common *cis* or *trans*-elements that potentially regulate expression of multiple P450 genes in

resistant plants.

Distinct promoter sequences for *CYP72A219_4284* were found in R pseudo-F2 plants

compared to S, while no such variations were observed for *CYP81E8*. QTLs with high effects on

the herbicide resistance trait, localized in chromosomes four, eight, and fifteen were also

identified. These QTLs include three cytochrome P450 genes and ABC transporters. While these

are important genes and warrant further analysis, a gene responsible for metabolizing

tembotrione was identified through RNA-seq analysis and validated, suggesting that the new

P450s in the QTL mapping might have a small effect or even not be crucial for the resistance

mechanism of the NER population. The key findings from the QTL are the diverse set of

regulatory genes, including key transcription factors that could play a crucial role in the up-

regulation of *CYP72A219* in resistant plants. Transcription factors such as *DREB2A*, *WRKY,* and

*GATA16*, along with zinc finger family found in the mapping analysis are known for their

involvement in abiotic stress responses (Banerjee & Roychoudhury, 2015, Zhang *et al.*, 2015,

Seale, 2020). Significantly, it should be highlighted that the identified QTL on chromosome four

is located two to three Mbp upstream of the *CYP72A219_4284* gene. This proximity suggests

that the transcription factor (TF) within the QTL region may regulate the gene, potentially

influencing its expression. This regulatory effect could be particularly relevant due to the

presence of a distinct promoter sequence observed in resistant plants.

Promoters of P450s have been investigated in other weed species. Nucleotide

polymorphisms were observed in the promoters of *CYP81A12* and *CYP81A21* in the R alleles of

multiple herbicide resistant *Echinochloa phyllopogon* plants; however, these single nucleotide

polymorphisms did not exhibit a significant association with herbicide resistance (Iwakami *et al.*,

2014). Three SNPs were identified in the promoter region of *BsCYP81Q32* between the R and S

variants, spanning approximately 567 to 1207 base pairs upstream of the gene. These SNPs were

responsible for increased expression of *GUS* in tobacco leaves transformed with the gene,

suggesting the presence of a crucial sequence within that specific region (Wang *et al.*, 2023).

Additionally, the researchers identified a transcription factor, *BsTGAL6*, belonging to the bZIP

family, which demonstrated efficient binding to the promoter of *BsCYP81Q32* (Wang *et al.*,

2023). This finding suggests the involvement of a *trans*-element that regulates the expression of

*BsCYP81Q32* in resistant plants.

Our study successfully identified and validated the metabolic resistant gene,

*CYP72A219_4284*, in the tembotrione-resistant *Amaranthus palmeri* population NER. This gene

demonstrated the ability to metabolize tembotrione into the main metabolite hydroxy-

tembotrione. We have evidence that this gene is highly upregulated in various *A. palmeri*

populations across the United States, which poses a potential threat to weed management

strategies targeting this species. We also discovered important QTLs associated with the

resistance trait, warranting further investigation and validation for the CYP72A219 regulation.

Future research focusing on validating the identified transcription factors will shed light on the

regulation of P450 genes and enhance our understanding of their regulation in weed species.

**ACKNOWLEDGEMENTS**

This work was supported by funding from the Weed Resistance Competence Center at Bayer AG and USDA National Institute of Food and Agriculture, Hatch projects COL00783 and COL00785 to the Colorado State University Agricultural Experiment Station.

**AUTHOR CONTRIBUTIONS**

CAGR, AK, AP-J, PJT, RB, FED, and TAG designed the research. CAGR, AK, CS, FP, and SS performed experiments. AK, AP-J, PJT and RB contributed experimental materials. CAGR, FP, SS, and JM analyzed the data. CAGR, FED, and TAG wrote the paper. All authors contributed to editing and approving the paper.

**DATA AVAILABILITY**

The transcriptome and genome sequencing data that support the findings of this study are available at https://www.ncbi.nlm.nih.gov, with reference numbers GSE243128 and PRJNA1012341, respectively. The gene sequences are available at Genbank accession numbers OR596705- OR596711.

581 **Tab. 1**. Logistic equation parameters and resistance index (RI) for survival (% of untreated
582 control) in parental *Amaranthus palmeri* herbicide-susceptible (NES), herbicide-resistant (NER),
583 F1 cross A, and F1 cross B populations in response to tembotrione.

| Population | b | ED$_{50}$ Dose (g ha$^{-1}$) | Lower CI (95%) | Upper CI (95%) | SE | RI | p-value |
|---|---|---|---|---|---|---|---|
| NES | 2.4 | 9.3 | 7.9 | 10.6 | 0.7 | - | |
| NER | 2.1 | 37.3 | 31.2 | 43.4 | 3.1 | 4 | <0.001 |
| F1 cross A | 2.4 | 16.6 | 14.2 | 19.1 | 1.2 | 1.8 | 0.2 |
| F1 cross B | 2.6 | 23.6 | 20.15 | 27 | 1.7 | 2.5 | <0.001 |

584 b: slope; ED$_{50}$: herbicide dose that causes a 50% reduction in plant survival; CI: confidence
585 interval of parameter ED$_{50}$ ($\alpha = 0.05$); RI: resistance index = ED$_{50}$ ratio between populations
586 with NES.
587

589
**Fig. 2**. Analysis of differentially expressed genes (DEGs) in Pseudo-F2 *Amaranthus palmeri*
susceptible (S) and HPPD-resistant (R). (a) The Venn diagram of the up- and down-regulated

DEGs indicating unique and common DEGs for 18 different comparisons. (b) Heatmap of top
DEGs for the comparisons R vs S at 0, 6 and 12 h after herbicide treatment (HAT). (c) Volcano
plots displaying gene expression differences between *Amaranthus palmeri* and treatments of
contrast of R versus S before herbicide application (0 HAT), contrast of R versus S at 6 HAT,
and contrast of R versus S at 12 HAT. Differentially expressed genes with significant thresholds
of p-value <0.0001, adjusted p-value <0.05, log2 fold-change >1 or < -1 are represented by red
circles. Dashed lines represent significance thresholds of adjusted p-value<0.05 and log2 fold-
change >1 or < -1. (d) The log2 normalized counts of top four cytochrome P450 genes in
response to tembotrione in S and R population, MAKER_25717 (CYP72A219_4284),
MAKER_25718 (CYP72A219_4285), MAKER_29886 (CYP72A219_7285), and
MAKER_10107 (CYP81E8). Differently expressed genes had a threshold of adjusted p-value of
0.05 and log2 fold-change 1.

621
622 **Fig. 3**. LC-MS/MS chromatogram of metabolites formed in yeast transformed with P450 genes
623 from *A. palmeri*. Blue peak – parental tembotrione, read peak – hydroxy-tembotrione, green
624 peak, dihydroxy-tembotrione. (a) WAT11 and (b) WAT21 yeast strain carrying *Arabidopsis*
625 *thaliana* cytochrome P450 reductase 1 and 2, respectively. * at least 10-fold higher peak than the
626 background signal.
627

**Fig. 4**. MUSCLE v5 multiple alignment of *CYP81E8* (a) and *CYP72A219_4284* (b) promoter sequences. The promoter sequences were reverse-oriented from the reference genome sequence to enable viewing in + strand orientation. Black space indicates differences from the reference *A. palmeri* genome (id 55760).

654
655 **Fig. 5**. Dose-response of *Amaranthus palmeri* herbicide susceptible (NES), resistant (NER), F1
656 cross A and F1 cross B plants in response to tembotrione.

**Fig. 6**. Frequency distribution and genome scan analysis. (a) Distribution of fresh biomass in

plants used for QTL mapping of pseudo-F2 populations from cross A, B, and combined A+B.

Blue and red colors represent sensitive and resistant pseudo-F2 plants, respectively. Dashed line

indicates the average biomass for parental NES (blue) and parental NER (red). (b) Genome scan

of pseudo-F2 *A. palmeri* populations from cross A, B, and combined A+B. y-axis indicates LOD

score (logarithm of the odds) statistical estimate, x-axis indicates genome position on scaffolds

levels. The plants were treated with 77g ha$^{-1}$ of tembotrione, and fresh biomass was measured

after 28 d.

684
685 **Fig. 7**. Markers associated with QTLs for tembotrione resistance in *A. palmeri*. (a) – (d) Markers
686 corresponding to the QTL in scaffold 6 (M6). (e) – (h) Markers corresponding to the QTL in
687 scaffold 10 (M10). (i) Marker in scaffold 14 (M14). Boxplots represent the range of biomass
688 values, including the maximum and minimum values, lower and upper quartiles, and the median.
689 The diamond-shaped symbols indicate the average biomass for each genotype.
690

## REFERENCES

**Aarthy, T, Shyam, C & Jugulam, M. 2022**. Rapid metabolism and increased expression of *CYP81E8* gene confer high level of resistance to tembotrione in a multiple-resistant Palmer amaranth (*Amaranthus palmeri* S. Watson). *Frontiers in Agronomy, **4, 1010292***.

**Andersen, CL, Jensen, JL & Ørntoft, TF. 2004**. Normalization of real-time quantitative reverse transcription-PCR data: a model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Res,* **64**: 5245-50.

**Banerjee, A & Roychoudhury, A. 2015**. WRKY proteins: signaling and regulation of expression during abiotic stress responses. *Scientific World Journal,* **2015**: 807560.

**Berger, S, Madeira, PT, Ferrell, J, Gettys, L, Morichetti, S, Cantero, JJ & Nuñez, C. 2016**. Palmer amaranth (*Amaranthus palmeri*) identification and documentation of ALS-resistance in Argentina. *Weed Science,* **64**: 312-320.

**Bolger, AM, Lohse, M & Usadel, B. 2014**. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics,* **30**: 2114-20.

**Bourguet, D & Raymond, M. 1998**. The molecular basis of dominance relationships: the case of some recent adaptive genes. *Journal of Evolutionary Biology,* **11**: 103-122.

**Brazier-Hicks, M, Franco-Ortega, S, Watson, P, Rougemont, B, Cohn, J, Dale, R, Hawkes, TR, Goldberg-Cavalleri, A, Onkokesung, N & Edwards, R. 2022**. Characterization of cytochrome P450s with key roles in determining herbicide selectivity in maize. *ACS Omega.*

**Brazier, M, Cole, DJ & Edwards, R. 2002**. O-glucosyltransferase activities toward phenolic natural products and xenobiotics in wheat and herbicide-resistant and herbicide-susceptible black-grass (*Alopecurus myosuroides*). *Phytochemistry,* **59**: 149-156.

**Broman, KW, Gatti, DM, Simecek, P, Furlotte, NA, Prins, P, Sen, Ś, Yandell, BS & Churchill, GA. 2019**. R/qtl2: Software for Mapping Quantitative Trait Loci with High-Dimensional Data and Multiparent Populations. *Genetics,* **211**: 495-502.

**Chauhan, BS. 2020**. Grand challenges in weed management. *Frontiers in Agronomy,* **1**: 3.

**Chen, S, Zhou, Y, Chen, Y & Gu, J. 2018**. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics,* **34**: i884-i890.

**de Figueiredo, MRA, Barnes, H, Boot, CM, de Figueiredo, ABTB, Nissen, SJ, Dayan, FE & Gaines, TA. 2022**. Identification of a novel 2,4-D metabolic detoxification pathway in 2,4-D-resistant waterhemp (*Amaranthus tuberculatus*). *Journal of Agricultural and Food Chemistry,* **70**: 15380-15389.

**Dimaano, NG, Yamaguchi, T, Fukunishi, K, Tominaga, T & Iwakami, S. 2020**. Functional characterization of cytochrome P450 *CYP81A* subfamily to disclose the pattern of cross-resistance in *Echinochloa phyllopogon*. *Plant Molecular Biology,* **102**: 403-416.

**Doyle, JJ & Doyle, JL. 1990**. Isolation of plant DNA from fresh tissue. *Focus,* **12**: 13-15.

**Gaines, TA, Duke, SO, Morran, S, Rigon, CAG, Tranel, PJ, Küpper, A & Dayan, FE. 2020**. Mechanisms of evolved herbicide resistance. *Journal of Biological Chemistry,* **295**: 10307–10330.

**Gaines, TA, Lorentz, L, Figge, A, Herrmann, J, Maiwald, F, Ott, M-C, Han, H, Busi, R, Yu, Q, Powles, SB & Beffa, R. 2014**. RNA-Seq transcriptome analysis to identify genes involved in metabolism-based diclofop resistance in *Lolium rigidum. The Plant Journal,* **78**: 865-876.

Gaines, TA, Zhang, W, Wang, D, Bukun, B, Chisholm, ST, Shaner, DL, Nissen, SJ, Patzoldt, WL, Tranel, PJ, Culpepper, AS, et al. 2010. Gene amplification confers glyphosate resistance in *Amaranthus palmeri*. *Proceedings of the National Academy of Sciences, USA,* **107**: 1029-1034.

Giacomini, DA, Patterson, EL, Küpper, A, Beffa, R, Gaines, TA & Tranel, PJ. 2020. Co-expression clusters and allele-specific expression in metabolism-based herbicide resistance. *Genome Biology and Evolution,* **12**: 2267–2278.

González-Torralva, F & Norsworthy, JK. 2021. Understanding resistance mechanisms to trifluralin in an Arkansas Palmer amaranth population. *Genes,* **12** 1225.

Han, H, Yu, Q, Beffa, R, González, S, Maiwald, F, Wang, J & Powles, SB. 2021. Cytochrome P450 *CYP81A10v7* in *Lolium rigidum* confers metabolic resistance to herbicides across at least five modes of action. *The Plant Journal,* **105**: 79-92.

Hudson, WH & Ortlund, EA. 2014. The structure, function and evolution of proteins that bind DNA and RNA. *Nat Rev Mol Cell Biol,* **15**: 749-60.

Hwang, J-I, Norsworthy, JK, Carvalho-Moore, P, Barber, LT, Butts, TR & McElroy, JS. 2023. Exploratory analysis on herbicide metabolism and very-long-chain fatty acid production in metolachlor-resistant Palmer amaranth (*Amaranthus palmeri* S. Wats.). *Journal of Agricultural and Food Chemistry,* **71**: 6014-6022.

Iwakami, S, Endo, M, Saika, H, Okuno, J, Nakamura, N, Yokoyama, M, Watanabe, H, Toki, S, Uchino, A & Inamura, T. 2014. Cytochrome P450 *CYP81A12* and *CYP81A21* are associated with resistance to two acetolactate synthase inhibitors in *Echinochloa phyllopogon*. *Plant Physiology,* **165**: 618-629.

Jhala, AJ, Sandell, LD, Rana, N, Kruger, GR & Knezevic, SZ. 2014. Confirmation and control of triazine and 4-hydroxyphenylpyruvate dioxygenase-inhibiting herbicide-resistant Palmer amaranth (*Amaranthus palmeri*) in Nebraska. *Weed Technology,* **28**: 28-38.

Kato, S, Yokota, Y, Suzuki, R, Fujisawa, Y, Sayama, T, Kaga, A, Anai, T, Komatsu, K, Oki, N, Kikuchi, A & Ishimoto, M. 2020. Identification of a cytochrome P450 hydroxylase, *CYP81E22*, as a causative gene for the high sensitivity of soybean to herbicide bentazon. *Theoretical and Applied Genetics,* **133**: 2105-2115.

Kaundun, SS, Hutchings, SJ, Dale, RP, Howell, A, Morris, JA, Kramer, VC, Shivrain, VK & McIndoe, E. 2017. Mechanism of resistance to mesotrione in an *Amaranthus tuberculatus* population from Nebraska, USA. *PLoS One,* **12**: 22.

Kim, D, Paggi, JM, Park, C, Bennett, C & Salzberg, SL. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology,* **37**: 907-915.

Kohlhase, DR, O'Rourke, JA, Owen, MDK & Graham, MA. 2019. Using RNA-seq to characterize responses to 4-hydroxyphenylpyruvate dioxygenase (HPPD) inhibitor herbicide resistance in waterhemp (*Amaranthus tuberculatus*). *BMC Plant Biology,* **19**: 182.

Küpper, A, Peter, F, Zollner, P, Lorentz, L, Tranel, PJ, Beffa, R & Gaines, TA. 2018. Tembotrione detoxification in 4-hydroxyphenylpyruvate dioxygenase (HPPD) inhibitor-resistant Palmer amaranth (*Amaranthus palmeri* S. Wats.). *Pest Management Science,* **74**: 2325-2334.

Li, H & Durbin, R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics,* **25**: 1754-1760.

**Liao, Y, Smyth, GK & Shi, W. 2014**. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics,* **30**: 923-30.
**Lu, H, Liu, Y, Li, M, Han, H, Zhou, F, Nyporko, A, Yu, Q, Qiang, S & Powles, S. 2023**.
Multiple metabolic enzymes can be involved in cross-resistance to 4-
hydroxyphenylpyruvate-dioxygenase-inhibiting herbicides in wild radish. *Journal of*
*Agricultural and Food Chemistry,* **71**: 9302-9313.
**Lu, H, Yu, Q, Han, H, Owen, MJ & Powles, SB. 2019**. Evolution of resistance to HPPD-
inhibiting herbicides in a wild radish population via enhanced herbicide metabolism. *Pest*
*Management Science,* **76**: 1929-1937.
**Montgomery, JS, Giacomini, D, Waithaka, B, Lanz, C, Murphy, BP, Campe, R, Lerchl, J,**
**Landes, A, Gatzmann, F, Janssen, A, et al. 2020**. Draft genomes of *Amaranthus*
*tuberculatus*, *Amaranthus hybridus*, and *Amaranthus palmeri*. *Genome Biology and*
*Evolution,* **12**: 1988-1993.
**Montgomery, JS, Giacomini, DA & Tranel, PJ. 2021**. Molecular confirmation of resistance to
PPO inhibitors in *Amaranthus tuberculatus* and *Amaranthus palmeri*, and isolation of the
G399A PPO2 substitution in *A. palmeri*. *Weed Technology,* **35**: 99-105.
**Montgomery, JS, Morran, S, MacGregor, DR, McElroy, JS, Neve, P, Neto, C, Vila-Aiub,**
**MM, Sandoval, MV, Menéndez, AI, Kreiner, JM, et al. 2023**. The International Weed
Genomics Consortium: Community Resources for Weed Genomics Research. *bioRxiv*:
2023.07.19.549613.
**Murphy, BP, Beffa, R & Tranel, PJ. 2021**. Genetic architecture underlying HPPD-inhibitor
resistance in a Nebraska *Amaranthus tuberculatus* population. *Pest Management Science,*
**77**: 4884-4891.
**Oerke, EC. 2006**. Crop losses to pests. *Journal of Agricultural Science,* **144**: 31-43.
**Palmieri, VE, Alvarez, CE, Permingeat, HR & Perotti, VE. 2022**. A122S, A205V, D376E,
W574L and S653N substitutions in acetolactate synthase (ALS) from *Amaranthus*
*palmeri* show different functional impacts on herbicide resistance. *Pest Management*
*Science,* **78**: 749-757.
**Pan, G, Zhang, XY, Liu, KD, Zhang, JW, Wu, XZ, Zhu, J & Tu, JM. 2006**. Map-based
cloning of a novel rice cytochrome P450 gene *CYP81A6* that confers resistance to two
different classes of herbicides. *Plant Molecular Biology,* **61**: 933-943.
**Pan, L, Guo, Q, Wang, J, Shi, L, Yang, X, Zhou, Y, Yu, Q & Bai, L. 2022**. *CYP81A68*
confers metabolic resistance to ALS and ACCase-inhibiting herbicides and its epigenetic
regulation in *Echinochloa crus-galli*. *Journal of Hazardous Materials,* **428**: 128225.
**Poplin, R, Ruano-Rubio, V, DePristo, MA, Fennell, TJ, Carneiro, MO, Auwera, GAVd,**
**Kling, DE, Gauthier, LD, Levy-Moonshine, A, Roazen, D, et al. 2018**. Scaling
accurate genetic variant discovery to tens of thousands of samples. *bioRxiv*: 201178.
**Rigon, CAG, Gaines, TA, Küpper, A & Dayan, FE. 2020**. Metabolism-based herbicide
resistance, the major threat among the non-target site resistance mechanisms. *Outlooks on*
*Pest Management,* **31**: 162-168.
**Roberts, J & Florentine, S. 2022**. A review of the biology, distribution patterns and
management of the invasive species *Amaranthus palmeri* S. Watson (Palmer amaranth):
Current and future management challenges. *Weed Research,* **62**: 113-122.
**Saika, H, Horita, J, Taguchi-Shiobara, F, Nonaka, S, Nishizawa-Yokoi, A, Iwakami, S,**
**Hori, K, Matsumoto, T, Tanaka, T, Itoh, T, et al. 2014**. A novel rice cytochrome P450

829 gene, *CYP72A31*, confers tolerance to acetolactate synthase-inhibiting herbicides in rice
830 and arabidopsis. *Plant Physiology,* **166**: 1232-40.

831 **Salas-Perez, RA, Saski, CA, Noorai, RE, Srivastava, SK, Lawton-Rauh, AL, Nichols, RL &**
832 **Roma-Burgos, N. 2018**. RNA-seq transcriptome analysis of *Amaranthus palmeri* with
833 differential tolerance to glufosinate herbicide. *PLoS One,* **13**: e0195488.

834 **Seale, M. 2020**. Hot on the trail of DREB2A protein stability. *Plant Physiology,* **183**: 7-8.

835 **Shen, J, Yang, Q, Hao, L, Zhang, L, Li, X & Zheng, M. 2022**. The metabolism of a novel
836 cytochrome P450 (*CYP77B34*) in tribenuron-methyl-resistant *Descurainia sophia* L. to
837 herbicides with different mode of actions. *International Journal of Molecular Sciences,*
838 **23**: 5812.

839 **Sherman, BT, Hao, M, Qiu, J, Jiao, X, Baseler, MW, Lane, HC, Imamichi, T & Chang, W.**
840 **2022**. DAVID: a web server for functional enrichment analysis and functional annotation
841 of gene lists (2021 update). *Nucleic Acids Res,* **50**: 216-221.

842 **Shyam, C, Peterson, DE & Jugulam, M. 2022**. Resistance to 2,4-D in Palmer amaranth
843 (*Amaranthus palmeri*) from Kansas is mediated by enhanced metabolism. *Weed Science,*
844 **70**: 390-400.

845 **Siminszky, B, Corbin, FT, Ward, ER, Fleischmann, TJ & Dewey, RE. 1999**. Expression of a
846 soybean cytochrome P450 monooxygenase cDNA in yeast and tobacco enhances the
847 metabolism of phenylurea herbicides. *Proceedings of the National Academy of Sciences,*
848 *USA,* **96**: 1750-1755.

849 **Wang, H, Xu, D, Wang, S, Wang, A, Lei, L, Jiang, F, Yang, B, Yuan, L, Chen, R, Zhang, Y**
850 **& Fan, W. 2022**. Chromosome-scale *Amaranthus tricolor* genome provides insights into
851 the evolution of the genus *Amaranthus* and the mechanism of betalain biosynthesis. *DNA*
852 *Research,* **30**: 1-15.

853 **Wang, J, Lian, L, Qi, J, Fang, Y, Nyporko, A, Yu, Q, Bai, L & Pan, L. 2023**. Metabolic
854 resistance to acetolactate synthase inhibitors in *Beckmannia syzigachne*: identification of
855 *CYP81Q32* and its transcription regulation. *The Plant Journal,* **115**: 317-334.

856 **Ward, SM, Webster, TM & Steckel, LE. 2013**. Palmer amaranth (*Amaranthus palmeri*): a
857 review. *Weed Technology,* **27**: 12-27.

858 **Williams, MM & Pataky, JK. 2008**. Genetic basis of sensitivity in sweet corn to tembotrione.
859 *Weed Science,* **56**: 364-370.

860 **Zhang, C, Hou, Y, Hao, Q, Chen, H, Chen, L, Yuan, S, Shan, Z, Zhang, X, Yang, Z, Qiu, D,**
861 **et al. 2015**. Genome-wide survey of the soybean GATA transcription factor gene family
862 and expression analysis under low nitrogen stress. *PLoS One,* **10**: e0125174.

863 **Zheng, T, Yu, X, Sun, Y, Zhang, Q, Zhang, X, Tang, M, Lin, C & Shen, Z. 2022**. Expression
864 of a cytochrome P450 gene from Bermuda grass *Cynodon dactylon* in soybean confers
865 tolerance to multiple herbicides. *Plants,* **11**: 949.

**SUPPORTING INFORMATION**

**Fig. S1.** Scheme of crosses performed to generate Pseudo-F2 generation from *Amaranthus palmeri* population HPPD-resistant (NER) x susceptible (NES).

**Fig. S2.** Gene-wise dispersion estimates.

**Fig. S3.** Scatterplot of principal components.

**Fig. S4.** Distance hierarchical clustering heatmap.

**Fig. S5.** Phylogenetic analysis.

**Fig. S6.** Relative gene expression of candidate cytochrome P450 genes in parental NES and NER plants.

**Fig. S7.** Motifs analysis using MEME-suite tool.

**Fig. S8.** Motif sequences found in *Amaranthus palmeri CYP72A219_4284* promoter in sensitive and resistant pseudo-F2 plants.

**Fig. S9.** Dose response of tembotrione in different *Amaranthus palmeri* populations.

**Fig. S10.** Parental tembotrione and metabolites in HPPD resistant populations of *Amaranthus palmeri* over time after tembotrione application.

**Fig. S11.** Chemical structures of tembotrione and the main metabolites.

**Fig. S12.** Representative reverse-phase HPLC chromatogram for *Amaranthus palmeri* populations at 48 HAT with $^{14}$C-Tembotrione.

**Fig. S13.** Relative gene expression of candidate cytochrome P450 genes in different *Amaranthus palmeri* populations.

**Fig. S14.** *CYP72A219_4284* copy number relative to *ALS* in different populations of *Amaranthus palmeri*.

**Fig. S15.** Functional enrichment of genes from QTLs of scaffold 10, scaffold 6 and scaffold 14.

**Table S1.** Gene stability assessed by NormFinder algorithm.

**Table S2.** Primer sequences used for different experiments and their characteristic.

**Table S3.** Constitutively differentially expressed genes between tembotrione resistant and susceptible *Amaranthus palmeri*. Adjusted *P*-value <0.01.

**Table S4.** Logistic equation parameters and the resistance index (RI) of survival (% of untreated control) for different populations of *Amaranthus palmeri* subjected to different doses of tembotrione.

**Table S5.** Logistic equation parameters and the resistance index (RI) of shoot fresh weight (of untreated control) for different populations of *Amaranthus palmeri* subjected to different doses of tembotrione.

**Table S6.** Quantitative trait loci found in Pseudo-F2 *Amaranthus palmeri* in cross B and combined cross A + B for the trait of HPPD resistance.

**Methods S1** Plant material for QTL mapping and RNAseq experiment

**Methods S2.** Yeast transformation

**Methods S3.** Promoter amplification

**Methods S4.** Involvement of CYP72A219_4284 in different HPPD-resistant *A. palmeri* populations

**Methods S5.** Dose-response and segregation analysis

**Methods S6.** Statistical analysis

**Notes S1** Alignment of *CYP81E8* and *CYP72A219_4284* promoter between susceptible and resistant *Amaranthus palmeri*.