Cross-continental variation of herbivore resistance in a global plant invader

Peipei Cao[1#], Zhiyong Liao[2#], Shengyu Wang[1], Lei Zhang[1], Tiantian Lin[3], Madalin

Parepa[4], Jingwen Bi[1], Yujie Zhao[1], Yaolin Guo[1], Oliver Bossdorf[4], Christina L.

Richards[4,5], Stacy B. Endriss[6,7], Jihua Wu[1,8], Ruiting Ju[1], Bo Li[1,3]

[1] Ministry of Education Key Laboratory for Biodiversity Science and Ecological

Engineering, National Observations and Research Station for Wetland Ecosystems of

the Yangtze Estuary, Institute of Biodiversity Science and Institute of Eco-

Chongming, School of Life Sciences, Fudan University, Shanghai, China

[2] CAS Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical

Garden, Chinese Academy of Sciences, Mengla, China

[3] Ministry of Education Key Laboratory for Transboundary Ecosecurity of Southwest

China, Yunnan Key Laboratory of Plant Reproductive Adaptation and Evolutionary

Ecology and Centre for Invasion Biology, Institute of Biodiversity, School of Ecology

and Environmental Science, Yunnan University, Kunming, China

[4] Plant Evolutionary Ecology, Institute of Evolution & Ecology, University of

Tübingen, Tübingen, Germany

[5] Department of Integrative Biology, University of South Florida, Tampa, FL, USA

[6] Graduate Degree Program in Ecology, Colorado State University, Fort Collins, CO,

USA

[7] Department of Bioagricultural Sciences and Pest Management, Colorado State

University, Fort Collins, CO, USA

[8] State Key Laboratory of Herbage Improvement and Grassland Agro-ecosystems,

College of Ecology, Lanzhou University, Lanzhou, China

# These authors contributed equally.

*Corresponding Authors:

Bo Li

Tel: 0086-21-31246681

Email: bool@fudan.edu.cn

Total word count (excluding summary, references and legends):	4654	No. of figures:	4 (4 in colour)
Summary:	196	No. of Tables:	2
Introduction:	1032	No of Supporting Information files:	3 (Fig. S1; TableS1-S2）
Materials and Methods:	1615		
Results:	590		
Discussion:	1304		
Acknowledgements:	113		

Summary

● Successful plant invasions are often explained with adaptation to novel

environments. However, invasive species often occupy broad niches within their

native and introduced ranges, and a true understanding of microevolution during

invasion therefore requires broad sampling of ranges, ideally with a knowledge of

introduction history.

● We tested for genetic differentiation in herbivore resistance among 128

introduced (Europe, North America) and native (China, Japan) populations of the

invasive Japanese knotweed (*Reynoutria japonica*) in two common gardens in the

native range.

● In both common gardens we found that resistance traits of introduced populations

differed from most Chinese native populations, but not from populations in Japan,

the putative sources of introduction. Compared to Chinese populations,

introduced European populations had thicker leaves with a lower C:N ratio but

higher flavonoids contents. In the native range, variation in herbivore resistance

was much more strongly associated with climate of origin than in introduced

populations.

● Our results support the idea that founder effects played a key role in the invasion

of knotweed into Europe and North America, with introduction of particular

resistance phenotypes from Japan. Our study also demonstrates how knowledge

of introduction history can avoid drawing wrong conclusions from observed

biogeographic divergence.

Keywords: common garden, plant defense, plant invasion, rapid evolution,

Reynoutria japonica, secondary chemistry

Introduction

The number of invasive plant species has increased dramatically over the past two centuries (Seebens *et al.*, 2017), causing exceedingly negative impacts on environment (Powell *et al.*, 2013; Bellard *et al.*, 2016; Castro-Díez *et al.*, 2019) and economy (Bradshaw *et al.*, 2016; Diagne *et al.*, 2021). In the context of globalization and climate change, one-sixth of the world's land surface is expected to be highly vulnerable to invasion (Early *et al.*, 2016), and the number of new invasive species is likely to increase further (Seebens *et al.*, 2017, 2021) Thus, a comprehensive understanding of the mechanisms that underlie the successful invasion of alien species is an urgent issue in ecology and evolution (Kempel *et al.*, 2013; Pyšek *et al.*, 2020).

Many eco-evolutionary hypotheses proposed to explain increased performance of invasive populations are related to altered or novel abiotic or biotic environmental conditions in the introduced ranges (Cripps *et al.*, 2006; Mitchell *et al.*, 2006; Montti *et al.*, 2016). The enemy release hypothesis (ERH) states that plant species experience a decrease in regulation by herbivores and other natural enemies, resulting in increased performance (Keane & Crawley, 2002). Since resistance to enemies can be costly (Koricheva, 2002), shifts in enemy pressure may lead to rapid evolutionary changes in growth and defense allocation (Bossdorf *et al.*, 2005; Buswell *et al.*, 2011; Li *et al.*, 2022). The evolution of increased competitive ability (EICA) hypothesis therefore predicts that invasive populations may allocate fewer resources to herbivore defenses, and thus become more competitive than their native conspecifics (Blossey & Nötzold, 1995; Joshi & Vrieling, 2005; Lin *et al.*, 2015). However, despite strong evidence for enemy release (Keane & Crawley, 2002; Liu & Stiling, 2006; Castells *et al.*, 2013; Rotter *et al.*, 2019; Xiao et al., 2020), especially with regard to specialists, the evidence in support of EICA is so far mixed (Bossdorf *et al*., 2005; Felker-Quinn *et al*., 2013; Rotter & Holeski, 2018), with lower herbivore resistance observed in several invasive species (Siemann & Rogers, 2001; Jakobs *et al*., 2004; Huang & Ding, 2016), but similar or even higher resistance in others (Genton *et al*., 2005; Lewis *et al*., 2006; Alba *et al*., 2011; Bhattarai *et al*., 2017; Gruntman *et al*., 2017; Lin *et al*., 2019). Some of these mixed results have been attributed to the different costs of different types of resistance (Koricheva, 2002; Neilson *et al*., 2013), and to the changes in herbivore community composition in the introduced ranges of invasive plants (Müller-Schärer *et al*., 2004; SDH-shifting defence hypothesis; Cripps *et al*., 2006).

Not only biotic conditions, but also altered abiotic environmental conditions may

impose novel selection pressures and thus result in adaptive evolutionary changes in

introduced populations (Buswell *et al*., 2011; Alexander *et al*., 2012; Lee & Kotanen,

2015). For instance, climatic conditions often differ between native and introduced

ranges of invasive species (Early & Sax, 2014; Bocsi *et al*., 2016), and several

previous studies have already demonstrated climate-related trait differentiation

between native and introduced plant populations, suggesting rapid adaptation after

introduction (Etterson *et al*., 2008; Alexander, 2013; Woods & Sultan, 2022).

Furthermore, the existence of similar trait clines along climatic gradients in native and

introduced populations is often also considered as evidence for adaptive post-invasion

evolution (Hulme & Barrett, 2013; Bock *et al*., 2015), through multiple introductions

of pre-adapted genotypes (Neuffer & Hurka, 1999; Keller & Taylor, 2008) and

subsequent selection, i.e. 'sorting out' of standing genetic variation.

Climatic differences can also interact with the effects of herbivores. First, climate

can directly affect plant defenses through physiological constraints and altered

resource availability (Coley, 1985; Wright *et al*., 2004). Second, climatic conditions

also influence the abundance and diversity of herbivores in plant communities (Maron

et al., 2014; Anstett *et al*., 2016; Zhang *et al*., 2016a), thus indirectly driving selection

for particular plant traits (Loughnan & Williams, 2019). Therefore, to understand

invasive plant success and predict habitat vulnerability to future invasion, we need to

understand the impacts of both enemy release and climatic conditions on herbivore

resistance of invasive plant populations. Nevertheless, the relationships between

climate and herbivore resistance have so far rarely been compared between native and

introduced populations (Xiao *et al*., 2020).

Comparisons between native and introduced populations within common gardens

are often facing several challenges (van Kleunen *et al*., 2018; Woods & Sultan, 2022).

First, invasive species often occupy broad climatic niches both in their native and

introduced ranges (Brandenburger *et al*., 2019, 2020; Liu *et al*., 2020), and

comparisons of only few populations may lead to wrong conclusions, simply because

the studied populations do not represent their respective ranges well enough (Yang *et*

al.,2014). Second, when testing for evolution during invasion, the sources of

introduced populations are of course the appropriate reference to invasive

populations. Thus, a lack of knowledge of invasion history may further limit our

ability to correctly identify trait changes in invasive populations (Cano *et al*., 2009;

Bukovinszky *et al*., 2014). Third, because of widespread genotype-by-environment

interactions (Richards *et al*., 2006), the results of common garden studies may to

some extent be garden-specific. For instance, differences in abiotic and biotic

conditions (e.g. climate, interacting species) might influence not only absolute values

of growth and defense traits but also their relative differences between plant origins

(Maron *et al*., 2004; Moloney *et al*., 2009; Qin *et al*., 2013; Yang *et al*., 2021).

Here, we worked with the invasive Japanese knotweed (*Reynoutria japonica*)

and built on a previous cross-continental field survey of the species, which provided

plant materials, along with important metadata, that allowed us to establish two large

common gardens in different climatic zones of the native range, with 55 populations

of origin from native range in China and Japan, and 73 populations from the

introduced ranges in Europe and North America. The worldwide distribution and

relatively clear invasion history of invasive knotweed provided an excellent model to

explore biogeographic divergence in herbivore resistance and resistance traits (leaf

traits and leaf chemistry), as well as their relationships with herbivore pressure and

climatic factors at the collecting sites. We expected that (1) introduced populations

have lower herbivore resistance than native populations, and that (2) traits of native

populations are more strongly associated with climates of origin than those of

introduced populations.

Materials and methods

Study system

Reynoutria japonica (Japanese knotweed) is native to eastern Asia and was introduced

to Europe in the early 1840s and to North America in the 1870s as an ornamental

(Bailey & Conolly, 2000). Along with its sister species *R. sachalinensis* it has become

widely naturalized in both introduced ranges (Barney, 2006; Shimoda & Yamasaki,

2016; Del Tredici, 2017). *R. japonica*, *R. sachalinensis*, and their hybrid *R. ×*

bohemica spread rapidly along river banks and roadsides, often forming dense stands

of hundreds of square meters (Bímová *et al*., 2004; Tiébré *et al*., 2008; Rouifed *et al*.,

2014). Knotweed invasion seriously threatens the biodiversity and integrity of native

ecosystems (Stoll *et al*., 2012; Mincheva *et al*., 2014), which causes substantial

economic damage (Reinhardt *et al*., 2003; DEFRA, 2003). As a major environmental

threat in Europe and North America, Japanese knotweed is listed among the 100

world's worst invasive alien species by IUCN (Lowe *et al*., 2000).

Sample collection

The plant materials we used came from a cross-latitudinal survey of 150 Japanese knotweeds populations in the native range of China and the introduced ranges of North America and Europe (Fig. **1**, Table S1; Irimia *et al*., 2023). In China our survey ranged from the Guangdong province in the South to the Shandong province in the (North-)East, in Europe from Northern Italy to Central Sweden, and in the United States from Georgia to Maine (Fig. **1**, Table S1). In each range we surveyed 50 populations along a 2000 km transect (*c.* every 40 km). The surveys in Europe and North America were done in 2019, that in China in 2020. For full details on the field survey please see Irimia *et al*. (2023). Briefly, at each sampling site we confirmed the knotweed taxon based on morphological characters (Bailey *et al*., 2009), laid a 30 m transect for sampling, selected five knotweed stems at regular intervals along the transect, and collected rhizomes from these individuals.

In addition to the main field survey, we also collected six native *R. japonica* populations from around Nagasaki in Japan, the region believed to be a key source of European and North American introductions (Bailey & Conolly, 2000; Del Tredici, 2017). A recent phylogenetic analysis based on chloroplast DNA (Zhang *et al*., unpublished data) has confirmed this hypothesis. Between 28 April and 4 May 2021, we collected rhizomes from 5-8 individuals separated by at least 6 m in each Japanese population. The rhizomes from Japan, Europe and North America were imported and temporally grown in a quarantine glasshouse at Xishuangbanna Tropical Botanical Garden, whereas the Chinese rhizomes were only stored at 4 °C for 2 weeks and then planted in pots to minimize maternal effects before setting up the common gardens. Because of legal restrictions, only one rhizome per North American populations could be imported prior to setting up the experiment.

Common garden experiments

To test for heritable variation among knotweed plants from different ranges and populations, we set up two large common garden experiments, one at Fudan University in Shanghai (31°20′N, 121°30′E) and the other at Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences (21°41′N, 101°25′E) in the Yunnan province. The two common garden locations differed in their climatic conditions, with warmer and moister conditions in Xishuangbanna (average March-October

temperature 24.1 °C; average monthly precipitation 171 mm) compared to Shanghai

(average temperature 20.3 °C; monthly precipitation 116 mm) (http://www.nmic.cn/;

He *et al.*, 2001; Song *et al.*, 2010; Yu *et al.*, 2021).

Because of import restrictions and variable cultivation success during quarantine

conditions, we had to omit some introduced populations and ended up with a total of

128 *R. japonica* populations of origin: 55 populations from the native range (50 from

China, 5 from Japan) and 73 populations from the introduced ranges (46 from Europe,

27 from North America) (Fig. **1**, Table S1). Both gardens were set up in open areas on

flat ground covered with artificial grass mats.

Prior to planting the experiments, we cut rhizomes from each individual and kept

them at 4 °C for at least four weeks to facilitate sprouting success. We cut all

rhizomes to a size of 3-10 g (at least one intact node), removed the fine roots, and, as

a measure of initial size, determined the fresh weight of each cutting. On 11 March

2022, all rhizomes were treated with fungicide and planted separately into 10.8 L pots

filled with the same potting soil (Pindstrup substrate 0–10 mm, Pindstrup Mosebrug

A/S, Denmark) in both gardens. Because of limited rhizome availability, we

eventually planted a total of 518 *R. japonica* individuals in the Shanghai garden (27

individuals from 27 North Amercian populations; 218 individuals from 46 European

populations; 239 individuals from 50 Chines populations; 34 individuals from 5

Japanese populations), and 463 individuals in the Xishuangbanna garden (16

individuals from 16 North American populations; 190 individuals from 46 European

populations; 232 individuals from 50 Chinese populations; 25 individuals from 5

Japanese populations). In each garden, we arranged the plants in five blocks, ideally

with one individual from each of the populations in each block, random assignment to

blocks, and random order within blocks. To avoid aboveground interference, the

distances between pots were all at least 90 cm. To avoid nutrient depletion, we added

10 g Osmocote fertilizer (Osmocote plus 801, N: P: K 16:8:12, Everris International

B.V., Heerlen, Netherlands) to each pot at the beginning of the experiment, and once

again in the middle of the experiment. Throughout the experiment, we watered the

plants whenever the soil had become dry. To avoid losses of water and nutrients, all

pots were individually placed on plastic trays. The pots, soils, and fertilizers used in

the two common gardens were identical.

Herbivore damage and resistance traits

To quantify variation in plant herbivore resistance, we estimated the degree of

herbivore damage experienced by each individual in each garden, determined the

levels of leaf secondary metabolites (lignin, flavonoids, alkaloids), and measured

additional leaf traits that are often associated with palatability to herbivores (leaf

toughness, leaf thickness, leaf ratios C : N) (Feng *et al*., 2011; Lin *et al*., 2015). In the

middle of growing season (July 2022), we estimated herbivore damage as the

percentage of leaf area eaten on each plant individual, separately for beetle and

caterpillar damage (based on chewing modes and leaf notch types; Fig. S2). Two days

later, we sampled five fresh, fully developed leaves (leaves 1, 2, 4, 5 and 6 from the

top) from the tallest shoot of each plant individual, and measured the leaf thickness of

each plant with a digital micrometer (Digimatic Outside Micrometer, Mitutoyo,

Japan), and its toughness with a penetrometer (FA10, SAUTER, Balingen, Germany)

in the Shanghai garden, and with a mechanical testing machine (ZQ990A, Dongguan

Zhiqu Precision Instrument Co., Ltd, China) in the Xishuangbanna garden. For each

plant, we then estimated the leaf thickness and toughness as the averages of the five

measurements.

Finally, we dried all leaves at 60 °C for 72 h, and used these to analyze leaf

chemistry. After grinding samples to the required particle size with a ball mill

(MM400, Retsch, Germany), we measured total C and N with an organic elemental

analyzer (FlashSmart™ Elemental Analyzer, Thermo-Fisher Scientific, USA) via

thermal combustion and TCD/IR detection of CO_2/N_2, and also measured leaf lignin,

alkaloids and total flavonoids using the MZS-1-G, SWJ-1-Y and LHT-1-G test kits

(Suzhou Comin Biotechnology Co., Ltd., Suzhou, China), respectively. In total, we

analysed 981 plant samples, 518 from the Shanghai garden and 463 individuals from

the Xishuangbanna garden.

Statistical analyses

To test for range differences in plant traits and herbivore damage, we fitted linear

mixed models in R version 4.2.1 (R Core Team, 2022), with range (China, Japan,

Europe, North America) as fixed effect, and population and block as random effects.

To account for variation in initial size, our models included initial rhizome weight as

covariates. We assessed the significance of fixed effects through Type III Wald chi-

squared tests using the *car* package (Fox & Weisberg, 2018). For traits that displayed

a significant range effect ($P < 0.05$), we then conducted Tukey post-hoc tests with the

emmeans and *pairs* functions (*emmeans* package; Lenth, 2018). Where necessary, we

log-transformed the herbivore damage data to normalize the distribution of residuals.

To test for associations between leaf traits and herbivore damage, we calculated

population-level spearman correlation coefficients for each pairwise combination of

resistance trait (leaf traits and leaf chemistry) and herbivore damage, separately for

each garden, using the *Hmisc* package (Hauke & Kossowski, 2011).

Finally, we tested whether the climatic conditions of population origins were

associated with the resistance traits of native and invasive populations. For this, we

obtained eight bioclimatic variables from the WorldClim database (Fick & Hijmans,

2017) that seemed particularly meaningful for characterising growing and

overwintering conditions for knotweed plants: annual temperature (bio1), maximum

temperature of the warmest month (bio4), minimum temperature of the coldest month

(bio5), temperature seasonality (bio6), mean annual precipitation (bio12),

precipitation of the wettest month (bio13), precipitation of the driest month (bio14),

and precipitation seasonality (bio15). We simplified these climate data through a

principal component analyses (PCA) using the *prcomp* function. The first and second

principal component ('climate PC1' and 'climate PC2' hereafter) explained 54% and

21% of the variance in the eight climatic variables across all populations, respectively,

with climate PC1 mainly related to mean temperature as well as cold and precipitation

extremes, and climate PC2 related to temperature maxima and seasonality (Fig. S1,

Table S2). We then performed population-level linear regression analyses with either

climate PC1 or climate PC2 as explanatory variables, and the common-garden

averages of the resistance traits of native or introduced populations as dependent

variables (Moreira *et al*., 2018; Galmán *et al*., 2021).

Results

We found that on average the traits of invasive populations often differed from those

of native populations. However, these overall patterns were largely driven by the

native populations from China which were distinct from European and North

American populations, whereas the native populations from Japan were often similar

to the invasive ones.

Range differences in herbivore damage

The average levels of herbivore damage differed substantially between the two

common gardens: in Xishuangbanna the percentages of leaf area eaten by beetles and caterpillars were three and 19 times higher, respectively, than in Shanghai (Fig. **2**). In Shanghai, plants from Chinese populations experienced significantly higher levels of caterpillar herbivory than plants from Europe or North America (Fig. **2b**), and the same was true for beetle damage in Xishuangbanna, where the Chinese populations were also most strongly attacked (Fig. **2c**). Interestingly, Chinese populations experienced less caterpillar damage than European populations in Xishuangbanna, but we never found any significant differences between the two introduced ranges, or between Japanese populations and the invasive populations, in the two gardens.

Range differences in resistance traits

Plants from native Chinese populations were also distinct in their resistance traits. In both common gardens, Chinese plants had significantly thinner leaves than plants from invasive populations (Fig. **3a,d**), and in the Shanghai garden they also had higher leaf ratios C : N and lower leaf flavonoids than plants from both invasive ranges (Figs. **3c** and **4c**). Moreover, Chinese populations had tougher leaves than European populations in the Shanghai garden (Fig. **3b**) and higher leaf ratios C : N than North American populations in the Xishuangbanna garden (Fig. **3f**). We did not find any range differences in leaf lignin and leaf alkaloids between native and invasive populations Fig. **4a,b,d,e**), and there were no significant differences in leaf traits between native populations from Japan and the invasive populations (Figs. **3** and **4**).

Correlations between resistance traits and herbivore damage

In the Shanghai garden, leaf damage by caterpillars was negatively correlated with leaf thickness and leaf flavonoids, but positively with leaf ratios C : N (Table 1). In contrast, in the Xishuangbanna garden, beetle herbivory was negatively correlated with leaf thickness and flavonoids, but caterpillar herbivory was positively correlated with leaf thickness (Table 1).

Correlations between climates of origin and resistance traits

Significant correlations between climates of origin and resistance traits were frequent among native populations but much less common among introduced populations (10 vs. 3 correlations in native vs introduced plants; Table 2). In both common gardens,

leaf thickness was negatively correlated with climate PC1 but positively correlated
with climate PC2 for native but not invasive plants, and leaf ratios C : N of native
populations was negatively related to climate PC1. Only in the Xishuangbanna
garden, leaf ratios C : N of invasive populations was positively correlated to climate
PC2. For native populations, leaf flavonoids were positively related to climate PC1
but negatively to PC2 in both gardens, but for invasive populations we found a
positive climate PC1-flavonoid correlation only in Xishuangbanna, and a negative
correlation of flavonoids with climate PC2 only in Shanghai (Table 2).

In other words, native populations from warmer, humid sites with high
temperature seasonality tended to have thicker leaves with a higher ratios C : N and
lower flavonoids contents (Table 2), where for invasive populations climate
associations were rarer and more variable: in Shanghai, higher flavonoid contents of
invasive populations were associated with high summer temperature and low
seasonality whereas in Xishuangbanna lower flavonoids were associated with warm
and humid areas of origin (Table 2).

Discussion
Successful plant invasions are often explained by evolutionary changes in plant traits.
However, the power of previous common garden comparisons of native versus
introduced populations has often remained limited. Here, we tested for genetic
differentiation in herbivore resistance among 128 native and introduced populations
of invasive knotweed (*Reynoutria japonica*) in two common gardens in the native
range. We found that on average introduced populations from Europe and North
America differed in several resistance traits from native Asian populations, but that
these overall differences were driven by native populations from China, whereas the
resistance traits of native Japanese populations were generally similar to populations
from the introduced range. Our study supports – at the level of phenotypes – a
Japanese source of knotweed invasions, and that founder effects – rather than post-
invasion evolution – explains overall range differences in knotweed traits.

Range differences in herbivore resistance
The EICA hypothesis predicts that because of a release from native herbivores
introduced populations might have reduced herbivore defenses (Blossey & Notzold,
1995). We found that introduced European and North American populations were

indeed significantly different in many leaf traits from native Chinese populations, but that there were little differences between introduced populations and the putative source populations from Japan. Our results thus demonstrate the importance of accounting for invasion history when testing for evolution during invasions (Colautti & Lau, 2015; Brandenburger *et al.*, 2020).

We found no significant differences in many herbivore resistance traits such as leaf ratios C : N, leaf toughness, leaf flavonoids, lignin or alkaloids between introduced European and North American and native Japanese populations in both common gardens. There were also no differences in the herbivore damage of plants in both gardens, which represented two different natural herbivore environments in the native range. Our results thus did not support the EICA prediction that invasive populations have evolved lower herbivore resistance than their native sources (Blossey & Notzold, 1995). Several studies with other invasive species have shown no difference or even higher resistance of invasive populations in common garden experiments, e.g. with *Chromolaena odorata* (Liao *et al.*, 2014), *Brassica nigra* (Oduor *et al.*, 2011), or *Verbascum thapsus* (Endriss *et al.*, 2018). We are only aware of one previous common garden study with native and introduced populations of *R. japonica* (Rouifed *et al.* 2018). It compared only five native Japanese and five invasive French populations in a common garden in the introduced range, but it also found that introduced populations did not differ in their composition of secondary metabolites, stem stiffness and leaf thickness, but that they had even higher leaf toughness than native Japanese populations. As the Japanese populations used in our experiments – as well as those used in Rouifed *et al.* (2018) – were from the putative source region of knotweed introductions to Europe and North America, our results thus indicate that the introduction of plants with particular resistance profiles, rather than post-invasion evolution, underlies the trait divergence between the native and introduced range, and may have played a role in the invasion success of *R. japonica*.

We found that herbivore damage varied substantially between the two common gardens, and related to different leaf traits. For instance, in the Shanghai garden damage by caterpillars was negatively associated with leaf thickness and flavonoids, but positively with leaf ratios C : N, whereas in Xishuangbanna, the same type of damage was only positively associated with leaf thickness. Part of these divergent results might have resulted from the different herbivore communities in the two common gardens, e.g. if different herbivores have different feeding preferences (Van

der Meijden, 1996; Bossdorf *et al*., 2004; Yang *et al*., 2014). In Xishuangbanna, we found that damage by beetles was associated with different traits than damage by caterpillars: beetles preferred plants with thinner leaves and lower flavonoid levels, whereas caterpillars preferred plants with thicker leaves. Besides the herbivore communities, the two common gardens of course also differed in climatic conditions, including temperature, precipitation and solar radiation, and since leaf traits could also be related to other environmental drivers, e.g. influence drought or UV tolerance (Levitt & Lovett, 1985; Strauss & Agrawal, 1999; Harborne & Williams, 2000; Peter Constabel *et al*., 2014; Barton & Boege, 2017; Li *et al*., 2022), this may have further contributed to divergent herbivory-leaf trait relationships. More generally, our results demonstrate the environmental contingency of common garden results, including when testing for variation in herbivore resistance (Maron *et al*., 2004; Qin *et al*., 2013; Yang *et al*., 2014; Bossdorf *et al*., 2005; Yang *et al*., 2021), and hence the value of working with multiple common gardens.

Associations between climates of origin and herbivore resistance

We found that the climatic conditions at the collecting sites were significantly associated with resistance traits in both common gardens, but only for plants from the native range, whereas for plants from the introduced range there were only very few significant associations. Many previous studies of invasive plants have documented the development of parallel clinal patterns in ecological traits among populations of native and introduced ranges (Agrawal *et al*., 2015; Leger & Rice, 2007; Etterson *et al*., 2008; Rosche *et al*., 2019; Hodgins *et al*., 2020) and have usually interpreted this as evidence for rapid post-invasion evolution and adaptation. However, there are also studies of successful invasive plants with unparalleled (Bhattarai *et al*., 2017; Yang *et al*., 2021; Woods & Sultan, 2022) or no clinal patterns at all (Endriss *et al*., 2018; Sun & Roderick, 2019), and our results also provide little evidence for post-invasion genetic differentiation of introduced European or North American populations in relation to climate.

A simple reason for the observed lack of genetic differentiation could be that there was too little genetic variation for natural selection to act on. The majority of invasive *R. japonica* populations are thought to be descendants of a single introduced clone (Hollingsworth & Bailey, 2000; Richards *et al*., 2012; Gaskin *et al*., 2014; Groeneveld *et al*., 2014; Zhang *et al*., 2016b). However, so far this knowledge is

based on low-resolution molecular studies of limited numbers of populations, and we

clearly need broader and higher-resolution population genomic analyses to evaluate

this explanation. Moreover, even if little genetic variation was introduced in the first

place, novel mutations can sometimes be common enough to result in genetic

variation that can be exploited for adaptation, even within a relatively short period of

time after an invasion (Lynch & Conery, 2000, Ossowski *et al.*, 2010). In any case, an

alternative explanation for the lack of climate-related genetic differentiation could be

that adaptation did not occur *yet*. Although rapid evolutionary changes have

frequently been reported for invasive plants, examples from the invasion literature

show that geographic clines usually develop 50-150 generations after introduction

(Moran & Alexander, 2014), but many of these examples came from short-lived

plants. *Reynoutria japonica* was introduced to Europe *c.* 170 years ago and some

decades later to North America, but it began to expand rapidly only in the past 50-100

years (Bailey & Conolly, 2000), and it is a perennial, so it is a possibility that there

was not enough time. Again, genomic approaches will help to answer this question.

Conclusion

Our study combined replicate common gardens and a biogeographical approach with

samples spanning an exceptionally large climatic gradient from both introduced and

native ranges to understand evolution of herbivore resistance in the invasive species

R. japonica. We found that the resistance traits of introduced European and North

American populations differed from most Chinese native populations but were

generally similar to Japanese populations that are putative sources of the

introductions. Thus, we find little evidence for post-invasion evolution but that plants

with particular resistance profiles have been introduced to Europe and North America,

and may have played a role in the invasion success of *R. japonica*. Our study

demonstrates the importance of accounting for invasion history when testing for

evolution during invasion.

Acknowledgements

We thank Fatima Elkott, Christiane Karasch-Wittmann, Elodie Kugler, Rongjin Li,

Kaitao Liu, Rui Min, Jeannie Mounger, Julia Rafalski, Conner Richardson, Eva

Schloter, Sabine Silberhorn, Weihan Zhao, Wenchao Zhong, Zhiming Zhong and Xin

Zhuang for help with field collections, and Jieren Jin, Guangye Li, Shibo Tian,

Chaonan Wang and Niyao Xiang with common garden experiments. We also thank

the Center for Gardening and Horticulture and the Seed Bank of Tropical Plant

Resources at XTBG for providing quarantine facilities and cold rooms. This study

was supported by the National Natural Science Foundation of China (grant

31961133028), the German Federal Ministry of Education and Research (MOPGA

Project 306055) and the German Research Foundation (grant 431595342).

477   Conflict of interest statement

478   The authors declare no conflict of interests.

480   Author contributions

481   PC, ZL, MP, OB, CLR and BL designed the experiments. PC, ZL, SW, LZ, YZ and

482   JB conducted the research. PC and ZL analyzed the data and wrote the first draft. All

483   authors contributed to the revision and improvement of the manuscript.

486   Data availability statement

487   Data available from the Dryad Digital Repository (private for peer review):

488   https://datadryad.org/stash/share/EYcQ6ThtOzgWS6oC2dS3Xlj7ZhylK9veRFDySOd

489   EZQ0, DOI is 10.5061/dryad.qbzkh18r4.

Table 1 Spearman's rank correlations (r_s) between leaf traits and leaf herbivory, tested across 128 native and introduced populations of Japanese knotweed, when grown in two different common gardens in Shanghai and Xishuangbanna

	Thickness	Toughness	Ratios C : N	Lignin	Alkaloids	Flavonoids
Shanghai						
% Leaf area eaten by beetles	0.13	-0.06	-0.11	-0.11	0.10	-0.07
% Leaf area eaten by caterpillars	**-0.40*****	0.12	**0.34*****	-0.14	0.05	**-0.35*****
Xishuangbanna						
% Leaf area eaten by beetles	**-0.30****	-0.13	-0.06	0.06	0.00	**-0.31*****
% Leaf area eaten by caterpillars	**0.19***	-0.08	-0.03	0.15	0.03	0.11

Significant correlations are in bold. Significance levels: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$.

Table 2 Associations between climates of origin (first two PCs from a PCA of eight bioclimatic variables, see Table S2) and variation in leaf traits across native (N; China and Japan) and introduced (I; Europe and North America) Japanese knotweed populations, when grown in two common gardens in Shanghai and Xishuangbanna in China

		Climate PC1				Climate PC2		
		β	P	R^2		β	P	R^2
Shanghai								
Thickness	N:	**-0.01**	**0.013**	**0.09**		**0.02**	**0.001**	**0.18**
	I:	0.00	0.855	0.00		0.00	0.747	0.00
Ratios C : N	N:	**-1.29**	**0.022**	**0.08**		0.79	0.170	0.02
	I:	-0.40	0.157	0.01		-0.30	0.286	0.00
Flavonoids	N:	**1.57**	**0.002**	**0.16**		**-1.86**	**0.000**	**0.23**
	I:	-0.35	0.187	0.01		**-0.64**	**0.015**	**0.07**
Xishuangbanna								
Thickness	N:	**-0.01**	**0.016**	**0.09**		**0.02**	**0.000**	**0.22**
	I:	0.00	0.641	0.00		0.00	0.984	0.00
Ratios C : N	N:	**-2.00**	**0.000**	**0.28**		0.89	0.075	0.04
	I:	0.56	0.189	0.01		**1.38**	**0.001**	**0.16**
Flavonoids	N:	**2.73**	**0.000**	**0.30**		**-2.61**	**0.000**	**0.27**
	I:	**1.19**	**0.002**	**0.14**		0.41	0.298	0.00

The values are regression coefficients and their significance levels (*P* values < 0.05 in bold), and adjusted model R^2 from linear regressions.

Fig. 1. Geographic origins of the studied knotweed (*Reynoutria japonica*) clones, the two common gardens, and performance variation of different knotweed origins. (a) The geographic locations where rhizomes of *R. japonica* had been collected in the introduced ranges of North America (1; *n* = 27 populations) and Europe (2; *n* = 46), and in the native ranges of China (3; *n* = 50) and Japan (4; *n* = 5), and stars represent the locations of Shanghai (Fudan University Campus) and Xishuangbanna (Xishuangbanna Tropical Botanical Garden, Yunnan) common gardens. (b) Aerial photos of the two common garden experiments in Shanghai and Xishuangbanna. (c) Representative photos illustrating average growth differences between the two gardens and among differential ranges of origin in mid-July.

Fig. 2. Average levels of herbivores damage observed in *R. japonica* plants from China (CN), Japan (JA), North America (NA) and Europe (EU) when grown in common gardens in Shanghai (a, b) and Xishuangbanna (c, d). The values are adjusted means and SEs from ANOVA, with different letters above error bars indicating significant group differences based on Tukey's HSD post-hoc tests. *Chi-square* statistics and significance level of regression models are also shown. ns, not significant.

Fig. 3. Average levels of leaf traits observed in *R. japonica* plants from China (CN), Japan (JA), North America (NA) and Europe (EU) when grown in common gardens in Shanghai (a-c) and Xishuangbanna (d-f). The values are adjusted means and SEs from ANOVAs, with different letters above error bars indicating significant group differences based on Tukey's HSD post-hoc tests. *Chi-square* statistics and significance level of regression models are also shown. ns, not significant.

Fig. 4. Average levels of leaf chemistry observed in *R. japonica* plants from China (CN), Japan (JA), North America (NA) and Europe (EU) when grown in common gardens in Shanghai (a-c) and Xishuangbanna (d-f). The values are adjusted means and SEs from ANOVAs, with different letters anove error bars indicate significant group differences based on Tukey's HSD post-hoc tests. *Chi-square* statistics and significance level of regression models are also shown. ns, not significant.

References:

Agrawal AA, Hastings AP, Bradburd GS *et al*. 2015. Evolution of plant growth and defense in a continental introduction. *The American Naturalist* **186**: E1-E15.

Alba C, Deane Bowers M, Blumenthal D *et al*. 2011. Evolution of growth but not structural or chemical defense in *Verbascum thapsus* (common mullein) following introduction to North America. *Biological Invasions* **13**: 2379-2389.

Alexander JM, van Kleunen M, Ghezzi R *et al*. 2012. Different genetic clines in response to temperature across the native and introduced ranges of a global plant invader. *Journal of Ecology* **100**: 771-781.

Alexander JM. 2013. Evolution under changing climates: climatic niche stasis despite rapid evolution in a non-native plant. *Proceedings of the Royal Society B: Biological Sciences* **280**: 20131446.

Anstett DN, Nunes KA, Baskett C *et al*. 2016. Sources of controversy surrounding latitudinal patterns in herbivory and defense. *Trends in Ecology & Evolution* **31**: 789-802.

Bailey JP, Bímová K, Mandák B. 2009. Asexual spread versus sexual reproduction and evolution in Japanese Knotweed *s.l.* sets the stage for the "Battle of the Clones". *Biological Invasions* **11**: 1189-1203.

Bailey JP, Conolly AP. 2000. Prize-winners to pariahs-A history of Japanese Knotweed *s.l.* (Polygonaceae) in the British Isles. *Watsonia* **23**: 93-110.

Barney JN. 2006. North American history of two invasive plant species: phytogeographic distribution, dispersal vectors, and multiple introductions. *Biological Invasions* **8**: 703-717.

Barton KE, Boege K. 2017. Future directions in the ontogeny of plant defence: understanding the evolutionary causes and consequences. *Ecology Letters* **20**: 403-411.

Bellard C, Cassey P, Blackburn TM. 2016. Alien species as a driver of recent extinctions. *Biology Letters* **12**: 20150623.

Bhattarai GP, Meyerson LA, Anderson J *et al*. 2017. Biogeography of a plant invasion: genetic variation and plasticity in latitudinal clines for traits related to herbivory. *Ecological Monographs* **87**: 57-75.

Bímová K, Mandák B, Kašparová I. 2004. How does Reynoutria invasion fit the various theories of invasibility? *Journal of Vegetation Science* **15**: 495-504.

Blossey B, Notzold R. 1995. Evolution of increased competitive ability in invasive

nonindigenous plants: a hypothesis. *Journal of Ecology* **83**: 887-889.

Bock DG, Caseys C, Cousens RD *et al.* **2015.** What we still don't know about invasion genetics. *Invasion genetics: The Baker and Stebbins Legacy* **24**: 2277-2297.

Bocsi T, Allen JM, Bellemare J *et al.* **2016.** Plants' native distributions do not reflect climatic tolerance. *Diversity and Distributions* **22**: 615-624.

Bossdorf O, Auge H, Lafuma L *et al.* **2005.** Phenotypic and genetic differentiation between native and introduced plant populations. *Oecologia* **144**: 1-11.

Bossdorf O, Prati D, Auge H *et al.* **2004.** Reduced competitive ability in an invasive plant. *Ecology Letters* **7**: 346-353.

Bradshaw CJA, Leroy B, Bellard C *et al.* **2016.** Massive yet grossly underestimated global costs of invasive insects. *Nature Communications* **7**: 12986.

Brandenburger CR, Kim M, Slavich E *et al.* **2020.** Evolution of defense and herbivory in introduced plants—Testing enemy release using a known source population, herbivore trials, and time since introduction. *Ecology and Evolution* **10**: 5451-5463.

Brandenburger CR, Sherwin WB, Creer SM *et al.* **2019.** Rapid reshaping: the evolution of morphological changes in an introduced beach daisy. *Proceedings of the Royal Society B* **286**: 20181713.

Bukovinszky T, Gols R, Agrawal AA *et al.* **2014.** Reciprocal interactions between native and introduced populations of common milkweed, *Asclepias syriaca*, and the specialist aphid, *Aphis nerii*. *Basic and Applied Ecology* **15**: 444-452.

Buswell JM, Moles AT, Hartley S. 2011. Is rapid evolution common in introduced plant species? *Journal of Ecology* **99**: 214-224.

Cano L, Escarré J, Vrieling K *et al.* **2009.** Palatability to a generalist herbivore, defence and growth of invasive and native *Senecio* species: testing the evolution of increased competitive ability hypothesis. *Oecologia* **159**: 95-106.

Castells E, Morante M, Blanco-Moreno JM *et al.* **2013.** Reduced seed predation after invasion supports enemy release in a broad biogeographical survey. *Oecologia* **173**: 1397-1409.

Castro-Díez P, Vaz AS, Silva JS *et al.* **2019.** Global effects of non-native tree species on multiple ecosystem services. *Biological Reviews* **94**: 1477-1501.

Colautti RI, Lau JA. 2015. Contemporary evolution during invasion: evidence for differentiation, natural selection, and local adaptation. *Invasion Genetics: The Baker and Stebbins Legacy* **24**: 1999-2017.

Coley PD, Bryant JP, Chapin III FS. 1985. Resource availability and plant antiherbivore defense. *Science* **230**: 895-899.

Cripps MG, Schwarzländer M, McKenney JL *et al*. 2006. Biogeographical comparison of the arthropod herbivore communities associated with *Lepidium draba* in its native, expanded and introduced ranges. *Journal of Biogeography* **33**: 2107-2119.

Del Tredici P. 2017. The introduction of Japanese knotweed, *Reynoutria japonica*, into North America. *The Journal of the Torrey Botanical Society* **144**: 406-416.

Department of Environment, Food and Rural Affairs. 2003. Review of non-native species policy. London, UK: DEFRA Publications.

Diagne C, Leroy B, Vaissière AC *et al*. 2021. High and rising economic costs of biological invasions worldwide. *Nature* **592**: 571-576.

Early R, Bradley BA, Dukes JS *et al*. 2016. Global threats from invasive alien species in the twenty-first century and national response capacities. *Nature Communications* 7: 12485.

Early R, Sax DF. 2014. Climatic niche shifts between species' native and naturalized ranges raise concern for ecological forecasts during invasions and climate change. *Global Ecology and Biogeography* **23**: 1356-1365.

Endriss SB, Alba C, Norton AP *et al*. 2018. Breakdown of a geographic cline explains high performance of introduced populations of a weedy invader. *Journal of Ecology* **106**: 699-713.

Etterson JR, Delf DE, Craig TP *et al*. 2008. Parallel patterns of clinal variation in *Solidago altissima* in its native range in central USA and its invasive range in Japan. *Botany* **86**: 91-97.

Feng YL, Li YP, Wang RF *et al*. 2011. A quicker return energy-use strategy by populations of a subtropical invader in the non-native range: a potential mechanism for the evolution of increased competitive ability. *Journal of Ecology* **99**: 1116-1123.

Fick SE, Hijmans RJ. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology* **37**: 4302-4315.

Fox J, Weisberg S. 2018. *An R companion to applied regression*. Los Angeles, USA: Sage publications.

Galmán A, Abdala-Roberts L, Wartalska P *et al*. 2021. Elevational gradients in constitutive and induced oak defences based on individual traits and their

correlated expression patterns. *Oikos* **130**: 396-407.

Gaskin JF, Schwarzländer M, Grevstad FS *et al*. 2014. Extreme differences in population structure and genetic diversity for three invasive congeners: knotweeds in western North America. *Biological Invasions* **16**: 2127-2136.

Genton BJ, Kotanen PM, Cheptou PO *et al*. 2005. Enemy release but no evolutionary loss of defence in a plant invasion: an inter-continental reciprocal transplant experiment. *Oecologia* **146**: 404-414.

Groeneveld E, Belzile F, Lavoie C. 2014. Sexual reproduction of Japanese knotweed (Fallopia japonica *s.l.*) at its northern distribution limit: new evidence of the effect of climate warming on an invasive species. *American Journal of Botany* **101**: 459-466.

Gruntman M, Segev U, Glauser G *et al*. 2017. Evolution of plant defences along an invasion chronosequence: defence is lost due to enemy release–but not forever. *Journal of Ecology* **105**: 255-264.

Harborne JB, Williams CA. 2000. Advances in flavonoid research since 1992. *Phytochemistry* **55**: 481-504.

Hauke J, Kossowski T. 2011. Comparison of values of Pearson's and Spearman's correlation coefficients on the same sets of data. *Quaestiones Geographicae* **30**: 87-93.

He FF, Xue J, Mu HZ. 2001. Estimation of total solar radiation and its temporal and spatial distributions over Shanghai area. *Resources Science* **32**: 693-700.

Hodgins KA, Guggisberg A, Nurkowski K *et al*. 2020. Genetically based trait differentiation but lack of trade-offs between stress tolerance and performance in introduced *Canada thistle*. *Plant Communications* **1**: 100116.

Hollingsworth ML, Bailey JP. 2000. Evidence for massive clonal growth in the invasive weed *Fallopia japonica* (Japanese Knotweed). *Botanical Journal of the Linnean Society* **133**: 463-472.

Huang W, Ding J. 2016. Effects of generalist herbivory on resistance and resource allocation by the invasive plant, *Phytolacca americana*. *Insect Science* **23**: 191-199.

Hulme PE, Barrett SCH. 2013. Integrating trait-and niche-based approaches to assess contemporary evolution in alien plant species. *Journal of Ecology* **101**: 68-77.

Irimia RE, Zhao W, Cao P *et al*. 2023. Cross-continental shifts of ecological strategy in a global plant invader. Under review.

Jakobs G, Weber E, Edwards PJ. 2004. Introduced plants of the invasive *Solidago gigantea* (Asteraceae) are larger and grow denser than conspecifics in the native range. *Diversity and Distributions* **10**: 11-19.

Joshi J, Vrieling K. 2005. The enemy release and EICA hypothesis revisited: incorporating the fundamental difference between specialist and generalist herbivores. *Ecology Letters* **8**: 704-714.

Keane RM, Crawley MJ. 2002. Exotic plant invasions and the enemy release hypothesis. *Trends in Ecology & Evolution* **17**: 164-170.

Keller SR, Taylor DR. 2008. History, chance and adaptation during biological invasion: separating stochastic phenotypic evolution from response to selection. *Ecology Letters* **11**: 852-866.

Kempel A, Chrobock T, Fischer M *et al*. 2013. Determinants of plant establishment success in a multispecies introduction experiment with native and alien species. *Proceedings of the National Academy of Sciences* **110**: 12727-12732.

Koricheva J. 2002. Meta-analysis of sources of variation in fitness costs of plant antiherbivore defenses. *Ecology* **83**: 176-190.

Lee Y, Kotanen PM. 2015. Differences in herbivore damage and performance among *Arctium minus* (burdock) genotypes sampled from a geographic gradient: a common garden experiment. *Biological Invasions* **17**: 397-408.

Leger EA, Rice KJ. 2007. Assessing the speed and predictability of local adaptation in invasive California poppies (*Eschscholzia californica*). *Journal of Evolutionary Biology* **20**: 1090-1103.

Lenth RV. 2018. *Emmeans: estimated marginal means, aka least-squares means* (R package version 1.8.2).

Levitt J, Lovett JV. 1985. Alkaloids, antagonisms and allelopathy. *Biological Agriculture & Horticulture* **2**: 289-301.

Lewis KC, Bazzaz FA, Liao Q *et al*. 2006. Geographic patterns of herbivory and resource allocation to defense, growth, and reproduction in an invasive biennial, *Alliaria petiolata*. *Oecologia* **148**: 384-395.

Li YP, Feng YL, Li WT *et al*. 2022. Leaf trait association in relation to herbivore defense, drought resistance, and economics in a tropical invasive plant. *American Journal of Botany* **109**: 910-921.

Liao ZY, Zheng YL, Lei YB *et al*. 2014. Evolutionary increases in defense during a biological invasion. *Oecologia* **174**: 1205-1214.

Lin T, Doorduin L, Temme A *et al.* **2015.** Enemies lost: parallel evolution in structural defense and tolerance to herbivory of invasive *Jacobaea vulgaris*. *Biological Invasions* **17**: 2339-2355.

Lin T, Klinkhamer PGL, Pons TL *et al.* **2019.** Evolution of increased photosynthetic capacity and its underlying traits in invasive *Jacobaea vulgaris*. *Frontiers in Plant Science* **10**: 1016.

Liu H, Stiling P. 2006. Testing the enemy release hypothesis: a review and meta-analysis. *Biological Invasions* **8**: 1535-1545.

Liu W, Zhang Y, Chen X *et al.* **2020.** Contrasting plant adaptation strategies to latitude in the native and invasive range of *Spartina alterniflora*. *New Phytologist* **226**: 623-634.

Loughnan D, Williams JL. 2019. Climate and leaf traits, not latitude, explain variation in plant–herbivore interactions across a species' range. *Journal of Ecology* **107**: 913-922.

Lowe S, Browne M, Boudjelas S *et al.* **2000.** *100 of the world's worst invasive alien species: a selection from the global invasive species database*. Auckland, New Zealand: Invasive Species Specialist Group.

Lynch M, Conery JS. 2000. The evolutionary fate and consequences of duplicate genes. *Science* **290**: 1151-1155.

Maron JL, Baer KC, Angert AL. 2014. Disentangling the drivers of context-dependent plant–animal interactions. *Journal of Ecology* **102**: 1485-1496.

Maron JL, Vilà M, Arnason J. 2004. Loss of enemy resistance among introduced populations of St. John's wort (*Hypericum perforatum*). *Ecology* **85**: 3243-3253.

Mincheva T, Barni E, Varese GC *et al.* **2014.** Litter quality, decomposition rates and saprotrophic mycoflora in *Fallopia japonica* (Houtt.) Ronse Decraene and in adjacent native grassland vegetation. *Acta Oecologica* **54**: 29-35.

Mitchell CE, Agrawal AA, Bever JD *et al.* **2006.** Biotic interactions and plant invasions. *Ecology Letters* **9**: 726-740.

Moloney KA, Holzapfel C, Tielbörger K *et al.* **2009.** Rethinking the common garden in invasion research. *Perspectives in Plant Ecology, Evolution and Systematics* **11**: 311-320.

Montti L, Ayup MM, Aragón R *et al.* **2016.** Herbivory and the success of *Ligustrum lucidum*: evidence from a comparison between native and novel ranges. *Australian Journal of Botany* **64**: 181-192.

Moran EV, Alexander JM. 2014. Evolutionary responses to global change: lessons from invasive species. *Ecology Letters* **17**: 637-649.

Moreira X, Castagneyrol B, Abdala-Roberts L *et al*. 2018. Latitudinal variation in plant chemical defences drives latitudinal patterns of leaf herbivory. *Ecography* **41**: 1124-1134.

Müller-Schärer H, Schaffner U, Steinger T. 2004. Evolution in invasive plants: implications for biological control. *Trends in Ecology & Evolution* **19**: 417-422.

Neilson EH, Goodger JQD, Woodrow IE *et al*. 2013. Plant chemical defense: at what cost? *Trends in Plant Science* **18**: 250-258.

Neuffer B, Hurka H. 1999. Colonization history and introduction dynamics of *Capsella bursa-pastoris* (Brassicaceae) in North America: isozymes and quantitative traits. *Molecular Ecology* **8**: 1667-1681.

Oduor AMO, Lankau RA, Strauss SY *et al*. 2011. Introduced *Brassica nigra* populations exhibit greater growth and herbivore resistance but less tolerance than native populations in the native range. *New Phytologist* **191**: 536-544.

Ossowski S, Schneeberger K, Lucas-Lledó JI *et al*. 2010. The rate and molecular spectrum of spontaneous mutations in *Arabidopsis thaliana*. *Science* **327**: 92-94.

Peter Constabel C, Yoshida K, Walker V. 2014. Diverse ecological roles of plant tannins: plant defense and beyond. *Recent Advances in Polyphenol Research* **4**: 115-142.

Powell KI, Chase JM, Knight TM. 2013. Invasive plants have scale-dependent effects on diversity by altering species-area relationships. *Science* **339**: 316-318.

Pyšek P, Hulme P E, Simberloff D *et al*. 2020. Scientists' warning on invasive alien species. *Biological Reviews* **95**: 1511-1534.

Qin RM, Zheng YL, Valiente-Banuet A *et al*. 2013. The evolution of increased competitive ability, innate competitive advantages, and novel biochemical weapons act in concert for a tropical invader. *New Phytologist* **197**: 979-988.

R Core Team. 2022. *R: A language and environment for statistical computing.* Vienna, Austria: R Foundation for Statistical Computing.

Reinhardt F, Herle M, Bastiansen F *et al*. 2003. *Economic impact of the spread of alien species in Germany*. Berlin, Germany: Federal Environmental Agency.

Richards CL, Bossdorf O, Muth NZ *et al*. 2006. Jack of all trades, master of some? On the role of phenotypic plasticity in plant invasions. *Ecology Letters* **9**: 981-993.

Richards CL, Schrey AW, Pigliucci M. 2012. Invasion of diverse habitats by few

Japanese knotweed genotypes is correlated with epigenetic differentiation. *Ecology Letters* **15**: 1016-1025.

Rosche C, Hensen I, Schaar A *et al*. 2019. Climate outweighs native vs. nonnative range-effects for genetics and common garden performance of a cosmopolitan weed. *Ecological Monographs* **89**: e01386.

Rotter MC, Holeski LM. 2018. A meta-analysis of the evolution of increased competitive ability hypothesis: genetic-based trait variation and herbivory resistance trade-offs. *Biological Invasions* **20**: 2647-2660.

Rotter MC, Vallejo-Marin M, Holeski LM. 2019. A test of the evolution of increased competitive ability in two invaded regions. *Evolutionary Ecology* **33**: 713-735.

Rouifed S, Piola F, Spiegelberger T. 2014. Invasion by *Fallopia* spp. in a French upland region is related to anthropogenic disturbances. *Basic and Applied Ecology* **15**: 435-443.

Rouifed S, Puijalon S, Bardon C *et al*. 2018. Comparison of defence and performance traits between one widespread clone and native populations in a major invasive plant species. *Diversity and Distributions* **24**: 297-312.

Seebens H, Bacher S, Blackburn TM *et al*. 2021. Projecting the continental accumulation of alien species through to 2050. *Global Change Biology* **27**: 970-982.

Seebens H, Blackburn TM, Dyer EE *et al*. 2017. No saturation in the accumulation of alien species worldwide. *Nature Communications* **8**: 14435.

Siemann E, Rogers WE. 2001. Genetic differences in growth of an invasive tree species. *Ecology Letters* **4**: 514-518.

Song FQ, Zhao JB, Zhang YP *et al*. 2010. Effects of the regional climate change on the plant growth trend in Xishuangbanna. *Acta Botanica Yunnanica* **32**: 547-553.

Stoll P, Gatzsch K, Rusterholz HP *et al*. 2012. Response of plant and gastropod species to knotweed invasion. *Basic and Applied Ecology* **13**: 232-240.

Strauss SY, Agrawal AA. 1999. The ecology and evolution of plant tolerance to herbivory. *Trends in Ecology & Evolution* **14**: 179-185.

Sun Y, Roderick GK. 2019. Rapid evolution of invasive traits facilitates the invasion of common ragweed, *Ambrosia artemisiifolia*. *Journal of Ecology* **107**: 2673-2687.

Tiébré MS, Saad L, Mahy G. 2008. Landscape dynamics and habitat selection by the alien invasive *Fallopia* (Polygonaceae) in Belgium. *Biodiversity and Conservation* **17**: 2357-2370.

van der Meijden E. 1996. Plant defence, an evolutionary dilemma: contrasting effects of (specialist and generalist) herbivores and natural enemies. In *Proceedings of the 9th International Symposium on Insect-plant Relationships*. Springer Netherlands, 307-310.

van Kleunen M, Bossdorf O, Dawson W. 2018. The ecology and evolution of alien plants. *Annual Review of Ecology, Evolution, and Systematics* **49**: 25-47.

Woods EC, Sultan SE. 2022. Post-introduction evolution of a rapid life-history strategy in a newly invasive plant. *Ecology* **103**: e3803.

Wright IJ, Reich PB, Westoby M *et al*. 2004. The worldwide leaf economics spectrum. *Nature* **428**: 821-827.

Xiao L, Ding J, Zhang J *et al*. 2020. Chemical responses of an invasive plant to herbivory and abiotic environments reveal a novel invasion mechanism. *Science of the Total Environment* **741**: 140452.

Yang X, Huang W, Tian B *et al*. 2014. Differences in growth and herbivory damage of native and invasive kudzu (*Peuraria montana* var. *lobata*) populations grown in the native range. *Plant Ecology* **215**: 339-346.

Yang Y, Liu M, Pan Y *et al*. 2021. Rapid evolution of latitudinal clines in growth and defence of an invasive weed. *New Phytologist* **230**: 845-856.

Yu H, Song QH, Zhang YP *et al*. 2021. Long-term variation characteristics of radiation in the tropical seasonal rainforest in Xishuangbanna, southwestern China. *Journal of Beijing Forestry University* **43**: 56-67.

Zhang S, Zhang Y, Ma K. 2016a. Latitudinal variation in herbivory: hemispheric asymmetries and the role of climatic drivers. *Journal of Ecology* **104**: 1089-1095.

Zhang YY, Parepa M, Fischer M *et al*. 2016b. Epigenetics of colonizing species? A study of Japanese knotweed in Central Europe. In: Barrett SCH, Colautti RI, Dlugosch KM, Rieseberg LH, eds. *Invasion genetics: the baker and stebbins legacy.* Chichester, UK: John Wiley & Sons, 328–340.

Supporting Information

Table S1 Geographic origins of the 128 studied *Reynoutria japonica* populations and their uses in the Shanghai (SH) and Xishuangbanna (XSBN) common gardens.

Population ID	Site	Latitude	Longitude	Garden
China (Native)				
CN01	Guangdong	23.29	114.01	SH / XSBN
CN02	Guangdong	23.64	113.84	SH / XSBN
CN03	Guangdong	23.74	113.91	SH / XSBN
CN04	Guangdong	23.75	115.27	SH / XSBN
CN05	Guangdong	23.82	115.38	SH / XSBN
CN06	Guangdong	24.44	113.25	SH / XSBN
CN07	Guangdong	24.62	113.76	SH / XSBN
CN08	Guangdong	24.71	115.82	SH / XSBN
CN09	Jiangxi	24.96	115.61	SH / XSBN
CN10	Jiangxi	25.24	115.73	SH / XSBN
CN11	Jiangxi	25.53	115.84	SH / XSBN
CN12	Jiangxi	25.80	115.94	SH / XSBN
CN13	Jiangxi	26.03	116.16	SH / XSBN
CN14	Jiangxi	26.10	114.69	SH / XSBN
CN15	Jiangxi	26.27	116.32	SH / XSBN
CN16	Jiangxi	26.55	116.27	SH / XSBN
CN17	Jiangxi	26.85	116.37	SH / XSBN
CN18	Jiangxi	27.08	116.34	SH / XSBN
CN19	Jiangxi	27.38	116.45	SH / XSBN
CN20	Jiangxi	27.66	116.68	SH / XSBN
CN21	Jiangxi	27.87	116.78	SH / XSBN
CN22	Jiangxi	28.19	116.76	SH / XSBN
CN23	Jiangxi	28.54	117.05	SH / XSBN
CN24	Jiangxi	28.71	117.02	SH / XSBN
CN25	Jiangxi	29.10	117.12	SH / XSBN
CN26	Jiangxi	29.21	117.15	SH / XSBN
CN27	Jiangxi	29.31	117.85	SH / XSBN
CN28	Jiangxi	29.43	117.16	SH / XSBN

CN29	Jiangxi	29.64	117.22	SH / XSBN
CN30	Anhui	29.81	117.05	SH / XSBN
CN31	Anhui	29.98	116.97	SH / XSBN
CN32	Anhui	30.18	117.04	SH / XSBN
CN33	Anhui	30.44	116.23	SH / XSBN
CN34	Anhui	30.69	115.98	SH / XSBN
CN35	Anhui	30.82	116.05	SH / XSBN
CN36	Anhui	30.92	116.21	SH / XSBN
CN37	Anhui	31.09	116.10	SH / XSBN
CN38	Anhui	31.21	116.02	SH / XSBN
CN39	Anhui	31.40	115.93	SH / XSBN
CN40	Anhui	31.58	115.97	SH / XSBN
CN41	Henan	31.81	115.85	SH / XSBN
CN42	Jiangsu	32.85	118.44	SH / XSBN
CN43	Henan	34.27	115.68	SH / XSBN
CN44	Jiangsu	34.72	116.79	SH / XSBN
CN45	Shandong	35.06	116.31	SH / XSBN
CN46	Shandong	35.52	117.82	SH / XSBN
CN47	Shandong	35.74	117.64	SH / XSBN
CN48	Shandong	36.00	117.65	SH / XSBN
CN49	Shandong	36.38	117.66	SH / XSBN
CN50	Shandong	36.87	117.68	SH / XSBN
Japan (Native)				
JA01	Unzen	32.81	129.92	SH / XSBN
JA02	Saikai	32.99	129.77	SH / XSBN
JA03	Sasebo	33.21	129.68	SH / XSBN
JA04	Hirado	33.32	129.60	SH / XSBN
JA05	Hirado	33.33	129.53	SH / XSBN
North America (Introduced)				
NA01	Georgia	34.24	-83.46	SH / XSBN
NA02	NorthCarolina	35.10	-83.10	SH / XSBN
NA03	NorthCarolina	35.74	-82.68	SH / XSBN

NA04	NorthCarolina	36.21	-81.78	SH / XSBN
NA05	NorthCarolina	36.38	-81.38	SH / XSBN
NA06	Virginia	36.65	-80.92	SH / XSBN
NA07	Virginia	36.67	-80.57	SH
NA08	Virginia	37.37	-79.41	SH
NA09	Virginia	37.43	-79.16	SH
NA10	Virginia	38.46	-78.59	SH / XSBN
NA11	Virginia	38.65	-78.54	SH
NA12	Virginia	39.08	-78.09	SH / XSBN
NA13	Maryland	39.61	-76.68	SH / XSBN
NA14	NewJersey	41.01	-74.35	SH / XSBN
NA15	NewYork	41.41	-73.96	SH / XSBN
NA16	NewYork	41.77	-73.86	SH
NA17	Massachusetts	42.64	-72.91	SH / XSBN
NA18	Vermont	42.92	-72.77	SH / XSBN
NA19	Vermont	43.20	-72.45	SH / XSBN
NA20	Vermont	43.84	-72.19	SH
NA21	NewHampshire	43.94	-71.68	SH / XSBN
NA22	NewHampshire	44.12	-71.18	SH / XSBN
NA23	Maine	44.11	-70.69	SH
NA24	Maine	44.21	-70.31	SH
NA25	Maine	44.53	-69.89	SH
NA26	Maine	44.94	-68.99	SH
NA27	Maine	44.95	-68.64	SH
Europe (Introduced)				
EU01	Italy	44.67	7.29	SH / XSBN
EU02	Italy	44.75	7.48	SH / XSBN
EU03	Italy	44.88	7.69	SH / XSBN
EU04	Italy	45.19	8.04	SH / XSBN
EU05	Italy	45.64	8.37	SH / XSBN
EU06	Italy	45.80	8.87	SH / XSBN
EU07	Switzerland	46.27	9.00	SH / XSBN
EU08	Switzerland	46.51	8.7	SH / XSBN

EU09	Switzerland	47.16	8.56	SH / XSBN
EU10	Germany	47.62	8.22	SH / XSBN
EU11	Germany	47.81	7.61	SH / XSBN
EU12	Germany	47.94	7.88	SH / XSBN
EU13	Germany	48.28	8.11	SH / XSBN
EU14	Germany	48.37	8.56	SH / XSBN
EU15	Germany	48.47	8.92	SH / XSBN
EU16	Germany	48.56	8.39	SH / XSBN
EU17	Germany	48.79	8.32	SH / XSBN
EU18	Germany	49.59	8.73	SH / XSBN
EU19	Germany	49.90	8.83	SH / XSBN
EU20	Germany	50.07	8.48	SH / XSBN
EU21	Germany	50.31	7.79	SH / XSBN
EU22	Germany	50.54	7.08	SH / XSBN
EU23	Germany	51.14	6.78	SH / XSBN
EU24	Germany	51.43	7.29	SH / XSBN
EU25	Germany	51.87	7.55	SH / XSBN
EU26	Germany	52.15	7.62	SH / XSBN
EU27	Germany	52.39	7.94	SH / XSBN
EU28	Germany	52.72	8.26	SH / XSBN
EU29	Germany	53.01	8.70	SH / XSBN
EU30	Germany	53.22	9.57	SH / XSBN
EU31	Germany	53.45	10.08	SH / XSBN
EU32	Germany	53.99	10.25	SH / XSBN
EU33	Germany	54.31	10.13	SH / XSBN
EU34	Denmark	54.71	11.45	SH / XSBN
EU35	Denmark	55.46	12.19	SH / XSBN
EU36	Sweden	55.68	13.19	SH / XSBN
EU37	Sweden	56.14	13.76	SH / XSBN
EU38	Sweden	56.45	13.60	SH / XSBN
EU39	Sweden	56.83	13.96	SH / XSBN
EU40	Sweden	57.70	14.11	SH / XSBN
EU41	Sweden	58.17	14.58	SH / XSBN

EU42	Sweden	58.41	15.65	SH / XSBN
EU43	Sweden	58.67	16.20	SH / XSBN
EU44	Sweden	58.89	17.56	SH / XSBN
EU45	Sweden	59.32	18.02	SH / XSBN
EU46	Sweden	59.95	17.71	SH / XSBN

Table S2 Summary of PCA of eight climate variables. The first and second components explained 54% and 21% of climatic variation among native and introduced ranges' localities, respectively. Climate variables with stronger loadings (> 0.40) on climate PCs are shown in bold.

	PC1 climate	PC2 climate
Eigenvalues	4.378	1.731
% variance	54.7 %	21.6 %
Loadings		
Mean annual temperature (bio1)	**-0.461**	0.091
Maximum temperature of warmest month (bio4)	-0.063	**-0.728**
Minimum temperature of coldest month (bio5)	**-0.404**	-0.265
Temperature seasonality (bio6)	-0.323	**0.515**
Mean annual precipitation (bio12)	-0.360	0.188
Precipitation of the wettest month (bio13)	**-0.458**	0.002
Precipitation of the driest month (bio14)	0.170	0.176
Precipitation seasonality (bio15)	-0.385	-0.245

Fig. S1 Principal component analysis (PCA) for climate variables at sample collecting sites of *R. japonica* in common gardens. Colored dots indicate sample collecting sites. Percentages on the X- and Y-axes indicate the variation explained by each principal component.