1 **TITLE: A broad antibody class engages the influenza virus hemagglutinin head at its stem**
2 **interface**
3

4 **AUTHORS:**
5 Holly C. Simmons[1,2,a], Joel Finney[3,a,b], Ryutaro Kotaki[4], Yu Adachi[4], Annie Park Moseman[3], Akiko
6 Watanabe[3], Shengli Song[5], Lindsey R. Robinson-McCarthy[1,2], Valerie Le Sage[1,2], Masayuki Kuraoka[3],
7 E. Ashley Moseman[3], Garnett Kelsoe[3] , Yoshimasa Takahashi[4] and Kevin R. McCarthy[1,2,*]
8
AFFILIATIONS:
[1]Center for Vaccine Research, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA.
[2]Department of Microbiology and Molecular Genetics, University of Pittsburgh School of Medicine,
Pittsburgh, PA, USA.
[3]Department of Integrative Immunobiology, Duke University, Durham, North Carolina, USA
[4] Department of Immunology, National Institute of Infectious Diseases, Tokyo 162-8640, Japan
[5] Department of Surgery, Duke University, Durham, North Carolina 27710, USA
[a] Contributed equally
[b] Present address: Laboratory of Molecular Medicine, Boston Children's Hospital, Boston,
Massachusetts, USA
[*]krm@pitt.edu

Abstract:
Influenza infection and vaccination impart strain-specific immunity that fails to protect against both
seasonal antigenic variants and the next pandemic. However, antibodies directed to conserved sites
can confer broad protection. We identify and characterize a class of human antibodies that engage a
previously undescribed, conserved, epitope on the influenza hemagglutinin protein (HA). Prototype
antibody S8V1-157 binds at the normally occluded interface between the HA head and stem.
Antibodies to this HA head-stem interface epitope are non-neutralizing *in vitro* but protect against lethal
infection in mice. Their breadth of binding extends across most influenza A serotypes and seasonal
human variants. Antibodies to the head-stem interface epitope are present at low frequency in the
memory B cell populations of multiple donors. The immunogenicity of the epitope warrants its
consideration for inclusion in improved or "universal" influenza vaccines.

32
INTRODUCTION:
Influenza pandemics arise from antigenically novel animal influenza A viruses transmitted to humans
from zoonotic hosts. Historically, pandemic viruses have generally become endemic and have
continued to circulate as seasonal viruses. Sustained viral circulation comes from on-going antigenic
evolution that leads to escape from population-level immunity elicited by previous exposures [1].
Immunity elicited by infection, vaccination, or both is insufficient to confer enduring immunity against
seasonal variants or future pandemic viruses. Although antibodies provide the strongest protection
against infection, they also drive the antigenic evolution of the viral surface proteins [1]. Broadly
protective monoclonal antibodies that engage conserved sites have been isolated from human donors
[2]. Passive transfer of these antibodies to animal models imparts broadly protective immunity. Selective
elicitation of similar antibodies by a next generation influenza vaccine is predicted to confer protection
less susceptible to escape than protection from current vaccines [3-5].
45
The influenza hemagglutinin (HA) protein is the major target of protective antibodies [6]. HA facilitates
cell entry by attaching to cells via an interaction with its receptor, sialic acid, and by acting as a virus-
cell membrane fusogen. HA is synthesized as a polyprotein, HA0, which forms homotrimers that are
incapable of undergoing the full series of conformational rearrangements required for membrane fusion.
A fusion-competent trimer is produced by cleavage of HA0 into HA1 and HA2 domains by cellular
proteases often resident on the target cell [7-9]. HA1 includes the globular HA head domain that contains
the receptor binding site (RBS), while HA2 contains the helical stem regions that undergo a cascade of
rearrangements that ultimately fuse viral and cellular membranes. The requirement for HA to transit
through multiple conformations during fusion likely accounts for the intrinsic propensity of HA0 or HA1-

HA2 to transiently explore states that deviate from the stable prefusion form defined by static structures
determined by X-ray crystallography [10-13].
57
Large genetic and antigenic differences separate influenza A HA proteins. They are classified into two
groups comprising 16 sialic acid binding serotypes: group 1 (H1, H2, H5, H6, H8, H9, H11, H12, H13,
H16) and group 2 (H3, H4, H7, H10, H14, H15). Using a similar convention for a second glycoprotein,
neuraminidase (NA), influenza viruses are named by their HA and NA content (e.g., H2N2, H10N8).
Currently, divergent H1N1 and H3N2 viruses circulate as seasonal human influenza viruses. Humans
can produce antibodies that engage surfaces conserved between serotypes and thereby provide cross-
serotype protection. These epitopic regions include the HA RBS, stem, anchor, head interface, the so-
called "long alpha helix", and the HA2 beta-hairpin (PDB 8UDG reported in Finney et al, PNAS in
press) [2,14-17]
67
Here, we report a class of human antibodies directed to a previously unreported, widely conserved, site
at the base, or neck, of the HA head domain. In the stable prefusion structure of the HA trimer, the
epitope for prototype antibody S8V1-157 is at the interface of the head and stem, occluded in the
"resting state" prefusion conformation. Biochemical, cellular, and *in vivo* passive transfer experiments
indicate that this HA head-stem interface epitope is sufficiently exposed to allow antibody binding and
to confer strong protection against lethal influenza virus infection in murine challenge models. Many
humans harbor antibodies directed to this epitope; some of these antibodies bind HAs from divergent
influenza serotypes, groups, and >50 years of human seasonal virus antigenic variation. Immunogens
that elicit such antibodies might be included in influenza vaccines intended to confer broader protection.
77
RESULTS:
A class of antibodies engages an epitope at head-stem epitope of the HA head domain.
By culturing individual human memory B (Bmem) cells and then screening the culture supernatants
containing secreted IgGs, we identified (as reported previously [18]) 449 HA-reactive antibodies from four
donors (S1, S5, S8 and S9). These antibodies represent the Bmem cells circulating in the blood at the
time the donors were immunized with the TIV2015-2016 seasonal influenza vaccine (visit 1; V1), or 7
days later (visit 2; V2). From donors S1 and S8, we found four IgGs (S1V2-17, S1V2-60, S1V2-65 and
S8V1-157 that had a novel pattern of HA reactivity (Figure 1A). All bound seasonal H1s and H3s and
an H5 HA; they also bound recombinant HA constructs comprising only the HA head domain, indicating
the IgGs did not target stem epitopes (Figure 1A). These four antibodies competed with each other for
HA-binding, implying overlapping epitopes, but failed to compete with antibodies directed to conserved,
broadly protective epitopes in the HA head and stem (Figure 1B).
90
91 All four antibodies had light chains encoded by IGKV2-28*01 or IGKV2D-28*01 (Figure 1C), which have
92 identical germline coding sequences. The heavy chains are encoded by either IGHV1-24*01 or IGHV5-
93 51*01 and have short (11-13 amino acid) third complementarity-determining regions (HCDR3s). S1V2-
94 60 and S1V2-65 are clonally related, while S1V2-17 and S8V1-157 share a common IGHV1-24*01-
95 IGKV2-28*01 pairing, despite their derivation from different donors.
96
We determined the structure of S8V1-157 complexed with a monomeric HA head construct. Crystals
were only obtained using an A/American black duck/New Brunswick/00464/2010(H4N6) (H4-NB-2010)
HA (Figure 2A and Table S1). S8V1-157 engages an epitope at the base of the head, just where it
faces the stem [19]. Hence this HA1 epitope is occluded by the association of HA1 with HA2 (Figure 2A).
S8V1-157 contacts residues mediating the intra-protomer interaction, which are conserved across
divergent HAs (Figure 2B). Evolutionary constraints imposed by the requirement of stable association,
at neutral pH, of HA1 and HA2 likely account for the conservation of the head-stem interface epitope
and its respective breadth of binding of these antibodies. Antibody or B cell receptor engagement of the
HA this epitope, in the context of a prefusion HA molecule, would require displacement of the HA head
from the HA2 helical stem.
107

The compact HCDR3 of S8V1-157 packs against the heavy chain CDR and framework regions (FR),
HCDR1-FR1 and FR2-HCDR2, to produce a cleft between the heavy and light chains that
accommodates the head-stem interface epitope (Figure 2C). Residues N33, Y35 and Y37 within the
elongated LCDR1 make extensive contacts with a conserved HA surface. The δ-amide of N33 donates
and receives hydrogen bonds from the main chain of HA residue HA-C305. Its main chain amide also
donates a hydrogen bond to the carboxyl group of HA-D304. Y35 donates a hydrogen bond to the main
chain carbonyl of HA-T291 and participates in van der Walls interactions by stacking upon HA-P293.
Y37 receives a hydrogen bond from the main chain residue HA-K307, donates a hydrogen bond to HA-
P293, participates in van der Walls contacts with the aryl group of HA-F294 and the aliphatic side chain
of HA-K307.
118
The structure of the S8V1-157-H4-NB-2010 complex provides a rationale for the genetic features
common to this antibody class. The LCDR1 NXYXY motif is germline-encoded by a small subset of
IGKV-genes. Of these, only IGKV2-28/IGKV2D-28a also encode a hydrophobic residue at position 55.
L55 participates in van der Walls contacts with HA-P293 (Figure 2C) and contributes to a hydrophobic
surface on S8V1-157 that complements a hydrophobic patch within its epitope on the HA head (Figure
S1). Polar residues present at position 55 in otherwise similar light chains are predicted to be less
favorable in this local environment. Similar sequence features are not present in the IGLV locus.
Additionally, the short HCDR3 (common among these antibodies) creates shallow cavity that
accommodates the hydrophobic surfaces that would normally pack against HA2 in the prefusion HA
trimer.
129
Antibodies to the HA head-stem interface epitope are broadly binding.
We determined the binding of these antibodies to divergent HA serotypes by enzyme-linked
immunosorbent assay (ELISA) (Figures 3A and S2). The HA head-stem epitope antibodies had very
similar breadth: each bound all seasonal H1s (1977-2019) and H3s (1968-2020) assayed, and also
bound several other HA subtypes within groups 1 and 2. Typically, if an HA was bound by one head-
stem epitope antibody it was bound by the remaining three (Figure 3). Affinities and breadth of binding
were generally higher for group 2 HAs but extended to group 1 HAs, including seasonal H1, pandemic
H2, and pre-pandemic H5 HAs. Overall, these antibodies engaged HAs from 11 of the 16 non-bat
influenza A HA serotypes. Failure to engage specific HAs was not due to epitope inaccessibility, since
these antibodies also did not bind matched, soluble, HA head domains that present the HA head-stem
epitope without steric hindrance (Figures S3 and S4). A truncated HA head domain, lacking the S8V1-
157 epitope, was not bound by the four antibodies. All four are likely to engage a common, discrete,
epitope comprising the terminal boundaries of the HA head domain.
143
Using a flow cytometry-based assay, we demonstrated that HA head-stem epitope antibodies engage
divergent HAs present on the cell surface (Figure 4). HAs were efficiently bound by these antibodies in
a pattern consistent with our ELISA data. The intensity of labeling by HA head-stem epitope antibodies
was generally lower than for labeling by control antibodies directed to solvent-exposed epitopes on the
HA head, but comparable to labeling by the HA head-interface antibody S5V2-29 [18]. Together, these
observations suggest that HA interface epitopes are not as occupied by antibody as are fully exposed
epitopes. Epitope accessibility and/or differences in the number of IgG molecules bound per HA may
account for these differences. To determine whether S8V1-157 could engage HA0 present on the
surface of a cell, cells were transiently transfected with an A/Aichi/02/1968(H3N2)(X-31) HA expression
vector and assessed for antibody binding using flow cytometry, and HA processing by western blotting
cell lysates for its endogenous HA-tag in HA1 (Figure S4 A and B). HA0 was not processed in these
cells but was labeled by S8V1-157. The HA head-stem epitope is therefore exposed in biochemical and
cellular contexts.
157
Antibodies to the HA head-stem epitope protect against lethal influenza virus infection.
S8V1-157 failed to neutralize influenza virus *in vitro* (Figure S5). To determine if these antibodies
confer protection *in vivo* we performed a series of prophylactic studies in mice. In a pilot study passive
transfer of S8V1-157 antibody conferred moderate protection (Figure S6B). We expanded our studies

to include additional control antibodies and to directly assess the impact of antibody IgG subtype on
protection. Notably, unlike humans, in mice the antibody IgG2c isotype is capable of directing protective
Fc-dependent antibody effector functions, including antibody-dependent-cellular cytotoxicity (ADCC)
and complement deposition (ADCD). In these experiments we passively transferred 150μg of
musinized IgG1 (which has minimal Fc-effector activity) or IgG2c versions of S8V1-157, S5V2-65
(which competes for HA binding with S8V1-157), HC19[20] (potently neutralizing
A/Aichi/02/1968(H3N2)(X-31) specific antibody), S5V2-29[18] (an HA-head interface antibody that
protects against lethal infection and is enhanced by Fc-dependent mechanisms) and CR3022[21] (SARS-
CoV antibody) to mice prior to challenge with a lethal dose of A/Aichi/02/1968(H3N2)(X-31) (Figure S6).
The 150μg (~7.5 mg/kg) does is consistent with human monoclonal antibody therapeutics.
172
HC19 protected mice from any measurable, infection-induced weight loss, while the irrelevant antibody
CR3022 offered no protection against weight loss, requiring recipient mice to be ethically euthanized by
day 10 post-infection (Figure 5A). Animals administered antibodies to the HA head-stem or head-
interface epitope experienced mild weight loss, but nearly all recovered within 10 days post-infection.
IgG1 versions of the same antibodies offered slightly less protection than the corresponding IgG2cs. Fc
effector functions thus had a role in immune control of the infections (Figure 5B-D). Across our studies,
HA head-stem epitope antibodies did not potently inhibit influenza virus infection but did confer strong
protection against severe disease and mortality.
181
The HA head-stem epitope is immunogenic in humans.
We screened additional human Bmem cell cultures for S8V1-157-competing antibodies (Figure 6). The
samples were taken from seven donors, including the same subjects as before (S1, S5, S8, S9), but
vaccinated and sampled during subsequent flu seasons (see Materials and Methods for details);
subject S12, who was immunized and sampled at the same times; and subjects KEL01 and KEL03,
who were sampled after receiving TIV in 2014-2015. From 528 clonal cultures that produced HA-
binding IgG, we identified eight additional supernatants that inhibited S8V1-157 binding by >90%
(Figure 6A). In a screen of antibody reactivity, five of the eight competing antibodies reacted with group
1 and group 2 HAs.
191
Paired heavy and light chain sequences were recovered from seven of the eight S8V1-157-competing
antibodies (Figure S7). Two antibodies, S1V4-P4-C5 and S12V6-P7-H4, use an IGKV2-28 light chain
paired with IGHV1-24*01, like S8V1-157 and S1V2-17. All four antibodies have short, 11-12 amino acid
HCDR3s. These features, shared with S8V1-157, likely define a public antibody class present in at
least three human donors. The other newly identified S8V1-157 competitors have varied usage of Vh
and Vk genes and HCDR3 lengths (10-19 amino acids). These antibodies could either have footprints
that overlap the head-stem interface epitope or that prevent its exposure.
199
DISCUSSION:
We identified and characterized a human antibody response directed to a previously unrecognized,
widely conserved, cryptic epitope on the influenza HA head domain. These antibodies bind broadly to
divergent HA serotypes found in animals and seasonal antigenic variants of human viruses.
Prophylactic passive transfer of these antibodies to mice conferred protection against lethal influenza
virus disease. Overall, across our studies we find HA head-stem epitope antibodies comprise ~1% of
the circulating, HA-reactive Bmem cell repertoire. Cells with HA head-stem epitope BCRs are present in
multiple donors both before and after vaccination against seasonal influenza. These cells have a bias,
but not a restriction, for IGKV2-28 usage. Across the 12 known examples we find no additional
constraints. Humans are therefore likely to produce similar antibodies and have the capacity to mount
polyclonal, poly-epitope, broadly protective antibody responses.
211
212 HA head-stem epitope antibodies confer robust protection against lethal influenza challenge,
213 demonstrating that the HA head-stem epitope must be exposed on infected cells and/or virions.
214 Structural, biophysical, and computational approaches indicate that HA trimers transiently adopt

conformations that expose epitopes normally occluded in its defined prefusion form[10-13]. Such transient fluctuations might explain how B cells and antibodies recognize ordinarily occluded epitopes[16-18,22-24]. A non-mutually exclusive possibility is that head-stem epitope antibodies directly recognize postfusion HA on the surface of infected cells, as has been proposed for some HA stem antibodies, including LAH31[17] and m836[24] antibodies, whose conformational epitope is present only in the postfusion structure. It is plausible that HA0 is cleaved into HA1/HA2 by proteases and a requisite fraction accumulates in a post-fusion form that would expose this epitope. Nonetheless, in our biochemical assays, and staining of transiently transfected cells, HA head-stem epitope antibodies avidly bound HA0, which cannot fully transition to its stable post fusion structure[8].

Humans and mice mount antibody responses directed to seemingly difficult to engage HA epitopes [14-18,22-26]. Prevalence of such antibodies in immune repertoires indicates that these epitopes are immunogenic. In this instance, human head-stem interface antibodies typically have ~4-10% V gene mutation frequencies indicating that this epitope is presented, in sufficient abundance, over recurrent influenza exposures. Exposure of the HA head-stem epitope requires large-scale displacement of the HA head that are likely accompanied by other rearrangements of HA protomers. The immunogenicity of the HA head-stem epitope poses a critical question for vaccinology: how does the antigen presented to a B cell in a germinal center reaction relate to its form on infectious virions or to what has been defined in the laboratory in considerable biochemical and structural detail? Antibodies directed to interface/buried epitopes on other unrelated viral glycoproteins demonstrate that our lack of understanding is not specific to HA [27].

The most broadly protective human antibody classes are directed to conserved epitopes with limited antibody accessibility. Adjacent protomers and/or membranes hinder access to stem/anchor epitopes and interface epitopes are normally occluded [2,14,15,28]. Nevertheless, antibodies recognizing these epitopes protect against lethal influenza virus disease in small animal models. These antibodies are typically less potently neutralizing than antibodies directed to unhindered sites and their protection is often enhanced by antibody Fc-mediated effector functions [16-18,23,26,29,30]. Their resistance to human influenza antigenic evolution and ability to engage emerging, pre-pandemic, and other animal viruses make their selective elicitation a potential strategy to improve current influenza vaccines. Given the promise of these antibodies, a rigorous understanding of their potential to enhance human protective immunity and/or ameliorate disease is needed.

METHODS:

Human subjects.

Peripheral blood mononuclear cells (PBMCs) were obtained from human donors KEL01 (male, age 39) and KEL03 (female, age 39), under Duke Institutional Review Board committee guidelines. Written informed consent was obtained from all three subjects. KEL01 and KEL03 received the trivalent inactivated seasonal influenza vaccine (TIV) 2014-2015 Fluvirin, which contained A/Christchurch/16/2010, NIB-74 (H1N1), A/Texas/50/2012, NYMC X-223 (H3N2), and B/Massachusetts/2/2012, NYMC BX-51B. Blood was drawn on day 14 post-vaccination, and PBMCs isolated by centrifugation over Ficoll density gradients (SepMate-50 tubes, StemCell Tech) were frozen and kept in liquid nitrogen until use.

PBMCs were also obtained from human donors S1 (female, age 51-55), S5 (male, age 21-25), S8 (female, age 26-30), S9 (female, age 51-55), and S12, under Boston University Institutional Review Board committee guidelines. Written informed consent was obtained from all five subjects. Donors met all of the following inclusion criteria: between 18 and 65 years of age; in good health, as determined by vital signs [heart rate (<100 bpm), blood pressure (systolic ≤ 140 mm Hg and ≥ 90 mm Hg, diastolic ≤ 90 mm Hg), oral temperature (<100.0 °F)] and medical history to ensure existing medical diagnoses/conditions are not clinically significant; can understand and comply with study procedures, and; provided written informed consent prior to initiation of the study. Exclusion criteria included: 1) life-threating allergies, including an allergy to eggs; 2) have ever had a severe reaction after influenza vaccination; 3) a history of Guillain-Barre Syndrome; 4) a history of receiving immunoglobulin or other blood product within the 3 months prior to vaccination in this study; 5) received an experimental agent (vaccine, drug, biologic, device, blood product, or medication) within 1 month prior to vaccination in this study or expect to receive an experimental agent during this study; 6) have received any live licensed vaccines within 4 weeks or inactivated licensed vaccines within 2 weeks prior to the vaccination in this study or plan receipt of such vaccines within 2 weeks following the vaccination; 7) have an acute or chronic medical condition that might render vaccination unsafe, or interfere with the evaluation of humoral responses (includes, but is not limited to, known cardiac disease, chronic liver disease, significant renal disease, unstable or progressive neurological disorders, diabetes mellitus, autoimmune disorders and transplant recipients); 8) have an acute illness, including an oral temperature greater than 99.9°F, within 1 week of vaccination; 9) active HIV, hepatitis B, or hepatitis C infection; 10) a history of alcohol or drug abuse in the last 5 years; 11) a history of a coagulation disorder or receiving medications that affect coagulation. Subjects S1, S5, S8, S9, and S12 received seasonal influenza vaccination during three consecutive North American flu seasons (2015-2016, 2016-2017, 2017-2018), and had blood drawn on day 0 (pre-vaccination; visits 1, 3, and 5) and day 7 (post-vaccination, visits 2, 4, and 6) each year. During the 2015-2016 season (visits 1 and 2), the subjects received the TIV Fluvirin, which contained A/reassortant/NYMC X-181 (California/07/2009 x NYMC X-157) (H1N1), A/South Australia/55/2014 IVR-175 (H3N2), and B/Phuket/3073/2013. During the 2016-2017 season (visits 3 and 4), the subjects received the quadrivalent inactivated vaccine Flucelvax, containing A/Brisbane/10/2010 (H1N1), A/Hong Kong /4801/2014 (H3N2), B/Utah/9/2014, and B/Hong Kong/259/2010. During the 2017-2018 season (visits 5 and 6), the subjects received the quadrivalent inactivated vaccine Flucelvax, containing A/Singapore/GP1908/2015 IVR-180 (H1N1), A/Singapore/GP2050/2015 (H3N2), B/Utah/9/2014, and B/Hong Kong/259/2010.

Cell lines.

Human 293F cells were maintained at 37°C with 5-8% CO_2 in FreeStyle 293 Expression Medium (ThermoFisher) supplemented with penicillin and streptomycin. HA-expressing K530 cell lines [31] (*Homo sapiens*) were cultured at 37°C with 5% CO_2 in RPMI-1640 medium plus 10% FBS (Cytiva), 2-mercaptoethanol (55 μM; Gibco), penicillin, streptomycin, HEPES (10 mM; Gibco), sodium pyruvate (1 mM; Gibco), and MEM nonessential amino acids (Gibco). Madin-Darby canine kidney (MDCK) were maintained in Minimum Essential medium supplemented with 10% fetal bovine serum, 5 mM L-glutamine and 5 mM penicillin/streptomycin.

Recombinant Fab expression and purification.

The heavy and light chain variable domain genes for Fabs were cloned into a modified pVRC8400
expression vector, as previously described[32-34]. Fab fragments used in crystallization were produced
with a noncleavable 6xhistidine (6xHis) tag on the heavy chain C-terminus. Fab fragments were
produced by polyethylenimine (PEI) facilitated, transient transfection of 293F cells. Transfection
complexes were prepared in Opti-MEM (Gibco) and added to cells. 5 days post transfection, cell
supernatants were harvested and clarified by low-speed centrifugation. Fabs were purified by passage
over TALON Metal Affinity Resin (Takara) followed by gel filtration chromatography on Superdex 200
(GE Healthcare) in 10 mM tris(hydroxymethyl)aminomethane (tris), 150 mM NaCl at pH 7.5 (buffer A).
310
Single B cell Nojima cultures.
Nojima cultures were previously performed[18] and in Finney et al., in press at PNAS. Briefly, peripheral
blood mononuclear cells (PBMCs) were obtained from four human subjects S1 (female, age range 51-
55), S5 (male, age 21-25), S8 (female, age 26-30), and S9 (female, age 51-55). Single human Bmem
cells were directly sorted into each well of 96-well plates and cultured with MS40L-low feeder cells in
RPMI1640 (Invitrogen) containing 10% HyClone FBS (Thermo scientific), 2-mercaptoethanol (55 μM),
penicillin (100 units/ml), streptomycin (100 μg/ml), HEPES (10 mM), sodium pyruvate (1 mM), and
MEM nonessential amino acid (1X; all Invitrogen). Exogenous recombinant human IL-2 (50 ng/ml), IL-4
(10 ng/ml), IL-21 (10 ng/ml) and BAFF (10 ng/ml; all Peprotech) were added to cultures. Cultures were
maintained at 37° degrees Celsius with 5% CO_2. Half of the culture medium was replaced twice weekly
with fresh medium (with fresh cytokines). Rearranged *V(D)J* gene sequences for human Bmem cells
from single-cell cultures were obtained as described[18,32,35]. Specificity of clonal IgG antibodies in culture
supernatants and of rIgG antibodies were determined in a multiplex bead Luminex assay (Luminex
Corp.). Culture supernatants and rIgGs were serially diluted in 1 × PBS containing 1% BSA, 0.05%
NaN_3 and 0.05% Tween20 (assay buffer) with 1% milk and incubated for 2 hours at room temperature
with the mixture of antigen-coupled microsphere beads in 96-well filter bottom plates (Millipore). After
washing three times with assay buffer, beads were incubated for 1 hour at room temperature with
Phycoerythrin-conjugated goat anti-human IgG antibodies (Southern Biotech). After three washes, the
beads were re-suspended in assay buffer and the plates read on a Bio-Plex 3D Suspension Array
System (Bio-Rad).
331
Recombinant HA expression and purification.
Recombinant HA head domain constructs and Full-length HA ectodomains (FLsE) were expressed by
polyethylenimine (PEI) facilitated, transient transfection of 293F cells. To clone HA head domains,
synthetic DNA for the domain was subcloned into a pVRC8400 vector encoding a C-terminal rhinovirus
3C protease site and a 6xHis tag. To produce FLsE constructs, synthetic DNA was subcloned into a
pVRC8400 vector encoding a T4 fibritin (foldon) trimerization tag and a 6xHis tag. Transfection
complexes were prepared in Opti-MEM (Gibco) and added to cells. 5 days post transfection, cell
supernatants were harvested and clarified by low-speed centrifugation. HA was purified by passage
over TALON Metal Affinity Resin (Takara) followed by gel filtration chromatography on Superdex 200
(GE Healthcare) in buffer A. HA head domains used for crystallography underwent the following
additional purification steps. HA heads were cleaved using the Pierce 3C HRV Protease Solution Kit
(Ref 88947) and passed over TALON Metal Affinity Resin to capture cleaved tags. Cleaved HA heads
were then further purified by gel filtration chromatography on Superdex 200 (GE Healthcare) in buffer
A.
346
ELISA
Five hundred nanograms of rHA FLsE or HA head domain were adhered to high-capacity binding, 96
well-plates (Corning 9018) overnight in PBS pH 7.4 at 4°C. HA coated plates were washed with a PBS-
Tween-20 (0.05%v/v) buffer (PBS-T) and then blocked with PBS-T containing 2% bovine serum
albumin (BSA) for 1 hour at room temperature. Blocking solution was then removed, and 5-fold dilutions
of IgGs (in blocking solution) were added to wells. Plates were then incubated for 1 hour at room
temperature. Primary IgG solution was removed and plates were washed three times with PBS-T.
Secondary antibody, anti-human IgG-HRP (Abcam ab97225) diluted 1:10,000 in blocking solution, was
added to wells and incubated for 30 minutes at room temperature. Plates were then washed three

times with PBS-T. Plates were developed using 150µl 1-Step ABTS substrate (ThermoFisher,
Prod#37615). Following a brief incubation at room temperature, HRP reactions were stopped by the
addition of 100µl of 1% sodium dodecyl sulfate (SDS) solution. Plates were read on a Molecular
Devices SpectraMax 340PC384 Microplate Reader at 405 nm. KD values for ELISA were obtained as
follows. All measurements were performed in technical triplicate. The average background signal (no
primary antibody) was subtracted from all absorbance values. Values from multiple plates were
normalized to the FI6v3[30] standard (FluA20 was used for the HA head assays) that was present on
each ELISA plate. The average of the three measurements were then graphed using GraphPad Prism
(v9.0). KD values were determined by applying a nonlinear fit (One site binding, hyperbola) to these
data points. The constraint that Bmax must be greater than 0.1 absorbance units was applied to all KD
analysis parameters.
367
Recombinant IgG expression and purification.
The heavy and light chain variable domains of selected antibodies were cloned into modified
pVRC8400 expression vectors to produce full length human IgG1 heavy chains and human lambda or
kappa light chains. IgGs were produced by transient transfection of 293F cells as specified above. Five
days post-transfection supernatants were harvested, clarified by low-speed centrifugation, and
incubated overnight with Protein A Agarose Resin (GoldBio) at 4°C. The resin was collected in a
chromatography column and washed with one column volume of buffer A. IgGs were eluted in 0.1M
Glycine (pH 2.5) which was immediately neutralized by 1M tris (pH 8.5). Antibodies were then dialyzed
against phosphate buffered saline (PBS) pH 7.4.
377
Recombinant mIgG expression and endotoxin free purification.
The heavy and light chain variable domains of selected antibodies were cloned into respective modified
pVRC8400 expression vector to produce full length murine IgG1 and IgG2C heavy chains and murine
lambda or kappa light chains. IgGs were produced by transient transfection of 293F cells as specified
above. Five days post-transfection supernatants were harvested, clarified by low-speed centrifugation.
20% volume of 1M 2-morpholin-4-ylethanesulfonic acid (MES) pH 5 and Immobilized Protein G resin
(Thermo Scientific Prod#20397) were added to supernatant and sample was incubated overnight at
4°C. The resin was collected in a chromatography column and washed with one column volume of
10mM MES, 150 mM NaCl at pH 5. mIgGs were eluted in 0.1M Glycine (pH 2.5), which was
immediately neutralized by 1M tris (pH 8.5). Antibodies were then dialyzed against phosphate buffered
saline (PBS) pH 7.4. All mIgGs administered to mice were purified using buffers made with endotoxin-
free water (HyPure Cell Culture Grade Water, Cytiva SH30529.03) and were dialyzed into endotoxin-
free Dulbecco's PBS (EMD Millipore, TMS-012-A). Following dialysis, endotoxins were removed using
Pierce High Capacity Endotoxin Removal Spin Columns (Ref 88274).
392
Virus microneutralization assays.
Two-fold serial dilutions of 50 ug/mL of HC19, S8V1-157 or CR3022 were incubated with $10^{3.3}$ TCID$_{50}$ of
A/Aichi/02/1968 H3N2 (X-31) influenza virus for 1 hour at room temperature with continuous rocking.
Media with TPCK was added to 96-well plates with confluent MDCK cells before the virus:serum mixture
was added. After 4 days, CPE was determined and the neutralizing antibody titer was expressed as the
reciprocal of the highest dilution of serum required to completely neutralize the infectivity each virus on
MDCK cells. The concentration of antibody required to neutralize 100 TCID$_{50}$ of virus was calculated
based on the neutralizing titer dilution divided by the initial dilution factor, multiplied by the antibody
concentration.
402
Protection:
C57BL/6 female and male mice were obtained from the Jackson Laboratory. All mice were housed
under pathogen-free conditions at Duke University Animal Care Facility. Eight to 10-week old mice
were injected *i.p.* with 150 µg of recombinant antibody diluted to 200 µl in PBS. Three hours later, mice
were anesthetized by *i.p.* injection of ketamine (85 mg/kg) and xylazine (10 mg/kg) and infected
intranasally with 3×LD$_{50}$ (1.5×10^4 PFU) of A/Aichi/2/1968 X-31 (H3N2) in 40 µL total volume (20
µL/nostril). Mice were monitored daily for survival and body weight loss until 14 days post-challenge.

The humane endpoint was set at 20% body weight loss relative to the initial body weight at the time of
infection. All animal experiments were approved by the standards and guidance set forth by Duke
University IACUC.
413
Female C57BL/6J mice at 8 weeks old, purchased from Japan SLC, were i.p. injected with 150 µg of
the antibodies in 150 µL PBS. Three hours later, the mice were intranasally infected with $5LD_{50}$ of
A/Aichi/02/1968(H3N2)(X31), kindly gifted from Dr. Takeshi Tsubata (Tokyo Medical and Dental
University), under anesthesia with medetomidine-midazolam-butorphanol. Survival and body weight
were daily assessed for 14 days with a humane endpoint set as 25% weight loss from the initial body
weight. The experimental procedures were approved by the Animal Ethics Committee of the National
Institute of Infectious Diseases, Japan, and performed in accordance with the guidelines of the
Institutional Animal Care and Use Committee. Statistical significance of body weight change and
survival of mice after lethal influenza challenge was calculated by Two-way ANOVA test and Mantel-
Cox test, respectively, using GraphPad Prism (v10.0) software
424

Competitive inhibition assay
Competitive binding inhibition was determined by a Luminex assay, essentially as described[18,36].
Briefly, serially diluted human rIgGs or diluted (1:10) Bmem cell culture supernatants were incubated
with HA-conjugated Luminex microspheres for 2 h at room temperature or overnight at 4°C. S8V1-157
mouse IgG1 was then added at a fixed concentration (100 ng/ml final for competition with rIgGs, or 10
ng/ml final for competition with culture supernatants) to each well, and incubated with the competitor
Abs and HA-microspheres for 2 h at room temperature. After washing, bound S8V1-157 was detected
by incubating the microspheres with 2 µg/ml PE-conjugated rat anti-mouse IgG1 (SB77e,
SouthernBiotech) for 1 hr at room temperature, followed by washing and data collection. Irrelevant rIgG
or culture supernatants containing HA-nonbinding IgG were used as non-inhibiting controls.
435

Flow cytometry
Flow cytometry analysis of rIgG binding to HA-expressing K530 cell lines was performed essentially as
described [31]. Briefly, K530 cell lines were thawed from cryopreserved aliquots and expanded in culture
for ≥3 days. Pooled K530 cells were incubated at 4°C for 30 min with 0.4 µg/ml recombinant human
IgGs diluted in PBS plus 2% fetal bovine serum. After washing, cells were labeled with 2 µg/ml PE-
conjugated goat anti-human IgG (Southern Biotech) for 30 min at 4°C. Cells were then washed and
analyzed with a BD FACSymphony A5 flow cytometer. Flow cytometry data were analyzed with FlowJo
software (BD).
444

Flow cytometry analysis of rIgG binding to HA-expressing 293F cells was performed as follows. 293F
cells were transfected using PEI with either plasmid encoding full-length HA from
A/Aichi/02/1968(H3N2)(X31) or with empty vector. 36 hours post-transfection, cells were incubated at
4°C for one hour with 0.4 µg/ml recombinant human IgGs diluted in PBS plus 2% fetal bovine serum.
After washing, cells were labeled with BB515 mouse anti-human IgG (BD Biosciences) for 30 min at
4°C at the manufacturer's recommended concentration, followed by washing and fixation in 2%
paraformaldehyde. Cells were analyzed with a BD LSRFortessa flow cytometer. Flow cytometry data
were analyzed with FlowJo software (BD).
453

Western blots
293F cells were transfected using PEI with either plasmid encoding full-length HA from
A/Aichi/02/1968(H3N2)(X31) or with empty vector. 24 hours post-transfection, cells were lysed in RIPA
buffer (25mM Tris pH7.6, 150mM NaCl, 1% NP40 alternative, 1% sodium deoxycholate, 0.1% SDS).
Cell lysates were clarified of debris and boiled with Laemmli buffer with 2-mercaptoethanol. Samples
were run on 4-20% acrylamide gels and transferred to nitrocellulose membranes. Membranes were
blocked in 5% milk in PBS with 0.05% Tween 20 and probed with anti-HA tag antibody (Thermo-Fisher
A01244-100; 0.3 µg/ml), followed by IR800-conjugated goat anti-mouse immunoglobulin (LiCor 926-

32210; 0.1 μg/ml) and DyLight680-conjugated mouse anti-rabbit GAPDH antibody (BioRad
MCA4739D680; 0.3 μg/ml). Membranes were imaged using a LiCor Odyssey CLx imager.
464
Crystallization
S1V2-157 Fab fragments were co-concentrated with the HA-head domain of A/American black
duck/New Brunswick/00464/2010(H4N6) at a molar ratio of ~1:1.3 (Fab to HA-head) to a final
concentration of ~20 mg/ml. Crystals of Fab-head complexes were grown in hanging drops over a
reservoir solutions containing 0.1 M Lithium sulfate, 0.1 M Sodium chloride, 0.1 M 2-(N-
morpholino)ethanesulfonic acid (MES) pH 6.5 and 30% (v/v) poly(ethylene glycol) (PEG) 400. Crystals
were cryprotected with 30% (v/v) PEG 400, 0.12 M Lithium sulfate, 0.3 M Sodium chloride, and 0.06 M
MES pH 6.5. Cryprotectant was added directly to the drop, crystals were harvested, and flash cooled in
liquid nitrogen.
474
Structure determination and refinement
We recorded diffraction data at the Advanced Photon Source on beamline 24-ID-C. Data were
processed and scaled (XSCALE) with XDS[37]. Molecular replacement was carried out with PHASER[38],
dividing each complex into four search models (HA-head, Vh, Vl and constant domain). Search models
were 5XL2, 6EIK , 5BK5 and 6E56.We carried out refinement calculations with PHENIX[39] and model
modifications, with COOT[40]. Refinement of atomic positions and B factors was followed by translation-
liberation-screw (TLS) parameterization. All placed residues were supported by electron density maps
and subsequent rounds of refinement. Final coordinates were validated with the MolProbity server[41].
Data collection and refinement statistics are in Table S1. Figures were made with PyMOL (Schrödinger,
New York, NY.
485
486 **DATA AND SOFTWARE AVAILABILITY**:
487 Coordinates and diffraction data have been deposited at the PDB, accession number 8US0. Antibody
488 sequences have been deposited in NCBI GenBank, accession numbers OR825693-OR825700.
489

ACKNOWLEDGMENTS: We thank the many members of our Program Project Consortium for advice
and discussion. X-ray diffraction data were recorded at beamline ID-24-C (operated by the Northeast
Collaborative Access team: NE-CAT) at the Advanced Photon Source (APS, Argonne National
Laboratory). We thank NE-CAT staff members for advice and assistance in data collection. NE-CAT is
funded by NIH grant P30 GM124165. APS is operated for the DOE Office of Science by Argonne
National Laboratory under contract DE-AC02-06CH11357. The research was supported by NIAID
Program Project Grant P01 AI089618 (to G.H.K.), funds from the University of Pittsburgh Center for
Vaccine Research (to K.R.M), and Japan Agency for Medical Research and Development grant
JP22fk0108141 (to Y.A. and Y.T.)

499
500 **FIGURE LEGENDS:**
501
Figure 1: Identification of a novel, broadly binding, HA head-directed antibody class. A. Luminex
screening of Bmem-cell Nojima culture supernatants identified four antibodies that broadly react with
influenza A HA FLsEs and head domains. B. In a Luminex competitive binding assay, the four
antibodies from (A) that share a pattern of reactivity did not compete with antibodies that engage known
HA epitopes, but compete with each other for HA binding. Structures of Fab-HA complexes were
aligned on an HA trimer from A/American black duck/New Brunswick/00464/2010(H4N6) (PDB:
5XL2)[19]. Fab structures include HC19[20] (PDB 2VIR), S5V2-29[18] (PDB 6E4X), HC45[42] (PDB 1QFU),
CR9114[43] (PDB 4FQY), CR8020[44] (PDB 3SDY) and FI6v3[30] (PDB 3ZTJ). SARS-CoV antibody
CR3022[21] was used as an HA non-binding control. C. The cross-competing HA antibodies share
genetic signatures.
512
Figure 2: Human antibodies engage a recessed surface at the head-stem interface of the
influenza HA molecule. A. Structure of antibody S8V1-157 complexed with the HA head domain of

A/American black duck/New Brunswick/00464/2010(H4N6) colored in gray. The heavy chain is colored
darker blue and the light chain is lighter blue. Engagement of this site is incompatible with the defined
prefusion H4 HA trimer[19], colored in white (PDB: 5XL2) or with individual HA monomers. B. A surface
projection showing the degree of amino acid conservation among HAs engaged by this antibody class
(see Figure 3). The head-stem epitope is circumscribed in blue in the rightmost panel. Conservation
scores were produced using ConSurf [45,46]. C. Key S8V1-157 contacts. The orientation relative to panel
A is indicated.
522
523   **Figure 3: Breadth of HA binding by HA head-stem epitope antibodies.** A. Equilibrium dissociation
524   constants (K_d), determined by ELISA. Broadly binding influenza A HA antibody FI6v3 [30] and influenza B
525   HA antibody CR8071 [43] served as binding controls. B. Phylogenetic relationships of HAs used in our
526   panel. HAs bound by HA head-stem epitope antibodies are indicated. Binding data from Figure S3 are
527   incorporated into panel B.
528
Figure 4: HA head-stem epitope antibodies bind cell surface-anchored HA. Flow cytometry
histograms depict the fluorescence intensities of recombinant IgG binding to K530 cell lines expressing
recombinant, native HA on the cell surface. K530 cells were labeled with 400 ng/ml of the four head-
stem epitope antibodies or control antibodies targeting the HA receptor binding site (HC19[20], K03.12[36]
and H5.3[47]), the head interface (S5V2-29[18]), a lateral head epitope (HC45[42]), stem (FI6v3[30]), or SARS-
CoV spike protein (CR3022[21]). HA abbreviations correspond to: H1/SI06: A/Solomon
Islands/3/2006(H1N1), H1/CA09: A/California/04/2009(H1N1), H2/JP57: A/Japan/305/1957(H2N2),
H5/VN04: A/Viet Nam/1203/2004(H5N1), H6/TW13: A/Taiwan/2/2013(H6N1), H8/CA07: A/northern
shoveler/California/HKWF1204/2007(H8N4),H3/HK68: A/Aichi/02/1968(H3N2)(X31), H3/TX12:
A/Texas/50/2012(H3N2), H4/NB10:A/American black duck/New Brunswick/00464/2010, H7/TW13:
A/Taiwan/1/2017(H7N9), H10/JX13: A/Jiangxi/IPB13/2013(H10N8),H14/WI10:
A/mallard/Wisconsin/10OS3941/2010(H14N6).
541
Figure 5: HA head-stem epitope antibodies protect against lethal influenza virus infection and
severe disease. C57BL/6 mice (n = 7 per group) were intraperitoneally injected with 150 μg of
recombinant antibody via intraperitoneal injection three hours prior to intranasal challenge with 5xLD50
of A/Aichi/02/1968(H3N2)(X31). Mice were weighed daily and euthanized at a humane endpoint of 25%
loss of body weight. Antibodies passively transferred included musinized IgG1 and IgG2c versions of
HA head-stem epitope antibodies S8V1-157 and S1V2-65, neutralizing antibody HC19[20], head interface
antibody S5V2-29[18] and SARS-CoV antibody CR3022[21]. Mice injected with PBS were included as an
additional control. A. Post-infection survival rate. B-D. Body weight curves for infected mice
administered S8V1-157 (B), S1V2-65 (C), or S5V2-29 (D) antibodies, compared with controls. $*p <$
0.05 and $***p < 0.001$ compared with isotype control CR3022. Not significant (n.s.), p 0.05; † p < 0.05,
†† p < 0.01, and †††† p < 0.0001 IgG2c compared with IgG1.
553
554   **Figure 6: The HA head-stem epitope epitope is immunogenic in humans.** A. An additional 528
555   Nojima culture supernatants from donors K01, K03, S1, S5, S8, S9 and S12 were screened for
556   competition with a recombinant musinized S8V1-157 IgG1 for HA binding. Culture supernatants that
557   inhibited S8V1-157 binding by >90% are colored and specified. B. HA reactivity of S8V1-157-competing
558   Nojima culture supernatants, as determined by multiplex Luminex assay. N.D.: not determined.
559
560   **Supporting Figures:**
561
562   **Supporting Table 1: Data collection and refinement statistics.**
563
564   **Figure S1: Surface hydrophobicity on the HA molecule.** Surface hydrophobicity and the head-stem
565   epitope are shown on the HA trimer of A/American black duck/New Brunswick/00464/2010(H4N6)
566   (PDB: 5XL2) [19]. Coloring is based on the PyMOL Color h script that utilizes a normalized consensus
567   hydrophobicity scale[48]. Views and orientations match those in Figure 2. The head-stem epitope is
568   circled in the far right panel.

Figure S2: ELISA titrations of antibodies on HA coated plates. The broadly binding stem antibody
FI6v3[30] was used as a positive control and an influenza B specific head antibody, CR8071[43], as a
negative control for influenza A isolates. Data points represent the average of three technical replicates.
The standard error of the mean is shown for each point. KDs were calculated from the curves fit to
these data points.

Figure S3: Head-stem epitope antibody binding to HA head-only domains. A. Structures of HA
head domains used in these experiments the S8V1-157-HA head complex is shown for reference. A
compact head domain[33] , is shown for reference (PDB 7TRH). B. HA heads for HAs not bound or not
expressed in Figure 3 were produced alongside a positive control A/California/07/2009(H1N1)(X-181)
and a compact head version. Sequences of the HA head domains are aligned to the
A/Aichi/02/1968(H3N2)(X-31) reference sequence and numbered by H3 convention. C. Dissociation
constants from ELISA measurements of head-stem epitope antibodies to HA head domains. Head
interface antibody FluA-20[23] was used as a positive control and influenza B head antibody CR8071[43] as
a negative control.

**Figure S4: ELISA titrations of antibodies on HA head coated plates and characterization of cell
expressed HA.** A. Head interface antibody FluA-20[23] was used as a positive control and influenza B
head antibody CR8071[43] as a negative control. Data points represent the average of three technical
replicates. The standard error of the mean is shown for each point. KDs were calculated from the
curves fit to these data points. B. Lysates from 293F transfected with either HA from
A/Aichi/02/1968(H3N2)(X31) or empty vector were subjected to western blotting with either anti-HA tag
antibody, which recognizes an endogenous sequence in the H3 HA head (HA1), or anti-GAPDH
antibody. Molecular weights of unprocessed HA0 and processed HA1 are indicated with open or closed
arrowheads, respectively. C. 293F transfected with either HA from A/Aichi/02/1968(H3N2)(X31) or
empty vector were stained with the indicated antibodies and analyzed by flow cytometry. Gates
denoting cells bound by antibody are shown, with the percent of the total population indicated above.

Figure S5: S8V1-157 is a non-neutralizing antibody. Neutralization IC50 values for S8V1-157,
neutralizing antibody HC19[20] (positive control) and SARS-CoV antibody CR3022[21] (negative control).

Figure S6: S8V1-157 protects against lethal influenza virus infection. C57BL/6 mice received a
passive transfer of 100μg of recombinant antibody via intraperitoneal injection three hours prior to
challenge with a 3xLD50 of A/Aichi/02/1968(H3N2)(X31) administered intranasally. Mice were
monitored for survival (A) and weight loss (B). Mice were sacrificed at a humane endpoint of 25% loss
of body weight. Antibodies passively transferred include musinized IgG2c versions of HA head-stem
epitope antibody S8V1-157 (n=6), neutralizing antibody HC19[20] (n=4), and SARS-CoV antibody
CR3022[21] (n=4).

Figure S7: Antibody genetics of S8V1-157 competing antibodies. Antibody names, gene usage
and HCDR3 sequences for S8V1-157 competing antibodies identified in Figure 6.

REFERENCES:

1. Knossow, M., and Skehel, J.J. (2006). Variation and infectivity neutralization in influenza. Immunology *119*, 1-7. 10.1111/j.1365-2567.2006.02421.x.
2. Wu, N.C., and Wilson, I.A. (2020). Influenza Hemagglutinin Structures and Antibody Recognition. Cold Spring Harb Perspect Med *10*. 10.1101/cshperspect.a038778.
3. Krammer, F., Palese, P., and Steel, J. (2015). Advances in universal influenza virus vaccine design and antibody mediated therapies based on conserved regions of the hemagglutinin. Curr Top Microbiol Immunol *386*, 301-321. 10.1007/82_2014_408.
4. Krammer, F., Garcia-Sastre, A., and Palese, P. (2018). Is It Possible to Develop a "Universal" Influenza Virus Vaccine? Potential Target Antigens and Critical Aspects for a Universal Influenza Vaccine. Cold Spring Harb Perspect Biol *10*. 10.1101/cshperspect.a028845.
5. Erbelding, E.J., Post, D.J., Stemmy, E.J., Roberts, P.C., Augustine, A.D., Ferguson, S., Paules, C.I., Graham, B.S., and Fauci, A.S. (2018). A Universal Influenza Vaccine: The Strategic Plan for the National Institute of Allergy and Infectious Diseases. J Infect Dis *218*, 347-354. 10.1093/infdis/jiy103.
6. Cox, R.J. (2013). Correlates of protection to influenza virus, where do we go from here? Hum Vaccin Immunother *9*, 405-408. 10.4161/hv.22908.
7. Skehel, J.J., Bizebard, T., Bullough, P.A., Hughson, F.M., Knossow, M., Steinhauer, D.A., Wharton, S.A., and Wiley, D.C. (1995). Membrane fusion by influenza hemagglutinin. Cold Spring Harb Symp Quant Biol *60*, 573-580. 10.1101/sqb.1995.060.01.061.
8. Skehel, J.J., and Wiley, D.C. (2000). Receptor binding and membrane fusion in virus entry: the influenza hemagglutinin. Annu Rev Biochem *69*, 531-569. 10.1146/annurev.biochem.69.1.531.
9. Weissenhorn, W., Dessen, A., Calder, L.J., Harrison, S.C., Skehel, J.J., and Wiley, D.C. (1999). Structural basis for membrane fusion by enveloped viruses. Mol Membr Biol *16*, 3-9. 10.1080/096876899294706.
10. Benton, D.J., Gamblin, S.J., Rosenthal, P.B., and Skehel, J.J. (2020). Structural transitions in influenza haemagglutinin at membrane fusion pH. Nature *583*, 150-153. 10.1038/s41586-020-2333-6.
11. Das, D.K., Govindan, R., Nikic-Spiegel, I., Krammer, F., Lemke, E.A., and Munro, J.B. (2018). Direct Visualization of the Conformational Dynamics of Single Influenza Hemagglutinin Trimers. Cell *174*, 926-937 e912. 10.1016/j.cell.2018.05.050.
12. Casalino, L., Seitz, C., Lederhofer, J., Tsybovsky, Y., Wilson, I.A., Kanekiyo, M., and Amaro, R.E. (2022). Breathing and Tilting: Mesoscale Simulations Illuminate Influenza Glycoprotein Vulnerabilities. ACS Cent Sci *8*, 1646-1663. 10.1021/acscentsci.2c00981.
13. Garcia, N.K., Kephart, S.M., Benhaim, M.A., Matsui, T., Mileant, A., Guttman, M., and Lee, K.K. (2023). Structural dynamics reveal subtype-specific activation and inhibition of influenza virus hemagglutinin. J Biol Chem *299*, 104765. 10.1016/j.jbc.2023.104765.
14. Guthmiller, J.J., Han, J., Utset, H.A., Li, L., Lan, L.Y., Henry, C., Stamper, C.T., McMahon, M., O'Dell, G., Fernandez-Quintero, M.L., et al. (2022). Broadly neutralizing antibodies target a haemagglutinin anchor epitope. Nature *602*, 314-320. 10.1038/s41586-021-04356-8.
15. Benton, D.J., Nans, A., Calder, L.J., Turner, J., Neu, U., Lin, Y.P., Ketelaars, E., Kallewaard, N.L., Corti, D., Lanzavecchia, A., et al. (2018). Influenza hemagglutinin membrane anchor. Proc Natl Acad Sci U S A *115*, 10112-10117. 10.1073/pnas.1810927115.
16. Adachi, Y., Tonouchi, K., Nithichanon, A., Kuraoka, M., Watanabe, A., Shinnakasu, R., Asanuma, H., Ainai, A., Ohmi, Y., Yamamoto, T., et al. (2019). Exposure of an occluded hemagglutinin

epitope drives selection of a class of cross-protective influenza antibodies. Nat Commun *10*,
3883. 10.1038/s41467-019-11821-6.

17. Tonouchi, K., Adachi, Y., Suzuki, T., Kuroda, D., Nishiyama, A., Yumoto, K., Takeyama, H., Suzuki,
T., Hashiguchi, T., and Takahashi, Y. (2023). Structural basis for cross-group recognition of an
influenza virus hemagglutinin antibody that targets postfusion stabilized epitope. PLoS Pathog
19, e1011554. 10.1371/journal.ppat.1011554.

18. Watanabe, A., McCarthy, K.R., Kuraoka, M., Schmidt, A.G., Adachi, Y., Onodera, T., Tonouchi, K.,
Caradonna, T.M., Bajic, G., Song, S., et al. (2019). Antibodies to a Conserved Influenza Head
Interface Epitope Protect by an IgG Subtype-Dependent Mechanism. Cell *177*, 1124-1135
e1116. 10.1016/j.cell.2019.03.048.

19. Song, H., Qi, J., Xiao, H., Bi, Y., Zhang, W., Xu, Y., Wang, F., Shi, Y., and Gao, G.F. (2017). Avian-
to-Human Receptor-Binding Adaptation by Influenza A Virus Hemagglutinin H4. Cell Rep *20*,
1201-1214. 10.1016/j.celrep.2017.07.028.

20. Fleury, D., Wharton, S.A., Skehel, J.J., Knossow, M., and Bizebard, T. (1998). Antigen distortion
allows influenza virus to escape neutralization. Nat Struct Biol *5*, 119-123. 10.1038/nsb0298-
119.

21. ter Meulen, J., van den Brink, E.N., Poon, L.L., Marissen, W.E., Leung, C.S., Cox, F., Cheung, C.Y.,
Bakker, A.Q., Bogaards, J.A., van Deventer, E., et al. (2006). Human monoclonal antibody
combination against SARS coronavirus: synergy and coverage of escape mutants. PLoS Med *3*,
e237. 10.1371/journal.pmed.0030237.

22. Zost, S.J., Dong, J., Gilchuk, I.M., Gilchuk, P., Thornburg, N.J., Bangaru, S., Kose, N., Finn, J.A.,
Bombardi, R., Soto, C., et al. (2021). Canonical features of human antibodies recognizing the
influenza hemagglutinin trimer interface. J Clin Invest *131*. 10.1172/JCI146791.

23. Bangaru, S., Lang, S., Schotsaert, M., Vanderven, H.A., Zhu, X., Kose, N., Bombardi, R., Finn, J.A.,
Kent, S.J., Gilchuk, P., et al. (2019). A Site of Vulnerability on the Influenza Virus Hemagglutinin
Head Domain Trimer Interface. Cell *177*, 1136-1152 e1118. 10.1016/j.cell.2019.04.011.

24. Yu, F., Song, H., Wu, Y., Chang, S.Y., Wang, L., Li, W., Hong, B., Xia, S., Wang, C., Khurana, S., et
al. (2017). A Potent Germline-like Human Monoclonal Antibody Targets a pH-Sensitive Epitope
on H7N9 Influenza Hemagglutinin. Cell Host Microbe *22*, 471-483 e475.
10.1016/j.chom.2017.08.011.

25. McCarthy, K.R., Lee, J., Watanabe, A., Kuraoka, M., Robinson-McCarthy, L.R., Georgiou, G.,
Kelsoe, G., and Harrison, S.C. (2021). A Prevalent Focused Human Antibody Response to the
Influenza Virus Hemagglutinin Head Interface. mBio *12*, e0114421. 10.1128/mBio.01144-21.

26. Bajic, G., Maron, M.J., Adachi, Y., Onodera, T., McCarthy, K.R., McGee, C.E., Sempowski, G.D.,
Takahashi, Y., Kelsoe, G., Kuraoka, M., and Schmidt, A.G. (2019). Influenza Antigen Engineering
Focuses Immune Responses to a Subdominant but Broadly Protective Viral Epitope. Cell Host
Microbe *25*, 827-835 e826. 10.1016/j.chom.2019.04.003.

27. Wu, Y., and Gao, G.F. (2019). "Breathing" Hemagglutinin Reveals Cryptic Epitopes for Universal
Influenza Vaccine Design. Cell *177*, 1086-1088. 10.1016/j.cell.2019.04.034.

28. Otterstrom, J.J., Brandenburg, B., Koldijk, M.H., Juraszek, J., Tang, C., Mashaghi, S., Kwaks, T.,
Goudsmit, J., Vogels, R., Friesen, R.H., and van Oijen, A.M. (2014). Relating influenza virus
membrane fusion kinetics to stoichiometry of neutralizing antibodies at the single-particle level.
Proc Natl Acad Sci U S A *111*, E5143-5148. 10.1073/pnas.1411755111.

29. DiLillo, D.J., Tan, G.S., Palese, P., and Ravetch, J.V. (2014). Broadly neutralizing hemagglutinin
stalk-specific antibodies require FcgammaR interactions for protection against influenza virus in
vivo. Nat Med *20*, 143-151. 10.1038/nm.3443.

30.   Corti, D., Voss, J., Gamblin, S.J., Codoni, G., Macagno, A., Jarrossay, D., Vachieri, S.G., Pinna, D.,
710   Minola, A., Vanzetta, F., et al. (2011). A neutralizing antibody selected from plasma cells that
711   binds to group 1 and group 2 influenza A hemagglutinins. Science *333*, 850-856.
712   10.1126/science.1205669.

713   31.   Song, S., Manook, M., Kwun, J., Jackson, A.M., Knechtle, S.J., and Kelsoe, G. (2021). A cell-based
714   multiplex immunoassay platform using fluorescent protein-barcoded reporter cell lines.
715   Commun Biol *4*, 1338. 10.1038/s42003-021-02881-w.

716   32.   McCarthy, K.R., Raymond, D.D., Do, K.T., Schmidt, A.G., and Harrison, S.C. (2019). Affinity
717   maturation in a human humoral response to influenza hemagglutinin. Proc Natl Acad Sci U S A.
718   10.1073/pnas.1915620116.

719   33.   Schmidt, A.G., Xu, H., Khan, A.R., O'Donnell, T., Khurana, S., King, L.R., Manischewitz, J., Golding,
720   H., Suphaphiphat, P., Carfi, A., et al. (2013). Preconfiguration of the antigen-binding site during
721   affinity maturation of a broadly neutralizing influenza virus antibody. Proc Natl Acad Sci U S A
722   *110*, 264-269. 10.1073/pnas.1218256109.

723   34.   Schmidt, A.G., Therkelsen, M.D., Stewart, S., Kepler, T.B., Liao, H.X., Moody, M.A., Haynes, B.F.,
724   and Harrison, S.C. (2015). Viral receptor-binding site antibodies with diverse germline origins.
725   Cell *161*, 1026-1034. 10.1016/j.cell.2015.04.028.

726   35.   Kuraoka, M., Schmidt, A.G., Nojima, T., Feng, F., Watanabe, A., Kitamura, D., Harrison, S.C.,
727   Kepler, T.B., and Kelsoe, G. (2016). Complex Antigens Drive Permissive Clonal Selection in
728   Germinal Centers. Immunity *44*, 542-552. 10.1016/j.immuni.2016.02.010.

729   36.   McCarthy, K.R., Watanabe, A., Kuraoka, M., Do, K.T., McGee, C.E., Sempowski, G.D., Kepler,
730   T.B., Schmidt, A.G., Kelsoe, G., and Harrison, S.C. (2018). Memory B Cells that Cross-React with
731   Group 1 and Group 2 Influenza A Viruses Are Abundant in Adult Human Repertoires. Immunity
732   *48*, 174-184 e179. 10.1016/j.immuni.2017.12.009.

733   37.   Kabsch, W. (2010). Xds. Acta Crystallogr D Biol Crystallogr *66*, 125-132.
734   10.1107/S0907444909047337.

735   38.   McCoy, A.J., Grosse-Kunstleve, R.W., Adams, P.D., Winn, M.D., Storoni, L.C., and Read, R.J.
736   (2007). Phaser crystallographic software. J Appl Crystallogr *40*, 658-674.
737   10.1107/S0021889807021206.

738   39.   Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J., Hung,
739   L.W., Kapral, G.J., Grosse-Kunstleve, R.W., et al. (2010). PHENIX: a comprehensive Python-based
740   system for macromolecular structure solution. Acta Crystallogr D Biol Crystallogr *66*, 213-221.
741   10.1107/S0907444909052925.

742   40.   Emsley, P., and Cowtan, K. (2004). Coot: model-building tools for molecular graphics. Acta
743   Crystallogr D Biol Crystallogr *60*, 2126-2132. 10.1107/S0907444904019158.

744   41.   Chen, V.B., Arendall, W.B., 3rd, Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral, G.J., Murray,
745   L.W., Richardson, J.S., and Richardson, D.C. (2010). MolProbity: all-atom structure validation for
746   macromolecular crystallography. Acta Crystallogr D Biol Crystallogr *66*, 12-21.
747   10.1107/S0907444909042073.

748   42.   Bizebard, T., Gigant, B., Rigolet, P., Rasmussen, B., Diat, O., Bosecke, P., Wharton, S.A., Skehel,
749   J.J., and Knossow, M. (1995). Structure of influenza virus haemagglutinin complexed with a
750   neutralizing antibody. Nature *376*, 92-94. 10.1038/376092a0.

751   43.   Dreyfus, C., Laursen, N.S., Kwaks, T., Zuijdgeest, D., Khayat, R., Ekiert, D.C., Lee, J.H., Metlagel,
752   Z., Bujny, M.V., Jongeneelen, M., et al. (2012). Highly conserved protective epitopes on
753   influenza B viruses. Science *337*, 1343-1348. 10.1126/science.1222908.

44.   Ekiert, D.C., Friesen, R.H., Bhabha, G., Kwaks, T., Jongeneelen, M., Yu, W., Ophorst, C., Cox, F.,
Korse, H.J., Brandenburg, B., et al. (2011). A highly conserved neutralizing epitope on group 2
influenza A viruses. Science *333*, 843-850. 10.1126/science.1204839.
45.   Ashkenazy, H., Abadi, S., Martz, E., Chay, O., Mayrose, I., Pupko, T., and Ben-Tal, N. (2016).
ConSurf 2016: an improved methodology to estimate and visualize evolutionary conservation in
macromolecules. Nucleic Acids Res *44*, W344-350. 10.1093/nar/gkw408.
46.   Celniker, G., Nimrod, G., Ashkenazy, H., Glaser, F., Martz, E., Mayrose, I., Pupko, T., and Ben-Tal,
N. (2013). ConSurf: Using Evolutionary Data to Raise Testable Hypotheses about Protein
Function. Israel Journal of Chemistry *53*, 199-206. https://doi.org/10.1002/ijch.201200096.
47.   Winarski, K.L., Thornburg, N.J., Yu, Y., Sapparapu, G., Crowe, J.E., Jr., and Spiller, B.W. (2015).
Vaccine-elicited antibody that neutralizes H5N1 influenza and variants binds the receptor site
and polymorphic sites. Proc Natl Acad Sci U S A *112*, 9346-9351. 10.1073/pnas.1502762112.
48.   Eisenberg, D., Schwarz, E., Komaromy, M., and Wall, R. (1984). Analysis of membrane and
surface protein sequences with the hydrophobic moment plot. J Mol Biol *179*, 125-142.
10.1016/0022-2836(84)90309-7.
769
770

771
772
773
774
Figure 1: Identification of a novel, broadly binding, HA head-directed antibody class. A. Luminex
screening of Bmem-cell Nojima culture supernatants identified four antibodies that broadly react with
influenza A HA FLsEs and head domains. B. In a Luminex competitive binding assay, the four
antibodies from (A) that share a pattern of reactivity did not compete with antibodies that engage known
HA epitopes, but compete with each other for HA binding. Structures of Fab-HA complexes were
aligned on an HA trimer from A/American black duck/New Brunswick/00464/2010(H4N6) (PDB:
5XL2)[19]. Fab structures include HC19[20] (PDB 2VIR), S5V2-29[18] (PDB 6E4X), HC45[42] (PDB 1QFU),
CR9114[43] (PDB 4FQY), CR8020[44] (PDB 3SDY) and FI6v3[30] (PDB 3ZTJ). SARS-CoV antibody
CR3022[21] was used as an HA non-binding control. C. The cross-competing HA antibodies share
genetic signatures.
785
786

787
Figure 2: Human antibodies engage a recessed surface at the head-stem interface of the
influenza HA molecule. A. Structure of antibody S8V1-157 complexed with the HA head domain of
A/American black duck/New Brunswick/00464/2010(H4N6) colored in gray. The heavy chain is colored
darker blue and the light chain is lighter blue. Engagement of this site is incompatible with the defined
prefusion H4 HA trimer[19], colored in white (PDB: 5XL2) or with individual HA monomers. B. A surface
projection showing the degree of amino acid conservation among HAs engaged by this antibody class
(see Figure 3). The head-stem epitope is circumscribed in blue in the rightmost panel. Conservation
scores were produced using ConSurf [45,46]. C. Key S8V1-157 contacts. The orientation relative to panel
A is indicated.

797
798

		S1V2-17	S1V2-60	S1V2-65	S8V1-157	CR8071	FI6
H1N1	A/USSR/90/1977	7.0	1.2	2.2	2.7	>100	0.32
	A/Chile/01/1983	13	3.7	5.3	8.1	>100	0.33
	A/Texas/36/1991	2.9	0.73	1.0	1.2	>100	0.28
	A/Solomon Islands/3/2006	14	3.5	4.1	8.3	>100	0.67
	A/California/04/2009	6.9	0.86	1.0	3.8	>100	0.42
	A/Victoria/2570/2019(IVR-215)	2.8	0.39	0.40	2.1	<100	0.35
H2N2	A/Japan/305/1957	6.5	8.8	10	2.5	>100	0.36
H3N2	A/Aichi/02/1968(X31)	0.34	0.16	0.16	0.18	>100	0.31
	A/Philippines/02/1982	0.58	0.25	0.21	0.29	>100	1.2
	A/Sydney/05/1997	0.50	0.27	0.25	0.21	>100	3.0
	A/Wisconsin/67/2005	0.99	0.26	0.21	0.42	>100	3.5
	A/Hong Kong/4801/2014	0.85	0.26	0.21	0.49	>100	0.88
	A/Tasmania/503/2020(IVR-221)	0.62	0.28	0.21	0.22	>100	2.8
H4N6	A/American black duck/New Brunswick/00464/2010	0.55	0.17	0.20	0.32	>100	1.4
H5N1	A/Viet Nam/1203/2004	0.14	0.10	0.10	<0.1	>100	0.44
H6N1	A/Taiwan/2/2013	>100	>100	>100	>100	>100	0.47
H7N9	A/Taiwan/1/2017	0.11	0.13	0.12	<0.1	>100	0.24
H8N4	A/northern shoveler/California/HKWF1204/2007	9.2	0.77	1.7	4.5	>100	0.29
H9N2	A/turkey/Wisconsin/1/1966	>100	3.2	>100	>100	>100	0.37
	A/swine/Hong Kong/9/98	13	0.48	8.0	26	>100	0.40
H11N9	A/duck/Memphis/546/1974	>100	>100	>100	>100	>100	0.74
H12N5	A/duck/Alberta/60/1976	>100	>100	>100	>100	>100	0.44
H14N6	A/mallard/Wisconsin/10OS3941/2010	1.9	0.26	0.30	1.2	>100	1.4
H15N9	A/wedge-tailed shearwater/Western Australia/2576/1979	61	0.65	8.0	6.0	>100	2.5
H16N3	A/black-headed gull/Sweden/2/99	>100	20	>100	>100	>100	0.55
Influenza B	B/Brisbane/60/2008	>100	>100	>100	>100	0.35	>100

K_d in nM				
<1	1-5	5-25	25-100	>100

B. Group 1 HAs / Group 2 HAs

Bound by all 4 antibodies
Bound only by S1V2-60

0.5

Figure 3: Breadth of HA binding by HA head-stem epitope antibodies. A. Equilibrium dissociation constants (K_d), determined by ELISA. Broadly binding influenza A HA antibody FI6v3 [30] and influenza B HA antibody CR8071 [43] served as binding controls. B. Phylogenetic relationships of HAs used in our panel. HAs bound by HA head-stem epitope antibodies are indicated. Binding data from Figure S3 are incorporated into panel B.

808
Figure 4: HA head-stem epitope antibodies bind cell surface-anchored HA. Flow cytometry
histograms depict the fluorescence intensities of recombinant IgG binding to K530 cell lines expressing
recombinant, native HA on the cell surface. K530 cells were labeled with 400 ng/ml of the four head-
stem epitope antibodies or control antibodies targeting the HA receptor binding site (HC19[20], K03.12[36]
and H5.3[47]), the head interface (S5V2-29[18]), a lateral head epitope (HC45[42]), stem (FI6v3[30]), or SARS-
CoV spike protein (CR3022[21]).
815
816

Figure 5: HA head-stem epitope antibodies protect against lethal influenza virus infection and severe disease. C57BL/6 mice (n = 7 per group) were intraperitoneally injected with 150 μg of recombinant antibody via intraperitoneal injection three hours prior to intranasal challenge with 5xLD50 of A/Aichi/02/1968(H3N2)(X31). Mice were weighed daily and euthanized at a humane endpoint of 25% loss of body weight. Antibodies passively transferred included musinized IgG1 and IgG2c versions of HA head-stem epitope antibodies S8V1-157 and S1V2-65, neutralizing antibody HC19[20], head interface antibody S5V2-29[18] and SARS-CoV antibody CR3022[21]. Mice injected with PBS were included as an additional control. A. Post-infection survival rate. B-D. Body weight curves for infected mice administered S8V1-157 (B), S1V2-65 (C), or S5V2-29 (D) antibodies, compared with controls. *p < 0.05 and ***p < 0.001 compared with isotype control CR3022. Not significant (n.s.), p 0.05; † p < 0.05, †† p < 0.01, and †††† p < 0.0001 IgG2c compared with IgG1.

A.

Legend:
- S8V6 P5-C6
- S8V6 P7-H9
- S8V6 P10-G1
- S1V4 P4-C5
- S5V4 P1-A2
- S5V4 P2-D2
- S12V4 P1-F1
- S12V6 P7-H4
- S8V1-157r (recombinant Ab ctrl)

B.

Antibody	H1 2009 X	H1 2009	H1 2006	H3 1968	H3 2005	H3 2014[a]	H3 2014[b]	H5 2004	BYam2013	BVic 2004	IgG	Igκ	Igλ
S8V5 P5-C6	211,868	N.D.	108,575					N.D.			85,808	80,971	
S8V6 P7-H9		N.D.		57,232	41,063	17,839	35,872	N.D.			100,560	104,082	
S8V6 P10-G1	183,143	N.D.	27,673					N.D.			80,617		65,397
S1V4 P4-C5	11,528	2,664		26,677	1,387	13,996	7,592	2,315	200		64,575	78,170	
S5V4 P1-A2		1,555		53,091	67,485	29,348	111,273				64,266	74,232	
S5V4 P2-D2		2,711	200	93,375	105,075	79,837	166,328				82,873	91,355	
S12V4 P1-F1	42,514	19,363	600	9,458	3,555	22,409	5,793	16,648			59,983	73,887	
S12V6 P7-H4		1,225		600	18,737	11,681	1,041				66,471	68,493	

MFI				
<100	100-1,000	1,000-10,000	10,000-100,000	>100,000

Figure 6: The HA head-stem epitope epitope is immunogenic in humans. A. An additional 528 Nojima culture supernatants from donors K01, K03, S1, S5, S8, S9 and S12 were screened for competition with a recombinant musinized S8V1-157 IgG1 for HA binding. Culture supernatants that inhibited S8V1-157 binding by >90% are colored and specified. B. HA reactivity of S8V1-157-competing Nojima culture supernatants, as determined by multiplex Luminex assay. N.D.: not determined.

839 **Supporting Figures:**
840
841

842 **Supporting Table 1: Data collection and refinement statistics.**

	S8V1-157-A/American black duck/New Brunswick/00464/2010(H4N6) Fab-HA head complex
PDB ID	8US0
Data Collection	APS 24-ID-C
Number of datasets	1
Resolution, Å	49.17 - 3.704 (3.836 - 3.704)
Wavelength (Å)	0.9792
Space Group	P 1 21 1
Unit cell dimensions (a, b, c), Å	127.92 241.54 144.13
Unit cell angles (α, β, γ) °	90.00 101.62 90.00
I/σ	7.00 (1.20)
Rmeas	0.15 (1.12)
Rpim	0.10 (0.80)
Rmerge, %	0.10 (0.80)
CC*	1.00 (0.79)
CC½	0.99 (0.45)
Completeness, %	98.77 (98.49)
Number of reflections	175949 (17445)
Redundancy	2.0 (2.0)
Refinement	
Number of reflections:	
Working	89804 (8919)
Free	4506 (491)
Rwork, %	24.45 (38.84)
Rfree, %	29.14 (41.97)
Ramachandran plot, % (favored, disallowed)	94.83 (0.14)
Rmsd bond lengths, Å	0.002
Rmsd bond angles, °	0.49
Average B-factor	144.76

Rmerge, $\Sigma h,k,l \Sigma i |Ii(hkl) - \langle I(hkl)\rangle|/\Sigma h,k,l \Sigma i Ii(hkl)$, where I is an intensity that is observed i times; I/σ, signal-to-noise ratio (average observed intensity divided by average standard deviation of the observed intensity); Rwork, $\Sigma h,k,l ||Fobs| - |Fcalc||/\Sigma h,k,l |Fobs|$, where h, k, l covers the "working set" of observed structure factor amplitude (Fobs) reflections used in refinement (total reflections minus the test set) and Fcalc is the calculated structure factor amplitude; Rfree, calculated as for Rwork but on 5% of data excluded prior to refinement. Values in parentheses refer to highest-resolution shell. Related to Experimental Procedures.

843
844
845

846
847 **Figure S1: Surface hydrophobicity on the HA molecule.** Surface hydrophobicity and the head-stem
848 epitope are shown on the HA trimer of A/American black duck/New Brunswick/00464/2010(H4N6)
849 (PDB: 5XL2) [19]. Coloring is based on the PyMOL Color h script that utilizes a normalized consensus
850 hydrophobicity scale[48]. Views and orientations match those in Figure 2. The head-stem epitope is
851 circled in the far right panel.
852
853

854
855 **Figure S2: ELISA titrations of antibodies on HA coated plates.** The broadly binding stem antibody
856 FI6v3[30] was used as a positive control and an influenza B specific head antibody, CR8071[43], as a
857 negative control for influenza A isolates. Data points represent the average of three technical replicates.
858 The standard error of the mean is shown for each point. KDs were calculated from the curves fit to
859 these data points.
860
861

Figure S3

Figure S3: Head-stem epitope antibody binding to HA head-only domains. A. Structures of HA head domains used in these experiments the S8V1-157-HA head complex is shown for reference. A compact head domain[33] , is shown for reference (PDB 7TRH). B. HA heads for HAs not bound or not expressed in Figure 3 were produced alongside a positive control A/California/07/2009(H1N1)(X-181)

867 and a compact head version. Sequences of the HA head domains are aligned to the
868 A/Aichi/02/1968(H3N2)(X-31) reference sequence and numbered by H3 convention. C. Dissociation
869 constants from ELISA measurements of head-stem epitope antibodies to HA head domains. Head
870 interface antibody FluA-20[23] was used as a positive control and influenza B head antibody CR8071[43] as
871 a negative control.
872
873

Figure S4

A   Compact HA head domain

Full HA head domain

B

C

Figure S4: ELISA titrations of antibodies on HA head coated plates and characterization of cell expressed HA. A. Head interface antibody FluA-20[23] was used as a positive control and influenza B head antibody CR8071[43] as a negative control. Data points represent the average of three technical replicates. The standard error of the mean is shown for each point. KDs were calculated from the curves fit to these data points. B. Lysates from 293F transfected with either HA from A/Aichi/02/1968(H3N2)(X31) or empty vector were subjected to western blotting with either anti-HA tag antibody, which recognizes an endogenous sequence in the H3 HA head (HA1), or anti-GAPDH antibody. Molecular weights of unprocessed HA0 and processed HA1 are indicated with open or closed

883   arrowheads, respectively. C. 293F transfected with either HA from A/Aichi/02/1968(H3N2)(X31) or
884   empty vector were stained with the indicated antibodies and analyzed by flow cytometry. Gates
885   denoting cells bound by antibody are shown, with the percent of the total population indicated above.
886
887

	A/Aichi/02/1968(H3N2)(X-31)		
Antibody	HC19	S8V1-157	CR3022
IC50 µg/mL	<3.5	>3535.5	>3535.5

888
889 **Figure S5: S8V1-157 is a non-neutralizing antibody.** Neutralization IC50 values for S8V1-157,
890 neutralizing antibody HC19[20] (positive control) and SARS-CoV antibody CR3022[21] (negative control).
891
892

893
894
Figure S6: S8V1-157 protects against lethal influenza virus infection. C57BL/6 mice received a
passive transfer of 100μg of recombinant antibody via intraperitoneal injection three hours prior to
challenge with a 3xLD50 of A/Aichi/02/1968(H3N2)(X31) administered intranasally. Mice were
monitored for survival (A) and weight loss (B). Mice were sacrificed at a humane endpoint of 25% loss
of body weight. Antibodies passively transferred include musinized IgG2c versions of HA head-stem
epitope antibody S8V1-157 (n=6), neutralizing antibody HC19[20] (n=4), and SARS-CoV antibody
CR3022[21] (n=4).
902
903

Antibody	V_H	J_H	D	HCDR3	Length	V_L	J_L	LCDR3	LCDR3
S8V5_P5-C6	IGHV4-61*08	IGHJ6*02	IGHD3-10*01	C ARVSMVPGEGWFFGVDV W	17	IGKV3-20*01	IGKJ4*01	C QHYGGSLT F	8
S8V6_P7-H9	IGHV1-8*01	IGHJ4*02	IGHD6-13*01	C VRGDXAAAXY W	10	IGKV4-1*01	IGKJ2*01	C QQYYSTPMYT F	10
S8V6_P10-G1	unrecoverable	-	-	–	-	unrecoverable	-	–	-
S1V4_P4-C5	IGHV1-24*01	IGHJ6*03	IGHD4-23*01	C AITLSYYYMNV W	11	IGKV2-28*01 or IGKV2D-28*01	IGKJ1*01	C MQALQTPPWT F	10
S5V4_P1-A2	IGHV1-18*01	IGHJ3*02	IGHD3-3*01	C ARSKFGLVGRDVFDI W	15	unrecoverable	-	–	-
S5V4_P2-D2	IGHV1-18*01	IGHJ3*02	IGHD4-23*01	C ARSKLGLVGRDVFDI W	15	IGKV4-1*01	IGKJ4*01	C QQYYDTPPS F	9
S12V4_P1-F1	IGHV4-31*04	IGHJ6*02	IGHD1-7*01	C ARETGIISGSNFFYYGMDV W	19	IGKV3-15*01	IGKJ1*01	C QQYNNWPLRT F	10
S12V6_P7-H4	IGHV1-24*01	IGHJ4*02	IGHD3-3*02	C AINLGGVPTTIL W	12	IGKV2-28*01 or IGKV2D-28*01	IGKJ5*01	C MQALQTPIT F	9

904
905 **Figure S7: Antibody genetics of S8V1-157 competing antibodies.** Antibody names, gene usage
906 and HCDR3 sequences for S8V1-157 competing antibodies identified in Figure 6.
907
908