Phylogenomic Insights into the Evolution and Origin of Nematoda

Xue Qing[1] [#], Y. Miles Zhang[2#], Sidi Sun[1], Mohammed Ahmed[3], Wen-Sui Lo[4], Wim Bert[5],
Oleksandr Holovachov[6], Hongmei Li[1*]

[1] Department of Plant Pathology, Nanjing Agricultural University, Nanjing 210095, China

[2] Institute of Ecology and Evolution, University of Edinburgh, Edinburgh, UK

[3] School of Life Sciences, Liverpool University, Liverpool, UK

[4] Institute of Future Agriculture, Northwest A&F University, Yangling 712100, China

[5] Department of Biology, Ghent University, Ghent B-9000, Belgium

[6] Department of Zoology, Swedish Museum of Natural History, Stockholm SE-104 05,
Sweden

[#] These authors contributed equally to this work

* Corresponding author, e-mail: lihm@njau.edu.cn

Running title: Phylogenomic of Nematoda

Abstract:

The phylum Nematoda represents one of the most cosmopolitan and abundant metazoan
groups on Earth. In this study, we reconstructed the phylogenomic tree for phylum Nematoda.
A total of 60 genomes, belonging to eight nematode orders, were newly sequenced, providing
the first low-coverage genomes for the orders Dorylaimida, Mononchida, Monhysterida,
Chromadorida, Triplonchida, and Enoplida. The resulting phylogeny is well-resolved across
most clades, with topologies remaining consistent across various reconstruction parameters.
The subclass Enoplia is placed as a sister group to the rest of Nematoda, agrees with previous
published phylogenies. While the order Triplonchida is monophyletic, it is not well-supported,
and the order Enoplida is paraphyletic. Taxa possessing a stomatostylet form a monophyletic
group; however, the superfamily Aphelenchoidea does not constitute a monophyletic clade. The
genera *Trichinella* and *Trichuris* are inferred to have shared a common ancestor approximately
202 mya, a considerably later period than previously suggested. All stomatostylet-bearing
nematodes are proposed to have originated ~305 mya, corresponding to the transition from the

Devonian to the Permian period. The genus *Thornia* is placed outside of Dorylaimina and

Nygolaimina, disagreeing with its position in previous studies. Additionally, we tested the

whole genome amplification method and demonstrated that it is a promising strategy for

obtaining sufficient DNA for phylogenomic studies of microscopic eukaryotes. This study

significantly expanded the current nematode genome dataset, and the well-resolved phylogeny

enhances our understanding of the evolution of Nematoda.

Keywords:

Nematodes, worm, phylogeny, genome sequencing, whole genome amplification, diversity

Introduction

The phylum Nematoda represents one of the most cosmopolitan, abundant, and diverse

metazoans on Earth, with an estimated number of species extending to 10 million (Lambshead,

1993) (Figure 1). Free-living nematodes occupy all trophic levels in the food web and play a

pivotal role in carbon cycling and energy flows (Ferris, 2010; Ingham et al., 1985; van den

Hoogen et al., 2019), and the genera *Caenorhabditis* and *Pristionchus* are two popular

nematode models in molecular and evolutionary biology. Nematode parasites are a threat to not

only the health of plants (e.g. *Meloidogyne* spp., *Heterodera* spp.), but also that of animals and

humans (*e.g.* ascariasis, trichuriasis, and hookworm disease, filariasis). Resolving nematode

phylogeny is an important step in understanding the origin of parasitism and transition of

lifestyles, from basic evolutionary biology to pathogen control and development of anthelmintic

drugs (International Helminth Genomes Consortium, 2019). However, the high phenotypic

plasticity (Coomans, 2002), a near-complete lack of synapomorphic diagnostic characters, and

the absence of an informative fossil record have prevented scientists from deriving a consistent

evolutionary framework (Lorenzen, 1981; Coomans, 2000). The first molecular phylogeny was

reconstructed based on 18S rRNA of 53 species and led to the recognition of three major

lineages of nematodes: Dorylaimia, Enoplia, and Chromadoria (Blaxter et al.,1998; De Ley and

Blaxter et al., 2002). However, despite its popularity as a molecular marker, 18S rRNA lacks

the resolving power for deep phylogenies, i.e. the root of the nematode tree (Blaxter and

Koutsovoulos, 2015). As a result, currently all published rRNA phylogenies lack supporting

evidence for the "backbone" of the tree, and many lineages remain poorly resolved (Ahmed

and Holovachov, 2021).

By contrast, phylogenomic analyses have yielded statistically-robust and congruent results that have proven useful in resolving previously controversial phylogenetic relationships across evolutionary lineages (Misof et al., 2014; Telford et al., 2015; Williams et al., 2020). Advances in sequencing technology have also enabled the generation of inexpensive low-coverage genomes for phylogenomic studies (Zhang et al. 2019, Kim et al. 2024). However, difficulties in obtaining high-quality DNA have remained a major challenge for -omics studies on uncultured microscopic organisms. In the case of nematodes, pioneering work has been limited to studies on vertebrate parasitic taxa and model *Caenorhabditis* spp. (Blaxter and Koutsovoulos, 2015; International Helminth Genomes Consortium, 2019; Stevens et al., 2019), with little or no representation for species less than 1mm in length as well as for uncultured species. Recent advances in whole-genome amplification (WGA) have enabled single cells to generate sufficient DNA for sequencing, and these techniques have now been applied to various microscopic eukaryotic (Lepere et al.2011; Weisz et al., 2019; Sahraei et al., 2022; Lee et al., 2023), making it possible to sequence nematodes from a single individual at a phylum scale. The current paper focuses on sequencing the low-coverage genomes of 60 nematode species covering eight orders using WGA, aiming to provide finer resolution for phylogeny and improved knowledge about patterns and processes of nematode evolution.

Figure 1. Diversity of free-living and parasitic species in Nematoda. 1-27: the head region, 28-35: body habitus. 1, 40:*Acrobeloides*; 2: *Acrobeles*; 3: *Bicirronema*, 4: *Teratocephalus*; 5: *Tricirronema*; 6: *Pristionchus*; 7: *Steinernema*; 8: *Prismatolaimus*, 9: *Aphanolaimus*, 10: *Alaimus*; 11: *Aphelenchoides*; 12: *Diphtherophora*; 13: *Tripyla*; 14: *Miconchus*; 15: *Mylonchulus*; 16: *Ironus*; 17: *Aporcella*; 18, 38: *Helicotylenchus*; 19, 34: *Meloidogyne*; 20: *Trichodorus*; 21: *Sphaerolaimus*; 22, 35: *Greeffiella*; 23: *Eudorylaimus*; 24: *Oesophagostomum*; 25: *Ancylostoma*; 26: *Strongylus*; 27:

88 *Bunostomum*; 28: *Croconema*; 29: *Tylenchulus*; 30: Draconematidae; 31, 37: *Desmoscolex*; 32, 36:

89 *Tricoma*; 33: *Heterodera*; 39: *Etamphidelus*; 41: *Malenchus*. 7: insect parasite; 18-20, 29, 33, 34, 38:

90 plant parasites; 24-27: vertebrate parasites; 1-6, 8-17, 23, 39-41: terrestrial free-living species; 21, 22,

91 28, 30-32, 35-37 marine free-living species.

Material & Methods

DNA extraction, sequencing, and genome assembly

Nematodes were isolated from soil or marine sediments and extracted using a Baermann

funnel. One to five adult fresh worms were mounted on temporary slide and morphologically

identified under an Olympus BX51 microscope (Olympus Optical, Japan). The coverslip was

then removed, the worms were cut into pieces with a sterilized scalpel, and transferred with a

needle into a 200 μl PCR tube. Genomic DNA extraction and amplification were performed

with the REPLI-g Single Cell Kit (Qiagen, Germany) using 1–12 individual alive nematodes.

In addition to the whole genome amplification (WGA) method, DNA was directly extracted

from 20,000 fresh individuals for three species (*Aphelenchoides blastopthorus*, *Aphelenchoides*

smolae and *Ditylenchus* sp.) using the Ezup Column Animal Genomic DNA Purification Kit

(Sangon Biotech, China), in order to evaluate the performance of WGA. The TruSeq DNA

Sample Prep Kit (Illumina Inc. CA, USA) was used for library preparation according to the

manufacturer's instructions. Sequencing was preformed using the Illumina NovaSeq platform.

Quality control was performed in FASTP (Chen et al., 2018), reads less than 50 bp, Q-score

less than 20, or reads with N base number > 3 were removed. Filtered reads were assembled

using SPAdes v3.15.5 with the default parameters (Bankevich et al. 2012). We ran BlobToolKit

(Challis et al., 2020) on raw assemblies to manually screen for scaffolds derived from non-

target organisms. To avoid potential noise, we removed fragments with total length < 3k and

coverage < 15. The remaining sequences were subjected to a microbial contamination

prediction analysis using Kraken2 (Wood et al., 2019) to identify and eliminate any fragments

stemming from microbial contaminants. The raw sequences generated in this study were

submitted to GenBank SRA database with accession number PRJNA1128552.

Phylogenomic analysis

116   The phylogenetic hypothesis was reconstructed using newly generated data supplemented

117 with additional genomes and transcriptomes downloaded from GenBank (see Table S1), and a

118 total of 156 Nematoda and ten outgroups (two of each from Phyla Nematomorpha, Tardigrada,

119 Arthropoda, Onychophora, and Priapulida) were included. The complete and single copy

120 ortholog genes were extracted using BUSCO (Simão et al. 2015) using the Nematode ortholog

database OrthoDB v10 2024-01-08, and translated into amino acids (AA). We selected a 50% (USCO50, 1365 loci) and 70% complete (USCO70, 191 loci) matrices using custom script *matrix_generation.sh* from Du et al. (2023) for downstream analyses to balance between the number of loci and data completeness. The matrices were aligned using the L-INS-I option in MAFFT v7.490 (Katoh and Toh, 2008), trimmed using TrimAl (Capella-Gutiérrez, et al. 2009), and concatenated using FASconCAT-G v1.05.1 (Kück, 2014). We further removed potentially misaligned regions using SpruceUp (Borowiec, 2019) at 97% lognormal criterion. The remaining loci were further filtered using custom script *loci_filtering_alignment-based.sh* provided by Du et al. (2023) wrapper for PhyKIT v1.11.10 (Steenwyk et al., 2021) to remove loci with <100% parsimony-informative sites (Shen et al., 2016); loci with <0.35 relative composition variability (Phillips and Penny, 2003); and loci failing symmetry test (*p*-value 0.01-0.1) for stationarity and homogeneity (Naser-Khdour et al., 2019). The summary statistics of the final filtered matrices were calculated using AMAS v0.98 (Borowiec, 2016).

The following analyses were performed on both matrices using IQ-TREE v2.2.0 (Minh et al., 2020a): 1) Partitioned by locus with best model selected using ModelFinder (Kalyaanamoorthy et al. 2017), and with ultrafast bootstrap (Hoang et al. 2017) and SH-like approximate likelihood ratio test (SH-aLRT, Guindon et al. 2010) as support values; 2) Individual gene trees with best model selected using ModelFinder; 3) Gene and site concordance factors (gCF, sCF, respectively) using steps 1 and 2 following (Minh et al. 2020b); 4) GHOST model to account for heterotachy (Crotty et al. 2019); and finally, 5) the posterior mean site frequency model (PMSF) which has been designed to correct for long-branch-attraction by modeling site heterogeneity (Wang et al. 2018), the C20 model was only used for USCO50 matrix due to the computational limits required for such a large dataset, whereas the more complex C60 model was used for the USCO70 matrix.

We also analysed both datasets using the multispecies coalescent-based method (MSC) implemented in ASTRAL-III (Zhang et al. 2018), using gene trees generated in step 2 above as input. We collapsed the gene tree nodes with <10% support using Newick Utilities (Junier, 2010), and support was evaluated using local posterior probability (LPP, Sayyari and Mirarab, 2016). Detailed scripts and flags for both IQ-TREE and ASTRAL can be found in Table S2.

Molecular dating and ancestral state reconstruction

Divergence dating was performed using MCMCTree using PAML v4.9j (Yang, 2007) using a subset of the PMSF_C20 topology as guide tree and the USCO50 matrix, filtered using

the custom script *loci_filtering_tree-based.sh* provided by Du et al. (2023) based on the gene
trees built in step 2 above with the following: average bootstrap support of >80 (Salichos and
Rokas, 2013); signal-to-noise ratio (treeness/relative composition variability) of 1.5 (Phillips
and Penny, 2003); and degree of violation of the molecular clock of 0.2 (Liu et al., 2017); to
obtain a final subset matrix of 135 loci. Approximate likelihood calculation and ML estimation
of branch lengths to reduce the computational burden by calculating the Hessian matrices using
the LG substitution model and the independent rates clock mode. Given the controversial
relationships among Cryptovermes (Nematoida and Panarthropoda) within Ecdysozoa is
beyond the scope of this study, we chose not to include fossil outgroups from these lineages.
Instead we used fossil age from Phylum Priapulida dated at 518 mya as a calibration point for
Scalidophora which is the sister to Cryptovermes, and a root age of 636 mya following Howard
et al. (2022). Seven nematode fossils plus one Priapulida fossil were selected for calibration
based on existing fossil records (Table S3).

Two independent runs were performed with 50,000 generation burnins and 500,000
generations using the HKY85 model, and convergence was assessed using Tracer v1.7
(Rambaut et al. 2018). The tree was visualized using MCMCtreeR (Puttick, 2019) in R v4.2.0
(R Core Team 2022).

Results

Whole genome amplification and genome assembly

A total of 60 species from terrestrial and marine environments were newly sequenced. The
raw assemblies ranged in size from 43.67 Mb to 839.99 Mb, comprising contigs spanning from
15.99 Kb to 2.69 Mb. Genome sizes decreased after removing potential contaminations, with
lengths ranging from 2.90 MB to 189.63 Mb, and contig counts varying from 484 to 33537
(Table S4). The BUSCO analysis yielded completeness percentages ranging from 7.65% to
81.86% for the assembled genomes. We observed smaller size and more fragmented genomes
(higher contig numbers) when using the WGA method compared to direct DNA extraction
without amplification from fresh nematodes (Figures S1, S2). Further BUSCO evaluation
indicated varying performance results, *i.e.* both methods generated a similar genome
completeness in *Ditylenchus* sp., a more complete *A. blastopthorus* and *A. smolae* genome with
direct extraction and WGA, respectively (Figure S3). The assembled contigs were annotated
and plotted according to their GC content and coverage (Figure S4), revealing that direct
extraction introduced more bacterial contaminations than the WGA approach.

Evolution of Nematoda

From the assembled genomes, we extracted 1028 AA loci with 304,894 bp for the USCO50 matrix, and 115 loci with 23,069 bp for the USCO70 matrix. In general, the topologies inferred from USCO70 matrix were less supported when compared to those from the USCO50 matrix. The reconstructed phylogenies have similar topologies in most cases regardless of the metrics, datasets, and reconstruction methods (Figure 2). Nematoda is either sister to Nematomorpha (PMSF) or to a clade including Nematomorpha and Tardigrada (MSC and GHOST). Within Nematoda, the differences between topologies are summarised in the table of Figure 2. The subclass Enoplia primarily consists of taxa inhabiting marine environments. Placing the class Enoplia as the sister taxon to the rest of Nematoda in our analysis suggests a likely marine origin for the nematode ancestor. This subclass comprises two orders, Enoplida and Triplonchida. Notably, the representation of taxa in the latter group has significantly improved compared to transcriptomic analyses (Smythe et al. 2019; Ahmed et al. 2022). The monophyly of Triplonchida is well-supported. Enoplida is paraphyletic due to *Ironus dentifurcatus* being part of a lineage that is sister to the rest of Enoplida. This is consistent with the previous finding of Ahmed et al. (2022) who recovered Enoplida as paraphyletic, with *Bathylaimus* sp. (Triplonchida) and *Tobrilus* sp. (Enoplida) forming a sister branch to the rest of Enoplida.

There are five orders within Dorylaimia (clade I in Blaxter et al.,1998) represented in this analysis, all of which are terrestrial. A lineage consisting of the vertebrate parasitic orders Trichinellida and Dioctophymatida forms a sister branch with a clade comprising Dorylaimida, Mononchida and Mermithida. The orders Dorylaimida, Mononchida and Trichinellida, which are represented by more than one taxon, all form well-supported monophyletic groups, consistent with previous 18S rRNA-based and phylogenomic analyses (van Megen et al. 2009; Smythe et al. 2019; Ahmed et al. 2022). Apart from well-represented Rhabditida, four early-diverging Chromadoria orders are represented in this study, Chromadorida, Araeolaimida, Monhysterida and Plectida, with the first three being mostly marine. Within Rhabditida, all three suborders are strongly supported. The three infraorders representing Spirurina also received robust support. The Diplogasteromorpha within Rhabditina was likewise well-supported, but Rhabditomorpha was paraphyletic with genus *Poikilolaimus* being sister lineage to the clade of Diplogasteromorpha + Rhabditomorpha. The Bunonematomorpha was only represented by one genus *Bunonema*, and was placed either as sister taxa to all other members of Rhabditina (most cases) or to *Poikilolaimus* (GHOST model in IQ-TREE). Even though the analysis accounted for all three infraorders of Tylenchina, the majority of the taxa belonged to the infraorder Tylenchomorpha. In congruence with previous 18S rRNA and transcriptomic

analyses of Tylenchina, Strongyloidoidea, one of the superfamilies of Panagrolaimomorpha, diverged first from the rest of Tylenchina, with two separate branching events–first with Steinernematidae, and then the Alloionematidae + Strongyloididae lineage. The remaining crown members of the tree formed a well-supported branch, within which Panagrolaimidae, another family of Panagrolaimomorpha, nested. Consistent with other phylogenomic studies, but contradictory to 18S rRNA data, Tylenchomorpha was found to be monophyletic, supporting the single evolutionary origin of the stomatostylet (the tylenchid and aphelenchid feeding apparatus), although the superfamily Aphelenchoidea does not form a monophyletic clade within Tylenchomorpha.

Figure 2. Phylogeny of Nematoda inferred from amino acids matrix USCO50 implemented in IQ-TREE. Node labels show ultrafast bootstrap approximation (UFBoot)/gene concordance factor (gCFs)/site concordance factor (sCFs). Node supports from other analyses are given in the table at left top corner, with the corresponding node numbers indicated by the coloured squares at tree node. The slash in table suggests the topology is not supported. The nodes without corresponding nodes number suggest they were fully supported in all analyses. The newly sequenced species were indicated with star.

Molecular dating of Nematoda

The divergence time dating, based on the ten calibrated nodes (Figure S5), traces the divergence date for Nematoda from its sister group Nematomorpha during late Ediacaran to Ordovician period (620–455 mya) (Table 1). Within Nematoda, the most recent common ancestor of all Enoplia dated back to a period spanning the end of the Ediacaran to the mid-Devonian (573–378 mya), slightly later than the common ancestor of all Chromadoria (561–408 mya) in mean highest posterior distribution (HPD, 473 *vs* 482 mya). All Dorylaimia had their most recent common ancestor ~446 mya, during the beginning Cambrian through to the end of Devonian. Our analysis indicates that human parasite *Trichinella* and *Trichuris* shared a common ancestor ~202 mya, which is a considerably more recent compared ~275 mya and ~281 mya, respectively reported in Zarlenga et al. (2006) and Korhonen et al. (2016). The most recent common ancestor of all stomatostylet-bearing nematodes is suggested to have existed ~305 mya, coinciding with the transition from the Devonian to end of the Permian period.

Table 1. The Nematoda divergence time estimation performed by using MCMCTree on the amino acids matrix USCO50 PMSF C20 topology.

Clade	Time estimations (mya)		
	Mean HPD	95% Lower HPD	95% Upper HPD
Nematoda	531	455	620
Chromadoria	482	408	561
Spirurina (Clade III)	323	259	388
Spiruromorpha	261	198	322
Ascaridomorpha	228	168	295
Oxyuridomorpha	148	66	224
Tylenchina (Clade IV)	361	302	427
Cephalobomorpha	105	45	178
Tylenchomorpha	305	254	364
Rhabditina (Clade V)	336	277	403
Diplogasteromorpha	240	185	296
Strongyloidea	120	90	154
Plectida	294	181	405
Linhomoeina	291	193	399
Monhysterina	325	238	413
Chromadorida	348	242	454

Desmodorida	345	238	456
Dorylaimia (Clade I)	446	367	537
Dorylaimida	290	219	363
Mononchida	210	148	278
Trichinellida	202	120	292
Enoplia (Clade II)	473	378	573
Enoplida	403	305	501
Triplonchida	304	194	420

Discussion

Evolution of early-branching Nematoda

The general topology of the tree agrees well with the results of previously-published
analyses obtained using mostly transcriptomic datasets for different sets of species (Smythe et
al., 2019, Ahmed et al., 2022), with Enoplia (Clade II of Blaxter et al., 1998) diverging first
from the rest of the nematodes, and Dorylaimia (Clade I of Blaxter et al., 1998) being close
second, in full agreement with the scarce morphological and embryological evidence
summarised by Holterman et al. (2006). Within Enoplia, all included species are represented
by newly sequenced genomes, and although none of the taxa are identical to earlier datasets,
the resultant topology agrees in basic details with the transcriptome-only phylogenies (Smythe
et al., 2019, Ahmed et al., 2022). However, the bootstrap support of several nodes within
Enoplida is suboptimal: Triplonchida is resolved as monophyletic, while Enoplida as
paraphyletic. This agrees with morphological evidence for the monophyly of Triplonchida
suggested by De Ley and Blaxter (2002) but contradicts the phylogenies based on 18S rRNA
gene (summarised in Ahmed and Holovachov, 2021). Enoplia (Clade II) is one of the most
morphologically diverse groups of nematodes, so there are no easily testable morphology-based
phylogenies. Enoplia are also likely to comprise the first nematode lineages that made the
transition from marine to freshwater habitats in the Ediacaran-Devonian periods several times
independently (Table 1). Future sampling of additional genomes from all lineages within
Enoplia will not only help resolve relationships within the group but will also shed light on how
colonisation of freshwater habitats took place in the early Palaeozoic.

The tree topology within Dorylaimia (Clade I) not only agrees fully with previously
published datasets (Smythe et al., 2019, Ahmed et al., 2022), but a considerably expanded
dataset of Mononchida and especially Dorylaimida also allows further comparison with
morphology- and rRNA-based phylogenies. Within Dorylaimida, the genus *Thornia* branches
off first, which disagrees with morphological evidence (Peña-Santiago, 2014). The sole

available 18S rRNA sequence of *Thornia* is highly divergent and its position is therefore
unreliable (Fig. S1. in Holterman et al. 2019). The remaining phylogeny of Dorylaimida is more
in agreement with previous studies: Nygolaimina (represented here by *Paravulvus* and
Aetholaimus) and Dorylaimina (all other taxa) are sister clades. The two families represented
by more than one species are Actinolaimidae (*Paractinolaimus* and *Trachactinolaimus*) and
Qudsianematidae (*Eudorylaimus* and *Discolaimus*), are herein resolved as monophyletic and
polyphyletic respectively, in agreement with previous rRNA-based analyses (summarised in
Ahmed and Holovachov, 2021). The entire Clade I is represented by limno-terrestrial
(Mononchida and Dorylaimidae) and parasitic (Trichinellida, Dioctophymatida and
Mermithida) species. Of the latter, only Trichinellida includes considerable number of parasites
inhabiting marine vertebrate species. Molecular dating suggests that the Clade I diverged off
from the rest of Nematoda in the Cambrian-Devonian, but the phylogeny of the group is not
detailed enough to hypothesize about the transition between marine, freshwater, and terrestrial
habitats within this lineage. Mermithida, parasitoids of insects and other (mostly) terrestrial
arthropods are represented by single species in the current analysis; molecular dating suggests
a very broad timeframe for the origin of the clade, spanning from Devonian to Triassic, much
later than the period in which their arthropod hosts have diversified on land. Similarly, the plant-
parasitic lineage within the Clade I is represented by a single species, *Longidorus juglans*, and
molecular dating suggests it split off from non-parasitic dorylaims during the Permian-Jurassic,
at about the same time as their hosts, flowering plants. However, more data on the evolution of
both hosts and parasites is necessary to make in-depth cophylogenetic study.

Evolution of plant-parasitic Tylenchina

Taxa closely related to plant-parasitic lineages have long been a focal point of interest in
understanding the origins of plant parasitism (Bert et al., 2008; Ahmed et al., 2022). Most
rRNA-based phylogenies indicate that Cephalobomorpha shares the most recent common
ancestry with Tylenchomorpha (Blaxter et al., 1998; Holterman et al., 2006; Bert et al., 2008;
van Megen et al., 2009). However, our analyses are consistent with the analyses of Ahmed et
al. (2022), which positions both Cephalobomorpha and Panagrolaimomorpha as sister taxa to
Tylenchomorpha. The taxonomic position of aphelenchs (Aphelenchoidea: Aphelenchoididae
+ Aphelenchidae) relative to the tylenchs (Tylenchomorpha without aphelenchs) has been a
subject of especially extensive debate. Despite the evident morphological similarities between
Aphelenchidae and Aphelenchoididae, the latter has been placed outside the Tylenchomorpha
based on rRNA-based phylogenies. They were separated from Aphelenchidae, which were

found to be sister to the remaining members of Tylenchomorpha (van Megen et al., 2009). This placement was hypothesized as resulting from long branch attraction and/or elevated AT-content (De Ley and Blaxter, 2002; Holterman et al., 2006) and their monophyletic grouping could not be significantly rejected (Bert et al., 2008). In contrast, mitogenome trees have consistently shown aphelenchs as monophyletic and positioned away from other Tylenchomorpha (Kern et al., 2020). This analysis supports the hypothesis that is most plausible based on morphology, namely that taxa equipped with the stomatostylet form a monophyletic group, a supposition that is congruent with the findings of Ahmed et al. (2022). Nonetheless, in our analysis aphelenchs are still depicted as paraphyletic and Aphelenchidae are, similar to rRNA-based phylogenies, more closely related to tylenchs than to the morphologically more similar Aphelenchoidea.

Limitations of molecular dating of nematode phylogeny

Although the number of known nematode fossils is much larger than for some other soft-bodied invertebrate phyla (Poinar et al., 2011), their usefulness for molecular dating is limited. The large majority of known nematode fossils originate from various fossilised resins (amber and copal), are relatively recent and, due to the nature of the fossilisation, are strongly biased towards insect-associated and terrestrial nematodes. The next most common category also includes relatively recent fossils of vertebrate parasites found in coprolites and mummified remains. Only a few fossils date back to Palaeozoic, during which the majority of nematode lineages are thought to have diverged. Comparing the dates of origin of parasitic nematodes and their respective hosts we find both agreements and disagreements. The possible date of origin of Longidoridae, which is parasite of flowering plants, matches that of their hosts during the Jurassic-Cretaceous period, while plant-parasitic Tylenchomorpha has an earlier origin (Devonian to the Permian) than its hosts. The earliest possible divergence date of plant-parasitic Triplonchida, represented in our analysis by only *Trichodorus pakistanensis* (Carboniferous to early Cretaceous) predates the origin of their hosts, flowering plants (van der Kooi and Ollerton, 2020). Similarly, the possible date of divergence of animal parasitic Ascaridomorpha is underestimated in our analysis (early Permian to Jurassic) compared to cophylogenetic study using fossil data from their vertebrate hosts (Carboniferous) (Li et al., 2018). The dating of the split between *Trichinella* and *Trichuris* is also inconsistent between our and previous analyses (Zarlenga et al, 2006; Korhonen et al. 2016). These examples not only expose the limitations of the fossils when dating the nematode phylogeny, but highlight the need for a

multidisciplinary approach, combining fossil evidence, co-phylogenetic studies, and mutation accumulation rates.

Whole genome amplification and microscopic eukaryotes phylogenomic

Microscopic eukaryotes, such as Protozoa, Nematoda, Platyhelminthes, and Gastrotricha, are abundant, diverse and fill critical ecological roles across every ecosystem on Earth, and yet they were poorly known across subjects ranging from biodiversity and, evolution, to genomics (Wardle, 2006; Bik et al., 2012). The lack of omics studies in these groups can partly be explained by their small body size and low DNA quantity. Indeed, the high-throughput sequencing protocols require large amounts of input DNA (*e.g.* several micrograms), typically thousands of times more than that which is contained within a single microscopic individual. Zhang et al. (2019) proposed WGA in genome sequencing and phylogenomic of small animals, and three pioneering studies have used this method based on Illumina and PacBio platform to generate nematodes genomes (Roberts et al., 2024; Lee et al., 2023; Doyle et al., 2019), although no study had yet evaluated its performance using microscopic species. In this study, we demonstrated that WGA generated smaller assembly and more fragmented contigs, probably due to the bias in the amplification process (Sabina et al., 2015). Albeit reduced, the quality of results remains high enough to allow certain types of studies, *e.g.* our nematode phylogenomic analysis. As a result, our study supports WGA as being a powerful tool in phylogenomic, population genomic and metagenomic studies of microscopic eukaryotes.

Acknowledgements: X.Q. was funded by National Natural Science Foundation of China (32001876). Y.M.Z. was funded by the Oak Ridge Institute for Science and Education (ORISE) fellowship during part of this work. We thank Dr. Wilfrida Decraemer and Laboratory of Parasitology in Ghent University for providing nematode images, Dr. Kan Zhuo and Guokun Liu for providing nematode samples. We thank the Smithsonian Institution High Performance Cluster (https://doi.org/10.25572/SIHPC) for providing computational resources and support that have contributed to the research results reported in this publication.

Supplementary material:

Data available from the Dryad Digital Repository: https://doi.org/10.5061/dryad.b2rbnzsnt.
The raw genomes available in NCBI with accession number PRJNA1128552.

References

Ahmed M., Roberts N.G., Adediran F., Smythe A.B., Kocot K.M., Holovachov O. 2022. Phylogenomic analysis of the phylum Nematoda: conflicts and congruences with morphology, 18S rRNA, and mitogenomes. Front. Ecol. Evol. 9:769565.

Ahmed M., Holovachov O. 2021. Twenty years after De Ley and Blaxter—how far did we progress in understanding the phylogeny of the phylum Nematoda?. Animals 11:3479.

Bankevich A., Nurk S., Antipov D., Gurevich A.A., Dvorkin M., Kulikov A.S., Lesin V.M., Nikolenko S.I., Pham S., Prjibelski A.D. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J. Comput. Biol. 19:455–477.

Bert W., Leliaert F., Vierstraete A.R., Vanfleteren J.R., Borgonie G. 2008. Molecular phylogeny of the Tylenchina and evolution of the female gonoduct (Nematoda: Rhabditida). Mol. Phylogenet. Evol. 48:728–744.

Bik H.M., Porazinska D.L., Creer S., Caporaso J.G., Knight R., Thomas W.K. 2012. Sequencing our way towards understanding global eukaryotic biodiversity. Trends Ecol. Evol. 27:233–243.

Blaxter M., De Ley P., Garey J.R., Liu L.X., Scheldeman P., Vierstraete A., Vanfleteren J.R., Mackey L.Y., Dorris M., Frisse L.M., Vida J.T. 1998. A molecular evolutionary framework for the phylum Nematoda. Nature 392:71–75.

Blaxter M., Koutsovoulos G. 2015. The evolution of parasitism in Nematoda. Parasitology 142:S26–S39.

Borowiec M. L. (2016). AMAS: A fast tool for alignment manipulation and computing of summary statistics. PeerJ. 4: e1660.

Borowiec M.L. 2019. Spruceup: fast and flexible identification, visualization, and removal of outliers from large multiple sequence alignments. J. Open Source Softw. 4:1635.

Capella-Gutiérrez S., Silla-Martínez J.M., Gabaldón T. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972–1973.

Challis R., Richards E., Rajan J., Cochrane G., Blaxter M. 2020. BlobToolKit–interactive quality assessment of genome assemblies. G3: Genes, Genomes, Genetics 10:1361–1374.

Chen, M.Y., Liang, D., Zhang, P. 2017. Phylogenomic resolution of the phylogeny of laurasiatherian mammals: Exploring phylogenetic signals within coding and noncoding sequences. Genome Biol. Evol. 9(8): 1998–2012.

Chen S., Zhou Y., Chen Y., Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics 34: i884–i890.

Coomans A. 2000. Nematode systematics: past, present and future. Nematology 2:3–7.

Coomans A. 2002. Present status and future of nematode systematics. Nematology 4:573–582.

Crotty S.M., Minh B.Q., Bean N.G., Holland B.R., Tuke J., Jermiin L.S., von Haeseler A. 2020. GHOST: Recovering Historical Signal from Heterotachously Evolved Sequence Alignments. Syst. Biol. 69:249–264.

De Ley P., Blaxter M.L. 2002. Systematic position and phylogeny. In: Lee D.L., editor. The biology of nematodes. London: CRC Press, p. 1–30.

Doyle S.R., Sankaranarayanan G., Allan F., Berger D., Jimenez Castro P.D., Collins J.B., Crellen T., Duque-Correa M.A., Ellis P., Jaleta T.G., Laing R., Maitland K., McCarthy C., Moundai T., Softley B., Thiele E., Ouakou P.T., Tushabe J.V., Webster J.P., Weiss A.J., Lok J., Devaney E., Kaplan R.M., Cotton J.A., Berriman M., Holroyd N. 2019. Evaluation of DNA extraction methods on individual helminth egg and larval stages for whole-genome sequencing. Front. Genet. 10:826.

Du S., Ding Y., Li H., Zhang A., Luo A., Zhu C., Zhang, F. 2023. Construction of a phylogenetic matrix: Scripts and guidelines for phylogenomics. Zool. Syst. 460(48): 107–116.

Ferris H. 2010. Contribution of nematodes to the structure and function of the soil food web. J
Nematology, 42:63–67.
Guindon S., Dufayard J-F., Lefort V., Anisimova M., Hordijk W., Gascuel O. 2010. New
algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
performance of PhyML 3.0. Syst. Biol. 59: 307–321.
Hoang D.T., Chernomor O., von Haeseler A., Minh B.Q., Vinh L.S. 2018. UFBoot2: Improving
the Ultrafast Bootstrap Approximation. Mol. Biol. Evol. 35:518–522.
Holterman M., van der Wurff A., van den Elsen S., van Megen H., Bongers T., Holovachov O.,
Bakker J., Helder J. 2006. Phylum-wide analysis of SSU rDNA reveals deep phylogenetic
relationships among nematodes and accelerated evolution toward crown clades. Mol. Biol.
Evol. 23:1792–800.
Holterman M., Schratzberger M., Helder J. 2019. Nematodes as evolutionary commuters
between marine, freshwater and terrestrial habitats. Biol. J. Linn. Soc. 128:756–767.
Howard R.J., Giacomelli M., Lozano-Fernandez J., Edgecombe G.D., Fleming J.F., Kristensen
R.M., Ma X., Olesen J., Sørensen M.V., Thomsen P.F., Wills M.A., 2022. The Ediacaran
origin of Ecdysozoa: integrating fossil and phylogenomic data. Journal of the Geological
Society 179(4): jgs2021-107.
Ingham R.E., Trofymow J.A., Ingham E.R., Coleman D.C. 1985. Interactions of bacteria, fungi,
and their nematode grazers: effects on nutrient cycling and plant growth. Ecol.
Monographs 55:119–140.
International Helminth Genomes Consortium. 2019. Comparative genomics of the major
parasitic worms. Nat. Genet. 51:163–174.
Junier T., Zdobnov E.M. 2010. The Newick utilities: high-throughput phylogenetic tree
processing in the UNIX shell. Bioinformatics 26:1669–1670.
Kalyaanamoorthy S., Minh B.Q., Wong T.K., von Haeseler A., Jermiin L.S. 2017. ModelFinder:
fast model selection for accurate phylogenetic estimates. Nat. Methods. 14:587–589.
Katoh K., Toh H. 2008. Recent developments in the MAFFT multiple sequence alignment
program. Brief Bioinform. 9:286–298.
Kern E.M.A., Kim T., Park J.-K. 2020. The mitochondrial genome in nematode
phylogenetics. Front. Ecol. Evol. 8:250.
Kim B.Y., Gellert H.R., Church S.H., Suvorov A., Anderson S.S., Barmina O., Beskid S.G.,
Comeault A.A., Crown K.N., Diamond S.E., Dorus, S., 2024. Single-fly genome
assemblies fill major phylogenomic gaps across the Drosophilidae Tree of Life. PLoS
Biol. 22(7): e3002697.
Korhonen P.K., Pozio E., La Rosa G., Chang B.C., Koehler A.V., Hoberg E.P., Boag P.R., Tan
P., Jex A.R., Hofmann A., Sternberg P.W., Young N.D., Gasser R.B. 2016. Phylogenomic
and biogeographic reconstruction of the *Trichinella* complex. Nat. Commun. 7:10513.
Kück P., Longo G.C. 2014. FASconCAT-G: extensive functions for multiple sequence
alignment preparations concerning phylogenetic studies. Front. Zool. 11:81.
Lambshead P.J.D. 1993. Recent developments in marine benthic biodiversity research.
Oceanis 19:5–24.
Lee Y-C., Ke H-M., Liu Y-C., Lee H-H., Wang M-C., Tseng Y-C., Kikuchi T., Tsai I.J. 2023.
Single-worm long-read sequencing reveals genome diversity in free-living nematodes.
Nucl. Acids Res. 51:8035–8047.
Lepere C., Demura M., Kawachi M., Romac S., Probert I., Vaulot D. 2011. Whole-genome
amplification (WGA) of marine photosynthetic eukaryote populations. FEMS Microbiol.
Ecol. 76:513–523.
Li L., Lü L., Nadler S.A., Gibson D.I., Zhang L.-P., Chen H.-X., Zhao W.-T., Guo Y.-N. 2018.
Molecular phylogeny and dating reveal a terrestrial origin in the early carboniferous for
ascaridoid nematodes. Syst. Biol. 67:888–900.

Liu, L., Zhang, J., Rheindt, F.E., Lei, F., Qu, Y., Wang, Y., Zhang, Y., Sullivan, C., Nie, W., Wang, J., Yang, F., Chen, J., Edwards, S.V., Meng, J., Wu, S. 2017. Genomic evidence reveals a radiation of placental mammals uninterrupted by the KPg boundary. Proc. Natl. Acad. Sci. U. S. A. 114(35): E7282–E7290.

Lorenzen S. 1981. Entwurf eines phylogenetischen systems der freilebenden Nematoden (Outline of a phylogenetic system for the free-living nematodes). Veröff. Inst. Meeresforsch. Bremerh. Suppl. 7, 1–472.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., von Haeseler A., Lanfear R. 2020a. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the genomic era. Mol. Biol. Evol. 37:1530–1534.

Minh B.Q., Hahn M.W., Lanfear R. 2020b. New methods to calculate concordance factors for phylogenomic datasets. Mol. Biol. Evol. 37:2727–2733.

Misof B., Liu S., Meusemann K., Peters R.S., Donath A., Mayer C., Frandsen P.B., Ware J., Flouri T., Beutel R.G. Niehuis O. 2014. Phylogenomics resolves the timing and pattern of insect evolution. Science 346:763–767.

Naser-Khdour S., Minh B.Q., Zhang W., Stone E.A., Lanfear R. 2019. The prevalence and impact of model violations in phylogenetic analysis. Genome Biol. Evol. 11:3341–3352.

Peña-Santiago R. 2014. Order Dorylaimida Pearse, 1942. In: Schmidt-Rhaesa A., editor. Nematoda. Berlin: De Gruyter, p. 277–298.

Phillips, M.J., Penny, D. 2003. The root of the mammalian tree inferred from whole mitochondrial genomes. Molecular Phylogenetics and Evolution 28(2): 171–185.

Poinar G.O. 2011. The evolutionary history of nematodes as revealed in stone, amber and mummies. Leiden, The Netherlands : Koninklijke Brill NV, 429 p.

Puttick M.N. 2019. MCMCtreeR: functions to prepare MCMCtree analyses and visualize posterior ages on trees. Bioinformatics 35:5321–5322.

R Core Team. 2022. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available online at https://www.R-project.org/.

Rambaut A., Drummond A.J., Xie D., Baele G., Suchard M.A. 2018. Posterior summarization in Bayesian phylogenetics using Tracer 1.7. Syst. Biol. 67:901–904.

Roberts N.G., Gilmore M.J., Struck T.H., Kocot K. 2024. Multiple displacement amplification facilitates SMRT sequencing of microscopic animals and the genome of the Gastrotrich *Lepidodermella squamata* (Dujardin, 1841). bioRxiv, pp.2024-01. https://doi.org/10.1101/2024.01.17.576123

Sabina J., Leamon J.H. 2015. Bias in whole genome amplification: causes and considerations. Methods Mol. Biol. 1347:15–41.

Sahraei S.E., Sanchez-Garcia M., Montoliu-Nerin M., Manyara D., Bergin C., Rosendahl S., Rosling A. 2022. Whole genome analyses based on single, field collected spores of the arbuscular mycorrhizal fungus *Funneliformis geosporum*. Mycorrhiza 32:361–371.

Salichos, L., Rokas, A. 2013. Inferring ancient divergences requires genes with strong phylogenetic signals. Nature, 497: 327–331.

Sayyari E., Mirarab S. 2016. Fast coalescent-based computation of local branch support from quartet frequencies. Mol. Biol. Evol. 33:1654–1668.

Shen, X-X., Salichos, L., Rokas, A. 2016. A genome-scale investigation of how sequence, function, and tree-based gene properties influence phylogenetic inference. Genome Biology and Evolution 8(8): 2565–2580.

Simão F.A., Waterhouse R.M., Ioannidis P., Kriventseva E.V., Zdobnov E.M. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. Bioinformatics 31:3210–3212.

Smythe A.B., Holovachov O., Kocot K.M. 2019. Improved phylogenomic sampling of free-
living nematodes enhances resolution of higher-level nematode phylogeny. BMC Evol.
Biol. 19:1–5.
Steenwyk, J.L., Buida III, T.J., Labella, A.L., Li, Y., Shen, X.X. and Rokas, A., 2021. PhyKIT:
a broadly applicable UNIX shell toolkit for processing and analyzing phylogenomic data.
Bioinformatics 37(16): 2325–2331.
Stevens L., Félix M.A., Beltran T., Braendle C., Caurcel C., Fausett S., Fitch D., Frézal L.,
Gosse C., Kaur T., Kiontke K., Newton M.D., Noble L.M., Richaud A., Rockman M.V.,
Sudhaus W., Blaxter M. 2019. Comparative genomics of 10 new *Caenorhabditis* species.
Evol. Lett. 3:217–236.
Telford M.J., Budd G.E., Philippe H. 2015. Phylogenomic insights into animal evolution. Curr.
Biol. 25:R876–R887.
van den Hoogen J., Geisen S., Routh D., Ferris H., Traunspurger W., Wardle D.A., De Goede
R.G., Adams B.J., Ahmad W., Andriuzzi W.S., Bardgett R.D. 2019. Soil nematode
abundance and functional group composition at a global scale. Nature 572:194–198.
van der Kooi C.J., Ollerton J. 2020. The origins of flowering plants and pollinators. Science
368:1306–1308.
van Megen H., van den Elsen. S., Holterman M., Karssen G., Mooyman P., Bongers T.,
Holovachov O., Bakker J., Helder J. 2009. A phylogenetic tree of nematodes based on
about 1200 full-length small subunit ribosomal DNA sequences. Nematology 11:927–950.
Wang H.C., Minh B.Q., Susko E., Roger A.J. 2018. Modeling site heterogeneity with posterior
mean site frequency profiles accelerates accurate phylogenomic estimation. Syst.
Biol. 67:216–235.
Wardle D.A. 2006. The influence of biotic interactions on soil biodiversity. Ecol. Lett. 9:870–
886.
Weisz F., Lalle M., Nohynkova E., Sannella A.R., Dluhošová J., Cacciò S.M. Testing the
impact of Whole Genome Amplification on genome comparison using the polyploid
flagellated *Giardia duodenalis* as a model. Exp. Parasitol. 207:107776.
Williams T.A., Cox C.J., Foster P.G., Szöllősi G.J., Embley T.M. 2020. Phylogenomics
provides robust support for a two-domains tree of life. Nat. Ecol. Evol. 4:138–147.
Wood D.E., Lu J., Langmead B. 2019. Improved metagenomic analysis with Kraken 2. Genome
Biol. 20:257.
Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol. Biol.
Evol. 24:1586–1591.
Zarlenga D.S., Rosenthal B.M., La Rosa G., Pozio E., Hoberg E.P. 2006. Post-Miocene
expansion, colonization, and host switching drove speciation among extant nematodes of
the archaic genus *Trichinella*. Proc. Natl. Acad. Sci. U. S. A. 103:7354–7359.
Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. BMC Bioinformatics 19:153.
Zhang F., Ding Y., Zhu C.D., Zhou X., Orr M.C., Scheu S., Luan Y.X. 2019. Phylogenomics
from low-coverage whole-genome sequencing. Methods Ecol. Evol. 10:507–517.