1 # Molecular Speciation of Cyathostomins From Horses In Ireland

2 Orla Byrne, Disha Gangotia, Annetta Zintl, Liam Kiser, Olivia Boxall, Daniel McSweeney, Fiona
3 O'Neill, Stacey Dunne, Breanna Rose Lamb, Nicola Walshe, Grace Mulcahy.

4 UCD School of Veterinary Medicine and Conway Institute, University College Dublin, Dublin,
5 Ireland.
6

7 Corresponding author: orla.byrne1@ucdconnect.ie;
8

9 Keywords: Equine; Helminth; Cyathostomin; Molecular Speciation; Ireland
10

11 Graphical Abstract:
12

13
14

Abstract:

Cyathostomins are globally important equine parasites, responsible for both chronic and
acute pathogenic effects. The occurrence of mixed infections with numerous cyathostomin
species hinders our understanding of parasite epidemiology, host-parasite dynamics and
species pathogenicity. There have been few studies of cyathostomin species occurring in

horses in Ireland, where temperate climatic conditions with year-round rainfall provides

suitable conditions for infection of grazing animals with bursate nematodes. Here, we

amplified and sequenced the ITS-2 region of adult worms harvested at post-mortem from 11

adult horses between August 2018 and June 2020, and recorded species prevalence and

abundance of worms recovered from the caecum, right ventral colon and left dorsal colon.

Seasonal distribution, phylogenetic relationships and community composition were also

recorded and compared with other relevant studies, including a global meta-analysis. Overall,

our results agree with previous research, in that there does not seem to be a major difference

in cyathostomin species occurrence in equids in different geographical regions. We confirmed

the results of other studies in relation to the difficulties in discriminating between

Cylicostephanus calicatus and *Cor. coronatus* on the basis of ITS-2 sequences. We found a

positive correlation between the occurrence of *Cyathostomum pateratum* and *Cyc. calicatus*

($p < 0.01$). *Cor. coronatus*, the most abundant species, was only recovered from horses

sampled in autumn. Although this study is limited in terms of sample size, it provides a

platform for furthering understanding of the epidemiology and pathogenesis of equine

cyathostomin infections in temperate climates.

1. Introduction

Cyathostomins are a group of parasitic helminths that inhabit the gastrointestinal tract. They

are obligate parasites which are ubiquitous in equids and of significant pathogenic concern

(Boxell *et al*., 2004; Ghafar *et al*., 2022; Hinney *et al*., 2011; Love *et al*., 1999; Morariu *et al*.,

2016; Steinbach *et al*., 2006). Currently, there are over 50 recognised species with 40

recorded as infecting horses and mixed infections are the norm (Lichtenfels *et al*., 2008; Lyons

et al., 1999). Cyathostomins have a direct life cycle, with infection occurring through ingestion

of infective stage (L3) larvae on pasture. Following infection, cyathostomin larvae penetrate

into the mucosa of the large intestine, and can remain here in a dormant (hypobiotic) state,

emerging to mature and establish patent infection in the lumen after a variable pre-patent

period of up to three years (Gibson, 1953; Nielsen and Lyons, 2017). The hypobiotic state is

relevant to anthelmintic efficacy and to cyathostomin-associated disease (Boisseau *et al*.,

2023; Morariu *et al*., 2016). Typically, of the total worm burden in an individual animal, the

majority will comprise hypobiotic larvae in the mucosa (Dowdall *et al*., 2002; Duncan *et al*.,

1998; Gibson, 1953; Matthews, 2008). The stimuli responsible for hypobiotic larvae resuming

development and completing their life-cycle are unclear, but have been hypothesised to

involve cross-talk between adult and larval worms, along with interaction with the gut

microbiota, the host immune system and environmental disruption such as anthelmintic

treatment (Gibson, 1953; Love *et al*., 1999; Matthews, 2008; Walshe *et al*., 2019; Walshe *et*

al., 2021; Walshe *et al*., 2020). The nature of communication between adult and larval

cyathostomins, and the extent to which this can occur between species, is not well-

understood. The existence of a large number of species that are typically present in mixed

infections highlights the importance of understanding these dynamics, in efforts both to

understand the basic biology of cyathostomin infections and the provision of improved

control strategies (Bellaw and Nielsen, 2020; Hodgkinson *et al*., 2003).

Horses may tolerate light cyathostomin infection without the development of overt clinical

signs (Love *et al*., 1999; Morariu *et al*., 2016; Murphy and Love, 1997; Walshe *et al*., 2021).

However, high burdens can lead to ill-thrift and poor performance (Murphy and Love, 1997)

and in some cases a specific clinical presentation of acute larval cyathostominosis (ALC), which

develops in response to synchronous emergence of larvae from the mucosa and involves

severe local and systemic inflammation, with high morbidity and potential mortality

(Hodgkinson *et al*., 2003; Love *et al*., 1999; Walshe *et al*., 2020). The emergence of encysted

cyathostomin larvae is believed to be affected by many factors, including negative inhibition

from an established adult worm population, removed following anthelmintic treatment

(Steinbach *et al*., 2006; Walshe *et al*., 2019; Walshe *et al*., 2021). Administration of

anthelmintics has also been shown to have an adverse impact on the gut microbiota, either

transiently, in normal animals (Walshe *et al*., 2019) or more severely in ALC, with severe

typhlocolitis developing (Walshe *et al*., 2021). ALC has a 50% mortality rate recorded in

referral centres (Giles *et al*., 1985; Peregrine *et al*., 2006; Walshe *et al*., 2021).

In the past, characterisation of cyathostomin communities relied exclusively on

morphological identification of species (Kinsella, 2002; Lichtenfels *et al*., 2008). However, this

is laborious, time-consuming and highly specialised (Avramenko *et al*., 2015; Hodgkinson *et*

al., 2003; Lichtenfels *et al*., 2008), and it has largely been supplanted by molecular speciation,

either of individual worms, or using population analysis "Nemabiome" approaches

(Avramenko *et al*., 2015).

There are just two published studies on characterising cyathostomin species in horses in
Ireland. The first (Kinsella, 2002), used morphological speciation and the second (Elghryani,
2018) used reverse line blots of the intergenic spacer gene region. Elsewhere, internal
transcribed spacer 2 (ITS-2) primers as described by (Gasser *et al.*, 1993), have been used for
molecular diagnostics of equine nematode infections, reviewed by (Ghafar *et al.*, 2022).

We aimed to sequence the ITS-2 region from individual cyathostomins retrieved from
naturally infected horses in Ireland in a post-mortem study and to compare the community
composition to those described in previous studies in Ireland and internationally.

2. Materials and Methods

2.1. Ethical approval

This was a post-mortem study and did not require any specific ethical approval.

2.2. Study population

The study conducted an abattoir survey, undertaken between August 2018 and June 2020. A
total of eleven equine carcasses were selected at random for examination at two Irish
abattoirs. In accordance with food safety legislation, no horse had received anthelmintic
treatment within 32 days of slaughter. The horses were assessed for clinical abnormalities by
a veterinarian prior to inclusion in the study as part of the normal pre-mortem inspection at
the abattoir.

Samples were taken in September, October, February, March and June, with samples taken
in February designated as winter samples, March as spring, June as summer, and
September/October as autumn.

2.3. Sample processing

Each carcass was processed as per abattoir standard operating procedures. Sampling was
undertaken on the same day as euthanasia. To isolate the large intestine, the ileocecal
junction was located, tied off with cable ties and dissected. The liver, stomach, spleen,
pancreas, and small intestine were removed, and the large intestine was transferred from the
processing line to a dissection table.

Further cable ties were placed at the junction between the caecum and right ventral colon,
the junction between the right ventral and left ventral colon, the junction between the left
ventral colon and left dorsal colon, and the junction between the left dorsal and right dorsal
colon. An incision was made in each of the three intestinal segments (caecum, right ventral
colon (RVC) and left dorsal colon (LDC)), and the luminal contents were emptied separately
into individual 10L graduated buckets. For each section, the mucosa was gently rinsed, and
the liquid was added to the bucket containing luminal contents. The contents were then
stirred whilst water was added to make a homogeneous suspension to a volume of 10L. Two
100ml aliquots were taken from each section, each equating to 1% of the volume within each
container. Faeces were collected from the rectum of each horse and placed in a ziplock bag.
All samples were placed on ice for transfer to the laboratory.

2.4. Faecal Egg Counting
The Mini-FLOTAC technique (Cringoli, 2006) was utilised to count strongyle eggs in faeces.
Briefly, 45ml of saturated NaCL was added to 5g of faeces and the flotation chambers were
completely filled with the faecal suspension. After 10 minutes, the key was used to turn the
reading disc clockwise (about 90°), the sample was examined under the microscope and the
eggs per gram counted.

2.5. Luminal worm retrieval

Each duplicate 1% aliquot of the caecum, RVC and LDC was examined individually using a
modification of the Baermann method. The contents of each aliquot were enclosed in a
muslin parcel suspended in water in a conical flask and left to sediment for 24 hours. The
entire sediment was then examined in detail (using aliquots forming a thin layer in a petri

dish) under a dissecting microscope. The remaining sediment was also examined under a dissecting microscope.

The total number of worms present in each 100ml sample was enumerated and recorded. Individual worms were removed manually and washed prior to storage. Each worm was placed in a 2ml Eppendorf with >/=50ml PBS and stored at -20˚C. For each section a maximum of 50 worms (range 1-50) was collected for subsequent examination from combined duplicate samples

2.6. DNA Extraction and amplification

2.6.1. Genomic DNA extraction from individual nematodes

Individual worms were dissected into a minimum of three parts using a sterile needle. DNA was then extracted from each worm using the Qiagen Powerfecal Pro extraction kit as per the standard operating protocol, with minor adaptations. Specifically, samples were pulse vortexed for a minimum of 45 minutes, the dry spin after solution C5 was at 16,600 x G for three minutes, and solution C6 of the kit was put through the filter membrane twice to increase elution yield (Bredtmann *et al.*, 2019). After extraction, samples were stored at -20°C until PCR amplification.

2.6.2. PCR amplification

PCR reactions were performed in 50µl reaction volumes using ITS-2 primers as described by Gasser *et al.*, (1993), (NC1 forward; 5'ACGTCTGGTTCAGGGTTGTT3' and NC2 reverse; 5'TTAGTTTCTTTTCCTCCGCT3').

The reaction mixture 10µl of Promega GoTaq Flexi Buffer, 1.5mM of Promega MgCL2 solution, 0.2mM Promega DNTP PCR nucleotide mix, 1µM of each forward and reverse primer (Eurofins Genomics) 1.25U of Promega GoTaq G2 Flexi DNA Polymerase, 10µl template DNA and 15.75µl nuclease free water. Amplification steps included initial denaturation at 95°C for 2

minutes, then cycling steps starting at 95°C for 1 minute, annealing at 55°C for 1 minute and

extension at 72°C for 1 minute, for 40 cycles, followed by extension at 72°C for 5 minutes.

For each run, a negative control containing 10µl nuclease-free water in place of DNA template

was included.

Amplicons were purified using the Qiagen Qiaquick PCR purification kit and sequenced in both

directions using the PCR primers (Eurofins Genomics).

2.7. Molecular speciation

Consensus sequences were constructed from forward and reverse reads and compared with

published nematode sequences in the NCBI database using Clustal Omega (Clustal Omega <

Multiple Sequence Alignment < EMBL-EBI).

Samples that did not yield a reliable BLAST result, E value > E^{-10}, low query cover and

percentage identity <95%, were removed from the dataset.

2.8. Data analysis

Data was analysed using SPSS (Version 27) for descriptive statistics and to evaluate statistical

significance. Significance was determined using T-tests and ANOVA. The Kruskal-Wallis test

was utilised for non-parametric data including faecal egg counts (FEC), numbers of worms and

species distribution. Spearman rank correlation coefficient was calculated and created using

R functions, Excel and Inkscape (v.1.3, https://inkscape.org/release/inkscape-1.3/). The

ggplot2 package in R (version 4.3.1 - https://cran.r-project.org/bin/windows/base/old/),

Excel and Inkscape were used for data visualisation. MEGA11, IQtree web server

(http://iqtree.cibiv.univie.ac.at/) and iTOL (https://itol.embl.de/) were used for phylogenetic

tree creation. SPSS (V27) was used for Chi Squared, T-tests and ANOVA analyses. MEGA11

was used for MUSCLE alignment and trimming of sequences. A Maximum Likelihood tree was

created using MEGA, IQtree and iTOL to create a tree with 1,000 ultrafast bootstrap

replications and pairwise deletion for gaps/missing data. In order to construct a simplified

tree to compare multiple species, representative sequences were selected for each species.

3. Results

The genetic data from this study has been uploaded to GenBank under the accession numbers OR941129 - OR941242.

3.1. Study cohort and parasitism levels

Table 1 provides the details of horses included in the study, together with their strongyle FEC and number of worms recovered in a 1% sample of the total luminal contents. The horses included in the study ranged from yearlings to twenty-three years old with a mean of 7.7 years of age, a median of 6 years. There were four females and seven males. Five were thoroughbred, and six Irish Sport Horse. FEC ranged from 0-1,670 eggs per gram (epg). Nematodes were retrieved from two horses with a FEC of 0. Conversely, high FEC did not always correlate with high luminal burden of nematodes. Measurement of worm burdens revealed a negative binomial distribution. Five horses were sampled in autumn and two horses were sampled in spring, summer and winter, respectively.

3.2. Species recovered, abundance and prevalence

Table 2 shows the number of worms recovered in each intestinal compartment in this study, and compared to those from a previous study on horses in Ireland (Kinsella, 2002). In both studies, numbers recovered from the caecum were significantly lower than from the colonic compartments (our study: P=0.009, Kinsella P=0.05). The LDC yielded the highest number of worms in our study. There was no significant difference between the numbers recovered from the RVC and LDC.

Table 3 shows the numbers of worms speciated from each intestinal compartment for each horse. Up to 14 individual worms from each intestinal compartment, per horse, were successfully speciated. For all but one horse, worms from two or three compartments were speciated.

The species abundance in each of the three intestinal compartments is shown in Figure 1.

Coronocyclus coronatus was the most abundant species recovered (29%). Between them,

Cor. coronatus, *Cylicocyclus nassatus* and *Cylicostephanus longibursatus* accounted for more

than half (54%) of the entire dataset. Although there was no significant difference in species

distribution across the three intestinal compartments, *Cylicocyclus insigne* was equally

abundant to *Cor. coronatus* in the LDC.   This compartment also had the most diverse range

of species (n=11), with the caecum having the least diversity (n=8).

The prevalence of individual species and the richness of cyathostomin communities in this

study population is illustrated in Figure 2. *Cor. coronatus* and *Cyc. longibursatus* were the

most prevalent, being found in seven of the eleven horses (63.63% prevalence), *Cyc. nassatus*

was present in six horses (54.54%) and *Cylicostephanus goldi* in five horses (45.45%). *Cyc.*

insigne, *Cylicostephanus calicatus*, *Cylicocyclus ashworthi* and *Cyathostomum catinatum*

were each recorded in four horses, at a prevalence of 36.36%. *Cyathostomum pateratum* was

found in three horses (27.27% prevalence) with *Cylicostephanus minutus, Coronocyclus.*

labiatus and *Cylicocyclus leptostomum* in one animal each, comprising 9.09% prevalence.

Species richness ranged from two to seven species per individual, with a mean of 4.27 +/-

0.604 (SEM). The modal species richness was two and five species equally, with a median

species richness of five within this dataset. Over 50% (54.5%) of the horses sampled had >5

cyathostomin species present. One large strongyle, *Strongylus vulgaris* was identified (Horse

5) but not included in the dataset.

3.3. Community composition

Figure 3 is a heatmap (Spearman correlation) of species co-occurrence. The analysis identified

significant positive correlation (0.86) between *Cya. pateratum* and *Cyc. calicatus* ($p < 0.01$). A

negative correlation was observed between *Cyc. insigne* and *Cyc. calicatus,* although this did

not reach significance ($p = 0.08$). *Cor. coronatus* showed trends towards positive co-

occurrences with *Cyc. nassatus* (0.46), *Cyc. goldi* (0.51), *Cyc. calicatus* (0.55), *Cyc. ashworthi*

(0.46), *Cya. pateratum* (0.33) and *Cyc. minutus* (0.41). *Cor. labiatus* had the most negative

correlations with seven negative interactions recorded to *Cyc. minutus* (-0.1), *Cya. pateratum*

(-0.19), *Cya. catinatum* (-0.23), *Cyc. calicatus* (-0.23), *Cyc. goldi* (-0.27), *Cyc.nassatus* (-0.28),

and *Cyc. nassatus* (-0.36).

Next, (Figure 4) we compared the prevalence of cyathostomin species found in our study with

that found in a previous study in Ireland (Kinsella, 2002) and a global meta-analysis (Bellaw

and Nielsen, 2020). Twelve cyathostomin species were common to our study, Kinsella, 2002,

and the global meta-analysis. There were no species identified in Ireland that were not

identified in the latter. Broadly, the most prevalent species in the meta-analysis and in

Kinsella, 2002, were also highly represented in our study. Prevalence as reported by Bellaw

and Nielsen, 2020, was highest for *Cya. catinatum* (94%), *Cyc. nassatus* (91%), and *Cyc.*

longibursatus (89%). However, we found *Cor. coronatus* and *Cyc. nassatus* to have the joint

highest prevalence, occurring in 64% (n=7) of the horses sampled. Kinsella, 2002, recorded

four species at 100% prevalence, namely, *Cya. catinatum, Cyc. nassatus, Cyc. calicatus* and

Cyc. longibursatus. Of the 35 species recorded in the meta-analysis, our study identified 34%

of species, while Kinsella, 2002, identified 54%. The ten most prevalent species in both Irish

studies matched those outlined in the global meta-analysis, albeit with some differences in

the ranking of the top ten species. The other Irish study (Elghryani, 2018) did not have

sufficient prevalence data for comparative purposes. The most notable absence in our study

from the top species in the meta-analysis is *Cor. labratus*, which was the 12[th] most prevalent

species recorded globally but has not been identified in any of the three Irish studies

mentioned here.

3.4. Seasonal variation

Figure 5 shows the seasonal variation in abundance recorded in our study. Notably, the most

highly abundant species, *Cor. coronatus*, was recorded only during horses sampled during

autumn. The greatest number of species (nine) were also identified in autumn samples. *Cyc.*

longibursatus was the only species recorded in all four seasons. Seven of the species were

observed in only one season.

3.5. Phylogenetic analysis of species

Sequences were used for construction of a Maximum Likelihood phylogenetic tree (Figure 6A)

which was rooted at the midpoint. This unveiled some interspecies occurrences within

branches, notably between *Cyc. calicatus* and *Cor. coronatus*.

Figure 6B shows a simplified phylogenetic tree, where consensus sequences for individual

cyathostomins sequenced at the ITS-2 region were used for phylogenetic analysis. *Cya.*

pateratum and *Cyc. goldi* branched together, as did *Cor. labiatus* and *Cyc. Insigne,* and *Cyc.*

leptostomum with *Cyc. Nassatus. Cyc. calicatus* and *Cor. coronatus* branch off together from

the root of the tree.

4. Discussion

4.1. Strongyle egg counts

There was no direct relationship between strongyle faecal egg count and worm burden, which

was not unexpected (Chapman *et al*., 2003a; Chapman *et al*., 2003b; Nielsen *et al*., 2010) but

indicates the importance of understanding the potential differences between individual

cyathostomin species in terms of differences in biotic potential.   We recovered worms from

horses with a 0 EPG count, which has been well-documented previously (Gibson, 1953; Love

et al., 1999; Matthews, 2008; Walshe *et al*., 2019; Walshe *et al*., 2020). As expected, we

observed a negative binomial distribution in the analysis of the egg shedding in our

population.

4.2. Species abundance

This study detected twelve of the thirteen most common species globally (Figure 4), as

compared to Bellaw and Nielsen's meta- analysis (Bellaw and Nielsen, 2020). Among the

nineteen species identified in the morphological study by Kinsella (2002), we detected twelve.

Notably, *Cya. Catinatum's* prevalence was 36% in our study, contrasting to 100% prevalence

in Kinsella's work and 94% in the global meta-analysis. Elghryani (2018), found *Cyc. nassatus*

as the most prevalent among the twelve species identified. The high abundance and

prevalence of *Cyc. nassatus* is consistent among all three Irish studies and the global meta-

analysis. Overall, the prevalence findings between the three studies from Ireland broadly reflect one another (Figure 3). A notable observation was the absence of the twelfth most prevalent species as identified by (Bellaw and Nielsen, 2020), *Cor. labratus,* which was absent from the Irish studies. This species has been recorded in the UK and France (Cwiklinski *et al*., 2012; Peachey *et al*., 2018).

4.3. Phylogenetic analysis

 Cyc. calicatus and *Cor. coronatus* branch off together at the root of the tree 6B and are also interdispersed in the ML tree (Figure 6A). This reflects their close relationship, as previously identified, and the limitations of using the ITS-2 locus to differentiate the subsets of these species (Courtot *et al*., 2023; Hung *et al*., 2000; Louro *et al*., 2021).

4.4. Characterisation of cyathostomin communities, and correlation with other properties in our study population.

4.4.1. Distribution across intestinal compartments

Across the three intestinal compartments examined in this study, the number of worms retrieved from the caecum was significantly lower than that recovered from LDC (P =0.009). (Ogbourne, 1976), assessed abundance of cyathostomins postmortem and found 50% in the ventral colon and 45% in the dorsal colon. The worm burden in the caecum represented only 5% of the overall cyathostomin burden. Similarly, (Stancampiano *et al*., 2010), observed that the majority of small strongyles were concentrated in the ventral colon (67.34%) and less frequently in the caecum (27.52%) and dorsal colon (5.14%). While this distribution was also reflected by Kinsella (2002), the converse was found for encysted stages of cyathostomins, as recorded by (Nielsen *et al*., 2021) who documented 40% of EL3/LL3/L4 in the caecum, and 38%/22% in the ventral and dorsal colon respectively. This recurrent recording of biological preference of luminal cyathostomins for the colon over the caecum, and the inverse for encysted larvae may reflect the slower transit time of the colon and the natural peristalsis of

food through the digestive tract, enabling passive transport and a longer exposure to the
intestinal wall necessary to facilitate larval encystment (Geor *et al.*, 2013; Miyaji *et al.*, 2008).

4.4.2. Community composition

Over half of the horses in this study had more than five cyathostomin species recovered from
them (Figure 2). We found a significant positive correlation between the occurrence of *Cyc.*
nassatus and *Cyc. insigne* (Figure 3), which supports similar findings by (Sallé *et al.*, 2018).

4.4.3. Seasonality

Samples were taken from each quarter of the year to account for seasonal factors of
cyathostomin distribution. *Cyc. longibursatus* was consistently recovered in all four seasons.
The most abundant species, *Cor. coronatus*, was identified only in autumn, as was the case
for *Cyc. calicatus*, which share a close relationship at the ITS-2 region (Hung *et al.*, 2000; Louro
et al., 2021). Increased abundance of *Cor. coronatus* in autumn was also noted in a recent
study in Saskatchewan which found a significantly higher proportional abundance of *Cor.*
coronatus in the autumn compared to the spring (Saworski, 2023). Our findings also reflect
those of Kinsella (2002), who found significantly more cyathostomin species recorded in
spring and autumn than in other seasons. A Louisiana based study also reflected these
seasonal findings, with autumn recording the greatest species diversity (Chapman *et al.*,
2003b). Variability of seasonal patterns among species is an understudied facet of
cyathostomin biology which may be crucial for comprehensive characterisation of
pathogenesis and epidemiology of cyathostomin-related disease, as this tends to have a
seasonal distribution of presentation (Reid *et al.*, 1995).

5. Limitations

The most important limitation of this study is sample size. This is a descriptive study, which
affects the potential for statistical power. Conclusions from observations within any study

with a limited sample size must take this into account. Results from Table 3 are considered in

the context that only two nematodes were retrieved for sampling from horse eight. Notably,

it is clear from this research and other studies that the ITS-2 locus cannot resolve all isolates

of *Cor. coronatus* and *Cyc. calicatus,* pointing to the need for a different locus for further

molecular work, such as COX1, or a multi-locus approach (Courtot *et al*., 2023; Ghafar *et al*.,

2022).

6. Conclusions

This study has produced new insights into cyathostomin species occurrence, interactions,

community composition and seasonality in horses in Ireland, with comparisons to previous

studies in this locality a global meta-analysis.

It is likely that communication occurs not only between cyathostomin species and stages, but

also between the helminths, gut microbiota and the host, as observed within other systems

within the gastrointestinal tract (Gieryńska *et al*., 2022). Post-mortem analysis and

identification in naturally-infected hosts, as conducted here, will continue to have a role to

play in advancing understanding of cyathostomin biology and pathogenesis, in parallel with

population-based approaches. More extensive studies involving both population and post-

mortem studies are warranted in order to enhance risk assessment, biomarker-based

diagnostic tools and treatment protocols (Walshe *et al*., 2020). It is also useful in investigating

potentially cryptic species as seen in this study with *Cor. coronatus* and *Cyc. calicatus.*

Studies on cyathostomin population dynamics are a crucial step in understanding the

complex picture of cyathostomin infections in the face of pervasive anthelmintic resistance

and their potential to advance mechanistic understanding of gut health in horses.

Author contributions

388   Conceptualization, G.M., N.W., O.B.; writing- original draft preparation, O.B. – review and

389   editing, G.M., N.W., O.B. and A.Z.; laboratory work and bioinformatics- O.B., D.G.; student

390   laboratory contributions, L.K., O.BX., D.McS., F.ON., S.D., BR.L. All authors have read and

391   agreed to the published version of this manuscript.

Funding:

This research was funded by a Clinical Primer grant to Nicola Walshe from the Wellcome
Institutional Strategic Support fund, and by UCD's own funds.

Acknowledgements:

We would like to acknowledge the valuable input and assistance of Amanda Lawlor, Senior
Technical Officer, UCD School of Veterinary Medicine, and Dr. Nagwa Elghyrani.

Conflict of interest:

The authors declare no conflict of interest

References:

Avramenko, R.W., Redman, E.M., Lewis, R., Yazwinski, T.A., Wasmuth, J.D., Gilleard, J.S., 2015. Exploring the gastrointestinal "nemabiome": deep amplicon sequencing to quantify the species composition of parasitic nematode communities. PLoS One 10, e0143559.

Bellaw, J.L., Nielsen, M.K., 2020. Meta-analysis of cyathostomin species-specific prevalence and relative abundance in domestic horses from 1975–2020: emphasis on geographical region and specimen collection method. Parasites & Vectors 13, 1-15.

Boisseau, M., Mach, N., Basiaga, M., Kuzmina, T., Laugier, C., Sallé, G., 2023. Patterns of variation in equine strongyle community structure across age groups and gut compartments. Parasites & Vectors 16, 64.

Boxell, A., Gibson, K., Hobbs, R., Thompson, R., 2004. Occurrence of gastrointestinal parasites in horses in metropolitan Perth, Western Australia. Australian Veterinary Journal 82, 91-95.

Bredtmann, C.M., Krücken, J., Murugaiyan, J., Balard, A., Hofer, H., Kuzmina, T.A., von Samson-Himmelstjerna, G., 2019. Concurrent proteomic fingerprinting and molecular analysis of cyathostomins. Proteomics 19, 1800290.

Chapman, M., Kearney, M., Klei, T., 2003a. Equine cyathostome populations: accuracy of species composition estimations. Veterinary parasitology 116, 15-21.

Chapman, M.R., French, D.D., Klei, T.R., 2003b. Prevalence of strongyle nematodes in naturally infected ponies of different ages and during different seasons of the year in Louisiana. Journal of Parasitology 89, 309-314.

Courtot, É., Boisseau, M., Dhorne-Pollet, S., Serreau, D., Gesbert, A., Reigner, F., Basiaga, M., Kuzmina, T., Lluch, J., Annonay, G., 2023. Comparison of two molecular barcodes for the study of equine strongylid communities with amplicon sequencing. PeerJ 11, e15124.

Cringoli, G., 2006. FLOTAC, a novel apparatus for a multivalent faecal egg count technique. Parassitologia 48, 381-384.

Cwiklinski, K., Kooyman, F., Van Doorn, D., Matthews, J., Hodgkinson, J., 2012. New insights into sequence variation in the IGS region of 21 cyathostomin species and the implication for molecular identification. Parasitology 139, 1063-1073.

Dowdall, S., Matthews, J., Mair, T., Murphy, D., Love, S., Proudman, C., 2002. Antigen-specific IgG (T) responses in natural and experimental cyathostominae infection in horses. Veterinary parasitology 106, 225-242.

Duncan, J., Bairden, K., Abbott, E., 1998. Elimination of mucosal cyathostome larvae by five daily treatments with fenbendazole. Veterinary Record 142, 268-271.

Elghryani, N., 2018. The Prevalence and Anthelmintic Sensitivity of Gastrointestinal Nematodes of Horses in Ireland. University College Dublin,

Gasser, R.B., Chilton, N.B., Hoste, H., Beveridge, I., 1993. Rapid sequencing of rDNA from single worms and eggs of parasitic helminths. Nucleic acids research 21, 2525.

Geor, R.J., Harris, P., Coenen, M., 2013. Equine applied and clinical nutrition: health, welfare and performance. Elsevier Health Sciences.

Ghafar, A., Abbas, G., Beasley, A., Bauquier, J., Wilkes, E.J., Jacobson, C., McConnell, E., El-Hage, C., Carrigan, P., Cudmore, L., 2022. Molecular diagnostics for gastrointestinal helminths in equids: Past, present and future. Veterinary Parasitology, 109851.

Gibson, T., 1953. The effect of repeated anthelmintic treatment with phenothiazine on the faecal egg counts of housed horses, with some observations on the life cycle of Trichonema spp. in the horse. Journal of Helminthology 27, 29-40.

Gieryńska, M., Szulc-Dąbrowska, L., Struzik, J., Mielcarska, M.B., Gregorczyk-Zboroch, K.P., 2022. Integrity of the intestinal barrier: The involvement of epithelial cells and microbiota—A mutual relationship. Animals 12, 145.

454 Giles, C., Urquhart, K., Longstaffe, J., 1985. Larval cyathostomiasis (immature trichonema-induced
455   enteropathy): A report of 15 clinical cases. Equine veterinary journal 17, 196-201.

456 Hinney, B., Wirtherle, N.C., Kyule, M., Miethe, N., Zessin, K.-H., Clausen, P.-H., 2011. Prevalence of
457   helminths in horses in the state of Brandenburg, Germany. Parasitology research 108, 1083-
458   1091.

459 Hodgkinson, J.E., Lichtenfels, J.R., Mair, T., Cripps, P., Freeman, K., Ramsey, Y., Love, S., Matthews, J.,
460   2003. A PCR–ELISA for the identification of cyathostomin fourth-stage larvae from clinical
461   cases of larval cyathostominosis. International Journal for Parasitology 33, 1427-1435.

462 Hung, G.-C., Chilton, N.B., Beveridge, I., Gasser, R.B., 2000. A molecular systematic framework for
463   equine strongyles based on ribosomal DNA sequence data. International Journal for
464   Parasitology 30, 95-103.

465 Kinsella, J.L., 2002. Species Composition, Distribution, and Abundance, with Monthly Variations of
466   Cyathostominea Within the Equid in Ireland, 1999/2000, and a Preliminary Analysis of
467   Proteolytic Activity of Excretory Secretory Products from Cyathostominea. University College
468   Dublin,

469 Lichtenfels, J.R., Kharchenko, V.A., Dvojnos, G.M., 2008. Illustrated identification keys to strongylid
470   parasites (Strongylidae: Nematoda) of horses, zebras and asses (Equidae). Veterinary
471   parasitology 156, 4-161.

472 Louro, M., Kuzmina, T.A., Bredtmann, C.M., Diekmann, I., de Carvalho, L.M.M., von Samson-
473   Himmelstjerna, G., Krücken, J., 2021. Genetic variability, cryptic species and phylogenetic
474   relationship of six cyathostomin species based on mitochondrial and nuclear sequences.
475   Scientific Reports 11, 8245.

476 Love, S., Murphy, D., Mellor, D., 1999. Pathogenicity of cyathostome infection. Veterinary parasitology
477   85, 113-122.

478 Lyons, E.T., Tolliver, S., Drudge, J., 1999. Historical perspective of cyathostomes: prevalence,
479   treatment and control programs. Veterinary Parasitology 85, 97-112.

480 Matthews, J., 2008. An update on cyathostomins: anthelmintic resistance and worm control. Equine
481   Veterinary Education 20, 552-560.

482 Miyaji, M., Ueda, K., Nakatsuji, H., Tomioka, T., Kobayashi, Y., Hata, H., Kondo, S., 2008. Mean
483   retention time of digesta in the different segments of the equine hindgut. Animal Science
484   Journal 79, 89-96.

485 Morariu, S., Mederle, N., Badea, C., Dărăbuş, G., Ferrari, N., Genchi, C., 2016. The prevalence,
486   abundance and distribution of cyathostomins (small stongyles) in horses from Western
487   Romania. Veterinary Parasitology 223, 205-209.

488   Murphy, D., Love, S., 1997. The pathogenic effects of experimental cyathostome infections in ponies.
489   Veterinary parasitology 70, 99-110.

490   Nielsen, M.K., Baptiste, K.E., Tolliver, S., Collins, S., Lyons, E., 2010. Analysis of multiyear studies in
491   horses in Kentucky to ascertain whether counts of eggs and larvae per gram of feces are
492   reliable indicators of numbers of strongyles and ascarids present. Veterinary Parasitology 174,
493   77-84.

494   Nielsen, M.K., Lyons, E.T., 2017. Encysted cyathostomin larvae in foals–progression of stages and the
495   effect of seasonality. Veterinary parasitology 236, 108-112.

496   Nielsen, M.K., Martin, A.N., Scare, J.A., Steuer, A.E., 2021. Precision and spatial variation of
497   cyathostomin mucosal larval counts. Veterinary Parasitology 290, 109349.

498   Ogbourne, C.P., 1976. The prevalence, relative abundance and site distribution of nematodes of the
499   subfamily Cyathostominae in horses killed in Britain. Journal of Helminthology 50, 203-214.

500   Peachey, L., Molena, R., Jenkins, T., Di Cesare, A., Traversa, D., Hodgkinson, J., Cantacessi, C., 2018.
501   The relationships between faecal egg counts and gut microbial composition in UK
502   Thoroughbreds infected by cyathostomins. International journal for parasitology 48, 403-412.

503   Peregrine, A.S., McEwen, B., Bienzle, D., Koch, T.G., Weese, J.S., 2006. Larval cyathostominosis in
504   horses in Ontario: an emerging disease? The Canadian Veterinary Journal 47, 80.

505   Reid, S., Mair, T., Hillyer, M., Love, S., 1995. Epidemiological risk factors associated with a diagnosis of
506   clinical cyathostomiasis in the horse. Equine Veterinary Journal 27, 127-130.

507   Sallé, G., Kornaś, S., Basiaga, M., 2018. Equine strongyle communities are constrained by horse sex
508   and species dipersal-fecundity trade-off. Parasites & vectors 11, 1-11.

509   Saworski, T., 2023. Fecal Egg Counts, Nemabiomes, and Management Practices in Saskatchewan
510   Horses. University of Saskatchewan,

511   Stancampiano, L., Gras, L.M., Poglayen, G., 2010. Spatial niche competition among helminth parasites
512   in horse's large intestine. Veterinary Parasitology 170, 88-95.

513   Steinbach, T., Bauer, C., Sasse, H., Baumgärtner, W., Rey-Moreno, C., Hermosilla, C., Damriyasa, I.M.,
514   Zahner, H., 2006. Small strongyle infection: Consequences of larvicidal treatment of horses
515   with fenbendazole and moxidectin. Veterinary Parasitology 139, 115-131.

516   Walshe, N., Duggan, V., Cabrera-Rubio, R., Crispie, F., Cotter, P., Feehan, O., Mulcahy, G., 2019.
517   Removal of adult cyathostomins alters faecal microbiota and promotes an inflammatory
518   phenotype in horses. International journal for parasitology 49, 489-500.

519   Walshe, N., Mulcahy, G., Crispie, F., Cabrera-Rubio, R., Cotter, P., Jahns, H., Duggan, V., 2021. Outbreak
520   of acute larval cyathostominosis–A "perfect storm" of inflammation and dysbiosis. Equine
521   Veterinary Journal 53, 727-739.

522   Walshe, N., Mulcahy, G., Hodgkinson, J., Peachey, L., 2020. No Worm is an island; the influence of

523   commensal gut microbiota on cyathostomin infections. Animals 10, 2309.

Tables

Table 1: Horse details, FEC and number of worms recovered. RVC= Right Ventral Colon and LDC = Left Dorsal Colon. * = P < 0.05. Mean calculations are +/- the SEM (standard error of the mean). The number of worms recovered accounts for a 1% subsection of the total luminal contents of each animal sampled. Numbers of worms recovered from the caecum were lower than those recovered from LDC (P=0.009). No significant difference was recorded between RVC or from composite RVC and LDC colon to the caecum (P=0.291). FEC represents eggs per gram (epg) of faecal material).

		Horse Details					Number of worms recovered	
Horse	Age	Breed	Sex	Sampling Date	FEC (epg)	RVC	LDC	Caecum
1	16	ISH	F	Sep-18	1550	79	188	10
2	8	ISH	M (gelding)	Sep-18	0	11	3	1
3	1	TB	M (colt)	Sep-18	1670	205	155	12
4	10	ISH	F	Oct-18	25	201	245	23
5	1	TB	M (colt)	Oct-18	905	590	33	16
6	6	ISH	M (colt)	Feb-20	225	2	26	0
7	2	TB	M (colt)	Feb-20	800	1	249	2
8	12	ISH	M	Mar-20	50	0	2	0
9	23	TB	F	Mar-20	85	9	24	1
10	3	TB	F	Jun-20	715	1410	3328	33
11	3	ISH	M	Jun-20	0	40	56	53
Total	-	-	-	-	-	2548	4309	151
Mean	7.7+/- 2.124	-	-	-	547.73 +/- 188.83	231.63 +/- 129.21	391 +/- 295*	13.73 +/- 5.1*
Range	1-23	-	-	-	0-1670	0-1410	2-3328	0-53

Table 2: Comparison of numbers of cyathostomins recovered from the caecum, RVC and LDC in our study and with a previous study in horses in Ireland from Kinsella (2002). In both studies, the number of worms recovered from the caecum was significantly lower, (our study: P=0.009, Kinsella P=0.05). The greatest number of worms was recovered from RVC in the 2002 study, whereas LDC yielded the greatest number overall in our study. Total worms in our study refers to 1% sample aliquot which was taken from each section of the gastrointestinal tract sampled. Kinsella retrieved 250ml aliquots from 15 sections of the Caecum, ventral colon and dorsal colon.

Region	Range This Study	Total Worms This Study	% Of Total	Total Worms Kinsella 2002	% Of Total Kinsella 2002
Caecum	0-53	151	2.1	511	6.9
RVC	0-1410	2548	36.3	4064	55.5
LDC	2-3328	4309	61.4	2744	37.4
Total Across Compartments	0-3328	7008	-	7319	-

Table 3: Numbers of worms speciated from each intestinal compartment for each horse. Mean numbers are +/- SEM). Up to 14 individual worms were speciated from each area, and in total, 116 individual worms were sequenced with sufficient data quality for speciation.

Horse	RVC	LDC	Caecum
1	2	9	-
2	5	-	-
3	8	2	4
4	3	13	-
5	3	-	4
6	1	4	-
7	-	4	1
8	-	2	-
9	3	3	-
10	8	5	14
11	4	5	9
Total	37	47	32
Mean	3.36 +/- 0.83	4.27 +/- 1.16	2.91 +/- 1.4
Range	0-8	0-13	0-14

Figure 1: Species abundance data from samples obtained from Right Ventral Colon (a) Left Dorsal Colon (b), Caecum (c) and across all three compartments (d) . *Cor. coronatus* was the most abundant species recovered, at 29% of total nematodes speciated. Between them, *Cor. coronatus*, *Cyc. nassatus* and *Cyc. longibursatus* accounted for more than half (54%) of the entire dataset. The LDC had the most diverse range of species (n=11), with the caecum having the least diversity (n=8).

Horse	Coronocyclus coronatus	Cylicocyclus nassatus	Cylicostephanus longibursatus	Cylicostephanus goldi	Cylicocyclus insigne	Cylicostephanus calicatus	Cylicocyclus ashworthi	Cyathostomum catinatum	Cyathostomum pateratum	Cylicostephanus minutus	Strongylus vulgaris	Coronocyclus labiatus	Cylicocyclus leptostomum
1	■		■	■		■			■				
2	■			■									
3	■	■				■	■			■			
4	■	■	■	■		■	■		■				
5	■		■			■		■	■		■		
6		■	■		■								
7			■				■					■	
8		■				■							
9			■					■					
10	■	■		■		■	■	■					
11	■	■	■	■				■					■

Figure 2: Prevalence of individual cyathostomin species and richness of cyathostomin communities in this study population. Filled sections depict species found in each horse. The mean species richness across samples was 4.27 +/- 0.604 (S.E.M). Richness ranged from 2-7 species per horse. There was a median of five species within this dataset. Over 50% (54.5%) of the horses sampled had >5 cyathostomin species. *Coronatus coronatus* and *Cylicostephanus longibursatus* were the most prevalent in 7 of 11 subjects (63.63% prevalence), *Cylicocyclus nassatus* was present in 6 horses (54.54%) and *Cylicostephanus goldi* was present in 5 horses (45.45%). The least prevalent were *Cylicostephanus minutus, Coronocyclus labiatus* and *Cylicocyclus leptostomum* were in one animal each, comprising 9.09% prevalence. One large strongyle, *Strongylus vulgaris* was identified but not included in the cyathostomin dataset.

Figure 3: Heatmap of species co-occurrence using Spearmann correlations. This heatmap indicated a significant positive correlation (0.86) between *Cyathostomum pateratum* and *Cylicostephanus calicatus* (p < 0.01). A negative correlation was observed between *Cylicocylus insigne* and *Cylicostephanus calicatus,* although this did not reach significance (p = 0.08).

Figure 4: Prevalence of cyathostomin species as compared with a previous study in Ireland (Kinsella 2002) and a global meta-analysis (Bellaw and Nielsen, 2020). We identified twelve cyathostomin species recorded in both the previous Irish study and the global meta-analysis. There were no species identified in either study in Ireland that were not identified within the global meta-analysis. Generally, the most prevalent species in the two previous studies were also highly prevalent in our study.

Figure 5: Seasonal variation observed in the recovery of cyathostomin species from the large intestine (n=116). Winter= Dec-Feb, Spring= Mar-May, Summer= June-Aug, Autumn= Sept-Nov. Sampling months include February, March, June, September and October. The greatest diversity of species (nine) was recovered in the Autumn. *Cylicostephanus. longibursatus* was the only species recovered in all four seasons. Seven species were observed in one season only.

Figure 6A: Full phylogenetic tree of sequences from recovered worms. Consensus sequences for individual cyathostomins were used for phylogenetic analysis. MEGA11 was used for MUSCLE alignment and trimming of sequences. A tree was created using the Maximum likelihood algorithm with 1,000 bootstrap replications and pairwise deletion for gaps. This tree unveiled species that located within groupings of other species, notably *Cylicostephanus calicatus* and *Coronocyclus coronatus.*

Figure 6B: A species-specific tree was created (5B) and a representative sequence was selected by creating ML trees of each species and selecting the midpoint sequence. *Cya. pateratum* and *Cyc. goldi* branched together, as did *Cor. labiatus* with *Cyc. insigne* and *Cyc. leptostomum* with *Cyc.nassatus. Cyc. calicatus* and *Cor. coronatus* diverge together at the root of the tree, reflecting their close relationship as identified by Louro (2021) and the discrepancy speciating the two using the ITS-2 locus. The asterix (*) denotes a bootstrapping value of more than 80%, indicating confidence that the species within this branch are truly more closely related to each other than to the species on the rest of the tree.