Molecular Species Determination of Cyathostomins from Horses in Ireland

Orla Byrne, Disha Gangotia, John Crowley, Annetta Zintl, Liam Kiser, Olivia Boxall, Daniel

McSweeney, Fiona O'Neill, Stacey Dunne, Breanna Rose Lamb, Nicola Walshe, Grace

Mulcahy.

UCD School of Veterinary Medicine, University College Dublin, Dublin, Ireland.

Corresponding author: orla.byrne1@ucdconnect.ie; +353873285717

Keywords: Equine; Helminth; Cyathostomin; Molecular Species Determination; Ireland

Abstract:

Cyathostomins are globally important equine parasites, responsible for both chronic and

acute pathogenic effects. The occurrence of mixed infections with numerous cyathostomin

species hinders our understanding of parasite epidemiology, host-parasite dynamics and

species pathogenicity. There have been few studies of cyathostomin species occurring in

horses in Ireland, where temperate climatic conditions with year-round rainfall provide

suitable conditions for infection of grazing animals with bursate nematodes. Here, we

amplified and sequenced the ITS-2 region of adult worms harvested at post-mortem from 11

adult horses between August 2018 and June 2020, and recorded species prevalence and

abundance of worms recovered from the caecum, right ventral colon and left dorsal colon,

using both BLAST and IDTAXA for taxonomic attribution. Phylogenetic relationships and

community composition were also recorded and compared with other relevant studies,

including a global meta-analysis. Overall, our results agree with previous studies that there

does not seem to be a major difference in cyathostomin species occurrence in equids in

different geographical regions. We confirmed the results of other workers in relation to the

difficulties in discriminating between *Cylicostephanus calicatus* and *Coronocyclus coronatus*

on the basis of ITS-2 sequences.

1. Introduction

Cyathostomins are a group of parasitic helminths that inhabit the gastrointestinal tract. They

are obligate parasites which are ubiquitous in equids and of significant pathogenic concern

(Boxell *et al.*, 2004; Ghafar *et al.*, 2022; Hinney *et al.*, 2011; Love *et al.*, 1999; Morariu *et al.*,

2016; Steinbach *et al.*, 2006). Currently, there are over 50 recognised species with 40

recorded as infecting horses, and mixed infections are the norm (Lichtenfels *et al.*, 2008;

Lyons *et al.*, 1999). Cyathostomins have a direct life cycle, with infection occurring through

ingestion of infective stage (L3) larvae on pasture. Following infection, cyathostomin larvae

penetrate into the mucosa of the large intestine, and can remain here in a dormant

(hypobiotic) state, emerging to mature and establish patent infection in the lumen after a

variable pre-patent period of up to three years (Gibson, 1953; Nielsen and Lyons, 2017). The

hypobiotic state is relevant to anthelmintic efficacy and to cyathostomin-associated disease

(Boisseau *et al.*, 2023; Morariu *et al.*, 2016). Typically, of the total worm burden in an

individual animal, the majority will comprise hypobiotic larvae in the mucosa (Dowdall *et al.*,

2002; Duncan *et al.*, 1998; Gibson, 1953; Matthews, 2008). The stimuli responsible for

hypobiotic larvae resuming development and completing their life-cycle are unclear, but are

thought to involve cross-talk between adult and larval worms, along with interaction with the

gut microbiota, the host immune system and environmental disruption such as anthelmintic

treatment (Gibson, 1953; Love *et al*., 1999; Matthews, 2008; Walshe *et al*., 2019; Walshe *et

al*., 2021; Walshe *et al*., 2020). The existence of a large number of species that are typically

present in mixed infections highlights the importance of understanding these dynamics, in

efforts both to understand the basic biology of cyathostomin infections and the provision of

improved control strategies (Bellaw and Nielsen, 2020; Hodgkinson *et al*., 2003).

Horses may tolerate light cyathostomin infection without the development of overt clinical

signs (Love *et al*., 1999; Morariu *et al*., 2016; Murphy and Love, 1997; Walshe *et al*., 2021).

However, high burdens can lead to ill-thrift and poor performance (Murphy and Love, 1997)

and in some cases a specific clinical presentation of acute larval cyathostominosis (ALC), which

develops in response to synchronous emergence of larvae from the mucosa and involves

severe local and systemic inflammation, with high morbidity and potential mortality

(Hodgkinson *et al*., 2003; Love *et al*., 1999; Walshe *et al*., 2020). The emergence of encysted

cyathostomin larvae is believed to be affected by many factors, including negative inhibition

from an established adult worm population, removed following anthelmintic treatment

(Steinbach *et al*., 2006; Walshe *et al*., 2019; Walshe *et al*., 2021). ALC has a 50% mortality rate

recorded in referral centres (Giles *et al*., 1985; Peregrine *et al*., 2006; Walshe *et al*., 2021).

In the past, characterisation of cyathostomin communities relied exclusively on

morphological identification of species (Kinsella, 2002; Lichtenfels *et al*., 2008). However, this

is laborious, time-consuming and highly specialised (Avramenko *et al*., 2015; Hodgkinson *et

al*., 2003; Lichtenfels *et al*., 2008), and it has largely been supplanted by molecular species

determination, either of individual worms, or using population analysis "Nemabiome"

approaches (Avramenko *et al.*, 2015).

There are just two previously published studies on characterising cyathostomin species in

horses in Ireland. The first (Kinsella, 2002), used morphological species determination and the

second (Elghryani, 2018) used reverse line blots of the intergenic spacer gene region.

Elsewhere, internal transcribed spacer 2 (ITS-2) primers as described by (Gasser et al., 1993)

have been widely used as taxonomic markers for cyathostomins.   The focus of recent studies

have utilised community analysis tools, such as 'Nemabiome' application of ITS-2 sequencing

(Avramenko et al., 2015; Courtot et al., 2023; Halvarsson et al., 2023; Nielsen et al., 2022) .

We aimed to sequence the ITS-2 region from individual cyathostomins retrieved from

naturally infected horses in Ireland in a post-mortem study and to compare the community

composition from the colon and caecum to those described in previous studies in Ireland and

internationally. We also aimed to explore any difficulties in distinguishing species based on

ITS-2 sequencing, to investigate any trends in seasonality in the data, and to compare and

contrast two different methods of inferring taxonomy based on ITS-2 sequences, using BLAST

and IDTAXA methods.

2. Materials and Methods

2.1. Ethical approval

This was a post-mortem study and did not require any specific ethical approval.

 2.2. Study population

We conducted an abattoir survey between August 2018 and June 2020. A total of eleven

equine carcasses were selected at random for examination at two Irish abattoirs. In

accordance with food safety legislation, no horse had received anthelmintic treatment within

32 days of slaughter. The horses were assessed for clinical abnormalities by a veterinarian

prior to inclusion in the study as part of the normal pre-mortem inspection at the abattoir.

2.3. Sample processing

Each carcass was processed as per abattoir standard operating procedures. Sampling was

undertaken on the same day as euthanasia. To isolate the large intestine, the ileocecal

junction was located, tied off with cable ties and dissected. The liver, stomach, spleen,

pancreas, and small intestine were removed, and the large intestine was transferred from the

processing line to a dissection table.

Further cable ties were placed at the junction between the caecum and right ventral colon,

the junction between the right ventral and left ventral colon, the junction between the left

ventral colon and left dorsal colon, and the junction between the left dorsal and right dorsal

colon. An incision was made in each of the three intestinal segments (caecum, right ventral

colon (RVC) and left dorsal colon (LDC), and the luminal contents were emptied separately

into individual 10L graduated buckets. For each section, the mucosa was gently rinsed, and

the liquid was added to the bucket containing luminal contents. The contents were then

stirred whilst water was added to make a homogeneous suspension to a volume of 10L. Two

100ml aliquots were taken from each section, equating to 2% of the total volume. Faeces

were collected from the rectum of each horse and placed in a ziplock bag. All samples were

placed on ice for transfer to the laboratory.

2.4. Faecal Egg Counting

The Mini-FLOTAC technique (Cringoli, 2006) was utilised to count strongyle eggs in faeces.

Briefly, 45ml of saturated NaCL was added to 5g of faeces and the flotation chambers were

completely filled with the faecal suspension. After 10 minutes, the key was used to turn the

reading disc clockwise (about 90°), the sample was examined under the microscope and the

eggs per gram calculated. The sensitivity limit for the Mini-FLOTAC is 5 epg.

2.5. Luminal worm retrieval

Each aliquot representing 2% of the contents of the caecum, RVC and LDC was examined

individually using a modification of the Baermann method by enclosing them in a muslin

parcel suspended in water in a conical flask and leaving to sediment for 24 hours. The entire

sediment was then examined in detail (using aliquots forming a thin layer in a petri dish) under

a dissecting microscope. The remaining sediment was also examined under a dissecting

microscope.

The total number of worms present in each 200ml sample was enumerated and recorded.

Individual worms were removed manually and washed prior to storage. Each worm was

placed in a 2ml Eppendorf with >/=50ml PBS and stored at -20˚C. For each section a maximum

of 50 worms (range 1-50) was retained for subsequent examination from combined duplicate

samples. IBM SPSS (Version 27) was used for descriptive statistics. The Kruskal-Wallis test

was utilised for non-parametric data including faecal egg counts (FEC), numbers of worms and

species distribution

2.6. DNA Extraction and amplification

2.6.1. Genomic DNA extraction from individual nematodes

Individual worms were dissected into a minimum of three parts using a sterile needle. DNA

was then extracted from each worm using the Powerfecal Pro extraction kit (Qiagen, Product

number 51804), as per the standard operating protocol, with minor adaptations. Specifically,

samples were pulse vortexed, as part of the bead beating step, for a minimum of 45 minutes,

the dry spin after solution C5 was at 16,600 x G for three minutes, and solution C6 of the kit

was put through the filter membrane twice to increase elution yield (Bredtmann *et al.*, 2019).

After extraction, samples were stored at -20°C until PCR amplification.

2.6.2. PCR amplification

PCR reactions were performed in 50µl reaction volumes using ITS-2 primers as described by

Gasser *et al.*, (1993), (NC1 forward; 5'ACGTCTGGTTCAGGGTTGTT3' and NC2 reverse;

5'TTAGTTTCTTTTCCTCCGCT3').

The reaction mixture consisted of 10µl of Promega GoTaq Flexi Buffer,(product no. M780A)

1.5mM of Promega MgCL2 solution, 0.2mM Promega DNTP PCR nucleotide mix, (product

number U151A) 1µM of each forward and reverse primer (Eurofins Genomics) 1.25U of

Promega GoTaq G2 Flexi DNA Polymerase, (product no. M780A) 10µl template DNA and

15.75µl nuclease free water (Merck Millipore 693520). Amplification steps included initial

denaturation at 95°C for 2 minutes, then cycling steps starting at 95°C for 1 minute, annealing

at 55°C for 1 minute and extension at 72°C for 1 minute, for 40 cycles, followed by extension

at 72°C for 5 minutes.

For each run, a negative control containing 10µl nuclease-free water in place of DNA template

was included.

Amplicons were purified using the Qiagen Qiaquick PCR purification kit (Product.No. 28106)

and analysed through Sanger sequencing in both directions using the PCR primers.

Sequencing was outsourced to Eurofins Genomics, (Sandyford, Dublin 18, Ireland) and FASTA

files were returned to the research group for analysis.

2.7. Molecular taxonomy attribution

Aligned sequences were constructed from forward and reverse reads and compared with

published nematode sequences in the NCBI database using Clustal Omega.

Samples that did not yield a reliable BLAST result, (E value > E^{-10}, low query cover and

percentage identity <95%,) were removed from the dataset.

As we did not carry out morphological examination of recovered worms, all sequences are

uploaded as 'unidentified cyathostomin' on NCBI database and a note accompanying each

species outlining its closest hit on BLAST. These nucleotide sequence data is available in the

GenBank™ database under the accession numbers: **OR941129 - OR941242.**

2.7.1 Analysis of sequences using Nematode ITS2 database and IDTAXA

Sequences obtained from individual worms were also compared with data in an opensource

ITS2 rDNA database for both parasitic and free living nematodes (Wokentine et al., 2020). To

infer taxonomy, IDTAXA (Murali et al., 2018) as a component of the DECIPHER (2.22.0)

package in R (v4.1.2), and phyloseq package version (1.38.0)   through R studio

(v2023.12.0.369) was used within the DADA 2 pipeline as described at www.nemabiome.ca .

2.8 Phylogenetic analiysis

MEGA11, and iTOL (https://itol.embl.de/) were used for phylogenetic tree creation. MEGA11

was used for MUSCLE alignment and trimming of sequences. A tree was created using the

Maximum likelihood algorithm with 1,000 bootstrap replications and pairwise deletion for

gaps. (Tamura et al., 2021). The evolutionary history was inferred by using the Maximum

Likelihood method and Tamura-Nei model(Tamura and Nei, 1993). The bootstrap consensus

tree inferred from 1000 replicates (Felsenstein, 1985) is taken to represent the evolutionary

history of the taxa analyzed. Branches corresponding to partitions reproduced in less than

50% bootstrap replicates are collapsed. Initial tree(s) for the heuristic search were obtained

automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise

distances estimated using the Tamura-Nei model, and then selecting the topology with

superior log likelihood value. Seven sequences were removed at the alignment section of the

phylogenetic analysis as they were too short when aligned with other species. There were a

total of 203 positions in the final dataset. Visualisation of the phylogenetic tree was

conducted using iTOL V6.8.1(Letunic and Bork, 2021)

3.   Results

3.1. Study cohort and worm recovery

Table 1 provides the details of horses included in the study, together with their strongyle FEC.

number of worms, and the areas from which worms for molecular identification were

retained. The horses included in the study ranged from yearlings to twenty-three years old

with a mean of 7.7 years of age, and a median of 6 years. There were four females and seven

males. Five were Thoroughbred, and six Irish Sport Horse. FEC ranged from 0-1,670 eggs per

gram (epg).   Nematodes were retrieved from two horses with a FEC of 0. Conversely, high

FEC did not always correlate with high luminal burden of nematodes. Numbers of worms

recovered from the caecum were lower than those recovered from LDC (P=0.009). No other

significant differences in worm recovery between compartments was noted.   Up to 14

individual worms from each intestinal compartment, per horse, had taxonomy inferred to

species level by BLAST and IDTAXA.   For all but one horse, worms from two or three

compartments were included in the taxonomic analysis.

3.2. BLAST analysis of ITS-2 amplified sequences

A total of 116 samples were extracted, amplified and compared using the BLAST algorithm

with sequences in the NCBI database.   Another 43 samples did not provide adequate

sequence information for analysis Of the 116 sequences analysed, 43 samples had a 100%

identity score with their presumed species. The remaining samples had taxonomy inferred

based on similarity scores of > 95% similarity, and E,= 10^{-10} . The amount of samples that fell

within the percentage similarity values outlined previously (Mitchell et al., 2019) are outlined

as follows: >99% n= 46, >98% n= 16, >97% n= 5, >96% n= 3.. (Supplementary File).

3.3 IDTAXA analysis of amplified sequences and comparison with BLAST results.

A total of 103 of the original 116 consensus Sanger sequences gave valid outputs from the

nemabiome.ca workflow using the ITS2 database using IDTAXA to infer taxonomy.   Thirteen

sequences did not pass quality control cut-offs for the IDTAXA algorithm because they fell

below the 60% taxonomic classification confidence threshold. Of the 103 sequences meeting

the threshold, 83 (80.5%) were assigned to species level by IDTAXA and gave the same result

as BLAST using >95% similarity, (Table 2). Of the remainder, 13 were identified only to genus

level by IDTAXA, but as the same genus as BLAST. A further six were presumptively identified

as *Cylicostephanus calicatus* by BLAST and *Coronocylus coronatus* by IDTAXA.   One sequence

identified on BLAST as *Cor. coronatus* was assigned as unclassified *Cyathostomum* spp. using

IDTAXA.

3.3. Species recovered, abundance and prevalence

The species abundance in each of the three intestinal compartments is shown in Figure 1.

Cor. coronatus was the most abundant species recovered (29%).   Between them, *Cor.*

coronatus, *Cylicocyclus nassatus* and *Cylicostephanus longibursatus* accounted for more than

half (54%) of the entire dataset. Although there was no significant difference in species

distribution across the three intestinal compartments, *Cylicocyclus insigne* was equally

abundant to *Cor. coronatus* in the LDC.   This compartment also had the most diverse range

of species (n=11), with the caecum having the least diversity (n=8).

Figure 2 outlines the composition per horse of presumed species. It outlines the species found

in each of the 11 horses, represented as a percentage of the total samples amplified from

each individual. The actual number of worms that each segment represents is also included

in the figure, overlaying each segment. This figure provides an insight into the species

composition differences between individuals from the samples that were obtained in this

study. For the purposes of this analysis, *Strongylus vulgaris* (horse 5) was included in this

analysis.

The prevalence of individual species and the richness of cyathostomin communities in this

study population is illustrated in Figure 3. *Cor. coronatus* and *Cyc. longibursatus* were the

most prevalent, found in seven of the eleven horses (63.63% prevalence), *Cyc. nassatus* was

present in six horses (54.54%) and *Cylicostephanus goldi* in five horses (45.45%). *Cyc. insigne,*

Cylicostephanus calicatus, Cylicocyclus ashworthi and *Cyathostomum catinatum* were each

recorded in four horses, at a prevalence of 36.36%. *Cyathostomum pateratum* was found in

three horses (27.27% prevalence) with *Cylicostephanus minutus, Coronocyclus. labiatus* and

Cylicocyclus leptostomum in one animal each, comprising 9.09% prevalence.

Minimal richness from the samples taken ranged from two to seven species per individual,

with a mean of 4.27 +/- 0.604 (SEM). The modal species richness was two and five species

equally, with a median species richness of five within this dataset. Over 50% (54.5%) of the

horses sampled had >5 cyathostomin species present. One large strongyle, *Strongylus*

vulgaris, was identified (Horse 5) but not included in the dataset.

3.4 Analysis in relation to previous local and global data

Next, (Figure 4) we compared the composition of cyathostomin species found in our study

with data from a previous study in Ireland (Kinsella, 2002) and a global meta-analysis (Bellaw

and Nielsen, 2020). Twelve cyathostomin species were common to our study, Kinsella, 2002,

and the global meta-analysis. Notably, *Cor. labratus* was not identified in either of the Irish

datasets, but features in the global meta-analysis. Our study presumptively identified 12 of

the 13 most prevalent global species, (Bellaw and Nielsen, 2020), using two different

methods of inferring taxonomy from ITS-2 sequences.

3.6. Phylogenetic analysis of species

Sequences were used for construction of a Maximum Likelihood phylogenetic tree (Figure 5)

which was rooted at the midpoint. Figure 5 unveiled some interspecies occurrences within

branches, notably between *Cyc. calicatus* and *Cor. coronatus*.

4. Discussion

There was no direct relationship between strongyle faecal egg count and worm burden in the

carcases examined, which was not unexpected (Chapman *et al*., 2003a; Chapman *et al*.,

2003b; Nielsen *et al*., 2010).   We recovered worms from horses with a 0 EPG count, which

has been well-documented previously (Gibson, 1953; Love *et al*., 1999; Matthews, 2008;

Walshe *et al*., 2019; Walshe *et al*., 2020).

We did not attempt morphological identification of the worms recovered in our study,

because of the inherent difficulties in doing so and our lack of expertise in this area. Our

taxonomic classifications, therefore, are inferred solely on the basis of sequence information.

Clearly, reliance on molecular data only for this purpose raises the possibility of errors due to

errors in sequence databases, judgement calls when there is similarity to more than one

species, and the inability to distinguish between some species on the basis of one locus (in

this case ITS-2) alone. In order to validate and provide for greater reliability from our results,

we used two distinct methods for inferring taxonomy from sequence information, BLAST, and

IDTAXA.

IDTAXA is a method of taxonomic classification based on sequences of marker genes from

organisms found in microbiome samples or other sources. It involves assigning sequences to

a reference taxonomy containing sequences from known members of a reference dataset,

and avoids overclassification of sequences from organisms which may not be present in the

reference set (Murali et al., 2018).   A database of ITS2 rDNA sequences from both parasitic

and free-living nematodes is publicly available, (Workentine et al., 2020) and is now widely

used in "nemabiome" studies in order to characterise a mixed population of parasitic

nematodes within a single host, or host population, including in horses (Ghafar et al., 2022;

Halvarsson et al., 2023; Poissant et al., 2021; Sargison et al., 2022) following high-throughput

parallel sequencing and metabarcoding.   However, the same bioinformatics pipeline using

DADA 2 can be used to assign taxonomy to data obtained by conventional Sanger sequencing.

The results obtained in our study from BLAST of individual sequences and from the DADA 2

IDTAXA output gave identical results to species level for 82 (80.6%) of worms and results

consistent to genus level for a further 12 (12.6%). For the remaining seven sequences, (6.8%)

there was inconsistency in distinguishing between *Cor. coronatus* and *Cyl. calicatus*, reflecting

difficulties in using ITS-2 sequences to distinguish between these potentially cryptic species

in other studies (Bredtmann et al., 2019,(Louro et al., 2021).

Overall, given the consistency between taxonomic assignment by BLAST, and by IDTAXA, in

this study, together with the difficulty and potential for error in morphological identification

of strongyle worms of horses, we suggest that comparison of two sequence-based methods

of taxonomic classification is useful in further determining the taxonomic relationships

between these parasites.   The characterisation of additional loci may also be useful in some

cases (Courtot et al., 2023)

This study detected sequences similar to twelve of the thirteen most common species globally

(Figure 4), as compared to Bellaw and Nielsen's meta-analysis (Bellaw and Nielsen, 2020). It

should be noted that prevalence as outlined in this study is based on a 2% sample of contents

of each intestinal compartment sampled and thus is likely to underestimate prevalence.

Among the nineteen species identified in the morphological study by Kinsella (2002), we

detected twelve (Figure 4). Notably, the prevalence of *Cya. catinatum* was 36% in our study,

in contrast to 100% prevalence in Kinsella's work and 94% in the global meta-analysis.

Elghryani (2018), found *Cyc. nassatus* as the most prevalent among the twelve species

identified. The high abundance and prevalence of *Cyc. nassatus,* occurring in 64% (n=7) of the

horses sampled, is consistent among all three Irish studies and the global meta-analysis. The

prevalence findings between the three studies from Ireland broadly reflect one another. A

notable observation was the absence of the twelfth most prevalent species as identified by

(Bellaw and Nielsen, 2020), *Cor. labratus,* which was absent from the Irish studies, and has

not been detected either in some studies in similar climates-Wales and Scotland(Mitchell et

al., 2019; Sargison et al., 2022). This species has however, been recorded in England and France (Cwiklinski *et al.*, 2012; Peachey *et al.*, 2018).

Cyc. calicatus and *Cor. coronatus* are interspersed within the ML tree. This reflects their close relationship, as previously identified, and the limitations of using the ITS-2 locus to differentiate the subsets of these species (Courtot *et al.*, 2023; Hung *et al.*, 2000; Louro *et al.*, 2021).

Across the three intestinal compartments examined in this study, the number of worms retrieved from the caecum was significantly lower than that recovered from LDC (P =0.009). This was also observed in a previous Irish study(Kinsella, 2002). (Ogbourne, 1976), assessed abundance of cyathostomins postmortem and found 50% in the ventral colon and 45% in the dorsal colon. The worm burden in the caecum represented only 5% of the overall cyathostomin burden. Similarly, (Stancampiano *et al.*, 2010), observed that the majority of small strongyles were concentrated in the ventral colon (67.34%) and less frequently in the caecum (27.52%) and dorsal colon (5.14%). While this distribution was also reflected by Kinsella (2002), the converse was found for encysted stages of cyathostomins, as recorded by (Nielsen *et al.*, 2021) who documented 40% of EL3/LL3/L4 in the caecum, and 38%/22% in the ventral and dorsal colon respectively. This recurrent recording of biological preference of luminal cyathostomins for the colon over the caecum, and the inverse for encysted larvae may reflect the slower transit time of the colon and the natural peristalsis of food through the digestive tract, enabling passive transport and a longer exposure to the intestinal wall necessary to facilitate larval encystment (Geor *et al.*, 2013; Miyaji *et al.*, 2008).

Over half of the horses in this study had more than five cyathostomin species recovered from

them (Figure 2).(Abbas et al., 2023; Reid et al., 1995)

5. **Limitations**

The most important limitation of this study is sample size. This is a descriptive study, which

affects the potential for statistical power. Conclusions from observations within any study

with a limited sample size must take this into account. Results are considered in the context

that only two nematodes were retrieved for sampling from horse eight. Notably, it is clear

from this research and other studies that the ITS-2 locus cannot resolve all isolates of *Cor.*

coronatus and *Cyc. calicatus,* pointing to the need for a different locus for further molecular

work, such as COX1, or a multi-locus approach (Courtot *et al*., 2023; Ghafar *et al*., 2022). We

acknowledge also the value of examining seasonality in relation to cyathostomin species

prevalence and abundance, as well as potential interactions, positive or negative between

species. Variability of seasonal patterns among species, in particular, is an understudied facet

of cyathostomin biology which may be crucial for comprehensive characterisation of

pathogenesis and epidemiology of cyathostomin-related disease, as this tends to have a

seasonal distribution of presentation(Abbas et al., 2023; Reid et al., 1995).   Other studies,

(Salle et al., 2018) have begun to explore interactions among cyathostomin communities, and

this also has a potential bearing on the pathogenesis and epidemiology of cyothostomin-

related disease. Our study was too small to make reliable observations on these points.

6. Conclusions

This study has produced new insights into cyathostomin species occurrence, interactions and, community composition in horses in Ireland, with comparisons to previous studies in this locality and a global meta-analysis. It also displays a high level of agreement in assigning taxonomy from ITS-2 sequences, based on BLAST and IDTAXA, which we believe is useful in increasing the reliability of studies using molecular information to study cyathostomin communities.

Post-mortem analysis and identification in naturally-infected hosts, as conducted here, will continue to have a role to play in advancing understanding of cyathostomin biology and pathogenesis, in parallel with population-based approaches. More extensive studies involving both population and post-mortem studies are warranted in order to enhance risk assessment, biomarker-based diagnostic tools and treatment protocols (Walshe *et al.*, 2020). It is also useful in investigating potentially cryptic species as seen in this study with *Cor. coronatus* and *Cyc. calicatus.*

Studies on cyathostomin population dynamics are a crucial step in understanding the complex picture of cyathostomin infections in the face of pervasive anthelmintic resistance and their potential to advance mechanistic understanding of gut health in horses.

Author contributions

Conceptualization, G.M., N.W., O.B.; writing- original draft preparation, O.B. – review and editing, G.M., N.W., O.B. and A.Z.; laboratory work and bioinformatics- O.B., D.G., J.C.; laboratory contributions, L.K., O.BX., D.McS., F.ON., S.D., BR.L. All authors have read and agreed to the published version of this manuscript.

Funding:

This research was funded by a Clinical Primer grant to Nicola Walshe from the Wellcome

Institutional Strategic Support fund, and by UCD's own funds.

Acknowledgements:

We would like to acknowledge the valuable input and assistance of Amanda Lawlor, Senior

Technical Officer, and Dr. Nagwa Elghyrani, Adjunct Research Fellow, UCD School of

Veterinary Medicine

Conflict of interest:

The authors declare no conflict of interest.

Bibliography:

Abbas, G., Ghafar, A., Bauquier, J., Beasley, A., Ling, E., Gauci, C.G., El-Hage, C., Wilkes, E.J., McConnell, E., Carrigan, P., 2023. Prevalence and diversity of ascarid and strongylid nematodes in Australian Thoroughbred horses using next-generation sequencing and bioinformatic tools. Veterinary parasitology 323, 110048.

Avramenko, R.W., Redman, E.M., Lewis, R., Yazwinski, T.A., Wasmuth, J.D., Gilleard, J.S., 2015. Exploring the gastrointestinal "nemabiome": deep amplicon sequencing to quantify the species composition of parasitic nematode communities. PLoS One 10, e0143559.

Bellaw, J.L., Nielsen, M.K., 2020. Meta-analysis of cyathostomin species-specific prevalence and relative abundance in domestic horses from 1975–2020: emphasis on geographical region and specimen collection method. Parasites & Vectors 13, 1-15.

Boisseau, M., Mach, N., Basiaga, M., Kuzmina, T., Laugier, C., Sallé, G., 2023. Patterns of variation in equine strongyle community structure across age groups and gut compartments. Parasites & Vectors 16, 64.

Boxell, A., Gibson, K., Hobbs, R., Thompson, R., 2004. Occurrence of gastrointestinal parasites in horses in metropolitan Perth, Western Australia. Australian Veterinary Journal 82, 91-95.

Bredtmann, C.M., Krücken, J., Murugaiyan, J., Balard, A., Hofer, H., Kuzmina, T.A., von Samson‐Himmelstjerna, G., 2019. Concurrent proteomic fingerprinting and molecular analysis of cyathostomins. Proteomics 19, 1800290.

Chapman, M., Kearney, M., Klei, T., 2003a. Equine cyathostome populations: accuracy of species composition estimations. Veterinary parasitology 116, 15-21.

Chapman, M.R., French, D.D., Klei, T.R., 2003b. Prevalence of strongyle nematodes in naturally infected ponies of different ages and during different seasons of the year in Louisiana. Journal of Parasitology 89, 309-314.

Courtot, É., Boisseau, M., Dhorne-Pollet, S., Serreau, D., Gesbert, A., Reigner, F., Basiaga, M., Kuzmina, T., Lluch, J., Annonay, G., 2023. Comparison of two molecular barcodes for the study of equine strongylid communities with amplicon sequencing. PeerJ 11, e15124.

Cringoli, G., 2006. FLOTAC, a novel apparatus for a multivalent faecal egg count technique. Parassitologia 48, 381-384.

Cwiklinski, K., Kooyman, F., Van Doorn, D., Matthews, J., Hodgkinson, J., 2012. New insights into sequence variation in the IGS region of 21 cyathostomin species and the implication for molecular identification. Parasitology 139, 1063-1073.

Dowdall, S., Matthews, J., Mair, T., Murphy, D., Love, S., Proudman, C., 2002. Antigen-specific IgG (T) responses in natural and experimental cyathostominae infection in horses. Veterinary parasitology 106, 225-242.

Duncan, J., Bairden, K., Abbott, E., 1998. Elimination of mucosal cyathostome larvae by five daily treatments with fenbendazole. Veterinary Record 142, 268-271.

Elghryani, N., 2018. The Prevalence and Anthelmintic Sensitivity of Gastrointestinal Nematodes of Horses in Ireland. University College Dublin,

Felsenstein, J., 1985. Confidence limits on phylogenies: an approach using the bootstrap. evolution 39, 783-791.

Gasser, R.B., Chilton, N.B., Hoste, H., Beveridge, I., 1993. Rapid sequencing of rDNA from single worms and eggs of parasitic helminths. Nucleic acids research 21, 2525.

Geor, R.J., Harris, P., Coenen, M., 2013. Equine applied and clinical nutrition: health, welfare and performance. Elsevier Health Sciences.

Ghafar, A., Abbas, G., Beasley, A., Bauquier, J., Wilkes, E.J., Jacobson, C., McConnell, E., El-Hage, C., Carrigan, P., Cudmore, L., 2022. Molecular diagnostics for gastrointestinal helminths in equids: Past, present and future. Veterinary Parasitology, 109851.

Gibson, T., 1953. The effect of repeated anthelmintic treatment with phenothiazine on the faecal egg counts of housed horses, with some observations on the life cycle of Trichonema spp. in the horse. Journal of Helminthology 27, 29-40.

Giles, C., Urquhart, K., Longstaffe, J., 1985. Larval cyathostomiasis (immature trichonema‐induced enteropathy): A report of 15 clinical cases. Equine veterinary journal 17, 196-201.

Halvarsson, P., Grandi, G., Hägglund, S., Höglund, J., 2023. Gastrointestinal parasite community structure in horses after the introduction of selective anthelmintic treatment strategies. Veterinary Parasitology, 110111.

Hinney, B., Wirtherle, N.C., Kyule, M., Miethe, N., Zessin, K.-H., Clausen, P.-H., 2011. Prevalence of helminths in horses in the state of Brandenburg, Germany. Parasitology research 108, 1083-1091.

Hodgkinson, J.E., Lichtenfels, J.R., Mair, T., Cripps, P., Freeman, K., Ramsey, Y., Love, S., Matthews, J., 2003. A PCR–ELISA for the identification of cyathostomin fourth-stage larvae from clinical cases of larval cyathostominosis. International Journal for Parasitology 33, 1427-1435.

Kinsella, J.L., 2002. Species Composition, Distribution, and Abundance, with Monthly Variations of Cyathostominea Within the Equid in Ireland, 1999/2000, and a Preliminary Analysis of Proteolytic Activity of Excretory Secretory Products from Cyathostominea. University College Dublin,

Letunic, I., Bork, P., 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. Nucleic acids research 49, W293-W296.

Lichtenfels, J.R., Kharchenko, V.A., Dvojnos, G.M., 2008. Illustrated identification keys to strongylid parasites (Strongylidae: Nematoda) of horses, zebras and asses (Equidae). Veterinary parasitology 156, 4-161.

Louro, M., Kuzmina, T.A., Bredtmann, C.M., Diekmann, I., de Carvalho, L.M.M., von Samson-Himmelstjerna, G., Krücken, J., 2021. Genetic variability, cryptic species and phylogenetic relationship of six cyathostomin species based on mitochondrial and nuclear sequences. Scientific Reports 11, 8245.

Love, S., Murphy, D., Mellor, D., 1999. Pathogenicity of cyathostome infection. Veterinary parasitology 85, 113-122.

Lyons, E.T., Tolliver, S., Drudge, J., 1999. Historical perspective of cyathostomes: prevalence, treatment and control programs. Veterinary Parasitology 85, 97-112.

Matthews, J., 2008. An update on cyathostomins: anthelmintic resistance and worm control. Equine Veterinary Education 20, 552-560.

Mitchell, C.J., O'SULLIVAN, C.M., Pinloche, E., Wilkinson, T., Morphew, R.M., McEwan, N.R., 2019. Using next-generation sequencing to determine diversity of horse intestinal worms: identifying the equine 'nemabiome'. Journal of equine science 30, 1-5.

Miyaji, M., Ueda, K., Nakatsuji, H., Tomioka, T., Kobayashi, Y., Hata, H., Kondo, S., 2008. Mean retention time of digesta in the different segments of the equine hindgut. Animal Science Journal 79, 89-96.

Morariu, S., Mederle, N., Badea, C., Dărăbuş, G., Ferrari, N., Genchi, C., 2016. The prevalence, abundance and distribution of cyathostomins (small stongyles) in horses from Western Romania. Veterinary Parasitology 223, 205-209.

Murphy, D., Love, S., 1997. The pathogenic effects of experimental cyathostome infections in ponies. Veterinary parasitology 70, 99-110.

Nielsen, M.K., Baptiste, K.E., Tolliver, S., Collins, S., Lyons, E., 2010. Analysis of multiyear studies in horses in Kentucky to ascertain whether counts of eggs and larvae per gram of feces are reliable indicators of numbers of strongyles and ascarids present. Veterinary Parasitology 174, 77-84.

Nielsen, M.K., Lyons, E.T., 2017. Encysted cyathostomin larvae in foals–progression of stages and the
effect of seasonality. Veterinary parasitology 236, 108-112.

Nielsen, M.K., Martin, A.N., Scare, J.A., Steuer, A.E., 2021. Precision and spatial variation of
cyathostomin mucosal larval counts. Veterinary Parasitology 290, 109349.

Nielsen, M.K., Steuer, A.E., Anderson, H.P., Gavriliuc, S., Carpenter, A.B., Redman, E.M., Gilleard, J.S.,
Reinemeyer, C.R., Poissant, J., 2022. Shortened egg reappearance periods of equine
cyathostomins following ivermectin or moxidectin treatment: morphological and molecular
investigation of efficacy and species composition. International Journal for Parasitology 52,
787-798.

Ogbourne, C.P., 1976. The prevalence, relative abundance and site distribution of nematodes of the
subfamily Cyathostominae in horses killed in Britain. Journal of Helminthology 50, 203-214.

Peachey, L., Molena, R., Jenkins, T., Di Cesare, A., Traversa, D., Hodgkinson, J., Cantacessi, C., 2018.
The relationships between faecal egg counts and gut microbial composition in UK
Thoroughbreds infected by cyathostomins. International journal for parasitology 48, 403-412.

Peregrine, A.S., McEwen, B., Bienzle, D., Koch, T.G., Weese, J.S., 2006. Larval cyathostominosis in
horses in Ontario: an emerging disease? The Canadian Veterinary Journal 47, 80.

Poissant, J., Gavriliuc, S., Bellaw, J., Redman, E.M., Avramenko, R.W., Robinson, D., Workentine, M.L.,
Shury, T.K., Jenkins, E.J., McLoughlin, P.D., 2021. A repeatable and quantitative DNA
metabarcoding assay to characterize mixed strongyle infections in horses. International
Journal for Parasitology 51, 183-192.

Reid, S., Mair, T., Hillyer, M., Love, S., 1995. Epidemiological risk factors associated with a diagnosis of
clinical cyathostomiasis in the horse. Equine Veterinary Journal 27, 127-130.

Sargison, N., Chambers, A., Chaudhry, U., Júnior, L.C., Doyle, S.R., Ehimiyein, A., Evans, M., Jennings,
A., Kelly, R., Sargison, F., 2022. Faecal egg counts and nemabiome metabarcoding highlight
the genomic complexity of equine cyathostomin communities and provide insight into their
dynamics in a Scottish native pony herd. International Journal for Parasitology 52, 763-774.

Stancampiano, L., Gras, L.M., Poglayen, G., 2010. Spatial niche competition among helminth parasites
in horse's large intestine. Veterinary Parasitology 170, 88-95.

Steinbach, T., Bauer, C., Sasse, H., Baumgärtner, W., Rey-Moreno, C., Hermosilla, C., Damriyasa, I.M.,
Zahner, H., 2006. Small strongyle infection: Consequences of larvicidal treatment of horses
with fenbendazole and moxidectin. Veterinary Parasitology 139, 115-131.

Tamura, K., Nei, M., 1993. Estimation of the number of nucleotide substitutions in the control region
of mitochondrial DNA in humans and chimpanzees. Molecular biology and evolution 10, 512-
526.

Tamura, K., Stecher, G., Kumar, S., 2021. MEGA11: molecular evolutionary genetics analysis version
11. Molecular biology and evolution 38, 3022-3027.

Walshe, N., Duggan, V., Cabrera-Rubio, R., Crispie, F., Cotter, P., Feehan, O., Mulcahy, G., 2019.
Removal of adult cyathostomins alters faecal microbiota and promotes an inflammatory
phenotype in horses. International journal for parasitology 49, 489-500.

Walshe, N., Mulcahy, G., Crispie, F., Cabrera‐Rubio, R., Cotter, P., Jahns, H., Duggan, V., 2021.
Outbreak of acute larval cyathostominosis‐A "perfect storm" of inflammation and dysbiosis.
Equine Veterinary Journal 53, 727-739.

Walshe, N., Mulcahy, G., Hodgkinson, J., Peachey, L., 2020. No Worm is an island; the influence of
commensal gut microbiota on cyathostomin infections. Animals 10, 2309.

Workentine, M.L., Chen, R., Zhu, S., Gavriliuc, S., Shaw, N., Rijke, J.d., Redman, E.M., Avramenko, R.W.,
Wit, J., Poissant, J., 2020. A database for ITS2 sequences from nematodes. BMC genetics 21,
1-4.

Table 1: Horse details, FEC and number of worms recovered and speciated from each intestinal
compartment. RVC= Right Ventral Colon and LDC = Left Dorsal Colon. * = P < 0.05. Mean calculations
are +/- the SEM (standard error of the mean). The number of worms recovered accounts for a 1%
subsection of the total luminal contents of each animal sampled. Numbers of worms recovered from
the caecum were lower than those recovered from LDC (P=0.009). No significant difference was
recorded between RVC or from composite RVC and LDC colon to the caecum (P=0.291). FEC
represents eggs per gram (epg) of faecal material).

Horse Details						Number of worms recovered			Speciated Worms		
Horse	Age	Breed	Sex	Sampling Date	FEC (epg)	RVC	LDC	Caecum	RVC	LDC	Caecum
1	16	ISH	F	Sep-18	1550	79	188	10	2	9	-
2	8	ISH	M (gelding)	Sep-18	0	11	3	1	5	-	-
3	1	TB	M (colt)	Sep-18	1670	205	155	12	8	2	4
4	10	ISH	F	Oct-18	25	201	245	23	3	13	-
5	1	TB	M (colt)	Oct-18	905	590	33	16	3	-	4
6	6	ISH	M (colt)	Feb-20	225	2	26	0	1	4	-
7	2	TB	M (colt)	Feb-20	800	1	249	2	-	4	1
8	12	ISH	M	Mar-20	50	0	2	0	-	2	-
9	23	TB	F	Mar-20	85	9	24	1	3	3	-
10	3	TB	F	Jun-20	715	1410	3328	33	8	5	14
11	3	ISH	M	Jun-20	0	40	56	53	4	5	9
Total	-	-	-	-	-	2548	4309	151	37	47	32
Mean	7.7+/-2.124				547.73 +/- 188.83	231.63 +/- 129.21	391 +/- 295*	13.73 +/- 5.1*	3.36 +/- 0.83	4.27 +/- 1.16	2.91 +/- 1.4
Range	1-23	-	-	-	0-1670	0-1410	2-3328	0-53	0-8	0-13	0-14

571 **Table 2.** Comparison of BLAST and IDTAXA identification of ITS sequences. We compared the
572 degree to which BLAST, with >95% sequence identity, and the IDTAXA algorithm gave comparable
573 results using our ITS-2 sequences.

Sequences for which taxonomy inferred by BLAST	Agreement on Species using IDTAXA	Agreement on Genus level, IDTAXA	Inability to distinguish between *Cor. coronatus* and *Cyl. calicatus*	Species BLAST, different genus IDTAXA	Did not pass IDTAXA Quality Threshold
Cor. coronatus	26	2	7	1 (unclassified Cyathostomum).	5
Cyl. nassatus	15				
Cyl. goldi	11				
Cyl. insigne	11				
Cyl. longibursatus	9	1		3	3
Cyl. ashworthi	3	6			
Cya. pateratum	2	1			
Cya. catinatum	2	3		2	2
Cor. labiatus	1				
Cyl. leptostomus	1				
Cyl. minutus	1				
S. vulgaris	1				
Cyl. calicatus					2

Figure 1: Species abundance data from samples obtained from Right Ventral Colon (a) Left Dorsal Colon (b), Caecum (c) and across all three compartments (d). *Cor. coronatus* was the most abundant species recovered, at 29% of total nematodes speciated. Between them, *Cor. coronatus, Cyc. nassatus* and *Cyc. longibursatus* accounted for more than half (54%) of the entire dataset. The LDC had the most diverse range of species (n=11), with the caecum having the least diversity (n=8).

Figure 2: 100% stacked bar graph of the species distribution per horse. Species richness ranged from 2-7 per horse. Each horse represents a stacked column, identified on the X- axis. The species are represented by coloured sections illustrated as a percentage of the nematodes obtained from each horse. The actual number of samples from each species is also illustrated within each segment.

Figure 3: Prevalence of cyathostomin species as compared with a previous study in Ireland (Kinsella 2002) and a global meta-analysis (Bellaw and Nielsen, 2020). We identified sequences similar to twelve cyathostomin species recorded in both the previous Irish study and the global meta-analysis. There were no species identified in either study in Ireland that were not identified within the global meta-analysis. Generally, the most prevalent species in the two previous studies were also highly prevalent in our study.

Figure 4: Full phylogenetic tree of sequences from recovered worms. IDtaxa similarity sequences for individual cyathostomins were used for phylogenetic analysis. MEGA11 was used for MUSCLE alignment and trimming of sequences. A tree was created using the Maximum likelihood algorithm with 1,000 bootstrap replications and pairwise deletion for gaps. This tree unveiled species that located within groupings of other species, notably *Cylicostephanus calicatus* and *Coronocyclus coronatus.*